Stage-dependent behavioural changes but early castration induced by the acanthocephalan parasite *Polymorphus minutus* in its *Gammarus pulex* intermediate host.

Yann Bailly, Frank Cézilly, Thierry Rigaud

► **To cite this version:**

Yann Bailly, Frank Cézilly, Thierry Rigaud. Stage-dependent behavioural changes but early castration induced by the acanthocephalan parasite *Polymorphus minutus* in its *Gammarus pulex* intermediate host.. *Parasitology*, 2018, 145 (3), pp.260-268. 10.1017/S0031182017001457 . hal-01729162

HAL Id: hal-01729162

<https://hal.science/hal-01729162>

Submitted on 28 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Stage-dependent behavioural changes but early castration induced by the
2 acanthocephalan parasite *Polymorphus minutus* in its *Gammarus pulex*
3 intermediate host.

4

5

6 YANN BAILLY, FRANK CÉZILLY and THIERRY RIGAUD *

7

8 *Université Bourgogne Franche-Comté, Laboratoire Biogéosciences, UMR CNRS 6282, équipe Ecologie*
9 *Evolutive, 6 boulevard Gabriel, 21000 Dijon, France*

10

11 Running title : Castration and behavioural alteration in a gammarid

12

13 Corresponding author :

14 Thierry Rigaud
15 Université Bourgogne Franche-Comté,
16 Laboratoire Biogéosciences, UMR CNRS 6282,
17 Equipe Ecologie Evolutive,
18 6 boulevard Gabriel,
19 21000 Dijon,
20 France

21

22 thierry.rigaud@u-bourgogne.fr

23 +00 33 380 39 39 45

24

25

26 SUMMARY

27 Multidimensionality in parasite-induced phenotypic alterations (PIPA) has been observed in a large
28 number of host-parasite associations, particularly in parasites with complex life cycles. However, it is
29 still unclear whether such a syndrome is due to the successive activation of independent PIPAs, or
30 results from the synchronous disruption of a single mechanism. The aim of the present study was to
31 investigate the onset and progression of two PIPAs (a behavioural alteration: reversion of geotaxis,
32 and castration) occurring in the crustacean amphipod *Gammarus pulex* infected with the
33 acanthocephalan *Polymorphus minutus*, at different parasite developmental stages. Modifications of
34 geotaxis in hosts differed according to the parasite developmental stage. Whereas the cystacanth
35 stage induced a negative geotaxis (exposing the gammarid to predation by birds, the definitive
36 hosts), the acanthella stage, not yet infective for the definitive host, induced a stronger positive
37 geotaxis (presumably protecting gammarids from bird predation). In contrast, castration was almost
38 total at the acanthella stage, with no significant variation in the intensity according to parasite
39 maturation. Finally, no significant correlation was found between the intensity of behavioural
40 changes and the intensity of castration. We discuss our results in relation with current views on the
41 evolution of multidimensionality in PIPA.

42

43 Key words: Behavioural alterations, castration, infection syndrome, Amphipoda, Acanthocephala

44

45 KEY FINDINGS

- 46 • Acanthocephalans induce several phenotypic alterations in their amphipods hosts
- 47 • Alteration of host geotaxis and castration were studied simultaneously
- 48 • The two alterations were not parallel in time, nor were their intensities correlated accross
- 49 individuals
- 50 • Immature and mature parasites affected host geotaxis in opposite directions
- 51 • Although the two alterations appear to be unrelated in time, they may still depend on a
- 52 common mechanism
- 53
- 54

55 INTRODUCTION

56 Many parasite taxa have complex life cycles, with different host species harboring different
57 parasite developmental stages. Adult parasites typically mate and reproduce inside a definitive host,
58 from which eggs or larvae are released into the external environment. Parasites can then infect the
59 intermediate host(s), where they generally grow through several developmental stages before
60 becoming infective to the definitive host. Completion of the parasite's life cycle often occurs
61 trophically, through predation of the intermediate host by the definitive one. Complex life-cycles are
62 supposed to induce strong selective pressures for parasites to optimize their transmission from one
63 host to the other. Indeed, many parasites induce various alterations in the phenotype of their
64 intermediate hosts, some of them appearing to increase parasites' fitness at the expense of that of
65 their hosts. Such phenomena are regularly interpreted in the framework of the 'parasite
66 manipulation' hypothesis (Moore, 2002; Thomas *et al.* 2005; but see Cézilly *et al.* 2010). In particular,
67 numerous parasites induce phenotypic alterations (hereafter PIPA for Parasite-Induced Phenotypic
68 Alterations), including changes in the anti-predatory behaviour of their intermediate hosts, thereby
69 possibly increasing trophic transmission (Moore, 2002; Thomas *et al.* 2005; Poulin, 2010; see
70 however Perrot-Minnot *et al.* 2012).

71 Most often, numerous physiological and behavioural traits are simultaneously modified,
72 including some with no obvious consequences in terms of trophic transmission (Thomas *et al.*, 2010),
73 generating a parasite-induced infection syndrome (Cézilly and Perrot-Minnot, 2010; Cézilly *et al.*
74 2013). Among the traits belonging to such syndromes is parasitic castration (Baudoin, 1975). The
75 reduction or suppression of host reproduction by a parasitic infection is generally considered an
76 evolutionary strategy by which the parasite increases its own fitness at the expense of that of its
77 host (Baudoin, 1975; Jensen *et al.* 2006; Laferty and Kuris, 2009; but see Hurd, 2001 for other
78 interpretations).

79 Both behavioural and reproductive PIPA may then be constitutive of an infection syndrome.
80 Indeed, Cézilly and Perrot-Minnot (2010) highlighted that numerous parasite taxa inducing

81 behavioural alterations also induce castration. However, because these different PIPAs have been
82 studied independently, it is still unclear whether such a syndrome is due to the progressive
83 accumulation of several independent PIPAs during evolutionary time (Thomas et al. 2010), or results
84 from a cascade of effects following disruption of a major physiological mechanism (Cézilly and
85 Perrot-Minnot, 2010). As suggested by Cézilly and Perrot-Minnot (2010), evidence for a positive
86 correlation between the intensities of two distinct PIPA could be suggestive of some common
87 underlying mechanism (even if an absence of correlation would not be sufficient to conclude to the
88 absence of such a link).

89 Another way to provide evidence for associations between various PIPAs could be to
90 examine the temporal coincidence in the manifestations of various dimensions of an infection
91 syndrome. In addition to rare cases of direct reproductive organ consumption by parasites, castration
92 can be achieved through two main different ways, with different outcomes in their link with other
93 PIPA. Castration can result from a large neuroendocrine disorder including the perturbation of the
94 endocrine glands regulating gonadic development. In that case, links between parasites and
95 neuroendocrine modulators may play a crucial role in linking behavioural alterations and castration
96 (Hurd, 2009), and castration may coincide with behavioural alterations (Cézilly and Perrot-Minnot,
97 2010). Alternatively, an 'energy-draining' castration can be achieved through the uptake of essential
98 host resources by the parasite, independently of other effects of the parasite (Hall *et al.* 2007).
99 Therefore, under the energy-drain hypothesis, castration should occur early in the course of infection
100 (Hall *et al.* 2007) and should not coincide with behavioural modifications. However, as discussed by
101 Hall et al (2007), the feeding rate of parasites may not be constant, as it may vary as parasites grow
102 larger, or increase hyperbolically with food density, depending on the type of feeding functional
103 response. In addition, recent findings on parasite behavioural manipulation showed that this
104 phenomenon is dynamic. While parasite stages infective to the next host generally induce alterations
105 that seem to increase trophic transmission, the reverse might be observed in intermediate hosts
106 infected by parasites at an early developmental stage, still unable to infect the definitive host

(Dianne *et al.* 2011, Weinreich *et al.* 2013). This phenomenon might have been selected because trophic transmission occurring before the parasite has reached the infective stage would result in zero fitness (Parker *et al.* 2009). Ideally, then, an optimal temporal sequence of manipulations could consist of a 'protection-then-increased exposure' strategy.

The purpose of the present study was to test the time course and correlations between behavioural alteration and castration, two PIPAs occurring during the infection of the crustacean amphipod *Gammarus pulex* by the acanthocephalan parasite *Polymorphus minutus*. Adult *P. minutus* are intestinal parasites of various waterbird species, and rely on freshwater amphipods as intermediate hosts (Kennedy, 2006). Cystacanths (the developmental stage at which the parasite is able to infect its avian hosts) of *P. minutus* are known to alter the behaviour of *G. pulex* through reversing geotaxis (infected animals are attracted to the top of the water column, while uninfected ones tend to stay at the bottom of the water column), and increasing clinging behaviour (Cézilly *et al.* 2000; Bauer *et al.* 2005). Infected animals thus spend more time at the water surface where they cling on any material available, while uninfected ones spend most of their time at the bottom of the water column. Infected hosts exhibiting such altered behaviour have been shown to be more exposed to predation by mallards, one of the *P. minutus* definitive host, than uninfected ones (Hinsbo, 1972; Jacquin *et al.* 2014). Parallel to this, Bollache *et al.* (2001, 2002) evidenced a strong parasitic castration in *G. pulex* associated with the presence of *P. minutus* cystacanths. Infected males and females are almost never found in pre-copulatory mate guarding, and infected females lay no egg, following disruption of oocyte maturation. Little is known, however, on the time course of behavioural and reproductive PIPAs. Bethel and Holmes (1973) reported that *G. lacustris* infected by acanthellae (early developmental stage) of *P. paradoxus* did not express the behavioural alterations shown when they are infected by cystacanths of the same species. On the other hand, Rauque and Semenas (2009) found that infection with acanthellae of two acanthocephalan parasite species has no effect on the fecundity of their amphipod intermediate host, while Sparkes *et al.* (2006) found that behavioural reproductive suppression in freshwater isopod males infected with

Acanthocephalus dirus did not occur until the parasite had reached the cystacanth stage. Taken together, these observations suggest that reproductive and behavioural PIPAs might be synchronous, but, to the best of our knowledge, this has not yet been assessed within a single study.

MATERIAL AND METHODS

Gammarid sampling and maintenance

Gammarids were sampled between February and April 2016 in a tributary of the Suzon River, near Val-Suzon (Côte d’Or). Individuals were randomly collected using a hand net at all depths of the river (between ca 30 cm, the maximum depth, and the surface). Gammarids were sampled every three weeks and individuals from each sample spent the same amount of time in the laboratory. Within three days after sampling, gammarids were separated according to their infection status (uninfected, infected by an acanthellae, or infected by a cystacanth) and sex. These traits were determined through observations with a stereoscopic microscope, *P. minutus* infection and stage being easily detected through the translucent cuticle of the intermediate host. Sex of gammarids is easily distinguishable by examination of the second pair of gnathopods and, often, the presence of eggs in the female’s external broodpouch (Sutcliffe 1992). Gammarids were maintained in separate tanks (30*20*10cm) in a mix of river water and tap water (dechlorinated and oxygenated, renewed each week). Rocks and plants were added to the tank to approximate river conditions, and conditioned elm leaves were used regularly to feed gammarids. Room temperature was 15 ± 1°C, and photoperiod was 13:11h light:dark. Behavioural tests and reproductive traits were measured between day 4 and day 13 post-sampling. Each day, groups of 6 to 8 infected individuals and 6 to 8 uninfected ones were tested simultaneously.

Behavioural tests

157 *Geotaxis*

158 Based on the procedure described in Bauer *et al.* (2005), we tested the vertical distribution of
159 infected and uninfected gammarids. To that end, animals were placed individually in transparent
160 vertical glass columns (35cm height, 6cm diameter) filled with stock water and covered by a net to
161 allow gammarids to grasp at the water surface. This setup was placed in a large box, black inside,
162 with one side open for observation. Columns were vertically divided in six equal parts, numbered
163 from 0 (lower part) to 5 (upper part). Following an acclimatization period of 10 minutes, a score was
164 attributed to the individual depending on its vertical position in the column, every 30 seconds during
165 the next 10 minutes. The overall score for each individual was then calculated as the sum of each
166 score, such that a gammarid staying at the bottom of the column for 10 minutes scored 0 (strong
167 positive geotaxis), whereas one staying at the top for the same amount of time scored 100 (strong
168 negative geotaxis). The water column was emptied and filled again after each test. Data were
169 analyzed using non-parametric Wilcoxon or Kruskal-Wallis tests, followed by a Steel-Dwass non-
170 parametric post-hoc test.

171 *Inclination of males to pair*

172 During precopulatory mate guarding (or amplexus), male gammarids carry the females
173 underneath their own ventral surface, for several days before egg laying. We used a protocol similar
174 to that of Bollache *et al.* (2001) to assess the motivation of males to pair. After having been tested
175 for geotaxis, control and infected males were individually placed in small cups (3 cm height, 6 cm
176 diameter), filled with stock water, during 30 minutes of acclimatization. During this time, uninfected
177 females receptive to mating (i.e. paired female taken from the stock tank, carrying eggs at similar and
178 advanced development), were separated from their males and isolated. At the end of the
179 acclimatization time, a single receptive and uninfected female was added to each cup containing a
180 focal male. Cups were checked every 5 minutes over one hour, to see if individuals had entered
181 in amplexus. The experiment ended if amplexus did not occur after one hour, since, after such a time

period, males rarely engage in pairing (Bollache *et al.* 2001). Data were analyzed with a Generalized Linear Model (GLM), using a binomial distribution and a Logit link function, and using Firth Bias-adjusted Estimates, allowing the analysis of rare events.

At the end of the behavioural tests, each individual was kept separately in a small cup (3 cm height, 6 cm diameter) until dissection and measurement of reproductive parameters.

Reproductive traits

Dissections were made in Ringer, using a stereoscopic microscope (Nikon SMZ-10A) and a video measurement system (Nikon NIS – Elements D, v4.10). Gammarids were first anesthetized in carbonated water and their size determined by the distance from the fourth coxal plate basis to the dorsal limit of the individual (Bollache *et al.* 2002). They were then dissected, and the parasite developmental stage was confirmed in infected animals.

Reproductive traits in males

The second pair of gnathopods is used by males to grasp females during amplexus (Conlan, 1991). We therefore investigated whether parasitism affected this morphological trait associated with reproduction. The maximum length and maximum width of the right second gnathopod were measured to the nearest 0.01 mm, and their product was used as a proxy of gnathopod size. Since gnathopod size shows an allometric relationship with body size (see Cothran and Jeyasingh 2010), a linear regression was calculated between gnathopod size and the size of the fourth coxal plate, after log transformation. The residuals of the regression were used to compare infection groups using an analysis of variance. Sperm reserves were estimated by a method modified from Lemaître *et al.* (2009). One intact testis (including seminal vesicle) was isolated, and placed into 1000 µl of demineralized water. The gonad was fragmented in small pieces, and creation of sperm heaps was avoided by gently stirring the solution with forceps, to obtain a homogenous dilution of sperm in 1000 µl. Four sample of 10 µl were placed on a slide, and dried during 10 to 20 minutes at 37°C. Using a microscope (“Nikon Eclipse E400”, objective x100), the total number of spermatozooids for

each sample was counted, to obtain a total number of spermatozooids in 40 μ l, after having verified the repeatability between samples. Since spermatozoid number may depend on male body size, we used a GLM using a Poisson distribution and a Log link function, corrected for overdispersion, to estimate the link between spermatozoid count and the size of the fourth coxal plate after its log transformation. The residuals of this model were used to compare infection groups using an analysis of variance.

Reproductive traits in females

The total number of eggs or embryos present in the ventral brood pouch was counted to estimate female fecundity. To determine the effect of infection on oocyte maturation, gonads were dissected, and the number of developed oocytes and white (non-developing) oocytes were counted in one ovary, following Bollache *et al.* (2002). The number of 'blue oocytes' (developing but altered), which only occurred in *P. minutus*-infected individuals, was also counted (see Bollache *et al.* 2002). The proportion of females carrying eggs or embryos were analyzed with a GLM, using a binomial distribution and a Logit link function, and using Firth Bias-adjusted Estimates. Oocyte counts were analyzed using non-parametric Kruskal-Wallis tests, followed by a Steel-Dwass post-hoc test.

Correlations between geotaxis scores and reproductive traits were analyzed using non-parametric Spearman's rho, followed by a Bonferroni correction. All statistical tests were made using JMP v. 10.0.0 (©SAS Institute 2012).

RESULTS

During the three field sampling periods, the prevalence of infection with acanthellae progressively decreased from 63.0% in February to 0% in April (Figure 1). From this observation, as well as considering results from other longitudinal field studies of acanthocephalans (e.g. Sparkes *et al.* 2006, Kennedy 2006), we inferred that parasites were maturing progressively from acanthella to

cystacanth stages in the population. For measurements of geotaxis, we therefore compared similar parasite developmental stages among sampling dates.

Geotaxis measurements

At any date and within each infection status, geotaxis scores did not differ significantly between *G. pulex* males and females (Wilcoxon tests, all $P > 0.09$). Data were therefore pooled for further analyses. There was no significant variation in geotaxis score between sampling dates for both uninfected individuals (three sampling dates, $n = 60, 49$, and 13 , respectively, Kruskal-Wallis: $\chi^2 = 2.29$; 2 d.f., $P = 0.32$) and individuals infected by acanthellae (two sampling dates, $n = 34$ and 16 , respectively, Wilcoxon: $\chi^2 = 2.07$; 1 d.f., $P = 0.15$). Samples from different dates were therefore pooled for further analyses. However, geotaxis score varied according to sampling date for individuals infected with a cystacanth (three sampling dates, $n = 20, 25$ and 23 , respectively, Kruskal-Wallis: $\chi^2 = 8.31$; 2 d.f., $P = 0.016$). Infection with a cystacanth was associated with a lower score in February than in March and April (Steel-Dwass post-hoc test, $P = 0.04$ and $P = 0.03$, respectively), while the scores did not differ significantly between March and April ($P = 0.98$). Owing to the variation in cystacanth prevalence through time reported above, we distinguished two cystacanth categories, depending on whether they had been sampled in February or in March-April, for further analyses.

The geotaxis score of *G. pulex* varied significantly according to their infection status (Figure 2, Kruskal-Wallis: $\chi^2 = 30.19$; 3 d.f., $P < 0.0001$). Infections with cystacanths sampled in March-April were associated with geotaxis scores higher than those of uninfected gammarids, while infections by acanthellae were associated with geotaxis scores lower than those of uninfected individuals (Figure 2). Scores of individuals infected by cystacanths sampled in February were significantly lower than those recorded in March-April, but not significantly different from those of individuals infected by acanthellae and those of uninfected ones (Figure 2).

256

257 *Male reproductive traits*

258 About 40% of uninfected males initiated pairing with receptive females. This proportion significantly
259 differed from the proportions of infected males initiating a pair (maximum 10%), irrespective of the
260 parasite developmental stage (GLM: $\chi^2 = 25.73$, 3 d.f., $P < 0.0001$) (Figure 3a).

261 Sperm count, corrected for gammarid body size, was different among groups (ANOVA $F_{3,105} = 3.00$, P
262 $= 0.03$). Males infected with cyctacanth from the March-April sample had more sperm reserves than
263 uninfected males, but not males of other infection categories (Figure 3b).

264 The size of the second gnathopod, corrected for gammarid body size, did not differ according to the
265 infection status (ANOVA $F_{3,131} = 1.12$, $P = 0.34$; Figure 3c).

266

267 *Female reproductive traits*

268 The proportion of females carrying eggs or embryos in their ventral incubating pouch (i.e.
269 reproductive females) was 89.29 % in uninfected individuals. This proportion significantly differed
270 from that of reproductive infected females (maximum 13.6 %), irrespective of the parasite
271 developmental stage (GLM: $\chi^2 = 82.08$, 3 d.f., $P < 0.0001$) (Figure 4a).

272 Oocyte count was not analyzed in females infected with cystacanth sampled in February, because of
273 the small sample size of successful dissections ($n = 4$). In other categories, the number of oocytes per
274 ovary did not differ significantly among uninfected and infected females (Kruskal-Wallis: $\chi^2 = 3.74$, 2
275 d.f., $P = 0.15$) (Figure 4b). However, the proportion of developing oocytes was much higher in
276 uninfected females than in infected ones, irrespective of the developmental stage of parasites, in
277 which this proportion was virtually zero, with one single exception (Kruskal-Wallis: $\chi^2 = 54.44$, 2 d.f.,
278 $P < 0.0001$; Figure 4c). Indeed, the proportion of blue oocytes (i.e. showing abnormal development),
279 as well as the proportion of white oocytes (i.e. not developing oocytes) were higher in infected

females than in uninfected ones, irrespective of the developing stage of the parasite (Kruskal-Wallis: $\chi^2 = 37.15$, 2 d.f., $P < 0.0001$, Figure 4d, and $\chi^2 = 34.75$, 2 d.f., $P < 0.0001$, Figure 4e, respectively).

Correlations between geotaxis and reproductive traits

We considered categories of infections in two different ways. First, we distinguished acanthellae and the two cystacanths categories, as described above. Second, we grouped all the developmental stages into a single, “infected” category. Such a grouping was made to take into account a potential correlation between the time course of behavioural changes (as described before) and parasite castration. Only one of these correlations was found to be significant (Table 1): there was a positive correlation between geotaxis score and sperm number in infected males, after pooling data from all developmental stages (Figure 5).

DISCUSSION

We found that both the direction and extent of modification of geotaxis in *G. pulex* were highly dependent on the developmental stage of *P. minutus*, whereas castration was not.

Whereas the cystacanth stage induced a reversion of geotaxis (negative geotaxis, compared to positive geotaxis in uninfected animals), confirming previous observations (e.g. Cézilly *et al.* 2000; Bauer *et al.* 2005; Jacquin *et al.* 2014), the acanthella stage, not infective yet for the definitive host (Bethel and Holmes 1974), induced a stronger, positive geotaxis than that observed in uninfected gammarids. A previous study on the congeneric acanthocephalan *P. paradoxus* concluded that acanthellae induce no change in behaviour in gammarids (Bethel and Holmes, 1974). However, that study examined evasive behaviour and not geotaxis, such that further investigations are necessary to have a better appreciation of the time course of multidimensionality in manipulation in relation to parasite development. The present observations are however consistent with others showing that

immature stages of parasites may induce behavioural alterations opposite to those induced by the parasite stages infective for the next host (Hammerschmidt *et al.* 2009; Dianne *et al.* 2011, Gopko *et al.* 2015), or, at a finer scale, that parasites may alter host behaviour in a time-specific manner throughout the day, resulting in a protection from predation by non-suitable definitive hosts (Levri 1998). Such a phenomenon has been predicted by Parker *et al.* (2009) as an optimal ‘protection-then-increased exposure to predation’ strategy for trophically-transmitted parasites, and experiments have indeed revealed that such opposite behavioural alterations are associated with lower predation rates (Dianne *et al.* 2011; Weinreich *et al.* 2013). Whether infection with acanthellas of *P. minutus* is associated with lower vulnerability to predation by definitive hosts remains to be tested in the present case.

Our results further suggest that the transition between positive and negative geotaxis is progressive. Indeed, cystacanths sampled in February were associated with geotaxis scores intermediate between those of acanthellae and those of cystacanths that were sampled one month later. February cystacanths being on average younger than March-April cystacanths, we evidenced here a progressive shift in a parasite-induced behavioural change – ‘progressive’ meaning here that the change occurred over several days. This observation is consistent with the results obtained by Bethel and Holmes (1974) on *P. paradoxus*, for which 17 days, on average, separate the invagination of the acanthellae (defining the cystacanth stage) from the onset of altered behaviour in the intermediate host. This confirms that the intensity of behavioural changes increases with the course of acanthocephalan maturation after cystacanth stage (Franceschi *et al.* 2008).

Contrasting with what was observed for behavioural changes, we observed no variation in the intensity of castration as a function of parasite maturation. In both males and females the extent of castration at the acanthellae stage, did not differ from that observed at the cystacanth stage of infection. Even if acanthellae in this study were rather old (they were colored and visible through the host's cuticle), this result contrasts with those of Sparkes *et al.* (2006) and Rauque and Semenas

(2009) who observed, in three other acanthocephalan species inducing total or partial castration, that castration did not occur before parasites had reached the cystacanth stage. Therefore, castration induced by *P. minutus* seems to be triggered earlier during parasite development than in other castrating acanthocephalans.

As reported in Bollache *et al.* (2001), infected males paired at lower rates compared to uninfected ones. We further explored this phenomenon by considering whether this lower pairing success was due to gonadal dysfunction or morphological alteration. We found no evidence for an alteration of sperm reserves associated with infection by *P. minutus*, nor did we observe any alteration of gonadal structure. This is consistent with another study on acanthocephalans (Bierbower and Sparkes, 2007). Actually, *G. pulex* males infected with older cystacanths had higher sperm reserves. Since sperm tends to accumulate if males do not inseminate females (Lemaître *et al.* 2009), sperm accumulation after a longer period of sexual abstinence, due to a longer infection time, could explain this result. This is corroborated by the positive correlation between geotaxis score and sperm reserve when taking into account all stages of the *P. minutus* development (Figure 5): the older the parasites, the more they altered their host's geotaxis and the more the hosts accumulated sperm, probably as a consequence of behavioural castration. Due to our experimental procedure, we were unable to assess whether the sperm accumulated in infected males was functional or not. Such an assessment would not be straightforward, however, because sperm is not motile in gammarids, such that a comparison of insemination success between infected and uninfected males would be necessary to address the question. However, during our dissections, we did not observe any abnormal sperm structure. In addition, infection status had no influence on the size of the second gnathopod, the appendage used to grasp females during mate guarding. Therefore, alterations of reproductive function or morphology are unlikely to explain reduced male mating success in *G. pulex* infected with *P. minutus*. One possibility, then, is that infection with *P. minutus* has an effect on the nervous system of male *G. pulex* that disrupts their normal reaction towards sexually receptive

females. Such a possibility has been observed in an insect infected by another acanthocephalan species, where males cannot perceive normally sex pheromones (Carmichael *et al.* 1993).

No significant correlation was found between the intensity of behavioural changes and the intensity of castration, either when considering acanthellae- and cystacanth-induced behavioural changes as separated phenomena (i.e. correlations analyzed separately for the different developmental stages), or as two components of the same phenomenon (analyses made after grouping all parasite stages). This lack of correlation, as well as the differences in temporal dynamics between castration and geotaxis changes, does not argue in favour of a mechanistic link between these two PIPAs (Cézilly and Perrot-Minnot, 2010; Thomas *et al.* 2010). However, this result is, of course, not sufficient to conclude to the absence of such a link. Over the course of infection, products excreted by the parasite may accumulate in its host. In the same time, several by-products of the immune response of the host may accumulate too. However, the dose-response of castration and that of reversed geotaxis in relation to the concentration of a single or several key substances may not be the same. For instance, a behavioural or physiological dose-response may not be linear but sigmoid (Lau *et al.* 2000; Byers, 2013). It has recently been proposed that reversed geotaxis could be a behavioural response to hypoxia signaling, resulting either from the sustained hypoxic-like conditions induced by *P. minutus* cystacanths, or the release, from the excretion/secretion products of parasites, of lactate and succinate in the host brain (Perrot-Minnot *et al.* 2016). The problem remains that the two opposite geotaxis disruptions induced by the two parasite developmental stages may hardly been explained by such a phenomenon. Interestingly, however, some dose-responses can show bi-phasic shape, corresponding to a hormetic-like effect. In such cases, the exposition to a low dose of a substance may induce in an organism the opposite response to a very high dose (see for instance Rabhi *et al.* 2014). This could explain, from a mechanistic point of view, the contrasted effect of the developmental stages of *P. minutus* on the geotaxis of *G. pulex*. Independently, some studies have showed that natural or experimentally-induced hypoxia down-regulates the expression of vitellogenin, the major egg-yolk protein, in a number of crustacean and

380 fish (Brown-Peterson *et al.* 2011; Li and Brouwer, 2013; Lai *et al.* 2016 and references therein).
381 Therefore, under the hypothesis that hypoxia has an early and strong effect on reproduction but that
382 phototaxis behaviour shows a bi-phasic response to hypoxia, these two traits might actually be
383 synchronous. Now that the timing of expression of castration and geotaxis changes has been clarified
384 in *G. pulex* infected by *P. minutus*, unraveling the shape of hypoxia signaling dose-response on both
385 geotaxis and castration would be a promising avenue for testing the link between these different
386 PIPAs.

387

388 FINANCIAL SUPPORT

389 This study was supported by the Agence Nationale de la Recherche (ANR)(grant number ANR-13-
390 BSV7-0004-01).

391

References

- Baudoin M.** (1975). Host castration as a parasitic strategy. *Evolution* **29**, 335-352.
- Bauer, A., Haine, E. R., Perrot-Minnot, M.-J. and Rigaud, T.** (2005). The acanthocephalan parasite *Polymorphus minutus* alters the geotactic and clinging behaviours of two sympatric amphipod hosts: the native *Gammarus pulex* and the invasive *Gammarus roeseli*. *Journal of Zoology* **267**, 39-43.
- Bethel, W.M. and Holmes, J.C.** (1973). Altered evasive behavior and response to light in amphipods harboring acanthocephalan cystacanths. *Journal of Parasitology* **59**, 945-956.
- Bethel, W.M. and Holmes, J.C.** (1974). Correlation of development of altered evasive behavior in *Gammarus lactustris* (Amphipoda) harboring cystacanths of *Polymorphus paradoxus* (Acanthocephala) with the infectivity to the definitive host. *Journal of Parasitology* **60**, 272-274.
- Bierbower, S.M. and Sparkes, T. C.** (2007). Parasite-related pairing success in an intermediate host, *Caecidotea intermedius* (Isopoda): Effects of male behavior and reproductive physiology. *Journal of Parasitology* **93**, 445-449.
- Bollache, L., Gambade, G. and Cézilly, F.** (2001). The effects of two acanthocephalan parasites, *Pomphorhynchus laevis* and *Polymorphus minutus*, on pairing success in male *Gammarus pulex* (Crustacea: Amphipoda). *Behavioral Ecology and Sociobiology* **49**, 296-303.
- Bollache, L., Rigaud, T. and Cézilly, F.** (2002). Effects of two acanthocephalan parasites on the fecundity and pairing status of female *Gammarus pulex* (Crustacea: Amphipoda). *Journal of Invertebrate Pathology* **79**, 102-110.
- Brown-Peterson, N.J., Manning, C.S., Denslow, N.D. and Brouwer, M.** (2011). Impacts of cyclic hypoxia on reproductive and gene expression patterns in the grass shrimp: field versus laboratory comparison. *Aquatic Sciences* **73**, 127-141.
- Byers, J.** (2013). Modeling and regression analysis of semiochemical dose-response curves of insect antennal reception and behavior. *Journal of Chemical Ecology* **39**, 1081-1089.
- Carmichael, L.M., Moore, J. and Louis B. Bjostad L.B.** (1993). Parasitism and Decreased Response to Sex Pheromones in Male *Periplaneta Americana* (Dictyoptera: Blattidae). *Journal of Insect Behavior* **6**, 25-32.
- Cézilly, F. and Perrot-Minnot, M.-J.** (2010). Interpreting multidimensionality in parasite-induced phenotypic alterations: panselectionism versus parsimony. *Oikos* **119**, 1224-1229.
- Cézilly, F., Thomas, F., Médoc, V. and Perrot-Minnot, M.-J.** (2010). Host-manipulation by parasites with complex life cycles: adaptive or not? *Trends in Parasitology* **26**, 311-317.
- Cézilly, F., Grégoire, A. and Bertin, A.** (2000). Conflict between co-occurring manipulative parasites? An experimental study of the joint influence of two acanthocephalan parasites on the behaviour of *Gammarus pulex*. *Parasitology* **120**, 625-630.
- Cézilly, F., Favrat, A. and Perrot-Minnot, M.-J.** (2013). Multidimensionality in parasite-induced phenotypic alterations: ultimate versus proximate aspects. *Journal of Experimental Biology* **216**, 27-35.
- Conlan, K. E.** (1991). Precopulatory mating-behaviour and sexual dimorphism in the amphipod Crustacea. *Hydrobiologia* **223**, 255-282.

- 431 **Cothran, R.D. and Jeyasingh, P.D.** (2010). Condition dependence of a sexually selected trait in a
432 crustacean species complex: importance of the ecological context. *Evolution* **64**, 2535–2546.
- 433 **Dianne, L., Perrot-Minnot, M.-J., Bauer, A., Elsa, L. and Rigaud, T.** (2011). Protection first then
434 facilitation: a manipulative parasite modulates the vulnerability to predation of its intermediate host
435 according to its own developmental stage. *Evolution* **65**, 2692–2698.
- 436 **Franceschi, N., Bauer, A., Bollache, L. and Rigaud, T.** (2008). The effects of parasite age and intensity
437 on variability in acanthocephalan-induced behavioural manipulation. *International Journal for*
438 *Parasitology* **38**, 1161–1170.
- 439 **Gopko, M., Mikheev, V.N. and Taskinen, J.** (2015). Changes in host behaviour caused by immature
440 larvae of the eye fluke: evidence supporting the predation suppression hypothesis. *Behavioral*
441 *Ecology and Sociobiology* **69**, 1723–1730.
- 442 **Hall, S.R., Becker, C. and Cáceres, C.E.** (2007). Parasitic castration: a perspective from a model of
443 dynamic energy budgets. *Integrative and Comparative Biology* **47**, 295–309.
- 444 **Hammerschmidt, K., Koch, K., Milinski, M., Chubb, J. C. and Parker, G. A.** (2009). When to go:
445 optimization of host switching in parasites with complex life cycles. *Evolution* **63**, 1976–1986.
- 446 **Helluy, S. and Thomas, F.** (2003). Effects of *Microphallus papillorobustus* (Platyhelminthes :
447 Trematoda) on serotonergic immunoreactivity and neuronal architecture in the brain of *Gammarus*
448 *insensibilis* (Crustacea : Amphipoda). *Proceedings of the Royal Society Series B* **270**, 563–568.
- 449 **Hindsbo, O.** (1972). Effects of *Polymorphus* (Acanthocephala) on colour and behaviour of *Gammarus*
450 *lacustris*. *Nature* **238**, 333.
- 451 **Hurd, H.** (2001). Host fecundity reduction: a strategy for damage limitation? *Trends in Parasitology*
452 **17**, 363–368.
- 453 **Hurd, H.** (2009). Evolutionary drivers of parasite-induced changes in insect life-history traits: from
454 theory to underlying mechanisms. *Advances in Parasitology* **68**, 85 – 110.
- 455 **Jacquin, L., Mori, Q., Pause, M., Steffen, M., and Médoc, V.** (2014). Non-specific manipulation of
456 gammarid behaviour by *P. minutus* parasite enhances their predation by definitive bird hosts. *Plos*
457 *One* **9**, e101684.
- 458 **Jensen, K.H., Little, T., Skorpung, A. and Ebert, D.** (2006). Empirical support for optimal virulence in a
459 castrating parasite. *Plos Biology* **4**, e197.
- 460 **Kennedy, C.R.** (2006). *Ecology of the Acanthocephala*. Cambridge University Press, Cambridge, U.K.
- 461 **Lafferty, K.D. and Kuris, A.M.** (2009). Parasitic castration: the evolution and ecology of body
462 snatchers. *Trends in Parasitology* **25**, 564–572.
- 463 **Lai, K.P., Li, J.W., Chan, C.Y.S, Chan, T.F., Yuen, K.W.Y. and Chiu J.M.Y.** (2016). Transcriptomic
464 alterations in *Daphnia magna* embryos from mothers exposed to hypoxia. *Aquatic Toxicology* **177**,
465 454–463.
- 466 **Lau, C.E., Sun, L., Wang, K., Simpao, A. and Falk, J.L.** (2000). Oral cocaine pharmacokinetics and
467 pharmacodynamics in a cumulative-dose regimen: pharmacokinetic-pharmacodynamic modeling of
468 concurrent operant and spontaneous behavior within an operant context. *Journal of Pharmacology*
469 *and Experimental Therapeutics* **295**, 634–643.

- 470 **Lemaître, J. -F., Rigaud, T., Cornet, S. and Bollache, L.** (2009). Sperm depletion, male mating
471 behaviour and reproductive “time-out” in *Gammarus pulex* (Crustacea, Amphipoda). *Animal*
472 *Behaviour* **77**, 49–54.
- 473 **Levri, E.D.** (1998). The influence of non-host predators on parasite-induced behavioral changes in a
474 freshwater snail. *Oikos* **81**, 531–537.
- 475 **Li, T. and Brouwer, M.** (2013). Field study of cyclic hypoxic effects on gene expression in grass shrimp
476 hepatopancreas. *Comparative Biochemistry and Physiology, Part D* **8**, 309–316.
- 477 **Moore, J.** (2002). *Parasites and the Behavior of Animals*. Oxford University Press, Oxford.
- 478 **Parker, G.A., Ball, M.A., Chubb, J.C., Hammerschmidt, K. and Milinski, M.** (2009). When should a
479 trophically transmitted parasite manipulate its host? *Evolution* **63**, 448–458.
- 480 **Perrot-Minnot, M.-J., Madaleno, M., Balourdet, A. and Cézilly, F.** (2012). Host manipulation
481 revisited: no evidence for a causal link between altered photophobia and increased trophic
482 transmission of amphipods infected with acanthocephalans. *Functional Ecology* **26**, 1007–1014.
- 483 **Perrot-Minnot, M.-J., Sanchez-Thirion, K. and Cézilly F.** (2014). Multidimensionality in host
484 manipulation mimicked by serotonin injection. *Proceedings of the Royal Society Series B* **281**,
485 20141915.
- 486 **Perrot-Minnot, M.-J., Maddaleno, M. and Cézilly F.** (2016). Parasite-induced inversion of geotaxis in
487 a freshwater amphipod: a role for anaerobic metabolism? *Functional Ecology* **30**, 780–788.
- 488 **Poulin, R.** (2010). Parasite manipulation of host behavior: an update and frequently asked questions.
489 In *Advances in the Study of Behavior*, Vol. 41, (ed. Brockmann, H. J. et al.), pp. 151–186. Academic
490 Press, Burlington.
- 491 **Rabhi, K.K., Esancy, K., Voisin, A., Crespin, L., Le Corre, J., Tricoire-Leignel, H., Anton, S. and**
492 **Gadenne, C.** (2014). Unexpected effects of low doses of a neonicotinoid insecticide on behavioral
493 responses to sex pheromone in a pest insect. *PLoS One* **9**, e114411.
- 494 **Rauque, C. A. and Semenas, L.** (2009). Effects of two acanthocephalan species on the reproduction
495 of *Hyalella patagonica* (Amphipoda, Hyalellidae) in an Andean Patagonian Lake (Argentina). *Journal*
496 *of Invertebrate Pathology* **100**, 35–39.
- 497 **Sparkes, T.C., Weil, K.A., Renwick, D.T. and Talkington, J.A.** (2006). Development-related effects of
498 an acanthocephalan parasite on pairing success of its intermediate host. *Animal Behaviour* **71**, 439–
499 448.
- 500 **Sutcliffe, D.W.** (1992). Reproduction in *Gammarus* (Crustacea, Amphipoda): basic processes.
501 *Freshwater Forum* **2**, 102–128.
- 502 **Tain, L., Perrot-Minnot, M.-J. and Cézilly, F.** (2006). Altered host behaviour and brain serotonergic
503 activity caused by acanthocephalans: evidence for specificity. *Proceedings of the Royal Society Series*
504 *B* **273**, 3039–3045.
- 505 **Thomas, F., Adamo, S. and Moore, J.** (2005). Parasitic manipulation: where are we and where should
506 we go? *Behavioural Processes* **68**, 185–199.
- 507 **Thomas, F., Poulin, R. and Brodeur, J.** (2010). Host manipulation by parasites: a multidimensional
508 phenomenon. *Oikos* **119**, 1217–1223.

509 **Weinreich, F., Benesh, D. P. and Milinski, M.** (2013). Suppression of predation on the intermediate
510 host by two trophically-transmitted parasites when uninfected. *Parasitology* **140**, 129–135.

Figure captions

Figure 1: Proportion of acanthella developmental stage of *Polymorphus minutus* in their infected *G. pulex* hosts, according to sampling date. Other developmental stages were cystacanths.

Figure 2: Scores of geotaxis in *Gammarus pulex* according to their infection status, with categories as defined in the text. U: uninfected animals, A: infected with acanthellae, Cfeb: infected with cystacanths sampled in February, CMar-Apr: infected with cystacanths sampled in March and April (see text). Sample size are provided. Box-plots are median, interquartile and interdecile (and outliers). Boxed labelled with different letters denotes scores significantly different after a Steel-Dwass non parametric post-hoc test ($\alpha = 0.05$).

Figure 3: Reproductive traits in *Gammarus pulex* males, according to their infection status (U: uninfected animals, A: infected with acanthellae, Cfeb: infected with cystacanths sampled in February, CMar-Apr: infected with cystacanths sampled in March and April). (a) Proportion of males initiating pairing in one hour. Bars surrounded by different letters denotes proportions significantly different after Odd-ratios tests; (b) Sperm reserve in testis. Dots are means, bars are standard errors on the mean; Bar surrounded by a star denotes a mean significantly different for the uninfected control group after Dunnett post-hoc test ($P = 0.03$) (c) Size of second pair of gnathopods as a function of gammarid size and infection status. Sample size are provided within brackets below the x axes.

Figure 4: Reproductive traits in *Gammarus pulex* females, according to their infection status (U: uninfected animals, A: infected with acanthellae, Cfeb: infected with cystacanths sampled in February, CMar-Apr: infected with cystacanths sampled in March and April). Sample size are provided below the x axes. (a) Proportion of females with eggs in the brood pouch. Bars surrounded by different letters denotes proportions significantly different after Odd-ratios tests; (b) Number of oocytes counted in one ovary; (c) Proportion of developing oocytes (d) Proportion of blue oocytes, (e) Proportion of white oocytes. From (b) to (e), box-plots are median, interquartile and interdecile (and outliers). Bars surrounded by different letters denotes significant differences after a Steel-Dwass non parametric post-hoc test ($\alpha = 0.05$).

Figure 5: Correlation between geotaxis scores and sperm number in *Gammarus pulex* males infected with different stages of *Polymorphus minutus*. White dots: infected with acanthellae; grey dots infected with cystacanths sampled in February; black dots; infected with cystacanths sample in March-April.

Table 1: Correlations (Spearman’s rho) between geotaxis scores and major reproductive traits of *Gammarus pulex* males and females, according to their infection status.

Sex	Trait correlated with geotaxis	Infection category	N	Rho	P
Males	Gnathopod size*	U	64	-0.14	0.28
		A	28	0.001	0.97
		CFeb	13	0.47	0.11
		CMar-Apr	27	-0.20	0.31
		I	68	-0.001	0.99
	Sperm number**	U	53	-0.07	0.63
		A	22	0.40	0.07
		CFeb	10	0.30	0.39
		CMar-Apr	21	-0.07	0.75
		I	53	0.35	0.001
Females	% white oocytes	U	49	-0.05	0.70
		A	12	-0.16	0.61
		CFeb	n.a.		
		CMar-Apr	16	-0.15	0.55
		I	33	-0.18	0.29
	% blue oocytes	U	n.v.		
		A	12	-0.02	0.94
		CFeb	n.a.		
		CMar-Apr	16	0.007	0.98
		I	33	0.12	0.50

U: uninfected animals; A: infected with acanthellae; Cfeb: infected with cyctacanth's sampled in February; CMar-Apr: infected with cyctacanth's sampled in March and April; C: infected with cyctacanth's (all date grouped); I: infected animals (all developemental stages grouped).
*corrected for body size (residuals of the regression)
**corrected for body size (residuals of the Quasi-Poisson GLM)
n.a. sample size too low to be analyzed
n.v. no variation on this trait
Correlation in bold was significant even after Bonferroni correction.

Proportion acanthella among infected