



HAL
open science

Genetic Characterization of a French Cohort of GNE-mutation negative inclusion body myopathy patients with exome sequencing

Mathieu Cerino, Svetlana Gorokhova, Pascal Laforet, Rabah Ben Yaou, Emmanuelle Salort-Campana, Jean Pouget, Shahram Attarian, Bruno Eymard, Jean-François Deleuze, Anne Boland, et al.

► **To cite this version:**

Mathieu Cerino, Svetlana Gorokhova, Pascal Laforet, Rabah Ben Yaou, Emmanuelle Salort-Campana, et al.. Genetic Characterization of a French Cohort of GNE-mutation negative inclusion body myopathy patients with exome sequencing. *Muscle & Nerve*, 2017, 56, pp.993-997. 10.1002/mus.25638 . hal-01741741

HAL Id: hal-01741741

<https://hal.science/hal-01741741>

Submitted on 28 Mar 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Genetic characterization of a French cohort of GNE-mutation negative inclusion body myopathy patients using exome sequencing

Mathieu CERINO, PharmD,^{1,2} Svetlana GOROKHOVA, MD, PhD,¹ Pascal LAFORET, MD, PhD,³ Rabah BEN YAOU, PhD^{3,4} Emmanuelle SALORT-CAMPANA, MD, PhD,^{1,5} Jean POUGET, MD, PhD,^{1,5} Shahram ATTARIAN, MD, PhD,^{1,5} Bruno EYMARD, MD, PhD,³ Jean-François DELEUZE, PhD⁶ Anne BOLAND, PhD⁶ Anthony BEHIN, MD, PhD,³ Tanya STOJKOVIC, MD, PhD,³ Gisele BONNE, PhD,⁴ Nicolas LEVY, MD, PhD,^{1,2} Marc BARTOLI*, PhD^{1,2} Martin KRAHN*, MD, PhD,^{1,2}

¹ Aix Marseille Univ, Inserm, GMGF, Marseille, France

² APHM, Hôpital Timone Enfants, Département de Génétique Médicale, Marseille, France

³ APHP, G.H. Pitié Salpêtrière, Centre de Référence Maladies Neuromusculaires Paris-Est, Institut de Myologie, Paris, France.

⁴ Sorbonne Universités, UPMC Univ Paris 06, Inserm UMRS974, CNRS FRE3617, Center for Research in Myology, Institut de Myologie, G.H. Pitié Salpêtrière, Paris, France

⁵ APHM, Hôpital La Timone, Centre de référence des maladies neuromusculaires et de la SLA, Marseille, France

⁶ Centre National de Génotypage, Institut de Génomique, CEA, Evry, France.

* Authors with equal contribution.

Corresponding Authors:

Mathieu Cerino: mathieu.cerino@ap-hm.fr and Martin Krahn: martin.krahn@univ-amu.fr

INSERM AMU UMR_S910, Faculté de Médecine de Marseille, 4^e étage Aile Verte,
27 boulevard Jean Moulin, 13385 Marseille Cedex 05, FRANCE

TEL: 0033-(0)4 91 32 49 40

FAX: 0033-(0)4 91 80 43 19

Abstract word count: 150 ; Manuscript word count: 1335

We confirm that we have read the Journal's position on issues involved in ethical publication and affirm that this report is consistent with those guidelines.

The authors have no conflict of interest to report and no relevant financial relationships to disclose.

This study was supported by the France Génomique infrastructure (grant no. ANR-10-INBS-09) managed by the National Research Agency (ANR) part of the Investment for the Future program, Fondation Maladies Rares, FHU-MaRCHE, the APHM, Inserm, the "Bureau des PU-PH de l'Assistance Publique – Hôpitaux de Marseille (AP-HM)", and by a Grant FP7/2007–2013 from the European Community Seventh Framework Programme (Grant Agreement No. 2012–305121) "Integrated European omics research project for diagnosis and therapy in rare neuromuscular and neurodegenerative diseases" (NEUROMICS).

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as an 'Accepted Article', doi: 10.1002/mus.25638

Genetic characterization of a French cohort of GNE-mutation negative inclusion body myopathy patients using exome sequencing

(Abstract)

INTRODUCTION: Hereditary inclusion body myopathy (hIBM) refers to a group of clinically and genetically heterogeneous diseases. The overlapping histochemical features of hIBM with other genetic disorders lead to low diagnostic rates with targeted single-gene sequencing. This is true for the most prevalent form of hIBM, GNEpathy. Thus, we used whole exome sequencing (WES) to evaluate whether a cohort of clinically suspected GNEpathy patients undiagnosed by targeted *GNE* analysis could be genetically characterized.

METHODS: 20 patients with hIBM but undiagnosed by targeted *GNE* sequencing were analyzed using WES before data filtering on 306 genes associated with neuromuscular disorders.

RESULTS: 7 patients out of 20 were found to have disease-causing mutations in genes associated with hIBM, or genes allowing for hIBM in the differential diagnosis, or associated with unexpected diagnosis.

DISCUSSION: NGS is an efficient strategy in the context of hIBM, resulting in a molecular diagnosis for 35% of the patients initially undiagnosed by targeted *GNE* analysis.

Keywords: exome, hIBM, GNE, NGS, diagnosis, myopathy

INTRODUCTION

Hereditary inclusion body myopathies (hIBM) represent a heterogeneous group of muscular disorders defined by the relatively nonspecific criterion of rimmed vacuoles on muscle biopsy¹.

GNEpathy², caused by mutations in *GNE* (UDP-N-acetylglucosamine-2-epimerase/N-acetylmannosamine kinase, MIM*603824)³ is the most common form of hIBM, with many clinical features overlapping with other forms of hIBM, implicating other genes or forms with yet unknown underlying genetic defects. Targeted analysis of *GNE* in a large recently-described French cohort with suspected GNEpathy provides only a 20% diagnostic yield (32 of 164 patients)⁴.

In the present study, we evaluated the extent to which a cohort of clinically suspected GNEpathy patients undiagnosed by *GNE* targeted analysis may be genetically characterized, by implicating other genes previously known to cause neuromuscular disorders using whole exome sequencing (WES) associated with data filtering for 306 genes of interest.

METHODS

We selected 20 unrelated index cases (IC) with clinically suspected GNEpathy associated with rimmed vacuoles on muscle biopsy samples, but for which no *GNE* disease-causing mutation had been identified by direct targeted sequencing. Samples had been prepared and stored by the Center of Biological Resources, Department of Medical Genetics, La Timone Hospital, Marseille, and were used following the ethical recommendations of our institution and according to the Declaration of Helsinki. All included patients gave their written consent prior to the genetic study, in accordance with French law.

WES was performed using the SureSelect Human All Exon Kit version 5 (Agilent Technologies, Santa Clara, California) and the HiSeq 2000 (Illumina, San Diego, California).

Sequencing data were processed on the Illumina pipeline (CASAVA1.8.2) before using GATK⁵ variant calling and ANNOVAR⁶ annotation using the GRCh37/hg19 Human genome version, coverage statistics were computed using VarAFT (Variant Analysis and Filtration Tool ; <http://varaft.eu>, 2016), which uses BedTools⁷. VarAFT was also used to sort and filter the obtained variants.

Our initial analysis strategy focused on 306 genes previously reported to cause neuromuscular disorders, and selected from the Gene Table of Neuromuscular Disorders⁸ (including groups 1 to 5 and the main differential diagnosis genes) as previously described^{9,10}. A mean overall sequencing depth of 106X and a mean coverage of the coding exons of 95% (at 20X depth) and 91% (at 30X depth) was obtained for these 306 genes. Predicted pathogenicity of identified variants was determined using UMD-predictor¹¹, SIFT (Sort Intolerant From Tolerant human Protein)¹², PolyPhen-2 (Polymorphism Phenotyping v2)¹³ and HSF (Human Splicing Finder)¹⁴ softwares.

Regarding HSF¹⁴ in silico results, we defined four types of predicted splicing effects: 1) Probably damaging: associated with predicted strong splicing effect due to broken donor site (DS) or acceptor site (AS) and/or new DS/AS creation and/or strong possibility of broken Exonic Splicing Enhancer (ESE) site; 2) Possibly damaging: associated with predicted medium splicing effect relating to newly created DS/AS and/or medium possibility of broken ESE site; 3) Uncertain: associated with predicted mild splicing effect due to newly created DS/AS and/or low possibility of broken ESE site; and 4) Not affected: predicted weak or no splicing effect.

The overall pathogenicity score for each variant was determined according to the American College of Medical Genetics (ACMG) guidelines¹⁵. We established four groups of patients based on the degree of certainty of molecular diagnosis using the ACMG guidelines. The group with “definite diagnosis” consisted of the following patients: 1) Those carrying a homozygous variant classified as “pathogenic” using ACMG guidelines in a gene known to cause an autosomal recessive form of disease; 2) Compound heterozygotes carrying two variants classified as pathogenic; 3) Patients carrying one variant classified as pathogenic in a gene known to cause an autosomal dominant form of disease. The group with “probable diagnosis” was composed of patients carrying variants that were classified as “likely pathogenic” by ACMG guidelines. Patients carrying variants found to be pathogenic by certain prediction tools, but classified as “variants of uncertain significance” by ACMG guidelines were placed in the group with “possible diagnosis”. For those patients in the “no established diagnosis” group, no variant compatible with the patient’s phenotype was found.

All disease-causing variants identified by WES were confirmed using direct targeted sequencing (Genetic analyzer 3500XL; Thermo Fisher Scientific, Waltham, Massachusetts) and the following gene sequence references: *ACTA1* (NM_001100), *CAPN3* (NM_000070), *DES* (NM_001927), *FLNC* (NM_001458), *GYGI* (NM_004130), *MYH2* (NM_017534), *TARDBP* (NM_007375), *TTN* (NM_001267550) and *VCP* (NM_007126).

RESULTS

All phenotypic and mutational data are detailed in **Table 1**. A definite diagnosis was obtained for seven index cases (ICs). Patient P1 harbored a previously reported mutation in *TTN* (Titin, MIM*188840) associated with hereditary myopathy with early respiratory failure (HMERF)¹⁶. The homozygous status of this mutation is consistent with the parental

consanguinity. For Patients P2 and P5, the same heterozygous mutation in *VCP* (Valosin-Containing Protein, MIM *601023), previously described in the literature¹⁷, was discovered and associated with similar onset and clinical features (distal myopathy of upper and lower limbs). Compound heterozygous known mutations in *TTN*^{18,19} were found in patient P3, whereas patient P4 harbored a previously described heterozygous variant in *DES* (Desmin, MIM*125660)²⁰ leading to cardiomyopathy and myofibrillar abnormalities on the muscle biopsy, features that were retrieved in patient P4. For patient P6, a known *FLNC* (Filamin C, MIM *102565) mutation was found²¹. Surprisingly, we identified compound heterozygous mutations for the *GYGI* (Glycogenin 1, MIM*603942) gene in patient P7, associated with polyglucosan body myopathy type 2. In this patient, we found a previously described *GYGI* variant with a proven deleterious effect on splicing²², associated with a novel *GYGI* mutation leading to a frameshift of the reading frame and the introduction of a premature translation termination codon. Further investigations allowed additional clinical and histo-immunological features thus suggesting a polyglucosan body myopathy (data not shown). A probable diagnosis was obtained for patients P8 and P9, with novel compound heterozygous *TTN* mutations and a heterozygous *TARDBP* (Tar DNA-Binding Protein, MIM *605078) variant respectively while two novel heterozygous variants in the *FLNC* and the *ACTA1* (Actin, Alpha, skeletal muscle 1, MIM *102610) genes fulfilled the possible diagnosis overall pathogenicity score in patients P10 and P11 respectively.

Finally, 9 ICs remained without a molecular diagnosis following mutational analysis of the 306 genes of interest.

Considering only the first (definite) group of patients, the yield of diagnosed patients was 35% in this cohort (7/20).

DISCUSSION

Next-Generation Sequencing (NGS) is already used by many genetics laboratories and is being used with increasing frequency as the standard initial analysis for myopathies and other heterogeneous genetic disorders. The molecular diagnosis yield of 35% obtained in this study is consistent with other reports showing a range of 25 to 50 percent for rare genetic disorders diagnosis by WES^{23,24}.

Our study illustrates that a NGS approach is more efficient than the gene-by-gene strategy for several reasons. First, it allowed us to explore genes responsible for disorders within the differential diagnosis of hIBM, including *VCP* and *DES*. Second, our strategy permitted sequencing of large-sized genes, such as *TTN* and *FLNC*, which is not routinely performed, leading to the identification of variants in five index cases. Third, this approach allowed us to modify the incorrect diagnosis of hIBM in one patient, initially based on the presence of rimmed vacuoles on muscle biopsy, to a different muscle disorder caused by variants in the gene *GYGI*. Thus, NGS has the potential to alter a misdiagnosis due to a misleading muscle biopsy. Although using WES to explore a subset of genes might not provide as much target sequence coverage as a sequencing strategy specifically designed for these genes¹⁰, there are several advantages of using this approach. The sequencing results for samples where no pathogenic variants were identified in the initially explored genes can be reanalyzed to explore additional genes or all genes in the whole exome. In this way, further analyses are ongoing for the cases among our cohort that remain without genetic characterization. Another advantage of WES over targeted exome sequencing is its versatility and ability to be applied to many different diseases, as different sets of genes can be assessed without the need to develop and test a specific sequencing strategy^{25,26}.

In conclusion, the exome-based sequencing strategy described here is an efficient way to diagnose such genetically heterogeneous disorders as hIBMs.

Accepted Article

ABBREVIATIONS

ACMG: American College of Medical Genetics

AS: Acceptor Site

DS: Donor Site

ESE: Exonic Splicing Enhancer

hIBM: hereditary Inclusion Body Myopathies

HSF: Human Splicing Finder

IC: Index Case

NGS: Next-Generation Sequencing

PolyPhen-2: Polymorphism Phenotyping v2

SIFT: Sort Intolerant From Tolerant (amino acid substitutions)

VarAFT: Variant Analysis and Filtration Tool

WES: Whole Exome Sequencing

The authors sincerely thank Christel Castro, Jean-Pierre Desvignes, David Salgado, Christophe Bérout, Eric Salvo, Rafaele Bernard, Jocelyn Laporte, Johann Bohm and Mark Lathrop for their contributions to this work. We also thank the patients and their referring physicians for their participation.

REFERENCES

1. Askanas V, Engel WK. Sporadic inclusion-body myositis and hereditary inclusion-body myopathies: current concepts of diagnosis and pathogenesis. *Curr Opin Rheumatol*. 1998;10(6):530-542.
2. Huizing M, Carrillo-Carrasco N, Malicdan MC, Noguchi S, Gahl WA, Mitrani-Rosenbaum S, *et al*. *GNE* myopathy: New name and new mutation nomenclature. *Neuromuscul Disord* 2014;24(5):387–389.
3. Eisenberg I, Avidan N, Potikha T, Hochner H, Chen M, Olender T, Barash M, *et al*. The UDP-N-acetylglucosamine 2-epimerase/N-acetylmannosamine kinase gene is mutated in recessive hereditary inclusion body myopathy. *Nat Genet*. 2001;29(1):83-87.
4. Cerino M, Gorokhova S, Béhin A, Urtizberea JA, Kergourlay V, Salvo E, *et al*. Novel pathogenic variants in a French cohort widen the mutational spectrum of *GNE* myopathy. *Journal of Neuromuscular Diseases*. 2015;2(2):131-136. Published Online: 2015. DOI: 10.3233/JND-150074.
5. DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, *et al*. A framework for variation discovery and genotyping using next generation DNA sequencing data. *Nat Genet* 2011;43:491–498.
6. Wang K, Li M, Hakonarson H. ANNOVAR: Functional annotation of genetic variants from next-generation sequencing data. *Nucleic Acids Research*. 2010;38(16):164.
7. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics*. 2010;26(6):841-842.
8. Kaplan JC, Hamroun D. The 2015 version of the gene table of monogenic neuromuscular disorders (nuclear genome). *Neuromuscul Disord*. 2014;24(12):1123-1153.
9. Sevy A, Cerino M, Gorokhova S, Dionnet E, Mathieu Y, Verschuere A, *et al*. Improving molecular diagnosis of distal myopathies by targeted next-generation sequencing. *J Neurol Neurosurg Psychiatry*. 2016;87(3):340-342.
10. Gorokhova S, Cerino M, Mathieu Y, Courrier S, Desvignes JP, Salgado D, *et al*. Comparing targeted exome and whole exome approaches for genetic diagnosis of neuromuscular disorders. *Applied & Translational Genomics*. 2015;7:26–31. Published online: 2015. DOI:10.1016/j.atg.2015.07.006.
11. Salgado D, Desvignes JP, Rai G, Blanchard A, Miltgen M, Pinard A, *et al*. UMD-Predictor: a High Throughput Sequencing Compliant System for Pathogenicity Prediction of any Human cDNA Substitution. *Hum Mutat*. 2016. Published Online: 2016. DOI:10.1002/humu.22965.
12. Vaser R, Adusumalli S, Leng SN, Sikic M, Ng PC. SIFT missense predictions for genomes. *Nat Protoc*. 2016;11(1):1-9.

13. Adzhubei IA, Schmidt S, Peshkin L, Ramensky VE, Gerasimova A, Bork P, *et al.* A method and server for predicting damaging missense mutations. *Nat Methods*. 2010;7(4):248-249.
14. Desmet FO, Hamroun D, Lalande M, Collod-Bérout G, Claustres M, Bérout C. Human Splicing Finder: an online bioinformatics tool to predict splicing signals. *Nucleic Acids Res*. 2009;37:67.
15. Richards S, Aziz N, Bale S, Bick D, Das S, Gastier-Foster J, *et al.* Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med*. 2015;17(5):405-424.
16. Palmio J, Evilä A, Chapon F, Tasca G, Xiang F, Brådvik B, *et al.* Hereditary myopathy with early respiratory failure: occurrence in various populations. *J Neurol Neurosurg Psychiatry*. 2014;85(3):345-353.
17. Stojkovic T, Hammouda el H, Richard P, López de Munain A, Ruiz-Martinez J, Camaño P, *et al.* Clinical outcome in 19 French and Spanish patients with valosin-containing protein myopathy associated with Paget's disease of bone and frontotemporal dementia. *Neuromuscul Disord*. 2009;19(5):316-323.
18. Lange S, Xiang F, Yakovenko A, Vihola A, Hackman P, Rostkova E, *et al.* The kinase domain of titin controls muscle gene expression and protein turnover. *Science*. 2005;308(5728):1599-1603.
19. Hackman P, Marchand S, Sarparanta J, Vihola A, Pénisson-Besnier I, Eymard B, *et al.* Truncating mutations in C-terminal titin may cause more severe tibial muscular dystrophy (TMD). *Neuromuscul Disord*. 2008;18(12):922-928.
20. Bär H, Goudeau B, Wälde S, Casteras-Simon M, Mücke N, Shatunov A, *et al.* Conspicuous involvement of desmin tail mutations in diverse cardiac and skeletal myopathies. *Hum Mutat*. 2007;28(4):374-386.
21. Vorgerd M, van der Ven PF, Bruchertseifer V, Löwe T, Kley RA, Schröder R, *et al.* A mutation in the dimerization domain of filamin c causes a novel type of autosomal dominant myofibrillar myopathy. *Am J Hum Genet*. 2005;77(2):297-304.
22. Malfatti E, Nilsson J, Hedberg-Oldfors C, Hernandez-Lain A, Michel F, Dominguez-Gonzalez C, *et al.* A new muscle glycogen storage disease associated with glycogenin-1 deficiency. *Ann Neurol*. 2014;76(6):891-898.
23. Yang Y, Muzny DM, Reid JG, Bainbridge MN, Willis A, Ward PA, *et al.* Clinical whole-exome sequencing for the diagnosis of mendelian disorders. *N Engl J Med*. 2013;369(16):1502-1511.
24. Beaulieu CL, Majewski J, Schwartzentruber J, Samuels ME, Fernandez BA, Bernier FP, *et al.* FORGE Canada Consortium: outcomes of a 2-year national rare-disease gene-discovery project. *Am J Hum Genet*. 2014;94(6):809-817.

25. Lacoste C, Desvignes JP, Salgado D, Pecheux C, Villard L, Bartoli M, *et al.* Coverage Analysis of Lists of Genes involved in Heterogeneous Genetic Diseases following Benchtop Exome Sequencing using the Ion Proton. *J Genet.* 2016;95(1):203-208.
26. Dias C, Sincan M, Cherukuri PF, Rupps R, Huang Y, Briemberg H, *et al.* An analysis of exome sequencing for diagnostic testing of the genes associated with muscle disease and spastic paraplegia. *Hum Mutat.* 2012;33:614–626.
27. Conforti F.L., Sproviero W., Simone I.L., Mazzei R., Valentino P., Ungaro C., *et al.* TARDBP gene mutations in south Italian patients with amyotrophic lateral sclerosis. *J Neurol Neurosurg Psychiatry* 2011;82:587-588.
28. Solski J.A., Yang S., Nicholson G. A., Luquin N., Williams K.L., Fernando R. *et al.* A novel TARDBP insertion/deletion mutation in the flail arm variant of amyotrophic lateral sclerosis. *Amyotrophic Lateral Sclerosis Journal.* 2012;13(5):465-470.
29. Lattante S., Rouleau G.A. and Kabashi E. *TARDBP* and *FUS* Mutations Associated with Amyotrophic Lateral Sclerosis: Summary and Update. *Hum Mutat.* 2013;34(6):812–826.

Table 1: Pathogenicity assessment for the identified variants in patients with definite, probable and possible diagnoses

Patient	Gender	Phenotype / Genetic inheritance	Muscle biopsy	Genes/Variants	Status	UMD-predictor ¹¹	SIFT ¹²	PolyPhen-2 ¹³	ACMG Guidelines ¹⁵	Splicing prediction (HSF ¹⁴)	Pathogenic variants described in literature
Patients with definite diagnosis											
P1	F	PM of lower limbs at onset (40yo) evolving towards HMREF / AR	Rimmed vacuoles Cytoplasmic inclusions Disruption of the intermyofibrillar network	<i>TTN</i> : c.95195C>T (p.Pro31732Leu)	HOZ	Pathogenic	Damaging	Probably damaging	Pathogenic	NP	YES ¹⁶
P2	F	DM of upper and lower limbs (tibialis anterior muscle) No axial muscle weakness / AD	Rimmed vacuoles Disruption of the intermyofibrillar network	<i>VCP</i> : c.410C>T (p.Pro137Leu)	HTZ	Pathogenic	Damaging	Probably damaging	Pathogenic	Not affected	YES ¹⁷
P3	F	Early onset (childhood) DM of lower limbs (tibial muscular dystrophy) slowly evolving / AR	Rimmed vacuoles Dystrophic muscle biopsy	<i>TTN</i> : c.102271C>T (p.Arg34091Trp)	HTZ	Pathogenic	Damaging	Probably damaging	Pathogenic	NP	YES ¹⁸
				<i>TTN</i> : c.107647delT (p.Ser35883Glnfs*10)	HTZ	NP	NP	NP	Pathogenic	NP	YES ¹⁹
P4	M	Late onset (45yo) DM of upper and lower limbs with cardiac involvement / AD	Rimmed vacuoles Atrophic fibers Disorganized myofibrillar network	<i>DES</i> : c.1360C>T (p.Arg454Trp)	HTZ	Pathogenic	Damaging	Probably damaging	Pathogenic	Not affected	YES ²⁰
P5	M	DM of upper and lower limbs. Onset at 30yo / AD	Rare rimmed vacuoles (<5)	<i>VCP</i> : c.410C>T (p.Pro137Leu)	HTZ	Pathogenic	Damaging	Probably damaging	Pathogenic	Not affected	YES ¹⁷
P6	F	Late onset (45yo) DM of lower limbs and pelvic girdle myopathy / AD	Rimmed vacuoles	<i>FLNC</i> : c.8130G>A (p.Trp2710*)	HTZ	NP	NP	NP	Pathogenic	NP	YES ²¹
P7	F	Late onset (45yo) DM of upper and lower limbs with slow evolution / AR	Rimmed vacuoles (on the initial biopsy) recharacterized as polyglucosan bodies (on the second biopsy)	<i>GYGI</i> : c.143+3G>C (p.Asp3Glufs*4)	Comp. HTZ	NP	NP	NP	Pathogenic	Probably damaging	YES ²²
				<i>GYGI</i> : c.996_1005del10 (p.Tyr332fs*1)		NP	NP	NP	Pathogenic	NP	NO
Patients with probable diagnosis											
P8	M	Early onset (14yo) DM of lower limbs (tibial muscular dystrophy) evolving towards hamstring muscle with quadriceps sparing / AR	Rimmed vacuoles	<i>TTN</i> : c.15346C>T (p.Arg5116*)	HTZ	NP	NP	NP	Pathogenic	NP	NO
				<i>TTN</i> : c.107680G>A (p.Gly35894Arg)	HTZ	Pathogenic	Damaging	Probably damaging	Likely pathogenic	NP	NO
* P9	M	Late onset (50yo) DM of upper and lower limbs / AD	Rimmed vacuoles No inflammation Atrophic fibers	<i>TARDBP</i> : c.1127G>T (p.Gly376Val)	HTZ	Pathogenic	Tolerated	Benign	Likely pathogenic	Possibly damaging	† NO
Patients with possible diagnosis											
P10	M	Limb girdle muscular dystrophy / AD	Rare rimmed vacuoles (<5)	<i>FLNC</i> : c.6526C>T (p.Arg2176Cys)	HTZ	Pathogenic	Tolerated	Probably damaging	Uncertain significance	Not affected	NO
P11	M	DM of lower limbs with very slow evolution / AD	Rimmed vacuoles	<i>ACTA1</i> : c.437C>T (p.Ala146Val)	HTZ	Pathogenic	Tolerated	Probably damaging	Uncertain significance	Uncertain	NO

In bold: pathogenicity prediction strong and moderate; **NP:** not performed (UMD-predictor, SIFT and PolyPhen-2 algorithms do not provide a pathogenicity score for variants creating a stop or a frameshift); **UMD-predictor¹¹:** Universal Mutation Database predictor; **SIFT¹²:** *Sort Intolerant From Tolerant*; **PolyPhen-2¹³:** Polymorphism Phenotyping v2; **HSF¹⁴:** Human Splicing Finder; **ACMG¹⁵:** American College of Medical Genetics; **AD:** Autosomal Dominant; **AR:** Autosomal Recessive; **HMREF:** *Hereditary Myopathy with early REspiratory Failure*; **DM:** Distal Myopathy; **PM:** Proximal Myopathy; **yo:** years old; **HOZ:** homozygous; **HTZ:** heterozygous; **Comp. HTZ:** compound heterozygous (with confirmed segregation analysis).

Frequency in 1000G, ESP and ExAC databases of all the variants described in Table 1 is lower than 0.2%.

* **Additional variant** with uncertain significance found for patient P9: *MYH2*: c.2090A>G (p.His697Arg).

† **Variant affecting the same nucleic and amino acid positions** as another variant, c.1127G>A (p.Gly376Asp), previously described in the literature^{27,28,29} in two different familial amyotrophic lateral sclerosis cases with a similar phenotype presentation as patient P9 (upper and lower limb weakness with no cognitive impairment).