



Ammonia agricultural emissions over Europe as seen by IASI and impact on PM2.5 concentrations

Audrey Fortems-Cheiney, Gaelle Dufour, Matthias Beekmann, Sophie Générmont, Frédéric Meleux, Cathy Clerbaux, Pierre-François Coheur, Martin van Damme, Lieven Clarisse

► To cite this version:

Audrey Fortems-Cheiney, Gaelle Dufour, Matthias Beekmann, Sophie Générmont, Frédéric Meleux, et al.. Ammonia agricultural emissions over Europe as seen by IASI and impact on PM2.5 concentrations. EGU 2018, European Geophysical Union General Assembly, Apr 2018, Vienne, Austria. , 20, pp. EGU2018-12929-1, 2018, Geophysical Research Abstracts. hal-01765427

HAL Id: hal-01765427

<https://hal.science/hal-01765427>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License



Ammonia agricultural emissions over Europe as seen by IASI and impact on PM_{2.5} concentrations

Audrey Fortems-Cheiney (1), Gaelle Dufour (1), Matthias Beekmann (1), Sophie Genermont (2), Frederik Meleux (3), Cathy Clerbaux (4), Pierre-Francois Coheur (5), Martin van Damme (5), and Lieven Clarisse (5)

(1) LISA UMR7583, Creteil, France (audrey.cheiney@lisa.u-pec.fr), (2) INRA/AgroParisTech, Paris, France, (3) INERIS, Verneuil en Halatte, France, (4) LATMOS UMR8190, Paris, France, (5) ULB, Brussels, Belgium

Ammonia (NH₃), whose main source is agriculture, is an important gaseous precursor of atmospheric particulate matter (PM). However, NH₃ is the most poorly understood pollutant regulated by EU directives for air quality, as flux quantification is highly uncertain and measurement of this compound at the surface of the earth is difficult. Here, we derived daily European ammonia emissions using NH₃ total columns from the Infrared Atmospheric Sounding Interferometer (IASI) onboard Metop-A, at a relatively high spatial resolution (grid-cell of 0.5° x 0.5°) for the two years 2010 and 2011. This update of NH₃, in magnitude with a relatively high spatio-temporal variability, allows for a better comparison with independent PM_{2.5} measurements. It shows a different seasonal cycle in 2011 with a strong peak in March 2011, rather than in April in the EMEP prior inventory. It also reveals a peak in August 2011, not described in the EMEP inventory.

This preliminary study suggests that there are good promises for better quantifying NH₃ emissions by atmospheric inversions, and particularly with the 4D Bayesian variational inverse system PYVAR-CHIMERE -based on the adjoint model of CHIMERE- being adapted to NH₃.