

Determinants of mercury contamination in viperine snakes, *Natrix maura* , in Western Europe

Jérémy Lemaire, Paco Bustamante, Anthony Olivier, Olivier Lourdais, Bruno Michaud, Alexandre Boissinot, Pedro Galán, Francois Brischoux

► **To cite this version:**

Jérémy Lemaire, Paco Bustamante, Anthony Olivier, Olivier Lourdais, Bruno Michaud, et al.. Determinants of mercury contamination in viperine snakes, *Natrix maura* , in Western Europe. Science of the Total Environment, Elsevier, 2018, 635, pp.20 - 25. 10.1016/j.scitotenv.2018.04.029 . hal-01769909

HAL Id: hal-01769909

<https://hal.archives-ouvertes.fr/hal-01769909>

Submitted on 8 Jan 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Determinants of mercury contamination in viperine snakes, *Natrix maura*, in Western Europe

Jérémy Lemaire ^{a,b}, Paco Bustamante ^c, Anthony Olivier ^d, Olivier Lourdais ^a, Bruno Michaud ^a, Alexandre Boissinot ^a, Pedro Galán ^e, François Brischoux ^{a,□}

^aCentre d'Etudes Biologiques de Chizé, CEBC UMR 7372 CNRS-Université de La Rochelle, 79360 Villiers en Bois, France

^bRéserve Naturelle Nationale de Chérine, Maison de la Nature et de la Réserve, 36290 Saint-Michel-en-Brenne, France

^cLittoral Environnement et Sociétés (LIENSs), UMR 7266 CNRS-Université de La Rochelle, 2 rue Olympe de Gouges, 17000 La Rochelle, France


^dInstitut de recherche de la Tour du Valat, Le Sambuc, 13200 Arles, France

^eGrupo de Investigación en Biología Evolutiva (GIBE), Departamento de Biología, Facultad de Ciencias, Universidad de A Coruña, Campus da Zapateira, s/n., 15071 A Coruña, Spain

Corresponding author. E-mail address: francois.brischoux@gmail.com. (F. Brischoux).

How to cite : Lemaire J., Bustamante P., Olivier A., Lourdais O., Michaud B., Boissinot A., Galán P., Brischoux F. 2018. Determinants of mercury contamination in viperine snakes, *Natrix maura*, in Western Europe. *Sci Total Environ* 635:20–25. doi: 10.1016/j.scitotenv.2018.04.029, Accepted 3 April 2018

- Hg contamination in freshwater mesopredators has been largely overlooked.
- Hg concentrations were measured in scales of viperine snakes in France and Spain.
- Viperine snakes do accumulate Hg in their scales.
- Diet (fish versus amphibians) influenced accumulation rates of Hg.
- Highest values of Hg were found in individuals from a fish farm.



ABSTRACT

The effects of Hg contamination are presumably widespread across the components of aquatic ecosystems, but investigations have been mainly focused on freshwater fish, because this biota represents a major source of Hg for human populations. Yet, the possible bioaccumulation of Hg on other freshwater meso- and apex predators (e.g., amphibians, reptiles) has been largely overlooked, especially in Western Europe. In this study, the determinants of Hg concentrations were assessed for the viperine snake (*Natrix maura*) across 6 populations (>130 individuals sampled in 2016 and 2017) in France and Spain. Specifically, body size, sex, and diet were compared with Hg concentrations measured in ventral scales. Overall, *N. maura* accumulated Hg in their scales. Sex did not seem to influence Hg concentrations in this species. Significant differences in Hg concentrations were observed between study sites, and these differences were likely to be mediated by site-specific diet. Frog-eating individuals were characterized not only by lower mean values of Hg ($0.194 \pm 0.018 \mu\text{g}\cdot\text{g}^{-1}$ versus $0.386 \pm 0.032 \mu\text{g}\cdot\text{g}^{-1}$ for piscivorous individuals), but also by weaker slopes of the body size-Hg relationship as compared to fish-eating snakes, suggesting strong differences in accumulation rates due to food resources. Importantly, the highest slope of the body size-Hg relationship and the highest values of Hg were found in individuals foraging on trout raised by a fish farm, suggesting that fish farming may contribute to Hg contamination in inland freshwater systems. Finally, our results are compared with data on Hg concentrations in other species of aquatic snakes, in order to provide a comparative point for future studies.

1. Introduction

Mercury (Hg) is a well-known environmental contaminant which can originate from both natural and anthropogenic sources (Fitzgerald et al., 2007; Selin, 2009). Atmospheric and water circulation patterns tend to concentrate Hg in aquatic environments (Mason et al., 2012). In addition, anoxic conditions found in slow-moving water bodies promote the transformation of inorganic Hg in methyl-Hg by microorganisms (Compeau and Bartha, 1985; Jensen and Jernelöv, 1969), which is known for its toxic effects on humans and wildlife (e.g. Tan et al., 2009; Scheuhammer et al., 2008). Finally, Hg can be bioaccumulated within organisms and

biomagnified through the food chain, and concentrated in apex predators as fishes, birds or mammals (Mason et al., 1995; Atwell et al., 1998; Power et al., 2002).

Over the last decades, the potential deleterious effects of Hg have been investigated from both an ecological and public health perspective (Driscoll et al., 2013; Lavoie et al., 2013; Eley, 1997; Wolfe et al., 1998). Effects of Hg are multiple and cover a large spectrum of syndromes such as neurological dysfunction (Steuerwald et al., 2000; Basu et al., 2005; Clarkson and Magos, 2006; Scheuhammer and Sandheinrich, 2007; Depew et al., 2012), endocrine disorders (Wada et al., 2009; Meyer et al., 2014) or altered reproduction and offspring quality (Klaper et al., 2006; Burgess and Meyer, 2008; Bergeron et al., 2011; Hopkins et al., 2013a; Tartu et al., 2013). Heretofore, the comprehensive assessments of Hg contamination across complex environments and trophic webs are challenging. Indeed, although the toxic effects of Hg are presumably widespread across the components of aquatic ecosystems, investigations of Hg contamination have been mainly focused on freshwater fish (e.g., Depew et al., 2013; Åkerblom et al., 2014; Scheuhammer et al., 2014; Eagles-Smith et al., 2016), probably because this biota represents both a major source of protein for many human populations (Futsaeter and Wilson, 2013; Dong et al., 2015; Lepak et al., 2016; Fliedner et al., 2016). Aquatic birds have also attracted considerable scientific attention in this respect (Ackerman et al., 2016; Jackson et al., 2016; Blukacz-Richards et al., 2017; Sullivan and Kopec, 2018; Żarski et al., 2017; see also Whitney and Cristol, 2017 for a review). Yet, other aquatic vertebrates than fishes, and especially meso- and apex-predators, may well suffer from Hg contamination (e.g. Driscoll et al., 2007). Such overlooked organisms include amphibians (e.g., Bergeron et al., 2010; Todd et al., 2011), aquatic snakes (e.g., Burger et al., 2005; Drewett et al., 2013), and turtles (e.g., Meyer et al., 2014; Slimani et al., 2017). This is especially true in Western Europe where investigations of Hg contamination in aquatic tetrapods are very scarce as compared with other geographic areas such as Northern America. Nevertheless, inclusion of these organisms is of crucial importance if we are to globally assess Hg contamination worldwide, and in turn to monitor its effects on biodiversity and human health (Gustin et al., 2016).

In addition, some lineages of these overlooked meso- and apex predators also provide a unique set of features that make them useful biological tools to monitor Hg contamination in the wild (e.g., Burger et al., 2005; Slimani et al., 2017). For instance, as compared with highly mobile fish and birds, aquatic reptiles and amphibians are characterized by high levels of philopatry associated with relatively low capacities for large scale movements (Hillman et al., 2014). As a consequence, Hg concentrations in their tissues should strongly reveal those of their relatively small home ranges while highly mobile organisms such as fish and birds may provide information that integrates Hg contamination over large distances, and thus, different environments (Burger et al., 2007; Drewett et al., 2013; Slimani et al., 2017). Additionally, aquatic reptiles are situated relatively high in the trophic web, and as ectotherms, they display relatively low metabolic rates and relatively high tissue conversion rates of their food resources which should enhance their capacity to integrate long-term Hg contamination in their tissues. Characterized by an indeterminate growth, many aquatic reptile species display very wide size range between minute neonates and large adult individuals which allow access to bioaccumulation processes within a population. Finally, easily accessible tissues such as claws in turtles or scales in snakes, in which Hg tends to accumulate and bind to keratins (Hopkins et al., 2013b), provide a powerful opportunity to adopt a non-invasive technique in order to assess Hg contamination in these organisms (Schneider et al., 2015).

In this study, Hg concentrations were investigated in a widely distributed European semi-aquatic Natricinae, the viperine snake (*Natrix maura*). More than 130 individuals distributed across 6 populations situated in France and Spain were sampled. These populations were associated with contrasted ecological contexts and dominant trophic resources (mainly amphibians in 3 sites and fishes in 3 other sites, Table 1). A non-invasive technique (scale-clipping) was used in order to assay Hg concentrations across this wide geographic area. In addition, on a sub-sample of individuals, scale clipping was combined blood sampling to assess the relationship between Hg concentrations in the blood (reflecting short term Hg exposure, i.e., weeks) and in the scales (integrating Hg exposure over a longer time scale, i.e., months) in this species (as shown in other semi-aquatic snake species, Burger et al., 2005). Specifically, the aims of this study were:

- 1) to assess Hg contamination in this species across a large ecological context,
- 2) to investigate bioaccumulation rates across a wide range of body size,
- 3) and to examine the influence of sex and diet on Hg contamination and bioaccumulation rates.

2. Material and methods

2.1. Study species and study sites

The viperine snake (*Natrix maura*) is a semi-aquatic freshwater natricine widely distributed across Western Europe and Northern Africa, broadly from France to Morocco (Miras et al., 2015). This relatively small-sized species (up to ~80 cm total length) typically forages for fish and amphibians in aquatic environments such as streams, rivers, marshes, and lakes (Miras et al., 2015; Santos and Llorente, 2009). A skin-shedding occurs in *N. maura* at the onset of the activity period in early spring, while another shedding cycle is later associated with ovulation shortly before oviposition in females (June/July). Our sampling occurred in late spring between these two periods.

The six study sites were distributed in France and Spain and cover the different habitat types in which *N. maura* typically occurs (Table 1, see also graphical abstract). From north to south, the study sites were 1 - Réserve Naturelle de Chérine within the Brenne Natural Park, France (hereafter "Brenne"), 2 - Lac du Cébron, France (hereafter "Cébron"), 3 - The Boutonne river at Fontenille-Saint

Martin d'Entraigues, France (hereafter "Fontenille"), 4 - The Réserve Naturelle de Moëze-Oléron, France (hereafter "Moëze"), 5 - The Réserve Naturelle de la Tour du Valat, France (hereafter "Tour du Valat"), and 6 - Ons island, Spain (hereafter "Ons").

Table 1

Description of the study sites. Site number refers to their location in the graphical abstract. Diet was assessed through opportunistic regurgitations obtained in the field. Identified frog species were *Pelophylax* spp. for Brenne, *Pelophylax* spp. and *Hyla meridionalis* for Moëze and Tour du Valat. Identified fish species were *Perca fluviatilis* and *Gymnocephalus cernua* for Cébron, and *Blennius* spp. for Ons.

Number	Name	Coordinates	Habitat type	N (females/males)	Diet	Sampling period
1	Chérine	46°47'24.18"N, 1°12'2.79"E	Small lakes and ponds	11 (6/5)	Amphibians	May–June 2016
2	Cébron	46°46'5.47"N, 0°11'46.85"W	Artificial lake	22 (14/8)	Fish	May 2016
3	Fontenille	46°7'26.07"N, 0°8'23.98"W	Fish farm	27(16/11)	Fish	April–June 2016
4	Moëze	45°54'14.16"N, 1°4'22.33"W	Atlantic wetlands and channels	26(13/13)	Amphibians	April–June 2016
5	Tour du Valat	43°30'32.82"N, 4°39'49.14"E	Mediterranean wetlands and	29(17/12)	Amphibians	April–June 2016
6	Ons Island	42°22'32.59"N, 8°56'10.22"W	Oceanic island	13 (3/10)	Fish	June 2017

2.2. Field procedures and sampling

Snakes were captured between April and June 2016 and 2017 (Table 1) by hand either under artificial refuges deployed for snake captures or upon sighting. Snakes were measured on a flexible ruler (snoutvent length, SVL; and total length, TL, to 0.5 cm), weighted on a digital scale (to 0.1 g), and sexed by eversion of the hemipenis. Snakes were individually marked by ventral scale-clipping followed by heat branding using a surgical cautery (Bovie Medical Corporation). The scale clips obtained after marking were collected, dried on paper towel, and stored at ambient temperature in closed tubes for further Hg analyses (see below).

On a subsample of snakes from Tour du Valat (N = 8), we also collected blood samples in order to assess the relationships between blood and skin (scales) Hg in *N. maura*. Blood (~150 µl) was obtained through cardiocentesis with a heparinized 30-gauge needle and a 1 ml syringe. Whole blood was stored at -20 °C in sealed tubes until Hg analyses (see below).

At the end of the field procedures, all snakes were released at their place of capture usually within 30 min.

2.3. Mercury assays

Total Hg concentrations in skin and whole blood were determined using an atomic absorption spectrometer AMA-254 (Advanced Mercury Analyser-254; Altec®). At least two aliquots of 1–5 mg of subsamples for each individual were analysed. The accuracy and reproducibility of the Hg measurements was assessed by the analyses of certified reference material (CRM) TORT-2 (Lobster Hepatopancreas from the National Research Council of Canada; certified Hg concentration: 0.27 ± 0.06 µg·g⁻¹ dw) at the beginning and at the end of the analytical cycle, and by running CRM for every 10 samples (Bustamante et al., 2006). Measured values were 0.241 ± 0.014 µg·g⁻¹ dw, n = 17 (recovery 90 ± 3%). Mass of the CRM was adjusted to represent an amount of Hg similar to that in the samples. Blanks were analysed at the beginning of each set of samples and the limit of detection was 0.005 µg·g⁻¹ dry weight (dw). Hg concentrations in snake tissues further are expressed in µg·g⁻¹ dw.

3. Results

3.1. Scale versus blood Hg concentrations

Hg concentrations between blood and scale clips were positively correlated ($F_{1,6} = 8.87$, $r^2 = 0.59$, $p = 0.02$). However, blood Hg concentrations were 3 times higher than scale Hg concentrations (paired t-test, $t = 8.01$, $p < 0.0001$, $0.719 \pm 0.267 \mu\text{g}\cdot\text{g}^{-1}$ versus $0.237 \pm 0.195 \mu\text{g}\cdot\text{g}^{-1}$, respectively).

3.2. Determinants of Hg levels

Overall, a significant positive relationship was observed between snake size (SVL) and Hg concentration ($F_{1,130} = 43.95$, $r^2 = 0.25$, $p < 0.0001$). Within sites, the same trend was detectable (all $p < 0.005$, Fig. 2), except for Tour du Valat ($p = 0.16$, Fig. 1). The strength of the relationship was highly variable between sites, with three sites characterized by relatively weak slopes (Brenne, 0.009; Moëze, 0.007) and/or non-significant correlation (Tour du Valat), and three sites characterized by steeper slopes (Cébron, 0.019; Fontenille, 0.029; Ons, 0.015, Fig. 1). Interestingly, these two sets of sites differed according to the inferred diet of the snakes ($\chi^2 = 132$, $df = 5$, $p < 0.0001$, Figs. 1 & 2).

Accordingly, differences in Hg concentrations were found between sites (ANCOVA with the SVL as the covariate, $F_{5,124} = 11.81$, $p < 0.0001$, Fig. 2), and also a strong influence of inferred diet on Hg values with fish-eating snakes having higher Hg concentrations than frog-eating ones (ANOVA, $F_{1,129} = 30.63$, $p < 0.0001$, $0.386 \pm 0.032 \mu\text{g}\cdot\text{g}^{-1}$ versus $0.194 \pm 0.018 \mu\text{g}\cdot\text{g}^{-1}$ respectively, Fig. 2).

Finally, there were no sex related differences in Hg concentrations, either across all study sites ($F_{1,126} = 0.56$, $p = 0.46$), or when analyses were restricted to sites with balanced samples between males and females, and with N > 10 individuals in each category (Tour du Valat $F_{1,26} = 0.08$, $p = 0.78$; Moëze $F_{1,23} = 2.09$, $p = 0.16$; Fontenille $F_{1,25} = 0.18$, $p = 0.68$).

4. Discussion

This study provides an insight into Hg bioaccumulation in a freshwater mesopredator, the viperine snake (*N. maura*), over a large spatial range and across contrasting habitats. Overall, *N. maura* do accumulate Hg in their scales. Hg concentrations between blood (short-term Hg exposure, i.e., weeks) and scales (integrating Hg exposure over a longer time scale, i.e., months) were correlated, but that blood display higher Hg concentrations than scale clips. Sex did not seem to influence Hg concentrations in this species. Finally, strong differences of Hg concentrations between study sites that are likely to be mediated by sitespecific diet. Frog-eating individuals were characterized not only by lower mean values of Hg concentrations, but also by weaker slopes of the body size-Hg relationship as compared to fish-eating snakes, suggesting differences in Hg accumulation rates due to food resources. All of these findings are discussed sequentially below.

A positive correlation was observed between Hg concentrations quantified in skin (scale clips) and blood. As already demonstrated in other studies, this suggests that scale clipping (a non-invasive sampling technique) is a valuable method for monitoring Hg contamination in snake species (Burger et al., 2005) and that *N. maura* can be used as a bio indicator of Hg contamination. Interestingly, despite this broad relationship, significant differences were found in Hg values, with blood showing Hg concentrations three times higher than skin ($0.719 \pm 0.267 \mu\text{g}\cdot\text{g}^{-1}$ versus $0.237 \pm 0.195 \mu\text{g}\cdot\text{g}^{-1}$). Because these two tissues integrate Hg exposure at different time-scales (weeks for blood versus months for skin), this result suggests a possible seasonal shift in the feeding ecology of snakes from the Tour du Valat population, or less likely a seasonal shift in Hg content of their prey. It is likely that foraging ecology of this species, and especially prey availability and catchability shifts during the spring. Indeed, amphibians (with presumably lower Hg concentrations, see below) are more abundant in early spring (e.g., amphibian breeding season and activity peak), while we can posit that fish (with presumably higher Hg content, see below) become progressively more important in the diet of *N. maura* with increasing water temperatures that may allow better rates of fish capture

23




Figure 1 : Relationship between body size (snout-vent length) and Hg concentration measured in scale-clips. Left-hand side panels show data for frog-eating populations while right-hand side panels show data for fish-eating snakes.

Interestingly, the relatively high degree of variation of Hg concentrations between skin and blood in *N. maura* seem relatively uncommon in other species that share similar ecology (semi-aquatic snake species), where skin and blood Hg values tend to be very similar (Table 2). Importantly, the seasonal variation in Hg exposure suspected in *N. maura* from the Tour du Valat population may induce intense seasonal pulses in Hg contamination in this species, and the ecotoxicological consequences of these seasonal pulses (as compared to an exposure to constant values of Hg) clearly deserve further attention.

Although sex differences in Hg concentrations are expected if females transfer maternal Hg to their eggs during vitellogenesis (Hopkins et al., 2004), there was no influence of sex on Hg concentrations either across or within our study populations. Some of our sample sizes were modest, and may have hampered our ability to detect such difference. However, this was not the case for all of our study sites (see Table 1). Alternatively, we can hypothesise that the foraging ecology of males and females did not differ in our study populations, especially as body size were very similar between sexes in our sampled individuals. This result may also reflect the time period of the study. While Hg concentrations in the skin integrates Hg contamination occurring since the last shedding cycle, maternal transfer of Hg in eggs en-compasses the whole vitellogenesis process (i.e., from early spring to early summer when oviposition occurs).

Table 2

Review of Hg concentrations assessed in freshwater semi-aquatic snake species in Western Europe (this study) and Northern America. SVL stands for snout-vent length. Hg levels are given as the mean \pm SE ($\mu\text{g}\cdot\text{g}^{-1}$ dw).

Species	Site	Tissue	SVL (cm)	Hg ($\mu\text{g}\cdot\text{g}^{-1}$)	Reference	
<i>Natrix maura</i>	Brenne, France	Skin	32.3 f 12.0	0.145 f 0.071	This study	
	Cébron, France	Skin	35.6 f 8.8	0.289 f 0.218		
	Fontenille, France	Skin	35.9 f 6.9	0.439 f 0.261		
	Moëze, France	Skin	39.5 f 11.7	0.183 f 0.159		
	Tour du Valat, France	Skin	38.6 f 8.1	0.220 f 0.153		
		Blood	39.2 f 6.4	0.719 f 0.267		
	Ons, Spain	Skin	44.5 f 13.1	0.438 f 0.241		
		Savannah River Site, South Carolina, USA	Blood	57		0.1 f 0.04
	<i>Agkistrodon piscivorus</i>		Blood	51		0.400 f 0.047
		<i>Nerodia fasciata</i>	Skin	–		0.423 f 0.028
<i>Nerodia sipedon</i>	Blood		–	0.407 f 0.042	Burger et al., 2005	
	East Fork Poplar Creek, Tennessee, USA	Skin	61	0.500 f 0.045	Campbell et al., 2005	
Little River, Tennessee, USA		Blood	–	0.372 f 0.046	Campbell et al., 2005	
	Skin	57.5	0.365 f 0.032	Campbell et al., 2005		
Raritan Canal, New Jersey, USA		Blood	–	0.436 f 0.065	Campbell et al., 2005	
	Skin	~55	0.159 f 0.023	Burger et al., 2007		
South River in Waynesboro, Virginia, USA		Blood	59	0.128 f 0.026	Burger et al., 2007	
	<i>Nerodia taxispilota</i>	Blood	59	2.24 f 0.42	Drewett et al., 2013	
Savannah River Site, South Carolina, USA		Blood	59	0.7 f 0.015	Burger et al., 2006	

Therefore, if maternal transfer of Hg occurs in this species, we can expect that Hg concentrations in females would decrease during vitellogenesis, and that such diminution would be detectable after the second skin shedding cycle (early summer). Future studies would usefully explore temporal variation of Hg concentrations in the scale-clips of both males and females, in order to assess putative maternal transfer of Hg to the eggs, and to assess sex differences in Hg contamination rates.

The Hg concentrations quantified in *N. maura* are comparable to those recorded in other species sharing a similar ecology (semi-aquatic snake species feeding on both fish and amphibians, Table 2). Yet, in our study, *N. maura* was characterized by an overall smaller body size than the other species (Table 2), which suggest that accumulation rates may be higher in this species. Additionally, a strong influence of the study site on Hg concentrations in *N. maura* was found that seemed to be mediated by site-specific diet. Indeed, sites where *N. maura* consume predominantly amphibians were characterized not only by lower mean values of Hg, but also by relatively weaker slopes of the body size-Hg relationships, suggesting strong differences in accumulation rates between sites. Because amphibians (frogs) are situated at lower trophic levels than fishes (Vander Zanden et al., 1997), and because they rely mostly on airborne and terrestrial food sources (insects), they may display both lower Hg concentrations and higher proportion of inorganic Hg that fish that feed on aquatic food sources. Such probable difference in Hg concentrations of fish versus frogs likely explains the strong influence of diet on Hg values, with relatively lower concentrations in frog-eating as compared to fish-eating snakes. Yet, the possible influence of diet on Hg concentrations in *N. maura* we hypothesised will necessitate specific investigations in order to assess Hg values in prey (frogs versus fish) in our different study sites, as well as assessing the relationships between trophic levels of both the snakes and their prey (i.e., measured through stable isotopes) and Hg contamination.




Figure 2 : Mean Hg concentrations (\pm SE) measured in scale-clips. Grey symbols represent frog-eating populations, while black symbols show fish-eating snakes. For clarity, data are presented by ascending order

Finally, we emphasize the specific case of the Fontenille population of *N. maura*, which display both the highest slope of the body size-Hg concentration relationship and the highest value of size-specific Hg values with value reaching almost $1 \mu\text{g}\cdot\text{g}^{-1}$ in individuals <45 cm SVL, as compared to similar values in snakes >55 cm SVL in other populations (Figs.1 and 2, see also Table 2). This study site is situated on a fish farm, and *N. maura* from this population feed heavily on the fry and juveniles trouts raised by the fish farm. The relatively higher value of Hg concentrations and Hg accumulation rates in this population dovetail remarkably well with processes that have recently been described in highaltitude aquatic ecosystems (Hansson et al., 2017). In this study system, farmed trout raised on commercial pellets based on proteins from fishery products of marine origin show both high Hg concentrations and Hg isotopic signatures that are comparable to that of top-predator marine biota (Hansson et al., 2017). It is likely that the commercial pellets used to raise trouts at the Fontenille fish farm are also based on proteins from fishery products of marine origin, and that introduction of marine Hg through the fish food is responsible for the higher value of Hg concentrations and Hg accumulation rates in this *N. maura* population. Interestingly, both Hg concentrations and Hg accumulation rate of the Fontenille population are even higher than the Ons population which feed on marine fish (Galán, 2012). Future studies should usefully assess the relative contribution of fish farming on Hg contamination in aquatic ecosystems and its role as a vector of Hg of marine origin in inland freshwater systems (Hansson et al., 2017).

Acknowledgements

We thank all the people that were involved in snake surveys, and especially to Mélanie Bouyssou, Pierre Rousseau, Coralie Bossu, Mathieu Normand, Romain Lengagne, Fabrice Conort and Gustavo Cochon. We also thank the staff from the Natural Reserves in which some of our sampling occurred (Réserve Naturelle Nationale de Chérine and de Moëze-Oléron and Réserve Naturelle Régionale de la Tour du Valat in France and the Parque Nacional das Illas Atlánticas de Galicia in Spain), the staff from the Pisciculture de Lussay and the Conseil Départemental des Deux-Sèvres (Sabrina Lefebvre). We are also very grateful to Carine Churlaud and Maud Brault-Favrou from the plateforme Analyses Élémentaires of LIENSs for their advices during Hg analyses. We thank Mae Sexauer Gustin and two anonymous reviewers for their insightful comments on a previous version of our ms. Images for the graphical abstract were taken from phylopic.org. Funding was provided by the CNRS, the Agence de l'Eau Loire-Bretagne and the Agence de l'Eau Adour-Garonne. The IUF (Institut Universitaire de France) is acknowledged for its support to P. Bustamante as a Senior Member.

References

- Ackerman, J.T., Eagles-Smith, C.A., Herzog, M.P., Hartman, C.A., Peterson, S.H., Evers, D.C., Jackson, A.K., Elliott, J.E., Vander Pol, S.S., Bryan, C.E., 2016. Avian mercury exposure and toxicological risk across western North America: a synthesis. *Sci. Total Environ.* 568, 749–769.
- Åkerblom, S., Bignert, A., Meili, M., Sonesten, L., Sundbom, M., 2014. Half a century of changing mercury levels in Swedish freshwater fish. *Ambio* 43 (S1), 91–103.
- Atwell, L., Hobson, K.A., Welch, H.E., 1998. Biomagnification and bioaccumulation of mercury in an arctic marine food web: insights from stable nitrogen isotope analysis. *Can. J. Fish. Aquat. Sci.* 55 (5), 1114–1121.
- Basu, N., Scheuhammer, A.M., Grochowina, N.M., Klenavic, K., Evans, R.D., O'Brien, M., Chan, H.M., 2005. Effects of mercury on neurochemical receptors in wild river otters (*Lontra canadensis*). *Environ. Sci. Technol.* 39, 3585–3591.
- Bergeron, C.M., Bodinof, C.M., Unrine, J.M., Hopkins, W.A., 2010. Mercury accumulation along a contamination gradient and nondestructive indices of bioaccumulation in amphibians. *Environ. Toxicol. Chem.* 29 (4), 980–988.
- Bergeron, C.M., Hopkins, W.A., Todd, B.D., Hepner, M.J., Unrine, J.M., 2011. Interactive effects of maternal and dietary mercury exposure have latent and lethal consequences for amphibian larvae. *Environ. Sci. Technol.* 45 (8), 3781–3787.
- Blukacz-Richards, E.A., Visha, A., Graham, M.L., McGoldrick, D.L., de Solla, S.R., Moore, D.J., Arhonditsis, G.B., 2017. Mercury levels in herring gulls and fish: 42 years of spatiotemporal trends in the Great Lakes. *Chemosphere* 172, 476–487.
- Burger, J., Campbell, K.R., Campbell, T.S., Shukla, T., Jeitner, C., Gochfeld, M., 2005. Use of skin and blood as nonlethal indicators of heavy metal contamination in northern water snakes (*Nerodia sipedon*). *Arch. Environ. Contam. Toxicol.* 49 (2), 232–238.
- Burger, J., Murray, S., Gaines, K.F., Novak, J.M., Punshon, T., Dixon, C., Gochfeld, M., 2006. Element levels in snakes in South Carolina: differences between a control site and exposed site on the Savannah River site. *Environ. Monit. Assess.* 112, 35–52.
- Burger, J., Campbell, K.R., Murray, S., Campbell, T.S., Gaines, K.F., Jeitner, C., Shukla, T., Burke, S., Gochfeld, M., 2007. Metal levels in blood, muscle and liver of water snakes (*Nerodia* spp.) from New Jersey, Tennessee and South Carolina. *Sci. Total Environ.* 373, 556–563.
- Burgess, N.M., Meyer, M.W., 2008. Methylmercury exposure associated with reduced productivity in common loons. *Ecotoxicology* 17 (2), 83–91.
- Bustamante, P., Lahaye, V., Durnez, C., Churlaud, C., Caurant, F., 2006. Total and organic Hg concentrations in cephalopods from the North Eastern Atlantic waters: influence of geographical origin and feeding ecology. *Sci. Total Environ.* 368, 585–596.
- Campbell, K.R., Campbell, T.S., Burger, J., 2005. Heavy metal concentrations in northern water snakes (*Nerodia sipedon*) from East Fork Poplar Creek and the Little River, East Tennessee, USA. *Arch. Environ. Contam. Toxicol.* 49, 239–248.
- Clarkson, T.W., Magos, L., 2006. The toxicology of mercury and its chemical compounds. *Crit. Rev. Toxicol.* 36 (8), 609–662.
- Compeau, G.C., Bartha, R., 1985. Sulfate-reducing bacteria: principal methylators of mercury in anoxic estuarine sediment. *Appl. Environ. Microbiol.* 50 (2), 498–502.
- Depew, D.C., Basu, N., Burgess, N.M., Campbell, L.M., Devün, E.W., Drevnick, P.E., Wiener, J.G., 2012. Toxicity of dietary methylmercury to fish: derivation of ecologically meaningful threshold concentrations. *Environ. Toxicol. Chem.* 31 (7), 1536–1547.
- Depew, D.C., Burgess, N.M., Anderson, M.R., Baker, R., Bhavsar, S.P., Bodaly, R.A., Eckley, C.S., Evans, M.S., Gantner, N., Graydon, J.A., Jacobs, K., LeBlanc, J.E., St. Louis, V.L., Campbell, L.M., 2013. An overview of mercury concentrations in freshwater fish species: a national fish mercury dataset for Canada. *Can. J. Fish. Aquat. Sci.* 70 (3), 436–451.
- Dong, Z., Jim, R.C., Hatley, E.L., Backus, A.S.N., Shine, J.P., Spengler, J.D., Schaider, L.A., 2015. A longitudinal study of mercury exposure associated with consumption of freshwater fish from a reservoir in rural south central USA. *Environ. Res.* 136, 155–162.
- Drewett, D.V.V., Willson, J.D., Cristol, D.A., Chin, S.Y., Hopkins, W.A., 2013. Inter- and intraspecific variation in mercury bioaccumulation by snakes inhabiting a contaminated river floodplain. *Environ. Toxicol. Chem.* 32 (5), 1178–1186.
- Driscoll, C.T., Han, Y.-J., Chen, C.Y., Evers, D.C., Lambert, K.F., Holsen, T.M., Munson, R.K., 2007. Mercury contamination in forest and freshwater ecosystems in the northeastern United States. *Bioscience* 57 (1), 17–28.
- Driscoll, C.T., Mason, R.P., Chan, H.M., Jacob, D.J., Pirrone, N., 2013. Mercury as a global pollutant: sources, pathways, and effects. *Environ. Sci. Technol.* 47 (10), 4967–4983.
- Eagles-Smith, C.A., Ackerman, J.T., Willacker, J.J., Tate, M.T., Lutz, M.A., Fleck, J.A., Stewart, A.R., Wiener, J.G., Evers, D.C., Lepak, J.M., Davis, J.A., Pritz, C.F., 2016. Spatial and temporal patterns of mercury concentrations in freshwater fish across the Western United States and Canada. *Sci. Total Environ.* 568, 1171–1184.
- Eley, B.M., 1997. The future of dental amalgam: a review of the literature. Part 6: possible harmful effects of mercury from dental amalgam. *Br. Dent. J.* 182 (12), 455.
- Fitzgerald, W.F., Lamborg, C.H., Hammerschmidt, C.R., 2007. Marine biogeochemical cycling of mercury. *Chem. Rev.* 107 (2), 641–662.

- Füedner, A., Lohmann, N., Rüdell, H., Teubner, D., Wellmitz, J., Koschorreck, J., 2016. Current levels and trends of selected EU Water Framework Directive priority substances in freshwater fish from the German environmental specimen bank. *Environ. Pollut.* 216, 866–876.
- Futsaeter, G., Wilson, S., 2013. The UNEP global mercury assessment: sources, emissions and transport. *E3S Web of Conferences*. 1, p. 36001.
- Galán, P., 2012. *Natrix maura* en el medio marino de las Islas Atlánticas de Gaúcia. *Boletín de la Asociación Herpetológica Española*. 23, pp. 38–43.
- Gustin, M.S., Evers, D.C., Bank, M.S., Hammerschmidt, C.R., Pierce, A., Basu, N., Blum, J., Bustamante, P., Chen, C., Driscoll, C.T., Horvat, M., Jaffe, D., Pacyna, J., Pirrone, N., Selin, N., 2016. Importance of integration and implementation of emerging and future mercury research into the Minamata convention. *Environ. Sci. Technol.* 50 (6), 2767–2770.
- Hansson, S.V., Sonke, J., Galop, D., Bareille, G., Jean, S., Roux, G., 2017. Transfer of marine mercury to mountain lakes. *Sci. Rep.* 7 (1), 12719.
- Hillman, S.S., Drewes, R.C., Hedrick, M.S., Hancock, T.V., 2014. Physiological vagility and its relationship to dispersal and neutral genetic heterogeneity in vertebrates. *J. Exp. Biol.* 217, 3356–3364.
- Hopkins, W.A., Staub, B.P., Baionno, J.A., Jackson, B.P., Roe, J.H., Ford, N.B., 2004. Trophic and maternal transfer of selenium in brown house snakes (*Lamprophis fuliginosus*). *Ecotoxicol. Environ. Saf.* 58, 285–293.
- Hopkins, B.C., Willson, J.D., Hopkins, W.A., 2013a. Mercury exposure is associated with negative effects on turtle reproduction. *Environ. Sci. Technol.* 47, 2416–2422.
- Hopkins, B.C., Hepner, M.J., Hopkins, W.A., 2013b. Non-destructive techniques for biomonitoring of spatial, temporal, and demographic patterns of mercury bioaccumulation and maternal transfer in turtles. *Environ. Pollut.* 177, 164–170.
- Jackson, A., Evers, D.C., Eagles-Smith, C.A., Ackerman, J.T., Willacker, J.J., Elliott, J.E., Lepak, J.M., Vander Pol, S.S., Bryan, C.E., 2016. Mercury risk to avian piscivores across western United States and Canada. *Sci. Total Environ.* 568, 685–696.
- Jensen, S., Jernelöv, A., 1969. Biological methylation of mercury in aquatic organisms. *Nature* 223 (5207), 753–754.
- Klaper, R., Rees, C.B., Drevnick, P., Weber, D., Sandheinrich, M., Carvan, M.J., 2006. Gene expression changes related to endocrine function and decline in reproduction in fathead minnow (*Pimephales promelas*) after dietary methylmercury exposure. *Environ. Health Perspect.* 114 (9), 1337.
- Lavoie, R.A., Jardine, T.D., Chumchal, M.M., Kidd, K.A., Campbell, L.M., 2013. Biomagnification of mercury in aquatic food webs: a worldwide meta-analysis. *Environ. Sci. Technol.* 47 (23), 13385–13394.
- Lepak, J.M., Hooten, M.B., Eagles-Smith, C.A., Tate, M.T., Lutz, M.A., Ackerman, J.T., et al., 2016. Assessing potential health risks to fish and humans using mercury concentrations in inland fish from across western Canada and the United States. *Sci. Total Environ.* 571, 342–354.
- Mason, R.P., Reinfelder, J.R., Morel, F.M.M., 1995. Bioaccumulation of mercury and methylmercury. *Mercury as a Global Pollutant*, pp. 915–921.
- Mason, R.P., Choi, A.L., Fitzgerald, W.F., Hammerschmidt, C.R., Lamborg, C.H., Soerensen, A.L., Sunderland, E.M., 2012. Mercury biogeochemical cycling in the ocean and poúcy implications. *Environ. Res.* 119, 101–117.
- Meyer, E., Eagles-Smith, C.A., Sparling, D., Blumenshine, S., 2014. Mercury exposure associated with altered plasma thyroid hormones in the declining western pond turtle (*Emys marmorata*) from California Mountain streams. *Environ. Sci. Technol.* 48 (5), 2989–2996.
- Miras, J.A.M., Cheylan, M., Nouria, M.S., Joger, U., Sâ-Sousa, P., Pérez-Mellado, V., Schmidt, B., Meyer, A., Sindaco, R., Romano, A., Martínez-Solano, I., 2015. *Natrix maura*. IUCN Red List of Threatened Species.
- Power, M., Klein, G.M., Guiguer, K.R.R.A., Kwan, M.K.H., 2002. Mercury accumulation in the fish community of a sub-Arctic lake in relation to trophic position and carbon sources. *J. Appl. Ecol.* 39 (5), 819–830.
- Santos, X., Llorente, G.A., 2009. Decúne of a common reptile: case study of the viperine snake *Natrix maura* in a Mediterranean wetland. *Acta Herpetol.* 4 (2), 161–169.
- Scheuhammer, A.M., Sandheinrich, M.B., 2007. Recent advances in the toxicology of methylmercury in wildúfe. *Ecotoxicology* 17 (2), 67–68.
- Scheuhammer, A.M., Basu, N., Burgess, N.M., Elliott, J.E., Campbell, G.D., Wayland, M., Rodrigue, J., 2008. Relationships among mercury, selenium, and neurochemical parameters in common loons (*Gavia immer*) and bald eagles (*Haliaeetus leucocephalus*). *Ecotoxicology* 17 (2), 93–101.
- Scheuhammer, A.M., Braune, B., Chan, H.M., Frouin, H., Krey, A., Letcher, R., Loseto, L., Noël, M., Ostertag, S., Ross, P., Wayland, M., 2014. Recent progress on our understanding of the biological effects of mercury in fish and wildúfe in the Canadian Arctic. *Sci. Total Environ.* 509–510, 91–103.
- Schneider, L., Eggins, S., Maher, W., Vogt, R.C., Krikowa, F., Kinsley, L., Da Silveira, R., 2015. An evaluation of the use of reptile dermal scutes as a non-invasive method to monitor mercury concentrations in the environment. *Chemosphere* 119, 163–170.
- Seún, N.E., 2009. Global biogeochemical cycling of mercury: a review. *Annu. Rev. Environ. Resour.* 34 (1), 43–63.
- Súmani, T., El Hassani, M.S., El Mouden, E.H., Bonnet, M., Bustamante, P., Brischoux, F., Brault-Favrou, M., Bonnet, X., 2017. Large-scale geographic patterns of mercury contamination in Morocco revealed by freshwater turtles. *Environ. Sci. Pollut. Res.* 25 (3), 2350–2360.
- Steuerwald, U., Weihe, P., Jørgensen, P.J., Bjerve, K., Brock, J., Heinzow, B., Grandjean, P., 2000. Maternal seafood diet, methylmercury exposure, and neonatal neurologic function. *J. Pediatr.* 136 (5), 599–605.
- Sullivan, K.M., Kopec, A.D., 2018. Mercury in wintering American black ducks (*Anas rubripes*) downstream from a point-source on the lower Penobscot River, Maine, USA. *Sci. Total Environ.* 612, 1187–1199.
- Tan, S.W., Meiller, J.C., Mahaffey, K.R., 2009. The endocrine effects of mercury in humans and wildlife. *Crit. Rev. Toxicol.* 39, 228–269.
- Tartu, S., Goutte, A., Bustamante, P., Angelier, F., Moe, B., Clément-Chastel, C., Chastel, O., 2013. To breed or not to breed: endocrine response to mercury contamination by an Arctic seabird. *Biol. Lett.* 9 (4), 20130317.
- Todd, B.D., Willson, J.D., Bergeron, C.M., Hopkins, W.A., 2011. Do effects of mercury in larval amphibians persist after metamorphosis? *Ecotoxicology* 21 (1), 87–95.
- Vander Zanden, M.J., Cabana, G., Rasmussen, J.B., 1997. Comparing trophic position of freshwater fish calculated using stable nitrogen isotope ratios ($\delta^{15}\text{N}$) and úterature dietary data. *Can. J. Fish. Aquat. Sci.* 54, 1142–1158.
- Wada, H., Cristol, D.A., McNabb, F.M.A., Hopkins, W.A., 2009. Suppressed adrenocortical responses and thyroid hormone levels in birds near a mercury-contaminated river. *Environ. Sci. Technol.* 43 (15), 6031–6038.
- Whitney, M.C., Cristol, D.A., 2017. Impacts of sublethal mercury exposure on birds: a detailed review. *Rev. Environ. Contam. Toxicol.* 244, 113–163.
- Wolfe, M.F., Schwarzback, S., Sulaiman, R.A., 1998. Effects of mercury on wildúfe: a comprehensive review. *Environ. Toxicol. Chem.* 17, 146–160.
- Żarski, J.F., Skibniewski, M., Skibniewska, E., Żarski, T.P., Majdecka, T., 2017. The presence of mercury in the tissues of mallards (*Anas platyrhynchos* L.) from Włocławek reservoir in Poland. *Biol. Trace Elem. Res.* 176 (2), 384–390.