Changes in French weather pattern seasonal frequencies projected by a CMIP5 ensemble

Pierre Brigode, M. Gérardin, P. Bernardara, J. Gailhard, P. Ribstein

► To cite this version:

Pierre Brigode, M. Gérardin, P. Bernardara, J. Gailhard, P. Ribstein. Changes in French weather pattern seasonal frequencies projected by a CMIP5 ensemble. *International Journal of Climatology*, 2018, 10.1002/joc.5549 . hal-01773657

HAL Id: hal-01773657

<https://hal.science/hal-01773657>

Submitted on 23 Apr 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Changes in French weather pattern seasonal frequencies projected by a CMIP5 ensemble

Brigode P.^{1,2,*}, Gérardin, M.², Bernardara P.¹, Gailhard J.³ and Ribstein P.²

¹ LNHE, R&D, Electricité de France (EDF), Chatou, France.

² UMR 7619 Metis, Sorbonne Université, Paris, France.

³ DTG, DMM, Electricité de France (EDF), Grenoble, France.

* Now in: Université Côte d'Azur, CNRS, OCA, IRD, Géoazur.

Corresponding author: Pierre Brigode (pierre.brigode@unice.fr)

How to cite this article: Brigode P, Gérardin M, Bernardara P, Gailhard J, Ribstein P. Changes in French weather pattern seasonal frequencies projected by a CMIP5 ensemble. *Int J Climatol.* 2018; 1–16. <https://doi.org/10.1002/joc.5549>.

Keywords:

Weather pattern frequency; future weather pattern frequency; CMIP5; French climatology.

Abstract (up to 300 words):

Over the last decades, General Circulation Model (GCM) simulations have been regularly evaluated in terms of their ability to reproduce the historical frequency of significantly recurrent Weather Patterns (WP) observed at the regional scale. Thus, a good simulation of the frequency of these particular WP by the GCM is generally conditioning the good representation of the regional statistics of surface variables such as temperature and precipitation. In this paper, the seasonal frequency of eight particular WP have been calculated using the daily geopotential height fields simulated by an ensemble of 26 CMIP5 GCM. These WP are known as significantly influencing the French regional hydro-climatology in terms of both frequency of low flows and high precipitation events. Four different bias correction methods have been applied on the simulated geopotential height fields before the calculation of the seasonal WP frequencies. The GCM ensemble showed overall good performances in terms of the simulation of WP seasonal frequencies. The application of a spatially and temporally nonhomogenous correction of simulated geopotential height fields improved significantly the simulation of WP frequencies for the four seasons. Finally, the evolution of the WP frequencies over the next century has been quantified. Three WP (WP2, WP4 and WP8) have pronounced seasonal changes, with WP2 and WP4 being less frequent in summer and autumn seasons, respectively, while WP8 being more frequent over spring, summer and autumn seasons. The strong simulated frequency evolution of WP2 and WP8 is an interesting result, which predicts the climate to be drier with time for France. Thus, WP2 (western oceanic circulation), grouping rainy days over the northern France region, is simulated as less frequent in future summers, while WP8 (Anticyclonic situations), which groups non-rainy days over France, is simulated as more frequent in future summers.

1 INTRODUCTION

Since several decades, meteorologists and climatologists have worked on the identification of particular atmospheric circulations patterns that are recurrently observed at the regional scale and that are strongly affecting the spatial and the temporal variability of surface variables, such as temperature and precipitation (e.g. over Europe by Plaut & Simonnet, 2001 ; Cassou *et al.*, 2005 ; Boé & Terray 2008). Different classification methods are used for identifying a limited number of significantly recurrent circulation patterns (see Huth *et al.*, 2008 for a review), generally based on the analysis of atmospheric circulation variables such as Sea Level Pressure (SLP) or geopotential heights (Vautard, 1990; Michelangeli *et al.*, 1995). In the literature, the obtained recurrent circulations patterns have different names and are usually referenced as (weather, atmospheric, climate or circulation) regimes, types or patterns, based on their properties (Stephenson *et al.*, 2004). Philipp *et al.* (2007) opposed the weather regimes (WR) to the weather patterns (WP) in terms of their spatial and temporal characteristics. The WR are usually one to four “quasi-stationary states of the large-scale circulation system” defined at the 10-day timescale, while the WP are more numerous situations defined at a finer spatial scale since being devoted to the classification of the circulation system on a daily timescale. The utility of WP classifications for quantifying the influence of large-scale climate drivers on hydro-climatological variability has been discussed by several studies, mainly arguing that WP are defined at an intermediate spatial scale between WR and local climatic observations (e.g. Giuntoli *et al.*, 2013 ; Brigode *et al.*, 2013a, 2013b).

Recently, WP frequencies have thus been used as a conditioning variable for the statistical modelling of extreme hydro-climatological values (e.g. Planchon *et al.* 2009; Brigode *et al.*, 2013b). For example, several studies successfully linked WP (or circulation-related variables) to flood statistics at the regional scale (e.g. Nied *et al.*, 2014; Wilby & Quinn, 2013; Renard & Lall, 2014). In the framework of the statistical modelling of extreme precipitation, Garavaglia *et al.* (2010, 2011) proposed an eight WP classification for sub-sampling precipitation events over France in terms of their synoptic origin. Trambly *et al.* (2011, 2013) used the same WP classification as conditioning variables in a non-stationary extreme value frequency analysis and showed the soundness of using this framework for the statistical modelling of extreme precipitation in southern French regions. Furthermore, Giuntoli *et al.* (2013) found significant correlations between these WP and low flows over France, highlighting again the strong link between these WP and the seasonal French hydro-climatology.

These significant links are of particular interest in the context of the quantification of regional climate change impacts. Under the (strong) hypothesis that the statistical relationships observed over the historical period hold unchanged under climate changes, WP classifications can be used to downscale and transform changes of atmospheric circulations into changes of regional surface variables (Boé & Terray, 2008). Nevertheless, this approach is based on the other hypothesis that atmospheric circulations are better simulated by the Global Circulation Models (GCM) than surface variables such as precipitation. At the global scale, numerous studies showed that large-scale features of the atmospheric circulation are among the variables that GCM represent with the best robustness (e.g. Covey *et al.*, 2003; Räisänen, 2007), even though the ability of GCM to simulate changes in atmospheric circulation is recently questioned (e.g. Palmer, 1999; Corti *et al.*, 1999; Hsu & Zwiers, 2001; Gillett *et al.*, 2003; Shepherd, 2014; Hall, 2014; Murawski *et al.*, 2016).

WR and WP classifications have been widely used as a way to evaluate GCM outputs in terms of their ability to reproduce WR and WP frequencies over a target region (Sheridan & Lee, 2010). Frequencies of simulated WP have been evaluated over different regions, such as the Euro-Atlantic region (e.g. Demuzere *et al.*, 2009; Handorf & Dethloff, 2012 and Pastor & Casado, 2012), the Arctic region (e.g. Cassano *et al.*, 2006), the northwestern Iberian Peninsula (e.g. Lorenzo *et al.*, 2011) and the Northwest America region (e.g. McKendry *et al.*, 2006). Anagnostopoulou *et al.* (2008) used the 500 hPa geopotential height (Z_{500}) and the 1000-500 hPa thickness fields for WP classification over the eastern Mediterranean region and revealed several differences between observed WP frequencies and the frequencies simulated by the considered GCM (HadAM3P), with, for example, an overestimation of the summer frequency of anticyclonic situations. Sanchez-Gomez *et al.* (2009) evaluated the performances of an ensemble of 13 Regional Climate Models (RCM) forced by a reanalysis (ERA40) in terms of the simulation of WR over the Euro-Atlantic region and highlighted the good performances of the ensemble in terms of the simulation of the mean WR frequency. During the last years, the GCM outputs of the fifth Coupled Model Intercomparison Project (CMIP5, Taylor *et al.* (2012)) have been evaluated with the same methodology over different regions (e.g. by Belleflamme *et al.* (2013) over Greenland and by Dunn-Sigouin & Son (2013); Hertig & Jacobeit (2014) and Ullmann *et al.* (2014) over the Northern hemisphere), and the CMIP5 GCM appear to represent generally well the observed WR. Nevertheless, Belleflamme *et al.* (2014) and Santos *et al.* (2016) illustrated the limitations of two different GCM ensembles in simulating seasonal WP frequency over Europe for the recent decades. The ability of GCM to simulate WR and WP frequency seem thus to depend on the GCM considered, on the spatial scale considered and on the WR and WP classification method used.

The general aim of the paper is to quantify the frequency changes of the eight French WP previously defined by Garavaglia *et al.* (2010) simulated by an ensemble of 26 CMIP5 GCM over the 21st century. Before studying future WP frequency changes, the performances of the GCM ensemble will be quantified over the historical period (here 1979-2004), using four different bias correction method applied on the GCM geopotential heights fields.

2 DATASETS

2.1 Reanalysis of geopotential height fields

The ERA-Interim global atmospheric reanalysis (Dee *et al.*, 2011, noted ERAi hereafter) has been used as reference for geopotential height fields for the historical period. This gridded dataset is available from 1979, at a 0.75° spatial resolution and at a 12-hour temporal resolution. Two levels have been considered in this study, namely (i) the geopotential height field at 1000 hPa (noted Z_{1000} hereafter) and (ii) the geopotential height field at 700 hPa (noted Z_{700} hereafter), both at 0h for each day. The historical period considered in this study is 1979-2004.

2.2 GCM geopotential height field outputs

Daily Z_{1000} and Z_{700} outputs from 26 GCM have been considered in this study. All outputs have been generated within the fifth phase of the Coupled Model Intercomparison Project (CMIP5, Taylor *et al.*, 2012). The GCM studied are listed in table S1. The daily outputs of three different experiments have been used:

- The “historical” experiment, which consists of a simulation of the recent past (1950-2005 period) under the historical forcing;
- The “RCP4.5” experiment, which extends the “historical” experiment by simulating the future (2006-2100) under the RCP4.5 emission scenario (radiative forcing surplus of 4.5 W.m⁻² in 2100).
- The “RCP8.5” experiment, which extends the “historical” experiment by simulating the future (2006-2100) under the RCP8.5 emission scenario (radiative forcing surplus of 8.5 W.m⁻² in 2100).

The two emission scenarios considered here are members of the four Representative Concentration Pathway (RCP) scenarios considered in the CMIP5 experiments (Moss *et al.*, 2010).

Only the first member (named r1ip1) from the historical, RCP4.5 and RCP8.5 experiments have been considered for each GCM. The simulated Z_{1000} and Z_{700} fields have been spatially interpolated on the ERAi grid (0.75° grid spacing). The historical experiment outputs are considered over the 1979-2004 period (period of availability of the ERAi dataset) and the future experiment outputs are considered over the 2006-2098 period.

Several GCM outputs being produced by climate models with structural similarities (e.g. the bcc-csm1-1 outputs and the bcc-csm1-1-m outputs), a sub-sample of the 26 outputs has also been considered. Thus, the seasonal WP frequencies simulated using outputs produced by similar climate models (i.e. developed by the same institution) were averaged. The Table S1 presents the merging considered for the constitution of the second GCM ensemble (noted INS ensemble hereafter, where INS stands for “institution”), composed by 15 GCM outputs out of the 26 available.

2.3 Precipitation reanalysis

As an illustration of the regional precipitation patterns associated to each WP, the E-OBS daily precipitation reanalysis (Haylock *et al.*, 2008, version 16.0) is used in this study. This reanalysis provides daily precipitation amounts at a 0.25° resolution. This dataset has been used over Western Europe and the 1979-2004 period.

3 METHODOLOGY

This section aims at describing the WP classification considered, the bias correction methods used and the frequency analysis methodology, all performed in the R-project environment (R Core Team, 2016, <http://www.r-project.org/>).

3.1 The WP classification method

3.1.1 Average synoptic situation of the eight French WP

The WP considered in the paper are the eight French WP defined by Garavaglia *et al.* (2010). These WPs have been defined at the daily time step through a classification of “rainy days” over southern France for the 1956-1996 period (see Garavaglia *et al.*, 2010 and Brigode *et al.*, 2013b for more details on the rainy days classification method). For example, the first WP (WP1) is constituted by 668 rainy days of the 1956-1996 period.

The synoptic characteristics of each of the eight WP have then been calculated by averaging the synoptic situations of the different days constituting each WP. Thus, the WP1 synoptic situation is the average of the synoptic situations of the 668 rainy days previously identified as constituting the WP1. Four geopotential height fields are considered for the synoptic description of each day and thus of each WP:

1. the Z_{1000} field at 0h,
2. the Z_{1000} field at 24h,
3. the Z_{700} field at 0h,
4. the Z_{700} field at 24h.

In this study, the average synoptic situation of each WP has been estimated with the ERAi geopotential reanalysis (available from 1979, see section 2) and thus computed over the 1979-1996 period. Geopotential height fields have been considered over 504 ERAi grid points, centered over southern France (-7.50° to 12.75° of longitude and 37.50° to 50.25° of latitude). Figure 1 shows the average Z_{700} field at 24h of each WP, estimated over the 1979-1996 period, and the spatial coverage of the 504 ERAi grid points.

3.1.2 Classification of the historical period

In order to obtain a daily classification for the studied historical period, each past day is assigned to one of the eight WP by computing distances between the synoptic situation of the studied day and each WP. Teweles & Wobus (1954) distances (noted D_{TW} hereafter) are calculated to find the closest WP to each considered day (see Brigode *et al.*, 2013a, 2013b for more details). D_{TW} focuses on air flows since this distance involves gradients of absolute values of geopotential heights (Woodcock, 1980; Obled *et al.*, 2002; Bontron, 2004; Wetterhall *et al.*, 2005, Brigode *et al.*, 2016). The formula for estimating D_{TW} between two synoptic situations (Z1) and (Z2) characterized by geopotential height fields on a gridded domain oriented south-north (index i) and west-east (index j) is :

$$D_{TW} = \frac{\sum_{i,j} |e_G^i| + \sum_{i,j} |e_G^j|}{\sum_{i,j} |G_L^i| + \sum_{i,j} |G_L^j|} \times 100$$

where e_G is the difference, around a given grid point, between the Z1 and Z2 geopotential gradients, while G_L is the maximum of these two gradients in the direction considered (i or j):

$$|e_G^i| = |(Z1_{i,j} - Z1_{i+1,1}) - (Z2_{i,j} - Z2_{i+1,1})|$$

$$|e_G^j| = |(Z1_{i,j} - Z1_{i,j+1}) - (Z2_{i,j} - Z2_{i,j+1})|$$

$$|G_L^i| = \max(|Z1_{i,j} - Z1_{i+1,1}|, |Z2_{i,j} - Z2_{i+1,1}|)$$

$$|G_L^j| = \max(|Z1_{i,j} - Z1_{i,j+1}|, |Z2_{i,j} - Z2_{i,j+1}|)$$

D_{TW} ranges from 0 for two identical geopotential fields and 200 for two opposite geopotential fields. The final score between one day and one WP is the sum of four D_{TW} distances: (i) the D_{TW} between the Z_{700} fields at 0h, (ii) the D_{TW} between the Z_{1000} fields at 24h, (iii) the D_{TW} between the Z_{700} fields at 0h and (iv) the D_{TW} between the Z_{700} fields at 24h. The D_{TW} calculation is illustrated in figure 1, which shows the eight D_{TW} calculated between one particular day (21th September 1992) and the average synoptic situation of the eight French WPs for the Z_{700} fields at 24h. The minimal D_{TW} is found for the WP4 ($D_{TW}=48.4$). The three other calculations of D_{TW} for that particular day (Z_{1000} at 0h, Z_{1000} at 24h and Z_{700} at 0h) are summarized in the table 1. The sum of the four D_{TW} is minimal for the WP4 ($D_{TW}=205.5$): the 21th September 1992 is thus attributed to the WP4. Note that this synoptic situation has generated a considerable amount of precipitation in South-Eastern France, with more than 300 mm of precipitation observed locally that day (Benech *et al.*, 1993), generating a catastrophic and deadly flash flood of the Ouvèze river (Sénési *et al.*, 1996).

Table 1: D_{TW} distances calculated between the 21th September 1992 synoptic situation and the average synoptic situation of the eight French WPs for the Z_{1000} and Z_{700} fields at 0h and at 24h. In this case, the 21th September 1992 is finally attributed to the WP4.

	WP1	WP2	WP3	WP4	WP5	WP6	WP7	WP8
Z_{1000} at 0h	91.9	101.1	85.3	67.9	95.2	81.1	70.7	96.8
Z_{1000} at 24h	92.0	91.7	67.6	52.2	89.6	84.1	64.2	99.6
Z_{700} at 0h	62.5	73.3	49.7	36.9	63.4	63.6	46.1	70.4
Z_{700} at 24h	74.7	82.1	59.0	48.4	79.8	71.4	56.8	86.6
D_{TW} sums	321.1	348.2	261.6	205.5	328.1	300.3	237.7	353.4

Figure 1: Comparison between the Z_{700} synoptic situation of the 21th September 1992 (at 24h, panel I) and the Z_{700} situation of the eight French WP (panels A to H). The D_{TW} distances calculated between these geopotential fields are given on each panel.

3.1.3 Meteorological description of the eight French WP

The eight French WPs are described in the figures 2.1 and 2.2, in terms of average ERAi Z_{700} and Z_{1000} fields at 0h, E-OBS regional precipitation pattern and seasonal frequencies, calculated over the 1979-2004 period. The eight WP can be grouped in terms of their general atmospheric flow direction (cf. column (a) of the figures 2.1 and 2.2, showing average SLP for each WP):

- WP1 (Atlantic Wave), WP2 (Steady Oceanic) and WP3 (Southwest Circulation) correspond to westerly oceanic circulation, grouping days particularly rainy over the Alps, the Northwestern part of France and the Western part of France, respectively. The WP2 is one of the most frequent over the year, especially during the winter season.
- WP4 (South Circulation), WP6 (East Return) and WP7 (Central Depression) correspond to Mediterranean circulations, grouping days particularly rainy over the Southeastern part of France. WP6 and WP7 days are relatively rarely observed.
- The WP5 (Northeast Circulation) corresponds to continental circulations, grouping days not particularly rainy over France.

- Finally, the WP8 (Anticyclonic) corresponds to high pressure situations and thus groups non-rainy days. This French WP is the most frequent, especially in summer.

Figure 2.1: Description of the WP1 to WP4: (a) average ERAi SLP at 0h, (b) average ERAi Z_{700} at 0h, (c) average ERAi Z_{1000} at 0h, (d) ratio between the E-OBS mean precipitation amounts and the general precipitation amount (considering all WP) and (e) seasonal frequency of the WP, estimated over the 1979-2004 period.

Figure 2.2: Description of the WP5 to WP8: (a) average ERAi SLP at 0h, (b) average ERAi Z₇₀₀ at 0h, (c) average ERAi Z₁₀₀₀ at 0h, (d) ratio between the E-OBS mean precipitation amounts and the general precipitation amount (considering all WP) and (e) seasonal frequency of the WP, estimated over the 1979-2004 period.

3.2 Classification of the daily GCM outputs

The classification of the geopotential height fields simulated by the GCM has been done with the same methodology, i.e. by calculating D_{TW} between each simulated day (characterized by four synoptic situations: Z₁₀₀₀ at 0h, Z₁₀₀₀ at 24h, Z₇₀₀ at 0h and Z₇₀₀ at 24h) and the ERAi average situations of the eight French WP (also being characterized by four synoptic situations). The sum of the four D_{TW} is calculated for each simulated day and each WP, and the WP with the minimal D_{TW} sum is attributed to the studied day.

3.3 Geopotential height bias correction methods

While most of the studies on the GCM WP simulation used uncorrected GCM outputs, it is noteworthy that Demuzere *et al.* (2009) and Lorenzo *et al.* (2011) found better performances of GCM in terms of frequencies if the SLP fields used for the classification were bias-corrected before the classification procedure. Thus, the need for bias correction of GCM geopotential height fields before performing WP classification will be tested in this paper, by considering different bias correction methods.

D_{TW} distance values have been firstly computed without correcting the GCM geopotential height fields. The WP frequencies obtained by using these uncorrected GCM outputs are named D0 hereafter. Then, four different bias correction methods have been applied to the GCM outputs:

1. A spatially homogeneous correction of the geopotential height average values and standard deviations. The outputs of this bias correction method are named D1 hereafter. Note that since D_{TW} are estimated by considering synoptic circulation gradients, a spatially homogeneous correction of average values only is useless: lowering or rising the mean geopotential height fields has no effect on the D_{TW} values.
2. A spatially nonhomogeneous correction of the average values. The outputs of this bias correction method are named D2 hereafter.
3. A spatially nonhomogeneous correction of the geopotential height average values and standard deviations. The outputs of this bias correction method are named D3 hereafter.
4. A spatially nonhomogeneous correction of the monthly average values and standard deviations. The outputs of this bias correction method are named D4 hereafter.

To summarize, each GCM output is considered five times: firstly without any bias correction method (outputs named D0) and then after application of the D1, D2, D3 and D4 bias correction methods.

3.4 Seasonal WP frequencies and frequency variability

Seasonal frequencies of the eight French WP have been estimated over the 25-year historical period (01/03/1979-29/02/2004) and over 68 25-year periods extracted from the future RCP simulations (2006-2031, 2007-2032, ..., 2073-2098). Four 3-month seasons have been defined: the autumn season (September, October and November, noted SON hereafter), the winter season (December, January and February, noted DJF hereafter) the spring season (March, April and May, noted MAM hereafter) and the summer season (June, July and August, noted JJA hereafter). For each season and each WP, frequencies are defined as the percentage of days belonging to the considered WP. The reference observed seasonal frequencies of the eight WP are the seasonal frequencies represented in column (e) of figures 2.1 and 2.2.

In order to quantify the WP frequency variability within a time period, a non-parametric bootstrap resampling has been performed. Thus, for each 25-year time period considered, 100 samples of 15 randomly chosen years are constructed. Note that no replacements are allowed within this bootstrap resampling, and thus one particular year cannot be resampled twice. These 100 samples are then used to quantify the variability of the frequency by measuring the 90% confidence interval. The dispersion of observed WP frequencies are shown in figures 2.1 and 2.2 for the 1979-2004 period. The observed variability of the seasonal frequencies is limited, the highest variability is observed for the WP2, WP4 and WP8. For example, the summer frequency of WP8 ranges between 33.2% and 38.8%.

4 RESULTS

4.1 GCM simulations of WP average Z_{1000} and Z_{700}

The GCM ensemble outputs have been firstly evaluated in terms of the simulation of the average Z_{700} and Z_{1000} fields for the eight WP, over the 1979-2004 period. Figure 3 summarizes the spatial variability of the Z_{1000} and Z_{700} bias, calculated for each WP as the difference between the GCM ensemble mean fields and the average ERAi fields (presented in the figures 2.1 and 2.2). For Z_{700} , the GCM ensemble appears to overestimate the geopotential heights for the eight WP. The spatial distributions of these biases reveal a slighter bias over northwestern France (Atlantic Sea). The spatial distribution of Z_{1000} bias changes with WP. For WP1, WP5 and WP7, the GCM ensemble tends to underestimate the Z_{1000} values over northwestern France and to overestimate the Z_{1000} values over the southeastern part of the studied region. For the WP6, the Z_{1000} values are slightly underestimated over the entire studied region. For the other WP (WP2, WP3, WP4 and WP6), the GCM ensemble tends generally to overestimate the Z_{1000} values over the studied region. The spatial variability of biases justified the use of bias correction method that are spatially (D2 to D4) nonhomogeneous.

Figure 3: Z_{700} and Z_{1000} geopotential bias calculated for each WP as the difference between GCM ensemble mean and ERAi geopotential height fields.

4.2 Historical seasonal WP frequencies simulated by the GCM ensemble

The GCM ensemble has then been evaluated in terms of simulating historical WP frequencies for the four different seasons considered. Figure 4 presents the observed (ERAi) seasonal WP frequencies estimated over the 1979–2004 period and the WP frequencies simulated by the GCM ensemble (uncorrected and corrected). The seasonal variability of the WP frequencies are generally well simulated by the uncorrected GCM (except for the WP4). Nevertheless, the frequencies of the two most frequent WP (WP2 and WP8) are poorly simulated, with WP2 frequencies being strongly overestimated and WP8 strongly underestimated. Moreover, the dispersion of the GCM ensemble is large for these two WP as well as for the WP4. The impact of the different bias correction methods on the historical WP frequencies is not straightforward and needs further investigations.

Figure 4: Seasonal WP frequencies estimated over the 1979-2004 period. The grey rectangles present the observed (ERAi) WP frequencies. The boxplots are constructed with the WP frequencies simulated by the GCM ensembles (GCM and INS) without bias correction (blue boxplots) and with bias corrections (white boxplots, D1 to D4).

Figure 5 presents the distribution of Euclidean distances calculated, for each season and each bias correction method, between the vector of the 8 observed WP frequencies and the 26 vectors of 8 simulated WP frequencies. When no bias correction is applied (D0 method), the GCM ensemble is slightly less performant for DJF season (average distance around 19%) than for other seasons (average distance around 15.5%). The application of the D1 bias correction method (spatially homogeneous correction of the geopotential height standard deviations) appears to degrade the performance of the GCM ensemble in terms of seasonal WP frequencies. Nevertheless, the application of the D2 to D4 correction methods (all spatially inhomogeneous correction methods) improves the GCM ensemble performances for the DJF, MAM and SON seasons. For the JJA season, only the D4 bias correction

method improves the performance of the WP frequency simulation. Overall, the D4 bias correction method (only method implying a spatial and temporal nonhomogenous correction) has the best performances in terms of simulation of the ERAi WP frequencies on the historical period.

Figure 5: Distributions of seasonal Euclidean distances calculated, for each season and each bias correction method, between the vector of 8 observed WP frequencies and the 26 vectors of 8 simulated (GCM) WP frequencies over the 1979-2004 period. The blue boxplots are WP frequencies obtained without bias correction, and the white ones are obtained when the bias correction D1 to D4 are applied. Red values are mean values of Euclidean distances between WP frequency vectors and grey points are individual GCM distances.

4.3 Future evolutions of seasonal WP frequencies

In this section, the changes of WP frequencies simulated by the GCM ensemble are calculated. Regarding the performances obtained by the four different bias correction methods tested (see Section 4.2), only the best method (D4) has been considered for the estimation of future WP frequencies. Thus, future frequencies are calculated with both uncorrected GCM outputs (D0) and GCM outputs corrected with the D4 method.

Figure 6 presents the mean seasonal frequency evolutions simulated by the GCM ensemble, considering both RCP4.5 and RCP8.5 emission scenarios and both GCM outputs without bias correction and GCM outputs corrected with the D4 method. None particular temporal evolutions throughout seasons are simulated for WP1, WP5, WP6 and WP7. WP3 and WP4 appear to be slightly less frequent at the end of the century during JJA and SON seasons, respectively. These slight decreases seem to be more pronounced when considering the RCP8.5 simulations. WP2 and WP8 are the two WP with the most pronounced seasonal changes and are in opposition. Overall, WP2 is slightly less frequent (but its frequency evolution depends on the RCP outputs considered), while WP8 is highly more frequent. These conjoint evolutions are particularly notable for the JJA season. For this season, WP8 frequency is constantly increasing with time for RCP8.5 outputs, while the frequency increase is stopped around the 2060 years for RCP4.5. Oppositely, the WP2 frequency decrease for the same season is stronger when considering RCP8.5 outputs. For the SON season, a strong increase of WP8 is also notable, while a slight decrease of WP4 and WP2 frequencies is found (except for uncorrected RCP8.5 outputs). For the DJF and MAM seasons, the WP8 appears to be also more frequent, while no particular frequency evolution is found for WP2.

Figure 6: Seasonal WP frequency changes (in %) estimated with the GCM ensemble over 68 consecutive 25-year periods, considering RCP4.5 and RCP8.5 simulations, with no bias correction (D0) and with the D4 bias correction method. Changes are relative to the WP frequencies simulated by the GCM ensemble over the first 25-year period (2006-2031). Grey lines are individual GCM WP frequency changes.

Figure 7 summarizes the frequency changes calculated between the 2006-2031 period and the 2073-2098 period. A Student's t-Test has been performed for highlighting the significant changes of mean seasonal frequency (p -value < 5%). The dispersion of simulated changes is different depending on the WP, with larger dispersion for WP2, WP4 and WP8 frequencies. WP1 frequency appears to only slightly decrease in JJA months for the RCP8.5 simulations (-1.2%, 95% of the GCM ensemble simulating a decrease), when geopotential height bias are corrected with the D4 method. For WP2, the simulated changes are highly dependent on the season and on the application of a bias correction method. After bias correction, no clear frequency changes are obtained when considering RCP4.5 outputs. Nevertheless, the RCP8.5 outputs show a significant decrease of WP2 frequency for the JJA season with the bias correction method applied (-2.7%, 90% of the GCM ensemble simulating a decrease). WP3 appears to be slightly more frequent for DJF (significant increase of +1.3%, 75% of the GCM ensemble simulating an increase, with D4 correction method), while being less frequent for JJA season (significant decrease of -1.8%, 81% of the GCM ensemble simulating a decrease, with D4 correction method), for the RCP8.5 simulations. The summer decrease of the WP3 frequency is also found in the bias-corrected RCP4.5 simulations (significant decrease of -0.7%, 58% of the GCM ensemble simulating a decrease). WP4 are significantly less frequent in JJA season RCP8.5 (-1.2%, 81% of the GCM ensemble simulating a decrease) scenarios and in SON season for bias corrected RCP8.5 (-2.2%) scenarios. A significant decrease of -1.8% in the DJF season is found for WP5 for the bias corrected RCP8.5 simulations (decrease simulated by 81% of the GCM ensemble), while no significant changes are found for the other seasons and outputs. The bias corrected RCP8.5 outputs show significant less WP6 days for the different seasons, with a stronger decrease for the SON season (-1.5%, 95% of the GCM ensemble simulating a decrease). The WP7 is slightly less frequent for MAM, JJA and SON seasons, especially when considering RCP8.5 and bias-corrected outputs (-1.1%, -0.4% and -1.0% for these three seasons, decreases simulated by 81%, 67% and 81% of the GCM ensemble, respectively). Finally, the WP8 frequency is significantly increasing for MAM, JJA and SON seasons. These increases are stronger when considering bias corrected RCP8.5 outputs (+3.3%, +7.4% and +4.8%, increases simulated by 81%, 95% and 90% of the GCM ensemble respectively).

The summer (JJA) Z_{700} values simulated by the GCM ensemble is increasing with time over the studied domain. The GCM ensemble mean is 3137 m around 2018 and is 3183 m around 2085, considering the RCP8.5 scenario. This increase seems to be correlated with the decrease of the WP2 summer frequency and with the increase of the WP8 summer frequency (results not shown). Nevertheless, the WP classification was performed using the Teweles & Wobus (1954) distance (D_{TW}), metric calculated with the geopotential height gradients and not with the absolute values of the geopotential heights. Thus, the temporal increase of the geopotential height absolute values is not influencing the D_{TW} calculation and the simulated increase (decrease) of the seasonal WP8 (WP2) frequency are thus due to changes in general circulation over the studied domain and not due to the global increase of surface pressure in future climate.

Figure 7: Seasonal WP frequency changes estimated with the GCM ensemble between 2073-2098 period and 2006-2031 period, considering RCP4.5 and RCP8.5 simulations, with no bias correction (D0) and with the D4 bias correction. The significant changes of mean WP frequency are printed in red color and expressed as percentage of frequency change.

5 DISCUSSION AND CONCLUSION

An ensemble of 26 GCM used within the fifth Coupled Model Intercomparison Project (CMIP5) have been analyzed in order firstly to test its ability to reproduce observed seasonal geopotential height climatology over Western Europe, secondly to test its ability to reproduce the seasonal frequencies of eight French WPs previously defined (Garavaglia *et al.*, 2010) and finally to estimate the future seasonal WP frequencies.

Firstly, biases of simulated Z_{1000} and Z_{700} have been quantified over Western Europe for the historical period (here 1979-2004), relatively to the ERAi reanalysis. For Z_{1000} , the GCM ensemble biases have both a strong spatial variability and seasonal variability. For example, the GCM ensemble tends to underestimate the geopotential heights at the highest latitudes of the Western Europe in DJF and MAM seasons, while it tends to overestimate geopotential heights at the same latitudes for the JJA and SON seasons. For Z_{700} , the GCM ensemble tends to overestimate the geopotential heights for the four seasons, with a slighter bias over northern latitudes. These spatial and seasonal variabilities of geopotential biases could advocate the use of spatially and seasonally inhomogeneous bias correction methods on the geopotential height outputs.

Secondly, the ability of the GCM ensemble to reproduce historical WP frequencies has been quantified and revealed that the seasonal variability of the WP frequencies are generally well simulated by the uncorrected GCM ensemble (except for the WP4), with a slightly worse performance obtained for the DJF season. Nevertheless, the frequencies of the two most frequent WP (WP2 and WP8) are poorly simulated by the ensemble, with WP2 frequencies being strongly overestimated and WP8 strongly underestimated. Similar results have been obtained by Santos *et al.* (2016) by looking at WP frequencies simulated by an ensemble of 22 GCM over Western Europe and showing the overestimation by this other ensemble of the frequency of WP associated with zonal airflow. The use of four different bias correction methods showed that the application of a spatially and temporally nonhomogeneous correction of geopotential height fields (here the correction named D4) improved significantly the simulation of WP frequencies for the four seasons.

Finally, the evolution of the WP frequencies over the next century has been quantified, considering two emission scenarios (RCP4.5 and RCP8.5) and considering one bias correction method (D4) and no bias correction. The WP2, WP4 and WP8 have more pronounced seasonal changes, with WP2 and WP4 being less frequent in JJA and SON seasons, respectively, while WP8 being more frequent over MAM, JJA and SON season. The frequency changes calculated are higher for RCP8.5 simulations than for RCP4.5. Moreover, the temporal evolution of the WP frequencies appears to be constant over time for RCP8.5, while the evolution stops around the year 2060 for the RCP4.5 scenario. The use of a bias correction method is important in this context, since the significant mean changes of WP frequencies are all obtained with bias-corrected outputs (except the increase of WP8 summer frequency). Nevertheless, the analysis of the temporal evolution of WP frequencies (figure 6) and the distribution of simulated frequency changes (figure 7) showed that the bias-correction method used (D4) is not changing the sign of the mean frequency changes compared to the uncorrected GCM outputs. Note that the use of three other bias correction methods (D1 to D3) leads to the same change signs for the different WP and seasons (results not shown).

The strong simulated frequency evolution of WP2 and WP8 is an interesting result, which predicts the climate to be drier with time for France. Thus, WP2 (western oceanic circulation), grouping rainy days over the northern France region, is simulated as less frequent in future summers, while WP8 (Anticyclonic situations), which groups non-rainy days over France, is simulated as more frequent in future summers. These evolutions could have significant impacts on French low flows, since Giuntoli *et al.* (2013) recently highlighted strong correlations between these WP frequency and drought severity over France. An increase of the frequency of non-rainy WP is also found in the 22 GCM ensemble studied by Santos *et al.* (2016) when considering RCP8.5 outputs.

The bias of CMIP5 GCM in terms of the simulation of the eight French WP frequencies rises several questions, firstly about the WP classification methodology considered here. Thus, Vrac *et al.* (2007) showed that the WP classification method used has an impact on the identified patterns, and also that the choice of a given reanalysis as reference could lead to different WP classifications. Applying different classification methods on the same GCM ensemble and considering other reanalysis as reference would be an interesting perspective, in order to see if similar French WP are identified. In addition, the applied methodology assumes that only the WP frequencies are changing in the future, while WP structures are considered as constant in time. Küttel *et al.* (2010) thus highlighted large changes within type variations for European WP. The application of the methodology developed by Cattiaux *et al.* (2012) could be an interesting perspective in order to fully split the part of changes explained by WP frequencies and the part explained by WP structures, for example.

The second question to be raised is the spatial domain considered here for the definition of the WP classification and, consequently, for studying WP frequencies simulated by the GCM ensemble. This domain is rather small (-7.50° to 12.75° of longitude and 37.50° to 50.25° of latitude), thus concerning only a limited grid points for the GCM characterized by a limited atmospheric horizontal resolution. If the WP identified at this spatial scale have relevant characteristics in terms of spatial distribution of surface variables such as precipitation over France, it is possible that the horizontal resolution of GCM are too large for allowing them to identify such regional patterns. Nuissier *et al.* (2011), with a rather similar objective of determining WP leading to heavy precipitation events over southern France, used a larger spatial domain for the WP classification (-24° to $+39^{\circ}$ of longitude and 25.5° to 63° for latitude). Thus, trying to define French WP over a larger spatial domain is an interesting perspective, in order to quantify the future WP frequencies. The use of Regional Climate Model (RCM) could also be an interesting perspective, even though recent work questioned the ability of RCM to reproduce the daily weather regimes (e.g. Lucas-Picher *et al.*, 2016).

Finally, the use of spatially nonhomogeneous bias correction methods for geopotential height fields is questionable, since these corrections change the simulated circulation patterns, transforming the GCM outputs in order to be more similar to the reference ones. Moreover, the biases quantified over the historical period have to be assumed as being stationary over time in order to be applied over the future time period considered, which is a strong hypothesis.

The ultimate goal of this work was to use simulated future frequencies of particular WPs - known to potentially leading to heavy precipitation events (and then extreme floods) - in order to discuss future precipitation (floods) frequency. Within this framework, only the frequency of future rainfall events is studied, thus splitting frequency (related to circulation dynamic) and intensity (related to thermo-dynamic parameters) of such events. Nevertheless, recent studies highlighted that both

frequency and intensity have to be studied for discussing future extreme precipitation (e.g. Hertig *et al.*, 2013; Blenkinsop *et al.*, 2015) or climate extreme event attribution (e.g. Trenberth *et al.*, 2015).

Finally, another interesting future work would be to study the potential shift of WP persistence over the same region and with the same GCM ensemble, since the persistence of rainy or non-rainy WPs is a good indicator for flood or drought frequency and intensity.

6 ACKNOWLEDGEMENTS

We acknowledge the World Climate Research Programme's Working Group on Coupled Modelling, which is responsible for CMIP, and we thank the climate modeling groups (listed in table S1 of this paper) for producing and making available their model output. For CMIP the U.S. Department of Energy's Program for Climate Model Diagnosis and Intercomparison provides coordinating support and led development of software infrastructure in partnership with the Global Organization for Earth System Science Portals. We also acknowledge the E-OBS dataset from the EU-FP6 project ENSEMBLES (<http://ensembles-eu.metoffice.com>) and the data providers in the ECA&D project (<http://www.ecad.eu>). The authors thank the two reviewers, who provided constructive comments on an earlier version of the manuscript, which helped clarify the text.

7 BIBLIOGRAPHY

Anagnostopoulou C, Tolika K, Maheras P, Kutiel H, Flocas HA. 2008. Performance of the general circulation HadAM3P model in simulating circulation types over the Mediterranean region. *International Journal of Climatology* **28**(2): 185–203. DOI: 10.1002/joc.1521.

Belleflamme A, Fettweis X, Erpicum M. 2014. Do global warming-induced circulation pattern changes affect temperature and precipitation over Europe during summer? *International Journal of Climatology* **35**(7): 1484–1499. DOI: 10.1002/joc.4070.

Belleflamme A, Fettweis X, Lang C, Erpicum M. 2013. Current and future atmospheric circulation at 500 hPa over Greenland simulated by the CMIP3 and CMIP5 global models. *Climate Dynamics* **41**(7–8): 2061–2080. DOI: 10.1007/s00382-012-1538-2.

Benech B, Brunet H, Jacq V, Payen M, Rivrain J-C, Santurette P. 1993. La catastrophe de Vaison-La-Romaine et les violentes précipitations de septembre 1992: aspects météorologiques. *La Météorologie* (1): 72–90.

Blenkinsop S, Chan SC, Kendon EJ, Roberts NM, Fowler HJ. 2015. Temperature influences on intense UK hourly precipitation and dependency on large-scale circulation. *Environmental Research Letters* **10**(5): 054021. DOI: 10.1088/1748-9326/10/5/054021.

Boé J, Terray L. 2008. A Weather-Type Approach to Analyzing Winter Precipitation in France: Twentieth-Century Trends and the Role of Anthropogenic Forcing. *Journal of Climate* **21**(13): 3118–3133. DOI: 10.1175/2007JCLI1796.1.

- 531 Bontron G. 2004. Prvision quantitative des prcipitations: adaptation probabiliste par recherche
532 d'analogues. Utilisation des r-analyses NCEP/NCAR et application aux prcipitations du Sud-Est de la
533 France. Institut Polytechnique de Grenoble, PhD thesis, 276 pp.
- 534 Brigode P, Brissette F, Nicault A, Perreault L, Kuentz A, Mathevet T, Gailhard J. 2016. Streamflow
535 variability over the 1881–2011 period in northern Qubec: comparison of hydrological reconstructions
536 based on tree rings and geopotential height field reanalysis. *Climate of the Past* **12**(9): 1785–1804. DOI:
537 10.5194/cp-12-1785-2016.
- 538 Brigode P, Bernardara P, Gailhard J, Garavaglia F, Ribstein P, Merz R. 2013a. Optimization of the
539 Geopotential Heights Information Used in a Rainfall-Based Weather Patterns Classification over
540 Austria. *International Journal of Climatology* **33**(6): 1563–1573. DOI:10.1002/joc.3535.
- 541 Brigode P, Miovi Z, Bernardara P, Paquet E, Garavaglia F, Gailhard J, Ribstein P. 2013b. Linking ENSO
542 and heavy rainfall events over coastal British Columbia through a weather pattern classification.
543 *Hydrology and Earth System Sciences* **17**(4): 1455–1473. DOI: 10.5194/hess-17-1455-2013.
- 544 Cassano JJ, Uotila P, Lynch A. 2006. Changes in synoptic weather patterns in the polar regions in the
545 twentieth and twenty-first centuries, part 1: Arctic. *International Journal of Climatology* **26**(8): 1027–
546 1049. DOI: 10.1002/joc.1306.
- 547 Cassou C, Terray L, Hurrell JW, Deser C. 2004. North Atlantic Winter Climate Regimes: Spatial
548 Asymmetry, Stationarity with Time, and Oceanic Forcing. *Journal of Climate* **17**(5): 1055–1068. DOI:
549 10.1175/1520-0442(2004)017<1055:NAWCRC>2.0.CO;2.
- 550 Cassou C, Terray L, Phillips AS. 2005. Tropical Atlantic Influence on European Heat Waves. *Journal of*
551 *Climate* **18**(15): 2805–2811. DOI: 10.1175/JCLI3506.1.
- 552 Cattiaux J, Douville H, Ribes A, Chauvin F, Plante C. 2012. Towards a better understanding of changes
553 in wintertime cold extremes over Europe: a pilot study with CNRM and IPSL atmospheric models.
554 *Climate Dynamics* **40**: 2433. DOI: 10.1007/s00382-012-1436-7.
- 555 Corti S, Molteni F, Palmer TN. 1999. Signature of recent climate change in frequencies of natural
556 atmospheric circulation regimes. *Nature* **398**(6730): 799–802. DOI: 10.1038/19745.
- 557 Covey C, AchutaRao KM, Cubasch U, Jones P, Lambert SJ, Mann ME, Phillips TJ, Taylor KE. 2003. An
558 overview of results from the Coupled Model Intercomparison Project. *Global and Planetary Change*
559 **37**(1–2): 103–133. DOI: 10.1016/S0921-8181(02)00193-5.
- 560 Dee DP, Uppala SM, Simmons AJ, Berrisford P, Poli P, Kobayashi S, Andrae U, Balmaseda MA, Balsamo
561 G, Bauer P, Bechtold P, Beljaars ACM, van de Berg L, Bidlot J, Bormann N, Delsol C, Dragani R, Fuentes
562 M, Geer AJ, Haimberger L, Healy SB, Hersbach H, Hlm EV, Isaksen I, Kllberg P, Khler M, Matricardi
563 M, McNally AP, Monge-Sanz BM, Morcrette J-J, Park B-K, Peubey C, de Rosnay P, Tavolato C, Thpaut
564 J-N, Vitart F. 2011. The ERA-Interim reanalysis: configuration and performance of the data assimilation
565 system. *Quarterly Journal of the Royal Meteorological Society* **137**(656): 553–597. DOI: 10.1002/qj.828.
- 566 Demuzere M, Werner M, van Lipzig NPM, Roeckner E. 2009. An analysis of present and future ECHAM5
567 pressure fields using a classification of circulation patterns. *International Journal of Climatology* **29**(12):
568 1796–1810. DOI: 10.1002/joc.1821.
- 569 Dunn-Sigouin E, Son S-W. 2013. Northern Hemisphere blocking frequency and duration in the CMIP5
570 models. *Journal of Geophysical Research* **118**(3): 1179–1188. DOI: 10.1002/jgrd.50143.

- 571 Garavaglia F, Gailhard J, Paquet E, Lang M, Garçon R, Bernardara P. 2010. Introducing a rainfall
572 compound distribution model based on weather patterns sub-sampling. *Hydrology and Earth System*
573 *Sciences* **14**(6): 951–964. DOI: 10.5194/hess-14-951-2010.
- 574 Garavaglia F, Lang M, Paquet E, Gailhard J, Garçon R, Renard B. 2011. Reliability and robustness of
575 rainfall compound distribution model based on weather pattern sub-sampling. *Hydrology and Earth*
576 *System Sciences* **15**(2): 519–532. DOI: 10.5194/hess-15-519-2011.
- 577 Gillett NP, Zwiers FW, Weaver AJ, Stott PA. 2003. Detection of human influence on sea-level pressure.
578 *Nature* **422**(6929): 292–294.
- 579 Giuntoli I, Renard B, Vidal J-P, Bard A. 2013. Low flows in France and their relationship to large-scale
580 climate indices. *Journal of Hydrology* **482**: 105–118. DOI: 10.1016/j.jhydrol.2012.12.038.
- 581 Hall A. 2014. Projecting regional change. *Science* **346**(6216): 1461–1462. DOI:
582 10.1126/science.aaa0629.
- 583 Handorf D, Dethloff K. 2012. How well do state-of-the-art atmosphere-ocean general circulation
584 models reproduce atmospheric teleconnection patterns? *Tellus A* **64**(0). DOI:
585 10.3402/tellusa.v64i0.19777.
- 586 Haylock MR, Hofstra N, Tank AMG., Klok EJ, Jones PD, New M. 2008. A European daily high-resolution
587 gridded data set of surface temperature and precipitation for 1950–2006. *Journal of Geophysical*
588 *Research* **113**(D20): D20119. DOI: 10.1029/2008JD010201.
- 589 Hertig E, Jacobeit J. 2014. Variability of weather regimes in the North Atlantic-European area: past and
590 future. *Atmospheric Science Letters* **15**(4): 314–320. DOI: 10.1002/asl2.505.
- 591 Hertig E, Seubert S, Paxian A, Vogt G, Paeth H, Jacobeit J. 2013. Statistical modelling of extreme
592 precipitation indices for the Mediterranean area under future climate change. *International Journal of*
593 *Climatology* **34**(4): 1132–1156. DOI: 10.1002/joc.3751.
- 594 Hsu CJ, Zwiers F. 2001. Climate change in recurrent regimes and modes of northern hemisphere
595 atmospheric variability. *Journal of Geophysical Research* **106**(D17): 20145–20159. DOI:
596 10.1029/2001JD900229.
- 597 Huth R, Beck C, Philipp A, Demuzere M, Ustrnul Z, Cahynová M, Kyselý J, Tveito OE. 2008. Classifications
598 of Atmospheric Circulation Patterns. *Annals of the New York Academy of Sciences* **1146**(1): 105–152.
599 DOI: 10.1196/annals.1446.019.
- 600 Küttel M, Luterbacher J, Wanner H. 2010. Multidecadal changes in winter circulation-climate
601 relationship in Europe: frequency variations, within-type modifications, and long-term trends. *Climate*
602 *Dynamics* **36**(5–6): 957–972. DOI: 10.1007/s00382-009-0737-y.
- 603 Lorenzo MN, Ramos AM, Taboada JJ, Gimeno L. 2011. Changes in Present and Future Circulation Types
604 Frequency in Northwest Iberian Peninsula. *PLoS ONE* **6**(1). DOI: 10.1371/journal.pone.0016201.
- 605 Lucas-Picher P, Cattiaux J, Bougie A, Laprise R. 2016. How does large-scale nudging in a regional climate
606 model contribute to improving the simulation of weather regimes and seasonal extremes over North
607 America? *Climate Dynamics* **46**(3–4): 929–948. DOI: 10.1007/s00382-015-2623-0.

- 608 McKendry IG, Stahl K, Moore RD. 2006. Synoptic sea-level pressure patterns generated by a general
609 circulation model: comparison with types derived from NCEP/NCAR re-analysis and implications for
610 downscaling. *International Journal of Climatology* **26**(12): 1727–1736. DOI: 10.1002/joc.1337.
- 611 Michelangeli P-A, Vautard R, Legras B. 1995. Weather Regimes: Recurrence and Quasi Stationarity.
612 *Journal of Atmospheric Sciences* **52**: 1237–1256. DOI: 10.1175/1520-
613 0469(1995)052<1237:WRRASQ>2.0.CO;2.
- 614 Moss RH, Edmonds JA, Hibbard KA, Manning MR, Rose SK, Vuuren DP van, Carter TR, Emori S, Kainuma
615 M, Kram T, Meehl GA, Mitchell JFB, Nakicenovic N, Riahi K, Smith SJ, Stouffer RJ, Thomson AM, Weyant
616 JP, Wilbanks TJ. 2010. The next generation of scenarios for climate change research and assessment.
617 *Nature* **463**(7282): 747–756. DOI: 10.1038/nature08823.
- 618 Murawski A, Bürger G, Vorogushyn S, Merz B. 2016. Can local climate variability be explained by
619 weather patterns? A multi-station evaluation for the Rhine basin. *Hydrology and Earth System Sciences*
620 **20**(10): 4283–4306. DOI: 10.5194/hess-20-4283-2016.
- 621 Nied M, Pardowitz T, Nissen K, Ulbrich U, Hundecha Y, Merz B. 2014. On the relationship between
622 hydro-meteorological patterns and flood types. *Journal of Hydrology* **519**: 3249–3262. DOI:
623 10.1016/j.jhydrol.2014.09.089.
- 624 Nuissier O, Joly B, Joly A, Ducrocq V, Arbogast P. 2011. A statistical downscaling to identify the large-
625 scale circulation patterns associated with heavy precipitation events over southern France. *Quarterly*
626 *Journal of the Royal Meteorological Society* **137**(660): 1812–1827. DOI: 10.1002/qj.866.
- 627 Obled C, Bontron G, Garçon R. 2002. Quantitative precipitation forecasts: a statistical adaptation of
628 model outputs through an analogues sorting approach. *Atmospheric Research* **63**(3–4): 303–324. DOI:
629 10.1016/S0169-8095(02)00038-8.
- 630 Pastor MA, Casado MJ. 2012. Use of circulation types classifications to evaluate AR4 climate models
631 over the Euro-Atlantic region. *Climate Dynamics* **39**(7–8): 2059–2077. DOI: 10.1007/s00382-012-1449-
632 2.
- 633 Philipp A, Della-Marta PM, Jacobeit J, Fereday DR, Jones PD, Moberg A, Wanner H. 2007. Long-Term
634 Variability of Daily North Atlantic–European Pressure Patterns since 1850 Classified by Simulated
635 Annealing Clustering. *Journal of Climate* **20**(16): 4065–4095. DOI: 10.1175/JCLI4175.1.
- 636 Planchon O, Quénol H, Dupont N, Corgne S. 2009. Application of the Hess-Brezowsky classification to
637 the identification of weather patterns causing heavy winter rainfall in Brittany (France). *Natural*
638 *Hazards and Earth System Sciences* **9**(4): 1161–1173. DOI: 10.5194/nhess-9-1161-2009.
- 639 Plaut G, Simonnet E. 2001. Large-scale circulation classification, weather regimes, and local climate
640 over France, the Alps and Western Europe. *Climate Research* **17**(3): 303–324. DOI: 10.3354/cr017303.
- 641 R Core Team. 2016. *R A Language and Environment for Statistical Computing*. R Foundation for
642 Statistical Computing: Vienna, Austria.
- 643 Renard B, Lall U. 2014. Regional frequency analysis conditioned on large-scale atmospheric or oceanic
644 fields. *Water Resources Research* **50**(12): 9536–9554. DOI: 10.1002/2014WR016277.
- 645 Sanchez-Gomez E, Somot S, Déqué M. 2009. Ability of an ensemble of regional climate models to
646 reproduce weather regimes over Europe-Atlantic during the period 1961–2000. *Climate Dynamics*
647 **33**(5): 723–736. DOI: 10.1007/s00382-008-0502-7.

- 648 Santos JA, Belo-Pereira M, Fraga H, Pinto JG. 2016. Understanding climate change projections for
649 precipitation over Western Europe with a weather typing approach. *Journal of Geophysical Research*
650 **121**(3): 2015JD024399. DOI: 10.1002/2015JD024399.
- 651 Sénési S, Bougeault P, Chèze J-L, Cosentino P, Thepenier R-M. 1996. The Vaison-La-Romaine Flash
652 Flood: Mesoscale Analysis and Predictability Issues. *Weather and Forecasting* **11**(4): 417–442. DOI:
653 10.1175/1520-0434(1996)011<0417:TVLRFF>2.0.CO;2.
- 654 Sheridan SC, Lee CC. 2010. Synoptic climatology and the general circulation model. *Progress in Physical*
655 *Geography* **34**(1): 101–109. DOI: 10.1177/0309133309357012.
- 656 Stephenson DB, Hannachi A, O'Neill A. 2004. On the existence of multiple climate regimes. *Quarterly*
657 *Journal of the Royal Meteorological Society* **130**(597): 583–605. DOI: 10.1256/qj.02.146.
- 658 Taylor KE, Stouffer RJ, Meehl GA. 2012. An overview of CMIP5 and the experiment design. *Bulletin of*
659 *the American Meteorological Society* **93**(4): 485–498. DOI: 10.1175/BAMS-D-11-00094.1.
- 660 Teweles J, Wobus H. 1954. Verification of prognosis charts. *Bulletin of the American Meteorological*
661 *Society* **35**(10): 455–463.
- 662 Tramblay Y, Neppel L, Carreau J. 2011. Brief communication - Climatic covariates for the frequency
663 analysis of heavy rainfall in the Mediterranean region. *Natural Hazards and Earth System Sciences*
664 **11**(9): 2463–2468. DOI: 10.5194/nhess-11-2463-2011.
- 665 Tramblay Y, Neppel L, Carreau J, Najib K. 2013. Non-stationary frequency analysis of heavy rainfall
666 events in southern France. *Hydrological Sciences Journal* **58**(2): 280–294. DOI:
667 10.1080/02626667.2012.754988.
- 668 Trenberth KE, Fasullo JT, Shepherd TG. 2015. Attribution of climate extreme events. *Nature Climate*
669 *Change* **5**(8): 725–730. DOI: 10.1038/nclimate2657.
- 670 Ullmann A, Fontaine B, Roucou P. 2014. Euro-Atlantic weather regimes and Mediterranean rainfall
671 patterns: present-day variability and expected changes under CMIP5 projections. *International Journal*
672 *of Climatology* **34**(8): 2634–2650. DOI: 10.1002/joc.3864.
- 673 Vrac M, Hayhoe K, Stein M. 2007. Identification and intermodel comparison of seasonal circulation
674 patterns over North America. *International Journal of Climatology* **27**(5): 603–620. DOI:
675 10.1002/joc.1422.
- 676 Wetterhall F, Halldin S, Xu C. 2005. Statistical precipitation downscaling in central Sweden with the
677 analogue method. *Journal of Hydrology* **306**(1–4): 174–190. DOI: 10.1016/j.jhydrol.2004.09.008.
- 678 Wilby RL, Quinn NW. 2013. Reconstructing multi-decadal variations in fluvial flood risk using
679 atmospheric circulation patterns. *Journal of Hydrology* **487**: 109–121. DOI:
680 10.1016/j.jhydrol.2013.02.038.