



Local time for lattice paths and the associated limit laws

Cyril Banderier, Michael Wallner

► **To cite this version:**

Cyril Banderier, Michael Wallner. Local time for lattice paths and the associated limit laws. CEUR Workshop Proceedings, CEUR Workshop Proceedings, In press. <hal-01796320>

HAL Id: hal-01796320

<https://hal.archives-ouvertes.fr/hal-01796320>

Submitted on 19 May 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Local time for lattice paths and the associated limit laws

Cyril Banderier

CNRS & Univ. Paris Nord, France

<http://lipn.fr/~banderier/>

 <https://orcid.org/0000-0003-0755-3022>

Michael Wallner

LaBRI, Université de Bordeaux, France

<http://dmg.tuwien.ac.at/mwallner/>

 <https://orcid.org/0000-0001-8581-449X>

Abstract

For generalized Dyck paths (i.e., directed lattice paths with any finite set of jumps), we analyse their local time at zero (i.e., the number of times the path is touching or crossing the abscissa). As we are in a discrete setting, the event we analyse here is “invisible” to the tools of Brownian motion theory.

It is interesting that the key tool for analysing directed lattice paths, which is the kernel method, is not directly applicable here. Therefore, we introduce a variant of this kernel method to get the trivariate generating function (length, final altitude, local time): this leads to an expression involving symmetric and algebraic functions.

We apply this analysis to different types of constrained lattice paths (meanders, excursions, bridges, . . .). Then, we illustrate this approach on “basketball walks” which are walks defined by the jumps $-2, -1, 0, +1, +2$. We use singularity analysis to prove that the limit laws for the local time are (depending on the drift and the type of walk) the geometric distribution, the negative binomial distribution, the Rayleigh distribution, or the half-normal distribution (a universal distribution up to now rarely encountered in analytic combinatorics).

Keywords: Lattice paths, generating function, analytic combinatorics, singularity analysis, kernel method, generalized Dyck paths, algebraic function, local time, half-normal distribution, Rayleigh distribution, negative binomial distribution

1 Introduction

This article continues our series of investigations on the enumeration, generation, and asymptotics of directed lattice paths [ABBG18a, ABBG18b, BF02, BW14, BW17, BG06, BKK⁺17, Wal16]. Such lattice paths are a fundamental combinatorial structure ubiquitous in computer science (evolution of a stack, bijections with trees, permutations, . . .), probability theory (linked with random walks or queuing theory), and statistical mechanics (as basic building blocks for more general 2D models), to name a few.

Copyright © by the paper's authors. Copying permitted for private and academic purposes.

In: Luca Ferrari, Malvina Vamvakari (eds.): Proceedings of the conference GASCom 2018, Athens, Greece, 18-20 June 2018, published at <http://ceur-ws.org>





	ending anywhere	ending at 0
unconstrained (on \mathbb{Z})	 walk/path (\mathcal{W})	 bridge (\mathcal{B})
constrained (on \mathbb{N})	 meander (\mathcal{M})	 excursion (\mathcal{E})

Table 1: Four types of paths: walks, bridges, meanders, and excursions.

Let us give a definition of the lattice paths we consider:

Definition 1.1 (Jumps and lattice paths). A step set $\mathcal{S} \subset \mathbb{Z}^2$ is a finite set of vectors $\{(x_1, y_1), \dots, (x_m, y_m)\}$. An n -step lattice path or walk is a sequence of vectors (v_1, \dots, v_n) , such that v_j is in \mathcal{S} . Geometrically, it may be interpreted as a sequence of points $\omega = (\omega_0, \omega_1, \dots, \omega_n)$ where $\omega_i \in \mathbb{Z}^2$, $\omega_0 = (0, 0)$ (or another starting point) and $\omega_i - \omega_{i-1} = v_i$ for $i = 1, \dots, n$. The elements of \mathcal{S} are called steps or jumps. The length $|\omega|$ of a lattice path is its number n of jumps.

We restrict our attention to *directed paths* which are defined by the fact that, for each jump $(x, y) \in \mathcal{S}$, one must have $x \geq 0$. Lattice paths can have different additional constraints shown in Table 1.

Note that it is possible to encode lattice paths in words. Then, the constrained lattice paths we consider can be enumerated by context-free grammars [LY90, MRSV99, Duc00]. One drawback of the grammar approach is that it is not easy to get universal asymptotic results from it, even if it is possible to establish generic results on the associated critical exponents [BD15]. Also, the grammar approach quickly leads to heavy case-by-case computations (resultants of equations of huge degree) as soon as the set of jumps \mathcal{S} contains a large jump.

In this article, we show how to proceed for the enumeration and the asymptotics in these harder cases: our techniques are relying on the “kernel method” which (contrary to the context-free grammar approach) offers access to the *generic* structure of the final generating functions and the *universality* of their asymptotics via singularity analysis [BF02, FS09].

The following convenient notation, a variant of the Omega operator of MacMahon, will be another of our ingredients:

$$\{u^{>0}\} := \sum_{n>0} u^n [u^n] \quad \text{and} \quad \{u^{<0}\} := \sum_{n<0} u^n [u^n]$$

where $[u^n]g(u)$ stands for the coefficient of u^n in a (Laurent) power series $g(u)$ from $\mathbb{C}((u))$ or $\mathbb{C}[u, 1/u][[z]]$. MacMahon introduced this operator on rational functions, in order to get binomial identities or integer partition formulae [Mac15, Section VIII]. In the late 1990s, this operator experienced a strong revival, mostly by work/packages of Andrews, Paule, Riese, and Han [APR01, Han03]. Another nice application of the Omega operator is the proof of D-finiteness of some walks in the quarter plane by Bousquet-Mélou and Mishna [BMM10]. In this article we make use of this operator, not on sums or products of rational functions (like it is the case for integer partitions, or for quarter plane walks), but on the level of functional equations involving algebraic functions.

This equation can be read as “ $W =$ empty walk or the walk is at some altitude encoded by u^k and multiply by $P(u)$ to do all the jumps, then we remove the ones crossing the x -axis, and re-add them with a marker q .” This is conveniently rewritten as

$$(1 - zP(u)) \left(\sum_{k \in \mathbb{Z}} W_k u^k \right) = 1 - z(1 - q) \left(\sum_{k=-d+1}^{-1} \{u^{>0}\} (P(u)u^k) W_k + \sum_{k=1}^{c-1} \{u^{<0}\} (P(u)u^k) W_k \right).$$

Here, one may think that we could apply the classical kernel method (see e.g. [BF02]): one substitutes u by any root $u(z)$ of $1 - zP(u)$, as this will cancel the left-hand side, and thus doing that for all roots will lead to a system of $c + d$ unknowns, with $c + d$ independent relations. Hence, bingo, solving this system gives a closed-form formula for $W(z, u, q)$! Unfortunately, this is not working: indeed, in such an equation, if one substitutes a variable by another expression, one has to take care to stay in the ring of formal power series (in order to avoid non-trivial zero divisors, as exemplified by the phenomenon $(1 - u) \sum_{k \in \mathbb{Z}} u^k = 0$). In our equation, F belongs to $\mathbb{C}[u, 1/u][q][[z]]$. As the exponents of u range from $-\infty$ to $+\infty$, it is not legitimate to substitute u by our Puiseux series $u(z)$, because this would lead to arbitrary negative and positive powers of z . We need to adapt the kernel method and to transform it into what we call a “bilateral kernel method”:

1. Extract the positive part $\{u^{>0}\}$. This gives the following equation

$$\{u^{>0}\} \left((1 - zP(u)) \sum_{k=-d+1}^{\infty} W_k u^k \right) = -z(1 - q) \sum_{k=-d+1}^{-1} \{u^{>0}\} (P(u)u^k) W_k$$

which can be rewritten as

$$(1 - zP(u)) \sum_{k=c+1}^{\infty} W_k u^k = -z(1 - q) \sum_{k=-d+1}^{-1} \{u^{>0}\} (P(u)u^k) W_k - \{u^{>0}\} \left((1 - zP(u)) \sum_{k=-d+1}^c W_k u^k \right).$$

In this expression, it is legitimate to substitute u by the c roots $u_i(z)$ of $1 - zP(u)$ such that $u_i(z) \sim 0$ for $z \sim 0$. Thus we get c new equations:

$$0 = \left(zq \sum_{k=-d+1}^{-1} \{u^{>0}\} (P(u)u^k) W_k + z \{u^{>0}\} (P(u)) W_0 - \sum_{k=1}^c \{u^{>0}\} ((1 - zP(u))u^k) W_k \right)_{|u=u_i(z)}.$$

2. Extract the negative part $\{u^{<0}\}$. This leads to an equation in which it is legitimate to substitute u by the d roots $v_i(z)$ of $1 - zP(u)$ such that $|v_i(z)| \sim \infty$ for $z \sim 0$. Thus we get d new equations:

$$0 = \left(zq \sum_{k=1}^{c-1} \{u^{<0}\} (P(u)u^k) W_k + z \{u^{<0}\} (P(u)) W_0 - \sum_{k=-d}^{-1} \{u^{<0}\} ((1 - zP(u))u^k) W_k \right)_{|u=v_i(z)}.$$

3. Extract $[u^0]$. It gives one additional equation:

$$[u^0] \left((1 - zP(u)) \left(\sum_{k=-d}^c W_k u^k \right) \right) = 1.$$

All these equations involve (a subset of) of the yet unknown $c + d + 1$ auxiliary functions W_{-d}, \dots, W_c . This system fully allows to reconstruct the initial equation: W (the linear combination of the W_k 's) is also a solution of the initial functional equation, which was a contraction in the space of formal power series $\mathbb{C}[q, u, 1/u][[z]]$, therefore this functional equation had a unique solution in this ring, which we thus identified as being W , the only candidate for it. This explains why the system of equations is of full rank: it has unique power series solutions W_k , all of them expressible as a quotient of polynomials having the roots as variable. (We say more on this shape in the full version of this article: it requires an excursion (sic) via the Schur function world!) So, in all cases, one gets a closed form for the generating function W (and the W_k 's) in terms of the roots of the kernel. \square

Now, the local analysis of these roots at their branching points, as done in [BF02] allows to get the Puiseux behaviour of the functions $W(z, u, 1)$ and $W_0(z, u)$ near their dominant singularity (for any fixed value of u , this dominant singularity is still the same ρ as in [BF02]). It allows to rewrite locally these functions (for $z \sim \rho$) in the framework of the schemes developed in [Wal16], on which we comment more in the next section.

Let us now illustrate this approach on basketball walks (walks with jumps $-2, -1, 0, +1, +2$, see [BKK⁺17]). This leads to the simple linear system:

$$\begin{cases} W_0 = 1 + z(W_{-1} + W_{-2} + W_0 + W_1 + W_2) \\ qzW_{-1} + (zu_1 + z)W_0 + (zu_1^2 + zu_1 + z - 1)W_1 + (zu_1^3 + zu_1^2 + zu_1 + z - u_1)W_2 = 0 \\ qzW_{-1} + (zu_2 + z)W_0 + (zu_2^2 + zu_2 + z - 1)W_1 + (zu_2^3 + zu_2^2 + zu_2 + z - u_2)W_2 = 0 \\ qzW_1v_1^3 + (v_1 + 1)zW_0v_1^2 + ((z - 1)v_1^2 + zv_1 + z)v_1W_{-1} + (zv_1^3 + (z - 1)v_1^2 + zv_1 + z)W_{-2} = 0 \\ qzW_1v_2^3 + (v_2 + 1)zW_0v_2^2 + ((z - 1)v_2^2 + zv_2 + z)v_2W_{-1} + (zv_2^3 + (z - 1)v_2^2 + zv_2 + z)W_{-2} = 0 \end{cases}$$

Solving it gives a closed form expression for $W(z, u, q)$ lengthy to write. If one sets $u = 1$, the Ferrari–Bombelli formula can be used and gives the following closed form for the generating functions of the walks:

$$W(z, 1, q) = \frac{z(z+1)(q-1)^2(4q^2z^2 - 3qz^2 - 6qz + 4z^2 + q)\sqrt{z-1}}{2B\sqrt{5z-1}} + \frac{(q^2-1)\sqrt[4]{z-1}\sqrt{z}\sqrt{C-DE}}{\sqrt{2}B(5z-1)^{3/4}} - \frac{A}{2B}$$

where $E := \sqrt{(5z-1)(z-1)}$ and

$$\begin{aligned} A &:= 5(q^2 + 1)(4q^2 - 3q + 4)z^4 + (2q^4 - 49q^3 + 24q^2 - 49q + 2)z^3 - (8q^4 + 11q^3 - 20q^2 + 11q + 8)z^2 \\ &\quad + (2q^4 + 13q^3 + 16q^2 + 13q + 2)z - 2q(q+1)^2, \\ B &:= (31q^4 + q^3 + 61q^2 + q + 31)z^5 - (8q^4 + 83q^3 + 18q^2 + 83q + 8)z^4 - 2(7q^4 + 3q^3 - 15q^2 + 3q + 7)z^3 \\ &\quad + 2(4q^4 + 17q^3 + 20q^2 + 17q + 4)z^2 - (q^4 + 11q^3 + 19q^2 + 11q + 1)z + q^3 + 2q^2 + q, \\ C &:= (z-1)(5z-1)(z+1)\left((44q^4 - 26q^3 + 89q^2 - 26q + 44)z^5 + (4q^4 - 126q^3 + 44q^2 - 126q + 4)z^4\right. \\ &\quad \left.+ (-20q^4 - 14q^3 + 58q^2 - 14q - 20)z^3 + (4q^4 + 46q^3 + 24q^2 + 46q + 4)z^2 + (-8q^3 - 27q^2 - 8q)z + 4q^2\right), \\ D &:= 5(34q^4 - 6q^3 + 69q^2 - 6q + 34)z^7 + 2(22q^4 - 243q^3 + 67q^2 - 243q + 22)z^6 \\ &\quad - (132q^4 + 196q^3 - 81q^2 + 196q + 132)z^5 - 4(2q^4 - 81q^3 - 42q^2 - 81q + 2)z^4 \\ &\quad + (26q^4 + 58q^3 - 125q^2 + 58q + 26)z^3 - 2(2q^4 + 31q^3 + 29q^2 + 31q + 2)z^2 + q(8q^2 + 35q + 8)z - 4q^2. \end{aligned}$$

Note that the Cardan/Ferrari–Bombelli formulas are perhaps nice for the eyes (arguably!), but they are not the right way to handle these generating functions from a computer algebra point of view. It is indeed better to work directly with symmetric functions of the roots u_i 's (and this advantageously also allows to handle the cases of degree > 4): this is an efficient algorithmic way to use the Newton relations/Vieta's formulas between the roots of the kernel. Using the expression with the small roots is also the way to follow the Puiseux behaviour of these symmetric functions, as we detail in the forthcoming full version.

What is more, the above linear system shows that it is now routine to get in a few minutes thousands of coefficients of our generating functions (e.g. via Newton iteration, in any computer algebra software). Here are the first terms of the generating functions of x -axis crossings for walks:

$$\begin{aligned} W(z, 1, q) = & 1 + 5z + (2q + 23)z^2 + (2q^2 + 14q + 109)z^3 + \\ & (2q^3 + 16q^2 + 88q + 519)z^4 + (2q^4 + 18q^3 + 112q^2 + 504q + 2489)z^5 + \\ & (2q^5 + 20q^4 + 138q^3 + 700q^2 + 2776q + 11989)z^6 + \\ & (2q^6 + 22q^5 + 166q^4 + 930q^3 + 4150q^2 + 14896q + 57959)z^7 + \\ & (2q^7 + 24q^6 + 196q^5 + 1196q^4 + 5878q^3 + 23720q^2 + 78614q + 280995)z^8 + \\ & (2q^8 + 26q^7 + 228q^6 + 1500q^5 + 8004q^4 + 35518q^3 + 132264q^2 + 410046q + 1365537)z^9 + O(z^{10}). \end{aligned}$$

What is the asymptotic behaviour of these polynomials in q ? This is what we present in the next section.

3 Limit laws

Theorem 3.1 (Limit law for the local time at 0). *For a walk with step set encoded by $P(u)$, the limit laws for the local time at 0 depend on the drift $P'(1)$ of the walk (see Table 2):*

Type of the walk	Limit law
Excursion	Negative binomial distribution
Meander with drift < 0	Negative binomial distribution
Meander with drift ≥ 0	Geometric distribution
Walk with non-zero drift	Geometric distribution
Walk with zero drift	Half-normal distribution $\mathcal{H}(\lambda)$
Bridge	Rayleigh distribution $\mathcal{R}(\lambda)$

The parameters of these distributions are given in the proof and in Table 2: E.g. for bridges, the parameter of $\mathcal{R}(\lambda)$ is $\lambda = \sqrt{P''(1)/P(1)}$, for walks, the parameter of $\mathcal{H}(\lambda)$ is $\lambda = \tau/2\sqrt{P(1)/P''(1)}$, where τ is the unique real positive value such that $P'(\tau) = 0$.

What is more, if defined, the limit laws for the the number of x -axis crossings are the same as the ones of the local time (with an appropriate new value for the distribution parameters).

Proof (Sketch). Thanks to the Puiseux expansions following from Theorem 2.1 and [BF02], it is possible to derive the limit laws: we then get a shape on which we can apply the results of Drmota and Soria [DS97] on the Rayleigh and Gaussian distributions, and of [Wal16] on the half-normal distribution. These distributions are depicted in Table 2. Details are omitted in this extended abstract.

In the cases of excursions and meanders the local time is equal to the number of returns to zero. The results for excursions were derived by Banderier and Flajolet in [BF02, Theorem 5]. What is more, all ingredients for the case of meanders are also given in this paper, and the result follows the same lines. However, due to the drift dependent number of meanders (compare [BF02, Theorem 4]) three regimes need to be considered, leading to two different limit laws: negative binomial and geometric.





	<u>Geometric</u>	<u>Negative binomial</u>	<u>Half-normal</u>	<u>Rayleigh</u>
	Geom(p)	NB(m, p)	$\mathcal{H}(\lambda)$	$\mathcal{R}(\lambda)$
				
Support	$k \in \{0, 1, \dots\}$	$k \in \{0, 1, \dots\}$	$x \in \mathbb{R}_{\geq 0}$	$x \in \mathbb{R}_{\geq 0}$
PDF	$(1-p)^k p$	$\binom{m+k-1}{k} (1-p)^k p^m$	$\sqrt{\frac{2}{\pi\lambda^2}} \exp\left(-\frac{x^2}{2\lambda^2}\right)$	$\frac{x}{\lambda^2} \exp\left(-\frac{x^2}{2\lambda^2}\right)$
Mean	$\frac{1-p}{p}$	$\frac{m(1-p)}{p}$	$\lambda\sqrt{\frac{2}{\pi}}$	$\lambda\sqrt{\frac{\pi}{2}}$
Variance	$\frac{1-p}{p^2}$	$\frac{m(1-p)}{p^2}$	$\lambda^2\left(1 - \frac{2}{\pi}\right)$	$\lambda^2\left(2 - \frac{\pi}{2}\right)$

Table 2: A comparison of the geometric, negative binomial, half-normal, and Rayleigh distribution. They are the four distributions occurring for the local time of generalized Dyck paths. Feller [Fel68, Chapter 3] did the analysis for the simplest case of walks with jumps $-1, +1$, our article shows how to tackle the more difficult situation involving any set of jumps of amplitude ≥ 1 .

For the cases of walks and bridges the local time is equal to the number of returns to zero and the number of x -axis crossings. The results for the number of returns to zero of walks were derived by Wallner in [Wal16, Theorem 4.2] leading to a geometric or a half-normal distribution, depending whether the drift $P'(1)$ is non-zero or zero, respectively. As above, the result for the case of returns to zeros in bridges, follows also the same lines as the previous one and we omit the details in this extended abstract. In this case one uses the limit law of Drmota and Soria [DS97, Theorem 1] to prove the existence of a Rayleigh distribution.

It remains to consider the laws of x -axis crossings in the cases of bridges and walks. The proof of Theorem 2.1 gives access to closed forms of $W_0(z, q)$ and $W(z, 1, q)$, which are the generating functions of bridges and walks where crossings of the x -axis are marked by q . In order to treat both at the same time we abbreviate them until the end of this proof by $F(z, q)$. We want to apply either [DS97, Theorem 1] or [Wal16, Theorem 2.1]. Note that the technical conditions of these theorems are satisfied due to the closed forms in terms of small and large branches, and the fact, that due to the Weierstrass Preparation Theorem, the branches $u_1(z)$ and $v_1(z)$ (which are the real positive branches for $z > 0$ in the vicinity of 0, see Figure 2) satisfy the necessary conditions (compare the derivations in [Wal16]). In particular, as proven in [BF02], they satisfy a square root behaviour with the following expansion at $z = \rho$:

$$u_1(z) = \tau - \sqrt{2 \frac{P(\tau)}{P''(\tau)}} \sqrt{1 - z/\rho} + \dots,$$

$$v_1(z) = \tau + \sqrt{2 \frac{P(\tau)}{P''(\tau)}} \sqrt{1 - z/\rho} + \dots$$

Due to the methods derived in [BW17], we may assume without loss of generality, that our model is aperiodic. For such aperiodic models (like e.g. Motzkin and basketball walks) there exists a unique singularity $\rho > 0$ of $F(z, 1)$.

Now, the key fact is that our generating functions F have locally the following behaviour

$$\frac{1}{F(z, q)} = g(z, q) + h(z, q) \sqrt{1 - z/\rho},$$

for $|q - 1| < \varepsilon$ and $|z - \rho| < \varepsilon$ with $\arg(z - \rho) \neq 0$ where $\varepsilon > 0$ is some fixed real number, and $g(z, q)$ and $h(z, q)$ are analytic functions. Additionally, one has here that $g(\rho, 1) = 0$. Finally, in the case of bridges we get $g_q(\rho, 1) < 0$ and $h(\rho, 1) \neq 0$ yielding a Rayleigh law. Whereas, in the case of walks with zero drift we get $g_q(\rho, 1) = g_{qq}(\rho, 1) = 0$ and $h(\rho, 1) = 0$, but $g_z(\rho, 1) \neq 0$ and $h_q(\rho, 1) \neq 1$ giving a half-normal law. The respective parameters depend on the chosen step set. \square




Figure 2: The root u_1 and v_1 of the kernel $1 - zP(u)$ are crossing at $z = \rho$ with a square root behaviour. This implies the Puiseux expansions with a square root used in the proof above.




Figure 3: Distribution of X_n , the local time at 0 for bridges of length n . (Left: $n = 200$. Right: $n = 4000$. Step set $\mathcal{S} = [-2, -1, 0, 1, 2]$). One quickly observes a nearly perfect match of the distribution (in red) of X_n with the Rayleigh limit law (in green). This fast convergence is well explained by the error terms we get via analytic combinatorics.




Figure 4: Distribution of X_n , the local time at 0 for walks of length n . (Left: $n = 200$. Right: $n = 4000$. Step set $\mathcal{S} = [-2, -1, 0, 1, 2]$). One quickly observes a nearly perfect match of the distribution (in red) of X_n with the half-normal limit law (in green). Here again, this fast convergence is well explained by the error terms we get via analytic combinatorics.

4 Conclusion

In this article we showed how to derive the generating function for the local time at 0 of directed lattice paths. It completes the work of Banderier and Flajolet [BF02], who just handled the case of returns to zero of excursions. It is also extending the work of Feller [Fel68] and later Wallner [Wal16], who did the Dyck and Motzkin cases (for meanders/walks).

In order to solve the generating functions in the more general case, we used a mixture of the Omega operator and the kernel method. This leads to expressions from which we showed how to derive the limit law of the local time, for several models of constrained lattice paths. In the full version of this article, we give more closed form formulas, and we show that other parameters (like the number of “changes of signs”, or the number of jumps from positive to negative altitude) can also be analysed using our approach, and that they satisfy similar limit laws.

These parameters are very natural for discrete random walks, it is interesting that it is not possible to analyse them via a Brownian motion approach: indeed a Brownian motion can be seen as the limit after a rescaling of the amplitude by \sqrt{n} and the length by n (see [Mar03]). This rescaling implies that (discrete) jumps crossing the abscissa would be of amplitude 0, and are therefore completely erased. It is therefore nice that analytic combinatorics can get the asymptotics of this “discrete local time”, and the corresponding universal limit laws, while there are “invisible” via a Brownian motion approach. In a forthcoming article, we tackle further analysis of the height of discrete lattice paths; this allows to get the connection with the Brownian local time.

Acknowledgments: This work was started via collaboration funded by the SFB project F50 “Algorithmic and Enumerative Combinatorics” and the Franco-Austrian PHC “Amadeus”, and ended during the postdoctoral position of Michael Wallner at the University of Paris Nord, in September-December 2017, thanks to a MathStic funding. Michael Wallner is currently supported by the Erwin Schrödinger Fellowship of the Austrian Science Fund (FWF): J 4162-N35.

References

- [ABBG18a] Andrei Asinowski, Axel Bacher, Cyril Banderier, and Bernhard Gittenberger. [Analytic combinatorics of lattice paths with forbidden patterns: asymptotic aspects](#). *AofA proceedings (Analysis of Algorithms)*, 2018.
- [ABBG18b] Andrei Asinowski, Axel Bacher, Cyril Banderier, and Bernhard Gittenberger. [Analytic combinatorics of lattice paths with forbidden patterns: enumerative aspects](#). *LATA proceedings (Language and Automata Theory and Applications)*, 2018.
- [APR01] George E. Andrews, Peter Paule, and Axel Riese. [MacMahon’s partition analysis: the Omega package](#). *European J. Combin.*, 22(7):887–904, 2001.
- [BD15] Cyril Banderier and Michael Drmota. [Formulae and Asymptotics for Coefficients of Algebraic Functions](#). *Combinatorics, Probability, and Computing*, 24:1–53, 2015.
- [BF02] Cyril Banderier and Philippe Flajolet. [Basic analytic combinatorics of directed lattice paths](#). *Theoret. Comput. Sci.*, 281(1-2):37–80, 2002.
- [BG06] Cyril Banderier and Bernhard Gittenberger. [Analytic combinatorics of lattice paths: enumeration and asymptotics for the area](#). *Discrete Math. Theor. Comput. Sci. Proc.*, AG:345–355, 2006.
- [BKK⁺17] Cyril Banderier, Christian Krattenthaler, Alan Krinik, Dmitry Kruchinin, Vladimir Kruchinin, David Nguyen, and Michael Wallner. [Explicit formulas for enumeration of lattice paths: basketball and the kernel method](#). In *Lattice paths combinatorics and applications*, Developments in Mathematics Series. Springer, 2017. To appear.
- [BMM10] Mireille Bousquet-Mélou and Marni Mishna. [Walks with small steps in the quarter plane](#). In *Algorithmic probability and combinatorics*, volume 520 of *Contemp. Math.*, pages 1–39. Amer. Math. Soc., Providence, RI, 2010.
- [BW14] Cyril Banderier and Michael Wallner. [Some reflections on lattice paths](#). *AofA 2014, Paris*, 2014.

- [BW17] Cyril Banderier and Michael Wallner. [The kernel method for lattice paths below a rational slope](#). In *Lattice path combinatorics and applications*, Developments in Mathematics Series. Springer, 2017. To appear.
- [CV61] Endre Csáki and István Vincze. On some problems connected with the Galton-test. *Magyar Tud. Akad. Mat. Kutató Int. Közl.*, 6:97–109, 1961.
- [DS97] Michael Drmota and Michèle Soria. [Images and preimages in random mappings](#). *SIAM Journal on Discrete Mathematics*, 10(2):246–269, 1997.
- [Duc00] Philippe Duchon. [On the enumeration and generation of generalized Dyck words](#). *Discrete Math.*, 225(1-3):121–135, 2000. Formal power series and algebraic combinatorics (Toronto, 1998).
- [Fel68] William Feller. *An introduction to probability theory and its applications. Vol. I*. Third edition. Wiley & Sons, New York, 1968.
- [Fel71] William Feller. *An introduction to probability theory and its applications. Vol. II*. Second edition. John Wiley & Sons, Inc., New York-London-Sydney, 1971.
- [FS09] Philippe Flajolet and Robert Sedgewick. *Analytic Combinatorics*. Cambridge University Press, 2009.
- [Han03] Guo-Niu Han. [A general algorithm for the MacMahon omega operator](#). *Ann. Comb.*, 7(4):467–480, 2003.
- [Jai66] Gopi Chand Jain. [Joint distribution of intersections, \$\(\pm\)\$ waves and \$\(\pm\)\$ steps. I](#). *Proc. Nat. Inst. Sci. India Part A*, 32:460–471, 1966.
- [LY90] Jacques Labelle and Yeong-Nan Yeh. [Generalized Dyck paths](#). *Discrete Math.*, 82(1):1–6, 1990.
- [Mac15] Percy Alexander MacMahon. *Combinatory analysis. Volume 2*. Cambridge: Univ. Press, 1915.
- [Mar03] Philippe Marchal. [Constructing a sequence of random walks strongly converging to Brownian motion](#). In Cyril Banderier and Christian Krattenthaler, editors, *Discrete random walks (Paris, 2003)*, volume AC, pages 181–190. Discrete Math. Theor. Comput. Sci., 2003.
- [MRSV99] Donatella Merlini, Douglas G. Rogers, Renzo Sprugnoli, and M. Cecilia Verri. [Underdiagonal lattice paths with unrestricted steps](#). *Discrete Appl. Math.*, 91(1-3):197–213, 1999.
- [Wal16] Michael Wallner. [A half-normal distribution scheme for generating functions](#). *arXiv preprint*, arXiv:1610.00541, 2016.