induced magnetization in the present field and, second, that it could be a viscous remanent magnetization (VRM) acquired during prolonged exposure of the magnetic minerals to the planetary field and hence reflecting an unknown, but younger, age than that of the smooth plains. Although both of these physical processes are likely to operate, induced magnetizations cannot fully explain the observed HPF field strengths, and the net effect of VRM will be that our estimates of ancient field strength are lower bounds (10).

Within the range of uncertainty of crustal thickness (22–24) and magnetized layer source depths (10), most or all of the magnetization could reside within Mercury's crust (Fig. 4). We investigated whether such a scenario is consistent with thermal evolution models, given magnetizations acquired at ~4 Ga. We estimated the depth to T_c for a range of thermal gradients (Fig. 4). The Curie temperature was taken to be 325°C (that of pyrrhotite) as a conservatively low value for our calculations, and we used the maximum average daily surface temperature predicted for a range of Mercury's orbital eccentricities from 0 to 0.4 (10, 25). The results indicate that even for high thermal gradients at 4 Ga (26) the depth to T_c in the Suisei Planitia region is at least 20 km. For thermal gradients less than 8 K/km and upper limits on the crustal thickness in the region, the entire crust remains below T_c . These results imply that acquisition and subsequent preservation of an ancient crustal remanence by magnetic carriers with T_c values of at least 325°C are consistent with thermal models (10, 26–28), and for carriers with higher T_c some remanence may be carried by upper mantle material. Such a conclusion is predicated on the assumption that the surface temperature pole locations have remained stationary in a body-fixed coordinate system since the time that the remanent magnetization was acquired (10). The symmetry of the ancient field with respect to the present rotation axis supports such a presumption by suggesting that, since that epoch, there has been no substantial reorientation of the crust ("true polar wander") with respect to the planet's axis of greatest moment of inertia.

The simplest interpretation of the results presented here is that a core dynamo was present early in Mercury's history. If the dynamo was thermochemically driven [e.g., (6, 29)], this finding provides a strong constraint on models for the thermal evolution of Mercury's interior. In particular, the existence of a core dynamo at the time of smooth plains emplacement presents a new challenge to such models. An early core dynamo can be driven by superadiabatic cooling of the liquid core, but in typical thermal history models this phase has ended by 3.9 Ga. A later dynamo can be driven by the combined effects of cooling and compositional convection associated with formation of a solid inner core (26–28), but in most thermal history models inner core formation does not start until well after 3.7 Ga. Further progress in understanding the record of Mercury's ancient field can also be made with improved petrological constraints on crustal compositions [e.g., (30)], information

on the candidate magnetic mineralogies implied, and knowledge of their magnetic properties.

REFERENCES AND NOTES

- N. F. Ness, K. W. Behannon, R. P. Lepping, Y. C. Whang, K. H. Schatten, *Science* **185**, 151–160 (1974).
- B. J. Anderson *et al.*, *Science* **321**, 82–85 (2008).
- B. J. Anderson *et al.*, *Science* **333**, 1859–1862 (2011).
- B. J. Anderson *et al.*, *J. Geophys. Res.* **117** (E12), E00L12 (2012).
- C. L. Johnson *et al.*, *J. Geophys. Res.* **117** (E12), E00L14 (2012).
- H. Cao *et al.*, *Geophys. Res. Lett.* **41**, 4127–4134 (2014).
- L. C. Philpott *et al.*, *Geophys. Res. Lett.* **41**, 6627–6634 (2014).
- B. J. Anderson *et al.*, *Geophys. Res. Lett.* **41**, 7444–7452 (2014).
- H. Korth *et al.*, Modular model for Mercury's magnetospheric magnetic field confined within the average observed magnetopause. JGR Space Physics, doi:10.1002/2015JA021022 (2015).
- Materials and methods are available as supplementary materials on Science Online.
- B. J. Anderson, C. L. Johnson, H. Korth, *Geochim. Geophys. Geosyst.* **14**, 3875–3886 (2013).
- B. W. Denevi *et al.*, *J. Geophys. Res. Planets* **118**, 891–907 (2013).
- J. L. Whitten, J. W. Head, B. W. Denevi, S. C. Solomon, *Icarus* **241**, 97–113 (2014).
- P. K. Byrne *et al.*, *Nat. Geosci.* **7**, 301–307 (2014).
- S. Marchi *et al.*, *Nature* **499**, 59–61 (2013).
- M. E. Purucker, J. B. Nicholas, *J. Geophys. Res.* **115** (E12), E12007 (2010).
- M. A. Wieczorek, B. P. Weiss, S. T. Stewart, *Science* **335**, 1212–1215 (2012).
- L. R. Nittler *et al.*, *Science* **333**, 1847–1850 (2011).
- F. M. McCubbin, M. A. Riner, K. E. Vander Kaaden, L. K. Burkemper, *Geophys. Res. Lett.* **39**, L09202 (2012).
- L. G. Evans *et al.*, *J. Geophys. Res.* **117** (E12), E00L07 (2012).
- S. Z. Weider, L. R. Nittler, R. D. Starr, T. J. McCoy, S. C. Solomon, *Icarus* **235**, 170–186 (2014).
- E. Mazarico *et al.*, *J. Geophys. Res. Planets* **119**, 2417–2436 (2014).
- P. B. James, M. T. Zuber, R. J. Phillips, S. C. Solomon, *J. Geophys. Res. Planets* **120**, 287–310 (2015).
- S. Padovan, M. A. Wieczorek, J.-L. Margot, N. Tosi, S. C. Solomon, *Geophys. Res. Lett.* **42**, 1029–1038 (2015).
- A. C. M. Correia, J. Laskar, *Icarus* **201**, 1–11 (2009).
- J.-P. Williams, O. Aharonson, F. Nimmo, *Geophys. Res. Lett.* **34**, L21201 (2007).
- M. Grott, D. Breuer, M. Laneuville, *Earth Planet. Sci. Lett.* **307**, 135–146 (2011).
- N. Tosi, M. Grott, A.-C. Plesa, D. Breuer, *J. Geophys. Res. Planets* **118**, 2474–2487 (2013).
- S. Stanley, G. A. Glatzmaier, *Space Sci. Rev.* **152**, 617–649 (2010).
- K. E. Vander Kaaden, F. M. McCubbin, *J. Geophys. Res. Planets* **120**, 195–209 (2015).

ACKNOWLEDGMENTS

We thank the MESSENGER operations and engineering teams for enabling the low-altitude observations reported here. We are also grateful for the contributions of our friend and colleague M. H. Acuña whose expertise was critical to the Magnetometer development. The MESSENGER mission is supported by the NASA Discovery Program and the MESSENGER Participating Scientist Program. C.L.J. and L.C.P. also acknowledge support from the Natural Sciences and Engineering Research Council of Canada. Data from the MESSENGER mission are archived with the NASA Planetary Data System. We thank three reviewers for thoughtful comments that improved the manuscript.

SUPPLEMENTARY MATERIALS

www.sciencemag.org/content/348/6237/892/suppl/DC1
Supplementary Text
Figs. S1 to S11
Tables S1 and S2
References (31–53)

7 February 2015; accepted 28 April 2015
Published online 7 May 2015;
10.1126/science.aaa8720

CARBON CYCLE

The dominant role of semi-arid ecosystems in the trend and variability of the land CO₂ sink

Anders Ahlström,^{1,2*} Michael R. Raupach,^{3†} Guy Schurgers,⁴ Benjamin Smith,¹ Almut Arneth,⁵ Martin Jung,⁶ Markus Reichstein,⁶ Josep G. Canadell,⁷ Pierre Friedlingstein,⁸ Atul K. Jain,⁹ Etsushi Kato,¹⁰ Benjamin Poulter,¹¹ Stephen Sith,¹² Benjamin D. Stocker,^{13,14} Nicolas Viovy,¹⁵ Ying Ping Wang,¹⁶ Andy Wiltshire,¹⁷ Sönke Zaehle,⁶ Ning Zeng¹⁸

The growth rate of atmospheric carbon dioxide (CO₂) concentrations since industrialization is characterized by large interannual variability, mostly resulting from variability in CO₂ uptake by terrestrial ecosystems (typically termed carbon sink). However, the contributions of regional ecosystems to that variability are not well known. Using an ensemble of ecosystem and land-surface models and an empirical observation-based product of global gross primary production, we show that the mean sink, trend, and interannual variability in CO₂ uptake by terrestrial ecosystems are dominated by distinct biogeographic regions. Whereas the mean sink is dominated by highly productive lands (mainly tropical forests), the trend and interannual variability of the sink are dominated by semi-arid ecosystems whose carbon balance is strongly associated with circulation-driven variations in both precipitation and temperature.

Since the 1960s, terrestrial ecosystems have acted as a substantial sink for atmospheric CO₂, sequestering about one-quarter of anthropogenic emissions in an average year (1). This ecosystem service, which helps mitigate climate change by reducing the rate of increase of atmospheric greenhouse gases, is due to

an imbalance between the uptake of CO₂ through gross primary production (GPP, the aggregate photosynthesis of plants) and the release of carbon to the atmosphere by ecosystem respiration (R_{eco}) and other losses, including wildfires (C_{fire}). The net carbon flux (net biome production, NBP = GPP – R_{eco} – C_{fire}) results from the small

imbalance between the much larger uptake and release fluxes. Consequently, small fractional variations in either of these fluxes can cause substantial absolute variations in net carbon exchange with the atmosphere. These variations account almost entirely for year-to-year variations around the overall trend in atmospheric concentrations of CO₂ (2, 3).

Modeling studies suggest a large uncertainty of the future magnitude and sign of the carbon sink provided by terrestrial ecosystems (4–8). Robust projections are crucial to assessments of future atmospheric CO₂ burdens and associated climate change, and are therefore central to the effectiveness of future mitigation policies. Reducing the uncertainty of these projections requires better knowledge of the regions and processes governing the present sink and its variations. Inventories suggest that since the beginning of industrialization, the majority of carbon sequestered by the terrestrial biosphere has accumulated in forest ecosystems of the tropics and temperate zones (9). However, the relative contributions of ecosystems of different, climatically distinct, regions to variations in the land sink on interannual to multidecadal time scales are not well characterized. Here, we investigated relative regional contributions to the mean sink, to its trend over recent decades, and to the interannual variability (IAV) around the trend.

We used LPJ-GUESS (10–12), a biogeochemical dynamic global vegetation model, to simulate the geographic pattern and time course of NBP. LPJ-GUESS explicitly accounts for the dependency of plant production and downstream ecosystem processes on the demography (size structure) and composition of simulated vegetation. We forced

the model with historical climate (13) and CO₂ concentrations, accounting for emissions from land use change and carbon uptake due to regrowth after agricultural abandonment (14). We compared the results to an ensemble of nine ecosystem and land surface model simulations from the TRENDY model intercomparison project (12, 15) (hereinafter TRENDY models; table S1). The TRENDY ensemble is similarly based on historical climate and CO₂ but uses a static 1860 land use mask.

Global NBP, as simulated by LPJ-GUESS, shows strong agreement ($r^2 = 0.62$) with the Global Carbon Project (GCP) estimate of the net land CO₂ flux—an independent, bookkeeping-based estimate derived as the residual of emissions, atmospheric growth, and ocean uptake of CO₂ (1) (Fig. 1A). TRENDY models do not account for land use change. Relative to the GCP land flux estimate, they consequently predict a higher average NBP but similar interannual variation. Moreover, the offset between the TRENDY model

ensemble mean and the GCP land flux estimate is comparable to the GCP estimate of mean land use change emission flux for the period 1982–2011 (fLUC).

We divided the global land area into six land cover classes, following the MODIS MCD12C1 land cover classification (12, 16): tropical forests (Fig. 1B), extratropical forest, grasslands and croplands (here combined), semi-arid ecosystems (Fig. 1C), tundra and arctic shrub lands, and sparsely vegetated lands (areas classified as barren) (figs. S1 and S2).

When the global terrestrial CO₂ sink (average NBP and its trend (1982–2011) are partitioned among land cover classes, we find that tropical forests account for the largest fraction (26%, 0.33 Pg C year⁻¹) of the average sink over this period (1.23 Pg C year⁻¹) (Fig. 1D). In contrast, we find that semi-arid ecosystems dominate the positive global CO₂ sink trend (57%, 0.04 Pg C year⁻²; global, 0.07 Pg C year⁻²) (Fig. 1E). The TRENDY model ensemble shows a consistent pattern, with

Fig. 1. Global and regional NBP mean, trend, and variations (1982–2011). (A) Global NBP from LPJ-GUESS (red line) and GCP land flux time series (black line) with ± 0.8 Pg C uncertainty range (shaded gray area). TRENDY models mean (blue line) and first and third quartiles (shaded blue area) are plotted on a separate axis with a time-invariant offset corresponding to the time period average GCP fLUC estimate (1.2 Pg C year⁻¹). (B) Tropical forest NBP. LPJ-GUESS (red line) includes emissions from land use change. TRENDY models average (blue line) and first and third quartiles of the ensemble (shaded blue area) do not include emissions from land use change. (C) NBP of semi-arid ecosystems from LPJ-GUESS (including land use change emissions) and TRENDY models (excluding land use change emissions); colors and shading as in (B). (D) Contribution of land cover classes to global mean NBP (1982–2011) (mean NBP of land cover class as a proportion of mean global NBP). Horizontal lines in box plots show, from top to bottom, 95th, 75th, 50th, 25th, and 5th percentiles. (E) Contribution of land cover classes to global NBP trend (land cover class NBP trend as a proportion of global NBP trend). (F) Contribution of land cover classes to global NBP IAV (Eq. 1).

¹Department of Physical Geography and Ecosystem Science, Lund University, 223 62 Lund, Sweden. ²Department of Earth System Science, School of Earth, Energy and Environmental Sciences, Stanford University, Stanford, CA 94305, USA. ³Climate Change Institute, Australian National University, Canberra, ACT 0200, Australia. ⁴Department of Geosciences and Natural Resource Management, University of Copenhagen, 1350 Copenhagen, Denmark. ⁵Institute for Meteorology and Climate Research—Atmospheric Environmental Research, Karlsruhe Institute for Technology, 82476 Garmisch-Partenkirchen, Germany. ⁶Biogeochemical Intergration Department, Max Planck Institute for Biogeochemistry, 07745 Jena, Germany. ⁷Global Carbon Project, CSIRO Oceans and Atmospheric Flagship, Canberra, ACT, Australia. ⁸College of Engineering, Mathematics and Physical Sciences, University of Exeter, Exeter EX4 4QF, UK. ⁹Department of Atmospheric Sciences, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA. ¹⁰Institute of Applied Energy, 105-0003 Tokyo, Japan. ¹¹Institute on Ecosystems and the Department of Ecology, Montana State University, Bozeman, MT 59717, USA. ¹²College of Life and Environmental Sciences, University of Exeter, Exeter EX4 4RJ, UK. ¹³Department of Life Sciences, Imperial College, Ascot SL5 7PY, UK. ¹⁴Climate and Environmental Physics, Physics Institute and Oeschger Centre for Climate Change Research, University of Bern, Bern, Switzerland. ¹⁵Laboratoire des sciences du climat et de l'environnement, CEA Saclay, F-91191 Gif-sur-Yvette Cedex, France. ¹⁶CSIRO Ocean and Atmosphere Flagship, PMB 1, Aspendale, Victoria 3195, Australia. ¹⁷Met Office Hadley Centre, Fitzroy Road, Exeter EX1 3PB, UK. ¹⁸Department of Atmospheric and Oceanic Science and Earth System Science Interdisciplinary Center, University of Maryland, College Park, MD 20742, USA. *Corresponding author. E-mail: anders.ahlgren@nateko.lu.se †Deceased.

tropical forests dominating the mean sink (median 24%) and semi-arid ecosystems dominating the trend (median 51%). The predominance of semi-arid ecosystems in explaining the global land sink trend is consistent with widespread observations of woody encroachment over semi-arid areas (17) and increased vegetation greenness inferred from

satellite remote sensing over recent decades (17–19). Likewise, a recent study attributes the majority of the record land sink anomaly of 2011 to the response of semi-arid ecosystems in the Southern Hemisphere, Australia in particular, to an anomalous wet period; the study further postulates a recent increase in the sensitivity of carbon uptake

to precipitation for this region, which is attributed to vegetation expansion (20).

We further partitioned IAV in global NBP among land cover classes according to the contribution of individual regions (grid cells or land cover classes) to global NBP IAV (12). To this end, we adopted an index that scores individual geographic locations according to the consistency, over time, with which the local NBP flux resembles the sign and magnitude of global NBP (fig. S4):

$$f_j = \frac{\sum_t \frac{x_{jt} |X_t|}{|X_t|}}{\sum_t |X_t|} \quad (1)$$

where x_{jt} is the flux anomaly (departure from a long-term trend) for region j at time t (in years), and X_t is the global flux anomaly, so that $X_t = \sum_j x_{jt}$. By this definition f_j is the average relative anomaly x_{jt}/X_t for region j , weighted with the absolute global anomaly $|X_t|$. Regions receiving higher and positive average scores are inferred to have a larger contribution in governing global NBP IAV, as opposed to regions characterized by smaller or negative (counteracting) scores (fig. S3). The index we adopt does not characterize the variability of ecosystems of different regions, as, for example, the standard deviation would do (fig. S5); rather, it enables a comparison of their relative importance (contribution) in governing global IAV.

Semi-arid ecosystems were found to account for the largest fraction, 39%, of global NBP IAV, exceeding tropical forest (19%), extratropical forest (11%; all forest, 30%), and grasslands and croplands (27%) (Fig. 1F). The TRENDY model ensemble shows a similar partitioning, with semi-arid ecosystems accounting for 47% (median; tropical forests, 28%; extratropical forest, 6%; all forest, 35%). The overall contributions per land cover class are the sum of both positive and negative contributions that result from differences

Fig. 2. Climatic covariates of semi-arid ecosystem GPP variations. (A) Distribution by latitude of the empirical GPP product anomalies normalized by average standard deviation of GPP in semi-arid lands. The distribution is colored according to average local climatic covariates per latitude zone and distribution bin. (B) LPJ-GUESS GPP distribution calculated and colored as in (A). (C) Covariation of the multivariate ENSO index [MEI (31, 32)] anomalies with the empirical GPP product. (D) Covariation of MEI and modeled GPP anomalies per latitudinal zone. Note that the figure shows the covariates of latitudinal average local GPP anomalies, and not the average covariates based on GPP IAV contribution to NBP IAV.

Fig. 3. Climatic covariates of NBP extremes. (A) Climatic covariates of LPJ-GUESS negative NBP extremes (1st to 10th percentiles). (B) Mean climatic covariates of TRENDY models' negative NBP extremes (1st to 10th percentiles). (C) Covariates of LPJ-GUESS positive NBP extremes (90th to 99th percentiles). (D) Mean climatic covariates of TRENDY models' positive NBP extremes (90th to 99th percentiles).

in phase between IAV of individual grid cells compared with global IAV (fig. S4). The extent to which negative contributions reduce the overall land cover class contributions is minor for all regions except grasslands and croplands (fig. S6) (LPJ-GUESS, -13%; TRENDY median, -13%) because the latter are distributed widely across climate zones, and because both climate variations and the sensitivity of NBP to climate variations differ among regions.

To partition the global NBP IAV among component fluxes (GPP, R_{eco} , C_{fire}) and among land cover classes, we applied Eq. 1. We found that global NBP IAV is most strongly associated with variation in GPP; interannual GPP anomalies contribute 56% of the global NBP IAV in LPJ-GUESS and a median of 90% in the TRENDY model ensemble. Comparing different land cover classes, the GPP anomalies of semi-arid ecosystems alone contribute 39% in LPJ-GUESS and a median of 65% in the TRENDY model ensemble to global NBP IAV (fig. S7). Semi-arid vegetation productivity thus emerges clearly as the single most important factor governing global NBP IAV.

We used two complementary methods to attribute the variability in GPP—as the inferred primary driver of global NBP IAV—to its environmental drivers. First, we analyzed simulation results from LPJ-GUESS, linking output GPP anomalies to variability in the climatic input data. Second, we used a time-resolved gridded global GPP product derived from upscaled flux tower measurements (12, 21) (hereafter, empirical GPP product). This product uses an empirical upscaling of flux measurements and is thus entirely independent of the modeled GPP in our study.

The three main climatic drivers—temperature (T), precipitation (P), and shortwave radiation (S)—are interdependent and correlated. To account for the combined effects of these drivers, we adopted an analysis of GPP variations from an “impact perspective” (22–24): We first identified GPP anomalies and then extracted their climatic covariates. The primary challenge of such an analysis on an annual scale is to target climate indices that adequately characterize the “period of climatic influence” (e.g., growing season average, annual averages, minima or maxima of a given climatic forcing). To overcome this challenge, we used semiannual time series of climate drivers constructed via an optimization procedure that weights monthly anomalies of a given climate variable (T, P, or S), accounting for time lags of up to 24 months while making no additional prior assumptions as to the period of influence (12). For each GPP event, we extracted climatic covariates as z scores of the semiannual climatic drivers.

We evaluated the climatic covariates of GPP anomalies for semi-arid ecosystems from the empirical GPP product and modeled by LPJ-GUESS, focusing on T and P, and found similar responses of GPP to climate with both approaches across all latitude bands (Fig. 2, A and B). Negative GPP anomalies in semi-arid ecosystems are mainly driven by warm and dry (low rainfall) climatic events in most latitudes, suggestive of drought. By contrast, positive GPP anomalies are domi-

nated by cool and wet conditions. Averaging the distributions over latitudes (Fig. 2, A and B) and extracting the climatic covariates per percentile of the GPP distributions shows that GPP varies with climatic conditions on a straight line in T-P space (fig. S8), with a stronger covariation with P than with T. This implies that the full GPP distributions are driven by similar climatic patterns—that is, anomalies that differ in size and sign covary with corresponding differences in size and sign in the drivers. GPP extremes (the tails of the distribution of GPP among years) covary with El Niño–Southern Oscillation (ENSO) across all latitudes (Fig. 2, C and D). In both the model and the empirical GPP product, GPP anomalies are more strongly associated with the positive phase of ENSO (El Niño) than with the negative phase (La Niña); the sign of the relationship varies with latitude. Positive ENSO tends to coincide with negative GPP anomalies in the tropics (30°S to 20°N) and with positive GPP anomalies north of 20°N.

The agreement between climatic covariates of the data-based empirical GPP product and modeled GPP alongside the comparatively robust pattern of the covariation with climate suggests that GPP IAV for semi-arid ecosystems is mediated by climate. Because ENSO covaries with a considerable portion of the GPP distribution, we infer that ENSO is the dominating mode of global circulation variations driving GPP IAV over semi-arid ecosystems. Recent modeling studies have found that extreme El Niño events could become more common under climate change (25), which, together with an increased atmospheric demand for water associated with global warming, might exacerbate the impact of El Niño events over semi-arid ecosystems and further increase the role of semi-arid regions in driving global NBP IAV (26–28).

We repeated the calculation of climatic covariates to simulated NBP for LPJ-GUESS and each

of the TRENDY models. The resulting maps of covariates in T-P space are shown as average covariates of negative NBP extremes (Fig. 3, A and B) and positive NBP extremes (Fig. 3, C and D). In general, semi-arid ecosystems stand out as regions in which strong CO₂ uptake events are consistently associated with cool and moist conditions, and strong CO₂ release events with warm and dry conditions. In tropical forests, NBP covaries with both T and P as in semi-arid regions, but also with T alone. In high latitudes, wet or warm and wet conditions lead to negative NBP extremes, whereas dry or warm and dry conditions tend to lead to positive extremes, although the spatial heterogeneity of the covariates is large in this region (Fig. 3).

Our approach offers detailed spatial and temporal disaggregation of drivers and responses, which is important when analyzing drivers or covariates of global NBP IAV because of the high temporal and spatial variability in P (figs. S9 to S11). Using four upscaling levels with increasing spatial and temporal disaggregation [ranging from land surface mean P and T to semiannual P and T, averaged according to the spatial origin of each year's global NBP anomaly (eqs. S5 and S6)], we found that P and NBP IAV become more correlated at higher levels of disaggregation. At the highest disaggregation level, P is almost as strongly correlated with NBP IAV as T, suggesting a strong influence of soil moisture variations on global NBP IAV (28). This strong increase in P correlations with disaggregation resolves an apparent conflict between our findings and those of studies using regionally averaged drivers that emphasize the role of T in governing IAV in atmospheric CO₂ (28–30). For semi-arid ecosystems, T correlations with NBP IAV are slightly stronger than P correlations with NBP IAV (Fig. 4B), partly because of an asymmetric distribution of P and/or an asymmetric response of NBP to P IAV (fig. S12). The correlation of tropical forest P with

Fig. 4. Correlations between climatic drivers IAV (P and T) and global NBP IAV (mean of all 10 models). (A) Global P and T correlations to global NBP IAV. From black to white and left to right, bars represent annual P and T IAV correlations to global NBP IAV with increasing spatial and temporal disaggregation of P and T while averaging to global time series. Black bars represent averaged global land surface P and T weighted by grid cell area; dark gray bars represent P and T weighted by 30-year average contribution to global NBP IAV (Eq. 1 and fig. S4); light gray bars represent averaged P and T weighted by each year's contributions, thus accounting for the difference in the spatial distribution of contributions between years (eqs. S5 and S6); white bars represent semiannual climate drivers averaged to global time series using the annual spatial contributions (as for light gray bars), thereby accounting for the “period of climatic influence” and time lags of up to 24 months. (B) Correlations between P and T IAV and global NBP IAV for semi-arid ecosystems. Weights, where applicable, are based on contributions to global NBP IAV as in (A) but with P and T averaged over semi-arid ecosystems only. (C) Correlations between P and T IAV and global NBP IAV for tropical forest. Weights, where applicable, are based on contributions to global NBP IAV as in (A) but with P and T averaged over tropical forest only.

NBP IAV increases when we use the semiannual drivers, which suggests the importance of accounting for time lags and the “period of climatic influence” of P variations (12), but P correlations with NBP IAV are still weaker than T correlations with NBP IAV (Fig. 4C).

Our analysis provides evidence that semi-arid ecosystems, largely occupying low latitudes, have dominated the IAV and trend of the global land carbon sink over recent decades. Semi-arid regions have been the subject of relatively few targeted studies that place their importance in a global context. Our findings indicate that semi-arid regions and their ecosystems merit increased attention as a key to understanding and predicting interannual to decadal variations in the global carbon cycle.

REFERENCES AND NOTES

- C. Le Quéré et al., *Earth Syst. Sci. Data* **6**, 235–263 (2014).
- C. D. Keeling, T. P. Whorf, M. Wahlen, J. van der Plicht, *Nature* **375**, 666–670 (1995).
- C. Le Quéré et al., *Nat. Geosci.* **2**, 831–836 (2009).
- A. Ahlström, G. Schurgers, A. Arneth, B. Smith, *Environ. Res. Lett.* **7**, 044008 (2012).
- P. Friedlingstein et al., *J. Clim.* **19**, 3337–3353 (2006).
- A. D. McGuire et al., *Glob. Biogeochem. Cycles* **15**, 183–206 (2001).
- S. Schaphoff et al., *Clim. Change* **74**, 97–122 (2006).
- S. Sitch et al., *Glob. Change Biol.* **14**, 2015–2039 (2008).
- Y. Pan et al., *Science* **333**, 988–993 (2011).
- A. Ahlström, P. A. Miller, B. Smith, *Geophys. Res. Lett.* **39**, L15403 (2012).
- B. Smith, I. C. Prentice, M. T. Sykes, *Glob. Ecol. Biogeogr.* **10**, 621–637 (2001).
- See supplementary materials on Science Online.
- I. Harris, P. D. Jones, T. J. Osborn, D. H. Lister, *Int. J. Climatol.* **34**, 623–642 (2014).
- G. Hurrell et al., *Clim. Change* **109**, 117–161 (2011).
- S. Sitch et al., *Biogeosciences* **12**, 653–679 (2015).
- M. A. Friedl et al., *Remote Sens. Environ.* **114**, 168–182 (2010).
- N. Andela, Y. Y. Liu, A. I. J. M. van Dijk, R. A. M. de Jeu, T. R. McVicar, *Biogeosciences* **10**, 6657–6676 (2013).
- R. J. Donohue, T. R. McVicar, M. L. Roderick, *Glob. Change Biol.* **15**, 1025–1039 (2009).
- R. Fensholt et al., *Remote Sens. Environ.* **121**, 144–158 (2012).
- B. Poulter et al., *Nature* **509**, 600–603 (2014).
- M. Jung et al., *J. Geophys. Res.* **116**, G00J07 (2011).
- J. Zscheischler et al., *Environ. Res. Lett.* **9**, 035001 (2014).
- M. Reichstein et al., *Nature* **500**, 287–295 (2013).
- M. D. Smith, *J. Ecol.* **99**, 656–663 (2011).
- W. Cai et al., *Nat. Clim. Change* **4**, 111–116 (2014).
- K. E. Trenberth et al., *Nat. Clim. Change* **4**, 17–22 (2014).
- A. Dai, *Nat. Clim. Change* **3**, 52–58 (2013).
- X. Wang et al., *Nature* **506**, 212–215 (2014).
- W. Wang et al., *Proc. Natl. Acad. Sci. U.S.A.* **110**, 13061–13066 (2013).
- P. M. Cox et al., *Nature* **494**, 341–344 (2013).
- K. Wolter, M. S. Timlin, in *Proceedings of the 17th Climate Diagnostics Workshop* (University of Oklahoma, Norman, OK, 1993), pp. 52–57; www.esrl.noaa.gov/psd/enso/mei/WT1.pdf.
- K. Wolter, M. S. Timlin, *Weather* **53**, 315–324 (1998).

ACKNOWLEDGMENTS

This paper is dedicated to the memory of Michael Robin Raupach (1950–2015), whose scientific integrity and novel contributions leave a long-lasting legacy in the field of carbon cycle sciences. The MODIS MOD12C1 land cover product was obtained through the online Data Pool at the NASA Land Processes Distributed Active Archive Center (LP DAAC), USGS/Earth Resources Observation and Science (EROS) Center, Sioux Falls, South Dakota (https://lpdaac.usgs.gov/data_access). Supported by the Royal Physiographic Society in Lund (Birgit and Hellmuth Hertz

Foundation), Swedish Research Council grant 637-2014-6895, and the Mistra-SWECIA program (A. Ahlström); EC FP7 grant LUC4C (603542) (A. Arneth); OCE Distinguished Visiting Scientist to the CSIRO Ocean and Atmosphere Flagship, Canberra (B.S.); EC FP7 grant EMBRACE (282672) (A. Arneth, M.R., and B.D.S.); the Australian Climate Change Science Program (J.G.C.); NSF grant AGS 12-43071, U.S. Department of Energy grant DE-SC0006706, and NASA LCLUC program grant NNX14AD94G (A.K.J.); the Environmental Research and Technology Development Fund (S-10) of the Ministry of Environment of Japan (E.K.); CSIRO strategic research funds (Y.P.W.); and NOAA grants NA100AR4310248 and NA09NES4400006 and NSF grant AGS-1129088 (N.Z.). This

study is a contribution to the Lund Centre for Studies of Carbon Cycle and Climate Interactions (LUCCI) and the strategic research areas MERGE and BECC.

SUPPLEMENTARY MATERIALS

www.sciencemag.org/content/348/6237/895/suppl/DC1
Materials and Methods
Figs. S1 to S12
References (33–56)

26 October 2014; accepted 24 April 2015
10.1126/science.aaa1668

GLACIER MASS LOSS

Dynamic thinning of glaciers on the Southern Antarctic Peninsula

B. Wouters,^{1*} A. Martin-Español,¹ V. Helm,² T. Flament,³ J. M. van Wessem,⁴ S. R. M. Ligtenberg,⁴ M. R. van den Broeke,⁴ J. L. Bamber¹

Growing evidence has demonstrated the importance of ice shelf buttressing on the inland grounded ice, especially if it is resting on bedrock below sea level. Much of the Southern Antarctic Peninsula satisfies this condition and also possesses a bed slope that deepens inland. Such ice sheet geometry is potentially unstable. We use satellite altimetry and gravity observations to show that a major portion of the region has, since 2009, destabilized. Ice mass loss of the marine-terminating glaciers has rapidly accelerated from close to balance in the 2000s to a sustained rate of -56 ± 8 gigatons per year, constituting a major fraction of Antarctica's contribution to rising sea level. The widespread, simultaneous nature of the acceleration, in the absence of a persistent atmospheric forcing, points to an oceanic driving mechanism.

Ice shelves have been identified as sensitive indicators of climate change (1). Their retreat along the coast of the Northern Antarctic Peninsula has been noted over recent decades (2) and associated with a sudden and prolonged increase in discharge of the inland grounded ice (3–5), especially for those glaciers overlying deep troughs (6). The potential future contribution to sea-level rise of these glaciers relatively modest because their catchments are small compared with those further south (7). The Southern Antarctic Peninsula (SAP), including Palmer Land and the Bellinghausen Coast, rests on bedrock below sea level with a retrograde slope (deeper inland) (8), which is believed to be an inherently unstable configuration (9), permitting rapid grounding line retreat and mass loss to the ocean. Recent modeling results suggest that this marine ice sheet instability may have already been initiated for part of West Antarctica (10, 11).

The SAP is home to a number of fast flowing, marine terminating glaciers, many of which are still unnamed. Laser [ICESat, 2003–2009 (12)] and radar [Envisat, 2003–2010 (13)] altimetry identified moderate surface-lowering con-

centrated within a narrow strip along the coast, in particular near the grounding line of the Ferrigno Ice Stream (14), contrasted by widespread thickening further inland. Observations from the Gravity Recovery and Climate Experiment (GRACE) mission show that these opposing signals compensated each other, resulting in a near-zero mass balance for 2002–2010 (15).

The Cryosat-2 satellite, launched in April 2010, provides elevation measurements of land and sea ice at a high spatial resolution up to a latitude of 88°. In contrast to conventional altimetry missions such as Envisat, Cryosat-2's dual antenna and Doppler processing results in improved resolution and geolocation of the elevation measurement (16). Because of the long satellite repeat period of 369 days, it has a dense track spacing in our region of interest, which is a major advantage compared with the roughly 10-times-coarser ICESat track spacing. Two recent studies using Cryosat-2 data observed thinning along the coast of the Bellinghausen Sea (17, 18). Such elevation changes may result from either a decrease in surface mass balance (SMB) (accumulation minus ablation), compaction of the firn column, or an increase in the ice flow speed (also termed dynamic thinning). Both studies attributed the surface-lowering to interannual changes in SMB, based on the strong accumulation variability observed in the Gomez ice core (70.36°W, 73.59°S) (18, 19). Here, we take SMB and firn compaction into account and show that the

¹Bristol Glaciology Centre, University of Bristol, Bristol, UK.

²Alfred-Wegener-Institut Helmholtz-Zentrum für Polar- und Meeresforschung, Bremerhaven, Germany. ³Laboratoire d'Études en Géophysique et Océanographie Spatiales, Toulouse, France. ⁴Institute for Marine and Atmospheric Research, Utrecht University, Netherlands.

*Corresponding author. E-mail: bert.wouters@bristol.ac.uk

The dominant role of semi-arid ecosystems in the trend and variability of the land CO₂ sink

Anders Ahlström, Michael R. Raupach, Guy Schurgers, Benjamin Smith, Almut Arneth, Martin Jung, Markus Reichstein, Josep G. Canadell, Pierre Friedlingstein, Atul K. Jain, Etsushi Kato, Benjamin Poulter, Stephen Sitch, Benjamin D. Stocker, Nicolas Viovy, Ying Ping Wang, Andy Wiltshire, Sönke Zaehle and Ning Zeng

Science **348** (6237), 895-899.
DOI: 10.1126/science.aaa1668

The difference is found at the margins

The terrestrial biosphere absorbs about a quarter of all anthropogenic carbon dioxide emissions, but the amount that they take up varies from year to year. Why? Combining models and observations, Ahlström *et al.* found that marginal ecosystems—semiarid savannas and low-latitude shrublands—are responsible for most of the variability. Biological productivity in these semiarid regions is water-limited and strongly associated with variations in precipitation, unlike wetter tropical areas. Understanding carbon uptake by these marginal lands may help to improve predictions of variations in the global carbon cycle.

Science, this issue p. 895

ARTICLE TOOLS

<http://science.sciencemag.org/content/348/6237/895>

SUPPLEMENTARY MATERIALS

<http://science.sciencemag.org/content/suppl/2015/05/20/348.6237.895.DC1>

REFERENCES

This article cites 54 articles, 2 of which you can access for free
<http://science.sciencemag.org/content/348/6237/895#BIBL>

PERMISSIONS

<http://www.sciencemag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)

Science (print ISSN 0036-8075; online ISSN 1095-9203) is published by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. The title *Science* is a registered trademark of AAAS.

Copyright © 2015, American Association for the Advancement of Science