Development and physiology of the brown alga *Ectocarpus siliculosus*: two centuries of research

Bénédicte Charrier, Susana Coelho, Aude Le Bail, Thierry Tonon, Gurvan Michel, Philippe Potin, Bernard Kloareg, Catherine Boyen, Akira F Peters, J. Mark Cock

► To cite this version:

Bénédicte Charrier, Susana Coelho, Aude Le Bail, Thierry Tonon, Gurvan Michel, et al.. Development and physiology of the brown alga *Ectocarpus siliculosus*: two centuries of research. *New Phytologist*, 2008, 177 (2), pp.319-332. 10.1111/j.1469-8137.2007.02304.x . hal-01806426

HAL Id: hal-01806426

<https://hal.science/hal-01806426>

Submitted on 16 Nov 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Development and physiology of the brown alga *Ectocarpus siliculosus*:
two centuries of research**

Journal:	<i>New Phytologist</i>
Manuscript ID:	NPH-TR-2007-05852.R1
Manuscript Type:	TR - Commissioned Material - Tansley Review
Date Submitted by the Author:	19-Sep-2007
Complete List of Authors:	Charrier, Benedicte; CNRS, Station Biologique Coelho, Susana; CNRS, Station Biologique Le Bail, Aude; CNRS, Station Biologique Tonon, Thierry; CNRS, Station Biologique Michel, Gurvan; CNRS, Station Biologique Potin, Philippe; CNRS, Station Biologique Kloareg, Bernard; CNRS, Station Biologique Boyen, Catherine; CNRS, Station Biologique Peters, Akira; CNRS, Station Biologique
Key Words:	brown alga , defense, development , Ectocarpus , metabolism, reproduction, pheophyceae , pheromone

Development and physiology of the brown alga *Ectocarpus siliculosus*: two centuries of research

Bénédicte Charrier*, Susana M. Coelho, Aude Le Bail, Thierry Tonon, Gurvan Michel, Philippe Potin, Bernard Kloareg, Catherine Boyen, Akira F. Peters, J. Mark Cock.

UMR7139, Station Biologique, Centre National de la Recherche Scientifique, Université Pierre & Marie Curie Paris VI, Place Georges Teissier, 29 682 Roscoff cedex, France.

* author for correspondence. Tel :33-2-98-29-23-62 ; fax : 33-2-98-29-23-24 ; email : charrier@sb-roscoff.fr

Nber of figures : 5

Word number :

- I- Introduction : 564
- II – Diversity and taxonomy, distribution and ecology : 708
- III- Development : 2 148
- IV- Metabolism : 1 120
- V- Interaction with the environment : 1 433
- VI- Conclusion : 277
- References : 3 823
- Figure legends : 382
- TOTAL: 10 455**

Summary

Brown algae share several important features with land plants, such as their photoautotrophic nature and their cellulose-containing wall, but the two groups are distantly related from an evolutionary point of view. The heterokont phylum, to which the brown algae belong, is a eukaryotic crown group that is phylogenetically distinct not only from the green lineage, but also from the red algae and the opisthokont phylum (fungi and animals). As a result of this independent evolutionary history, the brown algae exhibit many novel features and, moreover, have evolved complex multicellular development independently of the other major groups mentioned above. In 2004, a consortium of laboratories, including the Station Biologique in Roscoff and Genoscope, initiated a project to sequence the genome of *Ectocarpus siliculosus*, a small filamentous brown alga that is found in temperate, coastal environments throughout the globe. The *E. siliculosus* genome, which is currently being annotated, is expected to be the first completely characterised genome of a multicellular alga. In this review we look back over two centuries of work on this brown alga and highlight the advances that have led to the choice of *E. siliculosus* as a genomic and genetic model organism for the brown algae.

198 words

Keywords

Brown alga; phaeophyceae; *Ectocarpus* ; development ; reproduction; pheromone; metabolism; defense

Abbreviations:

ARA, arachidonic acid; BL, blue light; CE: chloroplast envelope; CER: chloroplast endoplasmic reticulum; Ci, inorganic carbon; EPA, eicosapentaenoic acid; DGDG, digalactosyldiacylglycerol; DPG, diphosphatidylglycerol; DGTA, diacylglycerylhydroxymethyl-*N,N,N*-trimethyl- β -alanine; DGTS, diacylglyceryltrimethylhomoserine; MGDG, monogalactosyldiacylglycerol; PC, phosphatidylcholine; PE, phosphatidylethanolamine; PI, phosphatidylinositol; PG, phosphatidylglycerol; SQGD, sulfoquinovosyldiacylglycerol.

1 I- Introduction

2
3 The brown algae belong to the division Heterokonta and are therefore only very distantly
4 related to the three most intensely studied eukaryotic groups, the animals, fungi and green
5 plants (Baldauf, 2003; Davis, 2004; Fig. 1a). This independent evolutionary history has
6 furnished brown algae with many novel metabolic, physiological, cellular and ecological
7 characteristics including a complex halogen metabolism, cell walls containing many unusual
8 polysaccharides and high resistance to osmotic stress. Developmental processes are
9 particularly interesting in this group, which evolved complex multicellularity independently
10 of the three other major groups listed above. From a more applied point of view, the
11 evolutionary history of the brown algae also underlies the high commercial value of several
12 members of the group in the sense that they have evolved novel biomolecules such as
13 polysaccharides and defence elicitors that have a wide range of applications in industry
14 (Klarzynski *et al.*, 2000; McHugh, 2003).

15 It is important to note, however, that whilst the independent evolutionary history of the
16 brown algae is the source of much of the interest of this group, it can also be seen as a
17 handicap because the well-developed model organisms from the plant and animal lineages are
18 of limited relevance to brown algal biology. Specialised brown algal models have been
19 developed in specific domains, for example members of the fucoids for cell biology
20 approaches (see references in Corellou *et al.*, 2005), but a polyvalent model organism that
21 allows access to a wide range of questions at the molecular level has been lacking. This
22 situation is changing with the development of genomic and genetic tools for the filamentous
23 brown alga *Ectocarpus siliculosus*. The sequencing of the genome of this alga has recently
24 been completed and the sequence is currently being annotated

(http://www.cns.fr/externe/English/Projets/Projet_KY/organisme_KY.html). It is therefore an opportune moment to look back at the emergence of *Ectocarpus* as a model organism.

Research on *Ectocarpus* began in the 19th century with descriptions of species and taxonomy, followed by studies aimed at unravelling reproduction and life history. Other major aspects that have been studied include the sexual pheromones and infection of *Ectocarpus* by viruses. Research has also been carried out on ultrastructure, photosynthesis and carbon uptake, gamete recognition and resistance to anti-fouling agents. Several eukaryotic parasites of *Ectocarpus* have been described. A proposition to adopt *Ectocarpus* as a general model organism for the brown algae was made in 2004 (Peters *et al.*, 2004a). This proposition was based partly on this alga's long history as an experimental organism but also took into account several features that make *Ectocarpus* an interesting model for genetic and genomic approaches. These features include its small size, the fact that the entire life cycle can be completed in Petri dishes in the laboratory (Müller *et al.*, 1998), its high fertility and rapid growth (the life cycle can be completed in 3 months), the ease with which genetic crosses can be carried out and the relatively small size of the genome (200 Mbp compared with 1095 and 640 Mbp for *Fucus serratus* and *Laminaria digitata* respectively; Le Gall *et al.*, 1993 and Peters *et al.*, 2004a).

Here we present an overview of the work that has been carried out on *Ectocarpus* over the last two centuries and discuss how the availability of a number of genomic tools, in particular the complete genome sequence, is expected to accelerate research in many domains of brown algal biology in the coming years.

II– Diversity and taxonomy, distribution and ecology

1 - Diversity and taxonomy

Dillwyn (1802-1809) published the first valid description of *Ectocarpus* (using the name *Conferva siliculosa*) based on material collected by W.J. Hooker on "rocks in the sea at Cromer and Hastings". These English localities lie in Norfolk and East Sussex, respectively. Type material, collected by Hooker in 1807, is housed at BM (BM000685585 and BM000685588) under the name *C. confervoides*. Lyngbye (1819) described the genus *Ectocarpus* based on material from Denmark and cited *C. siliculosa* Dillwyn as basionym. The correct nomenclature, therefore, is *E. siliculosus* (Dillwyn) Lyngbye (see Silva *et al.*, 1996 for further details). *E. siliculosus* is the type species of the order Ectocarpales which includes most of the smaller brown algae. Originally regarded as phylogenetically primitive, molecular systematics has shown the Ectocarpales to belong to a group of brown algal orders that evolved rather recently. They are closely related to the large and highly differentiated Laminariales which are major components of coastal marine floras (Rousseau & de Reviers, 1999; Draisma *et al.*, 2003; Cho *et al.*, 2004; Kawai *et al.*, 2007, Fig. 1b). Many species have been described in *Ectocarpus* (www.algaebase.org, April 2007, lists 392 taxa of *Ectocarpus* of whom 98 are flagged as "current"; numerous strains publicly available at the Culture Collection of Algae and Protozoa (CCAP) in Oban, U-K, Gachon *et al.*, 2007a). However, only *E. fasciculatus* Harvey (1841) is currently recognized as a second well-defined species, based on morphology (Russell 1966, 1967a), crossing studies and sequence analyses (Stache-Crain *et al.*, 1997). Crossing experiments have shown that the taxon *E. siliculosus* may represent a species complex (Stache-Crain *et al.*, 1997) and ongoing, refined analyses are expected to resolve this complex, increasing the number of recognised species. Identification of different species of *Ectocarpus* based on morphology is difficult due to the plasticity of the commonly examined features (habit, branching pattern, size of sporangia). In addition, the

two generations of a species may differ considerably (Müller, 1972a; Kornmann & Sahling, 1977).

2 – Distribution

E. siliculosus is distributed world-wide in temperate regions, but does not occur in the tropics and south of the Antarctic convergence (Stache, 1990; Wiencke & Clayton, 2002). It occurs in fully marine and in low-salinity habitats (e.g. 5 psu in Finland) and has even been recorded at a fresh-water site in Australia (West & Craft, 1996) and in a salt polluted river in Germany (Geissler, 1983). Records of *E. fasciculatus* are mainly from the North Atlantic but there are some from Korea, Chile, South Georgia and South Africa (www.algaebase.org, 2007). On the shore, *Ectocarpus* occurs from high inter-tidal pools to the sub-littoral. It is found on abiotic substrata (rocks, wood, plastic, ship hulls) and epiphytic on macrophytes or free-floating (Russell, 1967a,b; 1983a,b). As a result of its ability to grow on a range of abiotic substrates, *Ectocarpus* is a common fouling alga (Morris & Russell, 1974).

3 – Ecology

There have been a limited number of ecophysiological and ecological studies in *Ectocarpus*. Growth rate is dependent on temperature, and there is evidence that temperature also influences the life cycle, at least in some strains (Müller, 1963; Bolton, 1983; see the “sporophyte and gametophyte architecture” section). The thermo-sensitivity of different strains suggests that there is a genetic heterogeneity within the *Ectocarpus* genus (Bolton, 1983). A similar result was obtained for osmo-acclimation (Thomas & Kirst, 1991a,b; see the “Abiotic stress” section). In the field, *Ectocarpus* is a short-lived annual which may dominate the ectocarpoid flora on kelps (Russell, 1983a,b). Despite the commonness of *Ectocarpus*,

there are few data concerning phenology in the field (mostly to be extracted from floristic works) and nothing precise on seasonality or habitats of the two generations.

The availability of the *E. siliculosus* genome sequence is expected to facilitate the analysis of the ecology of this species by providing a basis for the development of molecular markers. Particularly important challenges in this respect will include the identification of the sex locus and of genes specifically expressed in the two generations, as these will provide molecular tools that can be used to investigate several aspects of the life cycle under field conditions. Molecular markers will also allow the exploration of genetic polymorphism amongst *Ectocarpus* species from multiple locations across the globe.

III- Development

1 – Life cycle and reproduction

Male and female gametes are morphologically identical in *Ectocarpus* (isogamy) but differ with respect to their physiology and their behaviour: female gametes settle sooner and produce a pheromone whilst male gametes swim for longer and are attracted to the pheromone produced by the female. Studies on the reproduction of *Ectocarpus* began with the observation of sexual fusions involving the attraction of male gametes to settled female gametes from field thalli of *E. siliculosus* at Naples, Italy (Berthold, 1881). These findings were hotly debated until Sauvageau (1896, 1897) and Oltmanns (1899) succeeded in repeating the experiment. Gamete fusions in *Ectocarpus* were later used by Hartmann (1934) to support his erroneous theory of relative sexuality (see Müller, 1976a for details). Knight (1929) identified the young, unilocular sporangium as the site of meiosis, Papenfuss (1935) and Kornmann (1956) published major contributions to the life history of *Ectocarpus*, and

Boalch (1961) developed refined culture techniques. The entire life history of *E. siliculosus* from Naples was finally unravelled by Müller (1964, 1966, 1967, 1972b) using clonal cultures and chromosome counts. It is schematised in Fig. 2. The basic life history of *E. siliculosus* involves an alternation between the sporophyte and dioecious gametophytes, and sex determination is genotypic (Müller, 1967). Male and female gametophytes are morphologically indistinguishable. One of the problems for understanding the life history was that sporophytes and gametophytes are difficult to distinguish morphologically. Another problem was that zooids from plurilocular reproductive organs have different functions according to the generation forming them: on sporophytes they contain asexual zoospores that directly reproduce the sporophyte, on gametophytes they contain gametes. Further complications include the parthenogenesis of unfused gametes, which develop into haploid parthenogenetic sporophytes morphologically indistinguishable from diploid sporophytes, and heteroblasty (different fates) of spores from unilocular sporangia, developing either into gametophytes or into sporophytes. Furthermore, life cycle generation is not determined rigidly by ploidy (Müller, 1967).

An important challenge for the future will be the characterisation of the genetic mechanisms that control life cycle progression in *Ectocarpus*. This will require the development of methodologies for positional cloning of mutated loci and of genome-wide methods to analyse gene expression throughout the life cycle. Work is currently ongoing in several groups to develop these techniques.

2 – Sporophyte and gametophyte architecture

E. siliculosus is a small filamentous alga that grows to about 30 cm in length in nature, but it may become fertile in the laboratory at 1-3 cm.

Sporophyte development is initiated with the germination of the zygote. The first division produces two cells of identical developmental fate (Peters *et al.*, 2004b). Subsequent mitoses lead to the formation of a basal (or prostrate) filamentous structure, defining the early sporophyte (Fig. 3a). Phaeophycean hairs, *i.e.* hyaline filaments devoid of plastids developing from a basal meristem, are absent in *Ectocarpus* but present in the sister genus *Kuckuckia*. However, in *Ectocarpus* the distal end of filaments, or of plurilocular sporangia, may be less pigmented and resemble a hair; such structures may be referred to as pseudo-hairs (Cardinal, 1964; Pedersen, 1989). If the growth conditions are favourable (Ravanko, 1970), erect filaments (called "upright" filaments) emerge after a few days, contributing to the establishment of an overall filamentous architecture (Fig. 3c).

The typical structure of a vegetative cell is illustrated in Fig. 4. Features common to all brown algal cells include a chloroplast surrounded by four membranes, arranged as two double-membraned envelopes (chloroplast envelope, CE). The second envelope is loosely associated with the chloroplast and forms part of the chloroplast endoplasmic reticulum (CER). The lamellae of the chloroplast are composed of three thylakoids, which are absent from the pyrenoid space (Bouck, 1965; Oliveira & Bisalputra, 1973). In *Ectocarpus* the chloroplast is ribbon-shaped. The size and number of chloroplasts may vary within the same organism (Ravanko, 1970). Other typical features of *Ectocarpus* include several prominent and pedunculated pyrenoids on the inner face of the chloroplast, which are used as a taxonomic marker for the Ectocarpales (Evans, 1966; Rousseau & de Reviers 1999). CER and CE envelopes also surround pyrenoids, this time being tightly adjacent. A third external envelope, called the pyrenoid sac, surrounds the pyrenoid but has no connection with the reticulum system (Bouck, 1965). The nuclear envelope is continuous with the CER, which, itself is in close vicinity to the Golgi apparatus (Bouck, 1965; Oliveira & Bisalputra, 1973). It has been hypothesised that these connections create a complex network of membranes

allowing photosynthates to be efficiently transferred from the chloroplast to the Golgi apparatus, the latter being also in direct contact with the CER (Oliveira & Bisalputra, 1973). Cytoplasmic ER is dispersed throughout the cytoplasm and is mainly rough (Oliveira & Bisalputra, 1973). Osmiophilic bodies (OSB) which are thought to contain lipids, are dispersed throughout the cytoplasm and probably originate from the CER (Oliveira & Bisalputra, 1973). They have been observed within the cell wall, and also external to it. Vacuoles can be either large structures occupying peripheral locations (Oliveira & Bisalputra, 1973) in fixed material, or most of the cellular space (as in land plant cells; Knight, 1929, confirmed by data from our laboratory, after staining with cresyl blue or neutral red, on sporophytic filaments). The nuclear region encompasses two centrioles, which are considered as a microtubule organising centre (MTOC; Katsaros *et al.*, 1991). The chromosome number in the haploid nucleus is estimated to be ca 25 (Peters *et al.*, 2004a). Mitochondria are preferentially peripherally located. They are maternally inherited (Peters *et al.*, 2004b). The cell wall consists of a fibrillar matrix (see the “Photosynthesis and carbohydrate metabolism” section) with several plasmodesmata distributed uniformly along the cross walls. Upon ageing, cell wall ingrowths occur, accompanied by a reduction in the size of the nucleus, mitochondria, ER and Golgi, followed by the disintegration of the chloroplasts and finally by autolysis of the cytoplasm (Oliveira & Bisalputra, 1977a,b).

Most of the reproductive organs are carried by upright filaments (Müller, 1964). Two types of reproductive organs are produced by the sporophyte: plurilocular and unilocular sporangia. Plurilocular sporangia are cone-shaped three-dimensional structures of variable size (Knight, 1929), composed of a large number of locules with different shapes (Baker & Evans, 1973b; Fig. 3e). These locules are generated by several successive mitoses and each gives rise to a single zoospore, which is released through the apex of the sporangium. The mito-spores,

1 which are bi-flagellate and competent for swimming shortly after their release (Müller, 1980),
 2 are a means of vegetative reproduction. Unilocular sporangia are born on the side of branches
 3 (Fig. 3f, Baker & Evans, 1973a). A single meiosis occurs within the single thick-walled locule
 4 (Baker & Evans, 1973a) and this is followed by several mitoses, which generate about a
 5 hundred meio-spores, half of which are female and half male (Müller, 1980). Müller (1963)
 6 and Ravanko (1970) reported that plurilocular sporangia were produced when the external
 7 temperature was relatively high ($\sim 20^{\circ}\text{C}$ or summer), whereas unilocular sporangia were
 8 produced when the temperature was lowered to 13°C (mimicking winter conditions).
 9 However, this temperature dependence is not observed in many strains of *Ectocarpus* (A.F.
 10 Peters, unpublished results). Meio-spores germinate to produce haploid, dioecious
 11 gametophytes (Fig. 3b,d). These are filamentous organisms similar to the sporophyte, but with
 12 two important differences. Firstly, the meio-spore germinates asymmetrically to produce a
 13 rhizoid and an upright filament, so no prostrate structure forms (A.F. Peters, unpublished
 14 results). Secondly, the thallus is more ramified than that of the sporophyte (Müller, 1980).
 15 Gametophytes produce only plurilocular gametangia and these are similar structurally to
 16 plurilocular sporangia on sporophytes. Gametes resemble mito-spores in terms of their size
 17 and motility.

18 The developmental patterning varies greatly across the *Ectocarpus* species complex and is
 19 also dependent on the environment, growth conditions and even the age of the algae for some
 20 features (Ravanko, 1970). This plasticity is observed for the branching frequency and the
 21 number, shape, structure and positioning of the reproductive organs on filaments (Knight,
 22 1929; Müller, 1980; Kim & Lee, 1992). Phytohormones, especially cytokinins, have also been
 23 reported to influence the development of *Ectocarpus* sporophytes (Pedersén, 1968, 1973).

As mentioned above, brown algae are interesting for developmental studies because they have independently evolved complex multicellularity. *Ectocarpus* is a relevant model to address this problem as it is closely related to complex algae such as the Laminariales and the Fucales. However, developmental processes in *Ectocarpus* are clearly simplified compared with its morphogenetically more complex sister families, which provides an advantage for the detailed dissection of these developmental processes. In particular the simple growth pattern of the uniseriate, branched filaments represents an ideal system for the combined application of genetic and mathematical modelling approaches to understanding developmental patterning and then addressing the issue of the evolution of development and multicellularity.

3 – Gametes and spores

Several electron microscopy studies of the motile cells (zooids) of *Ectocarpus* have been reported (Baker & Evans 1973a,b; Lofthouse & Capon, 1975; Maier, 1997a,b). The typical structure of an *Ectocarpus* zooid is illustrated fig. 5.

Ectocarpus zooids correspond to the “primitive” type of brown algal zooid according to Kawai (1992). Gametes and spores typically contain a single chloroplast with a pyrenoid (Baker & Evans, 1973a; Lofthouse & Clayton, 1975; Maier, 1997a). As in the vegetative cells, lamella are composed of three thylakoids and the nuclear envelope is in continuity with the chloroplast endoplasmic reticulum (Maier, 1997a). The nucleus of the male gamete is rich in heterochromatin. Several dictyosomes are present in gametes and spores (Baker & Evans, 1973b; Maier, 1997a). The Golgi apparatus of mitospores is very active both before and after release (Baker & Evans, 1973b). This secretory activity may have important functions in the biosynthesis of the adhesive required for gamete adhesion and for the synthesis of new cell wall compounds during germination.

Ectocarpus gametes and spores are characterized by two flagella with lateral insertion. One is oriented forward and equipped with mastigonemes (hairs), and propels the cell with meandering beats. The second is oriented obliquely backwards, and has no mastigonemes. Most of the time this posterior flagella is passively dragged but occasional lateral beats induce abrupt changes in direction of up to 180° (Geller & Müller, 1981). Gametes and spores have a concave depression at the level of the eyespot into which the swelling of the proximal part of the posterior flagellum fits (Baker & Evans, 1973b; Kreimer *et al.*, 1991; Maier, 1997a). The possible function of the eyespot is to reflect and focus incident light onto the site of photoreception (Kawai *et al.*, 1990; Kreimer *et al.*, 1991). Zoids are capable of positive phototaxis and their posterior flagellum shows strong autofluorescence when irradiated by blue light (450 nm). The photoreceptor pigment is a flavoprotein, which is periodically shaded by a carotenoid stigma (Müller *et al.*, 1987; Kawai, 1988). The acronema (the whiplash tip) is extremely sensitive to mechanical stress and plays an important role in establishing the initial sexual contact between gametes. The details of the flagellar apparatus of both female and male gametes have been studied by electron microscopy (Müller & Falk, 1973; Maier, 1997b). Despite their different behaviours, no difference in their fine structure has been detected (Müller & Falk, 1973).

The pheromone that attracts male gametes to female gametes is an unsaturated hydrocarbon. The substance initially identified was ectocarpene (all-cis-1-(cycloheptadiene-2',5'-yl)-butene-1) (Müller *et al.*, 1971; Müller, 1976b; Müller, 1978; Müller & Schmid, 1988) but more recently Boland *et al.* (1995) have shown that a thermally labile cyclopropyl precursor, pre-ectocarpene, is more active by three orders of magnitude and is thus the actual pheromone.

There is evidence that cell-to-cell recognition between *Ectocarpus* gametes is mediated by N-acetyl glucosamine residues exposed on the plasma membrane of the female gametes and

that these residues are specifically recognized by a receptor on the male gamete (Schmid, 1993; Schmid *et al.*, 1994a). In addition, the lectins concanavalin A (Con A) and *Aleuria aurantia* agglutinin (AAA) bind specifically to the anterior flagella of *Ectocarpus* gametes and the molecules with which they interact also may be involved in the gamete recognition process (Maier & Schmid, 1995).

Zoids from plurilocular and unilocular *Ectocarpus* sporangia share similar overall intracellular structures, but there are some important differences: plurilocular zoids are smaller and swim faster with more rapid changes of direction (Baker & Evans, 1973a). Zoids from unilocular sporangia are strikingly different from the other *Ectocarpus* cell types (both plurilocular zoids and vegetative cells) in that the nucleus is physically separated from the chloroplast. Moreover, secretory activity is lower than in zoids from plurilocular sporangia (Baker & Evans, 1973b).

Gametes of *Ectocarpus* have also been used to study chemotaxis (Boland *et al.*, 1983), receptor modelling (Boland *et al.*, 1989) and phototaxis (Kawai *et al.*, 1990; Kreimer *et al.*, 1991). Protocols for isolation and biochemical characterisation of plasma membrane and CER membrane have also been developed (Schmid *et al.*, 1992). Together with more recently developed molecular tools, these methods now offer access to a range of interesting biochemical events that take place during gamete interaction at fertilization such as chemoreception, cell-cell recognition and fusion processes.

IV- Metabolism

1- Photosynthesis and carbohydrate metabolism

Marine environment constrains several aspects of photosynthesis in brown algae. First, carbon dioxide is not the main source of inorganic carbon (Ci). Indeed, seawater in equilibrium with air contains only 13 μM CO_2 , but more than 2 mM anionic carbon, mainly in the form of HCO_3^- (Beer, 1994). Secondly, when seaweeds are immersed they do not receive the full light spectrum. Light absorption increases with water depth and varies according to the wavelength: red-light (650 nm) is first absorbed followed by purple (400 nm) and yellow- (550 nm) lights. In contrast green (500 nm) and blue (450 nm) lights display a strong penetration: at 10 m depth red light is almost fully absorbed whereas the absorption of green- and blue lights is not significant.

Adapted to these conditions, brown algae differ from land plants by the pigment composition of their light harvesting complexes (LHCs). Chlorophyll c (chl c) and the carotenoid fucoxanthin are indeed the main light-harvesting pigments of brown algae. Their presence broadens the absorption spectrum toward the green light relative to the chlorophytes. Brown algae features two different LHCs associated to photosystems I and II (PSI and PSII), but their pigment composition is controversial. Barrett and Anderson (1980) described a fucoxanthin-chl a/c protein and a violaxanthin enriched-chl a/c protein. Conversely Alberte *et al.* (1981) reported a chl a/c protein devoid of fucoxanthin and a second LHC containing chl a and fucoxanthin but not chlorophyll c. More recent analyses showed that the LHCs associated to PSI and PSII are in fact virtually identical with respect to their pigmentation and peptide composition. All complexes bound chl a, chl c and fucoxanthin in the proportion 6 : 2 : 7, but the LHC associated to PSI is significantly enriched in violaxanthin (De Martino *et al.*, 2000). As in plants, these accessory pigments transfer energy to chlorophyll a within the photosynthetic reaction centres (Grossman *et al.*, 1995). Violaxanthin does not participate in light harvesting but is involved in photoprotection (Demmig-Adams & Adams, 1992; Lemoine *et al.*, 1995).

Under saturating red light, photosynthesis of *E. siliculosus* follows a circadian rhythm with maxima at about noon and can be stimulated by a pulse of blue light (BL) (Schmid & Dring, 1992). This stimulation is also observed in other Phaeophytes, but mostly absent in green or red algae (Schmid *et al.*, 1994b). BL induces several responses: an acidification at the surface of *E. siliculosus* (Schmid & Dring, 1993), accompanied by bicarbonate uptake (Schmid, 1998). A plasma membrane H^+ -ATPase is thought to be activated, and the resulting acidification to increase the conversion of HCO_3^- to CO_2 in the extracellular space (Schmid & Dring, 1993). Recently, such BL-induced H^+ -ATPases have been identified in the brown alga *Laminaria digitata* (Klenell *et al.*, 2002). BL also triggers the mobilization of an internal carbon source, since photosynthesis is stimulated even in the absence of external Ci (Schmid & Dring, 1996). As a result, a C4-like metabolism was initially proposed to exist in *E. siliculosus* (Schmid & Dring, 1996) but not all the enzymes necessary for a C4 cycle were detected (Bush & Schmid, 2001) and the pool of intermediates seems to be too small to act as an organic carbon stock (Hillrichs & Schmid, 2001). The sequestration of a pool of inorganic carbon in the vacuole and its movement to the cytosol in response to BL is now the favoured hypothesis (Schmid & Hillrichs, 2001). Analysis of the sequence of the *E. siliculosus* genome will help to confirm whether or not a C4 pathway exists in the brown algae.

In contrast to the *Plantae* (Moreira *et al.*, 2000), the *Phaeophyceae* do not store the carbon assimilated by photosynthesis as insoluble starch granules, but instead as the soluble 1,3- β -glucan polymer (laminarin) localised in the cytosol (Craigie, 1974), and as mannitol, involved in osmo-acclimation (Davis *et al.*, 2003; see the “Abiotic stresses” section). Brown algae also produce complex polysaccharides which constitute their cell wall. They synthesize some neutral polysaccharides in common with land plants, such as cellulose (Carpita & McCann, 2000), but also unique anionic polysaccharides, such as alginates and sulphated fucans

(Kloareg & Quatrano, 1988). *Ectocarpus* has not been well studied in this respect, but preliminary analyses in our laboratory confirm that all of the polysaccharides typical of brown algae are present in this genus (Estelle Deniaud, pers. comm.). The biosynthetic pathways of these brown algal polysaccharides are essentially unknown and the genome of *E. siliculosus* will be a much anticipated asset to investigate these crucial metabolisms.

2- Lipid metabolism

Worldwide, at least 50 species of brown algae are used as human food. Their lipid content has therefore attracted considerable attention from the viewpoint of both nutrition and pharmacology. A number of studies have been conducted to profile and quantify the fatty acids and the different classes of lipids in these organisms, and to investigate whether the various lipid patterns correlate with the taxonomic position or any other characteristic of the brown algae. Brown algal polar lipids include several common glycolipids (MGDG, DGDG, SQGD, see list of abbreviations) and phospholipids (PC, PE, PI, PG and DPG). Interestingly, several reports have highlighted the high proportion of long chain polyunsaturated fatty acids (LC-PUFAs) (Eichenberger *et al.*, 1993). Moreover, most of the Phaeophyceae contain the betaine lipids DGTA and DGTS, and either contain the common phosphatidylcholine phospholipids in surprising low amounts or do not produce them at all (Eichenberger *et al.*, 1993). All Ectocarpales contain phosphatidylcholine, but exhibit interspecific variation in their DGTA content: *E. fasciculatus* strains contain this lipid while *E. siliculosus* strains do not (Müller & Eichenberger, 1995; Müller, 1995). DGTA may therefore be used as a taxonomic marker (Müller, 1995), although an *E. fasciculatus* strain deficient in DGTA biosynthesis has been described. Genetic analysis of such a strain identified an autosomal locus necessary for the biosynthesis of this lipid (Müller & Eichenberger, 1997).

A novel phosphoglyceride, designated PX, was first isolated from *E. siliculosus*. It was shown to account for 2-4% of total lipids (Schmid *et al.*, 1994c), and to accumulate mostly in the plasma membrane of gametes. Since PX is rich in 20:4n-6 (ARA) and 20:5n-3 (EPA), it has been suggested that it could represent a potential reservoir for pheromone precursors. PX was subsequently detected in other brown algae (Ectocarpales, Fucales and Sphacelariales; Schmid *et al.*, 1994c).

Lipid metabolism plays also major roles in the control of defense mechanisms. Therefore, together with the genome sequence, it is likely that the efforts to screen mutants with altered resistance to pathogen attacks, will lead to the phenotyping of plants impaired in lipid or fatty acid metabolic pathways, thereby potentially revealing novel specific traits of this metabolism in brown algae.

V- Interactions with the environment

I- Ectocarpus pathogens

Despite of their small size and ephemeral life stages, filamentous brown algae have been frequently reported to be plagued by various pathogens, including viruses (Müller *et al.*, 1998) and eukaryotic parasites of different phylogenetic lineages: oomycetes, chytrids and hyphochytrids (Andrews 1976; Küpper & Müller 1999; Müller *et al.* 1999) and by parasites related to the Plasmodiophorea (Karling, 1944; Maier *et al.*, 2000). In addition, numerous historical records described ectocarpoids with abnormal sporangia or vegetative cells suspected to contain unknown parasites (Ratray, 1885; Müller *et al.*, 1998).

The oomycete *Eurychasma dicksonii* has been described mainly in wild populations of *Pylaiella littoralis* (Küpper & Müller, 1999) but it displays a broad host range and infects

various brown algae including *Ectocarpus* (Müller *et al.*, 1999) in which it was initially described by Wright (1879). There is a current effort to set up a defined pathosystem using *E. siliculosus* and *E. dicksonii*, and *Ectocarpus* strains have been shown to exhibit differential susceptibility to a defined *Eurychasma* strain. Conversely, several *Eurychasma* strains exhibit different host specificities, suggesting co-evolution of the two species (Gachon *et al.*, 2007b). The molecular bases of resistance and virulence are under investigation.

Chytrids were described earlier by Petersen (1905) and the hyphochytrid *Anisolpidium ectocarpii* was described by Karling (1943) and Johnson (1957). Like *E. dicksonii*, *Chytridium polysiphoniae* (Chytridiomycota) is ubiquitous and can infect many hosts, including *E. siliculosus* and *E. fasciculatus* (Müller *et al.*, 1999). Interestingly, its negative effects on photosynthesis of its host was described at the cellular level in the related ectocarpoid *P. littoralis* using fluorescence kinetic microscopy (Gachon *et al.*, 2006). Recently, the 18S rRNA genes of *Chytridium polysiphoniae* and *Eurychasma dicksonii* were sequenced and used to clarify their phylogenetic affiliations (Küpper *et al.*, 2006). The plasmodiophorean *Maullinia ectocarpii* is an obligate intracellular parasite of *Ectocarpus* spp. (Maier *et al.*, 2000). However, the extent to which this infection occurs in nature and its effect on algal fitness are presently unknown.

Viral infections represent by far the most studied phenomenon in *E. siliculosus* (Müller, 1996; Müller & Knippers, 2001). Until the late eighties most reports of virus infections in brown algal tissues were based on electron microscopy studies, which sporadically described “virus-like particles” (VLP). Viruses were obtained in culture for the first time from a New Zealand strain of *E. siliculosus* after lysis of host cells allowing evaluation of their infection potential (Müller *et al.*, 1990; Müller, 1991). Virus infections were found in about 50% of the individuals of a given natural population (Dixon *et al.*, 2000; Müller *et al.*, 2000) and were

1 shown to occur worldwide in correlation with the cosmopolitan distribution of *E. siliculosus*
 2 (Müller, 1991; Sengco *et al.*, 1996).

3 The viruses that infect different ectocarpoid algae exhibit considerable variability in size
 4 and diameter and, in general, they display a high level of host specificity (Müller *et al.*, 1998).
 5 However, several instances of trans-specific infection have been described, for example
 6 between EsV-1 (*Ectocarpus siliculosus* virus-1) and *Kuckuckia kylinii* (Müller, 1992; Müller
 7 & Schmid, 1996) and also between EfasV-1 (*Ectocarpus fasciculatus* virus-1) and *E.*
 8 *siliculosus* (Müller *et al.*, 1996; Sengco *et al.*, 1996). Interestingly, EsV-1 and EfasV-1 are the
 9 most similar of the brown algal viruses in terms of their genome size (Müller *et al.*, 1996).

10 The EsV-1 virus specifically infects the single-celled gametes or spores, *i.e.* the only cells in
 11 the life history that lack a cell wall (Maier & Müller, 1998). Following infection, a single
 12 copy of the viral DNA appears to integrate into the host genome (Delaroque *et al.*, 1999). The
 13 viral DNA is then transmitted, *via* mitotic divisions, to all the cells of the developing alga.
 14 This has been confirmed by regenerating algae from protoplasts derived from virus-infected
 15 gametophytes (Kuhlenkamp & Müller, 1994). Despite the fact that they carry the integrated
 16 virus, vegetative cells do not produce viral particles (Müller *et al.*, 1998). Viral particles are
 17 only produced in reproductive organs (sporangia and gametangia) of mature algae from where
 18 they are released to infect a new generation of zoids. In addition to these cycles of re-
 19 infection, the viral genome can be also transmitted to progeny through meiosis, in which case
 20 it segregates as a Mendelian factor and is inherited by half of the progeny (Müller, 1991;
 21 Bräutigam *et al.*, 1995). The pathogenic character of viral infections has been unambiguously
 22 confirmed, but this association's main impact is on reproductive success. Plant sterility varies
 23 from partial (Müller *et al.*, 1990) to total (Müller & Frenzer, 1993) but no significant
 24 difference in photosynthesis, respiration and growth rate were observed in infected

gametophytes or sporophytes (Del Campo *et al.*, 1997). This contrasts with the reduced photosynthetic performance of *Feldmannia* species infected with FsV (Robledo *et al.*, 1994).

The EsV-1 genome is a circular DNA molecule of a relatively large size (335 Kbp) for a phycodnavirus (Van Etten & Meints, 1999; Van Etten *et al.*, 2002) with double-stranded regions interrupted by single-stranded regions (Lanka *et al.*, 1993; Klein *et al.*, 1994). Both EsV-1 and the related *Feldmannia irregularis* virus (FirrV-1) have been sequenced (Delaroque *et al.*, 2001; Delaroque *et al.*, 2003). EsV-1 contains approximately 231 genes with a wide range of predicted functions including DNA metabolism, signalling, transposition, DNA integration and polysaccharide metabolism (Delaroque *et al.*, 2000a,b; Delaroque *et al.*, 2003). It has also been proposed that the ability of the virus to integrate into its host's genome could be exploited to develop a transformation vector for a wide range of brown algae, including *E. siliculosus* (Henry & Meints, 1994, Delaroque *et al.*, 1999). However, the complex integration pattern of the virus into the algal genome will considerably complicate this task (N. Delaroque, pers. comm.). A microarray has been constructed to analyse EsV-1 gene expression (Declan Schroeder, pers. comm.) and it will be particularly interesting in the future to couple the analysis of viral and genome-wide host gene expression during viral infection.

The development of genomic tools provides a new context to investigate the possible genetic basis of the co-evolution between some pathogens and brown algae. The search for inducers of defense responses and resistance against parasites is also still opened as, in contrary with kelps, *Ectocarpus* does not react with an oxidative burst upon recognition of alginate fragments (Küpper *et al.*, 2002a).

2- Abiotic stresses

1 *E. siliculosus* is able to exploit a wide range of habitats and environmental conditions (see
2 the “Distribution” section). This feature seems likely to be based at least as much as on a high
3 intrinsic genetic variability as on a general physiological toughness, as illustrated by work
4 carried out on copper and saline stress responses.

5 Interspecific variations in copper tolerance have been observed between different
6 strains of *E. fasciculatus* and *E. siliculosus*, with the latter being the most tolerant (Morris,
7 1974). Differences have also been observed among *E. siliculosus* strains that are differently
8 exposed to copper in their natural habitat (Russell & Morris, 1970; Hall, 1981). Cu^{2+}
9 interferes with the general process of photosynthesis in brown algae, and particularly in *E.*
10 *siliculosus*, by competing with magnesium for metal binding sites in the chlorophyll
11 molecules (Küpper *et al.*, 2002b). A study of the mechanism of tolerance to copper and other
12 heavy metals suggested a co-tolerance to copper, cobalt and zinc, and provided evidence for
13 an exclusion mechanism to explain the particularly low sensitivity of *E. siliculosus* copper
14 tolerant strains (Hall *et al.*, 1979; Hall, 1980, 1981). However, as yet there is no clear
15 explanation for the intra-specific variation with respect to this trait within this species.

16 Ability of some *E. siliculosus* strains to tolerate copper has been considered to
17 develop bioassays in which this alga is used for monitoring marine antifouling characteristics
18 of copper-based materials (Hall and Baker, 1985, 1986). In addition, copper chloride was
19 used to inhibit *E. siliculosus* infestations in tank cultures of *Gracilaria gracilis* (Van Heerden
20 *et al.*, 1997).

21
22 Russell and Bolton (1975) reported the occurrence of salinity ecotypes within *E. siliculosus*.
23 This study was extended by Thomas and Kirst (1991a,b) who showed that large differences in
24 photosynthesis, accumulation of osmotically active compounds (mannitol; Davis *et al.*, 2003)
25 and vitality occur between *E. siliculosus* isolates from different geographic locations

1 following changes in salinity. They also observed that sporophytes were more salt tolerant
2 than gametophytes, irrespective of their level of ploidy.

3 Detailed investigations are necessary to decipher the physiological and cellular bases of salt
4 and heavy metal tolerance in *E. siliculosus*. Mutagenesis and transcriptomic approaches will
5 thus help to better understand the mechanisms involved in osmotic and oxidative adaptation,
6 and to explain how these algae can cope with such a wide range of environmental conditions.

9 VI- Conclusion

11 Taken together, the above sections illustrate the broad range of phenomena that have been
12 studied in *Ectocarpus* and provide an indication of the domains that could be further explored
13 in the future. Notably, a large proportion of this past work, covering many diverse aspects of
14 *Ectocarpus* biology, was carried out in Dieter Müller's laboratory in Konstanz, and the efforts
15 of this group have therefore laid the foundations for the development of *Ectocarpus* as a
16 model organism.

17 The *Ectocarpus* genome project has federated a consortium of laboratories with an interest
18 in this organism and these laboratories are currently developing several molecular tools.
19 These include mutant screens, genetic transformation and genome-scale analysis of gene
20 expression. Several developmental mutants have been isolated and positional cloning of some
21 of the affected genes should be feasible in the near future. The availability of the genome
22 sequence together with the ability to analyse gene function by forward and reverse genetic
23 approaches will make it possible to address additional questions, many of which have been
24 evoked in this review. Examples include the biosynthesis of diverse, brown-algal-specific
25 metabolites, such as lipids and complex cell wall components, and the genetic basis of

resistance to biotic and abiotic aggression. The genome will also be an invaluable aide for the study of the ecology of *Ectocarpus*, by serving as the base for the development of neutral and selected molecular markers for the analysis of field isolates. In conclusion, the *Ectocarpus* genome sequence and the tool development associated with this project are providing access to a relatively unexplored branch of the eukaryotic tree and some exciting discoveries can be expected in this domain in the coming years.

References

- Alberte RS, Friedman AL, Gustafson DL, Rudnick MS, Lyman H. 1981.** Light-harvesting systems of brown algae and diatoms. Isolation and characterization of chlorophyll a/c and chlorophyll a/fucoanthin pigment-protein complexes. *Biochimica et Biophysica Acta* **635**: 304-316.
- Andrews JA. 1976.** The pathology of marine algae. *Biological Reviews* **51**: 211-253.
- Baker JRJ, Evans LV. 1973a.** The ship fouling alga *Ectocarpus*. I. Ultrastructure and cytochemistry of plurilocular reproductive stages. *Protoplasma* **77**: 1-13.
- Baker JRJ, Evans LV. 1973b.** The ship fouling alga *Ectocarpus*. II. Ultrastructure of unilocular reproductive stages. *Protoplasma* **77**: 181-189.
- Baldauf SL. 2003.** The deep roots of eukaryotes. *Science* **300**: 1703 - 1706.
- Barrett J, Anderson JM. 1980.** The P-700-chlorophyll alpha-protein complex and two major light-harvesting complexes of *Acrocarpia paniculata* and other brown seaweeds. *Biochimica et Biophysica Acta* **590**: 309-323.
- Beer S. 1994.** Mechanisms of inorganic carbon acquisition in marine macroalgae. *Progress in Phycological Research* **10**: 179-207.

- Berthold G. 1881.** Die geschlechtliche Fortpflanzung der eigentlichen Phaeosporeen. *Mitteilungen aus der Zoologischen Station zu Neapel* **2**: 401-413.
- Boalch GT. 1961.** Studies on *Ectocarpus* in culture. *Journal of the Marine Biological Association*. **41**: 279-304.
- Boland W, Marner FJ, Jaenicke L. 1983.** Comparative receptor study in gamete chemotaxis of the seaweeds *Ectocarpus siliculosus* and *Cutleria multifida* – An approach to interspecific communication of algal gametes. *European Journal of Biochemistry* **134**: 97-103.
- Boland W, König WA, Krebber R, Müller DG. 1989.** Separation of enantiomeric algal pheromones and related hydrocarbons by gas-liquid chromatography on modified cyclodextrins as chiral stationary phases. Biosynthetic relevance of racemic by-products. *Helvetica Chimica Acta* **72**: 1288-1292.
- Boland W, Pohnert G, Maier I. 1995.** Biosynthesis of algae pheromones. 4. Pericyclic-reaction in nature – Spontaneous cope rearrangement inactivates algae pheromones. *Angewandte Chemie* **34**: 1602-1604.
- Bolton JJ. 1983.** Ecoclinical variation in *Ectocarpus siliculosus* (Phaeophyceae) with respect to temperature growth optima and survival limits. *Marine Biology* **73**: 131-138.
- Bouck GB. 1965.** Fine structure and organelle associations in brown algae. *Journal of Cell Biology* **26**: 523-537.
- Bräutigam M, Klein M, Knippers R, Müller DG. 1995.** Inheritance and meiotic elimination of a virus genome in the host *Ectocarpus siliculosus* (Phaeophyceae). *Journal of Phycology* **31**: 823-827.
- Busch S, Schmid R. 2001.** Enzymes associated with β -carboxylation in *Ectocarpus siliculosus* (Phaeophyceae): are they involved in net carbon acquisition? *European Journal of Phycology* **36**, 61-70.

- 1 **Cardinal A. 1964.** Étude sur les Ectocarpacées de la Manche. *Nova Hedwigia* **15**: 1-86.
- 2 **Carpita N, McCann M. 2000.** The cell wall. In: Buchanan B, Gruissem W, Jones R, eds.
- 3 *Biochemistry and molecular biology of plants*. 52-108.
- 4 **Cho, GY, Lee SH, Boo SM. 2004.** A new brown algal order, Ishigeales (Phaeophyceae),
- 5 established on the basis of plastid protein-coding *rbcL*, *psaA*, and *psbA* region
- 6 comparisons. *Journal of Phycology* **40**: 921–936.
- 7 **Corellou F, Coelho SM, Bouget FY, Brownlee C. 2005.** Spatial re-organisation of cortical
- 8 microtubules in vivo during polarisation and asymmetric division of *Fucus* zygotes.
- 9 *Journal of Cell Science*. **118** :2723-2734.
- 10 **Craigie JS. 1974.** Storage products. In: Stewart WPD, eds. *Algal Physiology and*
- 11 *Biochemistry*. Berkeley, University of California press, USA 206-235.
- 12 **Davis TA, Volesky B, Mucci A. 2003.** A review of the biochemistry of heavy metal
- 13 biosorption by brown algae. *Water research* **37**: 4311-4330.
- 14 **Davis, RH. 2004.** The age of model organisms. *Nature Reviews Genetics* **5**: 69-75.
- 15 **Del Campo E, Ramazanov Z, Garcia-Reina G, Müller DG. 1997.** Photosynthetic responses
- 16 and growth performance of virus-infected and noninfected *Ectocarpus siliculosus*
- 17 (Phaeophyceae). *Phycologia* **36**: 186-189.
- 18 **Delaroque N, Maier I, Knippers R, Müller DG. 1999.** Persistent virus integration into the
- 19 genome of its algal host, *Ectocarpus siliculosus* (Phaeophyceae). *Journal of General*
- 20 *Virology* **80**: 1367-1370.
- 21 **Delaroque N, Wolf S, Müller DG, Knippers R. 2000a.** The brown algal virus EsV-1
- 22 particle contains a putative hybrid histidine kinase. *Virology* **273**: 383-390.
- 23 **Delaroque N, Wolf S, Müller DG, Knippers R. 2000b.** Characterization and
- 24 immunolocalization of major structural proteins in the brown algal virus EsV-1. *Virology*
- 25 **269**: 148-155.

- Delaroque N, Müller DG, Bothe G, Pohl T, Knippers R, Boland W. 2001.** The complete DNA sequence of the *Ectocarpus siliculosus* Virus EsV-1 genome. *Virology* **287**: 112-132.
- Delaroque N, Boland W, Müller DG, Knippers R. 2003.** Comparisons of two large phaeoviral genomes and evolutionary implications. *Journal of Molecular Evolution* **57**: 613-622.
- De Martino A, Douady D, Quinet-Szely M, Rousseau B, Crépineau F, Apt K, Caron L. 2000.** The light-harvesting antenna of brown algae: highly homologous proteins encoded by a multigene family. *European Journal of Biochemistry* **267**: 5540-5549.
- Demmig-Adams B, Adams III WW. 1992.** Photoprotection and other responses to high light stress. *Annual Review of Plant Physiology and Plant Molecular Biology* **43**: 599-626.
- Dixon NM, Leadbeater BSC, Wood KR. 2000.** Frequency of viral infection in a field population of *Ectocarpus fasciculatus* (Ectocarpales, Phaeophyceae). *Phycologia* **39**: 258-263.
- Draisma SGA, Peters AF, Fletcher RL. 2003.** Evolution and taxonomy in the Phaeophyceae: effects of the molecular age on brown algal systematics. In: Norton TA, eds *Out of the Past*. Collected reviews to celebrate the jubilee of the British Phycological Society. The British Phycological Society, Belfast 87-102.
- Eichenberger W, Araki S, Müller DG. 1993.** Betaine lipid and phospholipids in brown algae. *Phytochemistry* **34**: 1323-1333.
- Evans LV. 1966.** Distribution of pyrenoids among some brown algae. *Journal of Cell Science* **1**: 449-454.
- Gachon CMM, Küpper H, Küpper FC, Šetlík I. 2006.** Witnessing effects of a pathogen on photosynthesis of its host at the cellular level: chlorophyll fluorescence kinetic

- 1 microscopy of *Pylaiella littoralis* (Phaeophyceae) infected by *Chytridium polysiphoniae*
 2 (Chytridiomycota). *European Journal of Phycology* **41**: 395-403.
- 3 **Gachon CM, Day JG, Campbell CN, Proschold T, Saxon RJ, Küpper FC. 2007a.** Culture
 4 Collection of Algae and Protozoa (CCAP): A biological resource for protistan genomics.
 5 Gene. In Press.
- 6 **Gachon CM, Müller DG, Gaj G, Küpper FC. 2007b.** Confronting disease: not all algae are
 7 equal. *The Phycologist* **75**, 15, abstract.
- 8 **Geller A, Müller DG. 1981.** Analysis of the flagellar beat pattern of male *Ectocarpus*
 9 *siliculosus* gametes (Phaeophyta) in relation to chemotactic stimulation by female cells.
 10 *Journal of Experimental Biology* **92**: 53-666.
- 11 **Geissler U. 1983.** Die salzbelastete Flußstrecke der Werra - ein Binnenlandstandort für
 12 *Ectocarpus confervoides* (Roth) Kjellman. *Nova Hedwigia* **37**: 193-217.
- 13 **Grossman AR, Bhaya D, Apt KE, Kehoe DM. 1995.** Light-harvesting complexes in
 14 oxygenic photosynthesis: diversity, control, and evolution. *Annual Review of Genetics* **29**:
 15 231-288.
- 16 **Hall A, Fielding AH, Butler M. 1979.** Mechanism of copper tolerance in the marine fouling
 17 alga *Ectocarpus siliculosus* - evidence for an exclusion mechanism. *Marine Biology* **54**:
 18 195-199.
- 19 **Hall A. 1980:** Heavy-metal co-tolerance in a copper-tolerant population of the marine fouling
 20 alga, *Ectocarpus siliculosus* (DILLW) LYNGBYE. *New Phytologist* **85**: 73-78.
- 21 **Hall A. 1981.** Copper accumulation in copper-tolerant and non-tolerant populations of the
 22 marine fouling alga *Ectocarpus siliculosus* (Dillw.) Lyngbye. *Botanica Marina* **24**: 223-
 23 228.
- 24 **Hall A, Baker AJM. 1985:** Settlement and growth of copper-tolerant *Ectocarpus siliculosus*
 25 (DILLW.) LYNGBYE on different copper-based antifouling surfaces under laboratory

conditions.1. Corrosion trials in seawater and development of an algal culture system.
Journal of Materials Science **20**: 1111-1118.

Hall A, Baker AJM. 1986: Settlement and growth of copper-tolerant *Ectocarpus siliculosus* (DILLW.) LYNGBYE on different copper-based antifouling surfaces under laboratory conditions. 2. A comparison of the early stages of fouling using light and electron microscopy. *Journal of Materials Science* **21**: 1240-1252.

Hartmann M. 1934. Untersuchungen über die Sexualität von *Ectocarpus siliculosus*. *Archiv für Protistenkunde* **83**: 110-163.

Henry EC, Meints RH. 1994. Recombinant viruses as transformation vectors of marine macroalgae. *Journal of Applied Phycology* **6**: 247-253.

Hillrichs S, Schmid R. 2001. Activation by blue light of inorganic carbon acquisition for photosynthesis in *Ectocarpus siliculosus*: organic acid pools and short-term carbon fixation. *European Journal of Phycology* **36**: 71-79.

Johnson TW. 1957. Resting spore development in the marine phycomycete *Anisolpidium ectocarpii*. *American Journal of Botany* **44**: 875-878.

Karling JS. 1943. The life history of *Anisolpidium ectocarpii* gen. nov. et sp. nov., and a synopsis and classification of other fungi with anteriorly uniflagellate zoospores. *American Journal of Botany* **30**: 637-648.

Karling JS. 1944. *Phagomyxa algarum* n. gen.n. sp., an unusual parasite with plasmodiophoralean and proeomyxean characteristics. *American Journal of Botany* **31**: 38-52.

Katsaros C, Kreimer G, Melkonian M. 1991. Localization of tubulin and a centrin-homologue in vegetative cells and developing gametangia of *Ectocarpus siliculosus* (Dillw.) Lyngb. (Phaeophyceae, Ectocarpales). *Botanica Acta* **104**: 87-92.

- 1 **Kawai H. 1988.** A flavin-like autofluorescent substance in the posterior flagellum of golden
2 and brown algae. *Journal of Phycology* **24**: 114-117.
- 3 **Kawai H, Müller DG, Fölster E, Häder DP. 1990.** Phototactic responses in the gametes of
4 the brown alga, *Ectocarpus siliculosus*. *Planta* **182**: 292-297.
- 5 **Kawai H. 1992.** A summary of the morphology of chloroplast and flagellated cells in the
6 Phaeophyceae. *Korean Journal of Phycology* **7**: 33-43.
- 7 **Kawai H, Hanyuda T, Draisma SGA, Müller DG. 2007.** Molecular phylogeny of
8 *Discosporangium mesarthrocarpum* (Phaeophyceae) with a reinstatement of the order
9 Discosporangiales. *Journal of Phycology* **43**: 186-194.
- 10 **Kim HS, Lee IK. 1992.** Morphotaxonomic studies on the Korean Ectocarpaceae
11 (Phaeotphyta) I. Genus *Ectocarpus* Lyngbye. *Korean Journal of Phycology* **7**(2): 225-242.
- 12 **Klarzynski O, Plesse B, Joubert J-M, Yvin J-C, Kopp M, Kloareg B, Fritig B. 2000.**
13 Linear β -1,3 glucans are elicitors of defense responses in tobacco. *Plant Physiology* **124**:
14 1027-1037.
- 15 **Klein M, Lanka STJ, Müller DG, Knippers R. 1994.** Single-stranded regions in the
16 genome of the *Ectocarpus siliculosus* virus. *Virology* **202**: 1076-1078.
- 17 **Klenell M, Snoeijis P, Pedersén M. 2002.** The involvement of a plasma membrane H^+ -
18 ATPase in the blue-light enhancement of photosynthesis in *Laminaria digitata*
19 (Phaeophyta). *Journal of Phycology* **38**: 1143-1149.
- 20 **Kloareg B, Quatrano RS. 1988.** Structure of the cell walls of marine algae and
21 ecophysiological functions of the matrix polysaccharides. *Oceanography and Marine*
22 *Biology. An Annual Review.* **26**: 259-315.
- 23 **Knight M. 1929.** Studies in the Ectocarpaceae. II. The life-history and cytology of
24 *Ectocarpus siliculosus*, Dillw. *Transactions of the Royal Society of Edinburgh* **56**: 307-
25 332, 6 plates.

- Kornmann P. 1956.** Über die Entwicklung einer *Ectocarpus confervoides*-Form. *Pubblicazioni della stazione zoologica di Napoli* **28**: 32-43.
- Kornmann P, Sahling PH. 1977.** Meeresalgen von Helgoland. Benthische Grün-, Braun- und Rotalgen. *Helgoländer wissenschaftliche Meeresuntersuchungen* **29**: 1-289.
- Kreimer G, Kawai H, Müller DG, Melkonian M. 1991.** Reflective properties of the stigma in male gametes of *Ectocarpus siliculosus* (Phaeophyceae) studied by confocal laser scanning microscopy. *Journal of Phycology* **27**: 268-276.
- Kuhlenkamp R, Müller DG. 1994.** Isolation and regeneration of protoplasts from healthy and virus-infected gametophytes of *Ectocarpus siliculosus* (Phaeophyceae). *Botanica Marina* **37**: 525-530.
- Küpper FC, Müller DG. 1999.** Massive occurrence of the heterokont and fungal parasites *Anisolpidium*, *Eurychasma* and *Chytridium* in *Pylaiella littoralis* (Ectocarpales, Phaeophyceae). *Nova Hedwigia* **69**: 381-389.
- Küpper FC, Müller D, Peters A, Kloareg B, Potin P. 2002a.** Oligoalginat recognition and oxidative burst play a key role in natural and induced resistance of sporophytes of Laminariales. *Journal of Chemical Ecology* **28**: 2057-2081.
- Küpper H, Šetlík I, Spiller M, Küpper FC, Prášil O. 2002b.** Heavy metal-induced inhibition of photosynthesis: Targets of in vivo heavy metal chlorophyll formation. *Journal of Phycology* **38**: 429-441.
- Küpper FC, Maier I, Müller DG, Loiseaux-de Goër S, Guillou L. 2006.** Phylogenetic affinities of two eukaryotic pathogens of marine macroalgae, *Eurychasma dicksonii* and *Chytridium polysiphoniae*. *Cryptogamie Algologie* **27**: 165-184.
- Le Gall Y, Brown S, Marie D, Mejjad M, Kloareg B. 1993.** Quantification of nuclear DNA and G-C content in marine macroalgae by flow cytometry of isolated nuclei. *Protoplasma* **173**: 123-132.

- 1 **Lanka STJ., Klein M, Ramsperger U, Müller DG, Knippers R. 1993.** Genome structure of
2 a virus infecting the marine brown alga *Ectocarpus siliculosus*. *Virology* **193**: 802-811.
- 3 **Lemoine Y, Harker M, Rmiki NE, Rousseau B, Berkaloﬀ C, Duval JC, Young AJ,**
4 **Britton G. 1995.** Xanthophyll cycle operation and photoprotection in brown algae: effects
5 of high light and desiccation. *In Photosynthesis, from Light to Biosphere* (Mathis, P., ed.),
6 vol. IV, 119-122. Kluwer, Dordrecht.
- 7 **Lofthouse PF , Capon B. 1975.** Ultrastructural changes accompanying mitosporeogenesis in
8 *Ectocarpus parvus*. *Protoplasma* **84**: 83-99.
- 9 **Maier I, Schmid CE. 1995.** An immunofluorescence study on lectin binding sites in gametes
10 of *Ectocarpus siliculosus* (Ectocarpales, Phaeophyceae). *Phycological Research* **43**: 33-
11 42.
- 12 **Maier I. 1997a.** The fine structure of the male gamete of *Ectocarpus siliculosus*
13 (Ectocarpales, Phaeophyceae). I. General structure of the cell. *European Journal of*
14 *Phycology* **32**: 241-253.
- 15 **Maier I. 1997b.** The fine structure of the male gamete of *Ectocarpus siliculosus*
16 (Ectocarpales, Phaeophyceae). II. The flagellar apparatus. *European Journal of Phycology*
17 **32**: 255-266.
- 18 **Maier I, Müller DG. 1998.** Virus binding to brown algal spores and gametes visualized by
19 DAPI fluorescence microscopy. *Phycologia* **37**: 60-63.
- 20 **Maier I, Parodi E, Westermeier R, Müller DG. 2000.** *Maullina ectocarpii* gen. et sp. nov.
21 (Plasmodiophorea), an intracellular parasite in *Ectocarpus siliculosus* (Ectocarpales,
22 Phaeophyceae) and other filamentous brown algae. *Protist* **151**: 225-238.
- 23 **McHugh DJ. 2003.** A guide to the seaweed industry. FAO Fisheries Technical Paper No.
24 441. FAO, Rome: 105 pp.

- 1 **Moreira D, Le Guyader H, Philippe H. 2000.** The origin of red algae and the evolution of
2 chloroplasts. *Nature* **405**: 69-72.
- 3 **Morris OP. 1974.** Inter-specific differences in responses to copper by natural populations of
4 *Ectocarpus*. *British Phycology Journal* **9**: 269-272.
- 5 **Morris OP, Russel G. 1974.** Inter-specific differences in responses to copper by natural
6 populations of *Ectocarpus*. *British Phycology Journal* **9**: 269-272.
- 7 **Müller DG. 1963.** Die Temperaturabhängigkeit der Sporangienbildung bei *Ectocarpus*
8 *siliculosus* von verschiedenen Standorten. *Pubblicazioni della stazione zoologica di*
9 *Napoli* **33**: 310-314.
- 10 **Müller DG. 1964.** Life-cycle of *Ectocarpus siliculosus* from Naples, Italy. *Nature* **26**: 1402.
- 11 **Müller DG. 1966.** Untersuchungen zur Entwicklungsgeschichte der Braunalge *Ectocarpus*
12 *siliculosus* aus Neapel. *Planta* **68**: 57-68.
- 13 **Müller DG. 1967.** Generationswechsel, Kernphasenwechsel und Sexualität der Braunalge
14 *Ectocarpus siliculosus* im Kulturversuch. *Planta* **75**: 39-54.
- 15 **Müller DG, Jaenicke L, Donike M, Akintobi T. 1971.** Sex attractant in a brown alga:
16 chemical structure. *Science* **171**: 815-817.
- 17 **Müller DG. 1972a.** Life cycle of the brown alga *Ectocarpus fasciculatus* var. *refractus*
18 (Kütz.) Ardis. (Phaeophyta, Ectocarpales) in culture. *Phycologia* **11**: 11-13.
- 19 **Müller DG. 1972b.** Studies on reproduction in *Ectocarpus siliculosus*. *Société Botanique*
20 *Française, Mémoires*: 87-98.
- 21 **Müller DG, Falk H. 1973.** Flagellar structure of the gametes of *Ectocarpus siliculosus*
22 (Phaeophyta) as revealed by negative staining. *Archiv für Mikrobiologie* **91**: 313-322.
- 23 **Müller DG. 1976a.** Sexual isolation between a European and an American population of
24 *Ectocarpus siliculosus* (Phaeophyta). *Journal of Phycology* **12**: 252-254.

- 1 **Müller DG. 1976b.** Quantitative evaluation of sexual chemotaxis in two marine brown algae.
2 *Zeitschrift für Pflanzenphysiologie* **80**: 120-130.
- 3 **Müller DG. 1978.** Locomotive responses of male gametes to the species specific sex
4 attractant in *Ectocarpus siliculosus* (Phaeophyta). *Archiv für Protistenkunde* **120**: 371-
5 377.
- 6 **Müller DG. 1980.** Entwicklung von *Ectocarpus siliculosus* (Phaeophyta). Film C 1308 des
7 IWF, Göttingen 1979. *Publikationen zu wissenschaftlichen Filmen, Sektion Biologie*,
8 Serie 13 11/C 1308: 1-15.
- 9 **Müller DG, Maier I, Müller H. 1987.** Flagellum autofluorescence and photoaccumulation in
10 heterokont algae. *Photochemistry and Photobiology* **46**: 1003-1008.
- 11 **Müller DG, Schmid CE. 1988.** Qualitative and quantitative determination of pheromone
12 secretion in female gametes of *Ectocarpus siliculosus* (Phaeophyceae). *Biological*
13 *Chemistry Hoppe-Seyler* **369**: 647-653.
- 14 **Müller DG, Kawai H, Stache B, Lanka STJ. 1990.** A virus infection in the marine brown
15 alga *Ectocarpus siliculosus* (Phaeophyceae). *Botanica Acta* **103**: 72-82.
- 16 **Müller DG. 1991.** Mendelian segregation of a virus genome during host meiosis in the
17 marine brown alga *Ectocarpus siliculosus*. *Journal of Plant Physiology* **137**: 739-743.
- 18 **Müller DG. 1992.** Intergeneric transmission of a marine plant DNA virus.
19 *Naturwissenschaften* **79**: 37-39.
- 20 **Müller DG, Frenzer K. 1993.** Virus infection in three marine brown algae: *Feldmannia*
21 *irregularis*, *F. simplex*, and *Ectocarpus siliculosus*. *Hydrobiologia* **260-261**: 37-44.
- 22 **Müller DG. 1995.** Taxonomic value of a betaine lipid in *Ectocarpus* (Phaeophyceae) and the
23 first record of a sexual population of *E. fasciculatus* for the Pacific Ocean. *Phycological*
24 *Research* **43**: 175-177.

- 1 **Müller DG, Eichenberger W. 1995.** Crossing experiments, lipid composition, and the
2 species concept in *Ectocarpus siliculosus* and *E. fasciculatus* (Phaeophyceae,
3 Ectocarpales). *Journal of Phycology* **31**: 173-176.
- 4 **Müller DG. 1996.** Host-virus interactions in marine brown algae. *Hydrobiologia* **327**: 21-28.
- 5 **Müller DG, Schmid CE. 1996.** Intergeneric infection and persistence of *Ectocarpus* virus
6 DNA in *Kuckuckia* (Phaeophyceae, Ectocarpales). *Botanica Marina* **39**: 401-405.
- 7 **Müller DG, Sengco M, Bräutigam M, Schmid CE, Kapp M, Knippers R. 1996.**
8 Comparison of two DNA viruses infecting the marine brown algae *Ectocarpus siliculosus*
9 and *E. fasciculatus*. *Journal of General Virology* **77**: 2329-2333.
- 10 **Müller DG, Eichenberger W. 1997.** Mendelian genetics in brown algae: inheritance of a
11 lipid defect mutation and sex alleles in *Ectocarpus siliculosus* (Ectocarpales,
12 Phaeophyceae). *Phycologia* **36**: 79-81.
- 13 **Müller DG, Kapp M, Knippers R. 1998.** Viruses in marine brown algae. *Advances in Virus*
14 *Research* **50**: 50-67.
- 15 **Müller DG, Küpper FC, Küpper H. 1999.** Infection experiments reveal broad host ranges
16 of *Eurychasma dicksonii* (Oomycota) and *Chytridium polysiphoniae* (Chytridiomycota),
17 two eukaryotic parasites in marine brown algae (Phaeophyceae). *Phycological Research*
18 **47** : 217-223.
- 19 **Müller DG, Westermeier R, Morales J, Garcia Reina G, Del Campo E, Correa JA,**
20 **Rometsch E. 2000.** Massive prevalence of viral DNA in *Ectocarpus* (Phaeophyceae,
21 Ectocarpales) from two habitats in the North Atlantic and South Pacific. *Botanica Marina*
22 **43**: 157-159.
- 23 **Müller DG, Knippers R. 2001.** Phaeovirus, Phycodnaviridae. In: Tidona CA, Darai G, eds.
24 *The Springer Index of Viruses*. Springer, Berlin, Heidelberg, New York, 732-736.

- 1 **Oliveira L, Bisalputra T. 1973.** Studies in the brown alga *Ectocarpus* in culture. *Journal of*
2 *Submicrobial Cytology* **5**: 107-120.
- 3 **Oliveira L, Bisalputra T. 1977a.** Ultrastructural studies in the brown alga *Ectocarpus* in
4 culture: autolysis. *New Phytologist* **78**: 139-145.
- 5 **Oliveira L, Bisalputra T. 1977b.** Ultrastructural studies in the brown alga *Ectocarpus* in
6 culture: ageing. *New Phytologist* **78**: 131-138.
- 7 **Oltmanns F. 1899.** Ueber die Sexualität der Ectocarpeen. *Flora* **86**: 86-99.
- 8 **Papenfuss GF. 1935.** Alternation of generations in *Ectocarpus siliculosus*. *Botanical Gazette*
9 **96**: 421-446.
- 10 **Pedersén M. 1968.** *Ectocarpus fasciculatus*: marine brown alga requiring kinetin. *Nature*
11 **218**: 776.
- 12 **Pedersén M. 1973.** Identification of a cytokinin, 6-(3 methyl-2-butenylamino) purine, in sea
13 water and the effect of cytokinins on brown algae. *Physiologia Plantarum* **28**: 101-105.
- 14 **Pedersén PM. 1989.** Studies on *Kuckuckia spinosa* (Fucophyceae, Sorocarpaceae): life
15 history, temperature gradient experiments, and synonymy. *Nordic Journal of Botany* **9**:
16 443-447.
- 17 **Peters AF, Marie D, Scornet D, Kloareg B, Cock JM. 2004a.** Proposal of *Ectocarpus*
18 *siliculosus* (Ectocarpales, Phaeophyceae) as a model organism for brown algal genetics
19 and genomics. *Journal of Phycology* **50**: 1079-1088.
- 20 **Peters AF, Scornet D, Müller DG, Kloareg B, Cock JM. 2004b.** Inheritance of organelles
21 in artificial hybrids of the isogamous multicellular chromist alga *Ectocarpus siliculosus*
22 (Phaeophyceae). *European Journal of Phycology* **39**: 235-242.
- 23 **Petersen HE. 1905.** Contributions à la connaissance des phycomycètes marins (Chytridinae
24 Fischer). *Oversigt over det kgl. Danske Videnskabernes Selskats Forhandling* **5**: 439-
25 488.

- 1 **Ratray J. 1885.** Note on *Ectocarpus*. *Transactions of the Royal Society of Edinburgh* **32**:
2 589-602.
- 3 **Ravanko O. 1970.** Morphological, developmental, and taxonomic studies in the *Ectocarpus*
4 complex (Phaeophyceae). *Nova Hedwigia* **20**: 79-252.
- 5 **Robledo DR, Sosa PA, Garcia-Reina G, Müller DG. 1994.** Photosynthetic performance of
6 healthy and virus-infected *Feldmannia irregularis* and *F. simplex* (Phaeophyceae).
7 *European Journal of Phycology* **29**: 247-251.
- 8 **Rousseau F, de Reviers B. 1999.** Circumscription of the order Ectocarpales (Phaeophyceae):
9 bibliographical synthesis and molecular evidence. *Cryptogamie, Algologie* **20**: 5-18.
- 10 **Russell G. 1966.** The genus *Ectocarpus* in Britain. I. The attached forms. *Journal of the*
11 *marine biological Association of the United Kingdom* **46**: 267-294.
- 12 **Russell G. 1967a.** The genus *Ectocarpus* in Britain. II. The free-living forms. *Journal of the*
13 *marine biological Association of the United Kingdom* **47**: 233-250.
- 14 **Russell G. 1967b.** The ecology of some free-living Ectocarpaceae. *Helgoländer*
15 *wissenschaftliche Meeresuntersuchungen* **15**: 155-162.
- 16 **Russell G, Morris OP. 1970.** Copper tolerance in the marine fouling alga *Ectocarpus*
17 *siliculosus*. *Nature* **228**: 288-289.
- 18 **Russell G, and Bolton JJ. 1975.** Euryhaline ecotypes of *Ectocarpus siliculosus* (Dillw.)
19 Lyngb. *Estuarine and Coastal Marine Science* **3**: 91-94.
- 20 **Russell G. 1983a.** Parallel growth patterns in algal epiphytes and *Laminaria* blades. *Marine*
21 *Ecology Progress Series* **13**: 303-304.
- 22 **Russell G. 1983b.** Formation of an ectocarpoid epiflora on blades of *Laminaria digitata*.
23 *Marine Ecology Progress Series* **11**: 181-187.
- 24 **Sauvageau C. 1896.** Sur la conjugaison des zoospores de l'*Ectocarpus siliculosus*. *Comptes*
25 *rendus de l'Académie des Sciences* **123**: 431-433.

- 1 **Sauvageau C. 1897.** La copulation isogamique de l'*Ectocarpus siliculosus* est-elle apparente
2 ou réelle ? *Mémoires de la Société nationale des sciences naturelles de Cherbourg* **30**:
3 293-304.
- 4 **Schmid R, Dring MJ. 1992.** Circadian rhythm and fast responses to blue light of
5 photosynthesis in *Ectocarpus* (Phaeophyta, Ectocarpales). *Planta* **187**: 53-59.
- 6 **Schmid CE, Schroer N, Kawai H, Müller DG. 1992.** Isolation and biochemical
7 characterization of different gamete membranes in the chromophyte alga *Ectocarpus*
8 *siliculosus* (Phaeophyceae). *Plant Physiology and Biochemistry* **30**: 703-712.
- 9 **Schmid CE. 1993.** Cell-cell-recognition during fertilization in *Ectocarpus siliculosus*
10 (Phaeophyceae). *Hydrobiologia* **260-261**: 437-443.
- 11 **Schmid R, Dring MJ. 1993.** Rapid, blue-light-induced acidifications at the surface of
12 *Ectocarpus* and other marine macroalgae. *Plant Physiology* **101**: 907-913.
- 13 **Schmid CE, Schroer N, Müller DG. 1994a.** Female gamete membrane glycoproteins
14 potentially involved in gamete recognition in *Ectocarpus siliculosus* (Phaeophyceae).
15 *Plant Science* **102**: 61-67.
- 16 **Schmid R, Dring MJ, Forster RM. 1994b.** Kinetics of blue light stimulation and circadian
17 rhythmicity of light-saturated photosynthesis in brown algae: a species comparison.
18 *Journal of Phycology* **30**: 612-621.
- 19 **Schmid CE, Müller DG, Eichenberger W. 1994c.** Isolation and characterization of a new
20 phospholipid from brown algae. Intracellular localization and site of biosynthesis. *Journal*
21 *of Plant Physiology* **143**: 570-574.
- 22 **Schmid R, Dring MJ. 1996.** Influence of carbon supply on the circadian rhythmicity of
23 photosynthesis and its stimulation by blue light in *Ectocarpus siliculosus*: clues to the
24 mechanism of inorganic carbon acquisition in lower brown algae. *Plant, Cell and*
25 *Environment* **19**: 373-382.

- 1 **Schmid R. 1998.** Photosynthesis of *Ectocarpus siliculosus* in red light and after pulses of blue
2 light at high pH - evidence for bicarbonate uptake. *Plant, Cell and Environment* **21**: 523-
3 529.
- 4 **Schmid R, Hillrichs S. 2001.** Uptake and accumulation of inorganic carbon in *Ectocarpus*
5 *siliculosus* and its relation to blue light stimulation of photosynthesis. *European Journal*
6 *of Phycology* **36**: 257-264.
- 7 **Sengco M, Bräutigam M, Kapp M, Müller DG. 1996.** Detection of virus DNA in
8 *Ectocarpus siliculosus* and *E. fasciculatus* (Phaeophyceae) from various geographic areas.
9 *European Journal of Phycology* **31**: 73-78.
- 10 **Silva PC, Basson P, Moe RL. 1996.** Catalogue of the benthic marine algae of the Indian
11 Ocean. *University of California Publications in Botany* **79**: 1-1259.
- 12 **Stache, B. 1990.** Sexual compatibility and species concept in *Ectocarpus siliculosus*
13 (Ectocarpales, Phaeophyceae) from Italy, North Carolina, Chile, and New Zealand. In:
14 Garbary, D. J. & R. G. South (ed). *Evolutionary Biogeography of the Marine Algae of the*
15 *North Atlantic*. Springer Vlg., Berlin: 173-186.
- 16 **Stache-Crain B, Müller DG, Goff LJ. 1997.** Molecular systematics of *Ectocarpus* and
17 *Kuckuckia* (Ectocarpales, Phaeophyceae) inferred from phylogenetic analysis of nuclear
18 and plastid-encoded DNA sequences. *Journal of Phycology* **33**: 152-168.
- 19 **Thomas DN, Kirst GO. 1991a.** Salt tolerance of *Ectocarpus siliculosus* (Dillw.) Lyngb.:
20 comparison of gametophytes, sporophytes and isolates of different geographic origin.
21 *Botanica Acta* **104**: 26-36.
- 22 **Thomas DN, Kirst GO. 1991b.** Differences in osmoacclimation between sporophytes and
23 gametophytes of the brown alga *Ectocarpus siliculosus*. *Physiologia Plantarum* **83**: 281-
24 289.

- 1 **Van Etten JL, Meints RH. 1999.** Giant viruses infecting algae. *Annual Review of*
2 *Microbiology* **53**: 447-494.
- 3 **Van Etten JL, Graves MV, Müller DG, Boland W, Delaroque N. 2002.** Phycodnaviridae -
4 large DNA algal viruses. *Archives of Virology* **147**: 1479-1516.
- 5 **Van Heerden PDR, Robertson BL, De Kock L. 1997.** Inhibition of *Ectocarpus siliculosus*
6 infestations with copper chloride in tank cultures of *Gracilaria gracilis*. *Journal of*
7 *Applied Phycology* **9**: 255-259.
- 8 **West JA, Kraft GT. 1996.** *Ectocarpus siliculosus* (Dillwyn) Lyngb. from Hopkins River
9 Falls, Victoria - the first record of a freshwater brown alga in Australia. *Muelleria* **9**: 29-
10 33.
- 11 **Wiencke C, Clayton MN. 2002.** Antarctic seaweeds. Synopses of the Antarctic benthos, vol.
12 9. Gantner, Ruggell, Liechtenstein. 159 pp.
- 13 **Wright EP. 1879.** On a species of *Rhizophyidium* parasitic on species of *Ectocarpus*, with
14 notes on the fructification of the Ectocarpi. *Transactions of the Royal Irish Academy* **26**:
15 369-379.

17 **Figure legends:**

19 **Figure 1: Phylogeny of brown algae and Ectocarpales.**

20 **a:** Position of brown algae within the eukaryotes (adapted from Baldauf, 2003). Brown algae
21 belong to the heterokont phylum, which is phylogenetically distant from land plants and the
22 green and red algae. Photosynthetic organisms are framed. **b:** Position of the Ectocarpales (in
23 bold) within the brown algae (adapted from Kawai *et al.*, 2007).

Figure 2: Life cycle of *Ectocarpus siliculosus*.

Diploid sporophytes produce meio-spores (by meiosis) in unilocular sporangia (UL). Meio-spores grow into male or female gametophytes (dioecism). Gametophytes produce gametes in plurilocular gametangia (PL). Fusion of gametes produces a zygote that grows into a diploid sporophyte, completing the sexual cycle. Unfused gametes may grow parthenogenetically and form a parthenosporophyte, which is indistinguishable from the diploid sporophyte. Both sporophytes and parthenosporophytes can reproduce themselves asexually by the production of mito-spores in plurilocular sporangia.

Figure 3: Morphology of *Ectocarpus siliculosus*.

Photographies of one week old vegetative sporophyte (a) and gametophyte (b) and schemes representing the whole body of the mature sporophyte (c) and gametophyte (d) after six weeks of growth, are shown. e: Plurilocular sporangium or gametangium (occurring on the sporophyte and the gametophyte, respectively) before (left) and after (right) release of zoids. f: Unilocular sporangium from sporophyte. Sporangia and gametangia can be either sessile or pedicellate (Kim & Lee, 1992).

Figure 4: General ultrastructure of a vegetative cell of *Ectocarpus siliculosus*.

The general ultrastructure of a vegetative cell is similar in both prostrate and erect filaments (Oliveira & Bisalputra, 1973). The different compartments of the cell are illustrated (see text for details). Lines represent membranes and define sub-cellular compartments, except for thylakoids, drawn as a thick black line. Depending on their type and age, vegetative cell size varies from 10 to 35 μm in length, and 5 to 15 μm in width (in laboratory culture conditions).

Figure 5: General intracellular structure of *Ectocarpus siliculosus* zoids.

1 A scheme representing the different compartments of an *Ectocarpus siliculosus* zoid cell from
2 a plurilocular sporangium or gametangium is presented (see text for details). Legends are the
3 same as in fig. 4.

4

5 **Acknowledgements**

6 We are grateful to N. Delaroque (Max Plank Institute in Jena) for authorisation to
7 communicate unpublished information about viral integration, and to Declan Schroeder
8 (Marine Biological Association in Plymouth) for authorisation to mention the microarray for
9 the EsV-1 virus.

Figure 1

Figure 1: Phylogeny of brown algae and Ectocarpales. a: Position of brown algae within the eukaryotes (adapted from Baldauf, 2003). Brown algae belong to the heterokont phylum, which is phylogenetically distant from land plants and the green and red algae. Photosynthetic organisms are framed. b: Position of the Ectocarpales (in bold) within the brown algae (adapted from Kawai et al., 2007).

275x190mm (300 x 300 DPI)

Figure 2

Figure 2: Life cycle of *Ectocarpus siliculosus*. Diploid sporophytes produce meio-spores (by meiosis) in unilocular sporangia (UL). Meio-spores grow into male or female gametophytes (dioecism). Gametophytes produce gametes in plurilocular gametangia (PL). Fusion of gametes produces a zygote that grows into a diploid sporophyte, completing the sexual cycle. Unfused gametes may grow parthenogenetically and form a parthenosporophyte, which is indistinguishable from the diploid sporophyte. Both sporophytes and parthenosporophytes can reproduce themselves asexually by the production of mito-spores in plurilocular sporangia.

254x190mm (300 x 300 DPI)

Figure 3

Figure 3: Morphology of *Ectocarpus siliculosus*. Photographies of one week old vegetative sporophyte (a) and gametophyte (b) and schemes representing the whole body of the mature sporophyte (c) and gametophyte (d) after six weeks of growth, are shown. e: Plurilocular sporangium or gametangium (occurring on the sporophyte and the gametophyte, respectively) before (left) and after (right) release of zoids. f: Unilocular sporangium from sporophyte. Sporangia and gametangia can be either sessile or pedicellate (Kim & Lee, 1992).

254x190mm (300 x 300 DPI)

Figure 4: General ultrastructure of a vegetative cell of *Ectocarpus siliculosus*. The general ultrastructure of a vegetative cell is similar in both prostrate and erect filaments (Oliveira & Bisalputra, 1973). The different compartments of the cell are illustrated (see text for details). Lines represent membranes and define sub-cellular compartments, except for thylakoids, drawn as a thick black line. Depending on their type and age, vegetative cell size varies from 10 to 35 μ m in length, and 5 to 15 μ m in width (in laboratory culture conditions).

254x190mm (300 x 300 DPI)

Figure 5: General intracellular structure of *Ectocarpus siliculosus* zoids. A scheme representing the different compartments of an *Ectocarpus siliculosus* zoid cell from a plurilocular sporangium or gametangium is presented (see text for details). Legends are the same as in Fig. 4.

254x190mm (300 x 300 DPI)