Attacking hepatitis B virus cccDNA—The holy grail to hepatitis B cure

J. Lucifora, U. Protzer

► To cite this version:

J. Lucifora, U. Protzer. Attacking hepatitis B virus cccDNA—The holy grail to hepatitis B cure. Journal of Hepatology, 2016, 64 (1), pp.S41-S48. 10.1016/j.jhep.2016.02.009 . hal-01812699

HAL Id: hal-01812699

<https://hal.science/hal-01812699>

Submitted on 30 Aug 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Attacking Hepatitis B Virus cccDNA - the Holy Grail to Hepatitis B Cure

Julie Lucifora^{1, 2, 3} and Ulrike Protzer^{4, 5, #}

¹ Cancer Research Center of Lyon (CRCL), Lyon, 69008, France;

² INSERM U1052, CNRS UMR-5286, Lyon, 69008, France;

³ University of Lyon, Université Claude-Bernard (UCBL), 69008 Lyon, France;

⁴ Institute of Virology, Technische Universität München / Helmholtz Zentrum München, Trogerstrasse 30, 81675 Munich, Germany;

⁵ German Center for Infection Research (DZIF), Munich site

Contact Information:

Prof. Ulrike Protzer, MD

Institute of Virology

Technische Universität München / Helmholtz Zentrum München

Trogerstr. 30, D-81675 München

Germany

Tel: +498941406886

Fax: +498941406823

protzer@tum.de

word count: 3212 (excluding references)

figure numbers: 3

table number: 1

list of abbreviations: Hepatitis B virus (HBV), relaxed circular DNA (rcDNA), primary human hepatocyte (PHH), differentiated HepaRG (dHepaRG), Na⁺-taurocholate cotransporting peptide (NTCP), humanized liver-chimeric (HuHEP), Rolling circle amplification (RCA), protein-free rcDNA (pf-rcDNA), tyrosyl-DNA-phosphodiesterase (TDP), disubstituted sulfonamides (DSS), interferon alfa (IFN- α), tumor necrosis factor (TNF), lymphotoxin beta receptor (LT β R), APOBEC3A (A3A), APOBEC3B (A3B), interferon gamma (IFN γ), TAL effector nucleases (TALENs).

key words: hepatitis B virus, cccDNA, liver

no conflict of interest

financial support: Work in the authors' laboratories was supported by the European Association of the Study of Liver Diseases (EASL), the German Center for Infection Research (DZIF), the French Agency for Research on AIDS and Hepatitis Viruses (ANRS), the FINOVI Foundation, the French Association for Research on Cancer (ARC), the Helmholtz Association, the German Ministry for Research and Education (BMBF), the German Ministry for Economics (BMWi) and the German Research Foundation (DFG).

Abstract

HBV deposits a covalently closed circular DNA form, called cccDNA, in the nucleus of infected cells. As the central transcription template, the cccDNA mini-chromosome is a key intermediate in the HBV life cycle. Being hard to target in the nucleus by antivirals, cccDNA is responsible for chronicity of HBV infection. While little is known about the mechanisms involved in cccDNA formation, data are currently accumulating on control mechanisms involved in regulating transcription from cccDNA, and the first potential targeting approaches have already been reported. This review will summarize our knowledge about cccDNA biology and latest advances in strategies with the potential to target cccDNA and to finally achieve an HBV cure.

Introduction

Hepatitis B virus (HBV) has optimized its life cycle for long-term persistence in the liver. To achieve this, the virus establishes a plasmid-like covalently closed circular DNA form, the so-called cccDNA. HBV cccDNA serves as template for transcription of all HBV RNAs and production of progeny virus. HBV cccDNA resides in the nucleus of infected cells as an episomal (i.e. non-integrated) DNA associated to histones. It drives production of viral antigens such as HBeAg and HBsAg, and progeny virions. In contrast to cccDNA, the DNA contained in infectious virions is an only partially double stranded, relaxed circular DNA (rcDNA) that results from reverse transcription of a pregenomic RNA.

Available antivirals cannot cure hepatitis B because they target reverse transcription as a late step in the HBV life cycle and do not affect the cccDNA transcription template. Nucleos(t)ide analogues such as lamivudine, entecavir or tenofovir can efficiently control progeny release and thus viremia, but neither cccDNA persistence nor HBsAg or HBeAg expression and secretion. Similarly, neither entry inhibitors nor capsid assembly modifiers as novel antivirals under clinical development target HBV cccDNA (see *"New antiviral targets for novel treatment concepts for HBV"*, Durantel and Zoulim). Therefore, long-term treatment is required, which is expensive, challenges patient adherence and may lead to concomitant resistance [1]. Interferon (IFN)- α , which is also licensed for hepatitis B therapy and has direct antiviral as well as immune modulatory effects, may result in HBsAg loss and even virus clearance in a proportion of patients, but dosing is limited by severe side effects [2]. Therefore, efficient removal of cccDNA from infected hepatocytes is regarded as the "holy grail" to achieve HBV cure. This review will summarize the different experimental models and detection assays available to study cccDNA, our current knowledge on cccDNA biology as well as potential therapeutic approaches to attack cccDNA.

Experimental models to study the cccDNA fate

For a long time, *in vitro* and *in vivo* models to study the fate of HBV cccDNA were rather limited (see *Experimental models (cell culture and animals) for studies of human HBV pathobiology and new treatment concepts*, M Dandri). Only primary tupaia or human hepatocytes (PHH) and differentiated HepaRG (dHepaRG) were used allowing HBV infection

and thus replication of HBV from its natural transcription template, the cccDNA. These models, however, are very demanding. The recent identification of the Na⁺-taurocholate cotransporting peptide (NTCP) as a crucial HBV receptor [3] allowed establishing new cell lines that support HBV infection and the study of cccDNA *in vitro*.

While short-term experiments were enabled by an improved HBV infection rate (and consequently higher levels of cccDNA) in NTCP-overexpressing HuH7, HepaRG and in particular HepG2 cells [3, 4], experiments to study the fate and regulation of cccDNA over time remain challenging. Only HepaRG cells can be kept infected in culture for more than a month, but they differentiate into two different cell types (biliary like and hepatocyte like cells) and establish only low levels of cccDNA (see **Table 1**). Other cell lines keep dividing and cell division results in loss of cccDNA (see below). This technically challenges our research. It is furthermore important to note that despite recent improvements, *in vitro* HBV infections are still rather inefficient, requiring a huge excess of virus (100-10,000 virions/cell depending on the virus inoculum, the cell type and infection protocol used) and polyethylene glycol to facilitate virus fusion. Moreover, there is no evidence of HBV spread in cell culture – unlike for other viruses.

In vivo models are limited by the strict species specificity of HBV that only infects human and humanoid primates. Studies using chimpanzees largely contributed to our understanding of the virus-host interaction [5] but have since been banned [6]. While chimpanzee hepatocytes support HBV replication [7-9], monkeys such as macaques or baboons frequently used for research cannot be infected by HBV due to differences in the NTCP protein sequence [10]. Macaques expressing the human NTCP will, however, constitute a very promising experimental model. Mouse hepatocytes grafted with NTCP support HBV entry but do not allow cccDNA establishment [11], probably because they lack key factors involved in nuclear transport of rcDNA or rcDNA to cccDNA conversion. Humanized liver-chimeric (HuHEP) mice allow long-term study of cccDNA [13] but are immune deficient and due to technical and cost issues cannot not be used for large-scale experiments. Understanding cccDNA biology, regulation and fate would thus greatly benefit from further improvements of the HBV infection models.

The challenging detection of HBV cccDNA

With low copy numbers of cccDNA *in vitro* (Table 1) and *in vivo*, detection of cccDNA is a huge challenge for the HBV research field. Mean values from 0.01 to 1 copies per cell in livers of HBV-infected patients [14] or HuHEP mice [13] have been reported. This means that only a portion of the cells were actually infected by HBV, but methods to detect cccDNA on a per-cell level are lacking. Moreover, true discrimination of cccDNA from almost identical viral linear DNA or rcDNA increases the challenge, because these exist in vast excess in virions and in viral capsids in the cytoplasm of infected cells. While Southern blot analyses can be used to distinguish between relaxed rcDNA, linear and supercoiled cccDNA, the sensitivity of this technique is rather low. To increase sensitivity, cccDNA selective qPCR methods have been developed that are based on the use of primers (and probes) spanning the nick in rcDNA and hybridizing in its “gap region” (Figure 1). With the amazing technical development in PCR-based methods over the past years [15], the sensitivity of cccDNA detection is no longer an issue. Specificity of the cccDNA PCR methods, however, remains a big concern and correct quantification of cccDNA is hampered by false positive detection of rcDNA HBV genomes that are present in vast excess in infected cells. Rolling circle amplification (RCA) coupled or not with nested PCR has been described to increase specificity of cccDNA detection [16-18] but has not been widely accepted. Extraction of nuclear DNA only [19], plasmid safe DNase that digests linear viral DNA, but not rcDNA [20] and T5 exonuclease that degrades rcDNA but should leave cccDNA molecules intact (Xia et al, Hepatitis B Virus: Methods and Protocols, in press) can also be used to increase specificity. Of note, these exonucleases have different efficacy depending on the type and amount of “contaminating” DNA. So far, laboratories apply different protocols and a consensus or standard has yet to be reached. This, however, should be of high priority given that also the protocol of DNA extraction may influence cccDNA detection by generating nicks in cccDNA molecules that may render them susceptible to exonuclease treatment, may prevent RCA detection and may change the running behavior in gels before Southern blotting.

Establishment of HBV cccDNA in the nucleus

Upon infection, cccDNA is formed as a plasmid-like episome in the host cell nucleus from the

rcDNA genome in incoming virions that is linked to the viral polymerase protein. Alternatively, cccDNA may be established from capsid rcDNA that can be reimported into the nucleus after it has been newly formed in infected cells. cccDNA was recently detected in a mouse hepatocyte cell line immortalized by stable expression of the human transforming growth factor alpha [12] supporting the fact that human host factors are essential for cccDNA formation and more precisely for destabilization of mature nucleocapsids. Nuclear localization signals within the arginine-rich C-terminal domain of the HBV core protein [21] trigger transport of the rcDNA-containing HBV capsid to the nucleus where the capsid attaches to nucleoporin 153 and is disintegrated [22]. Upon translocation into the nucleus, the conversion of the rcDNA genome into cccDNA takes place via a multi-step process that remains largely obscure although it has been linked to DNA repair mechanisms [23].

Firstly, the rcDNA needs to be detached from the HBV polymerase and its RNA primer that are remnants from reverse transcription that took place within the viral capsid. Hereby, a protein-free rcDNA form (pf-rcDNA, equivalent to DP-rcDNA for deproteinized rcDNA) is generated. Pf-rcDNA may persist in the nucleus before conversion into cccDNA, and has even been observed in the cytoplasm [24-26]. Secondly, the incompleteness of the rcDNA (+) strand (the “gap”) and the terminal redundancy in its (-) strand (the “nick”) must be repaired (**Figure 2**). It has recently been demonstrated that tyrosyl-DNA-phosphodiesterase (TDP) 2 can specifically cleave the Tyr-DNA bond to release the polymerase protein from HBV rcDNA [27]. The fact that TDP2 knockout HepG2-NTCP cells engineered by a CRISPR/Cas9-based approach still proved susceptible to HBV infection [28] indicated that alternative pathway(s) exist involving nucleolytic cleavage by (a) still elusive endonuclease(s). Comparison of HBV and duck hepatitis B virus cccDNA formation revealed an unexpected contribution of the virus itself to cccDNA formation. DHBV but not HBV rcDNA converted efficiently into cccDNA in the same cells [20]. Linear HBV DNA, a probably accidental by-product of reverse transcription, cannot be transcribed per se but may form cccDNA-like molecules by non-homologous end joining. Most of these, however, will be functionally defective [23].

Structure and regulation of the HBV cccDNA

Once formed, cccDNA persists as an episome in the nucleus of infected cells and viral mRNAs are transcribed by the cellular RNA polymerase II. The cccDNA molecule is organized into a

chromatin-like structure that displays the typical beads-on-a-string arrangement in electron microscopy [29, 30]. Attached proteins are host histone but also non-histone proteins [31]. During the last years major efforts have been made to demonstrate that the transcriptional activity of cccDNA is subject to the “histone code” both *in vitro* and *in vivo* [19, 32, 33]. Using ChIP approaches, the recruitment of H3 and H4 histones, as well as that of several cellular transcription factors (including CREB, ATF, STAT1/2) and histone modifying enzymes (such as CBP, p300, PCAF/GCN5, HDAC1, hSirt1, PRMT1) has been demonstrated and was associated with modulation of cccDNA transcriptional activity [19, 32, 34]. Using a chromosome immunoprecipitation approach followed by next generation sequencing, the first genome-wide map of histone posttranslational modifications in cccDNA-associated chromatin was established [35]. This revealed that HBV chromatin shares some basic features with cellular chromatin, in particular marks linked to active transcription, but lacks significant levels of posttranslational histone modifications associated with transcriptional repression, which, in cellular chromatin, are indispensable for transcriptional control [35]. When applied to *de novo* infected HepG2-NTCP cells or primary human hepatocytes or to HBV-infected liver tissue the analysis revealed general features of cccDNA chromatin organization as well as features that are unique to the respective source of cccDNA [35]. This emphasizes the importance to repeat experiments and to confirm results using different infection models.

Despite being named a “minichromosome”, HBV cccDNA unlike other chromosomes does neither contain an origin of replication nor possess a nuclear retaining signal. It thus cannot replicate by itself in infected hepatocytes and should theoretically be excluded from the nucleus after cell division. cccDNA persistence therefore relies on its stability in the nucleus and on its replenishment via reimport of rcDNA from cytoplasmic capsids after cell division. The half-life of HBV cccDNA has not been exactly defined and varies with inflammatory activity resulting in hepatocyte turnover. Rebound of HBV replication upon withdrawal from nucleos(t)ide analogue treatment [36] and traces of HBV-DNA remaining detectable years after clinical recovery from acute hepatitis [37], however, indicate that HBV cccDNA is very long-lived in the human liver and can persist for decades. *In vitro*, cccDNA also appears to be very stable [38, 39] independently of its transcriptional activity as demonstrated in HepaRG cells infected with an HBx-deficient HBV unable to transcribe mRNA [40]. Since it is not self-maintained, cccDNA should theoretically be lost by cell division e.g. in case of liver regeneration. However, cccDNA may survive through mitosis even though it has been

1 observed that induction of cell division results in a dramatic decrease of cccDNA [41-45]. This
2 may be due to a “refill” via reimport of rcDNA from newly formed viral capsids into the
3 nucleus.
4
5
6

7 A combination of *in vitro* approaches, including electromobility shift and DNA protection
8 assays, has shown that a number of transcription factors bind to HBV regulatory elements
9 [46]. Among them, HNF1 α and HNF4 α are the crucial nuclear factors for HBV transcription
10 needed at sufficient levels [47, 48]. Viral proteins are also very important in cccDNA
11 structure and control. In particular, the non-structural protein HBx is associated with cccDNA
12 [32] and essential to initiate and maintain transcription from cccDNA [40]. Indeed, using
13 dHepaRG cells in which HBx is expressed in a tetracyclin-regulated fashion and infecting
14 them with an HBx-deficient HBV mutant, we showed that transcription of viral RNA exactly
15 followed the switching on and off of HBx expression [40]. Hereby it appeared that HBV
16 transcription is repressed in the absence of HBx rather than HBx functioning as a typical viral
17 transactivator. This repression is characterized by a more compact structure of cccDNA [49]
18 and associated with repressive epigenetic marks [32, 49] on cccDNA that were absent in the
19 presence of HBx [35]. Interestingly, it was shown that HBx supports HBV gene expression by
20 a conserved mechanism that acts specifically on episomal DNA templates and requires HBx
21 binding to the cellular DDB1 E3 ubiquitin ligase [50]. Altogether these data suggest that HBx
22 may block cellular factor(s) inducing mechanisms that inhibit transcription from cccDNA such
23 as SETDB1 [49], Spindlin1 [51] as well as essential other factors that were reported at the
24 recent international HBV meeting and will be published soon. In addition to HBx, the HBV
25 core protein (HBc) binds to cccDNA resulting in a rearrangement of the nucleosomal spacing
26 of the HBV nucleoprotein complexes [31]. Whether HBc also regulates transcription from
27 cccDNA is still debated. On one hand, it was shown that HBc alters the cccDNA methylation
28 profile and therefore maintains a permissive epigenetic state in a critical region of the HBV
29 cccDNA [52]. On the other hand, a decreased recruitment of HBc on HBVx- cccDNA was
30 recently described but did not account for the failure in cccDNA transcription [49]. The origin
31 of HBc attached to cccDNA (i.e. whether it stems from the incoming virus or from viral
32 replication), the kinetic of its recruitment onto cccDNA and its stability once bound to
33 cccDNA remain unresolved questions that would certainly help to understand its role on
34 cccDNA regulation. Noteworthy, as HBc is important for cccDNA structure and/or control,
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

capsid assembly modifiers (CpAMs) that are currently in clinical development were hypothesized to also have an effect on cccDNA [62]. This, however, needs to be proven.

Targeting HBV cccDNA to achieve HBV cure

Different options to target cccDNA to achieve affect HBV persistence are available: inhibition of its formation, inhibition of its function or inducing its degradation (**Figure 3**).

Two structurally related disubstituted sulfonamides (DSS) were shown to interfere with rcDNA conversion into cccDNA [38]. Given the proposed contribution of the DNA-damage response in this process, viral proteins or substances interfering with the DNA-damage response should also interfere with rcDNA to cccDNA conversion. Considering the long half-life of cccDNA, however, these compounds should only have an effect if given during the establishment of HBV infection or during phases of high hepatocyte turnover.

Once cccDNA is formed its transcription can be inhibited by cytokines. For instance, we demonstrated that IL-6 inhibits HBV transcription from cccDNA by reducing the levels of the essential transcription factors HNF1 α and HNF4 α through activation of the MAP kinase pathway [53]. This results in a loss of HNF1 α and HNF4 α binding to the cccDNA, which is accompanied by a redistribution of STAT3 binding from cccDNA to cellular IL-6 target genes [48]. It has also been shown that cccDNA transcription can be silenced by the use of IFN- α or small molecules targeting chromatin modifying enzymes [34, 54-56]. Genome-wide mapping of histone modifications revealed that transcription and active modifications in HBV chromatin can be reduced when innate immunity pathways are activated, or by a small molecule epigenetic modifying agent [35]. This supports the idea that chromatin-based regulation of cccDNA transcription could be of therapeutic use. A better understanding of the function of HBx protein and its interaction with DDB1 in the E3 ubiquitin ligase complex [57], that potentially results in degradation of host proteins able to control HBV, may allow to identify new targets for antiviral drug discovery. Because these strategies that aim at inhibiting transcription from cccDNA will likely only have a transient effect, they may help to achieve at least a functional cure of HBV.

Purging cccDNA from the nucleus of infected cells is a much more promising approach since this will result in a sustained control of HBV or even in HBV cure. In chimpanzees, it has been

1 shown that clearance of HBV replication, but also HBV cccDNA, occurs before full-blown
2 cytotoxicity of T cells hits the liver in acute hepatitis B [5]. This indicates that the cccDNA
3 pool can be affected by non-cytolytic mechanisms involving cytokines such as IFNs and
4 tumor necrosis factor (TNF) [5]. Recently, we showed that induction of intrahepatic antiviral
5 immune responses (through high doses of IFN- α or LT β R-agonists) is able to trigger non-
6 cytolytic degradation of cccDNA from infected cells *in vitro* [58]. Hereby, activation of
7 nuclear deaminases such as APOBEC3A (A3A) and APOBEC3B (A3B) results in cccDNA
8 deamination and a significant reduction of cccDNA [58]. Although A3A or A3B-mediated
9 deamination was essential, the exact pathways and enzymes involved in downstream
10 degradation of HBV-DNA still need to be deciphered. They are likely contributing to base-
11 excision or nucleotide-excision repair pathways. It should also be determined if cccDNA can
12 be fully purged or if a portion might be refractory to the treatments. Next we examined
13 whether the mechanism of A3A/B-mediated deamination might contribute to T cell-
14 mediated loss of cccDNA. We found that T-cell derived cytokines IFN γ and TNF induce A3A,
15 and to a minor extend also A3B in a synergistic fashion [59]. To study whether these and
16 eventually also other T cell-derived cytokines contribute to cccDNA loss in a non-cytolytic
17 fashion, we set up a transwell system, which allowed us to separate cytolytic and non-
18 cytolytic T cell functions. Hereby, we took advantage of chimeric antigen receptor (S-CAR)
19 grafted T cells [60, 61] that recognize HBsAg on the surface of infected cells but could
20 confirm the findings when T cells carried the natural T cell receptor. While upon direct
21 contact in co-culture T cells killed HBV infected cells, they were only able to release
22 cytokines when separated by the transwell. cccDNA loss, however, was observed in both
23 settings and clearly A3A/A3B dependent [59]. Thus, HBV-specific T cells were able to
24 eliminate HBV in a cytolytic and in a non-cytolytic fashion. This shows that cccDNA
25 degradation in a non-cytolytic fashion is possible and may support HBV cure approaches by
26 therapeutic vaccination, T cell redirection or other immune cell activation strategies such as
27 e.g. PD-1 or PD-ligand1 blockade (see “*New antiviral targets for novel treatment concepts for*
28 *HBV*”, Durantel and Zoulim).

29 Finally, as an alternative to immune activation that may result in tissue damage on the long
30 term if not successful in eliminating HBV, DNA cleaving enzymes, including homing
31 endonucleases or meganucleases, zinc-finger nucleases, TAL effector nucleases (TALENs),
32 and CRISPR-associated system 9 (Cas9) proteins, specifically targeted to the HBV cccDNA are

currently being engineered [63-69]. Clinical development of these interesting strategies, however, is limited by potential off-target effects resulting in cellular DNA damage. In addition, efficient delivery strategies targeting all infected hepatocytes are lacking and therapeutic application would currently require a gene therapy approach.

Taken together, further basic research and more detailed molecular studies are needed to evaluate the translational potential of the novel antiviral strategies mentioned to finally achieve a true HBV cure.

Acknowledgments

The authors thank Judith Fresquet for help with figure 1.

References

Author names in bold designate shared co-first authorship

- [1] Trepo C, Chan HL, Lok A. Hepatitis B virus infection. *Lancet* 2014.
- [2] Wursthorn K, Lutgehetmann M, Dandri M, Volz T, Buggisch P, Zollner B, et al. Peginterferon alpha-2b plus adefovir induce strong cccDNA decline and HBsAg reduction in patients with chronic hepatitis B. *Hepatology* 2006;44:675-684.
- [3] **Yan H, Zhong G**, Xu G, He W, Jing Z, Gao Z, et al. Sodium taurocholate cotransporting polypeptide is a functional receptor for human hepatitis B and D virus. *eLife* 2012;1:e00049.
- [4] Ni Y, Lempp FA, Mehrle S, Nkongolo S, Kaufman C, Falth M, et al. Hepatitis B and D viruses exploit sodium taurocholate co-transporting polypeptide for species-specific entry into hepatocytes. *Gastroenterology* 2014;146:1070-1083.
- [5] Guidotti LG, Rochford R, Chung J, Shapiro M, Purcell R, Chisari FV. Viral clearance without destruction of infected cells during acute HBV infection. *Science* 1999;284:825-829.
- [6] Harrington M. State of the (research) chimp. *Lab animal* 2012;41:31.
- [7] Gheit T, Sekkat S, Cova L, Chevallier M, Petit MA, Hantz O, et al. Experimental transfection of *Macaca sylvanus* with cloned human hepatitis B virus. *J Gen Virol* 2002;83:1645-1649.
- [8] **Lucifora J, Vincent IE**, Berthillon P, Dupinay T, Michelet M, Protzer U, et al. Hepatitis B virus replication in primary macaque hepatocytes: crossing the species barrier toward a new small primate model. *Hepatology* 2010;51:1954-1960.
- [9] **Dupinay T, Gheit T**, Roques P, Cova L, Chevallier-Queyron P, Tasahsu SI, et al. Discovery of naturally occurring transmissible chronic hepatitis B virus infection among *Macaca fascicularis* from Mauritius Island. *Hepatology* 2013;58:1610-1620.
- [10] Yan H, Peng B, He W, Zhong G, Qi Y, Ren B, et al. Molecular determinants of hepatitis B and D virus entry restriction in mouse sodium taurocholate cotransporting polypeptide. *J Virol* 2013;87:7977-7991.

- [11] Li H, Zhuang Q, Wang Y, Zhang T, Zhao J, Zhang Y, et al. HBV life cycle is restricted in mouse hepatocytes expressing human NTCP. *Cellular & molecular immunology* 2014;11:175-183.
- [12] Cui X, Guo JT, Hu J. Hepatitis B Virus Covalently Closed Circular DNA Formation in Immortalized Mouse Hepatocytes Associated with Nucleocapsid Destabilization. *J Virol* 2015;89:9021-9028.
- [13] Allweiss L, Volz T, Lutgehetmann M, Giersch K, Bornscheuer T, Lohse AW, et al. Immune cell responses are not required to induce substantial hepatitis B virus antigen decline during pegylated interferon-alpha administration. *J Hepatol* 2014;60:500-507.
- [14] Werle-Lapostolle B, Bowden S, Locarnini S, Wursthorn K, Petersen J, Lau G, et al. Persistence of cccDNA during the natural history of chronic hepatitis B and decline during adefovir dipivoxil therapy. *Gastroenterology* 2004;126:1750-1758.
- [15] Mu D, Yan L, Tang H, Liao Y. A sensitive and accurate quantification method for the detection of hepatitis B virus covalently closed circular DNA by the application of a droplet digital polymerase chain reaction amplification system. *Biotechnol Lett* 2015;37:2063-2073.
- [16] Margeridon S, Carrouee-Durantel S, Chemin I, Barraud L, Zoulim F, Treppe C, et al. Rolling circle amplification, a powerful tool for genetic and functional studies of complete hepatitis B virus genomes from low-level infections and for directly probing covalently closed circular DNA. *Antimicrob Agents Chemother* 2008;52:3068-3073.
- [17] **Zhong Y, Han J**, Zou Z, Liu S, Tang B, Ren X, et al. Quantitation of HBV covalently closed circular DNA in micro formalin fixed paraffin-embedded liver tissue using rolling circle amplification in combination with real-time PCR. *Clin Chim Acta* 2011;412:1905-1911.
- [18] **Zhong Y, Hu S, Xu C**, Zhao Y, Xu D, Zhao Y, et al. A novel method for detection of HBVcccDNA in hepatocytes using rolling circle amplification combined with in situ PCR. *BMC Infect Dis* 2014;14:608.
- [19] Pollicino T, Belloni L, Raffa G, Pediconi N, Squadrito G, Raimondo G, et al. Hepatitis B virus replication is regulated by the acetylation status of hepatitis B virus cccDNA-bound H3 and H4 histones. *Gastroenterology* 2006;130:823-837.
- [20] Kock J, Rosler C, Zhang JJ, Blum HE, Nassal M, Thoma C. Generation of covalently closed circular DNA of hepatitis B viruses via intracellular recycling is regulated in a virus specific manner. *PLoS Pathog* 2010;6:e1001082.
- [21] **Li HC, Huang EY**, Su PY, Wu SY, Yang CC, Lin YS, et al. Nuclear export and import of human hepatitis B virus capsid protein and particles. *PLoS Pathog* 2010;6:e1001162.
- [22] **Schmitz A, Schwarz A**, Foss M, Zhou L, Rabe B, Hoellenriegel J, et al. Nucleoporin 153 arrests the nuclear import of hepatitis B virus capsids in the nuclear basket. *PLoS Pathog* 2010;6:e1000741.
- [23] Nassal M. HBV cccDNA: viral persistence reservoir and key obstacle for a cure of chronic hepatitis B. *Gut* 2015;64:1972-1984.
- [24] Guo H, Jiang D, Zhou T, Cuconati A, Block TM, Guo JT. Characterization of the intracellular deproteinized relaxed circular DNA of hepatitis B virus: an intermediate of covalently closed circular DNA formation. *J Virol* 2007;81:12472-12484.
- [25] Guo H, Mao R, Block TM, Guo JT. Production and function of the cytoplasmic deproteinized relaxed circular DNA of hepadnaviruses. *J Virol* 2010;84:387-396.
- [26] **Guo H, Xu C**, Zhou T, Block TM, Guo JT. Characterization of the host factors required for hepadnavirus covalently closed circular (ccc) DNA formation. *PLoS One* 2012;7:e43270.

- [27] Koniger C, Wingert I, Marsmann M, Rosler C, Beck J, Nassal M. Involvement of the host DNA-repair enzyme TDP2 in formation of the covalently closed circular DNA persistence reservoir of hepatitis B viruses. *Proc Natl Acad Sci U S A* 2014;111:E4244-4253.
- [28] **Cui X, McAllister R**, Boregowda R, Sohn JA, Ledesma FC, Caldecott KW, et al. Does Tyrosyl DNA Phosphodiesterase-2 Play a Role in Hepatitis B Virus Genome Repair? *PLoS One* 2015;10:e0128401.
- [29] Newbold JE, Xin H, Tencza M, Sherman G, Dean J, Bowden S, et al. The covalently closed duplex form of the hepadnavirus genome exists in situ as a heterogeneous population of viral minichromosomes. *J Virol* 1995;69:3350-3357.
- [30] Bock CT, Schranz P, Schroder CH, Zentgraf H. Hepatitis B virus genome is organized into nucleosomes in the nucleus of the infected cell. *Virus Genes* 1994;8:215-229.
- [31] Bock CT, Schwinn S, Locarnini S, Fyfe J, Manns MP, Trautwein C, et al. Structural organization of the hepatitis B virus minichromosome. *Journal of molecular biology* 2001;307:183-196.
- [32] Belloni L, Pollicino T, De Nicola F, Guerrieri F, Raffa G, Fanciulli M, et al. Nuclear HBx binds the HBV minichromosome and modifies the epigenetic regulation of cccDNA function. *Proc Natl Acad Sci U S A* 2009;106:19975-19979.
- [33] Benhenda S, Ducroux A, Riviere L, Sobhian B, Ward MD, Dion S, et al. Methyltransferase PRMT1 is a binding partner of HBx and a negative regulator of hepatitis B virus transcription. *J Virol* 2013;87:4360-4371.
- [34] Belloni L, Allweiss L, Guerrieri F, Pediconi N, Volz T, Pollicino T, et al. IFN-alpha inhibits HBV transcription and replication in cell culture and in humanized mice by targeting the epigenetic regulation of the nuclear cccDNA minichromosome. *J Clin Invest* 2012;122:529-537.
- [35] Tropberger P, Mercier A, Robinson M, Zhong W, Ganem DE, Holdorf M. Mapping of histone modifications in episomal HBV cccDNA uncovers an unusual chromatin organization amenable to epigenetic manipulation. *Proc Natl Acad Sci U S A* 2015;112:E5715-5724.
- [36] Zoulim F. Hepatitis B virus resistance to antiviral drugs: where are we going? *Liver Int* 2011;31 Suppl 1:111-116.
- [37] Rehmann B, Ferrari C, Pasquinelli C, Chisari FV. The hepatitis B virus persists for decades after patients' recovery from acute viral hepatitis despite active maintenance of a cytotoxic T-lymphocyte response. *Nat Med* 1996;2:1104-1108.
- [38] Cai D, Mills C, Yu W, Yan R, Aldrich CE, Saputelli JR, et al. Identification of disubstituted sulfonamide compounds as specific inhibitors of hepatitis B virus covalently closed circular DNA formation. *Antimicrob Agents Chemother* 2012;56:4277-4288.
- [39] **Zhou T, Guo H**, Guo JT, Cuconati A, Mehta A, Block TM. Hepatitis B virus e antigen production is dependent upon covalently closed circular (ccc) DNA in HepAD38 cell cultures and may serve as a cccDNA surrogate in antiviral screening assays. *Antiviral Res* 2006;72:116-124.
- [40] **Lucifora J, Arzberger S**, Durantel D, Belloni L, Strubin M, Levrero M, et al. Hepatitis B virus X protein is essential to initiate and maintain virus replication after infection. *J Hepatol* 2011;55:996-1003.
- [41] **Lutgehetmann M, Volz T**, Kopke A, Broja T, Tigges E, Lohse AW, et al. In vivo proliferation of hepadnavirus-infected hepatocytes induces loss of covalently closed circular DNA in mice. *Hepatology* 2010;52:16-24.
- [42] Guo JT, Pryce M, Wang X, Barrasa MI, Hu J, Seeger C. Conditional replication of duck hepatitis B virus in hepatoma cells. *J Virol* 2003;77:1885-1893.

- [43] **Reaiche-Miller GY, Thorpe M**, Low HC, Qiao Q, Scougall CA, Mason WS, et al. Duck hepatitis B virus covalently closed circular DNA appears to survive hepatocyte mitosis in the growing liver. *Virology* 2013;446:357-364.
- [44] Summers J, Jilbert AR, Yang W, Aldrich CE, Saputelli J, Litwin S, et al. Hepatocyte turnover during resolution of a transient hepadnaviral infection. *Proc Natl Acad Sci U S A* 2003;100:11652-11659.
- [45] Allweiss L, Volz T., Giersch K., Lohse A.W., Petersen J. , Lütgehetmann M. , Dandri M. . 127 PROLIFERATION OF HEPATITIS B VIRUS INFECTED HUMAN HEPATOCYTES INDUCES SUPPRESSION OF VIRAL REPLICATION AND RAPID cccDNA DECREASE IN HUMANIZED MICE. *Journal of Hepatology* 2013;58:S56-S57.
- [46] Quasdorff M, Protzer U. Control of hepatitis B virus at the level of transcription. *J Viral Hepat* 2010;17:527-536.
- [47] Quasdorff M, Hosel M, Odenthal M, Zedler U, Böhne F, Gripon P, et al. A concerted action of HNF4alpha and HNF1alpha links hepatitis B virus replication to hepatocyte differentiation. *Cell Microbiol* 2008.
- [48] **Palumbo GA, Scisciani C**, Pediconi N, Lupacchini L, Alfalate D, Guerrieri F, et al. IL6 Inhibits HBV Transcription by Targeting the Epigenetic Control of the Nuclear cccDNA Minichromosome. *PLoS One* 2015;10:e0142599.
- [49] Riviere L, Gerossier L, Ducroux A, Dion S, Deng Q, Michel ML, et al. HBx relieves chromatin-mediated transcriptional repression of hepatitis B viral cccDNA involving SETDB1 histone methyltransferase. *J Hepatol* 2015;63:1093-1102.
- [50] **van Breugel PC, Robert EI, Mueller H**, Decorsiere A, Zoulim F, Hantz O, et al. Hepatitis B virus X protein stimulates gene expression selectively from extrachromosomal DNA templates. *Hepatology* 2012;56:2116-2124.
- [51] Ducroux A, Benhenda S, Riviere L, Semmes OJ, Benkirane M, Neuveut C. The Tudor domain protein Spindlin1 is involved in intrinsic antiviral defense against incoming hepatitis B Virus and herpes simplex virus type 1. *PLoS Pathog* 2014;10:e1004343.
- [52] Guo YH, Li YN, Zhao JR, Zhang J, Yan Z. HBc binds to the CpG islands of HBV cccDNA and promotes an epigenetic permissive state. *Epigenetics* 2011;6:720-726.
- [53] Hosel M, Quasdorff M, Wiegmann K, Webb D, Zedler U, Broxtermann M, et al. Not interferon, but interleukin-6 controls early gene expression in hepatitis B virus infection. *Hepatology* 2009;50:1773-1782.
- [54] Protzer U, Nassal M, Chiang PW, Kirschfink M, Schaller H. Interferon gene transfer by a hepatitis B virus vector efficiently suppresses wild-type virus infection. *Proc Natl Acad Sci U S A* 1999;96:10818-10823.
- [55] Belloni L, Palumbo G.A., Valente S., Rotili D., Pediconi N., , Mai A., Levrero M. Mimicking Interferon- α (IFN α) inhibitory activity on hepatitis B virus (HBV) transcription and replication by targeting the epigenetic control of nuclear cccDNA minichromosome with epigenetic small molecules. *Hepatology* 2012;56:369A.
- [56] Liu F, Campagna M, Qi Y, Zhao X, Guo F, Xu C, et al. Alpha-interferon suppresses hepadnavirus transcription by altering epigenetic modification of cccDNA minichromosomes. *PLoS Pathog* 2013;9:e1003613.
- [57] **Li T, Robert EI**, van Breugel PC, Strubin M, Zheng N. A promiscuous alpha-helical motif anchors viral hijackers and substrate receptors to the CUL4-DDB1 ubiquitin ligase machinery. *Nature structural & molecular biology* 2010;17:105-111.

- [58] **Lucifora J, Xia Y**, Reisinger F, Zhang K, Stadler D, Cheng X, et al. Specific and nonhepatotoxic degradation of nuclear hepatitis B virus cccDNA. *Science* 2014;343:1221-1228.
- [59] **Xia Y, Stadler D**, Lucifora J, Reisinger F, Webb D, Hosel M, et al. Interferon-gamma and Tumor Necrosis Factor-alpha Produced by T Cells Reduce the HBV Persistence Form, cccDNA, Without Cytolysis. *Gastroenterology* 2015.
- [60] Bohne F, Chmielewski M, Ebert G, Wiegmann K, Kurschner T, Schulze A, et al. T cells redirected against hepatitis B virus surface proteins eliminate infected hepatocytes. *Gastroenterology* 2008;134:239-247.
- [61] **Krebs K, Bottinger N**, Huang LR, Chmielewski M, Arzberger S, Gasteiger G, et al. T Cells Expressing a Chimeric Antigen Receptor That Binds Hepatitis B Virus Envelop Proteins Control Virus Replication in Mice. *Gastroenterology* 2013.
- [62] Belloni L, Li, L., Palumbo, G.A., Chirapu, S.R., Calvo, L., Finn, Mg., Lopatin, U., Zlotnick, A., Levvero, M. HAPs hepatitis B virus (HBV) capsid inhibitors block core protein interaction with the viral minichromosome and host cell genes and affect cccDNA transcription and stability. *Hepatology* 2013;54:277A.
- [63] Cradick TJ, Keck K, Bradshaw S, Jamieson AC, McCaffrey AP. Zinc-finger nucleases as a novel therapeutic strategy for targeting hepatitis B virus DNAs. *Molecular therapy : the journal of the American Society of Gene Therapy* 2010;18:947-954.
- [64] Schiffer JT, Swan DA, Stone D, Jerome KR. Predictors of hepatitis B cure using gene therapy to deliver DNA cleavage enzymes: a mathematical modeling approach. *PLoS computational biology* 2013;9:e1003131.
- [65] Chen J, Zhang W, Lin J, Wang F, Wu M, Chen C, et al. An efficient antiviral strategy for targeting hepatitis B virus genome using transcription activator-like effector nucleases. *Molecular therapy : the journal of the American Society of Gene Therapy* 2014;22:303-311.
- [66] Bloom K, Ely A, Mussolino C, Cathomen T, Arbuthnot P. Inactivation of hepatitis B virus replication in cultured cells and in vivo with engineered transcription activator-like effector nucleases. *Molecular therapy : the journal of the American Society of Gene Therapy* 2013;21:1889-1897.
- [67] Seeger C, Sohn JA. Targeting Hepatitis B Virus With CRISPR/Cas9. *Mol Ther Nucleic Acids* 2014;3:e216.
- [68] **Kennedy EM, Bassit LC**, Mueller H, Kornepati AV, Bogerd HP, Nie T, et al. Suppression of hepatitis B virus DNA accumulation in chronically infected cells using a bacterial CRISPR/Cas RNA-guided DNA endonuclease. *Virology* 2015;476:196-205.
- [69] **Dong C, Qu L**, Wang H, Wei L, Dong Y, Xiong S. Targeting hepatitis B virus cccDNA by CRISPR/Cas9 nuclease efficiently inhibits viral replication. *Antiviral Res* 2015;118:110-117.

Figure and table legends

Table 1: cccDNA levels and estimated time of maintenance of HBV infection in different *in vitro* and *in vivo* models.

Figure 1: cccDNA qPCR primers spanning the “nick region” of the HBV relaxed circular DNA (rcDNA). Examples of qPCR primers designed to selectively amplify cccDNA over rcDNA.

Primers in purple are used with DNA intercalating reagents; primers in blue and green are used in combination with probe detection.

Figure 2: Schematic representation of rcDNA conversion to cccDNA. (i) Removal of viral polymerase linked to the 5' end of the (–) DNA strand and of one of the redundant sequences, (ii) removal of the RNA primer linked to the 5' end of the (+) strand, (iii) fill in and completion of the viral (+) DNA strand, and (iv) ligations of DNA extremities for both (+) and (–) strands. (v) Once formed, cccDNA is supercoiling and associating with proteins, such as histones, to form the so-called cccDNA mini-chromosome.

Figure 3: Targeting cccDNA to achieve HBV cure.

*EMA = epigenetic modifying agent; **HBP = HBx binding partner activation

Table

	% of infected cells	cccDNA levels (average copies/ nucleus)	Maintenance of infection
Primary human hepatocytes	20 - 100 %	1 to 2	2-3 weeks ⁽⁵⁸⁾
dHepaRG	5 - 20 %	0.2 to 0.5	> 6 months ⁽⁴⁰⁾
HepG2-NTCP	50 – 100 %	1 to 5	10-15 days ⁽⁴⁾
HuHep mice	100 %	1 to 2	> 4 months ⁽¹³⁾

Figure 1

Figure 2

Figure 3