



Infinite families of inequivalent real circle actions on affine four-space

Lucy Moser-Jauslin

► To cite this version:

Lucy Moser-Jauslin. Infinite families of inequivalent real circle actions on affine four-space. *Épigraphique*, 2019, Volume 3, 11 p. 10.46298/epiga.2019.volume3.4685 . hal-01826458v2

HAL Id: hal-01826458

<https://hal.science/hal-01826458v2>

Submitted on 28 Feb 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License

Infinite families of inequivalent real circle actions on affine four-space

L. Moser-Jauslin

Abstract. The main result of this article is to construct infinite families of non-equivalent equivariant real forms of linear $\mathbb{C}^*$ -actions on affine four-space. We consider the real form of $\mathbb{C}^*$ whose fixed point is a circle. In [F-M] one example of a non-linearizable circle action was constructed. Here, this result is generalized by developing a new approach which allows us to compare different real forms. The constructions of these forms are based on the structure of equivariant $O_2(\mathbb{C})$ -vector bundles.

Keywords. Real forms; circle actions; non-linearizable actions; real affine varieties

2010 Mathematics Subject Classification. 14L30; 14R20; 14P99

[Français]

Titre. Sur des familles infinies d'actions non équivalentes du cercle réel sur l'espace affine de dimension 4

Résumé. Le résultat principal de cet article est de construire des familles infinies de formes réelles équivariantes, non équivalentes entre elles, d'actions linéaires de $\mathbb{C}^*$ sur l'espace affine de dimension 4. L'article [F-M] construisait un exemple d'action du cercle non linéarisable. Ici nous généralisons ce résultat en développant une nouvelle approche qui nous permet de comparer les différentes formes réelles. Les constructions de ces formes réelles s'appuient sur la structure de $O_2(\mathbb{C})$ -fibrés vectoriels équivariants.

Received by the Editors on July 16, 2018, and in final form on October 23, 2018.

Accepted on December 6, 2018.

L. Moser-Jauslin

Université de Bourgogne Franche-Comté, Institut de Mathématiques de Bourgogne – UMR 5584 du CNRS, 9 avenue Alain Savary, BP 47870, Dijon 21078, France

e-mail: moser@u-bourgogne.fr

This work was supported by the French “Investissements d’Avenir” program, project ISITE-BFC (contract ANR-IS-IDEX-OOOB)

Contents

1. Introduction	2
2. Description of real forms	2
3. Main result : the $(2, 2m + 1)$ case	3
4. Real forms of equivariant vector bundles	7
5. $O_2(\mathbb{C})$ -vector bundles	8
6. The $(1, 2)$ case	9
7. Questions	10

1. Introduction

In this article, we construct infinite families of inequivalent real forms of linear $\mathbb{C}^*$ -actions on complex affine four-space. Consider the real form σ of $\mathbb{C}^*$ whose fixed point set is a circle. That is, $\sigma(t) = \bar{t}^{-1}$ for all $t \in \mathbb{C}^*$. Now consider the linear action of $\mathbb{C}^*$ on the four dimensional affine space Y with weights $2, -2, n, -n$ for n an odd integer ≥ 3 . We study equivariant real forms of this $\mathbb{C}^*$ -variety, that is, real forms μ of Y which are compatible with the real form σ of $\mathbb{C}^*$. The construction of these real forms uses a classification of equivariant $O_2(\mathbb{C})$ -vector bundles where the base (resp. zero fiber) is the irreducible $O_2(\mathbb{C})$ -module for which $\mathbb{C}^*$ acts with weights 2 and -2 (resp. n and $-n$). For $n \geq 5$, it is known that one can find a family for which the actions of $O_2(\mathbb{C})$ on the total spaces are all inequivalent. By adapting these ideas to real forms, we will show in particular that, for $n \geq 5$, there are infinitely many distinct real forms of Y .

It was shown in [K-R] that, over any field of characteristic zero all actions of any form of a $\mathbb{G}_m$ -action on affine three-space is linearizable. The result from [F-MJ] shows that this result does not hold in dimension 4 for the field $\mathbb{R}$. Here, we describe a systematic approach to find many non-linearizable actions which are all pairwise inequivalent.

In Section 2, we recall basic definitions of real forms of equivariant varieties. Then in Section 3, we state and give the proof of the main results. In Sections 4 and 5, we describe the method of constructing the examples, using equivariant vector bundles for the group $O_2(\mathbb{C})$. We then show in Section 6 how the analogous result for another case fails to hold. In the final section, we state two open questions related to these results.

2. Description of real forms

If G is a complex linear algebraic group, a real form of G is given by an antiholomorphic involution σ on G where σ is a group automorphism.

Given a G -variety Y , a (G, σ) -real form of Y , or an *equivariant real form of Y compatible with σ* , is given by an antiholomorphic involution μ of Y such that $\mu(gy) = \sigma(g)\mu(y)$ for all $g \in G$ and $y \in Y$. Two real forms μ_1 and μ_2 are equivalent if there exists a regular G -automorphism ψ of Y such that $\mu_2 = \psi \circ \mu_1 \circ \psi^{-1}$.

Suppose that at least one real (G, σ) -form on Y , which we denote by μ_0 , exists. Then we will show that the set of all equivalence classes of real forms is determined by a cohomology set. Let $\Gamma = \{1, \gamma\}$ be the Galois group of $\mathbb{C}/\mathbb{R}$, and let $A = \text{Aut}_G(Y)$. Consider the action of Γ on A defined by

$$\gamma\psi = \mu_0 \circ \psi \circ \mu_0 = \mu_0 \psi \mu_0.$$

If μ is any real (G, σ) -form on Y , then $\varphi := \mu \circ \mu_0 = \mu\mu_0 \in A$, and $\gamma\varphi = \mu_0\mu\mu_0^2 = \mu_0\mu = (\mu\mu_0)^{-1} = \varphi^{-1}$ since μ and μ_0 are of order 2. Conversely, if $\varphi \in A$ satisfies the condition that $\gamma\varphi = \varphi^{-1}$, then $\varphi\mu_0$ is a

real form of Y , since $(\varphi\mu_0)^2 = \varphi^\gamma\varphi = \varphi\varphi^{-1} = id$. In other words, the map which sends μ to $\mu\mu_0$ defines a bijection between the set of (G, σ) -real forms of Y , and the set $Z^1(\Gamma, A) := \{\varphi \in A \mid \varphi^\gamma\varphi = \varphi^{-1}\}$.

Now suppose that $\varphi_1\mu_0$ and $\varphi_2\mu_0$ are two real forms (with $\varphi_1, \varphi_2 \in Z^1(\Gamma, A)$). They are equivalent if and only if there exists an automorphism $\psi \in A$ such that $\psi\varphi_1\mu_0\psi^{-1} = \varphi_2\mu_0$, that is, $\psi\varphi_1^\gamma\psi^{-1} = \varphi_2$. Thus we define the equivalence relation on $Z^1(\Gamma, A)$ by $\varphi_1 \sim \varphi_2$ if there exists $\psi \in A$ such that $\varphi_2 = \psi\varphi_1^\gamma(\psi)^{-1}$. The bijection above induces a bijection between the equivalence classes of (G, σ) -real forms of Y and $H^1(\Gamma, A) := Z^1(\Gamma, A)/\sim$.

3. Main result : the $(2, 2m+1)$ case

3.A. Construction of circle forms

The case considered here is $G = \mathbb{C}^*$, and $\sigma(t) = \bar{t}^{-1}$. The fixed points of σ form the circle S^1 of elements of norm 1 in $\mathbb{C}^*$.

If Y is a $\mathbb{C}^*$ -variety, a real form μ on Y as a $\mathbb{C}^*$ -variety which is compatible with σ is called a *real circle form*, and the action of (G, σ) on (Y, μ) is called a *circle action on Y* or a *real circle action on Y* .

Let W_k be the two-dimensional $\mathbb{C}^*$ -module with weights $(k, -k)$. In other words, $t(x, y) = (t^k x, t^{-k} y)$ for $t \in \mathbb{C}^*$.

Fix $m \geq 1$, and $n = 2m+1$. In this section, consider the following $\mathbb{C}^*$ -variety $Y = W_2 \times W_n$. The following defines a real circle form on Y :

$$\mu_0\left(\begin{pmatrix} a \\ b \end{pmatrix}, \begin{pmatrix} x \\ y \end{pmatrix}\right) = \left(\begin{pmatrix} \bar{b} \\ \bar{a} \end{pmatrix}, \begin{pmatrix} \bar{y} \\ \bar{x} \end{pmatrix}\right).$$

We call this form “a linear circle action on affine space”. Now we will construct a family of other real circle forms on Y . There might be many others, but we restrict the study to only some cases (where equivariant vector bundle methods can be used for constructing and distinguishing different forms).

Later, we will explain how these real forms come from non-equivalent $O(2)$ -actions found by G. Schwarz in [SI].

Let $T = ab$. Given $h \in \mathbb{R}[T]$, let $M_h \in GL_2(\mathbb{C}[a, b])$ be the matrix given by

$$M_h = \begin{pmatrix} 1 - Th^2 & a^n h^n \\ -b^n h^n & \sum_{j=0}^{n-1} (Th^2)^j \end{pmatrix},$$

Let φ_h be the automorphism of Y :

$$\varphi_h\left(\begin{pmatrix} a \\ b \end{pmatrix}, \begin{pmatrix} x \\ y \end{pmatrix}\right) = \left(\begin{pmatrix} a \\ b \end{pmatrix}, M_h \begin{pmatrix} x \\ y \end{pmatrix}\right).$$

Note that φ_h is $\mathbb{C}^*$ -equivariant: $\varphi_h \in \text{Aut}_{\mathbb{C}^*}(Y)$.

We now state the main theorem of this article.

Theorem 3.1. *Let $h \in \mathbb{R}[T]$ be a real polynomial, and let $\mu_h = \varphi_h\mu_0$.*

- (i) μ_h defines a real circle form on $Y = W_2 \times W_{2m+1}$;
- (ii) μ_h is equivalent to $\mu_{h'}$ if and only if there exists $r \in \mathbb{R}^*$ such that

$$h(T) \equiv rh'(r^2 T) \pmod{(T^m)}.$$

Note that, if one forgets the action, and one considers μ_h as a real form of Y as a variety, then they are all equivalent.

3.B. Two special cases

- (1) As a consequence, for $m = 1$, we find two inequivalent real forms. For $h = 0$, we find the linear circle action, and for $h = 1$, we have another, non-equivalent form. This case is precisely the action described in [F-MJ].
- (2) For $m = 2$, if $h = c_0 + c_1 T$ and $h' = c'_0 + c'_1 T$, with all coefficients being real, the theorem implies that the real forms $\varphi_h \circ \mu_0$ and $\varphi_{h'} \circ \mu_0$ are equivalent if and only if :
 - (i) $c_0 = c'_0 = c_1 = c'_1 = 0$;
 - (ii) $c_0 = c'_0 = 0, c_1 c'_1 \neq 0$;
 - (iii) $c_1 = c'_1 = 0, c_0 c'_0 \neq 0$; or
 - (iv) $c_0 c'_0 c_1 c'_1 \neq 0$ and $(c'_0/c_0)^3 = c'_1/c_1$.

In particular one finds an infinite family of inequivalent real circle forms of the same $\mathbb{C}^*$ -linear action.

3.C. Three linear subgroups of $A = \text{Aut}_{\mathbb{C}^*} Y$

In order to prove this theorem, we will use certain subgroups of $A = \text{Aut}_{\mathbb{C}^*} Y$. They are related to vector-bundle automorphisms described in general in the following sections.

- (1) Consider the subgroup $\Lambda \subset GL_2(\mathbb{C}[a, b])$ defined as follows. Set $T = ab$.

$$\Lambda = \left\{ M = \begin{pmatrix} P(T) & a^n Q(T) \\ b^n S(T) & R(T) \end{pmatrix} \mid P, Q, R, S \in \mathbb{C}[T], \det(M) = \Delta \in \mathbb{C}^* \right\}.$$

We define an action of Γ on the group Λ . If

$$M = \begin{pmatrix} P(T) & a^n Q(T) \\ b^n S(T) & R(T) \end{pmatrix}, \quad \text{then} \quad {}^\gamma M = \begin{pmatrix} \overline{R}(T) & a^n \overline{S}(T) \\ b^n \overline{Q}(T) & \overline{P}(T) \end{pmatrix}.$$

Now consider the subgroup $A_\Lambda \subset A$ of automorphisms of the form

$$\varphi_M \left(\begin{pmatrix} a \\ b \end{pmatrix}, \begin{pmatrix} x \\ y \end{pmatrix} \right) = \left(\begin{pmatrix} a \\ b \end{pmatrix}, M \begin{pmatrix} x \\ y \end{pmatrix} \right)$$

where $M \in \Lambda$.

By construction, ${}^\gamma \varphi_M = \varphi_{{}^\gamma M}$, and $\varphi_M^{-1} = \varphi_{M^{-1}}$.

- (2) We consider also a subgroup of automorphisms of Y coming from the linear circle action. More precisely, if $\omega \in \mathbb{C}$ is of norm 1, we construct the automorphism ρ_ω by

$$\rho_\omega \left(\begin{pmatrix} a \\ b \end{pmatrix}, \begin{pmatrix} x \\ y \end{pmatrix} \right) = \left(\begin{pmatrix} \omega^2 a \\ \omega^{-2} b \end{pmatrix}, \begin{pmatrix} \omega^n x \\ \omega^{-n} y \end{pmatrix} \right).$$

- (3) Finally, if $r \in \mathbb{R}$, we consider a subgroup of automorphisms that act only on the first factor:

$$\ell_r \left(\begin{pmatrix} a \\ b \end{pmatrix}, \begin{pmatrix} x \\ y \end{pmatrix} \right) = \left(\begin{pmatrix} ra \\ rb \end{pmatrix}, \begin{pmatrix} x \\ y \end{pmatrix} \right).$$

Proof the theorem, part (i). It suffices to note that $\varphi_h \in A_\Lambda$, and that $M_h^{-1} = {}^\gamma M_h$. This implies that the cocycle condition ${}^\gamma \varphi_h^{-1} = \varphi_h$ is verified, and therefore $\mu_h = \varphi_h \mu_0$ is an anti-holomorphic involution on Y , and it is compatible with σ , and therefore is a real circle form. $\square$

3.D. Equivalences

The first step in proving the equivalence relation of part (ii) of the theorem is to reduce the problem of equivalence to a computation using automorphisms in A_Λ .

Proposition 3.2. *The real forms μ_h and $\mu_{h'}$ are equivalent if and only if there exists $\psi \in A_\Lambda$ and $r \in \mathbb{R}^*$ such that $\psi \circ \varphi_h \circ \gamma \psi^{-1} = \varphi_{h'}$ where $h''(T) = rh'(r^2 T)$.*

This result is a special case of the general Proposition 4.2, stated later. The proof for this case is given completely here.

Proof. First note that $\ell_r \varphi_{h'} \gamma \ell_r^{-1} = \ell_r \varphi_{h'} \ell_r^{-1} = \varphi_{h''}$. Thus one direction of the equivalence is clear.

Now suppose that μ_h is equivalent to $\mu_{h'}$. This means that there is an automorphism $\psi \in A$ such that $\psi \circ \varphi_h \circ \gamma \psi^{-1} = \varphi_{h'}$.

We will now follow an argument from Masuda and Petrie [MP2] to create a vector bundle automorphism which has the same property. Let $H = \{\pm 1\}$ be the subgroup of $\mathbb{C}^*$ of two elements. Since ψ is equivariant, it stabilizes the fixed point set of H . In other words, ψ stabilizes the subvariety B defined as the zero set of x and y . Also, by equivariance, and from the hypothesis that $\psi \circ \varphi_h \circ \gamma \psi^{-1} = \varphi_{h'}$, we see that there exists $\lambda \in \mathbb{C}^*$ such that

$$\psi\left(\begin{pmatrix} a \\ b \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \end{pmatrix}\right) = \left(\begin{pmatrix} \lambda a \\ \bar{\lambda} b \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \end{pmatrix}\right).$$

By choosing an appropriate $r \in \mathbb{R}$ and $\omega \in \mathbb{C}$ of norm 1, one can construct $\psi_1 = \ell_r^{-1} \rho_\omega \psi$ such that ψ_1 fixes pointwise B . Note that $\psi_1 \circ \varphi_h \circ \gamma \psi_1^{-1} = \varphi_{h''}$.

Finally Y is the total space of the normal bundle of B in Y . Thus ψ_1 induces an automorphism ψ_2 on the normal bundle, which gives therefore an automorphism in A_Λ .

By the fact that φ_h and $\varphi_{h'}$ are also in A_Λ , we have that $\psi_2 \circ \varphi_h \gamma \psi_2^{-1} = \varphi_{h''}$, and the proposition is proven. $\square$

Now we will show that the real circle actions μ_h which were constructed are all equivalent over a certain open set of Y . Then part (ii) of the theorem can be proven using this result and Proposition 3.2.

Notation: $\mathbb{V}(ab) \subset Y$ is the closed subvariety Y defined by the equation $ab = 0$.

Note first that $U = Y \setminus \mathbb{V}(ab)$ is an open G -subvariety, and all the real forms μ_h restrict to real circle forms on U . We will start by showing that they are all equivalent on U .

Consider first the analogous setting that was used on Y . Let $A' = \text{Aut}_{\mathbb{C}^*} U$. Let $\Lambda' \subset GL_2(\mathbb{C}[a, b, T^{-1}])$ be defined as follows.

$$\Lambda' = \left\{ M = \begin{pmatrix} P(T) & a^n Q(T) \\ b^n S(T) & R(T) \end{pmatrix} \mid P, Q, R, S \in \mathbb{C}[T, T^{-1}], \det(M) = \Delta = cT^k, c \in \mathbb{C}^*, k \in \mathbb{Z} \right\}.$$

As before, define an action of Γ on the group Λ' . If

$$M = \begin{pmatrix} P(T) & a^n Q(T) \\ b^n S(T) & R(T) \end{pmatrix}, \quad \text{then} \quad \gamma M = \begin{pmatrix} \bar{R}(T) & a^n \bar{S}(T) \\ b^n \bar{Q}(T) & \bar{P}(T) \end{pmatrix}.$$

Consider the subgroup A'_Λ of A' of automorphisms of the form

$$\varphi_M\left(\begin{pmatrix} a \\ b \end{pmatrix}, \begin{pmatrix} x \\ y \end{pmatrix}\right) = \left(\begin{pmatrix} a \\ b \end{pmatrix}, M \begin{pmatrix} x \\ y \end{pmatrix}\right)$$

where $M \in \Lambda'$.

By construction, $\gamma \varphi_M = \varphi_{\gamma M}$, and $\varphi_M^{-1} = \varphi_{M^{-1}}$.

For $h \in \mathbb{R}[T]$, construct the matrix $K_h \in \Lambda'$ by

$$K_h = \begin{pmatrix} 1 & \frac{a^n h}{T^m} \\ \frac{b^n h \sum_{j=0}^{m-1} (Th^2)^j}{T^m} & \sum_{j=0}^m (Th^2)^j \end{pmatrix}$$

and let $\psi'_h \in A'_{\Lambda'}$ be the corresponding automorphism of U :

$$\psi'_h \left(\begin{pmatrix} a \\ b \end{pmatrix}, \begin{pmatrix} x \\ y \end{pmatrix} \right) = \left(\begin{pmatrix} a \\ b \end{pmatrix}, K_h \begin{pmatrix} x \\ y \end{pmatrix} \right).$$

Lemma 3.3. *Let $U = Y \setminus \mathbb{W}(T)$. The real forms μ_h restricted to U are all equivalent. More precisely, we have $\mu_h = \psi'_h \mu_0 \psi'^{-1}_h$.*

Proof. It suffices to check that $\det(K_h) = 1$ and that $K_h {}^\gamma K_h^{-1} = M_h$. $\square$

Lemma 3.4. *Let $\Psi \in \Lambda'$ be a matrix such that ${}^\gamma \Psi = \Psi$ and such that $\det(\Psi)$ is a non-zero complex constant. Then Ψ is of the form*

$$\Psi = \begin{pmatrix} \alpha & 0 \\ 0 & \bar{\alpha} \end{pmatrix}$$

for some $\alpha \in \mathbb{C}^*$.

Proof. Set $\Psi = \begin{pmatrix} P(T) & a^n Q(T) \\ b^n S(T) & R(T) \end{pmatrix}$ where $P, Q, R, S \in \mathbb{C}[T, T^{-1}]$, with determinant $\Delta \in \mathbb{C}^*$. The first coefficient of $\Psi {}^\gamma \Psi^{-1}$ is $(P\bar{P} - T^n Q\bar{Q})/\bar{\Delta}$. Therefore, we have the condition that $P\bar{P} - T^n Q\bar{Q} = \bar{\Delta}$. Consider $P\bar{P}$ and $Q\bar{Q}$ as Laurent polynomials. The highest monomial in $P\bar{P}$ is of even degree, and the highest monomial in $T^n Q\bar{Q}$ is of odd degree. Thus they do not cancel. The same holds for the lowest degree monomials. Thus the only possibility to have the given equality is that $Q = 0$ and P is of degree 0. Set $P = \alpha$. We find $\bar{\Delta} = \alpha \bar{\alpha}$, thus Δ is real. By considering the other coefficients of Ψ , we find similarly that $S = 0$, and that Ψ is of the given form. $\square$

Proof of Theorem 3.1. By Proposition 3.2, μ_h is equivalent to $\mu_{h'}$ if and only if there exists $r \in \mathbb{R}^*$ and $N \in \Lambda$ such that $N M_h {}^\gamma N^{-1} = M_{h'}$, where $h''(T) = r h'(r^2 T)$. This means

$$N K_h {}^\gamma K_h^{-1} {}^\gamma N^{-1} = K_{h''} {}^\gamma K_{h''}^{-1}.$$

Set $\Phi = K_{h''}^{-1} N K_h \in \Lambda'$. N satisfies the given equation if and only if ${}^\gamma \Phi = \Phi$. Also, since $\det(K_h) = \det(K_{h''}) = 1$, and since $\det N$ is a constant, the same holds of Φ . By Lemma 3.4, there exists $\alpha \in \mathbb{C}^*$ such that Φ is diagonal with terms α and $\bar{\alpha}$.

The coefficients of N are all polynomials in T . Note that $K_h = \begin{pmatrix} 1 & a^n q_h \\ b^n s_h & r_h \end{pmatrix}$ where $q_h = h/T^m$, $s_h = h(\sum_{j=0}^{m-1} Th^2)/T^m$, and $r_h = \sum_{j=0}^m (Th^2)^j$. The coefficients for N are all polynomials in $\mathbb{C}[a, b]$ if and only if:

$$\alpha s_h r_{h''} - \bar{\alpha} r_h s_{h''} \in \mathbb{C}[T]$$

and

$$\alpha q_{h''} - \bar{\alpha} q_h \in \mathbb{C}[T].$$

First of all, if $h \equiv h'' \pmod{T^m}$, then one can choose $\alpha = 1$, and the two conditions are verified. That is, one can construct the matrix $N \in \Lambda$ which induces the equivalence of φ_h and $\varphi_{h''}$.

For the converse, note that the second equation implies that $\alpha h \equiv \bar{\alpha} h'' \pmod{T^m}$.

Since h and h'' are real, if the real part of α is not zero, this implies that $h \equiv h'' \pmod{T^m}$. If α is purely imaginary, then $h \equiv -h'' \pmod{T^m}$. We also know that $\varphi_{h''}$ is equivalent to $\varphi_{-h''}$, by posing $r = -1$, which gives $r h''(r^2 T) = -h''(T)$. Thus the existence of an appropriate matrix N implies the condition in the theorem. $\square$

4. Real forms of equivariant vector bundles

The examples given here were constructed using techniques of equivariant $O_2(\mathbb{C})$ -vector bundles over $O_2(\mathbb{C})$ -modules. In [K-S], Kraft and Schwarz studied the following situation. Let G be a reductive complex group, and let B and F be two G -modules. The total space of an equivariant G -vector bundle over B with zero fiber F , as a variety, is isomorphic to affine space. If an equivariant bundle is not trivial, it can happen that the associated action on the total space is non-linearizable. In fact, all examples of algebraic non-linearizable actions on affine complex space known at this time are constructed in this way. Kraft and Schwarz studied the case where the algebraic quotient of B by G is the affine line. They showed that the set of equivalence classes of G -vector bundles with base B and zero fiber F forms a moduli space. Another approach to the study of these G -vector bundles was developed in [MP1], [MP2], [MMP] and [MJ], using the construction of equivariant bundles as subbundles of trivial ones. The method we used in Proposition 3.2 due to Masuda and Petrie was originally formulated to show how non-equivalent G -vector bundles can lead to non-equivalent actions on the total spaces.

In this section, we will compare the study of real forms of equivariant vector bundles and equivariant real forms on the total space. In the next section, we will describe how to use $O_2(\mathbb{C})$ -bundles to construct real forms of linear $\mathbb{C}^*$ -actions on four-space.

Let $\pi : E \rightarrow B$ be a G -vector bundle. This means that it is a vector bundle over B , the group G acts on E and B , π is equivariant, and G acts linearly on the fibers. That is, for any $b \in B$ and $g \in G$, the restriction of the action of g from $\pi^{-1}(b)$ to $\pi^{-1}(gb)$ is linear. If f is a G -automorphism of B , a G -automorphism of E over f is a G -automorphism φ of E such that $\varphi \circ \pi = \pi \circ f$, and such that φ is linear on the fibers, that is, for any $b \in B$, the restriction of φ from $\pi^{-1}(b)$ to $\pi^{-1}(f(b))$ is linear. A G -vector bundle automorphism of $E \rightarrow B$ is a G -automorphism φ of E over the identity map on B .

Suppose B is a complex G -variety, σ is a real form of G and μ_B is a fixed real form of B compatible with σ .

Definition 4.1. A real form μ of E is called a G -vector bundle real form compatible with σ and μ_B , if μ is an anti-holomorphic involution on E defining a real form compatible with σ and such that $\mu \circ \pi = \pi \circ \mu_B$, and μ is $\mathbb{R}$ -linear on the fibers. Two G -vector bundle real forms μ_1 and μ_2 are equivalent if there exists a G -vector bundle automorphism ψ which conjugates μ_1 to μ_2 .

Clearly, G -vector bundle real forms compatible with σ give real forms on the total space. Also, if two such real forms are equivalent as G -vector-bundle real forms, then they are equivalent as equivariant real forms on the total space. In general, the converse is not true. However, under some conditions, we can induce information of non-equivalence on the total space from non-equivalence as vector bundle real forms. The following proposition is essentially the result from Masuda-Petrie [MP2] which was used in Proposition 3.2.

Suppose $\pi : E \rightarrow B$ is a trivial G -vector bundle and f is a G -automorphism of B . Then f induces a G -automorphism on $E = B \times F$, which we also call f . Suppose now that σ is a real form of G , and μ_B is a real form of B compatible with σ , and μ is a G -vector bundle real form of E , compatible with σ and μ_B . If f is a G -equivariant automorphism of B which preserves μ_B , (that is, $f\mu_B f^{-1} = \mu_B$), then $f^*\mu := f\mu f^{-1}$ is another vector bundle real form of E compatible with σ and μ_B . By definition $f^*\mu$ and μ are equivalent as equivariant real forms of E .

Proposition 4.2. Let G be an algebraic complex group with real form σ , and let B be a G -variety with real form μ_B compatible with σ . Suppose $\pi : E \rightarrow B$ is a trivial G vector bundle.

Suppose that G has a subgroup H such that $E^H = B$, the zero section of the vector bundle.

Then, given two equivariant vector bundle real forms μ_1 and μ_2 on $E = B \times F$, they are equivalent as equivariant real forms of E if and only if there exists a G -equivariant automorphism f of B which preserves μ_B such that μ_1 and $f^*\mu_2$ are equivalent as G -vector bundle real forms.

Proof. Suppose ψ is a G -equivariant automorphism of E such that $\psi \circ \mu_1 \circ \gamma \psi^{-1} = \mu_2$. Since ψ is equivariant for H , $\psi(B) = B$, where B is identified with the zero section of the trivial vector bundle. $\psi_B = f$ is therefore a G -automorphism compatible with σ and μ_B . Extending f to $f = f \times id$ on $E = B \times F$, $(f^{-1}) \circ \psi$ is a G -equivariant automorphism of E which fixes pointwise the zero section B . This automorphism therefore induces a vector bundle automorphism ψ' of the normal bundle of B in the total space, which is G -isomorphic to E . Since μ_1 and μ_2 are linear on the fibers, $\psi' \mu_1 \gamma \psi'^{-1} = f^* \mu_2$. $\square$

5. $O_2(\mathbb{C})$ -vector bundles

In [S1] Schwarz gave a description of certain moduli spaces of G -vector bundle equivalence classes for $G = O_2(\mathbb{C}) = \mathbb{C}^* \rtimes \mathbb{Z}/2\mathbb{Z}$. The irreducible representations of G are given by the one-dimensional ones and, for every $k \in \mathbb{N} \setminus \{0\}$, the two dimensional representation W_k where $\mathbb{C}^*$ acts by weights k and $-k$, and G admits an involution which exchanges the two eigenspaces. Schwarz showed in particular that the set of equivalence classes $VEC(W_2; W_{2m+1})$ of G -vector bundles with base W_2 and zero fiber W_{2m+1} is a moduli space isomorphic to $\mathbb{C}^m$. All of these vector bundles come from considering the linear action of $\mathbb{C}^*$ on the total space, and then finding a compatible (regular) involution. One can then find non-linearizable G -actions on four-space.

In [F-MJ] it was shown that, for the case $m = 1$, one can use a certain involution from a non-linearizable action formed in this way to construct an equivariant real circle form for the $\mathbb{C}^*$ -action on the total space (see also [W]). In order to do this, one needed to use that the involution commutes with conjugation. More precisely, the regular involution coming from a non-linearizable action of G on affine four-space composed with conjugation defines an anti-holomorphic involution, since they commute. It was shown that the action given there was not equivalent to the linear circle action.

In the present article, the cases treated come from families of non-equivalent vector bundles. To show that two real forms are not equivalent in general is more difficult than simply showing they are not equivalent to the linear one. The key to this calculation was to use that they are all equivalent on the open set U constructed in Section 3.D.

We describe now how to find examples of real circle actions on complex four-space with a linear action using G -vector bundles. Denote by $s \in G$ an order two element of G which is not in the normal subgroup $\mathbb{C}^* : G = \mathbb{C}^* \rtimes \{1, s\}$.

A G -vector bundle $W_k \times W_n$ is constructed by defining the linear $\mathbb{C}^*$ -action with weights $(k, -k, n, -n)$, and an action of the element s respecting the conditions that $s^2 = 1$ and $st = t^{-1}s$. This restriction is very similar to that of finding real circle forms.

For the trivial G -vector bundle $W_k \times W_n$ the action of s is given by the involution τ_0 which exchanges the coordinates in the base and the zero fiber:

$$t \left(\begin{pmatrix} a \\ b \end{pmatrix}, \begin{pmatrix} x \\ y \end{pmatrix} \right) = \left(\begin{pmatrix} t^k a \\ t^{-k} b \end{pmatrix}, \begin{pmatrix} t^n x \\ t^{-n} y \end{pmatrix} \right)$$

for $t \in \mathbb{C}^*$, and

$$\tau_0 \left(\begin{pmatrix} a \\ b \end{pmatrix}, \begin{pmatrix} x \\ y \end{pmatrix} \right) = \left(\begin{pmatrix} b \\ a \end{pmatrix}, \begin{pmatrix} y \\ x \end{pmatrix} \right).$$

Any other G -vector bundle with base W_k and zero fiber W_n is equivalent to one where the $\mathbb{C}^*$ -action remains the same, and the action of s is given by an involution τ is of the form $\tau = \varphi \circ \tau_0$, where φ is a $\mathbb{C}^*$ -vector bundle automorphism, and τ is an involution. Analogously to the study of real forms, one can define an action of $\{1, s\}$ on the group of $\mathbb{C}^*$ -vector bundle automorphisms on the trivial $\mathbb{C}^*$ -vector bundle $W_k \times W_n$ by conjugation with $\tau_0 : {}^s\varphi = \tau_0 \circ \varphi \circ \tau_0$. The condition for τ to define a G -vector bundle is then that ${}^s\varphi^{-1} = \varphi$. Also, two G -vector bundles defined by the involution $\tau_1 = \varphi_1 \circ \tau_0$ and $\tau_2 = \varphi_2 \circ \tau_0$ are equivalent if and only if there exists a $\mathbb{C}^*$ -vector bundle automorphism ψ such that $\psi \varphi_1 {}^s\psi^{-1} = \varphi_2$.

Note that the real form μ_0 which defines a circle action on the total space is simply $\tau_0 \circ \text{conj} = \text{conj} \circ \tau_0$, where conj is the standard complex conjugation on $\mathbb{C}^4$. Now suppose that one restricts to vector bundle automorphisms φ which commute with conjugation. Then $\mu = \varphi \circ \mu_0$ defines another $(\mathbb{C}^*, \sigma)$ equivariant real form. The question of equivalence, however, is not identical. If ψ can be constructed to give equivalence of $O_2(\mathbb{C})$ -vector bundles in such a way that it commutes with conjugation, then it gives an equivalence of the corresponding real circle forms. However, if ψ does not commute with conjugation, the situation can become quite different. In the next section, such an example is treated.

The explicit automorphisms used in this article were obtained by calculating the non-equivalent vector bundles using the descriptions from [K-S] and from [MPI].

6. The (1, 2) case

There are also non-trivial $O_2(\mathbb{C})$ -vector bundles whose base is W_1 . These, however, do not in general lead to non-equivalent real circle forms of $\mathbb{C}^4$ with the corresponding $\mathbb{C}^*$ actions. First of all, the argument of Masuda and Petrie does not apply. But more than that, there are explicit examples where an involution coming from a non-trivial equivariant vector bundle leads to a linearizable real circle action. (See also [F-MJ].)

We present here the following case. Consider the $O_2(\mathbb{C})$ -vector bundle whose total space in $\mathbb{C}^4$ with weights $(1, -1, 2, -2)$ for the $\mathbb{C}^*$ -action.

Consider the involution τ defined by

$$\tau\left(\begin{pmatrix} a \\ b \end{pmatrix}, \begin{pmatrix} x \\ y \end{pmatrix}\right) = \left(\begin{pmatrix} b \\ a \end{pmatrix}, \Phi\left(\begin{pmatrix} y \\ x \end{pmatrix}\right)\right)$$

where

$$\Phi = \begin{pmatrix} 1 - T & a^4 \\ -b^4 & 1 + T + T^2 + T^3 \end{pmatrix}.$$

(Here, as before, $T = ab$.) It is known that this involution together with the $\mathbb{C}^*$ linear action defines an $O_2(\mathbb{C})$ -vector bundle which is non-trivial. (It is not known whether the $O_2(\mathbb{C})$ -action on the total space is linearizable or not.)

Let τ_0 be the involution which exchanges a and b , and exchanges x and y . Then $\tau = \varphi \circ \tau_0$, where φ is a vector bundle involution: $\varphi\left(\begin{pmatrix} a \\ b \end{pmatrix}, \begin{pmatrix} x \\ y \end{pmatrix}\right) = \left(\begin{pmatrix} a \\ b \end{pmatrix}, \Phi\left(\begin{pmatrix} x \\ y \end{pmatrix}\right)\right)$.

Now consider the total space Y as a $\mathbb{C}^*$ -variety. Since φ commutes with complex conjugation, $\varphi \circ \mu_0 = \mu$ defines a circle form on Y .

Proposition 6.1. *The circle form $\mu = \varphi \mu_0$ on $Y = W_1 \times W_2$ is equivalent to the linear circle action μ_0 .*

Proof. Let $N \in GL_2(\mathbb{C}[a, b])$ be the matrix:

$$N = \begin{pmatrix} 1 - \frac{(1-i)}{2}T - \frac{(1+i)}{4}T^2 & \frac{a^4}{4}(1-i) \\ \frac{-b^4}{4}(3-i+(1+i)T) & 1 + \frac{1-i}{4}(2T + T^2 + T^3) \end{pmatrix}.$$

Then

$$\psi\left(\begin{pmatrix} a \\ b \end{pmatrix}, \begin{pmatrix} x \\ y \end{pmatrix}\right) = \left(\begin{pmatrix} a \\ b \end{pmatrix}, N\left(\begin{pmatrix} x \\ y \end{pmatrix}\right)\right)$$

is a $\mathbb{C}^*$ -automorphism.

Also $\psi \mu_0 \psi^{-1} = \mu$, since $N \nu N^{-1} = \Phi$. Therefore, μ_0 is equivalent to μ . □

Note that the coefficients of N are not real, that is, ψ does not commute with complex conjugation.

7. Questions

7.A. Quotients

An interesting question to consider is the real forms of the quotients. Given $Y = W_2 \times W_n$, the algebraic quotient $Z = Y//(\mathbb{C}^*)$ is a singular threefold. Set $T = ab$, $W = xy$, $U = a^n y^2$ and $V = b^n x^2$. These polynomials generate the invariant ring, and they satisfy the equation $UV - T^n W^2 = 0$. More precisely, $Z \cong \mathbb{V}(UV - T^n W^2) \subset \mathbb{C}^4$. Any circle form on Y induces a real form on Z . Thus it is natural to ask if inequivalent circle forms on Y induce inequivalent real forms on Z . In fact, in Theorem 1.12 of [S2]¹, Schwarz showed that any automorphism of Z lifts to an automorphism of Y which is either equivariant or quasi-equivariant for the automorphism $t \rightarrow t^{-1}$ of the group $\mathbb{C}^*$. This is an important step in understanding how to approach this question. However, it is not clear when an equivalence of real forms on Z would lift to an equivalence of equivariant real forms on Y .

Note also, that one can show that the real forms are all equivalent in the diffeomorphic category.

7.B. Other examples

In this article, all the real forms that are considered come from a very particular type. They are constructed from regular involutions of non-trivial equivariant vector bundles, and these involutions all commute with complex conjugation $\mathbb{C}^4$ (as a variety).

It would be interesting to consider other cases. In particular, it is possible that, in comparison to the case presented in Section 6, there are examples of equivalent equivariant $O_2(\mathbb{C})$ -bundles that lead to inequivalent real circle forms. The techniques applied using $O_2(\mathbb{C})$ -vector bundles would have to be adapted to study this case.

References

- [F-MJ] G. Freudenburg and L. Moser-Jauslin, *Real and rational forms of certain $O_2(\mathbb{C})$ -actions, and a solution to the weak complexification problem*, Transform. Groups **9** (2004), no. 3, 257–272. [MR-2079135](#)
- [K-R] M. Koras and P. Russell, *Separable forms of $\mathbb{G}_m$ -actions on $\mathbb{A}^3$* , Transform. Groups **18** (2013), no. 4, 1155–1163. [MR-3127990](#)
- [K-S] H. Kraft and G. Schwarz, *Reductive group actions with one-dimensional quotient*, Inst. Hautes Études Sci. Publ. Math. **76** (1992), 1–97. [MR-1215592](#)
- [MP1] M. Masuda and T. Petrie, *Algebraic families of $O(2)$ -actions on affine space $\mathbb{C}^4$* . In: Algebraic groups and their generalizations: classical methods (University Park, PA, 1991), pp. 347–354, Proc. Sympos. Pure Math., vol. 56, part 1, Amer. Math. Soc., Providence, RI, 1994. [MR-1278718](#)
- [MP2] M. Masuda and T. Petrie, *Stably trivial equivariant algebraic vector bundles*, J. Amer. Math. Soc. **8** (1995), no. 3, 687–714. [MR-1303027](#)
- [MMP] M. Masuda, L. Moser-Jauslin, and T. Petrie, *Equivariant algebraic vector bundles over cones with smooth one-dimensional quotient*, J. Math. Soc. Japan **50** (1998), no. 2, 379–414. [MR-1613156](#)
- [MJ] L. Moser-Jauslin, *Algebraic equivariant vector bundles and the linearity problem*. In: Algebraic groups and their generalizations: classical methods (University Park, PA, 1991), pp. 355–364, Proc. Sympos. Pure Math., vol. 56, part 1, Amer. Math. Soc., Providence, RI, 1994. [MR-1278719](#)
- [SI] G. Schwarz, *Exotic algebraic group actions*, C. R. Acad. Sci. Paris Sér. I Math. **309** (1989), no. 2, 89–94. [MR-1004947](#)

¹ ↑ I thank the referee for pointing out this reference.

- [S2] G. Schwarz, *Quotients, automorphisms and differential operators*, J. Lond. Math. Soc. (2) **89** (2014), no. 1, 169–193. [MR-3174739](#)
- [W] J. Winkelmann, *Nonlinearizable actions of commutative reductive groups*, Transform. Groups **14** (2009), no. 2, 483–491. [MR-2504931](#)