

# Chemical equilibrium for the reactive system propionic acid + ethanol + ethyl propionate + water at 303.15 and 313.15 K

Maria Toikka, Anna Sadaeva, Artemiy Samarov, Alexandra Golikova, Maya Trofimova, Nataliya Shcherbakova, Alexander Toikka

### ► To cite this version:

Maria Toikka, Anna Sadaeva, Artemiy Samarov, Alexandra Golikova, Maya Trofimova, et al.. Chemical equilibrium for the reactive system propionic acid + ethanol + ethyl propionate + water at 303.15 and 313.15 K. Fluid Phase Equilibria, 2017, 451, pp.91-95. 10.1016/j.fluid.2017.08.010 . hal-01838714

## HAL Id: hal-01838714 https://hal.science/hal-01838714

Submitted on 13 Jul 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



### **Open Archive Toulouse Archive Ouverte**

OATAO is an open access repository that collects the work of Toulouse researchers and makes it freely available over the web where possible

This is an author's version published in: http://oatao.univ-toulouse.fr/n° 20410

Official URL: <u>http://doi.org/10.1016/j.fluid.2017.08.010</u>

**To cite this version:** Toikka, Maria and Sadaeva, Anna and Samarov, Artemiy and Golikova, Alexandra and Trofimova, Maya and Shcherbakova, Nataliya<sup>20</sup> and Toikka, Alexander *Chemical equilibrium for the reactive system propionic acid + ethanol + ethyl propionate + water at 303.15 and 313.15 K.* (2017) Fluid Phase Equilibria, 451. 91-95. ISSN 0378-3812

Any correspondence concerning this service should be sent to the repository administrator: <u>tech-oatao@listes-diff.inp-toulouse.fr</u>

### Chemical equilibrium for the reactive system propionic acid + ethanol + ethyl propionate + water at 303.15 and 313.15 K

Maria Toikka <sup>a</sup>, Anna Sadaeva <sup>a</sup>, Artemiy Samarov <sup>a</sup>, Alexandra Golikova <sup>a</sup>, Maya Trofimova <sup>a</sup>, Nataliya Shcherbakova <sup>b</sup>, Alexander Toikka <sup>a, \*</sup>

<sup>a</sup> Department of Chemical Thermodynamics and Kinetics, Saint Petersburg State University, Universitetskiy Prospect 26, Peterhof, St. Petersburg 198504, Russia

<sup>b</sup> ENSIACET-INP Toulouse, Laboratoire de Génie Chimique, 4 allée Emile Monso, Toulouse, 31432, France

#### ABSTRACT

Chemical equilibrium for the quaternary system propionic acid + ethanol + ethyl propionate + water was experimentally studied at 303.15 and 313.15 K and atmospheric pressure. The chemically equilibrium compositions were determined by gas chromatography and nuclear magnetic resonance analytical methods. It is shown that chemical equilibrium is reached both in homogeneous and heterogeneous area of composition of reactive mixture. The liquid – liquid equilibrium data for the surface of chemical equilibrium were obtained. The thermodynamic constants of chemical equilibrium at 303.15 and 313.15 K were determined.

Keywords: Chemical equilibrium Ethyl propionate Phase equilibrium Reactive system

#### 1. Introduction

Heterogeneous systems with chemical reactions are complex objects and their study gives new thermodynamic information, allows establishing new physical and chemical regularities. At the same time it is obvious not only fundamental, but also the practical importance of the problem of the splitting of liquid reaction mixtures. Such effect fundamentally changes the technological process of chemical synthesis. The object of our study is a quaternary reacting system with ethyl propionate synthesis reaction. Ethyl propionate is widely used in the food processing, flavor industry and biofuel production [1,2]. This ester is an important reagent in pharmaceutical industries, e.g. for the production of some antimalarial drugs including pyrimethamine. Ethyl propionate has also other promising areas of application, for example, in the manufacture of lithium batteries [3]. The esterification is one of the main industrial methods of ester synthesis and the data on phase and chemical equilibria are necessary for the process design. Unfortunately despite a practical importance of ethyl propionate the data equilibrium properties of system propionic sets on

acid + ethanol + ethyl propionate + water are very limited.

The only experimental data related to the properties of propionic acid + ethanol + ethyl propionate + water system were presented by Macedo and Rasmussen in 1982 [4]. The vapor – liquid equilibrium (VLE) was determined for the ethyl propionate – propionic acid mixture at 358 and 368 K and pressure 134.28–573.05 mm Hg for 20 compositions of binary mixtures. The VLE data have been correlated by UNIFAC model.

The aim of our work is the study of CE in propionic acid + ethanol + ethyl propionate + water system at 303.15 and 313.15 K and atmospheric pressure in a wide range of compositions including homogeneous and heterogeneous areas of reactive solutions. Accordingly we determined the liquid – liquid equilibrium (LLE) for chemically equilibrium states, i.e. simultaneous LLE and CE. The experimental data allow expanding the database on the phase and chemical equilibria in reacting systems and give the additional opportunities for the design of the process of ethyl propionate synthesis.

### 2. Experimental

#### 2.1. Materials

Propionic acid (>0.98 mol fraction, BASF, Germany), ethanol

<sup>\*</sup> Corresponding author. E-mail address: a.toikka@spbu.ru (A. Toikka).

 Table 1

 The purities of the chemicals.

Substance	Purity, initial mole fraction	Purity, final mole fraction <sup>a</sup>
Propionic acid	0.980	0.990
Ethanol	0.960	0.960 <sup>b</sup>
Ethyl propionate	0.980	0.997
Water	0.999	0.999

 $^{a}$  The uncertainty is estimated to be ±0.002 mol fraction.

<sup>b</sup> Ethanol sample contains 0.040 mol fraction of water.

(>0.96 mol fraction, Vekton, Russia) and ethyl propionate (>0.98 mol fraction, Vekton, Russia) were purified by distillation, water was bidistilled. Purity was determined by gas chromatography (GC) and is given in Table 1. All physical-chemical constants of pure substances are in agreement with the NIST data [5]. The fact that water is one of the components of the system under the study, gives the ability to use ethanol containing a known amount of water instead of absolute alcohol: this facilitated the experimental procedure. The amount of water remaining in ethanol after distillation was taken into account in preparation of initial mixtures.

#### 2.2. CE determination

The study of the CE at 303.15 and 313.15 K was carried out using <sup>1</sup>H NMR spectroscopy (for homogeneous solutions) and GC analysis (for heterogeneous area) by methods describing in Refs. [6,7].

Initial quaternary mixtures of known overall compositions were prepared by gravimetric method in NMR-tubes (178 mm  $\times$  4,2 mm i.d.) using an analytical balance Shinko VIBRA HT-120CE (Japan) with the accuracy of 0.001 g. Hydrochloric acid was used as a catalyst in amount of 0.003 mol fraction. NMR-tubes were placed in the liquid thermostat (303.15 and 313.15 K). The temperature uncertainty was ±0.05 K. The CE was reached in no longer than 4 days (303.15 and 313.15 K). All samples were analyzed using a 500 MHz Bruker AVANCE III NMR spectrometer, equipped with a BBI probe head with inner coil for <sup>1</sup>H nuclei. The spectra were acquired with an acquisition time of 3 s, a relaxation delay of 1 s, and a pulse with 30° flip angle. The 16 scans were accumulated. The spin-lattice NMR relaxation times T<sub>1</sub> for all molecular groups of all compounds were measured. Observed T<sub>1</sub> values are in range from 3 to 5 s. The relaxation delay was high enough for quantitative analysis of peak integrals since the relaxation times had close values and the flip angle of pulse was relatively small. The error introduced by saturation was below 1%. The processing of the acquired spectra was carried out using Bruker TopSpin software. The phase correction was done manually. Polynomial baseline correction was done automatically. The integration region of 250-500 Hz which was over 20 times wider than the linewidth of analyzed peak was chosen. The constancy of composition confirmed the achievement of CE. The uncertainty of the determination of the peak areas introduced by the processing is estimated to be 3%.

In the case of the splitting of chemically equilibrium mixtures the compositions were determined using GC analysis. Gas chromatograph "Chromatec Crystal 5000.2" (Russia) with thermal conductivity detector (TCD) and packed column Porapak R  $(1 \text{ m} \times 3 \text{ mm i.d.})$  was used. The TCD was chosen because of the presence of water. Helium with the flow rate of 60 ml/min was used as a carrier gas. Operating temperature of column, vaporizing injector and TCD were 483 K, 503.15 K and 513 K, respectively. Initial quaternary mixtures were prepared by gravimetric method in glass vessels (5 ml) using an analytical balance Shinko VIBRA HT-120CE (Japan) with the accuracy of 0.001 g. The CE was reached in no longer than 4 days (303.15 and 313.15 K). After reaching of CE and phase equilibrium heterogeneous samples were taken from both phases with chromatographic syringe ("Hamilton", USA) and analyzed by GC. To avoid the splitting of samples directly into the needle the chromatographic syringe was preliminary heated. The methods of internal standard and relative calibration were used to determine the compositions of CE. Propionic acid was accepted as linking component. Uncertainty of GC analysis averaged  $\pm 0.005$  mol fraction. Accuracy of initial solutions preparing was 0.001 g. The temperature uncertainty of the liquid thermostat was  $\pm 0.05$  K. The reproducibility of the experimental data was confirmed by the repeated analysis of samples. All chemically equilibrium mixtures were analyzed 3-5 times and the deviations of the results were in the frame of GC analysis uncertainties.

To check the consistency between two analytical methods (GC and NMR), we also determined few compositions in homogeneous area by GC analysis. The discrepancy does not exceed 0.005 mol fractions. In general both of methods enable to accurately determine the compositions of mixtures, but NMR method takes less time in comparison with GC analysis [6].

#### 3. Results and discussions

#### 3.1. Experimental data

The data on CE in homogeneous area of composition in propionic acid + ethanol + ethyl propionate + water system at 303.15 and 313.15 K and atmospheric pressure (NMR method) are presented in Table 2.

The data of Table 2 gives the opportunity to calculate the values of so-called concentration constants (pseudoreaction equilibrium constants)

$$K_{x} = \frac{x_{ethyl \ propionate} x_{water}}{x_{propionic \ acid} x_{ethanol}}$$

which depends on the composition and should be not considered as real thermodynamic constant of CE. On the other hand the relative constancy of  $K_x$  can be used to check the consistency of the data. The average  $K_x$  values for homogeneous area of CE are 5.5  $\pm$  1.0 at 303.15 K and 5.0  $\pm$  0.9 at 313.15 K.

The consistency of the CE data is also confirmed by the fact that in all cases the initial and final (CE) compositions belong to the same stoichiometric line (within standard uncertainties u(x)).

The data on CE and LLE for the heterogeneous region at 303.15 and 313.15 K and atmospheric pressure (GC analysis) are presented in Table 3.

The compositions of chemically equilibrium mixtures in the case of ester synthesis reaction



Table 2

The experimental CE data for the quaternary system propionic acid (1) + ethanol (2) + ethyl propionate (3) + water (4) at 303.15 and 313.15 K in the homogeneous region of compositions analyzed by <sup>1</sup>H NMR spectroscopy (mole fractions  $x_i$ , atmospheric pressure<sup>a</sup>).

<i>x</i> <sub>1</sub>	<i>x</i> <sub>2</sub>	<i>x</i> <sub>3</sub>	<i>x</i> <sub>1</sub>	<i>x</i> <sub>2</sub>	<i>x</i> <sub>3</sub>
303.15 K					
0.165	0.131	0.475	0.064	0.299	0.156
0.151	0.127	0.548	0.071	0.303	0.197
0.121	0.181	0.244	0.067	0.276	0.259
0.096	0.182	0.158	0.081	0.295	0.318
0.224	0.028	0.064	0.056	0.294	0.390
0.253	0.061	0.191	0.071	0.282	0.394
0.270	0.064	0.249	0.066	0.281	0.453
0.270	0.064	0.207	0.410	0.007	0.039
0.286	0.063	0.356	0.465	0.015	0.103
0.281	0.057	0.430	0.447	0.021	0.151
0.250	0.063	0.466	0.473	0.020	0.184
0.036	0.172	0.044	0.424	0.026	0.230
0.055	0.205	0.085	0.466	0.022	0.271
0.066	0.209	0.113	0.447	0.023	0.367
0.074	0.223	0.152	0.436	0.023	0.392
0.081	0.234	0.191	0.011	0.367	0.035
0.101	0.220	0.234	0.025	0.372	0.000
0.095	0.221	0.234	0.034	0.333	0.037
0.103	0.232	0.356	0.046	0.387	0.168
0.098	0.221	0.388	0.053	0.379	0.209
0.090	0.200	0.489	0.053	0.382	0.301
0.324	0.023	0.096	0.055	0.368	0.336
0.332	0.032	0.144	0.058	0.348	0.361
0.337	0.034	0.170	0.516	0.005	0.037
0.339	0.041	0.217	0.526	0.009	0.070
0.353	0.039	0.238	0.520	0.012	0.100
0.337	0.045	0.319	0.526	0.013	0.134
0.331	0.043	0.302	0.518	0.016	0.188
0.342	0.042	0.551	0.509	0.017	0.220
0.044	0.283	0.089	0.516	0.016	0.301
0.038	0.444	0.272	0.520	0.015	0.336
0.010	0.449	0.037	_	_	_
313.15 K					
0.156	0.126	0.496	0.330	0.044	0.410
0.142	0.118	0.577	0.320	0.042	0.459
0.176	0.107	0.501	0.045	0.277	0.089
0.048	0.128	0.042	0.058	0.270	0.119
0.050	0.187	0.048	0.000	0.270	0.100
0.064	0.214	0.000	0.086	0.263	0.232
0.075	0.225	0.149	0.078	0.301	0.270
0.084	0.237	0.186	0.082	0.281	0.316
0.098	0.224	0.228	0.088	0.256	0.356
0.100	0.228	0.246	0.080	0.261	0.373
0.100	0.246	0.295	0.073	0.275	0.438
0.103	0.236	0.347	0.443	0.020	0.143
0.103	0.223	0.379	0.458	0.021	0.169
0.100	0.214	0.464	0.417	0.028	0.221
0.323	0.042	0.215	0.420	0.029	0.240
0.338	0.043	0.243	0.432	0.031	0.306
0.334	0.047	0.297	0.428	0.029	0.375
0.330	0.049	0.321	0.014	0.373	0.033
0.327	0.046	0.362	0.024	0.372	0.067
0.525	0.014	0.130	0.026	0.375	0.099
0.521	0.016	0.148	0.037	0.349	0.131
0.521	0.015	0.182	0.043	0.391	0.167
0.524	0.017	0.203	0.051	0.300	0.205
0.500	0.019	0.235	0.051	0.392	0.249
0.506	0.020	0.317	0.050	0.377	0.335
0.011	0.448	0.036	0.050	0.365	0.360
0.015	0.454	0.061	0.516	0.005	0.034
0.023	0.453	0.086	0.529	0.009	0.070
0.035	0.454	0.169	0.519	0.012	0.103
0.041	0.430	0.192	0.037	0.444	0.273
0.040	0.433	0.233	_	_	_

<sup>a</sup> Standard uncertainties u(x) = 0.005, u(T) = 0.05.

#### Table 3

The experimental CE and LLE data for the quaternary system propionic acid (1) + ethanol (2) + ethyl propionate (3) + water (4) in the heterogeneous region of compositions at 303.15 and 313.15 K (mole fractions  $x_i$ , atmospheric pressure<sup>a</sup>).

Aqueous phase			Organic phase		
<i>x</i> <sub>1</sub>	<i>x</i> <sub>2</sub>	<i>x</i> <sub>3</sub>	$\overline{x_1}$	<i>x</i> <sub>2</sub>	<i>x</i> <sub>3</sub>
303.15 K					
0.081	0.148	0.059	0.165	0.154	0.440
0.081	0.122	0.055	0.197	0.130	0.449
0.097	0.098	0.046	0.211	0.105	0.451
0.111	0.080	0.042	0.236	0.090	0.430
0.134	0.068	0.044	0.263	0.074	0.412
313.15 K					
0.081	0.154	0.069	0.162	0.161	0.424
0.086	0.129	0.057	0.191	0.137	0.427
0.098	0.103	0.051	0.210	0.111	0.430
0.112	0.089	0.049	0.241	0.094	0.412
0.127	0.071	0.047	0.256	0.080	0.393

<sup>a</sup> Standard uncertainties u(x) = 0.005, u(T) = 0.05.

belong to the surface in composition tetrahedron. Four binary subsystems (propionic acid + ethyl propionate, propionic acid + water, ethanol + ethyl propionate, ethanol + water) are nonreactive and should be considered as borders of the surface of CE (edges of tetrahedron). Two reacting binary systems (propionic acid + ethanol and ethyl propionate + water) and all four ternary subsystems are in chemically nonequilibrium states and do not belong to this surface. Fig. 1 presents the CE surface of the quaternary propionic acid + ethanol + ethyl propionate + water system at 313.15 K (the CE surface for 303.15 K has approximately the same view). The LLE tie-lines in Fig. 1 give only a schematic representation of liquid-liquid envelope on this surface. It should be noted that LLE tie-lines on the surface of CE are analogous to unique reactive tie-lines which were introduced by Ung and Doherty for heterogeneous CE in ternary mixtures [8]. In quaternary system the set of unique reactive tie-lines forms the linear surface in composition tetrahedron [7,9].

The square of transformed composition variables  $\alpha_i$  [10,11] gives more visual representation of liquid-liquid envelope at CE surface. In the case of quaternary CE these variables  $\alpha_i$  are determined as following:

Ν.	_	Υ.	_	v
$\alpha_1$	_	A1	+	л,

 $\alpha_2 = x_2 + x_4$ 

 $\alpha_3 = x_3 - x_4$ 

The LLE tie-lines on the surface of CE 313.15 K, area of immiscibility and experimental homogeneous CE composition at 313.15 K



**Fig. 1.** The CE surface of propionic acid + ethanol + ethyl propionate + water system at 313.15 K ■■■ - surface of CE; ● — ● - experimental liquid-liquid tie-lines on the surface of CE.



**Fig. 2.** The area of immiscibility at the surface of CE in propionic acid + ethanol + ethyl propionate + water system at 313.15 K:  $\_$  –  $\_$  – liquid-liquid tie-lines at 313.15 K; ---- – the area of intersection of CE and LLE surfaces;  $\Box$  – composition points corresponding to CE in the homogeneous area (experimental results).

are presented in Fig. 2. Figs. 1 and 2 also display the mutual disposition of CE and binodal surfaces. The area of intersection of these surfaces in the composition tetrahedron corresponds to simultaneous CE and LLE.

#### 4. Calculation of thermodynamic constant of CE

In this work, the original UNIFAC model [12] was used for the prediction of the activity coefficients. The activity coefficient of component i in a multicomponent mixture according to UNIFAC model determines by the following equation:

 $ln\gamma_i = ln\gamma_i^C + ln\gamma_i^R,$ 

where an activity coefficient consists of two parts, a combinatorial part  $\gamma_i^{c}$  and a residual part  $\gamma_i^{R}$ .

The combinatorial part can be expressed as:

$$ln\gamma_i^C = ln\left(\frac{\phi_i}{x_i}\right) + \frac{z}{2}q_i ln\left(\frac{\theta_i}{\phi_i}\right) + l_i - \frac{\phi_i}{x_i}\sum_j x_j l_j$$

wherein:

$$\theta_i = \frac{q_i x_i}{\sum_j q_j x_j}$$
$$\phi_i = \frac{r_i x_i}{\sum_j r_j x_j}$$

$$l_i = \frac{z}{2}(r_i - q_i) - (r_i - 1)$$

where z is the lattice coordination number which is equal to 10. Parameters  $q_i$  and  $r_i$  are calculated as the sum of the group volume and region parameters  $Q_k$  and  $R_k$ :

$$q_i = \sum_k \nu_k^{(i)} Q_k$$
$$r_i = \sum_k \nu_k^{(i)} R_k$$

where  $v_k^{(i)}$  is an integer that indicates the number of times that group *k* appears in molecule *i*; surface region and group volume parameters  $Q_k$  and  $R_k$  were taken from Ref. [13] due to their applicability for LLE calculation.

Following expression describes another part of the activity coefficient:

$$ln\gamma_i^R = \sum_k^G \nu_k^{(i)} \left( ln\Gamma_k - ln\Gamma_k^{(i)} \right)$$

where  $\Gamma_k$  is the group residual activity coefficient and  $\Gamma_k$ <sup>(i)</sup> is the residual activity coefficient of group *k* in a reference solution containing only molecules of type *i*.

$$ln\Gamma_{k} = M_{k}Q_{k}\left[1 - ln \quad \sum_{m} \Theta_{m}\Psi_{mk}\right) - \sum_{m}\left(\frac{\Theta_{m}\Psi_{mk}}{\sum_{m}\Theta_{m}\Psi_{mk}}\right)\right]$$

where  $\Theta_m$  is calculated in the same way as for  $\theta_i$ ;  $\Psi_m$  is the group-interaction parameter. They are calculated according to the equation:

$$\Theta_m = \frac{\Theta_m X_m}{\sum_G^n \Theta_n X_n}$$
$$\Psi_{mn} = exp\left(-\frac{U_{mn} - U_{nn}}{RT}\right) = exp\left(-\frac{a_{mn}}{T}\right)$$

where  $X_m$  is the mole fraction of group m in the solution;  $U_{mn}$  is the energy of interaction between groups m and n. The group interaction parameters  $a_{mn}$  were taken from the literature [13].

The UNIFAC model was used for calculations of activities and thermodynamic constant of CE,  $K_a$ . The value of  $K_a$  could be expressed in terms of activities of species:

$$K_a = \prod a_i^{\nu_i},$$

where  $\alpha_i$  – activity,  $\nu_i$  – stoichiometric coefficients of reactants and products that are negative and positive, respectively. The values of  $K_a$  were calculated on the basis of the experimental data for homogeneous area of composition. The obtained constants of CE have the value 18.4 ± 2.7 at 303.15 K and 17.2 ± 2.3 at 313.15 K.

#### 5. Conclusions

Chemical equilibrium data for the quaternary reacting system propionic acid + ethanol + ethyl propionate + water were obtained at 303.15 and 313.15 K and atmospheric pressure. The existence of liquid-liquid envelope on the surface of CE is experimentally proved. The set of new experimental data enables to present surface of CE with the area of its intersection with LLE surface in composition tetrahedron. Using experimental data and UNIFAC model the values of thermodynamic constants of CE at 303.15 and 313.15K were determined as  $18.4 \pm 2.7$  and  $17.2 \pm 2.3$  respectively.

#### Acknowledgments

Maria Toikka is grateful for the scholarship of President of

Russian Federation (SP-2140.2016.1) and support of Saint Petersburg State University (travel-grant 12.42.735.2017). Artemiy Samarov acknowledges SPbSU for the technical support (12.40.534.2017). Alexander Toikka is also grateful to SPbSU for the technical support of this study (grant SPbSU 12.40.518.2017). Nataliya Shcherbakova acknowledges INPT (Institut National Polytechnique de Toulouse) and LGC (Laboratoire de Génie Chimique, UMR 5503 CNRS) for the support of the Russian-French research project "Geometry of separation processes of multicomponent mixtures". The authors acknowledge Resource Center of Saint Petersburg State University "Magnetic Resonance Research Centre" for the help in analytical investigations. Authors are also grateful to NIST ThermoData Engine group for the data on the properties of considered system.

#### References

- F. Contino, F. Foucher, C. Mounaïm-Rousselle, et al., Experimental characterisation of ethyl acetate, ethyl propionate, and ethyl butanoate in a homogeneous charge compression ignition engine, Energy fuels.. 25 (2011) 998–1003, http://dx.doi.org/10.1021/ef101602q.
- [2] T. Badawy, J. Williamson, H. Xu, Laminar burning characteristics of ethyl propionate, ethyl butyrate, ethyl acetate, gasoline and ethanol fuels, Fuel 183 (2016) 627–640, http://dx.doi.org/10.1016/j.fuel.2016.06.087.
- [3] M.Z. Kufian, S.R. Majid, Performance of lithium-ion cells using 1 M LiPF6 in EC/ DEC (v/v=1/2) electrolyte with ethyl propionate additive, Ionics 16 (2010) 409–416, http://dx.doi.org/10.1007/s11581-009-0413-6.
- [4] E.A. Macedo, P. Rasmussen, Vapor liquid equilibrium for the binary systems

ethyl acetate – acetic acid and ethyl proponate – propionic acid, J. Chem. Eng. Data 27 (1982) 463–465, http://dx.doi.org/10.1021/je00030a028.

- [5] M. Frenkel, R.D. Chirico, V. Diky, A.F. Kazakov, C.D. Muzny, J.W. Magee, I. Abdulagatov, K. Kroenlein, J.W. Kang, NIST ThermoData Engine, NIST Standard Reference Database 103b-Pure Compounds, Binary Mixtures, and Chemical Reactions, Version 5.0; Standard Reference Data Program, National Institute of Standards and Technology, Gaithersburg, MD, 2010.
- [6] A. Golikova, A. Samarov, M. Trofimova, S. Rabdano, M. Toikka, O. Pervukhin, A. Toikka, Chemical equilibrium for the reacting system acetic acid – ethanol – ethyl acetate – water at 303.15 K, 313.15 K and 323.15 K, J. Sol. Chem. 46 (2017) 374–387, http://dx.doi.org/10.1007/s10953-017-0583-1.
- [7] A. Samarov, P. Naumkin, A. Toikka, Chemical equilibrium for the reactive system acetic acid + n-butanol + n-butyl acetate + water at 308.15 K, Fluid Phase Equilib. 403 (2015) 10–13, http://dx.doi.org/10.1016/ j.fluid.2015.06.001.
- [8] S. Ung, M.F. Doherty, Vapor-liquid phase-equilibrium in systems with multiple chemical reactions, Chem. Eng. Sci. 50 (1995) 23–48, http://dx.doi.org/ 10.1016/0009-2509(94)00180-Y.
- [9] M. Toikka, A. Toikka, Peculiarities of phase diagrams of reactive liquid–liquid systems, Pure Appl. Chem. 85 (2013) 277–288, http://dx.doi.org/10.1351/ PAC-CON-12-05-07.
- [10] V.T. Zharov, Open evaporation of solutions of reacting substances, Zh. Fiz. Khim. J. Phys. Chem. USSR) 44 (1970) 1967–1974 (in Russian).
- [11] D. Barbosa, M.F. Doherty, A new set of composition variables for the representation of reactive-phase diagrams, Proc. R. Soc. Lond. A 413 (1987) 459–464, http://dx.doi.org/10.1098/rspa.1987.0126.
- [12] A. Fredenslund, R.L. Jones, J.M. Prausnitz, Group-contribution estimation of activity coefficients in nonideal liquid mixtures, AIChE J. 21 (1975) 1086–1099, http://dx.doi.org/10.1002/aic.690210607.
- [13] T. Magnussen, P. Rasmussen, A. Fredenslund, UNIFAC parameter table for prediction of liquid-liquid equilibriums, Ind. Eng. Chem. Process Des. Dev. 20 (1981) 331–339, http://dx.doi.org/10.1021/i200013a024.