Non decomposable connectives of linear logic

Roberto Maieli

► To cite this version:

| Roberto Maieli. Non decomposable connectives of linear logic. 2018. hal-01864488

HAL Id: hal-01864488

<https://hal.science/hal-01864488>

Preprint submitted on 30 Aug 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Non decomposable connectives of linear logic

Roberto Maieli

*Department of Mathematics and Physics, "Roma Tre" University
Largo San Leonardo Murialdo 1, 00146 Rome, Italy
maieli@mat.uniroma3.it*

Abstract

This paper studies the so-called generalized multiplicative connectives of linear logic, focusing on the question of finding the “non-decomposable” ones, i.e., those that may not be expressed as combinations of the default binary connectives of multiplicative linear logic, $\otimes$ (*tensor*) and $\wp$ (*par*). In particular, we concentrate on generalized connectives of a surprisingly simple form, called “entangled connectives”, and prove a characterization theorem giving a criterion for identifying the undecomposable ones.

Keywords: linear logic, partitions sets, proof nets, sequent calculus.

2010 MSC: 03F52, 03B47.

Contents

1	Introduction	2
2	Partitions and orthogonality	6
3	Generalized connectives	8
3.1	Sequent calculus syntax for generalized connectives	9
3.1.1	Decomposable sequential connectives	10
3.2	Proof nets syntax for generalized connectives	11
3.2.1	Decomposable graphical connectives	12
3.2.2	Proof structures with generalized links	13
3.3	Sequentialization of decomposable connectives	15
3.4	Undecomposable connectives	16

4	Entangled connectives	18
4.1	Entangled types	18
4.2	Entangled connectives	30
4.3	Decomposable <i>normal form</i> of entangled connectives	31
4.4	Undecomposable entangled connectives	34
5	Sequentialization of undecomposable connectives	35
6	Conclusions and future works	36

1. Introduction

Generalized connectives for the pure (units free) multiplicative fragment of linear logic (MLL, [6]) were introduced by Girard in his seminal paper [5] in terms of permutations but most of the results known after then are essentially due to Danos and Regnier [3] who replaced permutations by partitions of finite sets. A generalized multiplicative connective may be defined by a pair of dual sets of partitions of a same finite domain $\{1, 2, \dots, n\}$, dual in the sense that they are pairwise orthogonal where orthogonality is defined by a topological condition: the bipartite graph obtained by linking together classes of each partition sharing an element is acyclic and connected (ACC). E.g., partition $\{(1, 2), (3)\}$ is not orthogonal to partition $\{(1, 2, 3)\}$ since the bipartite graph $\mathcal{G}_1$ contains a cycle while the two sets of partitions, $\{\{(1, 2), (3)\}\}$ and $\{\{(1, 3), (2)\}, \{(1), (2, 3)\}\}$, are orthogonal since they are pairwise so (their respective bipartite graphs, $\mathcal{G}'_2$ and $\mathcal{G}''_2$, are ACC) as illustrated below:

There are two ways to interpret logical connectives (formulas) of MLL by means of pairs of orthogonal sets of partitions: one based on the *sequent calculus syntax* and the other one based on the *proof nets syntax*, according to the spirit of linear logic. In the *sequential syntax*, a partition describes a sequent calculus *rule* for producing the generalized formula: the domain of the partition is the set of principal sub-formulas occurrences of the given formula and each class describes one premise of the rule; so, a multiplicative rule for an n -ary connective $F(A_1, \dots, A_n)$ is completely characterized by the *organization*

of its principal sub-formulas $(A_1, \dots, A_n)$; indeed, since multiplicatives rules are *unconditional* about the context, any rule can be simply described by a *partition* over its principal sub-formulas occurrences (their indexes), omitting the contexts. E.g., the two orthogonal sets of partitions (or organizations sets), $\mathcal{O}_F = \{\{(1, 2), (3)\}\}$ and $\mathcal{O}_{F^\perp} = \{\{(1, 3), (2)\}, \{(1), (2, 3)\}\}$, are compact representations of, respectively, the sequent generalized rule below (when $F(1, 2, 3) = (A_1 \wp A_2) \otimes A_3$)

$$\frac{\frac{(1, 2)}{F(1, 2, 3)} \quad (3)}{r(F)} \equiv \frac{\frac{(1, 2)}{(1 \wp 2)} \wp \quad (3)}{F = (A_1 \wp A_2) \otimes A_3} \otimes$$

and the next two sequent n -ary rules (when $F^\perp(1, 2, 3) = (A_1 \otimes A_2) \wp A_3$):

$$\frac{\frac{(1, 3)}{F^\perp(1, 2, 3)} \quad (2)}{r_1(F^\perp)} \equiv \frac{\frac{(1, 3)}{(1 \otimes 2), 3} \otimes}{F^\perp = (A_1 \otimes A_2) \wp A_3} \wp \quad \frac{(1) \quad (2, 3)}{F^\perp(1, 2, 3)} r_2(F^\perp) \equiv \frac{\frac{(1) \quad (2, 3)}{(1 \otimes 2), 3} \otimes}{F^\perp = (A_1 \otimes A_2) \wp A_3} \wp$$

In the *graphical syntax*, a partition describes the “effects” of a *Danos-Regnier switching* over the frontier of the syntactical tree of a MLL formula. We may associate to each switching of a formula tree $F(A_1, \dots, A_n)$ a partition of its top frontier $A_1, \dots, A_n$ in to classes of connected components. The partitions set of the border leaves of a formula tree F , induced by all switchings, is called the *pre-type* of F (denoted $\mathcal{P}_F$). E.g., the pre-types the two formula trees below

are resp., the partitions set $\mathcal{P}_F = \{\{(1, 3), (2)\}, \{(1), (2, 3)\}\}$ (pre-type of $F = (A_1 \wp A_2) \otimes A_3$) and the partitions set $\mathcal{P}_{F^\perp} = \{\{(1, 2), (3)\}\}$ (pre-type of $F^\perp = (A_1 \otimes A_2) \wp A_3$).

These two syntaxes describing usual MLL formulas (i.e., definable by means of the basic binary connectives $\otimes$ and $\wp$) are shown to be dual. To be precise, the organizations set of a MLL formula F (denoted $\mathcal{O}_F$) is exactly the dual of the pre-type of F that is, $\mathcal{O}_F = \mathcal{P}_F^\perp$ so, $\mathcal{O}_F$ is a *type* (i.e. it is equal to its bi-orthogonal, $\mathcal{O}_F = \mathcal{O}_F^{\perp\perp}$), thus every sequential n -ary connective of MLL, $\mathcal{O}_F$ and $\mathcal{O}_{F^\perp}$, is reflexive that is $\mathcal{O}_F^\perp = \mathcal{O}_{F^\perp}$ and $\mathcal{O}_{F^\perp}^\perp = \mathcal{O}_F$.

We can generalize this construction in a natural way. Given two partitions sets we wonder whether they define a generalized multiplicative connective according to the two dual syntaxes:

- in the sequent calculus syntax it is sufficient to describe a connective¹ as a pair of orthogonal organizations sets $\mathcal{O}_C \perp \mathcal{O}_{C^\perp}$; each organizations set is a *type* describing *all rules* that allow to derive that connective; orthogonality is enough to ensure cut elimination;
- in the proof nets syntax, since partitions are interpreted as switchings associated to a connective and since switchings are in some sense *dense sub-sets*² we may define a connective as a pairs of partitions sets P and Q s.t. $P \perp Q$ and $P^\perp \perp Q^\perp$: the first condition $P \perp Q$ ensures (ACC) correctness of proof structures while the second condition $P^\perp \perp Q^\perp$ ensures the stability of correctness under cut reduction.

Then, given a pair of partitions sets, P and Q , as describing a connective (in sequential or graphical syntax), we wonder whereas this connective is definable by means of the basic binary connective of MLL ($\wp$ and $\otimes$) in which case it is called *decomposable*. Not all connectives are decomposable. To date there only exists one instance of non (binary) decomposable connective firstly discovered by Girard [5] in terms of permutations and later reformulated by Danos-Regnier [3] as a pair of orthogonal sets of partitions, G_4 and $G_4^\perp$, over the same domain $\{1, 2, 3, 4\}$:

$$G_4 = \{(1, 2), (3, 4)\}, \{(2, 3), (4, 1)\}\} \quad \text{and} \quad G_4^\perp = \{(1, 3), (2), (4)\}, \{(2, 4), (1), (3)\}\}.$$

Unfortunately, none (uniform) characterization of undecomposable connectives is known up to now. Under this respect, this work represents a first step forward: it defines a class of generalized connectives that are not decomposable, neither in sequential nor in graphical syntax, the so-called *undecomposable entangled connectives*. A pair of distinct partitions sets with same domain and same weight (cardinality) is entangled when each partition contains only binary or unary classes (Definition 21). Then a connective, P and Q , is entangled whenever P or Q is an entangled pair. Entangled pairs have interesting properties that are investigated throughout Section 4. The first one (Theorem 32, one of the main results of the paper), being that every entangled pair of partitions sets is a type. This result has several consequences: decomposable entangled pairs (and so, decomposable connectives) have a very simple characterization (the *normal form* of Theorem 36) which can be further

¹Every connective is immediately given together with its dual.

²"Switchings should be seen as a dense subset of para-proofs", page 41 of [7].

used to prove that given an undecomposable entangled connective, P and Q , where P is an entangled pair, then P cannot be embedded into a decomposable type $T \supsetneq P$, s.t. T and Q become decomposable (see Corollary 40).

Although our notion of entanglement doesn't solve the admittedly, difficult general problem ("try to find an uniform characterization of the full class of undecomposable connectives"), it is far from being an "ad hoc" condition. Intuitively, an entangled type is naturally obtained as soon as we "superpose" (i.e., we sum) the pre-types of two *bipoles*³ having the same "skeleton" (i.e., the same abstract syntactical tree) up to cyclic permutation of their frontier (the border leaves; see Remarks 41). This fact is a novelty since the union of types is not in general a type while the intersection of types is always a type (Property 3 of Section 2). Indeed, entangled types are the smallest types (w.r.t. the number of partitions), if we exclude the trivial singleton types. So, entangled connectives can be considered, in some sense, "elementary connectives", since they are the "smallest" generalized multiplicative connectives (w.r.t. the number of switchings or the number of sequential rules), if we exclude, of course, the basic ones ($\wp$ and $\otimes$). However, the class of undecomposable entangled connectives is quite special and we already discovered examples of undecomposable connectives falling outside of it like e.g. G_9 below (we omit the dual $G_9^\perp$):

$$G_9 = \{ \begin{array}{l} \{(1, 2, 3), (4, 5, 6), (7, 8, 9)\}, \\ \{(2, 3, 4), (5, 6, 7), (8, 9, 1)\}, \\ \{(3, 4, 5), (6, 7, 8), (9, 1, 2)\} \end{array} \}.$$

We are currently working on a more general characterization⁴ of the full class of primitive non decomposable connectives i.e., those ones that cannot be defined by means of other connectives, neither binary nor entangled.

More generally, non decomposable generalized connectives witness a deep asymmetry between proof nets and sequent proofs since the former ones allow us to express a kind of parallelism of proofs that the latter ones cannot do: actually, there exist proof nets in *η -expanded form*, built on non decomposable connectives, that have no correspondence with any sequential proof, if we

³Naively, a *bipole* B is a special MLL formula, introduced by Andreoli in [1], with only two layers of connectives: a generalized $\wp$ of generalized $\otimes$ -sub-formulas. Bipoles have the nice feature that their pre-types are already types.

⁴For this scope we consider partitions over cyclic permutations of linearly ordered sequences $1 < 2 < \dots < n$; see Section 6.

exclude the identity axioms $\vdash G, G^*$. This significant fact, lying at the core of linear logic, was already remarked in [3]:

“We saw with some surprise that the realm of multiplicatives became quite complex, even handled by a careful generalization. Yet the generalization seems more natural in the non-sequential framework [...]. Maybe we witness here the limits of sequential presentations of logic”.

2. Partitions and orthogonality

Definition 1 (partitions and orthogonality). A **partition** of a finite set $X = \{1, \dots, n\}$ (also called support) is a set of nonempty subsets of X , called classes (or parts), such that every element $i \in X$ is in exactly one of these subsets (i.e., X is a disjoint union of the subsets).

If p and p^* are two partitions of $X = \{1, \dots, n\}$, then:

- the **induced graph of incidence** of p and p^* , denoted $G(p, p^*)$, is the bipartite (undirected) graph which has for vertices the classes of p and p^* , two of them being linked iff they share an element $i \in X$;
- p and p^* are **orthogonal** (denoted, $p \perp p^*$) iff the induced graph $G(p, p^*)$ is acyclic and connected (i.e., it is a topological tree; shortly, ACC).

If P and Q are two sets of partitions, we say that P and Q are orthogonal (denoted $P \perp Q$) if they are pointwise orthogonal i.e., $\forall p \in P, \forall q \in Q, p \perp q$. $P^\perp$ denotes the set of partitions orthogonal to all the elements of P .

Example 2. Given two partitions, $p_1 = \{(1, 2)\}$ and $p_1^* = \{(1), (2)\}$ then, $p_1 \perp p_1^*$ since the induced graph $\mathcal{G}_1(p_1, p_1^*)$ is a tree; conversely, partitions $p_2 = \{(1), (2, 3)\}$ and $p_2^* = \{(1, 2, 3)\}$ are not orthogonal ($p_2 \not\perp p_2^*$) since $\mathcal{G}_2(p_2, p_2^*)$ contains a cycle as illustrated in Figure 1. Moreover, observe that w.r.t the sets of partitions given below, P_1, P_2, Q_1 and Q_2 , we only have $P_1 \perp Q_1$ and $P_2 \perp Q_2$ while e.g. $P_1 \not\perp Q_2$ and $P_2 \not\perp Q_1$:

$$\begin{aligned} P_1 &= \{\{(1, 2), (3, 4)\}\}, & Q_1 &= \{\{(1, 3), (2), (4)\}, \{(1, 4), (2), (3)\}, \{(2, 3), (1), (4)\}, \{(2, 4), (1), (3)\}\} \\ P_2 &= \{\{(2, 3), (4, 1)\}\}, & Q_2 &= \{\{(2, 4), (3), (1)\}, \{(2, 1), (3), (4)\}, \{(3, 4), (2), (1)\}, \{(3, 1), (2), (4)\}\}. \end{aligned}$$

Figure 1: bipartite graphs of incidence induced by pairs of partitions

The **degree** of a class x is the cardinality of x ; the **weight** of a partition p is the cardinality of p . Given a set of partition P , we say that:

- P has weight w if every partition $p \in P$ has weight w ;
- the **size** of P is the cardinality of P ;
- the **dimension** of P is the cardinality of its domain X .

As notation, we use: variables $X, Y, Z, \dots$, for finite sets $\{1, \dots, n\}$, variables $a, b, c, \dots$, for arbitrary elements of sets, variables $p, q, r, \dots$, for partitions of a set, variables $x, y, z, \dots$, for classes of a partition and variables $A, B, P, Q, \dots$, for sets of partitions.

Next facts on partitions sets of (Property 3) can be found in the literature on multiplicatives (see e.g. [9] for detailed proofs).

Property 3 (partitions and types). *Let A, B and A_i be sets of partitions over $\{1, \dots, n\}$. Then the following facts hold:*

1. $A \perp B$ iff $A \subseteq B^\perp$ iff $B \subseteq A^\perp$;
2. $A \subseteq B$ implies $B^\perp \subseteq A^\perp$;
3. $A \subseteq A^{\perp\perp}$;
4. $A \perp B$ iff $A^{\perp\perp} \perp B^{\perp\perp}$;
5. $A^\perp = A^{\perp\perp\perp}$;
6. $(A \perp B \text{ and } A^\perp \perp B^\perp)$ iff $(A^\perp = B^{\perp\perp})$ iff $(B^\perp = A^{\perp\perp})$;
7. In case $A \perp B$ and $A^\perp \perp B^\perp$:
 $(B^\perp = A \text{ and } B = A^\perp)$ iff $(A^{\perp\perp} = A \text{ and } B = B^{\perp\perp})$;
8. $A = A^{\perp\perp}$ iff $\exists B : A = B^\perp$; then, A is called **type**;
9. $(\bigcup_i A_i)^\perp = \bigcap_i A_i^\perp$;
10. $(\bigcap_i A_i)^\perp \supseteq \bigcup_i A_i^\perp$;
11. if A is a singleton (i.e., $A = \{p\}$ where p is a partition) then A is type;
12. if $A \perp B$, with A and B nonempty, then all partitions in A (resp., in B) have the same weight.

Let us prove case 6 of Property 3: assume $(A \perp B)$ and $(A^\perp \perp B^\perp)$ then, by case 1, $(A^\perp \perp B^\perp) \Rightarrow (A^\perp \subseteq B^{\perp\perp})$ and $(A \perp B) \Rightarrow (A \subseteq B^\perp)$ which implies, by case 2, $B^{\perp\perp} \subseteq A^\perp$; so $B^{\perp\perp} = A^\perp$. Vice-versa assume $B^{\perp\perp} = A^\perp$ then, by case 1, $(A^\perp \subseteq B^{\perp\perp}) \Rightarrow (A^\perp \perp B^\perp)$ and, by case 3, $(B^{\perp\perp} \subseteq A^\perp) \Rightarrow (B \subseteq A^\perp)$ from which, by case 1, $(A \perp B)$.

We draw attention to the fact that the *intersection of types is always a type* (case 9) while the *union of types is not always a type* (case 10): instances of these facts can be found in Example 5.

Case 11 follows by Lemma 4 of [9]. Finally, case 12 follows by the next well known graph theoretical property (see e.g. [8], pages 250-251).

Property 4 (Euler-Poincaré invariance). *Given a graph $G = (V, E)$ then, $(|V| - |E|) = (|CC| - |Cy|)$ where $|V|$, $|E|$, $|CC|$ and $|Cy|$ denotes, resp., the number of vertices, edges, connected components and primitive cycles of G .*

Example 5. *Assume two sets of partitions P_1 and P_2 over $\{a, b, c, d\}$ as below then, we can calculate their respective orthogonal types, by case 9 of Property 3, as follows:*

1. *if $P_1 = \{p_1 : \{(a, c), (b, d)\}, p_2 : \{(a, d), (b, c)\}\}$ then,
 $P_1^\perp = \{p_1\}^\perp \cap \{p_2\}^\perp = \{\{(d, c), (a), (b)\}, \{(a, b), (d), (c)\}\};$*
2. *if $P_2 = \{p_1 : \{(a, d, c), (b)\}, p_2 : \{(d, b, c), (a)\}\}$ then,
 $P_2^\perp = \{p_1\}^\perp \cap \{p_2\}^\perp = \{\{(b, a), (c), (d)\}\}.$*

Concerning case 10 of Property 3, we just observe here that although every singleton $\{p_i\}$ is a type (with $i = 1, 2$, by case 11 of Property 3), only P_1 is a type while P_2 is not so, since $P_2 \subsetneq P_2^{\perp\perp} = (P_1 \cup P_2)$. We will show in Section 4.1 that as soon as we restrict to consider pairs of singleton sets of partitions, $\{p\}$ and $\{q\}$, having same domain and weight and whose classes have degree $1 \leq d \leq 2$ then, their union $\{p\} \cup \{q\}$ is a (n entangled) type.

3. Generalized connectives

Partitions sets allow to express generalized multiplicative connectives in a very abstract, uniform and compact way. A generalized (or n -ary) multiplicative connective $C(A_1, \dots, A_n)$ may be defined, as in [3], by a pair of orthogonal sets of partitions, $P \perp Q$, over a same domain $\{1, \dots, n\}$ that is the set of principal sub-formulas indexes of C . There are indeed two dual ways to interpret (orthogonal pairs of) sets of partitions as defining logical connectives: the *sequential* (Section 3.1) and the *graphical* one (Section 3.2).

3.1. Sequent calculus syntax for generalized connectives

Following [3], we may define a connective by means of its sequential rules which are *conservative* w.r.t. the atoms i.e, the premises of the rule have exactly the same atoms as in the conclusion. Hence a multiplicative rule for an n -ary (i.e., generalized) connective $C(A_1, \dots, A_n)$ is completely characterized by the *organization* of its principal sub-formulas occurrences $(A_1, \dots, A_n)$ as described in the l.h.s. picture of Figure 2. Since multiplicative rules are *unconditional* about the context, any rule can be simply described by a *partition* over its principal sub-formulas indexes, omitting the contexts $\Gamma_1, \dots, \Gamma_p$, as in the r.h.s. picture of Figure 2.

$$\frac{\vdash \Gamma_1, A_1, \dots, A_{i_1} \quad \dots \quad \vdash \Gamma_p, A_p, \dots, A_{i_p}}{\vdash \Gamma_1, \dots, \Gamma_p, C(A_1, \dots, A_n)} C \qquad \frac{(1, \dots, i_1) \quad \dots \quad (p, \dots, i_p)}{C(1, \dots, n)} C$$

Figure 2: generalized sequential rule

Definition 6 (generalized rules as partitions). *If R is an n -ary rule, then the corresponding partition p is defined over $\{1, \dots, n\}$ as follows: $i, j \in \{1, \dots, n\}$ belong to a same class of p if formulas occurrences A_i and A_j belong to the same premise of R . Conversely, given a partition p , we can check if a rule R satisfies it. Renumber the principal formulas from left to right in the conclusions, and ask if the induced partition is really p .*

Definition 7 (sequential multiplicative generalized connectives). *A sequential multiplicative generalized connective C consists of two sets of orthogonal partitions, $\mathcal{O}_C \perp \mathcal{O}_C^*$: a set $\mathcal{O}_C$ of partitions representing the right rules and a set $\mathcal{O}_C^*$ of partitions representing the left rules handled by the dual connective C^* . By duality, the set $\mathcal{O}_C^*$ of left rules for C are exactly the set of right rules for the dual connective C^* (i.e., $\mathcal{O}_C^* = \mathcal{O}_{C^*}$).*

E.g., the two orthogonal sets of partitions, $\{(1), (2)\}$ and $\{(1, 2)\}$, correspond to the *primitive* (or *basic*) binary multiplicative connectives “ $\wp$ ” and “ $\otimes$ ” of MLL [6]:

$$\frac{\vdash \Gamma, A_1 \quad \vdash \Delta, B_2}{\vdash \Gamma, \Delta, \otimes(A_1, B_2)} \otimes \qquad \frac{\vdash \Gamma, A_1, B_2}{\vdash \Gamma, \wp(A_1, B_2)} \wp$$

Orthogonality ensures cut elimination; e.g., $\mathcal{O}_D = \{p : \{(1, 2), (3)\}\}$ and $\mathcal{O}_{D^*} = \{p^* : \{(1, 2, 3)\}\}$ do not define a sequential connective since $\mathcal{O}_D \not\perp \mathcal{O}_{D^*}$;

we cannot actually reduce the cut between $D(A_1, A_2, A_3)$ and $D^*(A_1^\perp, A_2^\perp, A_3^\perp)$, introduced just above by rules p and p^* , by means of cuts between each couple of dual sub-formulas⁵, A_i and $A_i^\perp$:

$$\pi : \frac{\frac{\vdots}{\vdash \Gamma, A_1, A_2} \quad \frac{\vdots}{\vdash \Delta, A_3} p \quad \frac{\vdots}{\vdash \Sigma, A_1^\perp, A_2^\perp, A_3^\perp} p^*}{\vdash \Gamma, \Delta, D(A_1, A_2, A_3)} \quad \frac{\vdash \Sigma, D^*(A_1^\perp, A_2^\perp, A_3^\perp)}{\vdash \Gamma, \Delta, \Sigma} cut \quad \not\rightarrow$$

If we denote MLL^+ the standard MLL sequent calculus of Figure 3 extended with the sequential general multiplicative rules of Figure 2 then, MLL^+ still satisfies cut elimination⁶.

$$\frac{}{A, A^\perp} id \quad \frac{\Gamma, A \quad \Delta, A^\perp}{\Gamma, \Delta} cut \quad \frac{\Gamma, A \quad \Delta, B}{\Gamma, \Delta, A \otimes B} \otimes \quad \frac{\Gamma, A, B}{\Gamma, A \wp B} \wp$$

Figure 3: Standard MLL Sequent Calculus

3.1.1. Decomposable sequential connectives

Definition 8 (sequential decomposable connectives). A sequential connective, $\mathcal{O}_C$ and $\mathcal{O}_{C^*}$, is (binary) decomposable iff there exists an MLL formula F (built by means of $\otimes$ and $\wp$) and assigned to C s.t. the two following propositions hold:

1. if $p \in \mathcal{O}_C$ (resp., $q \in \mathcal{O}_{C^*}$) then F (resp., $F^\perp$) is provable in MLL, from p (resp., from q), by only means of the binary rules, $\wp$ and $\otimes$;
2. if p (resp., q) is a partition from which we can derive F in MLL (resp., $F^\perp$) by only means of binary rules, then $p \in \mathcal{O}_C$ (resp., $q \in \mathcal{O}_{C^*}$).

Example 9. The generalized connective $(\mathcal{O}_Y, \mathcal{O}_{Y^*})$ below is decomposable for $Y = (1 \wp 2) \otimes 3 \otimes 4$ and $Y^* = (1 \otimes 2) \wp 3 \wp 4$:

$$\mathcal{O}_Y = \{p : \{(1, 2), (3), (4)\}\}$$

$$\mathcal{O}_{Y^*} = \{q_1 : \{(1, 3, 4), (2)\}, q_2 : \{(2, 3, 4), (1)\}, q_3 : \{(1, 3), (2, 4)\}, q_4 : \{(1, 4), (2, 3)\}\}$$

$$\frac{\vdash 1, 2 \quad \vdash 3 \quad \vdash 4}{\vdash Y(1, 2, 3, 4)} p \quad \frac{\vdash 1, 3, 4 \quad \vdash 2}{\vdash Y^*(1, 2, 3, 4)} q_1 \quad \frac{\vdash 2, 3, 4 \quad \vdash 1}{\vdash Y^*(1, 2, 3, 4)} q_2 \quad \frac{\vdash 1, 3 \quad \vdash 2, 4}{\vdash Y^*(1, 2, 3, 4)} q_3 \quad \frac{\vdash 1, 4 \quad \vdash 2, 3}{\vdash Y^*(1, 2, 3, 4)} q_4$$

⁵This is known as the *key-step* of cut-reduction: “a key-step between two partitions succeeds iff they are orthogonal” (see Lemma 1 of [3]).

⁶See Theorem 3 of [3].

Remarks 10 (The packaging problem). “Partitions of a decomposable connective X and $X^\perp$ do not describe all proofs involving it, but only those ones that are independent from the context” [3]. E.g., assume we want to prove a sequent S with only two conclusions, $D = (A_1 \wp B_2) \otimes C_3$ and $D^* = (A_1^\perp \otimes B_2^\perp) \wp C_3^\perp$ then, either we introduce D (resp., D^*) only by its sequential rules $\mathcal{O}_D = \{(1, 2), (3)\}$ (resp., $\mathcal{O}_{D^*} = \{\{(1), (2, 3)\}, \{(1, 3), (2)\}\}$) or we build S stepwise by exploiting the decomposition of D (resp., D^*). Clearly the latter method is more powerful since there are proofs, like the one below, that cannot be derived by the former one. Last rule for D (reps., for D^*) cannot be binary (i.e., with two premises) because any attempt to derive S would build a premise with the other conclusion $D^\perp$ (resp., D) together with only some (not all) of the principal formulas, A, B and C (resp., $A^\perp, B^\perp, C^\perp$): such sequents are never derivable from atomic logical axioms.

$$\frac{\frac{\frac{\vdash A, A^\perp \quad \vdash B, B^\perp}{\vdash A, B, (A^\perp \otimes B^\perp)} \otimes}{\vdash (A \wp B), (A^\perp \otimes B^\perp)} \wp \quad \vdash C, C^\perp}{\vdash (A \wp B) \otimes C, (A^\perp \otimes B^\perp), C^\perp} \otimes \quad \equiv \quad \frac{\frac{\frac{\vdash A, A^\perp \quad \vdash B, B^\perp}{\vdash A, B, (A^\perp \otimes B^\perp)} \otimes \quad \vdash C, C^\perp}{\vdash (A \wp B) \otimes C, (A^\perp \otimes B^\perp), C^\perp} D}{\vdash (A \wp B) \otimes C, (A^\perp \otimes B^\perp) \wp C^\perp} \wp$$

In other words, all the binary rules yielding a decomposable connective X (resp., $X^\perp$) in the sequent calculus can be packed in an single rule iff these rules can be permuted in order to appear consecutively (as a “package”).

3.2. Proof nets syntax for generalized connectives

Dually w.r.t. the sequential view, we may define a connective by means of its internal behavior that is, by means of the the set of partitions induced over its border by the set of *switchings* (i.e., internal tests) as in [3].

Definition 11 (graphical generalized connectives). A generalized multiplicative connective in graphical syntax consists of two sets of partitions, P and Q , over the same domain $\{1, \dots, n\}$, s.t. $P \perp Q$ and $P^\perp \perp Q^\perp$.

Dually to the sequential case, $P_\wp = \{\{(a), (b)\}\}$ and $Q_\otimes = \{\{(a, b)\}\}$ denote the basic connectives “ $\wp$ ” and “ $\otimes$ ”. In the following, unless differently declared, the expression “generalized connectives” means “generalized connectives in the graphical setting” according to Definition 11.

3.2.1. Decomposable graphical connectives

Definition 12 (Danos-Regnier switchings, pre-types and types). Let F be a MLL formula built by the binary multiplicative connectives $\wp$ and $\otimes$ from its top border X (i.e. the literals of F); given the parse (or syntactical) tree of F , a Danos-Regnier-switching [3] of F is the graph obtained after the mutilation of one of the two premises for each $\wp$ node of F ; then the pre-type of F , denoted by $\mathcal{P}_F$, is given by the set of partitions over X induced by all DR-switchings of the parse tree of F . After the mutilation, the elements of the border belonging to a same connected component, constitute a class of the induced partition. We denote $\mathcal{T}_F$ the type $\mathcal{P}_F^{\perp\perp}$ of F .

Definition 13 (decomposable connectives). A connective (P, Q) , is MLL binary decomposable (or definable) iff there exists a formula F , only built with binary connectives of MLL, s.t. P is the pre-type of the parse tree of F and Q is the pre-type of the tree of the dual formula $F^\perp$.

In the following we say that a set of partitions P is decomposable if there exists a formula F , s.t. P is the pre-type of F (i.e., $P = \mathcal{P}_F$). Next Fact 14 on decomposable sets of partitions corresponds to Theorem 7 of [9].

Fact 14 (orthogonal of a decomposable set of partitions). If F is a MLL formula then, $\mathcal{P}_F^\perp = \mathcal{T}_{F^\perp}$.

Remarks 15 (expressiveness of the graphical syntax). Connectives defined according to Definition 11 are very expressive. Given a set of partitions P there exists a connective (P, Q) for every Q s.t. $Q^{\perp\perp} = P^\perp$, by case 6 of Property 3. E.g., given (the type) $P = \{p_1 = \{(1, 2), (3, 4), (5)\}, p_2 = \{(1, 3), (2, 4), (5)\}\}$, there exist at least two graphical connectives, as follows:

1. $(P, P^\perp)$ where, by case 9 of Property 3, $P^\perp = \{p_1\}^\perp \cap \{p_2\}^\perp$ is the set of flagged partitions below:

$\{p_1\}^\perp$:	(145) (3) (2) ✓	(145) (2) (3) ✓
	(135) (4) (2)	(125) (4) (3)
	(235) (4) (1) ✓	(325) (4) (1) ✓
	(245) (3) (1)	(345) (2) (1)
	(45) (13) (2)	(45) (12) (3)
	(25) (13) (4)	(35) (12) (4)
	(25) (14) (3) ✓	(35) (14) (2) ✓
	(35) (14) (2) ✓	(25) (14) (3) ✓
	(15) (23) (4) ✓	(15) (32) (4) ✓
	(15) (24) (3)	(15) (34) (2)
	(45) (23) (1) ✓	(45) (32) (1) ✓
	(35) (24) (1)	(25) (34) (1)

2. (P, R) with $R = \{\{(1, 4, 5), (2), (3)\}, \{(3, 2, 5), (4), (1)\}\} \subsetneq P^\perp$ (s.t. $R^{\perp\perp} = P^\perp$).

3.2.2. Proof structures with generalized links

A *generalized multiplicative proof structure* is a graph which has for edges formulas of MLL^+ and for vertices *links* depicted on the l.h.s. of Figure 4 and defined as follows:

- an *axiom link*, is a vertex with only a pair of dual atomic conclusions (incident edges), $\{(a, a^\perp)\}$;
- a *cut link*, is a vertex with only a pair of dual premises (incident edges), $\{(A, A^\perp)\}$;
- a *generalized link* P , denoted λ_P , is a vertex with premises $A_1, \dots, A_n$ and exactly one conclusion $F(A_1, \dots, A_n)$ (also denoted by the special symbol $\sharp$) where (P, Q) is a generalized connective with domain $1, \dots, n$ and a bijection $\varphi : i \mapsto A_i$; λ_P is immediately given together with its dual link λ_Q . We assume there exactly is a generalized link for each connective.

Figure 4: axiom, cut, generalized links (on l.h.s.) and switching (on r.h.s.)

In a proof structure, each formula is conclusion of exactly one link and premise of at most one link; in the special case of the *conclusions* of a proof structure, i.e., formulas that are not premises of any link, we add an handling vertex (in order to have a graph). Not all proof structures are *correct* that is, correspond (sequentialize) to derivable sequent proofs. In order to characterize those sequentializable among all proof structures we need an intrinsic (i.e. non inductive) *correctness criterion*. First, we extend the usual notion of DR-switching to generalized links. For the switch of a generalized link λ_P

Figure 5: reduction step of a generalized cut link

(where P is a generalized connective), chose a partition $p = \{x_1, \dots, x_h\} \in P$, then chose a class $x_i \in p$, that is elect the upper edges belonging to this class and adjacent to the corresponding vertex of λ_P , disconnect all other upper edges, then for each remaining class of $p \setminus \{x_i\}$ connect together the pending edges (disconnected before) to a new vertex as in the rightmost side picture of Figure 4. Then, the correctness criterion for generalized proof structures remains the same as the Danos-Regnier's one for MLL proof nets: *a generalized proof structure π is correct (it is a generalized proof net) iff for each choice of the generalized switches, the associated graph is acyclic and connected*. So, a generalized proof net is an *MLL proof net* when each link is an axiom, a cut or a basic binary $\wp$ and $\otimes$ link. We also compactly describe a basic link by the set of its switches interpreted as partitions over the whole border, i.e., the top border A, B together with the bottom border $\sharp = (A \otimes B)$ or $\sharp = (A \wp B)$: $\lambda_{\wp} = \{\{(A, \sharp), (B)\}, \{(A), (B, \sharp)\}\}$ and $\lambda_{\otimes} = \{\{(D, \sharp, C)\}\}$.

Remarks 16 (on cut reduction). The “computational meaning” of generalized connectives is assured by the fact that reducing in a proof net a cut between (dual) generalized connectives, C and C^* , preserves the correctness criterion [3]. This fact follows by Definition 11 of a graphical connective where unlike the sequential syntax, requiring only orthogonality between sets of partitions, $P \perp Q$, is not enough for getting the stability of correctness under cut reduction: orthogonality must “pass to the contexts” $P^\perp \perp Q^\perp$; thus, the first condition $P \perp Q$ ensures (ACC) correctness of proof structures while the second condition $P^\perp \perp Q^\perp$ ensures the stability of correctness under cut reduction. A pictorial view is given in Figure 5. E.g., the sequential generalized connective $P_X = \{\{(1, 2), (3)\}\}$ and $Q_{X^*} = \{\{(1, 3), (2)\}\}$ is not a connective in proof net syntax since, by Definition 11, $P_X^\perp = \{\{(1, 3), (2)\}, \{(2, 3), (1)\}\}$ is not orthogonal to $Q_{X^*}^\perp = \{\{(1, 2), (3)\}, \{(2, 3), (1)\}\}$. This intuitively means

Figure 6: a cut reduction between “pseudo orthogonal” links

that if we admit the generalized links λ_X and λ_{X^*} (corresponding, resp., to connectives P_X and Q_{X^*}) then, the proof net π of Figure 6 (obtained by cutting π_X , with conclusion $\sharp_X$, against π_{X^*} , with conclusion $\sharp_{X^*}$) does not reduce to a correct proof net: actually after the reduction of the cut link (X, X^*) , the reductum π' will contain both a cycle and a pair of disconnected components. For simplicity reasons, we assumed in Figure 6 that π_X (resp., π_{X^*}) is the proof net built by gluing two orthogonal generalized links (modules) λ'_X and λ_X (resp. λ'_{X^*} and λ_{X^*}) through their common border $\{1, 2, 3\}$.

3.3. Sequentialization of decomposable connectives

The most natural way to compare the two syntaxes, sequential and graphical, is through their *sequentialization*, i.e. a way to set a precise correspondence between sequential proofs and proof nets. Here we focus only on the comparison w.r.t. decomposable connectives (we refer to Section 5 the comparison w.r.t the undecomposable ones discussed in Section 3.4).

There is a strong link between sequential and parallel decomposable connectives as exemplified by the basic cases $\mathcal{O}_{\otimes} = \{\{(1, 2)\}\}^\perp = \mathcal{P}_{\otimes}^\perp$ and $\mathcal{O}_{\wp} = \{\{(1), (2)\}\}^\perp = \mathcal{P}_{\wp}^\perp$. Actually, the two syntaxes are orthogonal views of a same decomposable connective, as stated in Proposition 17 proved in [3].

Proposition 17 (Danos-Regnier sequentialization). *Assume X is a decomposable connective s.t. $\mathcal{P}_X$ is its pre-type (in proof net syntax) and $\mathcal{O}_X$ is its set of right rules (the organizations set in the sequential syntax) then: $\mathcal{O}_X \subseteq \mathcal{P}_X^\perp$ (de-sequentialization part) and $\mathcal{P}_X^\perp \subseteq \mathcal{O}_X$ (sequentialization part).*

E.g., assume $\mathcal{P}_X = \{\{(A, C), (B)\}, \{(B, C), (A)\}\}$ and $\mathcal{O}_X = \{\{(A, B), (C)\}\}$ with $X = (A \wp B) \otimes C$, then $\mathcal{O}_X = \mathcal{P}_X^\perp$ as illustrated in Figure 7.

Next fact immediately follows by Proposition 17 and case 8 of Property 3.

Fact 18 (organizations sets are types). *The organizations set $\mathcal{O}_F$ of a decomposable formula F is a type.*

Figure 7: an instance of sequentialization

Corollary 19 (reflexive connectives). *A connective (P, Q) is reflexive when $P^\perp = Q$ and $Q^\perp = P$. All decomposable connectives are reflexive in the sequential syntax.*

Proof. 1) let X and $X^\perp$ be a decomposable connective; 2) by Property 3, $(P_X \perp P_{X^\perp}) \Leftrightarrow (P_X^{\perp\perp} \perp P_{X^\perp}^{\perp\perp})$; 3) by Proposition 17, $(\mathcal{O}_X = P_X^\perp) \wedge (\mathcal{O}_{X^\perp} = P_{X^\perp}^\perp)$; 4) by 1, $\mathcal{O}_X \perp \mathcal{O}_{X^\perp}$ then, by 2 and 3, $\mathcal{O}_X^\perp \perp \mathcal{O}_{X^\perp}^\perp$; 5) by 4 and Property 3 (case 7), it finally follows $(\mathcal{O}_X = \mathcal{O}_{X^\perp}^\perp) \wedge (\mathcal{O}_{X^\perp} = \mathcal{O}_X^\perp)$ since by 3, $(\mathcal{O}_X = \mathcal{O}_{X^\perp}^\perp) \wedge (\mathcal{O}_{X^\perp} = \mathcal{O}_X^\perp)$. $\square$

By reflexivity, it is enough to indicate only one of the two organizations sets of a sequential connective, $\mathcal{O}_C$ or $\mathcal{O}_{C^*}$, while for a graphical connective we have to indicate a pair of pre-types.

Example 20. *We may now interpret the two pairs of orthogonal sets of partitions, (P_1, Q_1) and (P_2, Q_2) of Example 2, as two decomposable connectives, according to the two dual points of view seen above: the sequential one as in Figure 8 and the graphical one as in Figure 9 where the decomposable graphical connectives are displayed as binary trees enclosed in dotted boxes:*

$$P_1 = \{(1, 2), (3, 4)\} \text{ and } Q_1 = \{(1, 3), (2, 4)\}, \{(1, 4), (2, 3)\}, \{(2, 3), (1, 4)\}, \{(2, 4), (1, 3)\}$$

$$P_2 = \{(2, 3), (4, 1)\} \text{ and } Q_2 = \{(2, 4), (3, 1)\}, \{(2, 1), (3, 4)\}, \{(3, 4), (2, 1)\}, \{(3, 1), (2, 4)\}.$$

3.4. Undecomposable connectives

Not all generalized connectives are binary decomposable; e.g. the sequential connective $\mathcal{O}_X = \{(1, 2), (3)\}$ and $\mathcal{O}_{X^*} = \{(1, 3), (2)\}$ seen in Remark 16 is undecomposable w.r.t. the basic connectives $\wp$ and $\otimes$ (it is neither a connective in the graphical syntax, indeed). Anyway, if we complete

$$\begin{array}{cc}
\frac{(1,2) \quad (3,4)}{(1 \wp 2) \otimes (3 \wp 4)} p_1 & \frac{(1,3) \quad (2) \quad (4)}{(1 \otimes 2) \wp (3 \otimes 4)} q_1^1 \quad \frac{(1,4) \quad (2) \quad (3)}{(1 \otimes 2) \wp (3 \otimes 4)} q_1^2 \\
& \frac{(2,3) \quad (1) \quad (4)}{(1 \otimes 2) \wp (3 \otimes 4)} q_1^3 \quad \frac{(2,4) \quad (1) \quad (3)}{(1 \otimes 2) \wp (3 \otimes 4)} q_1^4 \\
\\
\frac{(2,3) \quad (4,1)}{(2 \wp 3) \otimes (4 \wp 1)} p_2 & \frac{(2,4) \quad (3) \quad (1)}{(2 \otimes 3) \wp (4 \otimes 1)} q_2^1 \quad \frac{(2,1) \quad (3) \quad (4)}{(2 \otimes 3) \wp (4 \otimes 1)} q_2^2 \\
& \frac{(3,4) \quad (2) \quad (1)}{(2 \otimes 3) \wp (4 \otimes 1)} q_2^3 \quad \frac{(3,1) \quad (2) \quad (4)}{(2 \otimes 3) \wp (4 \otimes 1)} q_2^4
\end{array}$$

Figure 8: generalized decomposable connectives in sequent calculus syntax

Figure 9: generalized decomposable connectives in proof net syntax

$\mathcal{O}_{X^*}$, by adding the extra partition $\{(2,3), (1)\}$, we get the decomposable connective $\mathcal{O}_D$, $\mathcal{O}_{D^*}$ seen in Remark 10. Similarly, $\mathcal{O}_Z = \{\{(1,2), (3), (4)\}\}$ and $\mathcal{O}_{Z^*} = \{\{(1,3,4), (2)\}, \{(2,3,4), (1)\}\}$, is not a decomposable connectives but, as soon as we complete $\mathcal{O}_{Z^*}$ by adding a couple of extra partitions, $\{(1,3), (2,4)\}$ and $\{(1,4), (2,3)\}$, we get the decomposable connective $\mathcal{O}_Y$ and $\mathcal{O}_{Y^*}$ of Example 9.

A more serious example is the famous *Girard's connective* $(G_4, G_4^\perp)$:

$$G_4 = \{\{(1,2), (3,4)\}, \{(2,3), (4,1)\}\} \quad G_4^\perp = \{\{(1,3), (2), (4)\}, \{(2,4), (1), (3)\}\}$$

along with some other instances below (we omit their orthogonal)

$$\begin{array}{ccc}
G_6' = \{ \{(1,2), (3,4), (5,6)\}, & G_6'' = \{ \{(1,2,3), (4,5,6)\}, & G_9 = \{ \{(1,2,3), (4,5,6), (7,8,9)\}, \\
\{(2,3), (4,5), (6,1)\} \} & \{(2,3,4), (5,6,1)\}, & \{(2,3,4), (5,6,7), (8,9,1)\}, \\
& \{(3,4,5), (6,1,2)\} \} & \{(3,4,5), (6,7,8), (9,1,2)\} \}.
\end{array}$$

As we will see in Section 4.3, the pregnancy of Girard's connectives is given by the fact that, unlike the two previous ones, (X, X^*) and (Z, Z^*) , there is no chance to complete them in such a way they become decomposable

connectives neither in the sequential nor in the graphical setting. That is consequence of the fact that connectives like $(G_4, G_4^\perp)$ are described by pairs of types. Moreover, sequents like $\vdash G_4, G_4^\perp$ are not provable *tout court* in MLL^+ (if we exclude the trivial non atomic logical axioms) because of the “packaging problem” seen in Remark 10. Each rule for G_4 (resp., for $G_4^\perp$) is binary (resp., ternary) so any attempt to derive G_4 (resp., $G_4^\perp$) would build a premise (of sequent $\vdash G_4, G_4^\perp$) with the other conclusion together with only some (not all) of the principal formulas of the applied rule; so, sequents like $\vdash G_4, G_4^\perp$ are never logically provable from atomic logical axioms. Finally, since G_4 (resp., $G_4^\perp$) is not decomposable and since there is no chance to embed it into a decomposable connective, sequent $\vdash G_4, G_4^\perp$ is not even stepwise derivable.

The study of a “minimal” class of undecomposable connectives will be the subject of next sections.

4. Entangled connectives

In this section we introduce a special class of sets of partitions: *entangled pairs* (Definition 21) which is then exploited to define the class of *entangled connectives* (Definition 33). Entangled pairs have interesting properties that are investigated throughout this Section 4. The first one (Theorem 32, one of the main results of the paper), being that every entangled pair of partitions sets is a type. This result has several consequences: decomposable entangled pairs (and so, decomposable connectives) have a very simple characterization (the *normal form* of Theorem 36), which can be further used to prove that given an undecomposable entangled connective, P and Q , where P is an entangled pair, P cannot be embedded into a decomposable type $T \supsetneq P$, s.t. T and Q become decomposable (Corollary 40).

4.1. Entangled types

Definition 21 (entangled pairs of partitions). *Let P be a pair of nonempty distinct partitions, p_1 and p_2 , on the same (finite) domain $X = \{1, \dots, n\}$. P is an entangled pair if it satisfies the following two conditions:*

1. p_1 and p_2 have same weight;
2. each class belonging to p_1 or p_2 has degree 1 or 2.

Definition 22 (restriction of entangled pairs). *Given a set of partitions P on a finite domain X and given an element $a \in X$, the **restriction of P***

by point a (also a -restriction of P), denoted $P^{(\downarrow a)}$, is the set of partitions obtained after erasing a from each partition of P . A restriction of P by a is **conservative** when P and $P^{(\downarrow a)}$ have the same weight.

We say that a restriction of an entangled pair P by a point a is **admissible** when $P^{(\downarrow a)}$ is in turn an entangled pair; hence, an entangled pair P is **restrictable** if there exists an admissible restriction of P by a point.

We say that an entangled pair P is **restrictable only in one point** (also P **admits only one restriction by a single point**) iff $\exists a$ s.t. $P^{(\downarrow a)}$ is an entangled pair and $\neg \exists b \neq a$ s.t. $P^{(\downarrow b)}$ is an entangled pair.

We can extend the notion of restriction to sets of points of the support, i.e., $P^{(\downarrow Y)}$ where $Y \subseteq X$.

Example 23. Both P_1 and $P_1^\perp$ of Example 5 are entangled pairs while neither P_2 nor $P_2^\perp$ is so. Sets P_3 , P_4 and P_4' below are not restrictable entangled pairs; P_5 and P_6 below are both conservative restrictable entangled pairs: in particular, P_5 admits two restrictions, one by point c and another one by point d , while P_6 admits only one restriction, that one by single point a :

$$\begin{aligned} P_3 &= \{ \{(a,b),(c)\}, \{(a,c),(b)\} \} & P_5 &= \left\{ \begin{array}{l} \{(a,c),(b),(d,e),(f)\}, \\ \{(b,c),(a),(d,f),(e)\} \end{array} \right\} \\ P_4 &= \{ \{(a,b),(c),(d)\}, \{(c,d),(a),(b)\} \} & P_6 &= \left\{ \begin{array}{l} \{(a,b),(c),(d,e),(f),(g)\}, \\ \{(a,c),(b),(f,g),(d),(e)\} \end{array} \right\} \\ P_4' &= \left\{ \begin{array}{l} \{(a,b),(c,d),(e),(f),(g),(h)\}, \\ \{(e,f),(g,h),(a),(b),(c),(d)\} \end{array} \right\}. \end{aligned}$$

Lemma 24 (orthogonality of non restrictable entangled pairs). *If P is a non restrictable entangled pair then, its orthogonal $P^\perp$ is not empty.*

Proof. Since $P = \{p_1, p_2\}$ is an entangled pair, every element of the support, $a \in X$, must occur in each p_i either as singleton class (a) or in a pair class (a,b) for some $b \in X$ with $a \neq b$. Moreover, since P is not restrictable it cannot be the case that the same point a occurs:

1. as singleton class (a) in both p_1 and in p_2 , or
2. in a pair class in both p_1 and p_2 (that is, it cannot be the case that e.g., $(a,b) \in p_1$ and $(a,c) \in p_2$) except when P is in form (1) below, in which case we have $P^\perp = \{\{(a),(b,c)\}\}$ by calculation.

$$P = \left\{ \begin{array}{l} p_1 : \{(a,b),(c)\}, \\ p_2 : \{(a,c),(b)\} \end{array} \right\}; \quad (1)$$

In other words, if P is a non restrictable pair distinct from (1) then $\forall a \in X$ one of the following two cases holds:

- (i) in case the singleton class (a) occurs in p_1 (resp., occurs in p_2) then a must occur in a pair class (a, b) of p_2 (resp., in a pair class (a, b) of p_1);
- (ii) in case a occurs in a pair class (a, b) of p_1 (resp., of p_2) then a must occur as singleton class (a) in p_2 (resp., in p_1);

otherwise the erasing of point a in a non restrictable P would produce two restricted sets, $p_1^{(\downarrow a)}$ and $p_2^{(\downarrow a)}$, that result to be either equal ($p_1^{(\downarrow a)} = p_2^{(\downarrow a)}$), in case P is in form (1), or with distinct weights, $w_1 \neq w_2$, contradicting Definition 21. This means that P must appear as a generalization of P_4 or P'_4 of Example 23, that is, the domain X of P must have dimension $2n$ and form $\{a_1, \dots, a_n\} \uplus \{b_1, \dots, b_n\}$; in other words, X can be thought as the disjoint sum of two segments of points, $S_1 = \{a_1, \dots, a_n\}$ and $S_2 = \{b_1, \dots, b_n\}$:

- in the first one, S_1 (resp., in the last one, S_2), we find all points that occur both inside pair classes in p_1 and inside singleton classes in p_2 ;
- in the second one (resp., in the first one) we find all points that occur both inside singleton classes in p_1 and inside pair classes in p_2 .

So P is in form (2) below (where n is an even integer):

$$P = \left\{ \begin{array}{l} p_1 : \{(a_1, a_2), (a_3, a_4), \dots, (a_{n-1}, a_n), (b_1), \dots, (b_n)\}, \\ p_2 : \{(b_1, b_2), (b_3, b_4), \dots, (b_{n-1}, b_n), (a_1), \dots, (a_n)\} \end{array} \right\} \quad (2)$$

that is, every a_i (resp., b_i) must occur together with an other point a_j (resp., b_j) in a class of p_1 (resp., p_2) moreover, a_i and a_j (resp., b_i and b_j) must occur as singleton classes, (a_i) and (a_j) (resp., (b_i) and (b_j)), in p_2 (resp., p_1). Observe that P has weight $n/2 + n$, by the way we build form (2).

We can now build a partition $q = \{y_1, \dots, y_m\}$ with weight $m = n/2 + 1$ as follows (also displayed both on the top and on the bottom side of Figure 10):

- first elect a class, e.g. y_1 , containing all the elements of the domain s.t. they do not form pairwise a class of p_i ; the degree of y_1 will be n ;
- then, chose each remaining class, y_j with $2 \leq j \leq m$, in such a way that it consists of only two elements: one belonging to a class of p_i with two elements and the other one belonging to a class of p_i with a single element, as the top (resp., bottom) side picture of Figure 10; there are $n/2$ such classes.

By calculation partition q is orthogonal to each p_i since by the Euler-Poincaré law, $|V| - |E| = 1$ where V and E are resp., vertexes and edges of each induced graph $\mathcal{G}_i(q, p_i)$ with $|V| = n/2 + n + n/2 + 1 = 2n + 1$ and $|E| = 2n$ as depicted in Figure 10; therefore $P^\perp \neq \emptyset$. $\square$

Figure 10: orthogonal of non restrictable entangled pair

Proposition 25 (restrictable entangled pairs). *Let $P = \{p_1, p_2\}$ be an entangled pair that is restrictable by point a ; assume the orthogonal of the a -restriction of P , i.e., $(P^{\downarrow a})^\perp = \{q_1, \dots, q_m\}$, is nonempty, then the set of partitions $P^\sharp$ built according to the two cases below is orthogonal to P .*

1. *If a is singleton class in both p_1 and p_2 , i.e., P is in form (3)*

$$P = \left\{ \begin{array}{l} \{(a), x_2, \dots, x_n\}, \\ \{(a), y_2, \dots, y_n\} \end{array} \right\} \quad (3)$$

then, $P^\sharp$ is built as follows: for each partition q_i , $1 \leq i \leq m$, insert point a in a class x_i^j of q_i ; repeat this insertion of point a for each class x_i^j in q_i (each insertion produces a “pumped” partition also denoted $q_i^a \in P^\sharp$).

2. *Otherwise, if a belongs to a class with two elements, both in p_1 and p_2 , i.e. P is in form (4) below*

$$P = \left\{ \begin{array}{l} \{(a, b), x_2, \dots, x_n\}, \\ \{(a, c), y_2, \dots, y_n\} \end{array} \right\} \quad (4)$$

then, $P^\sharp$ is built as follows: insert (or “pump”) the singleton class (a) in each partition $q_i \in P^{(\downarrow a)^\perp}$; so $P^\sharp$ is as in form (5) below (each insertion produces a “pumped” partition also denoted $q_i^a \in P^\sharp$):

$$P^\sharp = \{q_1 \cup \{(a)\}, \dots, q_i \cup \{(a)\}, \dots, q_m \cup \{(a)\}\} \quad (5)$$

Proof. By Euler-Poincaré Property 4, every pumped partition $q_i^a \in P^\sharp$ is orthogonal to every partition $p \in P$ (the induced graph $\mathcal{G}(p, q_i^a)$ is ACC). $\square$

Fact 26 (entangled pair of dimension 3). *If the dimension of an entangled pair P is 3 (i.e. $|X| = 3$) then P is in form (1) and its weight is 2.*

Proof. It follows by calculation: actually, in order to have an entangled pair, P must have at least weight $w \geq 2$ and in case weight w is exactly 2 then, at least one class must have degree 2; thus, if $|X| = 3$ then P must be in form (1). $\square$

Lemma 27 (orthogonality of entangled pairs). *The orthogonal $P^\perp$ of an entangled pair P is not empty.*

Proof. By induction on the dimension $|X|$ of P .

The **base of induction** is when $|X| = 3$ then P must be in form (1), by Fact 26, so its weight w is 2. Thus, P is non restrictable and so by Lemma 24 its orthogonal is not empty.

For the **induction step** $|X| > 3$ we split our reasoning in two main cases:

1. if P is **restrictable by point** a , it is in form (3) or (4), then the orthogonal of the restriction $P^{(\downarrow a)}$, let us say $(P^{(\downarrow a)})^\perp = \{q_1, \dots, q_m\}$, is not empty by hypothesis of induction, so by Proposition 25, $P^\sharp \subseteq P^\perp$;
2. otherwise, if P is **not restrictable**, then we conclude by Lemma 24.

$\square$

Theorem 28 (non restrictable entangled types). *If P is a non restrictable entangled pair (i.e., $\neg \exists a \in X$ s.t. $P^{(\downarrow a)}$ is an entangled pair) then P is a type.*

Proof. If $P = \{p_1, p_2\}$ is a non restrictable entangled pair then, by Lemma 24, either P has form (1) or P has form (2). In case P has form (1) then it is a type (simply by calculation). Otherwise, if P has the form (2) then, we reason by absurdum, assuming that P is not a type; so, let $p_3 \neq p_{i=1,2}$ be a

partition belonging to $P^{\perp\perp}$. We reason on the classes of p_3 . Observe that p_3 cannot contain only singleton classes, otherwise p_3 will have a weight $(2n)$ greater than the weight of p_1 and p_2 , contradicting case 12 of Property 3 (by Lemma 27, $P^\perp$ is nonempty); thus p_3 must contain at least a class with at least two points. Moreover, since p_3 differs from p_1 and p_2 , it must satisfy at least one of the following three cases:

1. p_3 contains a class x with two points, a_i and a_j (resp., b_i and b_j) which belong to two distinct classes of p_1 (resp., p_2);
2. p_3 contains a class x with two points a_i and b_j s.t. $a_i \in \{a_1, \dots, a_n\}$ and $b_j \in \{b_1, \dots, b_n\}$;
3. p_3 contains two classes x_1 and x_2 s.t. x_1 belongs to p_1 and x_2 belongs to p_2 (resp., x_1 belongs to p_2 and x_2 belongs to p_1).

We show now that for each of these three cases we get a contradiction. Indeed, since P has the form (2) then, for each p_3 satisfying (at least) one of cases 1–3, we can build a partition $q = \{y_1, \dots, y_m\}$ in $P^\perp$, with weight $m = n/2 + 1$ like e.g. partition q depicted on the top (resp., bottom) side of Figure 10 in the proof of Lemma 24. We show that $p_3 \not\leq q$.

1. Case 1 is not consistent ($p_3 \not\leq q$) because we can chose q in such a way that class y_1 contains two points a_i and a_j (resp., b_i and b_j), occurring resp., in two distinct classes of p_1 (resp., p_2), like e.g. class $(a_1, \dots, a_{i-1}, \dots, a_{n-1}, b_1, \dots, b_{i-1}, \dots, b_{n-1})$ of Figure 10.
2. Case 2 is not consistent ($p_3 \not\leq q$) because we can chose q in such a way that a class y_j contains exactly two points, a_i and b_j , like e.g., class (a_2, b_2) , (a_i, b_i) , (a_n, b_n) or $(a_1, \dots, a_{i-1}, \dots, a_{n-1}, b_1, \dots, b_{i-1}, \dots, b_{n-1})$ in Figure 10.
3. Finally, case 3 is not consistent ($p_3 \not\leq q$) because if we assume two classes like e.g. (a_1, a_2) and (b_1, b_2) belonging to p_3 , then we can chose q containing two classes like e.g. $(a_1, \dots, a_{i-1}, \dots, a_{n-1}, b_1, \dots, b_{i-1}, \dots, b_{n-1})$ and (a_2, b_2) in Figure 10 (thus, the graph of incidence $\mathcal{G}(p_3, q_i)$ will contain a cycle).

□

Lemma 29 (restrictable entangled pairs). *Let $P = \{p_1, p_2\}$ be a restrictable entangled pair and let $p_3 \neq p_{i=1,2}$ be a partition in the type $P^{\perp\perp}$. If some a -restriction of P is a type (i.e., $P^{\downarrow a} = \{p_1^{(\downarrow a)}, p_2^{(\downarrow a)}\} = (P^{(\downarrow a)})^{\perp\perp}$ for some $a \in X$) then, the restriction $p_3^{(\downarrow a)}$ belongs to $P^{\downarrow a}$.*

Proof. Assume P admits a **non conservative restriction** by a point $a \in X$ that occurs as singleton class (a) both in p_1 and in p_2 i.e., P is in form (3), $P = \{p_1 = \{(a), x_2, \dots, x_n\}, p_2 = \{(a), y_2, \dots, y_n\}\}$, and let $p_3 \neq p_{i=1,2}$ be a partition belonging to the type $P^{\perp\perp}$. Point a must also occur in p_3 as singleton class (a) , i.e. $p_3 = \{(a), z_2, \dots, z_n\}$, otherwise we can find a partition $q_i^a \in P^\# \subseteq P^\perp$ (by Proposition 25 and Lemma 27) s.t. the induced graph $G(p_3, q_i^a)$ contains a cycle (where q_i^a denotes a partition of $\{q_1, \dots, q_m\} = (P^{(\downarrow a)})^\perp$ in which point a has been “pumped” as in case (1) of Proposition 25). This implies $p_3^{(\downarrow a)} \perp q_i$, $\forall q_i \in (P^{(\downarrow a)})^\perp$, i.e., $\{p_3^{(\downarrow a)}\} \perp (P^{(\downarrow a)})^\perp$, since, by Euler-Poincaré Property 4, $\mathcal{G}(p_3^{(\downarrow a)}, q_i)$ is ACC iff $\mathcal{G}(p_3, q_i^a)$ is ACC. Now, by assumption $P^{(\downarrow a)} = (P^{(\downarrow a)})^{\perp\perp}$ so, $\{p_3^{(\downarrow a)}\} \subseteq P^{(\downarrow a)}$, by case 1 of Property 3.

Otherwise, assume P admits only **conservative restrictions** by a point, i.e., assume P is a -restrictable as in form (4), $P = \{p_1 = \{(a, b), x_2, \dots, x_n\}, p_2 = \{(a, c), y_2, \dots, y_n\}\}$ for some a in X . Then, point a cannot occur in p_3 as singleton class (a) , otherwise, as before, we can find a partition $q_i^a \in P^\# \subseteq P^\perp$ (by Proposition 25 and Lemma 27) s.t. graph $G(p_3, q_i^a)$ has a disconnected component (where q_i^a denotes a partition of $\{q_1, \dots, q_m\} = (P^{(\downarrow a)})^\perp$ in which the singleton class (a) has been “pumped” as in case (2) of Proposition 25). So p_3 is in form $\{(a, d), z_2, \dots, z_n\}$ and this means that $p_3^{(\downarrow a)}$ is orthogonal to every $q_i \in (P^{(\downarrow a)})^\perp$, i.e. $\{p_3^{(\downarrow a)}\} \perp (P^{(\downarrow a)})^\perp$ (reasoning on the ACC induced graphs as before). Now, since by assumption $P^{(\downarrow a)}$ is a type i.e., $P^{(\downarrow a)} = (P^{(\downarrow a)})^{\perp\perp}$, we conclude that $\{p_3^{(\downarrow a)}\} \subseteq P^{(\downarrow a)}$, by case 1 of Property 3. $\square$

Proposition 30 (conservative single point restrictable pairs). *Assume $P = \{p_1, p_2\}$ is a conservative single point restrictable entangled pair that is, P is an entangled pair s.t. (see Definition 22):*

1. *it admits only one restriction by a single point a and*
2. *the restriction $P^{(\downarrow a)}$ is conservative.*

Then:

1. *the support X of P is given by the disjoint sum of next three segments*

$$S_0 : \{a, b, c\} \uplus S_1 : \{a_1, \dots, a_n\} \uplus S_2 : \{b_1, \dots, b_n\}$$

where S_1 and S_2 are defined as in the proof of Lemma 24, moreover

2. *P is obtained by merging the entangled pair $P_1 = \{p_1^1, p_2^1\}$ of form (1) together with the entangled pair $P_2 = \{p_1^2, p_2^2\}$ of form (2) that is,*

$P = \{p_1^1 \uplus p_1^2, p_2^1 \uplus p_2^2\}$ is in form (6) below:

$$P \left\{ \begin{array}{l} p_1 : \{(a, b) (c) (a_1, a_2) \dots (a_{i-1}, a_i) \dots (a_{n-1}, a_n) (b_1) (b_2) \dots (b_{i-1}) (b_i) \dots (b_{n-1}) (b_n)\} \\ p_2 : \{(a, c) (b) (b_1, b_2) \dots (b_{i-1}, b_i) \dots (b_{n-1}, b_n) (a_1) (a_2) \dots (a_{i-1}) (a_i) \dots (a_{n-1}) (a_n)\} \end{array} \right\} \quad (6)$$

where the restriction $P^{(\downarrow S_0)}$ is exactly that one in form (2).

Proof. Since the a -restriction is conservative a must occur both in one class of p_1 together with a point b and in one class of p_2 together with a point c of the support that is, $(a, b) \in p_1$ and $(a, c) \in p_2$ for some $b, c \in X$; moreover since P is only restrictable by the single point a , we have that:

1. $b \neq c$ and
2. b and c cannot occur together with other points of the support in some class of p_2 resp., p_1 otherwise P would be restrictable also in these two points, contradicting the assumption P is only restrictable by a .

This means that if $(a, b) \in p_1$ and $(a, c) \in p_2$ then $(b) \in p_2$ and $(c) \in p_1$. Now, since P is restrictable, p_1 and p_2 cannot consist of only these two pairs of classes that is, P cannot be reduced in form (1). Thus P must be in form (7)

$$P = \left\{ \begin{array}{l} p_1 = \{(a, b), (c)\} \uplus Q_1 \\ p_2 = \{(a, c), (b)\} \uplus Q_2 \end{array} \right\}. \quad (7)$$

where Q_1 and Q_2 are two sets of partitions over the support $X \setminus S_0$ with $S_0 = \{a, b, c\}$. Finally, since P is restrictable only by point a , Q_1 and Q_2 must be a non restrictable pair in form (2) over $S_1 \uplus S_2$; therefore P is an entangled pair in form (6) with support $S_0 : \{a, b, c\} \uplus S_1 : \{a_1, \dots, a_n\} \uplus S_2 : \{b_1, \dots, b_n\}$. $\square$

Lemma 31 (conservative single point restrictable entangled pairs).

Assume $P = \{p_1, p_2\}$ is a entangled pair that admits only one conservative restriction by point a , so P is in form (8) below with $b \neq c$:

$$P = \left\{ \begin{array}{l} p_1 : \{x_1 = (a, b), x_2, \dots, x_n\}, \\ p_2 : \{y_1 = (a, c), y_2, \dots, y_n\} \end{array} \right\}. \quad (8)$$

Let $p_3 \neq p_{i=1,2}$ be a partition in $P^{\perp\perp}$ then, p_3 must contain either class $x_1 = (a, b)$ or class $y_1 = (a, c)$ of P in form (8).

Proof. Assume P admits only one conservative restriction by point a then, by Proposition 30, P is as in form (6). Now, consider the next two partitions, q_1 and q_2 :

$q_1 : \{z_1 = (a), z_2 = (b, c)\} \in P_1^\perp$
 $q_2 : \{y_1 = (a_1, \dots, a_{i-1}, \dots, a_{n-1}, b_1, \dots, b_{i-1}, \dots, b_{n-1}), \dots, y_m = (a_n, b_n)\} \in P_2^\perp$
 where $P_1^\perp$ and $P_2^\perp$ are resp., the orthogonal of P_1 in form (1) and P_2 in form (2), by Proposition 30, as displayed on top and bottom side of Figure 11. Then build a partition q of the form $q_1 q_2(i, j)$, for some $1 \leq i, j \leq (2n + 3)$,

Figure 11: a conservative single point restrictable entangled pair

where the notation means that we concatenate q_1 and q_2 by mixing classes $z_h \in q_1$ and $y_k \in q_2$ containing, resp., points i and j . E.g., partition $q = \{z_1 \uplus y_1, z_2, y_2, \dots, y_m\}$, displayed in the bottom side of Figure 12, is obtained by concatenating q_1 and q_2 and mixing (by union) classes z_1 and y_1 . By Property 4, $q \in P^\perp$, since each induced graph $\mathcal{G}(p_i, q)$ for $i = 1, 2$ is ACC.

Now, assume by absurdum that a occurs in a class of $p_3 \neq p_{i=1,2}$ of $P^{\perp\perp}$ that is distinct from both classes (a, b) and (a, c) of (8). By Proposition 25 (case 2), a cannot occur in p_3 as singleton class (a) . So, assume a belongs to a class $(a, d, \dots)$ with $d \neq b$ and $d \neq c$. Then it is easy to find in $P^\perp$ a partition like q or q' , displayed resp., in the bottom and top sides of Figure 12, s.t. the induced graph $\mathcal{G}(p_3, q)$ resp., $\mathcal{G}(p_3, q')$, contains a cycle. Case $(a, b, c) \in p_3$ is also excluded because of partitions like q or q' , containing class $z_2 = (b, c)$. $\square$

Theorem 32 (entangled types). *Every entangled pair of partitions sets $P = \{p_1, p_2\}$, with domain $X = \{1, \dots, n\}$, is a type ($P = P^{\perp\perp}$) thus, entangled pairs are called entangled types.*

Figure 12: orthogonal of a conservative only single point restrictable entangled pair

Proof. By **induction on the dimension** of P , i.e. on the cardinality of its domain $X = \{1, \dots, n\}$.

The **base of induction** is when $n = 3$ (there is no entangled pair of dimension 2) then, by Fact 26, P is in form (1), so it is not restrictable; therefore it is a type, by Theorem 28.

For the **induction step**, assume P has dimension $n > 3$. If P is not restrictable then, by Theorem 28, P is a type. Otherwise, when P is restrictable, we split our reasoning in two main cases, depending on whether P admits or not a conservative restriction.

Case 1. Assume P admits a **non conservative restriction** by a point a , then P has the form below

$$P = \{p_1 : \{(a), x_2, \dots, x_n\}, p_2 : \{(a), y_2, \dots, y_n\}\}.$$

Assume by absurdum $p_3 \in P^{\perp\perp}$ with $p_3 \neq p_{i=1,2}$. By Proposition 25 (case 1), a must also occur in p_3 as singleton class, i.e. $p_3 = \{(a), z_2, \dots, z_n\}$ (otherwise, we get a cycle in the induced graph $\mathcal{G}(q_i^a, p_3)$ for some pumped $q_i^a \in P^\perp$). By hypothesis of induction, $P^{\downarrow a}$ is a type so, by Lemma 29, either $p_3 = p_1$ or $p_3 = p_2$.

Case 2. Assume P admits only **conservative restriction** by a point as e.g. that one by point a in form (9) below

$$P = \left\{ \begin{array}{l} p_1 : \{x_1 = (a, b), x_2, \dots, x_n\}, \\ p_2 : \{y_1 = (a, c), y_2, \dots, y_n\} \end{array} \right\}. \quad (9)$$

Assume by absurdum $p_3 \in P^{\perp\perp}$ with $p_3 \neq p_{i=1,2}$. By Proposition 25 (case 2), a cannot occur as singleton class (a) in p_3 (otherwise, we get a disconnection in the induced graph $\mathcal{G}(q_i^a, p_3)$ for some pumped $q_i^a \in P^\perp$), therefore the restriction of p_3 by a must be conservative (i.e., $p_3^{(\downarrow a)}$ and p_3 have the same weight). Now, by hypothesis of induction, the restriction $P^{(\downarrow a)}$ is a type thus, by Lemma 29, either $p_3^{(\downarrow a)} = p_1^{(\downarrow a)}$ or $p_3^{(\downarrow a)} = p_2^{(\downarrow a)}$; so, **a cannot occur in a class of p_3 containing more than three elements** (including a). Then, we split our reasoning in two sub-cases depending on whether b is or not equal c in P of form (9).

Case 2.1. If $b = c$ then form (9) of P becomes that below:

$$P = \left\{ \begin{array}{l} p_1 : \{x_1 = (a, b), x_2, \dots, x_n\}, \\ p_2 : \{y_1 = (a, b), y_2, \dots, y_n\} \end{array} \right\}.$$

By Proposition 25 (case 2) a cannot occur as singleton class (a) in p_3 . So, in case a does not also occur together with b in a class of p_3 then, by Lemma 29, b must occur alone as singleton class (b) in p_3 (by hypothesis of induction, $P^{(\downarrow a)}$ is a type). Now, by Proposition 25 (case 2) there exists at least a partition q_i in $P^\perp$ containing the singleton class (b), so $\mathcal{G}(p_3, q_i)$ contains a disconnected component (contradicting $p_3 \in P^{\perp\perp}$). Otherwise, in case a occurs together with b in a class of p_3 then, we get a contradiction, since, by Lemma 29, either $p_3 = p_1$ or $p_3 = p_2$ (by hypothesis of induction, $P^{(\downarrow a)}$ is a type).

Case 2.2. In case $b \neq c$ (w.r.t. P in form (9)), we distinguish two main sub-cases, depending on whether a belongs to a class of p_3 containing two or three elements of the domain.

Case 2.2.1. Assume **a occurs in a class of p_3 containing only two elements**; hence p_3 has the form below

$$p_3 = \{(a, d), z_2, \dots, z_n\};$$

then, $d \neq b$ and $d \neq c$ otherwise p_3 will be equal to p_1 or p_2 , by Lemma 29 (via the hypothesis of induction, $P^{(\downarrow a)}$ is a type). This implies that P must be restrictable by at least one more point than a otherwise, by Lemma 31, we get either $d = b$ or $d = c$. So, by Lemma 29, it does not matter whether d is or not an admissible restriction for P , it must be either $d = b$ or $d = c$, contradicting the assumption $p_3 \neq p_{i=1,2}$.

Case 2.2.2. Assume that **a occurs in a class of p_3 containing three elements**; hence p_3 has the form (10) below

$$p_3 = \{(a, d, e), z_2, \dots, z_n\}. \tag{10}$$

By Lemma 31, P must be restrictable by at least one more point than a ; therefore, w.r.t. all possible admissible restrictions of P , there are only three cases involving points d and e .

1. **Neither d nor e is an admissible restriction** of P . Assume P is restrictable by a point $f \neq a$. By hypothesis of induction the restriction $P^{(\downarrow f)}$ is a type, so by Lemma 29, either $p_3^{(\downarrow f)} = p_1^{(\downarrow f)}$ or $p_3^{(\downarrow f)} = p_2^{(\downarrow f)}$, that is absurdum, since $p_3^{(\downarrow f)}$ (p_3 has form (10)) contains a class (a, d, e) .
2. **Both d and e are admissible restrictions** of P . This means that, by Lemma 29 (via hypothesis of induction, $P^{(\downarrow a)}$ is a type and p_3 has form (10)) $d = b$ and $e = c$ (resp., $d = c$ and $e = b$); then, by playing with admissible restrictions, we get that p_1 (resp., p_2) will be inconsistent since it will contain by assumption class (a, b) (resp., class (a, c)) together with class (b, c) , this latter occurring as consequence, by Lemma 29, of the restriction of p_3 by point a , i.e.:

$$\begin{aligned} &\text{either } p_1 = \{(a, b), (b, c), x_3, \dots, x_n\} \in P \\ &\text{or } p_2 = \{(a, c), (b, c), y_3, \dots, y_n\} \in P. \end{aligned}$$

3. **Only one between d and e is an admissible restriction** of P . Assume d is an admissible restriction of P (in addition to that one by a). By reasoning on restrictions (since by Lemma 29, via hypothesis of induction, the restriction of p_3 in (10) by d is equal either to $p_1^{(\downarrow d)}$ or to $p_2^{(\downarrow d)}$), it must be either $e = b$ or $e = c$. Assume $e = c$ then, p_3 has form (11) below

$$p_3 = \{(a, d, c), z_2, \dots, z_n\}. \quad (11)$$

Since class (a, c) belongs to partition p_2 and since P is not restrictable by c (by hypothesis), then c must occur as singleton class (c) in p_1 ; so, form (9) of P will become like the one below:

$$P = \left\{ \begin{array}{l} p_1 : \{(a, b), (c), x_3, \dots, x_n\}, \\ p_2 : \{(a, c), y_2, y_3, \dots, y_n\} \end{array} \right\}.$$

By Lemma 29, via hypothesis of induction, the restriction of p_3 in form (11) by a must be equal either to $p_1^{(\downarrow a)}$ or to $p_2^{(\downarrow a)}$; this means, in particular, that class (d, c) of $p_3^{(\downarrow a)}$ must also occur either in p_1 or in p_2 : that is inconsistent since, in both cases, c will simultaneously occur in two classes both belonging either to p_1 or to p_2 , that is either $p_1 = \{(a, b), (c), (d, c), x_4, \dots, x_n\} \in P$ or $p_2 = \{(a, c), (d, c), y_3, \dots, y_n\} \in P$.

□

4.2. Entangled connectives

By Lemma 24, as soon as P is an entangled pair, then its orthogonal is nonempty, so it makes sense to define the class of *entangled connectives* as follows.

Definition 33 (entangled connectives). *A generalized connective (P, Q) , in graphical or sequential syntax, is called entangled whenever P or Q is an entangled pair of partitions sets.*

Next Fact 34 is consequence of case 6 of Property 3 and Theorem 32.

Fact 34. *If P is an entangled type then, (P, Q) is an entangled generalized connective iff $P = Q^\perp$.*

The **smallest** (w.r.t. the dimension of the domain) **entangled decomposable connective** is $P_X = \{\{(1, 3), (2)\}, \{(2, 3), (1)\}\}$ and $Q_{X^*} = \{\{(1, 2), (3)\}\}$, with $X = (1 \wp 2) \otimes 3$ and $X^* = (1 \otimes 2) \wp 3$ in graphical syntax.

If $P = \{p_1, p_2\}$ is an entangled pair over $X = \{1, \dots, n\}$ then, each singleton $\{p_i\}$ can be interpreted as the pre-type of a special decomposable formula with border $\{1, \dots, n\}$, called *bipole* [1]. A **bipole** B is a MLL formula with only two layers of connectives: a generalized $\wp$ of $(i-1) \otimes i$ sub-formulas (with $1 < i \leq n$) that is, $B = \wp(\hat{\otimes}_1, \dots, \hat{\otimes}_m)$ as in the l.h.s. picture of Figure 13. Dually, $B^\perp = \hat{\otimes}(\hat{\wp}_1, \dots, \hat{\wp}_n)$ is called *anti-bipole*, that is a MLL formula with only two layers of connectives: a generalized $\otimes$ of $(i-1) \wp i$ sub-formulas (with $1 < i \leq n$) as in the r.h.s. picture of Figure 13. Every bipole satisfies the property that its pre-type is a type (see Lemma 4 of [9]). To be precise, observe that in case an entangled pair P is interpreted as the (pre-)type of a bipole B then some $\otimes$ -subtrees of B may be unary.

Figure 13: a *bipole* skeleton on l.h.s. and an *anti-bipole* skeleton on r.h.s.

In the following we characterize, by means of Theorem 36 and Theorem 39, the *class of non binary decomposable entangled connectives*. We then show that these connectives can neither be approximated to decomposable connectives (Corollary 40) nor sequentialized in the MLL^+ sequent calculus (Section 5).

4.3. Decomposable normal form of entangled connectives

Definition 35 (decomposable normal form). Let $P = \{p_1, p_2\}$ be an entangled pair of partitions then: P is decomposable iff it is the pre-type of the parse tree of a multiplicative formula; moreover, P is in (decomposable) normal form iff P is the pre-type of a formula tree F with literals indexes as in the l.h.s picture of Figure 14 that is, P is obtained by a $\wp$ between:

- the smallest decomposable entangled connective, $\{\{(1, 3), (2)\}, \{(2, 3), (1)\}\}$, enclosed by dashed line, and
- the possibly empty bipole with possibly unary classes, $\{\{(4, 5), \dots, (n-1, n)\}\}$, enclosed by dotted line;

hence P in the following normal form (12).

$$P : \left\{ \begin{array}{l} p_1 = \{x_1 : (1, 3), x_2 : (2), x_3 : (4, 5), \dots, x_{m \geq 2} : (n-1, n)\}, \\ p_2 = \{y_1 : (2, 3), y_2 : (1), y_3 : (4, 5), \dots, y_{m \geq 2} : (n-1, n)\} \end{array} \right\}. \quad (12)$$

Figure 14: normal form of decomposable entangled connectives

E.g., $P = \{\{(a, b), (c, e), (d)\}, \{(a, b), (d, e), (c)\}\}$ is an entangled pair in decomposable normal form since $P = \mathcal{P}_F$ with $F = (a \otimes b) \wp ((c \wp d) \otimes e)$ while the entangled pair $Q = \{\{(a, c), (b), (d, e), (f)\}, \{(b, c), (a), (d, f), (e)\}\}$ is not in normal form (it is not even decomposable). Observe that normal form is defined up to commutativity of MLL.

Theorem 36 (decomposable normal form). An entangled pair of partitions $P = \{p_1, p_2\}$ is decomposable iff P is in normal form (12).

Proof. Assume P is a decomposable entangled pair of partitions, hence by Definition 13, there exists a formula F s.t. pre-type $\mathcal{P}_F$ of its syntactical tree is P . Since P is entangled (its size is 2 and its weight is ≥ 2), the

syntactical tree of F must contain at least a $\wp$ -node otherwise (by reasoning on switchings of F) the pre-type of F consists of only one partition with a single class (contradicting the assumption P is entangled); for similar reason F cannot consist of only $\wp$ -nodes otherwise the (pre-)type of F contains exactly one partition (contradicting the assumption P is entangled). So, F must contain at least a $\otimes$ -node and a $\wp$ -node. Now it must exist at least a $\wp$ -node L_1 above a $\otimes$ -node L_2 in the tree otherwise F is a bipole and therefore its (pre-)type would consist of a single partition, contradicting the assumption P is entangled pair. So, assume F is as in the r.h.s. picture of Figure 14, where t_1, t_2, t_3 and t_4 are sub-trees of the formula tree of F .

First observe that no other link can stay between L_1 and L_2 , otherwise such a link, let us say, $L' = \wp$ (resp., let us say $L' = \otimes$) would increase the size s of P (resp., the maximal degree $1 \leq \delta \leq 2$ allowed for each class of p_i) contradicting the assumption P is an entangled pair.

Then observe that every tree t_i , for $i = 1, 2, 3$, must be reduced to single points (i.e., each t_i is a tree consisting of a single node), otherwise:

1. in case t_i is a tree whose root is a $\otimes$ -node, we get a class with degree > 2 , contradicting the assumption P is entangled pair;
2. in case t_i is a tree whose root is a $\wp$ -node, by reasoning on the switchings of F (since there would be two $\wp$ -nodes dominating a $\otimes$ -node), we get the size of P is > 2 , contradicting the assumption P is entangled pair.

So, t_1, t_2 and 3 must be single-point trees; moreover t_4 (if any) must be a bipole like that one enclosed by dotted line in the l.h.s. of Fig. 14, otherwise by computing of all possible switchings of F we get that its pre-type $\mathcal{P}_F = P$ is not an entangled pair, that is contradiction.

Thus, F must be as in the l.h.s. picture of Figure 14, with the pre-type consisting of partitions p_1 and p_2 as in normal form (12).

Vice-versa, it is easy to calculate that an entangled pair in normal form (12) is decomposable. $\square$

Remarks 37 (normal form and semi-distributive law). *Informally we can say that an entangled pair P in decomposable normal form can be thought as the set of the pre-types (types indeed) of the two bipoles (reductum), B_1 and B_2 , that are obtained after one step of weak (or linear or semi-) distributive law [2] applied to the (redex) formula F (where $P = \mathcal{P}_F$) whose abstract skeleton tree is the one depicted on the l.h.s. of Figure 14. Actually, following [9], we may define a rewrite relation “ $\sim_{wd}$ ” on MLL formulas trees*

generated by associativity and commutativity of $\otimes$ and $\wp$ plus the **weak or semi-distributive law** synthetically defined as follows:

$$F: \begin{array}{c} A \quad B \\ \wp \\ \otimes \\ C \end{array} \rightsquigarrow_{wd} \left\{ \begin{array}{c} A \quad C \\ \otimes \\ \wp \\ B \end{array} F_1: \quad \wedge \quad \begin{array}{c} B \quad C \\ \otimes \\ \wp \\ A \end{array} F_2: \right\}$$

$$(A \wp B) \otimes C \vdash (A \otimes C) \wp B \quad \wedge \quad (A \wp B) \otimes C \vdash (B \otimes C) \wp A.$$

Then it is easy to calculate the equality $P = \mathcal{P}_F = \mathcal{P}_{B_1} \cup \mathcal{P}_{B_2}$ once F is as in the l.h.s. of Figure 14 and $F \rightsquigarrow_{wd} B_i$, for $i = 1, 2$, as below:

$$F: \begin{array}{c} 1 \quad 2 \\ \wp \\ \otimes \\ 3 \quad 4 \\ \wp \\ 5 \quad n-1 \\ \wp \\ n \end{array} \rightsquigarrow_{wd} \left\{ \begin{array}{c} 1 \quad 3 \\ \otimes \\ \wp \\ 2 \quad 4 \\ \wp \\ 5 \quad n-1 \\ \wp \\ n \end{array} B_1: \quad \wedge \quad \begin{array}{c} 2 \quad 3 \\ \otimes \\ \wp \\ 1 \quad 4 \\ \wp \\ 5 \quad n-1 \\ \wp \\ n \end{array} B_2: \right\}$$

Observe that, via Theorem 38, stated below and proved in [9], we can show that “**if P is a decomposable entangled pair then, P is a type**”. Actually, if P is a decomposable entangled pair then, by Theorem 36, P is the pre-type of the formula tree F in the l.h.s. of Figure 14 and so, by Theorem 38, P is a type. In other words, if P is a decomposable set of partitions then, by Theorem 38, both types $\mathcal{P}_F^{\perp\perp}$ and $\mathcal{P}_F^\perp$ (since, by Fact 14, $\mathcal{P}_F^\perp = \mathcal{T}_{F^\perp}$) can be computed by the exhaustive $\rightsquigarrow_{wd}^*$ -rewriting of F and $F^\perp$.

Theorem 38 (Maieli-Puite: type generation of a MLL formula tree). If F is a MLL formula with pre-type $\mathcal{P}_F$ then, its type $\mathcal{T}_F$ can be generated by the exhaustive $\rightsquigarrow_{wd}$ -rewriting of F , as follows:

$$\mathcal{T}_F = \bigcup_i \mathcal{P}_{F_i} \text{ s.t., } F \rightsquigarrow_{wd}^* F_i$$

where each F_i is a bipole, with type $\mathcal{P}_{F_i}$, obtained by transitive closure of $\rightsquigarrow_{wd}^*$ relation from F .

Proof. Given in [9]. □

4.4. Undecomposable entangled connectives

By Theorem 36, an entangled connective, P and Q , is binary decomposable iff P or Q is in normal form. We want now to make sure that if P (resp., Q) is a non decomposable entangled type, then neither Q (resp., P) will be decomposable (by Theorem 39), so P and Q is an undecomposable connective.

Theorem 39 (undecomposable entangled connectives). *Let E and Q be a generalized connective. If E is an entangled pair that is not binary decomposable then, Q is neither decomposable.*

Proof. Let (E, Q) be a generalized connective with E an undecomposable entangled pair. Suppose, for the sake of absurdity, that Q is decomposable, i.e. $Q = \mathcal{P}_A$ for some formula A . By Fact 34 and by Fact 14, we have

$$E =_{(\text{Fact 34})} Q^\perp = \mathcal{P}_A^\perp =_{(\text{Fact 14})} \mathcal{T}_{A^\perp}$$

that is, E is the bi-orthogonal of $\mathcal{P}_{A^\perp}$. Therefore, $\mathcal{P}_{A^\perp}$ is included in E , by case 3 of Property 3. Now, since E has only two elements, if $\mathcal{P}_{A^\perp}$ were strictly included in E , it would be either the empty set or a singleton set so, its bi-orthogonal would not be E ; therefore we have $E = \mathcal{P}_{A^\perp}$, a contradiction. $\square$

Next corollary is consequence of Fact 34 and Theorem 39.

Corollary 40 (completion). *Let E and Q be an entangled connective with E being a non (binary) decomposable entangled pair; then, E cannot be completed in such a way to become decomposable (i.e., $\neg \exists D \supsetneq E$ s.t. D is binary decomposable with D and Q being a connective).*

Remarks 41 (entangled connectives and bipoles). *Although our notion of “entanglement” doesn’t solve the (admittedly, difficult) general problem (“try to find a characterization of the full class of undecomposable connectives”), it is far from being an “ad hoc” condition. It is rather a natural condition that can be observed as soon as we “superpose” (sum) pairs of bipoles with the same “skeleton” (i.e., bipoles with the same abstract syntactical tree), like e.g. the two ones, F_1 and F_2 , enclosed by dotted lines in Figure 9 of Section 3.3. More concretely, an entangled connective, $P = \{p_1, p_2\}$ and $P^\perp$, can be interpreted as the union (resp., the intersection) of the types of two bipoles (resp., the types of two anti-bipoles) which are equivalent up to cyclic permutations of the literals indexes of the top border that is, bipoles (resp., anti-bipoles) s.t.:*

1. *they have the same syntactical tree skeleton;*

2. they have the same border up to cyclic permutation of literals indexes.

Formally, given two bipoles like F_1 and F_2 of Figure 9, with $\mathcal{T}_{F_1} = \{p_1\}$ and $\mathcal{T}_{F_2} = \{p_2\}$, then $P = \mathcal{T}_{F_1} \cup \mathcal{T}_{F_2}$ and $P^\perp = \mathcal{T}_{F_1^\perp} \cap \mathcal{T}_{F_2^\perp}$ is an entangled connective that is undecomposable, by Definition 35 and Theorem 36. E.g., consider the Girard's undecomposable connective G_4 and $G_4^\perp$ which is an entangled connective (by Definition 33): G_4 (resp., $G_4^\perp$) results by the union (resp., by the intersection) of the types of bipoles $F_1 = (1 \otimes 2) \wp (3 \otimes 4)$ and $F_2 = (2 \otimes 3) \wp (4 \otimes 1)^7$ (resp., of the types of anti-bipoles $F_1^\perp$ and $F_2^\perp$) of Figure 9 i.e., $G_4 = \mathcal{T}_{F_1} \cup \mathcal{T}_{F_2}$, (resp., $G_4^\perp = \mathcal{T}_{F_1^\perp} \cap \mathcal{T}_{F_2^\perp}$). This fact (Theorem 32) is a novelty since the union of types is not in general a type while the intersection of types is always a type (Property 3). Indeed, entangled types are the smallest types (w.r.t. the number of partitions), if we exclude the trivial singleton types (every set consisting of a single partition is a type by case 11 of Property 3). So, entangled connectives can be considered in some sense the “smallest” generalized multiplicative connectives (w.r.t. the number of switchings or the number of rules), if we exclude the basic ones, $\wp$ and $\otimes$, and bipoles.

5. Sequentialization of undecomposable connectives

The natural correspondence (sequentialization) between sequential decomposable connectives and graphical decomposable connectives is broken by non decomposable connectives! There exist proof nets, containing non decomposable links, without counterpart in the sequential calculus MLL^+ , if we exclude the trivial correspondence with (non atomic) logical axioms. Let G and G^* be an entangled non decomposable connective. By Theorem 39, neither G nor G^* is binary decomposable. There is indeed no η -expanded proof of $\vdash G, G^*$ since each rule for G , resp., G^* has at least two premises (see Section 3.4). Actually, sequents of non decomposable entangled formulas $\vdash G, G^*$, are not provable from atomic logical axioms in the extended MLL sequent calculus: that is because each rule for G (resp., G^*) is at least binary (it has at least two premises, by the entanglement) so any derivation of this sequent would build a premise with the other conclusion G^* (resp., G) together with only some (not all) of the principal formulas of the applied rule. This

⁷Notice that the respective literals indexes of the top borders of F_1 and F_2 are cyclic permutations of the linear sequence $1 < 2 < 3 < 4$.

situation is quite different with proof nets. Consider the proof structure π with only two conclusions, G and G^* , as in Figure 15, built as follows:

1. take the two non decomposable links λ_G and λ_{G^*} ;
2. label the elements of the top border $X = \{1, \dots, n\}$ of λ_G (resp., of λ_{G^*}) by n occurrences of literals, $a_1, \dots, a_n$ (resp., $a_1^\perp, \dots, a_n^\perp$);
3. put an axiom link for each matching pair of dual literals, a_i and $a_i^\perp$.

Clearly π is an η -expanded proof net: each global switching induces an acyclic and connected correction graph; nevertheless, π cannot be “sequentialized” in MLL^+ , if we exclude the trivial (non atomic) axiom $\vdash G, G^*$. Actually, any attempt of starting by sequentializing G (resp., G^*) will induce a single connected proof structure π_{G^*} (resp., π_G), with G^* (resp., G) among its conclusions, which is not correct: actually, there exist two switchings for π_{G^*} (resp., π_G) with at least two disconnected components (consequence of the entanglement conditions of Definition 21). Moreover, since neither G nor G^* is binary definable (by Corollary 40, neither G nor G^* can be completed to a decomposable connective), π is not even stepwise sequentializable. This

Figure 15: non sequentializable proof net with non decomposable conclusions G, G^*

fact witnesses an asymmetry between proof nets and sequent proofs since the former ones allow us to express a kind of parallelism of proofs that the latter ones cannot do.

6. Conclusions and future works

We gave the first characterization (via Theorems 32, 36 and 39) of a class of multiplicative undecomposable connectives: namely, the entangled connectives (Definition 33) that are not in (decomposable) normal form (Definition 35).

Anyway, there exist non decomposable connectives besides the entangled ones. We are currently working on a more general characterization of the full class of primitive non decomposable connectives (i.e., those ones non definable by means of other connectives). Naively our idea is that a non decomposable connective is rather defined by the set of partitions over the cyclic permutations of the linearly ordered support $1 < \dots < n$; that's the case with e.g. G_9 below (we omit its orthogonal $G_9^\perp$ for the sake of simplicity):

$$G_9 = \{ \begin{array}{l} \{(1, 2, 3), (4, 5, 6), (7, 8, 9)\}, \\ \{(2, 3, 4), (5, 6, 7), (8, 9, 1)\}, \\ \{(3, 4, 5), (6, 7, 8), (9, 1, 2)\} \end{array} \}.$$

Concerning larger (than MLL) fragments of linear logic, like e.g. MELL or MALL, we don't know at this moment about non definable connectives for such fragments. Certainly, if we restrict to consider the pure additive fragment (ALL) then, there does not exist any generalized connective that cannot be decomposed by the basic additive ones, $\&$ and $\oplus$. There is no “packaging problem” in ALL since contexts in which we derive generalized formulas are simply duplicated. Roughly speaking, non decomposability concerns rather the multiplicative partition of contexts than the additive superposition (better, *slicing*) of contexts.

The fact that one can compute by means of cut elimination, using such non decomposable connectives, is certainly a good starting point: nevertheless, the study of their connection with *concurrency* (typically, the Pi-Calculus [10]) rather than the *Curry-Howard correspondence* should be further developed⁸.

Finally, it would be also useful to have a *coherence semantics* for such non decomposable connectives. We could e.g. use the so-called *experiment method* of Girard [5] for the analysis of proof structures built on such non decomposable connectives. Examples of what can be done with these kind of semantic techniques can be found in [11] and more recently in [4].

⁸Naively, since, by the Curry-Howard isomorphism, “proofs are supposed to correspond to programs”, *what should be the program $[\pi]$ corresponding to the non decomposable proof net π of G, G^* (we met in Section 5)?* if such a program does exist, is there any correspondence between the non-sequentializability of π and the possibly *concurrent nature* of such a program π^* ? *Have we a chance to write “a sequential version” of such a “concurrent program” $[\pi]$?*

References

- [1] J.-M. Andreoli. Focussing and Proof Construction. *APAL* 107(1), pp. 131-163, 2001.
- [2] Cockett, J.R.B. and Seely, R.A.G.: Weakly Distributive Categories. *Journal of Pure and Applied Algebra*, vol. 114, pp. 133-173, 1997.
- [3] Danos, V. and Regnier, L.: The structure of multiplicatives. *AML*, 28, pp.181-203, 1989.
- [4] Ehrhard, T.: A new correctness criterion for MLL proof nets. *Proc. of the Twenty-Third Conf. CSL-LICS'14*. Vienna, July 14-18, 2014.
- [5] Girard, J.-Y.: Multiplicatives. *Rendiconti del Seminario Matematico dell'Univ. di Torino*, Special Issue on Logic and Computer Science, pp. 11-33, 1987.
- [6] Girard, J.-Y.: Linear Logic, *Theor. Comp. Sci.*, London Math. vol. 50(1), pp. 1-102, 1987.
- [7] J.-Y. Girard, On the meaning of logical rules I: syntax vs. semantics. *Computational Logic*, pp. 215-272, SV, Heidelberg, 1999.
- [8] Girard, J.-Y.: Le point aveugle. In: Hermann (ed.), *Vers la Perfection*, Paris, vol. I, 2006
- [9] Maieli, R. and Puite, Q.: Modularity of proof nets: generating the type of a module. *Archive for Math. Logic*, vol. 44(2), pp. 167-193, 2005.
- [10] Milner, R.: *Communicating and Mobile Systems: the Pi-Calculus*, Cambridge University Press, 1999.
- [11] Retoré, C.: A Semantic Characterisation of the Correctness of a Proof Net. *Math. Struct. in Comp. Sci.*, vol. 7(5), pp. 445-452, 1997.