MicroRNAs related to cholesterol metabolism affected by vegetable diet in rainbow trout (*Oncorhynchus mykiss*) from control and selected lines

Tengfei Zhu, Geneviève Corraze, Elisabeth Plagnes-Juan, Jérôme Montfort, Julien Bobe, Edwige Quillet, Mathilde Dupont-Nivet, Sandrine Skiba-Cassy

► To cite this version:

Tengfei Zhu, Geneviève Corraze, Elisabeth Plagnes-Juan, Jérôme Montfort, Julien Bobe, et al.. MicroRNAs related to cholesterol metabolism affected by vegetable diet in rainbow trout (*Oncorhynchus mykiss*) from control and selected lines. *Aquaculture*, 2019, 498, pp.132-142. 10.1016/j.aquaculture.2018.08.058 . hal-01870570

HAL Id: hal-01870570

<https://hal.science/hal-01870570>

Submitted on 26 Nov 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution| 4.0 International License

MicroRNAs related to cholesterol metabolism affected by vegetable diet in rainbow trout (*Oncorhynchus mykiss*) from control and selected lines

Tengfei Zhu^a, Geneviève Corraze^a, Elisabeth Plagnes-Juan^a, Jérôme Montfort^b, Julien Bobe^b, Edwige Quillet^c, Mathilde Dupont-Nivet^c, Sandrine Skiba-Cassy^{a,*}

^a INRA, Univ Pau & Pays Adour, E2S UPPA, UMR 1419, Nutrition Métabolisme Aquaculture, Saint Pée sur Nivelle F-64310, France

^b UMR1313 GABI, INRA, AgroParisTech, Université Paris-Saclay, F-78350 Jouy-en-Josas, France

^c UR1037 LPGP, INRA, F-35000 Rennes, France

For the sustainable development of aquaculture, vegetable ingredients have been used to replace the traditional ingredients fishmeal and fish oil in aquafeeds. On the other hand, selective breeding has also been implemented at research level to obtain strains that are adapted to feeding on the plant-based diet. To better understand the underlying mechanisms prompting the adaptation to plant-based diets in fish, we investigated the hepatic expression of several microRNAs (miRNAs) that are involved in the post-transcriptional regulation of cholesterol and lipid metabolism at 8 h and 16 h after the last meal in two lines of rainbow trout: one selected for better adaptation to plant-based diets and the corresponding control line. Both groups were fed either a fishmeal and fish oil based diet or a 100% plant-based diet. Results showed that the expression of miR-33a in liver was greatly elevated in trout fed the plant-based diet, while the expression of miR-122 and miR-128 was much higher in the selected line at 8 h after the last meal regardless of the diet. Furthermore, our results indicated that some genes involved in immune processes (caspase 6 apoptosis-related cysteine peptidase like 2, *casp6l2*) and cAMP signal transduction (phosphodiesterase 4B cAMP-specific a, *pde4ba*) were also potentially regulated by miRNAs. They were newly identified as putative direct targets of miRNAs and affected in trout fed the plant-based diet. Though further investigations are still needed to establish a valid relationship between miRNAs and their target genes, our study found miR-33a, miR-122 and miR-128 as potential candidates for further study and provided new perspectives to understand the role of miRNAs in the selective breeding for adaptation to the plant-based diets.

1. Introduction

Increasing demand for world fisheries production has resulted in a great expansion of aquaculture over the last decades, accounting for about half of the world global seafood production (FAO, 2016). As capture fisheries are limited and showed no increase since the mid-1980, the increasing demand for fish in the future will be inevitably supplied by aquaculture. However, the production of fishmeal and fish oil coming from wild fishery stocks and constituting important traditional ingredients in aquafeed is also going to keep stable or even decrease (Jackson, 2012). Therefore, the utilization of plant ingredients to replace fishmeal and fish oil has been considered as the best solution to support the sustainable development of aquaculture in recent years (Tacon et al., 2010) and evident progress has been made regarding the substitution of fishmeal and fish oil by plant raw materials in salmonids and marine fish (Benedito-Palos et al., 2008; Torstensen et al., 2008).

However, studies have indicated that total replacement of fishmeal and fish oil by the plant-based diets still reduced growth performances and flesh quality (Geay et al., 2011; Panserat et al., 2009). In order to further improve the use of plant food in carnivorous fish like salmonids, research efforts have been more recently focused on the selective breeding. Indeed, the selective breeding could be the alternative solution to enhance fish adaptation to the plant-based diets and has been well confirmed at a research level. For example, it has been demonstrated that survival rate and mean body weight were improved in the first generation of rainbow trout selected for the ability to adapt to a totally plant-based diet (Le Boucher et al., 2012). Moreover, in Atlantic salmon, the deposition and/or retention of dietary n-3 long-chain polyunsaturated fatty acids (LC-PUFA) in flesh, such as eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA), is a highly heritable trait (Leaver et al., 2011) and Nofima has also carried out profitable work in this respect (Horn et al., 2017), prompting further interest in

* Corresponding author.

E-mail address: sandrine.skiba@inra.fr (S. Skiba-Cassy).

exploring genotype-nutrient interactions.

The replacement of fishmeal and fish oil by plant ingredients caused great variations in nutrient supply, including the reduction of cholesterol and n-3 LC-PUFA. Cholesterol is derived from the animal organisms, so it is absent in plant ingredients (Tocher et al., 2008). In animals, cholesterol represents an important component of the membrane to modulate the fluidity and permeability and it is also the precursor of many biologically active compounds, including bile acids, steroid hormones and vitamin D (Goedeke and Fernández-Hernando, 2012).

To maintain cholesterol homeostasis, the expression of genes related to cholesterol metabolism is tightly controlled at both transcriptional and post-transcriptional level. The sterol regulatory element-binding proteins (SREBPs) and the liver X receptors (LXRs) are known to be well involved in the transcriptional regulation of cholesterol metabolism by targeting genes involved in the cholesterol synthesis (*hmgcr*, 3-hydroxy-3-methylglutaryl-CoA, reductase) and efflux (*abca1*, ATP-binding-cassette transporter-A1, *abcg5* and *abcg8*, ATP-binding-cassette transporter G5 and G8 and *cyp7a1*, cytochrome P450 cholesterol 7 α -hydroxylase), respectively (Sakakura et al., 2001; Zhao and Dahlman-Wright, 2010). At the post-transcriptional level, a class of small non-protein coding RNAs of 19–24 nucleotides, designated as microRNA (miRNA), has been found to regulate gene expression either by mRNA decay or translational repression (Iwakawa and Tomari, 2015). Among the wide numbers of miRNAs, miR-1 (Zhong et al., 2013), miR-33a (Gerin et al., 2010; Najafi-Shoushtari et al., 2010), miR-122 (Esau et al., 2006), miR-128 (Adlakha et al., 2013) and miR-223 (Vickers et al., 2014; Wang et al., 2013) were reported to be involved in the regulation of cholesterol and lipid metabolism in mammals.

The ability to grow on the plant-based diet has been proved to be genetically variable in fish (Le Boucher et al., 2011; Quinton et al., 2007). Although the growth potential has been effectively enhanced by selective breeding, the molecular mechanisms underlying this selection still remain unknown. In the present study, we used two lines of rainbow trout, the selected line which had better adaptation to the plant-based diet and the non-selected control line in order to investigate the potential role of miRNA, such as miR-1, miR-33a, miR-122, miR-128 and miR-223, in the adaptation to the plant-based diet. Therefore, the hepatic expression of these miRNAs and the genes related to cholesterol metabolism were analyzed. Furthermore, as miR-33a, miR-128 and miR-223 were found to be affected by the plant-based diet or selective breeding, we expanded our study to some genes that were identified in silico as potential direct targets of these miRNAs based on the complementarity between the seed sequence of the miRNAs and the 3' untranslated region (UTR) region of the rainbow trout transcripts.

2. Materials and methods

2.1. Ethics statement

Experimentation was conducted in the INRA experimental facilities (Peima facilities, Sizun, France and UMR Numéa, St-Pée-sur-Nivelle, France) authorized for animal experimentation by the French veterinary service which is the competent authority (B 29–277-02 and A 64–495-1). The experiments were in strict accordance with EU legal frameworks related to the protection of animals used for scientific research (Directive 2010/63/EU) and according to the National Guidelines for Animal Care of the French Ministry of Research (decree n°2013–118, February 1st, 2013). The scientists in charge of the experimentation received training and personal authorization (N° B64 10,003 and A29 102). In agreement with ethical committees “Comité d’Ethique Aquitaine Poissons Oiseaux” (C2EA-73) and “Comité d’Ethique Finistérien en Expérimentation Animale” (C2EA-74), the experiment reported here does not need approval by a specific ethical committee since it implies only classical rearing practices with all diets used in the experiment formulated to cover the nutritional requirements of rainbow trout (National Research Council, 2011). Fish were

monitored daily during the experiment. If any clinical symptoms (i.e. morphological abnormality, restlessness or uncoordinated movements) were observed, fish were sedated by immersion in 2% benzocaine solution and then euthanized by immersion in a 6% benzocaine solution (anesthetic overdose) during 3 min.

2.2. Fish, diet and experimental design

The experiment was conducted on two groups of rainbow trout: one line (S) which has been selected for better ability to grow and survive with a totally plant-based diet for three generations (Callet et al., 2017; Le Boucher et al., 2012) and a control line (C) maintained with high effective size and without artificial selection. 3600 eggs from S line and 3600 eggs from C line were obtained by within line mating (18 dams and 31 sires for each line). At 19 days post fertilization (dpf), the eyed eggs were randomly distributed into 12 tanks (0.25 m³) with 600 eyed eggs in each tank. The fish were reared in the 12 flow-through tanks (three tanks per treatment) in INRA experimental facilities (PEIMA, Sizun, France) at 11 °C and under artificial photoperiod condition (from 8 am to 8 pm). From the first-feeding stage, S and C fish were fed ad libitum with either a plant-based diet totally devoid of marine products (V diet) or a diet containing fishmeal and fish oil (M diet) during six months. Diets were manufactured in our experimental facilities of Donzacq (France) using a twin-screw extruder (Cletral, France). The M diet contained fishmeal and fish oil as main protein and lipid source, respectively. The V diet was formulated with a blend of vegetable oils (palm, rapeseed and linseed oils) and a blend of plant protein sources (corn and wheat gluten, soybean meal, soy protein concentrate, light white lupin, dehulled pea and extruded whole wheat). Diets were formulated to fulfill the requirements of rainbow trout according to NRC recommendations (National Research Council, 2011). Synthetic L-lysine, L-arginine, dicalcium-phosphate and soy-lecithin were added to the V diet to correct the deficiency in essential amino acids, phosphorous and phospholipid supply. Diet formulations and compositions are described in Table 1.

2.3. Sampling procedure

At the end of the feeding trial, fish were anesthetized with benzocaine (30 mg/L) and killed by a sharp blow to the head. Six fish were sampled from each tank at 8 h and 16 h after the last meal, respectively. Blood was removed from the caudal vein into EDTA syringes and centrifuged (1000 g, 10–15 min); then recovered plasma was immediately frozen in liquid nitrogen and kept at –80 °C for the measurement of cholesterol and triglycerides. Liver was dissected and weighed for hepato-somatic index determination and then immediately frozen in liquid nitrogen and kept at –80 °C. About 100 mg was cut from the whole liver for RNA extraction and then subsequent gene and miRNA expression analyses.

2.4. Diet and whole body composition analysis

Proximate analysis of the experimental diets was determined according to Association of Official Analytical Chemists (AOAC, 2000) as follows: dry matter was analyzed by drying the samples to constant weight at 105 °C for 24 h. Crude protein was determined using the Kjeldahl method after acid digestion and estimated by multiplying nitrogen by 6.25. Crude lipid was quantified by petroleum diethyl ether extraction using Soxhlet method. Gross energy content was determined in an adiabatic bomb calorimeter (IKA). Ash was examined by combustion in a muffle furnace at 550 °C for 16 h. Dietary sterols were assayed on a total lipid extract by means of the Liebermann-Burchard method (Stadtman, 1957).

Table 1
Ingredients and proximate composition of the diets.

Ingredients (%)	M	V
Fish meal	58.42	0.00
Corn gluten	0.00	16.03
Wheat gluten	0.00	17.72
Soybean meal	0.00	10.13
Soy protein concentrate	0.00	15.19
Light white lupin	0.00	6.75
Dehulled pea	0.00	4.30
Extruded whole wheat,	25.26	3.38
Fish oil	14.09	0.00
Vegetable oils ^a	0.00	15.61
Mineral premix ^b	1.12	1.18
Vitamin premix ^c	1.12	1.18
Soy lecithin	0.00	2.11
L-lysine	0.00	1.52
L-methionine	0.00	0.34
CaHPO ₄ ·2H ₂ O	0.00	3.21
Attractant Mix	0.00	1.35
Analytical composition		
Dry Matter (DM, %)	90.13	94.41
Crude protein (% DM)	46.98	50.01
Lipid (% DM)	18.80	19.49
Sterols (% DM)	0.77	0.35
Energy (kJ/g DM)	23.24	23.84
Ash (% DM)	8.17	6.25

^a Vegetable oils: palm oil (30%), rapeseed oil (55%), linseed oil (15%).

^b Mineral premix (g or mg/kg diet): calcium carbonate (40% Ca), 2.15 g; magnesium oxide (60% Mg), 1.24 g; ferric citrate, 0.2 g; potassium iodide (75% I), 0.4 mg; zinc sulfate (36% Zn), 0.4 g; copper sulfate (25% Cu), 0.3 g; manganese sulfate (33% Mn), 0.3 g; dibasic calcium phosphate (20% Ca, 18% P), 5 g; cobalt sulfate, 2 mg; sodium selenite (30% Se), 3 mg; KCl, 0.9 g; and NaCl, 0.4 g (UPAE, INRA).

^c Vitamin premix (IU or mg/kg diet): DL- α -tocopherol acetate, 60 IU; sodium menadione bisulphate, 5 mg; retinyl acetate, 15,000 IU; DL-cholecalciferol, 3000 IU; thiamin, 15 mg; riboflavin, 30 mg; pyridoxine, 15 mg; B12, 0.05 mg; nicotinic acid, 175 mg; folic acid, 500 mg; inositol, 1000 mg; biotin, 2.5 mg; calcium pantothenate, 50 mg; and choline chloride, 2000 mg (UPAE, INRA).

2.5. Analysis of plasma metabolites

Plasma metabolite concentration was measured using commercial kits, triglycerides (PAP 150, bioMérieux, France) and cholesterol (CHOD-PAP, Sobioda, France), adapted to microplate format, according to the recommendations of the manufacturer.

2.6. In silico identification of miRNA target gene

Putative miRNA targets were predicted using the miRanda algorithm (www.ncbi.nlm.nih.gov/pubmed/15502875) applied to studied mature miRNA sequences and rainbow trout 3'UTR regions. Rainbow trout 3'UTR regions were predicted using a set of 78,415 contigs originating from de novo assembled RNA-seq data publicly available in the PhyloFish online database (www.ncbi.nlm.nih.gov/pubmed/27189481). In order to focus on the 3'UTR of the contigs, zebrafish reference proteins obtained from ensembl (www.ensembl.org) were blasted onto the rainbow trout PhyloFish contigs. The region downstream of the alignment region on each contig was subsequently kept for further analysis using miRanda with default parameters.

2.7. Gene expression analysis

Quantitative RT-PCR gene expression analyses were performed on liver samples ($n = 6$) collected 8 h and 16 h after the last meal. Genes analyzed in the present study were: HMG-CoA reductase (*hmgcr*), HMG-CoA synthase (*hmgcs*), lanosterol 14 α -demethylase (*cyp51*) and 7-dehydrocholesterol reductase (*dhcr7*) for cholesterol biosynthesis, ATP-binding cassette transporter G5 (*abcg5*) and ATP-binding cassette

transporter G8 (*abcg8*) for cholesterol excretion, cholesterol 7 α -hydroxylase (*cyp7a1*) for bile acid synthesis, ATP-binding cassette transporter A1 (*abca1*) for plasma cholesterol efflux, sterol regulatory element-binding protein 2 (*srebp-2*) and liver X receptor α (*lxra*) as transcriptional factors, acyl-CoA cholesterol acyltransferase 1 (*acat1*) and acyl-CoA cholesterol acyltransferase 2 (*acat2*) for cholesterol esterification, fatty acid synthase (*fas*) and the transcription factor sterol regulatory element-binding protein 1c (*srebp-1c*) for lipogenesis and glucokinase (*gck*) for glycolysis although known as the first step of lipogenesis. Analysis was also expanded to genes identified as potential direct targets of miRNAs, including the genes targeted by miR-33a (retinol dehydrogenase 14a (*rdh14a*), caspase 6 apoptosis-related cysteine peptidase like 2 (*casp6l2*), leucine carboxyl methyltransferase 2 (*lcm2t2*)), the genes targeted by miR-122 (cytochrome P450 family 27 subfamily A polypeptide 7 (*cyp27a7*), methyltransferase like 5 (*mettl5*), histone deacetylase 3 (*hdac3*), hypoxia-inducible factor 1 α subunit a (*hif1aa*)), and the genes targeted by miR-128 (phosphodiesterase 4B cAMP-specific a (*pde4ba*) and RAR-related orphan receptor A paralog a (*rora*)). Total RNA was extracted as previously described (Mennigen et al., 2013) using the Trizol reagent (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions and was quantified by spectrophotometry (absorbance at 260 nm). The integrity of the samples was assessed using agarose gel electrophoresis. 1 μ g of total RNA was used for cDNA synthesis. The SuperScript III RNaseH-reverse transcriptase kit (Invitrogen) with oligo dT random primers (Promega, Charbonnières, France) was used to synthesize cDNA ($n = 6$ for cholesterol metabolism genes in 6 h). The primer sequences used for qRT-PCR analyses are listed in the Table 4 and Table 5. Quantitative RT-PCR assays were performed on the Roche LightCycler 480 II system (Roche Diagnostics, Neuilly sur Seine, France). The assays were carried out using a reaction mix of 6 μ L per sample containing 2 μ L of 76 times diluted cDNA, 0.24 μ L of each primer (10 μ M), 3 μ L of LightCycler 480 SYBR[®] Green I Master mix and 0.52 μ L DNase/RNase free water (5 Prime GmbH, Hamburg, Germany). The PCR protocol was initiated at 95 °C for 10 min for initial denaturation of the cDNA and hot-start Taq-polymerase activation, followed by 45 cycles of a three-step amplification program (15 s at 95 °C, 10 s at melting temperature T_m (60–65 °C), 15 s at 72 °C), according to the primer set used. Melting curves were systematically monitored (5 s at 95 °C, 1 min at 65 °C, temperature gradient at 0.11 °C/s from 65 to 97 °C) at the end of the last amplification cycle to confirm the specificity of the amplification reaction. Each PCR assay included replicate samples (duplicate of reverse transcription and PCR amplification) and negative controls (RT- and cDNA-free samples, respectively). As the expression of *18s* was not significantly different among treatments at 8 h (data not shown) and the expression of *ef1a* was not significantly different among treatments at 16 h (data not shown), the 18S ribosomal RNA (*18s*) and elongation factor 1 α (*ef1a*) was used for the normalization of the genes at 8 h and 16 h, respectively. Relative quantification of target gene expression was determined using the E-Method from the LightCycler 480 software (version SW 1.5; Roche Diagnostics). In all cases, PCR efficiency (E) measured by the slope of a standard curve with serial dilutions of cDNA ranged between 1.8 and 2.

2.8. miRNA expression analysis

Several miRNAs including miR-1-3p, miR-33a-5p, miR-122-5p, miR-128-3p and miR-223-3p were detected by the TaqMan[®] Advanced miRNA Assays (A25576, Applied Biosystems, ThermoFisher) in liver collected 8 h and 16 h after the last meal. The spike miR-39 (*Caenorhabditis. elegans*) was used as an exogenous control for normalization as previously described (Tay et al., 2017) and has been shown to be present at relatively constant levels among the four treatments. Total RNA in liver was obtained in the same way as that for gene expression analysis. 80 ng RNA were used for the poly(A) tailing, ligation and reverse transcription reactions to synthesize the cDNA of

all miRNAs followed by a miR-Amp reaction for cDNA pre-amplification according to the manufacturer's instruction. PCR was performed in a reaction mix of 6 µL containing 2 µL cDNA (200 times diluted for liver cDNA and 50 times diluted for plasma cDNA), 2.67 µL 2× Fast Advanced Master mix (Applied Biosystems, Foster city, California, USA), 0.27 µL TaqMan® Advanced miRNA Assay (20×) (Applied Biosystems, Foster city, California, USA) and 1.06 µL DNase/RNase free water (5 Prime GmbH, Hamburg, Germany). The sequences of the probes were listed in the Table 3. The PCR protocol was initiated at 95 °C for 20s for initial denaturation of the cDNA and the enzyme activation, followed by 50 cycles of a 2 steps amplification program (3 s at 95 °C for denaturation, 30s at 60 °C for annealing). Each PCR assay included replicates for each sample (duplicates of reverse transcription and PCR amplification) and negative controls (reverse transcriptase free and RNA free samples). Relative quantification of the target miRNA was determined using the E-Method from the LightCycler 480 software (version SW 1.5; Roche Diagnostics). PCR efficiency measured by the slope of a standard curve with serial dilutions of miRNA cDNA ranged between 1.8 and 2.

2.9. Statistical analysis

Results are expressed as means ± SD. Statistical analyses were carried out using two-way ANOVA (line, diet effects and line*diet interaction), followed by a Tukey test for post hoc analysis. Normality was assessed using the Shapiro-Wilk test, while homogeneity of variance was determined using Levene's test. For all statistical analysis, the level of significance was set at $P < .05$. Statistical analysis was performed using R software (Fox and Bouchet-Valat, 2016).

3. Results

3.1. Body weight, hepatosomatic index, cholesterol and triglyceride in plasma

After 6 months of feeding, the final body weight of rainbow trout fed the M diet was significantly higher than that fed the V diet. As expected, the fish from the S line reached a higher body weight than those in the C group regardless the dietary treatment. However, as already shown by Le Boucher et al. (2012) and Callet et al. (2017), the selection response was significantly higher when fish were fed the V diet than those fed the M diet. This means that when we selected fish fed 100% V diet for better survival and growth, two kinds of response occurred: 1) general selection for growth which will be expressed whatever the diet, 2) specific selection linked to the V diet. The hepatosomatic index was significantly higher in trout fed the M diet and was not affected by the lines (Table 2).

The cholesterol level in plasma was higher in trout fed the M diet at both 8 h and 16 h after the last meal. It was significantly affected by the line at 16 h after the last meal, with a significantly lower concentration of cholesterol in the plasma in the S line regardless of the diet. The level

Table 3

Sequences of the probes in TaqMan® Advanced miRNA Assays.

Assay	Assay ID	Probe sequences
miR-1-3p	477820_mir	TGGAATGTAAAGAAGTATGTAT
miR-33a-5p	478347_mir	GTGCATTGTAGITGCATTGCA
miR-39-3p	478293_mir	TCACCGGGGTGTAATCAGCTTG
miR-122-5p	477855_mir	TGGAGTGTGACAATGGTGTGTTG
miR-128-3p	477892_mir	TCACAGTGAACCGGTCTCTTT
miR-223-3p	477983_mir	TGTCAGTTTGTCAAATACCCCA
miR-451a	478107_mir	AAACCGTTACCATTACTGAGTT

of triglycerides in plasma was higher in trout fed the M diet than those fed the V diet but only at 16 h after the meal. No significant line effect was found for plasma triglycerides (Table 2).

3.2. Hepatic expression of miRNAs

Expression of miR-1-3p, miR-33a-5p, miR-122-5p, miR-128-3p and miR-223-3p was assessed in liver 8 h and 16 h after the last meal (Fig. 1). The expression of miR-1 and miR-223 was not affected by diet nor by line. Interestingly, miR-33a was expressed significantly lower in the liver of fish fed the M diet at both 8 h and 16 h. The expression of miR-122 and miR-128 significantly increased in the S line compared to the C line at 8 h after the meal. miR-122 was also found to be less expressed in fish fed the V diet but only at 16 h.

3.3. Hepatic expression of genes related to cholesterol and lipid metabolism

We analyzed the hepatic expression of genes related to cholesterol and lipid metabolism or potentially regulated by miR-1-3p, miR-33a-5p, miR-122-5p, miR-128-3p and miR-223-3p (Fig.2). With a high concordance among genes involved in cholesterol synthesis, the expression of *hmgcr*, *hmgcs*, *cyp51* and *dhcr7* significantly increased in the liver of fish fed the V diet at both 8 h and 16 h after the last meal. However, none of them were affected by the line. Correspondingly, the expression of their regulator *srebp-2* showed the same significant variation with higher expression in fish fed the V diet regardless of the lines at both 8 h and 16 h. The genes involved in cholesterol efflux were shown to be affected either by diet or line. The V diet significantly increased the expression of *abcg5* at 8 h, *cyp7a1* and *abca1* at both 8 h and 16 h after the meal. On the contrary, selective breeding for better adaptation to the plant-based diet led to decreased expression of *abcg8* at 16 h after the meal. The only gene not affected by the treatments was *lxra*, which was not affected neither by diet nor line. Regarding the genes involved in cholesterol esterification, a significant interaction between diets and lines was found for *acat1* which was higher expressed in the fish from the selected line fed the M diet. A significantly higher expression of *acat2* was observed in both lines fed the V diet at 8 h after the meal. Whatever the line, the expression of *fas*

Table 2

Morphometric parameters in rainbow trout from selected (S) line and control (C) line fed the marine (M) diet and vegetable (V) diet (C-M represents for the rainbow trout from C line fed the M diet, etc.);

		C-M		C-V		S-M		S-V		ANOVA: P values		
		Mean	SD	Mean	SD	Mean	SD	Mean	SD	Diet	Line	Diet × Line
BW (g)		42.70	5.61	30.23	6.99	54.65	8.33	40.00	4.46	0.000**	0.000**	0.618
		1.51	0.38	1.29	0.15	1.63	0.18	1.35	0.15	0.003**	0.258	0.758
Plasma												
Chol (g/L)	8 h	3.57	0.95	2.17	0.30	4.03	0.89	2.13	0.29	0.000**	0.456	0.379
	16 h	4.69	0.31	2.23	0.59	3.72	0.74	1.74	0.47	0.000**	0.004**	0.306
TG (g/L)	8 h	4.06	0.46	4.36	1.20	4.30	0.93	4.18	1.04	0.809	0.935	0.595
	16 h	4.92	0.99	2.62	1.13	4.27	1.37	2.67	1.14	0.001**	0.537	0.471

(BW: final body weight; HSI: hepatosomatic index; Chol: cholesterol; TG: triglyceride; n = 9 for BW and HSI, n = 6 for Chol and TG, two-way analysis of variance, two stars represent a P value < .01)

Table 4

Primer sequences of cholesterol metabolic and lipogenic genes.

Gene	Forward primer	Reverse primer	Genoscope ^a or Genbank accession numbers	
			Paralogue 1	Paralogue 2
<i>abca1</i>	CAGGAAGACGAGCACCTT	TCTGCCACCTCACACACTTC	GSONMG00078741001	GSONMG00074045001
<i>abcg5</i>	CACCGACATGGAGACAGAAA	GACAGATGGAAGGGGATGAA	GSONMG00075025001	/
<i>abcg8</i>	GATACCAAGGTTCCAGAGCA	CCAGAAACAGAGGGACCAGA	GSONMG00075024001	/
<i>acat1</i>	GGCAAGCCTGATGTGGTAGT	ACCGTGCCATTCTCCTTCTG	GSONMG00033938001	/
<i>acat2</i>	TGCTTGTGTCCCTGGGTTT	GTGTGGCTGTGACGTGTTTC	GSONMG00054446001	GSONMG00054447001
<i>cyp51</i>	CCCGTTGTGACGTTTACCA	GCATTGAGATCTTCGTTCTTGC	GSONMG00031182001	GSONMG00044416001
<i>cyp7a1</i>	ACGTCCGAGTGGCTAAAGAG	GGTCAAAGTGGAGCATCTGG	AB675933.1 GSONMG00066448001	AB675934.1 GSONMG00037174001
<i>dhcr7</i>	GTAACCCACAGACCAAGA	CCTCTCCTATGCAGCCAAC	GSONMG00025402001	GSONMG00039624001
<i>fas</i>	TGATCTGAAGGCCGTGTCA	GGGTGACGTTGCCGTGGTAT	GSONMG00062364001	
<i>gck</i>	GCACGGCTGAGATGCTCTTGT	GCCTTGAACCCCTTGGTCCAG	GSONMG00033781001	GSONMG00012878001
<i>hmgcr</i>	GACCATTTGGGAGCTTGTGT	GAACGGTGAATGTGCTGTGT	GSONMG00016350001	/
<i>hmgcs</i>	AGTGCCAAAGAGAGGGTGTG	TTCTGGTTGGAGACGAGGAG	GSONMG00010243001	/
<i>lxra</i>	TGCAGCAGCCGTATGTGGA	GCGGCGGGAGCTTCTTGTC	GSONMG00014026001	GSONMG00064070001
<i>srebp-1c</i>	CATGCGCAGGTTGTTTCTT	GATGTGTTGCTGTGGGACTG	XM_021624594.1	
<i>srebp-2</i>	TAGGCCCAAAGGGATAAAG	TCAGACACGACGAGCACAA	GSONMG00039651001	GSONMG00061885001
<i>18s</i>	CGGAGGTTTCGAAGACGATCA	TCGCTAGTTGGCATCGTTTAT		
<i>ef1a</i>	TCCTCTTGGTCTGTTTCGCTG	ACCCGAGGGACATCCTGTG	AF498320.1	/

^a <https://www.genoscope.cns.fr/trout>

and *srebp-1c* was higher in fish fed the V diet at both 8 h and 16 h after the meal, however, the expression of *gck* was much lower in fish fed the V diet at 8 h and 16 h. The expression of *gck* was also affected by the line, with much higher expression in the S line at 8 h. In summary, the expression of most genes related to cholesterol and lipid metabolism was affected by the V diet rather than the lines, showing a higher expression in both lines fed the V diet.

3.4. Expression of target genes by corresponding miRNAs in liver

Based on sequence complementarity between the seed sequences of miR-33a-5p, miR-122-5p and miR-128-3p and 3'UTR region of rainbow trout mRNAs, a list of potential direct targets of these previous cited miRNAs was established (Supplementary data S1–3). We analyzed the expression of *rdh14a*, *casp6l2* and *lcmt2* as potential direct targets of miR-33a (Fig. 3A). Among the three genes targeted by miR-33a, only the expression of *casp6l2* was detected to be affected by diet and line. The expression of *casp6l2* was significantly lower in fish fed the V diet at both 8 h and 16 h after the meal and significantly higher in S line only at 8 h after the meal. However, no significant difference was found in the expression of *rdh14a* and *lcmt2*. Regarding the genes targeted by miR-122 (*cyp27a7*, *mettl5*, *hdac3* and *hif1aa*) (Fig. 3B), we observed that feeding trout with the V diet resulted in decreased expression of *cyp27a7* at 8 h after the meal. A significant interaction between diet and line effects was found for *mettl5* at 16 h after the meal, showing a higher expression in the S line than the C line when fish were fed the M diet but a lower expression in the S line when fish were fed the V diet. On the other hand, the expression of *hdac3* and *hif1aa* was not affected by diets or lines at both time points. We finally analyzed *pde4ba* and *rora* as putative direct targets of miR-128 (Fig. 3C). Both genes expressed at

Table 6

Correlations between miRNAs and their potential target genes.

miRNAs	Potential target gene	Pearson value	
		8 h	16 h
miR-33a	<i>rdh14a</i>	−0.31	0.3051
	<i>casp6l2</i>	−0.47 [*]	−0.3813
	<i>lcmt2</i>	−0.06	−0.0878
miR-122	<i>cyp27a7</i>	0.42 [*]	0.0442
	<i>mettl5</i>	0.21	0.3137
	<i>hdac3</i>	−0.13	0.1321
	<i>hif1aa</i>	0.08	0.259
miR-128	<i>pde4ba</i>	−0.25	0.3258
	<i>rora</i>	−0.09	0.9701

Pearson correlation coefficient larger than zero is positive correlation, while Pearson correlation coefficient less than zero is negative correlation.

* $P < .05$ is statistically significant with one star.

much lower level in the fish fed the V diet at 8 h after the meal. This lowering effect of the V diet was maintained 16 h after the meal for *pde4ba*. Finally, results showed that the expression of *pde4ba* at 16 h after the meal was also affected by the line, with a lower expression in the S line.

3.5. Correlations between miRNAs and their putative target genes

Pearson correlation analysis showed that miR-33a negatively correlated with *casp6l2* and miR-122 positively correlated with *cyp27a7* significantly 8 h after the meal, however, other pairs of miRNAs and their target genes did not show any significant correlations (Table 6).

Table 5

Primer sequences of miRNA target genes.

Gene	F primer	R primer	Accession number
<i>rdh14a</i>	TCCAGGGCCTTCTCAAGAG	CAAAGTTCTCTGGCCACCTC	D2OM_RDH14A.2.2
<i>casp6l2</i>	TTCGTGTGCGCTCTTCTGAG	CACCTGGATCGTCATGCTTGT	D2OM_LOC100135902.1.1
<i>lcmt2</i>	CCGACAACCCCTTCACAGATT	GAGCTTTTCGTTCCCTACC	D2OM_LCMT1.1.1
<i>cyp27a7</i>	AGTGGTACAAGCTGCGGACT	GGACAGGTTGGTCAACAAGGT	D2OM_CYP27A7.3.3
<i>mettl5</i>	AAGTGGACGCTTTGAAGAA	CCAACACACAAACCTGCATC	D2OM_METL5.1.1
<i>hdac3</i>	GGCAACTCTTTAGCCAGTC	GGGGAATCCTGAAGCTTTTC	D2OM_HDAC3.2.3
<i>hif1aa</i>	CCCAACCCCTAGAGTGCTC	TGGTGAGTAAGGAAGCAGGG	D2OM_HIF1A.3.4
<i>pde4ba</i>	CAGCAATCCATGATGTGGAC	TTGGTCAGGCTCTGGAAGAT	D2OM_PDE4B.2.2
<i>rora</i>	TCAGTGGCTACATGGACAGC	GGGATGTCTCTCCGTGGTA	D2OM_contig_006818

Fig. 1. miRNAs expression involved in cholesterol metabolism at 8 h and 16 h including miR-1, miR-33a, miR-122, miR-128 and miR-223 in the liver of rainbow trout from selected (S) line and control (C) line fed the marine (M) diet and (V) vegetable diet. The selected line was selected for better survival and growth performance with a 100% plant-based diet during three generations. The control line was the common trout without any artificial selection. Expression values were normalized by spike miR-39. Values are means ($n = 6$), with their standard deviations represented by vertical bars. (Results of two-way analysis of variance are indicated with the P value). One star/circle represents a P value $< .05$ for diets/lines, respectively; two stars/circles represent a P value $< .01$ for diets/lines, respectively.

4. Discussion

Selective breeding in fish for better adaptation to plant-based diet represents a promising future in aquaculture when increasingly more fishmeal and fish oil are replaced by vegetable ingredients in aquafeeds. Recent selective breeding program developed in rainbow trout has proven that individual variability in the ability to survive and grow on the plant-based diet can be effectively used to promote fish production (Le Boucher et al., 2011). In order to decipher the molecular mechanisms underlying this adaptation to the plant-based diet totally devoid of marine ingredients, we examined the expression of five miRNAs potentially involved in the regulation of lipid and cholesterol metabolism in the liver of control and selected rainbow. We also analyzed the hepatic expression of putative indirect and direct target genes of these miRNAs.

Our study demonstrated that the hepatic expression of miRNAs could be affected either in fish fed the 100% plant-based diet or in line selected for better adaptation to the plant-based diet by selective breeding. The vegetable diet increased the hepatic expression of miR-33a, while selective breeding for better survival and growth on the totally plant-based diet enhanced the expression of miR-122 and miR-128. However, the miR-1 (reported to inhibit *lra* expression (Zhong et al., 2013), which may result in cellular cholesterol increase via reducing cholesterol efflux and conversion to bile acids) and miR-223 (known to reduce cellular cholesterol and promote cholesterol efflux by directly repressing *hmgcs* and indirectly increasing *abca1* and *cyp7a1*, respectively (Vickers et al., 2014)), were not modified by the diet nor the selective breeding.

One particularity of the miR-33a in human is that its gene is located in the intron of *srebp-2*, the gene encoding SREBP-2 protein (Marquart et al., 2010). This particularity seems to be conserved across animal species, since miR-33a is also positioned in an intron of the *srebp-2* gene in rainbow trout. The concomitance and co-expression of *srebp-2* and miR-33a could be the reason why the expression of both miR-33a and *srebp-2* were found to be concurrently higher in trout fed the plant-based diet. In mammals, miR-33a could repress the cholesterol efflux by targeting *abcg5* and *abcg8* (involved in direct cholesterol discharge into the bile), *cyp7a1* (key enzyme in bile acid synthesis) and *abca1* (contributing to hepatic cholesterol efflux to plasma) (Bommer and MacDougald, 2011; Gerin et al., 2010; Li et al., 2013). The increasing

expression of miR-33a might indicate the repression of cholesterol export and bile acid synthesis (Gerin et al., 2010; Li et al., 2013) when trout were fed the plant-based diet. However, the higher expression of cholesterol efflux genes (*abca1*, *cyp7a1* and *abcg5*) found in trout fed the V diet in the present study goes against this hypothesis. This discrepancy may be attributed to the variety of factors acting at transcriptional or post-transcriptional levels that orchestrate the regulation of gene expression together. For example, in mammals, it was reported that *srebp-2* positively regulated the expression of *abca1* (Wong et al., 2006) and the unusually long 3'UTR of *abca1* also harbors many binding sites of other miRNAs (Ramirez et al., 2011) that probably also contribute to the regulation of *abca1*. Therefore, in the present study, the higher expression of *abca1* in trout fed the V diet may be due to the higher expression of *srebp-2* and/or the involvement of other miRNAs counteracting the effect of the miR-33a.

miR-122, a liver specific miRNA, has been suggested to play roles in cholesterol and lipid metabolism (Esau et al., 2006; Krützfeldt et al., 2005). In contrast to most other miRNAs inhibiting gene expression, miR-122 works to promote the expression of genes involved in cholesterol and fatty acid synthesis through a mechanism still unknown (Aryal et al., 2017; Iliopoulos et al., 2010; Krützfeldt et al., 2005). In the present study, miR-122 expression was shown to be affected by the selective breeding with enhanced expression in the selected line compared to the control line at 8 h after the meal. However, the increase in miR-122 in the selected line was not associated with a concomitant enhanced expression of genes involved in cholesterol biosynthesis (*hmgcr*, *hmgcs*, *cyp51* and *dhcr7*) or lipogenesis (*fas* and *srebp-1c*). Similarly, the promoting effect of the vegetable diet on the expression of the foregoing genes was not accompanied by an enhanced expression of miR-122. Therefore, the present study confirms our previous results that in vivo inhibition of miR-122 by antagomir in trout did not affect the expression of genes involved in cholesterol synthesis (Mennigen et al., 2014).

Similar to miR-122, the expression of miR-128 increased in the selected line regardless of the diets. This increase was only observed 8 h after the meal, confirming our previous investigations that miRNAs could be under postprandial nutritional control in fish (Mennigen et al., 2012). miR-128 is considered as an actor to regulate cholesterol homeostasis by limiting cholesterol efflux (Adlakha et al., 2013). Cholesterol efflux is a mechanism implemented by cells to eliminate excess

cholesterol and prevent cellular cholesterol accumulation. This efflux is mediated by the ABCA1, of which expression has been shown to be inhibited at both mRNA and protein level by miR-128 through direct binding to the 3'UTR region of *abca1* mRNA (Adlakha et al., 2013). The positive effect of the selective breeding on the expression of miR-128 seems to be consistent with the lower plasma cholesterol level measured in the selected line at 16 h after the meal. However, the higher expression of miR-128 but lower plasma cholesterol in selected line was

inconsistent with the result of *abca1* expression at mRNA level, which was not affected by selective breeding. Therefore, further investigations are still needed to determine if *abca1* expression could be affected by the miR-128 at protein level rather than mRNA level.

In the present study, the expressions of miR-33a, miR-122 and miR-128 in liver were all affected either by diets or lines, while miR-1 and miR-223 were not affected by either line or diet. Therefore, we further utilized the 3'UTR of these three miRNAs to identify their putative

Fig. 2. Expression of genes involved in cholesterol synthesis (A), cholesterol efflux (B), cholesterol regulation (C), cholesterol esterification (D), and fatty acid synthesis (E) at 8 h and 16 h in the liver of rainbow trout from selected (S) line and control (C) line fed the marine (M) diet and (V) vegetable diet. The selected line was selected for better survival and growth performance with a 100% plant-based diet during three generations. The control line was the common trout without any artificial selection. Expression values were normalized by 18s at 8 h and ef1 α at 16 h. Values are means (n = 6), with their standard deviations represented by vertical bars. (Results of two-way analysis of variance are indicated with the P value). One star/circle represents a P value < .05 for diets/lines, respectively; two stars/circles represent a P value < .01 for diets/lines, respectively.

(C)

<i>srebp-2</i>	8h	16h
diet	0.00	0.00
line	0.60	0.13
diet x line	0.57	0.96

<i>lxra</i>	8h	16h
diet	0.20	0.08
line	0.16	0.30
diet x line	0.24	0.58

(D)

<i>acat1</i>	8h	16h
diet	0.31	0.10
line	0.07	0.76
diet x line	0.02	0.22

<i>acat2</i>	8h	16h
diet	0.00	0.09
line	0.99	0.64
diet x line	0.18	0.63

(E)

<i>fas</i>	8h	16h
diet	0.00	0.01
line	0.33	0.42
diet x line	0.56	0.87

<i>srebp-1c</i>	8h	16h
diet	0.04	0.00
line	0.58	0.68
diet x line	0.56	0.68

<i>gck</i>	8h	16h
diet	0.00	0.00
line	0.04	0.91
diet x line	0.39	0.12

Fig. 2. (continued)

direct target genes and then analyzed the expression of the suggested genes to see if they could be regulated by the corresponding miRNAs.

In contrary to the increased expression of miR-33a in trout fed the V diet, its putative direct target *casp6l2* showed significantly lower expression in trout fed the V diet regardless of time point, indicating the expression of *casp6l2* could be post-transcriptionally inhibited by miR-33a. The negative correlation between *casp6l2* and miR-33a is also consistent with this speculation. CASP6L2 belongs to the caspase family of cysteine proteases associated with immune responses and apoptosis. It is known that the activation of caspases proceeds to the production of pro-inflammatory cytokines and apoptosis (Siegel, 2006). As the immune function is affected in salmonids when marine ingredients are replaced by plant ingredients in aquafeeds (Martin and Król, 2017), the regulation of *casp6l2* by diet composition is not surprising.

Among the miR-122 putative target genes analyzed in the present study, only *cyp27a7* was found to be affected, with a lower expression

in fish fed the V diet. In the present study, the expression of miR-122 was subjected to an up-regulation linked to selective breeding 8 h after the last meal and a down-regulation by the plant-based diet 16 h after the last meal regardless of the fish line. As no significant effect of selective breeding was recorded for *cyp27a7* expression, its direct regulation by miR-122 remains questionable. Correspondingly, correlation analysis did not show the negative correlation between miR-122 and *cyp27a7*, either. The physiological role of CYP27A7 still remains to be determined. However, like the sterol 27-hydroxylase (CYP27A), a member of the same cytochrome family, CYP27A7 may be involved in the cholesterol and bile acid metabolism as well (Dubrac et al., 2005). In this respect, its lower expression in fish fed the V diet is in agreement with the impact of vegetable diet on cholesterol and bile acid metabolism.

Though the expression of miR-128 was up-regulated in the selected line 8 h after the last meal, the expression of *pde4ba* and *rora*, two

potential direct targets of miR-128, was not affected by selective breeding. This inconsistency may suggest that miR-128 is not involved in the regulation of these genes at mRNA level. The absence of correlation between miR-128 and *pde4ba* and *rora* also supports this speculation. On the other hand, the expression of these genes did, however, decrease in the trout fed the V diet. PDE4B belongs to a family of four genes (from PDE4A to D) that act to hydrolyze cAMP and switch off cAMP signaling cascades (Houslay and Adams, 2003). A recent study indicated that an increased hepatic expression of *pde4b* could alter cAMP signaling, impair fatty acid oxidation and predispose to liver steatosis (Avila et al., 2016). A current phenotypic trait observed in

trout fed the totally plant-based diet showed that the decrease in hepatosomatic index was related to the reduction of lipid content in the liver (Lazzarotto et al., 2015). The reduced expression of *pde4ba* observed in trout fed the plant-based diet is thus in agreement with the reduced accumulation of fat in the liver, which may be consistent with the role assigned to PDE4B in mammals. RORAA is a member of the retinoid-related orphan receptor (ROR) subfamily. These receptors play critical roles in development, immunity, circadian rhythm, and cellular metabolism (Jetten, 2009). It is reported that impaired expression of *rora* in natural sg/sf mutant mice leads to reduced plasma cholesterol level by limiting cholesterol efflux. These mice also exhibited reduced

(A)

(B)

Fig. 3. Expression of possible target genes by miR-33a (A), miR-122 (B) and miR-128 (C) identified in silico at 8 h and 16 h in the liver of rainbow trout from selected (S) line and control (C) line fed the marine (M) diet and (V) vegetable diet. The selected line was selected for better survival and growth performance with a 100% plant-based diet during three generations. The control line was the common trout without any artificial selection. Expression values were normalized by 18s at 8 h and *ef1a* at 16 h. Values are means (n = 6), with their standard deviations represented by vertical bars. (Results of two-way analysis of variance are indicated with the P value). One star/circle represents a P value < .05 for diets/lines, respectively; two stars/circles represent a P value < .01 for diets/lines, respectively.

(C)

Fig. 3. (continued)

expression of *srebp-1c* as well as of its target gene, fatty acid synthase (*fas*) (Lau et al., 2008). In the present study, because the lower expression of *rora* but higher expression of *abca1*, *abcg5*, *srebp-1c* and *fas* in fish fed the plant-based diet was inconsistent with the results in mice study, further investigations are needed to better understand the role of these retinoid-related orphan receptors in trout metabolism.

Among the genes that were chosen as putative targets of miR-33a, –122 and –128, some of them, including *rdh14*, *lcmt2*, *mttl5*, *hdac3* and *hif1aa*, were neither affected by fish lines nor diets. It has been established that miRNAs regulate gene expression via two main mechanisms depending on the degree of complementarity between miRNA and its target gene. When complementarity is high, mRNA is cleaved and degraded through the RNA-induced silencing complex (RISC). Conversely, low complementarity resulted in translational inhibition of the mRNA target and reduction of the abundance of the encoded protein (Zeng et al., 2003). Thus it is necessary to conduct additional studies at protein level to clearly evaluate the involvement of miR-33a, –122 or –128 in the regulation of these genes.

In conclusion, the present study demonstrated that hepatic expression of miRNAs may be affected by selective breeding of rainbow trout for better adaptation to plant-based diets as well as by total dietary replacement of fishmeal and fish oil by plant ingredients. The miR-33a was found to be greatly induced by plant-based diet whereas miR-122 and miR-128 were affected by selective breeding. This study opens new perspective to evaluate the role of miRNAs in fish nutrition and selective breeding. However, the present study is restricted to the investigation of gene expression at mRNA level instead of protein due to the unavailability of antibodies for Western blot in fish. Even though some relevant regulations were observed between miRNAs and their potential target genes, additional investigations necessitating the development of antibodies are essential to further establish the role of miRNAs in rainbow trout metabolism. Finally, the present study also highlighted some new genes affected by plant-based diet, thus linking this change in dietary composition to, among others, immune response, apoptosis and cAMP signaling pathways. The identification of these new molecular targets in the replacement of marine ingredients by plant products in rainbow trout may aid in opening new fields of investigation to better understand the mechanisms of adaptation to plant-based diet in carnivorous fish.

Acknowledgements

The authors thank F. Sandres and F. Terrier for manufacturing the experimental diets (INRA, Donzaq, France), L. Labbé and T. Kerneis for fish rearing and A. Herman for technical assistance in the laboratory.

Financial support

This work was carried out within the framework of the AGREINFISH project (project ANR-13-ADAP-001) funded by the French National Research Agency (ANR). T. Zhu gratefully acknowledges the financial assistance provided by the China Scholarship Council (CSC, File No. 201406330071) for his doctoral fellowship.

Conflict of interest

None.

Authorship

SS, GC: formulated research questions and designed the study. EQ, MDN: provided the lines. JM, JB: provided information on direct target gene of miRNA. TZ, EPJ: performed the analysis. TZ, SS, GC: analyzed the data. TZ, SS, GC wrote the paper.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.aquaculture.2018.08.058>.

References

- Adlakha, Y.K., Khanna, S., Singh, R., Singh, V.P., Agrawal, A., Saini, N., 2013. Pro-apoptotic miRNA-128-2 modulates ABCA1, ABCG1 and RXR α expression and cholesterol homeostasis. *Cell Death Dis.* 4, e780.
- AOAC, 2000. Official Methods of Analysis of AOAC International. Gaithersburg, MD USA.
- Aryal, B., Singh, A.K., Rotllan, N., Price, N., Fernández-Hernando, C., 2017. MicroRNAs and lipid metabolism. *Curr. Opin. Lipidol.* 28, 273–280.
- Avila, D.V., Barker, D.F., Zhang, J., McClain, C.J., Barve, S., Gobejishvili, L., 2016. Dysregulation of hepatic cAMP levels via altered Pde4b expression plays a critical role in alcohol-induced steatosis. *J. Pathol.* 240, 96–107.

- Benedito-Palos, L., Navarro, J.C., Sitjà-Bobadilla, A., Bell, J.G., Kaushik, S., Pérez-Sánchez, J., 2008. High levels of vegetable oils in plant protein-rich diets fed to gilthead sea bream (*Sparus aurata* L.): growth performance, muscle fatty acid profiles and histological alterations of target tissues. *Br. J. Nutr.* 100, 992–1003.
- Bommer, G.T., MacDougald, O.A., 2011. Regulation of lipid homeostasis by the bifunctional SREBF2-miR33a locus. *Cell Metab.* 13, 241–247.
- Callet, T., Médale, F., Larroquet, L., Surget, A., Aguirre, P., Kerneis, T., Labbé, L., Quillet, E., Geurden, I., Skiba-Cassy, S., Dupont-Nivet, M., 2017. Successful selection of rainbow trout (*Oncorhynchus mykiss*) on their ability to grow with a diet completely devoid of fishmeal and fish oil, and correlated changes in nutritional traits. *PLoS One* 12, e0186705.
- Dubrac, S., Lear, S.R., Ananthanarayanan, M., Balasubramanian, N., Bollineni, J., Shefer, S., Hyogo, H., Cohen, D.E., Blanche, P.J., Krauss, R.M., Batta, A.K., Salen, G., Suchy, F.J., Maeda, N., Erickson, S.K., 2005. Role of CYP27A in cholesterol and bile acid metabolism. *J. Lipid Res.* 46, 76–85.
- Esau, C., Davis, S., Murray, S.F., Yu, X.X., Pandey, S.K., Pear, M., Watts, L., Booten, S.L., Graham, M., McKay, R., Subramaniam, A., Propp, S., Lollo, B.A., Freier, S., Bennett, C.F., Bhanot, S., Monia, B.P., 2006. miR-122 regulation of lipid metabolism revealed by in vivo antisense targeting. *Cell Metab.* 3, 87–98.
- FAO, 2016. The State of World Fisheries and Aquaculture: 2016. Contributing to food security and nutrition for all. Rome.
- Fox, J., Bouchet-Valat, M., 2016. Rcmdr: R Commander. (R package version 2.3-0).
- Geay, F., Ferrarese, S., Zambonino-Infante, J.L., Bargelloni, L., Quentel, C., Vandeputte, M., Kaushik, S., Cahu, C.L., Mazurais, D., 2011. Effects of the total replacement of fish-based diet with plant-based diet on the hepatic transcriptome of two European sea bass (*Dicentrarchus labrax*) half-sibfamilies showing different growth rates with the plant-based diet. *BMC Genomics* 12, 522.
- Gerin, I., Clerbaux, L.A., Haumont, O., Lanthier, N., Das, A.K., Burant, C.F., Leclercq, I.A., MacDougald, O.A., Bommer, G.T., 2010. Expression of miR-33 from an SREBP2 intron inhibits cholesterol export and fatty acid oxidation. *J. Biol. Chem.* 285, 33652–33661.
- Goedeke, L., Fernández-Hernando, C., 2012. Regulation of cholesterol homeostasis. *Cell Mol. Life Sci.* 69, 915–930.
- Horn, S.S., Sonesson, A.K., Ruyter, B., 2017. Breeding for More Omega-3 in Salmon. *Nofima*. <https://nofima.no/en/nyhet/2017/11/breeding-for-more-omega-3-in-salmon/?lang=en>.
- Houslay, M.D., Adams, D.R., 2003. PDE4 cAMP phosphodiesterases: modular enzymes that orchestrate signalling cross-talk, desensitization and compartmentalization. *Biochem. J.* 370, 1–18.
- Iliopoulos, D., Drosatos, K., Hiya, Y., Goldberg, I.J., Zannis, V.I., 2010. MicroRNA-370 controls the expression of microRNA-122 and Cpt1alpha and affects lipid metabolism. *J. Lipid Res.* 51, 1513–1523.
- Iwakawa, H., Tomari, Y., 2015. The functions of microRNAs: mRNA decay and translational repression. *Trends Cell Biol.* 25, 651–665.
- Jackson, A., 2012. Fishmeal and fish oil and its role in sustainable aquaculture. *Int. Aquafeed* 15, 18–21.
- Jetten, A.M., 2009. Retinoid-related orphan receptors (RORs): critical roles in development, immunity, circadian rhythm, and cellular metabolism. *Nucl. Recept. Signal.* 7, Krützfeldt, J., Rajewsky, N., Braich, R., Rajeev, K.G., Tuschl, T., Manoharan, M., Stoffel, M., 2005. Silencing of microRNAs in vivo with 'antagomirs'. *Nature* 438, 685–689.
- Lau, P., Fitzsimmons, R.L., Raichur, S., Wang, S.M., Lechtken, A., Muscat, G.E.O., 2008. The orphan nuclear receptor, ROR α , regulates gene expression that controls lipid metabolism: staggerer (sg/sg) mice are resistant to diet-induced obesity. *J. Biol. Chem.* 283, 18411–18421.
- Lazzarotto, V., Corraze, G., Leprevost, A., Quillet, E., Dupont-Nivet, M., Médale, F., 2015. Three-year breeding cycle of rainbow trout (*Oncorhynchus mykiss*) fed a plant-based diet, totally free of marine resources: Consequences for reproduction, fatty acid composition and progeny survival. *PLoS One* 10, 1–17.
- Le Boucher, R., Quillet, E., Vandeputte, M., Lecalvez, J.M., Goardon, L., Chatain, B., Médale, F., Dupont-Nivet, M., 2011. Plant-based diet in rainbow trout (*Oncorhynchus mykiss* Walbaum): are there genotype-diet interactions for main production traits when fish are fed marine vs. plant-based diets from the first meal? *Aquaculture* 321, 41–48.
- Le Boucher, R., Dupont-Nivet, M., Vandeputte, M., Kerneis, T., Goardon, L., Labbe, L., Chatain, B.B., Bothaire, M.J.J., Larroquet, L., Médale, F., Labbé, L., Chatain, B.B., Bothaire, M.J.J., Larroquet, L., Médale, F., Quillet, E., 2012. Selection for adaptation to dietary shifts: towards sustainable breeding of carnivorous fish. *PLoS One* 7, e44898.
- Leaver, M.J., Taggart, J.B., Villeneuve, L., Bron, J.E., Guy, D.R., Bishop, S.C., Houston, R.D., Matika, O., Tocher, D.R., 2011. Heritability and mechanisms of n-3 long chain polyunsaturated fatty acid deposition in the flesh of Atlantic salmon. *Comp. Biochem. Physiol. Part D Genomics Proteomics* 6, 62–69.
- Li, T., Franc, J.M., Boehme, S., Chiang, J.Y.L., 2013. Regulation of cholesterol and bile acid homeostasis by the cholesterol 7 α -hydroxylase/steroid response element-binding protein 2/microRNA-33a axis in mice. *Hepatology* 58, 1111–1121.
- Marquart, T.J., Allen, R.M., Ory, D.S., Baldán, A., 2010. miR-33 links SREBP-2 induction to repression of sterol transporters. *Proc. Natl. Acad. Sci.* 107, 12228–12232.
- Martin, S.A., Król, E., 2017. Nutrigenomics and immune function in fish: new insights from omics technologies. *Dev. Comp. Immunol.* 75, 86–98.
- Mennigen, J.A., Panerat, S., Larquier, M., Plagnes-Juan, E., Médale, F., Seiliez, I., Skiba-Cassy, S., 2012. Postprandial regulation of hepatic microRNAs predicted to target the insulin pathway in rainbow trout. *PLoS One* 7, e38604.
- Mennigen, J.A., Skiba-Cassy, S., Panerat, S., 2013. Ontogenetic expression of metabolic genes and microRNAs in rainbow trout alevins during the transition from the endogenous to the exogenous feeding period. *J. Exp. Biol.* 216, 1597–1608.
- Mennigen, J.A., Martyniuk, C.J., Seiliez, I., Panerat, S., Skiba-Cassy, S., 2014. Metabolic consequences of microRNA-122 inhibition in rainbow trout, *Oncorhynchus mykiss*. *BMC Genomics* 15, 70.
- Najafi-Shoushtari, S.H., Kristo, F., Li, Y., Shioda, T., Cohen, D.E., Gerszten, R.E., Näär, A.M., 2010. MicroRNA-33 and the SREBP host genes cooperate to control cholesterol homeostasis. *Science* 328, 1566–1569.
- National Research Council, 2011. Nutrient Requirements of Fish and Shrimp. National academies press, Washington, DC.
- Panerat, S., Hortalon, G.A., Plagnes-Juan, E., Kolditz, C., Lansard, M., Skiba-Cassy, S., Esquerre, D., Geurden, I., Médale, F., Kaushik, S., Corraze, G., Esquerre, D., Geurden, I., Médale, F., Kaushik, S., 2009. Differential gene expression after total replacement of dietary fish meal and fish oil by plant products in rainbow trout (*Oncorhynchus mykiss*) liver. *Aquaculture* 294, 123–131.
- Quinton, C.D., Kause, A., Ruohonen, K., Koskela, J., 2007. Genetic relationships of body composition and feed utilization traits in European whitefish (L.) and implications for selective breeding in fishmeal- and soybean meal-based diet environments. *J. Anim. Sci.* 85, 3198–3208.
- Ramirez, C.M., Da, A., Goedeke, L., Salerno, A.G., Warrier, N., Cirera-Salinas, D., Sua, Y., Ferna, C., 2011. MicroRNA-758 regulates cholesterol efflux through posttranscriptional repression of ATP-binding cassette transporter A1. *Arterioscler. Thromb. Vasc. Biol.* 31, 2707–2714.
- Sakakura, Y., Shimano, H., Sone, H., Takahashi, A., Inoue, K., Toyoshima, H., Suzuki, S., Yamada, N., 2001. Sterol regulatory element-binding proteins induce an entire pathway of cholesterol synthesis. *Biochem. Biophys. Res. Commun.* 286, 176–183.
- Siegel, R.M., 2006. Caspases at the crossroads of immune-cell life and death. *Nat. Rev. Immunol.* 6, 308–317.
- Stadtman, T.C., 1957. [63] Preparation and assay of cholesterol and ergosterol. *Methods Enzymol.* 3, 392–394.
- Tacon, A.G.J., Hasan, M.R., Allan, G., El-Sayed, A.F.M., Jackson, A., Kaushik, S.J., Ng, W.K., Suresh, V., Viana, M.T., 2010. Aquaculture feeds: addressing the long-term sustainability of the sector. In: *Proceedings of the Global Conference on Aquaculture*, pp. 193–232.
- Tay, J.W., James, I., Hughes, Q.W., Tiao, J.Y., Baker, R.I., 2017. Identification of reference miRNAs in plasma useful for the study of oestrogen-responsive miRNAs associated with acquired Protein S deficiency in pregnancy. *BMC Res. Notes* 10, 312.
- Tocher, D.R., Bendiksen, E.Å., Campbell, P.J., Bell, J.G., 2008. The role of phospholipids in nutrition and metabolism of teleost fish. *Aquaculture* 280, 21–34.
- Torstensen, B.E., Espe, M., Sanden, M., Stubhaug, I., Waagbø, R., Hemre, G.-I., Fontanillas, R., Nordgarden, U., Hevrøy, E.M., Olsvik, P., 2008. Novel production of Atlantic salmon (*Salmo salar*) protein based on combined replacement of fish meal and fish oil with plant meal and vegetable oil blends. *Aquaculture* 285, 193–200.
- Vickers, K.C., Landstreet, S.R., Levin, M.G., Shoucri, B.M., Toth, C.L., Taylor, R.C., Palmisano, B.T., Tabet, F., Cui, H.L., Rye, K.-A., Sethupathy, P., Remaley, A.T., 2014. MicroRNA-223 coordinates cholesterol homeostasis. *Proc. Natl. Acad. Sci.* 111, 14518–14523.
- Wang, L., Jia, X.-J., Jiang, H.-J., Du, Y., Yang, F., Si, S.-Y., Hong, B., 2013. MicroRNAs 185, 96, and 223 repress selective high-density lipoprotein cholesterol uptake through posttranscriptional inhibition. *Mol. Cell. Biol.* 33, 1956–1964.
- Wong, J., Quinn, C.M., Brown, A.J., 2006. SREBP-2 positively regulates transcription of the cholesterol efflux gene, ABCA1, by generating oxysterol ligands for LXR. *Biochem. J.* 400, 485–491.
- Zeng, Y., Yi, R., Cullen, B.R., 2003. MicroRNAs and small interfering RNAs can inhibit mRNA expression by similar mechanisms. *Proc. Natl. Acad. Sci.* 100, 9779–9784.
- Zhao, C., Dahlman-Wright, K., 2010. Liver X receptor in cholesterol metabolism. *J. Endocrinol.* 204, 233–240.
- Zhong, D., Huang, G., Zhang, Y., Zeng, Y., Xu, Z., Zhao, Y., He, X., He, F., 2013. MicroRNA-1 and microRNA-206 suppress LXR α -induced lipogenesis in hepatocytes. *Cell. Signal.* 25, 1429–1437.