



HAL
open science

A decomposition theorem for smoothable varieties with trivial canonical class

Stéphane Druel, Henri Guenancia

► **To cite this version:**

Stéphane Druel, Henri Guenancia. A decomposition theorem for smoothable varieties with trivial canonical class. *Journal de l'École polytechnique - Mathématiques*, 2018, 5, pp.117 - 147. 10.5802/jep.65 . hal-01879001

HAL Id: hal-01879001

<https://hal.science/hal-01879001>

Submitted on 18 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A DECOMPOSITION THEOREM FOR SMOOTHABLE VARIETIES WITH TRIVIAL CANONICAL CLASS

STÉPHANE DRUEL AND HENRI GUENANCIA

ABSTRACT. In this paper we show that any smoothable complex projective variety, smooth in codimension two, with klt singularities and numerically trivial canonical class admits a finite cover, étale in codimension one, that decomposes as a product of an abelian variety, and singular analogues of irreducible Calabi-Yau and irreducible symplectic varieties.

CONTENTS

1.	Introduction	1
2.	Notation, conventions, and basic facts	3
3.	Kähler-Einstein metrics on smoothable spaces	5
4.	Relative Albanese morphism	7
5.	Holonomy and stability of the tangent sheaf	9
6.	Towards a decomposition theorem	11
7.	Proof of Theorem A	14
8.	Examples	16
	References	19

1. INTRODUCTION

The Beauville-Bogomolov decomposition theorem asserts that any compact Kähler manifold with numerically trivial canonical bundle admits an étale cover that decomposes into a product of a torus, and irreducible, simply-connected Calabi-Yau, and symplectic manifolds (see [Bea83]).

With the development of the minimal model program, it became clear that singularities arise as an inevitable part of higher dimensional life. If X is any complex projective manifold with Kodaira dimension $\kappa(X) = 0$, standard conjectures of the minimal model program predict the existence of a birational contraction $X \dashrightarrow X_{\min}$, where X_{\min} has terminal singularities and $K_{X_{\min}} \equiv 0$. This makes it imperative to extend the Beauville-Bogomolov decomposition theorem to the singular setting.

In the singular setting, several notions of irreducible Calabi-Yau varieties or irreducible symplectic varieties have been proposed but the results of [GKP16] and [GGK17] provide evidence that the following definition should be the correct one in view of a singular analogue of the Beauville-Bogomolov decomposition theorem.

Definition 1.1. Let X be a normal projective variety with canonical singularities such that $K_X \sim_{\mathbb{Z}} 0$.

- (1) We call X *irreducible Calabi-Yau* if $h^0(Y, \Omega_Y^{[q]}) = 0$ for all numbers $0 < q < \dim X$ and all finite covers $Y \rightarrow X$, étale in codimension one.
- (2) We call X *irreducible symplectic* if there exists a holomorphic symplectic 2-form $\sigma \in H^0(X, \Omega_X^{[2]})$ such that for all finite étale covers $f: Y \rightarrow X$, étale in codimension one, the exterior algebra of global reflexive forms is generated by $f^{[*]}\sigma$.

Let X be a normal projective variety of dimension at least 2 with $K_X \equiv 0$ and klt singularities. Suppose moreover that its tangent sheaf is strongly stable in the sense of Definition 2.3. In [GGK17, Theorem C], it is proved that X admits a finite cover, étale in codimension one, that is either an irreducible Calabi-Yau variety or an irreducible symplectic variety. That result, given the infinitesimal version of the Beauville-Bogomolov

Date: May 9, 2017.

2010 Mathematics Subject Classification. 14J32, 14E30.

decomposition theorem proved in [GKP16], is a strong indication that the notions of irreducible Calabi-Yau and symplectic varieties described in Definition 1.1 should be the good ones.

In [Dru16] the first author extends the Beauville-Bogomolov decomposition theorem to complex projective varieties of dimension at most five with klt singularities and numerically trivial canonical class. The main result of our paper is the following decomposition theorem for smoothable mildly singular spaces with numerically trivial canonical class.

Theorem A. *Let X be a normal complex projective variety with klt singularities and smooth in codimension two. Suppose that $K_X \equiv 0$. Suppose furthermore that there exists a flat projective holomorphic map with connected fibers $f: \mathcal{X} \rightarrow \Delta$ from a normal analytic space \mathcal{X} onto the complex open unit disk Δ such that $X \cong f^{-1}(0)$ and such that $f^{-1}(t)$ is smooth for $t \neq 0$. Then there exists a finite cover $Y \rightarrow X$, étale in codimension one, and a decomposition of Y into a product of an abelian variety and irreducible, Calabi-Yau and symplectic varieties.*

Remark 1.2. The assumptions of Theorem A imply that X does not have any quotient singularity. Indeed, a theorem of Schlessinger [Sch71, Theorem 2] (see also [Art76, Theorem 10.1]) shows that a germ of a quotient singularity (X, x) is rigid as soon as $\text{codim}\{x\} \geq 3$.

In addition to the smoothability condition, the strategy of proof of Theorem A requires us to assume that the $\text{codim}(X \setminus X_{\text{reg}}) \geq 3$. Let us briefly explain why. The idea of the proof is to consider a cover of the smooth generic fiber that splits off an abelian variety as well as irreducible, simply-connected Calabi-Yau and symplectic manifolds. A significant part of the paper is devoted to showing that one can take the flat limits of these irreducible pieces and recover the central fiber as product of those limits.

It is then tempting to believe that the flat limit X of irreducible and simply-connected, Calabi-Yau or symplectic manifolds admits a finite cover, étale in codimension one, which is an irreducible Calabi-Yau or symplectic variety. Unfortunately, this turns out to be false in general as we explain in Section 8.3. This makes it much harder to use the smoothability assumption in order to prove a decomposition theorem in full generality. However, we are able to prove that T_X is stable, see Theorem B below. To conclude the proof of Theorem A, we show that, in the setting of Theorem A, we must have $\pi_1^{\text{ét}}(X_{\text{reg}}) = \{1\}$ (see Theorem 7.1). Note that in Theorem A, we do not require \mathcal{X} to be \mathbb{Q} -Gorenstein as this condition is automatically satisfied, see Lemma 7.6.

As we explained above, it seems difficult to obtain a full decomposition theorem using our strategy without further assumptions on the singularities of X . However, we are still able to produce a splitting of some quasi-étale cover of X where each non-abelian factor has a stable tangent bundle, and possesses the same algebra of reflexive holomorphic forms as the one of an irreducible, simply-connected, Calabi-Yau or symplectic manifold of the same dimension.

Theorem B. *Let X be a normal complex projective variety with klt singularities. Suppose that $K_X \equiv 0$. Suppose furthermore that there exists a projective morphism with connected fibers $f: \mathcal{X} \rightarrow \Delta$ from a normal analytic space \mathcal{X} whose canonical divisor $K_{\mathcal{X}}$ is \mathbb{Q} -Cartier onto the complex open unit disk Δ such that $X \cong f^{-1}(0)$ and such that $f^{-1}(t)$ is smooth for $t \neq 0$. Then, there exists an abelian variety A as well as a projective variety X' with canonical singularities, a finite cover*

$$A \times X' \rightarrow X,$$

étale in codimension one, and a decomposition

$$X' \cong \prod_{i \in I} Y_i \times \prod_{j \in J} Z_j$$

of X' into normal projective varieties with trivial canonical class, such that the following holds.

- (1) The sheaf T_{Y_i} is stable with respect to any polarization and one has $h^0(Y_i, \Omega_{Y_i}^{[q]}) = 0$ for all numbers $0 < q < \dim Y_i$.
- (2) The sheaf T_{Z_j} is stable with respect to any polarization and there exists a reflexive 2-form $\sigma \in H^0(Z_j, \Omega_{Z_j}^{[2]})$ such that σ is everywhere non-degenerate on the smooth locus of Z_j , and such that the exterior algebra of global reflexive forms is generated by σ .

Remark 1.3. If $\pi : \tilde{Z}_j \rightarrow Z_j$ is a \mathbb{Q} -factorial terminalization of the variety Z_j from Theorem B, then it follows from [Nam06, Corollary 2] that \tilde{Z}_j is a *smooth* symplectic variety, but not necessarily irreducible, since the varieties Z_j are smoothable by construction (see Section 6). In particular, Z_j admits a symplectic resolution. Note that the existence of π is established in [BCHM10, Corollary 1.4.3].

In fact, a little more can be said about the factors of X' in the decomposition given by Theorem B above. We refer to Section 6.2 for partial results from the point of view of holonomy representation, and for a proof of Conjecture 1.4 below assuming that a weak analogue of Beauville-Bogomolov decomposition theorem holds.

Conjecture 1.4. *Let X be a normal complex projective variety with klt singularities and $K_X \equiv 0$. Suppose that T_X is stable with respect to some polarization. Then there exists a quasi-étale cover $Y \rightarrow X$ such that either Y is an abelian variety, or it splits as a product of copies of a single Calabi-Yau (resp. irreducible symplectic) variety.*

Note that a positive answer to the conjecture above implies that Theorem A holds without the assumption that X is smooth in codimension 2.

The following result is an immediate consequence of Theorem B.

Corollary C. *Let X be a normal complex projective variety with klt singularities and $K_X \equiv 0$. Suppose that $\pi_1^{\text{ét}}(X_{\text{reg}}) = \{1\}$. Suppose furthermore that there exists a projective morphism with connected fibers $f : \mathcal{X} \rightarrow \Delta$ from a normal analytic space \mathcal{X} whose canonical divisor $K_{\mathcal{X}}$ is \mathbb{Q} -Cartier onto the complex open unit disk Δ such that $X \cong f^{-1}(0)$ and such that $f^{-1}(t)$ is smooth for $t \neq 0$. Then there exists a decomposition of X into a product of irreducible, Calabi-Yau and symplectic varieties.*

Structure of the paper. Section 2 is mainly devoted to setting up the basic notation. We have also gathered a number of facts and basic results which will later be used in the proofs. Sections 3, 4 and 5 consist of technical preparations. In Section 3, we recall some results on deformations of Kähler-Einstein metrics on smoothable singular spaces with numerically trivial canonical class. In Section 4, we establish a structure result for families of mildly singular varieties with trivial canonical class. In Section 5, we use results from [GGK17] to analyze the stability of the tangent sheaf of smoothable singular spaces with numerically trivial canonical class. Section 6 is devoted to the proof of Theorem B. In Section 7, we prove Theorem A. Finally, in Section 8, we give examples of smoothable (irreducible) Calabi-Yau and symplectic varieties. We have also collected examples which illustrate to what extent our results are sharp.

Acknowledgements. The project started while the authors were visiting the Freiburg Institute for Advanced Studies. The authors would like to thank this institution for its support and Stefan Kebekus for the invitation. They would like to thank Benoît Claudon, Daniel Greb, Stefan Kebekus, Mihai Păun, and Song Sun for interesting discussions concerning the content of this paper.

The first author was partially supported by the project Foliage of Agence Nationale de la Recherche, under agreement ANR-16-CE40-0008-01.

The second author is partially supported by NSF Grant DMS-1510214.

2. NOTATION, CONVENTIONS, AND BASIC FACTS

2.1. Global Convention. Throughout the paper we work over the field \mathbb{C} of complex numbers.

A *variety* is a reduced and irreducible scheme of finite type over \mathbb{C} . An *analytic variety* is a reduced and irreducible analytic space. Given a scheme X , we denote by X^{an} the associated analytic space, equipped with the Euclidean topology.

Given a scheme or an analytic space X , we denote by X_{reg} its smooth locus.

2.2. Reflexive differential forms. Given a normal (analytic) variety X , we denote the sheaf of Kähler differentials by Ω_X^1 . If $0 \leq p \leq \dim X$ is any number, write $\Omega_X^{[p]} := (\Omega_X^p)^{**}$. The tangent sheaf will be denoted by $T_X := (\Omega_X^1)^*$.

2.3. Quasi-étale covers.

Definition 2.1. A morphism $\gamma: Y \rightarrow X$ between normal (analytic) varieties is called a *quasi-étale cover* if γ is finite and étale in codimension one.

Remark 2.2. Let $\gamma: Y \rightarrow X$ be a quasi-étale cover. By the Nagata-Zariski purity theorem, γ branches only on the singular set of X . In particular, we have $\gamma^{-1}(X_{\text{reg}}) \subset Y_{\text{reg}}$.

2.4. Stability. The word “stable” will always mean “slope-stable with respect to a given polarization”.

Definition 2.3 ([GKP11, Definition 7.2]). Let X be a normal projective (analytic) variety of dimension n , and let \mathcal{G} be a reflexive coherent sheaf. We call \mathcal{G} *strongly stable*, if for any quasi-étale cover $\gamma: Y \rightarrow X$, and for any choice of ample divisors H_1, \dots, H_{n-1} on Y , the reflexive pull-back $\gamma^{[*]}\mathcal{G}$ is stable with respect to (H_1, \dots, H_{n-1}) .

2.5. Smoothings. We will use the following notation.

Notation 2.4. Let $f: \mathcal{X} \rightarrow T$ be a morphism (resp. holomorphic map) of schemes (resp. analytic spaces). We will denote by \mathcal{X}_t the fiber of f over $t \in T$.

Definition 2.5. Let X be a compact analytic space. A *smoothing of X* is a flat proper holomorphic map $f: \mathcal{X} \rightarrow \Delta$, where \mathcal{X} is an analytic space and Δ is the complex open unit disk, such that $\mathcal{X}_0 \cong X$ and \mathcal{X}_t is smooth for any $t \neq 0$. A *smoothing of a proper scheme X* is a smoothing of the associated analytic space X^{an} . Let $f: \mathcal{X} \rightarrow \Delta$ be a smoothing of a compact analytic space or a proper scheme. We say that f is a *projective smoothing* if f is a projective map. If \mathcal{X} is normal, then we say that f is a *\mathbb{Q} -Gorenstein smoothing* if $K_{\mathcal{X}/\Delta}$ is \mathbb{Q} -Cartier.

Let X be a proper scheme. A *smoothing of X over an algebraic curve* is a flat proper morphism $f: \mathcal{X} \rightarrow C$, where \mathcal{X} is a scheme and C is a connected algebraic curve, such that $\mathcal{X}_{t_0} \cong X$ for some point t_0 on C and \mathcal{X}_t is smooth for any $t \neq t_0$. A smoothing $f: \mathcal{X} \rightarrow C$ of X over an algebraic curve is said to be *projective* if f is a projective morphism. If \mathcal{X} is normal, then we say that f is a *\mathbb{Q} -Gorenstein smoothing* if $K_{\mathcal{X}/C}$ is \mathbb{Q} -Cartier.

The following elementary facts will be used throughout.

Fact 2.6. Let \mathcal{X} be an analytic space (resp. a scheme), and let $f: \mathcal{X} \rightarrow T$ be a flat holomorphic map (resp. morphism) with connected fibers onto a smooth connected analytic (resp. algebraic) curve T . If \mathcal{X}_t is normal for any $t \in T$, then so is \mathcal{X} by [Gro65, Corollaire 5.12.7] and [DG67, Théorème 1.2]. In particular, \mathcal{X} is reduced and locally irreducible (resp. reduced and irreducible).

Fact 2.7. Let \mathcal{X} be a variety, and let $X \subset \mathcal{X}$ be a Cartier divisor. If X is regular at x , then so is \mathcal{X} .

Fact 2.8. In the setup of Fact 2.6, suppose that \mathcal{X} is a normal variety, and that $K_{\mathcal{X}}$ is \mathbb{Q} -Cartier. If \mathcal{X}_t has klt singularities for some point t on C , then \mathcal{X} has klt singularities in a neighborhood of \mathcal{X}_t by inversion of adjunction (see [Kol97, Corollary 7.6]). This implies that \mathcal{X} has rational singularities in a neighborhood of \mathcal{X}_t (see [KMM87, Theorem 1.3.6]).

Remark 2.9. In the setup of Fact 2.6, suppose that \mathcal{X} is a normal variety, and that $K_{\mathcal{X}}$ is \mathbb{Q} -Cartier. If \mathcal{X}_{t_0} has lc singularities for some point t_0 on C and \mathcal{X}_t has canonical singularities for $t \neq t_0$, then $\mathcal{X} \times_C C_1$ has canonical singularities for any finite morphism $C_1 \rightarrow C$ from a smooth algebraic curve C_1 by [Kar00, Theorem 2.5]. Note that the proof of [Kar00, Theorem 2.5] relies on [BCHM10, Theorem 1.2]. However, we will not need this stronger statement.

Let X be a normal projective variety. If X admits a projective \mathbb{Q} -Gorenstein smoothing over an algebraic curve then X obviously admits a projective \mathbb{Q} -Gorenstein smoothing. The main result of the present section is a partial converse to this observation. See Lemma 2.10 and Proposition 2.14 for precise statements.

Lemma 2.10. *Let X be a normal projective variety with Gorenstein singularities. If X admits a projective smoothing, then X admits a projective \mathbb{Q} -Gorenstein smoothing over an algebraic curve.*

Proof. Let $f: \mathcal{X} \rightarrow \Delta$ be a projective smoothing of X . We may assume without loss of generality that $\mathcal{X} \subset \mathbb{P}^N \times \Delta$ for some positive integer N . Let $\Delta \rightarrow \text{Hilb}(\mathbb{P}^N)^{\text{an}}$ be the universal holomorphic map, and let H any irreducible component of $\text{Hilb}(\mathbb{P}^N)$ such that H^{an} contains the image of Δ . Let also $U \subset \mathbb{P}^N \times H$

be the universal family. Note that $\dim H \geq 1$ and that the generic fiber of the natural morphism $U \rightarrow H$ is smooth. Replacing H by a Zariski open neighborhood of $[X]$, if necessary, we may assume that for any $t \in H$, U_t is normal and Cohen-Macaulay by [Gro66, Théorèmes 12.2.1 et 12.2.4]. Denote by $\omega_{U/H}$ the dualizing sheaf of the Cohen-Macaulay morphism $U \rightarrow H$ (see [Con00, 3.5]). By [Con00, Theorem 3.5.6], for any closed point $t \in H$, we have $\omega_{U/H}|_{U_t} \cong \omega_{U_t}$ where ω_{U_t} is the dualizing sheaf of $U_t \rightarrow \text{pt}$. Set $t_0 := [X] \in H$. Because $\omega_{U_{t_0}}$ is invertible by assumption, we see that $\omega_{U/H}$ is invertible in a Zariski open neighborhood of X . By shrinking H if necessary, we may therefore assume that $\omega_{U/H}$ is invertible. Let $C \rightarrow H$ be a smooth curve passing through $[X]$ and a general point, and set $\mathcal{Y} := C \times_H U$. It comes with a natural morphism $g: \mathcal{Y} \rightarrow C$. Note that g is a projective smoothing of X over C . By Fact 2.6, we know that \mathcal{Y} is normal. Applying [Con00, Theorem 3.5.6] again, we see that $\omega_{\mathcal{Y}/C}$ is invertible. Moreover, its restriction to the locus $\mathcal{Y}_{\text{reg}}^\circ \subset \mathcal{Y}_{\text{reg}}$ where g is smooth is $\omega_{\mathcal{Y}_{\text{reg}}^\circ/H}$. On the other hand, $\omega_{\mathcal{Y}_{\text{reg}}^\circ/H} \cong \mathcal{O}_{\mathcal{Y}_{\text{reg}}^\circ}(K_{\mathcal{Y}_{\text{reg}}^\circ/H}) \cong \mathcal{O}_{\mathcal{Y}}(K_{\mathcal{Y}/H})|_{\mathcal{Y}_{\text{reg}}^\circ}$. It follows that $\omega_{\mathcal{Y}/H} \cong \mathcal{O}_{\mathcal{Y}}(K_{\mathcal{Y}/H})$ since both are reflexive sheaves. Hence $K_{\mathcal{Y}/H}$ is Cartier, completing the proof of the lemma. \square

The same argument used in the proof of Lemma 2.10 above shows that the following holds.

Lemma 2.11. *Let X be a normal projective variety. If X admits a projective smoothing, then X admits a projective smoothing over an algebraic curve.*

The proof of [Laz04, Proposition 1.4.14] apply in the analytic setting to show that the following holds.

Lemma 2.12. *Let $f: \mathcal{X} \rightarrow T$ be a projective holomorphic map (resp. morphism) of analytic spaces (resp. schemes), and let \mathcal{L} be a line bundle on \mathcal{X} . Suppose furthermore that f is surjective. Then the set of points t on T such that $\mathcal{L}|_{\mathcal{X}_t}$ is not nef is a countable union of analytic subsets.*

We end the preparation for the proof of Proposition 2.14 with the following observation.

Lemma 2.13. *Let $f: \mathcal{X} \rightarrow T$ be a flat projective holomorphic map (resp. morphism) of analytic spaces (resp. schemes). Suppose that \mathcal{X} is normal, T is smooth, and that \mathcal{X}_t is connected with klt singularities for any point $t \in T$. Suppose furthermore that $K_{\mathcal{X}/T}$ is \mathbb{Q} -Cartier. If $K_{\mathcal{X}_{t_0}} \equiv 0$ for some point $t_0 \in T$, then there exists a Zariski open cover $(T^\alpha)_{\alpha \in A}$ of T such that $K_{\mathcal{X}^\alpha/T^\alpha}$ is torsion where $\mathcal{X}^\alpha := f^{-1}(T^\alpha)$. In particular, $K_{\mathcal{X}_t}$ is torsion for all $t \in T$.*

Proof. Let m_0 be a positive integer such that $m_0 K_{\mathcal{X}/T}$ is a Cartier divisor. Applying Lemma 2.12 in the analytic setting or [Laz04, Proposition 1.4.14] in the algebraic setting to $\pm m_0 K_{\mathcal{X}/T}$, we see that the set of points $t \in T$ such that $K_{\mathcal{X}_t} \not\equiv 0$ is a countable union of proper Zariski closed subsets. If $K_{\mathcal{X}_t} \equiv 0$, then $K_{\mathcal{X}_t}$ is torsion by [Nak04, Corollary V 4.9]. Note also that we have $m_0 K_{\mathcal{X}_t} \sim_{\mathbb{Z}} m_0 K_{\mathcal{X}/T}|_{\mathcal{X}_t}$ by the adjunction formula. Let now m be a positive integer. Because the functions $t \mapsto h^0(\mathcal{X}_t, \mathcal{O}_{\mathcal{X}_t}(\pm mm_0 K_{\mathcal{X}_t}))$ are upper semicontinuous in the Zariski topology on T (see [BS76, Chapter III, Theorem 4.12] in the analytic setting or [Har77, Theorem 12.8] in the algebraic setting), the set of points t on T such that $mm_0 K_{\mathcal{X}_t} \sim_{\mathbb{Z}} 0$ is closed. It follows that there exists a positive integer m_1 such that $m_1 m_0 K_{\mathcal{X}_t} \sim_{\mathbb{Z}} 0$ for all $t \in T$. From [BS76, Chapter III, Theorem 4.12] in the analytic setting or [Har77, Corollary 12.9] in the algebraic setting, we see that $f_* \mathcal{O}_{\mathcal{X}}(m_1 m_0 K_{\mathcal{X}/T})$ is a line bundle. Let $(T^\alpha)_{\alpha \in A}$ be a Zariski open affine cover of T such that $f_* \mathcal{O}_{\mathcal{X}}(m_1 m_0 K_{\mathcal{X}/T})|_{T^\alpha} \cong \mathcal{O}_{T^\alpha}$ for all $\alpha \in A$. Set $\mathcal{X}^\alpha := f^{-1}(T^\alpha)$, and let $t \in T^\alpha$. Because the formation of $f_* \mathcal{O}_{\mathcal{X}}(m_1 m_0 K_{\mathcal{X}/T})$ commutes with base change (see *loc. cit.*), any non-zero section of $m_1 m_0 K_{\mathcal{X}/T}|_{\mathcal{X}_t}$ extends to a global section of $m_1 m_0 K_{\mathcal{X}/T}|_{\mathcal{X}^\alpha}$ that is nowhere vanishing in a neighborhood of \mathcal{X}_t . Refining the cover, if necessary, we conclude that $m_1 m_0 K_{\mathcal{X}/T}|_{\mathcal{X}^\alpha} \cong \mathcal{O}_{\mathcal{X}^\alpha}$. This completes the proof of the lemma. \square

Proposition 2.14. *Let X be a normal projective variety with klt singularities. Suppose that $K_X \equiv 0$. If X admits a projective \mathbb{Q} -Gorenstein smoothing, then there exist a normal projective variety Y , a quasi-étale cover $Y \rightarrow X$, and a projective smoothing $\mathcal{Y} \rightarrow C$ of Y over an algebraic curve such that $K_{\mathcal{Y}/C} \sim_{\mathbb{Z}} 0$.*

Proof. Let $f: \mathcal{X} \rightarrow \Delta$ be a projective smoothing of X . Applying Lemma 2.13, we see that $K_{\mathcal{X}/\Delta}$ is torsion.

By [KM98, Definition 2.52 and Lemma 2.53], there exists a normal analytic variety \mathcal{Y} and a finite cover $\gamma: \mathcal{Y} \rightarrow \mathcal{X}$, étale over \mathcal{X}_{reg} , such that $K_{\mathcal{Y}/\Delta} \sim_{\mathbb{Z}} 0$. Note that $\gamma_t: \mathcal{Y}_t \rightarrow \mathcal{X}_t$ is étale for any $t \neq 0$. In particular, \mathcal{Y}_t is smooth if $t \neq 0$. Note also that \mathcal{Y}_0 is normal, and that $\gamma_0: \mathcal{Y}_0 \rightarrow \mathcal{X}_0$ is a quasi-étale

cover. Applying [Kol97, Proposition 3.16] to γ_0 , we see that $Y := \mathcal{Y}_0$ has klt singularities. Moreover, by the adjunction formula, we have $K_Y \sim_{\mathbb{Z}} 0$.

By Lemma 2.10, Y admits a projective \mathbb{Q} -Gorenstein smoothing over an algebraic curve. Arguing as above, we see that there exist a normal projective variety Y_1 as well as a quasi-étale cover $\gamma_1: Y_1 \rightarrow Y$, and a projective smoothing $\mathcal{Y}_1 \rightarrow C_1$ of Y_1 over an algebraic curve such that $K_{\mathcal{Y}_1/C_1} \sim_{\mathbb{Z}} 0$, completing the proof of the proposition. \square

3. KÄHLER-EINSTEIN METRICS ON SMOOTHABLE SPACES

In this section, we work in the following setting, referred to later as the analytic setting.

3.1. The analytic setting. Let $f: \mathcal{X} \rightarrow \Delta$ be a projective smoothing of a normal projective (analytic) variety X such that $K_{\mathcal{X}/\Delta} \sim_{\mathbb{Z}} 0$. Suppose moreover that X has canonical singularities. Let \mathcal{L} be a relatively ample line bundle on \mathcal{X} , and set $\mathcal{L}_t := \mathcal{L}|_{\mathcal{X}_t}$. Given $t \neq 0$, we denote by g_t the unique Ricci-flat Kähler metric on \mathcal{X}_t whose fundamental form ω_t satisfies $[\omega_t] = [c_1(\mathcal{L}_t)] \in H^2(\mathcal{X}_t, \mathbb{R})$. The existence of g_t is established in [Yau78]. We denote by g_0 the Ricci-flat Kähler metric on X_{reg} given by [EGZ09, Theorem 7.5] applied to $(\mathcal{X}_0, \mathcal{L}_0)$.

3.2. Smooth convergence on X_{reg} . In the setting described above, let $\Phi: X_{\text{reg}} \times \Delta \rightarrow \mathcal{X}$ be a smooth embedding such that $\Phi(x, t) \in \mathcal{X}_t$ for any $(x, t) \in X_{\text{reg}} \times \Delta$ and $\Phi|_{X_{\text{reg}} \times \{0\}} = \text{Id}_{X_{\text{reg}}}$ (see [RZ11b, p. 1547]). Let us write $\Phi_t := \Phi|_{X_{\text{reg}} \times \{t\}}: X_{\text{reg}} \rightarrow \mathcal{X}_t$. If I_t denotes the complex structure on \mathcal{X}_t for $t \neq 0$, and X_{reg} for $t = 0$, then it is not hard to see that $\Phi_t^* I_t$ converges to I_0 in the $\mathcal{C}_{\text{loc}}^\infty(X_{\text{reg}})$ -topology. The following theorem is due to Rong-Zhang (see [RZ11a, Theorem 1.4]).

Theorem 3.1 ([RZ11a]). *The family of Riemannian metrics $\Phi_t^* g_t$ converges to g_0 when $t \rightarrow 0$, in the $\mathcal{C}_{\text{loc}}^\infty(X_{\text{reg}})$ -topology.*

Sketch of proof. We recall the main arguments of the proof for the convenience of the reader.

The first step is to show that ω_t and $\omega_{\text{FS},t}$ differ by the dd^c of an uniformly bounded potential φ_t , that we can assume to be normalized by $\int_{X_t} \varphi_t \omega_t^n = 0$. Here, $\omega_{\text{FS},t} = \omega_{\text{FS}}|_{\mathcal{X}_t}$ under the embedding $\mathcal{X} \hookrightarrow \mathbb{P}^N \times \Delta$ given by sections of \mathcal{L} . This is proved in [RZ11a, Lemma 3.1] using Moser iteration given that the Sobolev and Poincaré constants of $(\mathcal{X}_t, \omega_t)$ are bounded, which in turn is a consequence of the diameter estimate [RZ11a, Theorems B.1 and 2.1].

The L^∞ -estimate on φ_t combined with Chern-Lu inequality implies that $\omega_t \geq C^{-1} \omega_{\text{FS},t}$ for some uniform $C > 0$ (see the proof of [RZ11a, Lemma 3.2]). This enables to get estimates at any order on $\Phi_t^* \varphi_t$ over compact subsets of X_{reg} . Therefore, one can extract smooth sequential limits φ_∞ of $\Phi_t^* \varphi_t$ over X_{reg} : φ_∞ is bounded on the whole X_{reg} and as $\Phi_t^* I_t$ converges to I_0 , φ_∞ satisfies the same Monge-Ampère equation as φ_0 on X_{reg} , where φ_0 is the normalized potential of ω_0 with respect to $\omega_{\text{FS},0}$. By the choice of the normalization, φ_∞ and φ_0 coincide, and therefore $\Phi_t^* \varphi_t$ converges locally smoothly to φ_0 , which concludes the proof. \square

Remark 3.2. An important observation is that if \mathcal{X}_t admits another complex structure J_t compatible with g_t (in the sense that J_t is unitary and parallel with respect to g_t), then one can extract sequences $t_j \rightarrow 0$ such that $\Phi_{t_j}^* J_{t_j}$ converges locally smoothly to a complex structure J_0 over X_{reg} which is compatible with respect to g_0 . Indeed, $\Phi_t^* J_t$ is almost g_0 -unitary and g_0 -parallel in the sense that the tensors $(\Phi_t^* J_t)^{*0} (\Phi_t^* J_t) - \text{Id}$ and $\nabla^{g_0} (\Phi_t^* J_t)$ converge to zero on X_{reg} , where $*_0$ denotes the adjoint with respect to g_0 . The first property gives order 0 estimates while the second one enables to get higher order ones. Arzela-Ascoli theorem combined with a diagonal argument yield the result.

3.3. Identification of the Gromov-Hausdorff limit. One can actually understand the global limit of (\mathcal{X}_t, g_t) , but this requires considerable more work and will not be used in the following. The next result is essentially contained in [SSY16, Section 3] (see also [LWX14, Section 4.2]). As the context is slightly different here since we deal with Calabi-Yau manifolds rather than log-Fano manifolds, we will recall their main arguments and point out the slight changes to operate.

Theorem 3.3 ([SSY16, LWX14]). *In the standard setting, the metric spaces $(\mathcal{X}_t, \omega_t)$ converge in the Gromov-Hausdorff sense to (X, d) when $t \rightarrow 0$ for some metric d on X . Moreover, the convergence is smooth on X_{reg} , and the restriction of d to X_{reg} is induced by the Riemannian metric g_0 .*

Sketch of proof. The fundamental results of [DS14] give the convergence in the sense of the statement above to a projective variety endowed with a singular Kähler-Einstein metric, but the key point is to identify that variety with the central fiber X of our family.

As in the proof of Theorem 3.1, one can write $\omega_t = \omega_{\text{FS},t} + dd^c \varphi_t$ with $\|\varphi_t\|_{L^\infty} \leq C$, C being independent of $t \in \Delta^*$. This has been seen to imply the estimate $\omega_t \geq C^{-1} \omega_{\text{FS},t}$. Also, it is easy but important to notice that the bound on the potential remains valid after Veronese re-embeddings (see [SSY16, Lemma 3.3]).

The next step consists in comparing the two different embeddings of \mathcal{X}_t into a large \mathbb{P}^M , one being given by a Veronese embedding (using a large power of \mathcal{L} , say k) and the other one being an embedding by L^2 sections of $\mathcal{L}_t^{\otimes k}$ with respect to ω_t . In the following, we may assume that the first re-embedding is the identity while the second one will be denoted by i_t . So in \mathbb{P}^M , we have two isomorphic varieties, namely \mathcal{X}_t and $i_t(\mathcal{X}_t)$. The variety $i_t(\mathcal{X}_t)$ can be obtained from \mathcal{X}_t by a transformation $g_t \in \text{PGL}(\mathbb{C}^{M+1})$ that sends a basis $(s_{0,t}, \dots, s_{M,t})$ of $H^0(\mathcal{X}_t, \mathcal{L}_t^{\otimes k})$ to a L^2 -orthonormal basis $(\sigma_{0,t}, \dots, \sigma_{M,t})$, where the first basis of sections is obtained by the standard embedding induced by $\mathcal{L}^{\otimes k}$. Then one can prove using the estimates on $\|\varphi_t\|_{L^\infty}, \int_{\mathcal{X}_t} \omega_t^n, \text{tr}_{\omega_t}(\omega_{\text{FS},t})$ that g_t evolves in a compact subset of $\text{PGL}(\mathbb{C}^{M+1})$ (see proof of [SSY16, Theorem 3.1]).

The last step invokes the main result of [DS14] that guarantees in this context that one can choose k and a sequence $t_j \rightarrow 0$ such that the re-embedding $i_{t_j}(X_{t_j})$ converges both in the sense of cycle and in the Gromov-Hausdorff sense to a projective variety W (for some metric on W). Up to extract a subsequence, one can assume that g_{t_j} converges to $g \in \text{PGL}(\mathbb{C}^{M+1})$ so that $W = g(X)$ as projective varieties. The rest is a consequence of [DS14]. \square

4. RELATIVE ALBANESE MORPHISM

The Albanese morphism, that is, the universal morphism to an abelian variety (see [Ser01]), is one important tool in the study of varieties with trivial canonical divisor. If X is a projective variety with rational singularities, recall from [Kaw85, Lemma 8.1] that $\text{Pic}^\circ(X)$ is an abelian variety, and that $\text{Alb}(X) \cong (\text{Pic}^\circ(X))^\vee$. Moreover, the Albanese morphism $a_X : X \rightarrow \text{Alb}(X)$ is induced by the universal line bundle. In particular, $\dim \text{Alb}(X) = h^1(X, \mathcal{O}_X)$. The following result of Kawamata describes the Albanese map of X (see also [Bri10, Section 3]).

Proposition 4.1 ([Kaw85, Proposition 8.3]). *Let X be a normal projective variety X with canonical singularities. Assume that K_X is numerically trivial. Then K_X is torsion, the Albanese map $a_X : X \rightarrow \text{Alb}(X)$ is surjective and has the structure of an étale-trivial fiber bundle. More precisely, the following holds. The neutral component A of the automorphism group $\text{Aut}(X)$ of X is an abelian variety. It acts on $\text{Alb}(X)$ by translation, compatibly with its action on X . The induced morphism $A \rightarrow \text{Alb}(X)$ is an isogeny and the fiber product over $\text{Alb}(X)$ decomposes as a product*

$$X \times_{\text{Alb}(X)} A \cong F \times A,$$

where F is a fiber of a_X .

In this section we extend Proposition 4.1 to the relative setting.

Proposition 4.2. *Let \mathcal{X} be a normal variety, and let $f : \mathcal{X} \rightarrow C$ be a flat projective morphism with connected fibers onto a smooth connected algebraic curve C . Suppose that \mathcal{X}_t has klt singularities for any point $t \in C$. Suppose furthermore that $K_{\mathcal{X}/C} \sim_{\mathbb{Z}} 0$.*

- (1) *The neutral component $\text{Aut}^0(\mathcal{X}_t)$ of the automorphism group of \mathcal{X}_t is an abelian variety, and the algebraic groups $\text{Aut}^0(\mathcal{X}_t)$ fit together to form an abelian scheme \mathcal{A} over C .*
- (2) *Suppose that f has a section, and consider the morphism*

$$a_{\mathcal{X}/C} : \mathcal{X} \rightarrow (\text{Pic}^\circ(\mathcal{X}/C))^\vee$$

induced by the universal line bundle. Let $0 : C \rightarrow (\text{Pic}^\circ(\mathcal{X}/C))^\vee$ denotes the neutral section, and set $\mathcal{Y} = a_{\mathcal{X}/C}^{-1}(0)$. Then $\mathcal{Y} \rightarrow C$ is a flat projective morphism with normal connected fibers, and there exists a relative isogeny $\mathcal{A} \rightarrow (\text{Pic}^\circ(\mathcal{X}/C))^\vee$ such that

$$\mathcal{X} \times_{(\text{Pic}^\circ(\mathcal{X}/C))^\vee} \mathcal{A} \cong \mathcal{Y} \times_C \mathcal{A}.$$

Before we give the proof of Proposition 4.2, we need the following auxiliary result.

Lemma 4.3. *Let $f: \mathcal{X} \rightarrow C$ be a flat projective morphism with connected fibers from a normal variety \mathcal{X} onto a smooth connected algebraic curve C . Suppose that \mathcal{X} has rational singularities. Then $h^i(\mathcal{X}_t, \mathcal{O}_{\mathcal{X}_t})$ is independent of $t \in C$ for any integer $i \geq 0$.*

Proof. Let \bar{C} be a smooth compactification of C , and let $\bar{\mathcal{X}}$ be a projective compactification of \mathcal{X} such that f extends to a morphism $\bar{f}: \bar{\mathcal{X}} \rightarrow \bar{C}$. Finally, let $\nu: \widehat{\mathcal{X}} \rightarrow \bar{\mathcal{X}}$ be a desingularization of $\bar{\mathcal{X}}$. Set $\widehat{f} := \bar{f} \circ \nu$. It follows from [Kol86, Corollary 3.9] that the sheaves $R^i \widehat{f}_* \mathcal{O}_{\widehat{\mathcal{X}}}$ are torsion free, and hence locally free. The relative Leray spectral sequence gives $R^i \widehat{f}_* \mathcal{O}_{\widehat{\mathcal{X}}|C} \cong R^i f_* \mathcal{O}_{\mathcal{X}}$ using the assumption that \mathcal{X} has rational singularities. This in turn implies that $h^i(\mathcal{X}_t, \mathcal{O}_{\mathcal{X}_t})$ is independent of $t \in C$ by a theorem of Grothendieck (see [Har77, Theorem 12.11]). \square

Proof of Proposition 4.2. Note that \mathcal{X}_t has canonical singularities since $K_{\mathcal{X}_t} \sim_{\mathbb{Z}} 0$ is a Cartier divisor by the adjunction formula. Write $n := \dim \mathcal{X} - 1$. From Lemma 4.3 and Fact 2.8, we see that $h^{n-1}(\mathcal{X}_t, \mathcal{O}_{\mathcal{X}_t})$ is independent of $t \in C$. On the other hand, we have $h^{n-1}(\mathcal{X}_t, \mathcal{O}_{\mathcal{X}_t}) = h^0(\mathcal{X}_t, \Omega_{\mathcal{X}_t}^{[n-1]})$ by [GKP11, Proposition 6.9]. It follows that $h^0(\mathcal{X}_t, T_{\mathcal{X}_t})$ is independent of $t \in C$ since $\Omega_{\mathcal{X}_t}^{[n-1]} \cong T_{\mathcal{X}_t}$. Note that $h^{n-1}(\mathcal{X}_t, \mathcal{O}_{\mathcal{X}_t}) = h^1(\mathcal{X}_t, \mathcal{O}_{\mathcal{X}_t})$ by Serre duality, so that $h^0(\mathcal{X}_t, T_{\mathcal{X}_t}) = h^1(\mathcal{X}_t, \mathcal{O}_{\mathcal{X}_t})$. By [GP11, Exposé VI_B, Proposition 1.6], the group scheme $\text{Aut}(\mathcal{X}/C)$ is then smooth over C . Note that the existence of $\text{Aut}(\mathcal{X}/C)$ is guaranteed by [Gro95b]. Let $t \in C$, and denote by $\text{Aut}^0(\mathcal{X}_t)$ the neutral component of the automorphism group $\text{Aut}(\mathcal{X}_t)$ of \mathcal{X}_t . We have $\dim \text{Aut}^0(\mathcal{X}_t) = h^0(\mathcal{X}_t, T_{\mathcal{X}_t}) = h^1(\mathcal{X}_t, \mathcal{O}_{\mathcal{X}_t})$. We claim that $\text{Aut}^0(\mathcal{X}_t)$ is an abelian variety. If not, by a theorem of Chevalley, $\text{Aut}^0(\mathcal{X}_t)$ contains an algebraic subgroup isomorphic either to \mathbb{G}_a or to \mathbb{G}_m , and hence, \mathcal{X} is uniruled. On the other hand, $\kappa(\mathcal{X}_t) = 0$ since $K_{\mathcal{X}_t} \sim_{\mathbb{Z}} 0$ and \mathcal{X}_t has canonical singularities, yielding a contradiction. Now, recall from [GP11, Exposé VI_B, Théorème 3.10] that the algebraic groups $\text{Aut}^0(\mathcal{X}_t)$ fit together to form a group scheme \mathcal{A} over C . Note that \mathcal{A} is quasi-projective over C (see [Gro95b]). Applying [Gro66, Corollaire 15.7.11], we see that \mathcal{A} is proper, and hence projective over C . In particular, \mathcal{A} is an abelian scheme over C .

Suppose from now on that f has a section $s: C \rightarrow \mathcal{X}$, and consider the relative Picard scheme $\text{Pic}(\mathcal{X}/C)$ whose existence is guaranteed by [Gro95a, Théorème 3.1]. By [Kaw85, Lemma 8.1], $\text{Pic}^\circ(\mathcal{X}_t)$ is an abelian variety for any point t on C . As above, the algebraic groups $\text{Pic}^\circ(\mathcal{X}_t)$ fit together to form a group scheme $\text{Pic}^\circ(\mathcal{X}/C)$ over C . Note that $\text{Pic}^\circ(\mathcal{X}/C) \subset \text{Pic}(\mathcal{X}/C)$ is an open subscheme, quasi-projective over C by [BLR90, Theorem 5]. Using [Gro66, Corollaire 15.7.11] again, we conclude that it is projective over C . Let $(\text{Pic}^\circ(\mathcal{X}/C))^\vee$ be the dual abelian scheme (see [MFK94, Corollary 6.8]), and consider the morphism

$$a_{\mathcal{X}/C}: \mathcal{X} \rightarrow (\text{Pic}^\circ(\mathcal{X}/C))^\vee$$

induced by the universal line bundle. Consider also the action $\varphi: \mathcal{A} \times_C \mathcal{X} \rightarrow \mathcal{X}$, and the second projection $p_2: \mathcal{A} \times_C \mathcal{X} \rightarrow \mathcal{X}$. By the rigidity lemma ([MFK94, Proposition 6.1]), there exists a morphism

$$\psi: \mathcal{A} \times_C (\text{Pic}^\circ(\mathcal{X}/C))^\vee \rightarrow (\text{Pic}^\circ(\mathcal{X}/C))^\vee$$

such that the square

$$\begin{array}{ccc} \mathcal{A} \times_C \mathcal{X} & \xrightarrow{\varphi} & \mathcal{X} \\ (\text{Id} \times a_{\mathcal{X}/C}) \downarrow & & \downarrow a_{\mathcal{X}/C} \\ \mathcal{A} \times_C (\text{Pic}^\circ(\mathcal{X}/C))^\vee & \xrightarrow{\psi} & (\text{Pic}^\circ(\mathcal{X}/C))^\vee \end{array}$$

is commutative. It follows that \mathcal{A} acts on $(\text{Pic}^\circ(\mathcal{X}/C))^\vee$ over C . By Proposition 4.1, the induced morphism

$$\mathcal{A} \rightarrow (\text{Pic}^\circ(\mathcal{X}/C))^\vee$$

is a relative isogeny. Let $0: C \rightarrow (\text{Pic}^\circ(\mathcal{X}/C))^\vee$ denotes the neutral section, and set $\mathcal{Y} := a_{\mathcal{X}/C}^{-1}(0)$. Using Proposition 4.1 again, one readily checks that

$$\mathcal{X} \times_{(\text{Pic}^\circ(\mathcal{X}/C))^\vee} \mathcal{A} \cong \mathcal{Y} \times_C \mathcal{A}.$$

The proposition then follows easily. \square

We end this section with an extension result for differential forms. We feel that this result might be of independent interest.

Proposition 4.4. *Let \mathcal{X} be a normal variety with rational singularities, and let $f: \mathcal{X} \rightarrow C$ be a flat projective morphism with connected normal fibers onto a smooth connected algebraic curve C . Let $0 \leq p \leq \dim \mathcal{X} - 1$ be an integer, and assume that $h^0(\mathcal{X}_t, \Omega_{\mathcal{X}_t}^{[p]}) = h^p(\mathcal{X}_t, \mathcal{O}_{\mathcal{X}_t})$ for any point t on C . Given $t_0 \in C$ and $\omega \in H^0(\mathcal{X}_{t_0}, \Omega_{\mathcal{X}_{t_0}}^{[p]})$, and replacing C by an open neighborhood of t_0 if necessary, there exists $\Omega \in H^0(\mathcal{X}, \Omega_{\mathcal{X}/C}^{[p]})$ such that $\Omega|_{(\mathcal{X}_{t_0})_{\text{reg}}} = \omega|_{(\mathcal{X}_{t_0})_{\text{reg}}}$.*

Remark 4.5. In the setup of Proposition 4.4 above, suppose moreover that \mathcal{X}_t has klt singularities for any $t \in C$. Then Hodge symmetry $h^0(\mathcal{X}_t, \Omega_{\mathcal{X}_t}^{[p]}) = h^p(\mathcal{X}_t, \mathcal{O}_{\mathcal{X}_t})$ holds by [GKP11, Proposition 6.9].

Proof of Proposition 4.4. Denote by $i: \mathcal{X}_{\text{reg}} \hookrightarrow \mathcal{X}$ the natural morphism, so that $\Omega_{\mathcal{X}/C}^{[p]} \cong i_* \Omega_{\mathcal{X}_{\text{reg}/C}}^{[p]}$ by [Har80, Proposition 1.6]. The sheaf $\Omega_{\mathcal{X}/C}^{[p]}$ is torsion free, and hence flat over C . This implies in particular that $f_* \Omega_{\mathcal{X}/C}^{[p]}$ is locally free of rank $h^0(\mathcal{X}_{t_1}, \Omega_{\mathcal{X}/C|_{\mathcal{X}_{t_1}}}^{[p]})$ where $t_1 \in C$ is a general point. On the other hand, if t_1 is general enough, then the sheaf $\Omega_{\mathcal{X}/C|_{\mathcal{X}_{t_1}}}^{[p]}$ is reflexive, and hence $\Omega_{\mathcal{X}/C|_{\mathcal{X}_{t_1}}}^{[p]} \cong \Omega_{\mathcal{X}_{t_1}}^{[p]}$.

Given $t \in C$, consider the exact sequence

$$0 \rightarrow \mathcal{O}_{\mathcal{X}_{\text{reg}}}(-\mathcal{X}_t \cap \mathcal{X}_{\text{reg}}) \rightarrow \mathcal{O}_{\mathcal{X}_{\text{reg}}} \rightarrow \mathcal{O}_{\mathcal{X}_t \cap \mathcal{X}_{\text{reg}}} \rightarrow 0,$$

and let $j_t: \mathcal{X}_t \cap \mathcal{X}_{\text{reg}} \hookrightarrow \mathcal{X}_t$ denotes the natural morphism. Tensoring the above exact sequence with the sheaf $\Omega_{\mathcal{X}_{\text{reg}/C}}^{[p]}$ and applying i_* yield an exact sequence

$$0 \rightarrow \Omega_{\mathcal{X}/C}^{[p]} \otimes \mathcal{O}_{\mathcal{X}}(-\mathcal{X}_t) \rightarrow \Omega_{\mathcal{X}/C}^{[p]} \rightarrow (j_t)_* \left(\Omega_{\mathcal{X}_{\text{reg}/C|_{\mathcal{X}_t \cap \mathcal{X}_{\text{reg}}}}^{[p]} \right).$$

Now, by [Gro05, Exposé VIII, Corollaire 2.3], $(j_t)_* \left(\Omega_{\mathcal{X}_{\text{reg}/C|_{\mathcal{X}_t \cap \mathcal{X}_{\text{reg}}}}^{[p]} \right)$ is a coherent sheaf of $\mathcal{O}_{\mathcal{X}_t}$ -modules. On the other hand, the complement of $\mathcal{X}_t \cap \mathcal{X}_{\text{reg}}$ in \mathcal{X}_t has codimension at least 2, and $\Omega_{\mathcal{X}_{\text{reg}/C|_{\mathcal{X}_t \cap \mathcal{X}_{\text{reg}}}}^{[p]}$ is locally free. It follows that $(j_t)_* \left(\Omega_{\mathcal{X}_{\text{reg}/C|_{\mathcal{X}_t \cap \mathcal{X}_{\text{reg}}}}^{[p]} \right)$ is reflexive, and thus $(j_t)_* \left(\Omega_{\mathcal{X}_{\text{reg}/C|_{\mathcal{X}_t \cap \mathcal{X}_{\text{reg}}}}^{[p]} \right) \cong \Omega_{\mathcal{X}_t}^{[p]}$ (see [Har80, Proposition 1.6]). Thus, we obtain an injective morphism of sheaves

$$\Omega_{\mathcal{X}/C|_{\mathcal{X}_t}}^{[p]} \hookrightarrow \Omega_{\mathcal{X}_t}^{[p]}$$

and hence

$$(4.1) \quad h^0(\mathcal{X}_t, \Omega_{\mathcal{X}/C|_{\mathcal{X}_t}}^{[p]}) \leq h^0(\mathcal{X}_t, \Omega_{\mathcal{X}_t}^{[p]}).$$

We claim that the inequality above is an equality, that is,

$$(4.2) \quad h^0(\mathcal{X}_t, \Omega_{\mathcal{X}/C|_{\mathcal{X}_t}}^{[p]}) = h^0(\mathcal{X}_t, \Omega_{\mathcal{X}_t}^{[p]}).$$

Indeed, let us start by observing that as the function $t \mapsto h^0(\mathcal{X}_t, \Omega_{\mathcal{X}/C|_{\mathcal{X}_t}}^{[p]})$ is upper semicontinuous on C , we also have for a general point t_1 on C

$$h^0(\mathcal{X}_t, \Omega_{\mathcal{X}/C|_{\mathcal{X}_t}}^{[p]}) \geq h^0(\mathcal{X}_{t_1}, \Omega_{\mathcal{X}/C|_{\mathcal{X}_{t_1}}}^{[p]}) = h^0(\mathcal{X}_{t_1}, \Omega_{\mathcal{X}_{t_1}}^{[p]}).$$

Now, recall from Lemma 4.3 that $h^p(\mathcal{X}_t, \mathcal{O}_{\mathcal{X}_t})$ is independent of $t \in C$. This implies that $h^0(\mathcal{X}_t, \Omega_{\mathcal{X}_t}^{[p]})$ is independent of $t \in C$ since $h^0(\mathcal{X}_t, \Omega_{\mathcal{X}_t}^{[p]}) = h^p(\mathcal{X}_t, \mathcal{O}_{\mathcal{X}_t})$ by assumption. It follows that

$$h^0(\mathcal{X}_t, \Omega_{\mathcal{X}/C|_{\mathcal{X}_t}}^{[p]}) \geq h^0(\mathcal{X}_t, \Omega_{\mathcal{X}_t}^{[p]}).$$

Combining this with (4.1), we obtain (4.2) hence the claim. Therefore, the quantity $h^0(\mathcal{X}_t, \Omega_{\mathcal{X}/C|_{\mathcal{X}_t}}^{[p]})$ is independent of $t \in C$. By a theorem of Grauert, it follows that the natural map

$$(f_* \Omega_{\mathcal{X}/C}^{[p]}) \otimes \mathbb{C}(t) \rightarrow H^0(\mathcal{X}_t, \Omega_{\mathcal{X}/C|_{\mathcal{X}_t}}^{[p]})$$

is an isomorphism (see [Har77, Corollary 12.9]). Finally, observe that the natural map $\Omega_{\mathcal{X}/C|_{\mathcal{X}_t}}^{[p]} \hookrightarrow \Omega_{\mathcal{X}_t}^{[p]}$ induces an isomorphism

$$H^0\left(\mathcal{X}_t, \Omega_{\mathcal{X}/C|_{\mathcal{X}_t}}^{[p]}\right) \cong H^0\left(\mathcal{X}_t, \Omega_{\mathcal{X}_t}^{[p]}\right)$$

by (4.2). This ends the proof of the proposition. \square

5. HOLONOMY AND STABILITY OF THE TANGENT SHEAF

In this section, we study the stability of the tangent sheaf of smoothable projective varieties with canonical singularities and trivial canonical divisor. The proof of Proposition 5.4 relies on results proved in [GGK17], which we recall first.

5.1. Singular Kähler-Einstein metrics, holonomy and stability. Let X be a normal projective analytic variety with canonical singularities such that $K_X \sim_{\mathbb{Z}} 0$, and let H be an ample Cartier divisor. By [EGZ09, Theorem 7.5], there exists a unique closed positive $(1, 1)$ -current ω with bounded potentials on X such that $[\omega] = [c_1(H)] \in H^2(X, \mathbb{R})$ and such that the restriction of ω to X_{reg} is a smooth Kähler form with zero Ricci curvature. Let g be the Riemannian metric associated with $\omega|_{X_{\text{reg}}}$ on X_{reg} . Given $x \in X_{\text{reg}}$, we view $(T_x X, g_x)$ as an euclidian vector space. We denote the holonomy group (resp. restricted holonomy group) of (X_{reg}, g) at x by $\text{Hol}(X_{\text{reg}}, g)_x$ (resp. $\text{Hol}(X_{\text{reg}}, g)_x^\circ$). It comes with a linear representation on $T_x X$. Recall that $\text{Hol}(X_{\text{reg}}, g)_x$ is a subgroup of $\text{SO}(T_x X, g_x)$, and that $\text{Hol}(X_{\text{reg}}, g)_x^\circ$ is the connected component of the Lie group $\text{Hol}(X_{\text{reg}}, g)_x$. Moreover, the complex structure I on X_{reg} is parallel with respect to g , so that the hermitian metric h_x on $T_x X_{\text{reg}}$ induced by g_x and I_x enables to realize the holonomy group as subgroup of $\text{U}(T_x X, h_x)$. We refer to [GGK17, Section 2.2] for more detailed explanations.

Irreducibility of the holonomy representation and stability of the tangent sheaf are related by the following.

Proposition 5.1 ([GGK17, Proposition 1.5]). *In the above setting, the tangent sheaf T_X is stable (resp. strongly stable) with respect to any polarization if and only if the holonomy representation $\text{Hol}(X_{\text{reg}}, g)_x \circlearrowleft T_x X$ (resp. the restricted holonomy representation $\text{Hol}(X_{\text{reg}}, g)_x^\circ \circlearrowleft T_x X$) is irreducible.*

5.2. Varieties with strongly stable tangent sheaf. The next result relates varieties strongly stable tangent sheaf to irreducible Calabi-Yau and irreducible symplectic varieties.

Theorem 5.2 ([GGK17, Proposition 1.4]). *In the setting of Proposition 5.1, suppose furthermore that X has dimension $n \geq 2$. If T_X is strongly stable, then one of the following two cases holds. In either case, the action of the restricted holonomy group on $T_x X$ is isomorphic to the standard representation.*

- (1) *The group $\text{Hol}(X_{\text{reg}}, g)_x^\circ$ is isomorphic to $\text{SU}(n)$, and X is Calabi-Yau.*
- (2) *The dimension of X is even, the group $\text{Hol}(X_{\text{reg}}, g)_x^\circ$ is isomorphic to $\text{Sp}(\frac{n}{2})$, and there exists a quasi-étale cover $Y \rightarrow X$ such that Y is irreducible symplectic.*

One of the crucial tools in the proof of Theorem 5.2 above is the so-called *Bochner principle*.

Theorem 5.3 (Bochner principle, [GGK17, Theorem A]). *In the setting of Proposition 5.1, let p and q be non-negative integers, and write $\mathcal{E} := (T_X^{\otimes p} \otimes (\Omega_X^1)^{\otimes q})^{**}$. Then the restriction to x induces a one-to-one correspondence between global sections of \mathcal{E} and $\text{Hol}(X_{\text{reg}}, g)_x$ -fixed points in \mathcal{E}_x .*

5.3. Smoothings and holonomy. In the analytic setting 3.1, assume that for $t \neq 0$, \mathcal{X}_t is simply-connected and either irreducible Calabi-Yau or irreducible symplectic. The following result then says that T_X is stable. If moreover $\pi_1^{\text{ét}}(X_{\text{reg}}) = \{1\}$, we conclude that T_X is strongly stable with respect to any polarization.

Proposition 5.4. *In the analytic setting 3.1, assume that for $t \neq 0$, \mathcal{X}_t is simply-connected and either irreducible Calabi-Yau or irreducible symplectic. We maintain notation of Proposition 5.1, and set $G := \text{Hol}(X_{\text{reg}}, g)_x$, $G^\circ := \text{Hol}(X_{\text{reg}}, g)_x^\circ$, and $V := T_x X$. Then the following holds.*

- (1) *The representation $G \circlearrowleft V$ is irreducible. Equivalently, T_X is stable with respect to any polarization.*
- (2) *$G \subset \text{SU}(V)$ if \mathcal{X}_t is irreducible Calabi-Yau and $G \subset \text{Sp}(V)$ if \mathcal{X}_t is irreducible symplectic.*
- (3) *If G° is trivial, then there exists a quasi-étale cover $A \rightarrow X$ where A is an abelian variety.*
- (4) *If G° is nontrivial, then G° is isomorphic to $\text{SU}(r) \times \cdots \times \text{SU}(r)$ or $\text{Sp}(r) \times \cdots \times \text{Sp}(r)$, for some positive integer r . In either case, the action of G° on V is isomorphic to the corresponding product of standard representations.*

Proof. Let us start with (2). In both cases, K_X is trivial so there exists a non-zero holomorphic n -form on X_{reg} . By the Bochner principle (see Theorem 5.3), this implies that $G \subset \text{SU}(V)$. In the symplectic case, Remark 3.2 enables to find two compatible complex structures J_0, K_0 on (X_{reg}, g_0) satisfying $I_0 J_0 K_0 = -\text{Id}$ and which are smooth limits of such complex structures on (\mathcal{X}_t, g_t) . In particular, setting $\sigma_0(X, Y) := g_0(J_0 X, Y) + i g_0(K_0 X, Y)$ defines a *non-degenerate* holomorphic 2-form σ_0 on X_{reg} (see [Bes87, Proposition 14.15]). Therefore, we must have $G \subset \text{Sp}(V)$ by the Bochner principle again.

Moving on to (1), assume that the representation $G \circlearrowleft V$ is reducible. Then one can decompose $V = \bigoplus_{i \in I} W_i$ and write $G = \prod_{i \in I} G_i$ where the action of G_i on W_j is irreducible if $j = i$ and trivial otherwise (see [Bes87, Theorem 10.38]). Consider the Calabi-Yau case first. We have $G_i \subset \text{SU}(W_i)$ by (2), which by the Bochner principle, provides a non-zero reflexive holomorphic form on X_{reg} of degree $\dim W_i$. By Proposition 4.4 and Remark 4.5, it follows that W_i is either zero or the whole V , which concludes. In the symplectic case, (2) and an elementary computation shows that W_i is even dimensional and that $G_i \subset \text{Sp}(W_i)$. In particular, one gets $h^0(X, \Omega_X^{[2]}) = \#I$, and Proposition 4.4 enables to conclude once again. Finally, the equivalence with stability is a consequence of Proposition 5.1 above.

Finally, let us prove (3) and (4). Set $V_0 := \{v \in V, \forall g \in G^\circ, g(v) = v\}$. As G° is normal in G , V_0 is fixed by G . By (1), this implies that V_0 is either zero or the whole V . Assume $V_0 = V$ to start with. Then G° is trivial, that is, (X_{reg}, g) is flat. This implies that $T_{X_{\text{reg}}}$ is holomorphically flat since the $(1, 0)$ -part of the Chern connection of g is holomorphic. By [GKP16, Corollary 1.16], we get the expected quasi-étale cover of X from an abelian variety. So one can now assume that $V_0 = \{0\}$. Therefore $G^\circ \circlearrowleft V$ decomposes as sum of non-trivial irreducible representations $V = \bigoplus_{j \in J} V_j$ along with $G^\circ = \prod_{j \in J} G_j^\circ$. By normality of G° in G , each element $g \in G$ permutes the V_j . As the representation $G \circlearrowleft V$ is irreducible, this implies that the representations $G_j^\circ \circlearrowleft V_j$ are pairwise isomorphic. Set $r = \dim V_j$. By the classification of restricted holonomy [GGK17, Proposition 5.4], we see that $G_j^\circ \circlearrowleft V_j$ is isomorphic to the standard representation of $\text{SU}(r)$ or $\text{Sp}(\frac{r}{2})$, completing the proof of the proposition. \square

6. TOWARDS A DECOMPOSITION THEOREM

6.1. Reduction to smoothings by simply-connected and irreducible manifolds. The main result of this section is Proposition 6.1 below. It reduces the proof of Theorem B to the case of a smoothing by simply-connected, irreducible Calabi-Yau or symplectic manifolds.

Proposition 6.1. *Let X be a normal projective variety with klt singularities. Suppose furthermore that $K_X \equiv 0$, and that X admits a projective \mathbb{Q} -Gorenstein smoothing. Then there exists an abelian variety A as well as a projective variety Y with canonical singularities, a quasi-étale cover*

$$A \times Y \rightarrow X,$$

and a decomposition

$$Y \cong \prod_{i \in I} Y_i$$

such that the following holds.

- (1) *The Y_i admit projective \mathbb{Q} -Gorenstein smoothings over algebraic curves by irreducible and simply-connected Calabi-Yau, or symplectic manifolds.*
- (2) *The sheaves T_{Y_i} are slope-stable with respect to any ample polarization on Y , with trivial determinants.*

We first provide technical tools for the proof of our result.

Lemma 6.2. *Let \mathcal{X} be a normal variety, and let $f: \mathcal{X} \rightarrow C$ be a flat projective morphism with connected fibers onto a smooth connected curve C . Suppose that $K_{\mathcal{X}/C}$ is \mathbb{Q} -Cartier. Suppose furthermore that \mathcal{X}_t has klt singularities for any point t on C . Let $\gamma: \mathcal{Y} \rightarrow \mathcal{X}$ be a quasi-étale cover with \mathcal{Y} normal. Then \mathcal{Y}_t is normal with klt singularities for any point t on C .*

Proof. Note first that $K_{\mathcal{Y}/C}$ is \mathbb{Q} -Cartier since we have $K_{\mathcal{Y}/C} \sim_{\mathbb{Z}} \gamma^* K_{\mathcal{X}/C}$. Moreover, by Fact 2.8, \mathcal{X} has klt singularities. Applying [Kol97, Proposition 3.16] to γ , we see that \mathcal{Y} has klt singularities as well. In particular, \mathcal{Y} is Cohen-Macaulay by [KMM87, Theorem 1.3.6], and hence so is \mathcal{Y}_t for all $t \in C$. By the Nagata-Zariski purity theorem, γ branches only over the singular set of \mathcal{X} . On the other hand, we know that the smooth locus of \mathcal{X}_t is contained in the smooth locus of \mathcal{X} . It follows that \mathcal{Y}_t is smooth in

codimension one. Now, from Serre's criterion for normality, we see that \mathcal{Y}_t is normal for any $t \in C$. Note also that $\gamma_t: \mathcal{Y}_t \rightarrow \mathcal{X}_t$ is a quasi-étale cover. By [Kol97, Proposition 3.16] applied to γ_t , we conclude that \mathcal{Y}_t has klt singularities. \square

Remark 6.3. In the setup of Lemma 6.2, let $\beta: B \rightarrow C$ be the Stein factorization of g . Then β is étale, and γ factors through the étale cover $B \times_C \mathcal{X} \rightarrow \mathcal{X}$.

Lemma 6.4. *Let $f: \mathcal{X} \rightarrow C$ be a flat projective morphism with connected fibers from a normal variety \mathcal{X} onto a smooth connected curve C . Suppose that $K_{\mathcal{X}/C}$ is \mathbb{Q} -Cartier. Suppose furthermore that \mathcal{X}_t has klt singularities for any point t on C . Let $C^\circ \subset C$ be a dense open set, and let $g^\circ: \mathcal{Y}^\circ \rightarrow C^\circ$ be a smooth projective morphism with connected fibers. Finally, let $\gamma^\circ: \mathcal{Y}^\circ \rightarrow f^{-1}(C^\circ)$ be a quasi-étale cover such that $g^\circ = f|_{f^{-1}(C^\circ)} \circ \gamma^\circ$. Then there exists a normal variety \mathcal{Y}_1 as well as a projective morphism $g_1: \mathcal{Y}_1 \rightarrow C_1$ with connected fibers onto a smooth curve C_1 , a finite morphism $\pi: C_1 \rightarrow C$, and a quasi-étale cover $\gamma_1: \mathcal{Y}_1 \rightarrow \mathcal{X} \times_C C_1$ such that the following holds. Write $C_1^\circ := \pi^{-1}(C^\circ)$.*

(1) *We have $\mathcal{Y}_1|_{C_1^\circ} \cong \mathcal{Y}^\circ \times_{C^\circ} C_1^\circ$ and $\gamma_1|_{g_1^{-1}(C_1^\circ)}$ is induced by γ° .*

(2) *Any fiber of g_1 has klt singularities.*

Proof. Let \mathcal{Y} be a reduced and irreducible variety, and let $g: \mathcal{Y} \rightarrow C$ be a projective morphism onto C such that $\mathcal{Y}_{g^{-1}(C^\circ)} \cong \mathcal{Y}^\circ$ as varieties over C° . We may also assume that γ° extends to a generically finite morphism $\gamma: \mathcal{Y} \rightarrow \mathcal{X}$. By a theorem of Kempf, Knudsen, Mumford, and Saint-Donat ([KKMSD73]), there exist a smooth curve C_1 , a finite morphism $\pi: C_1 \rightarrow C$, and a birational projective morphism $\widehat{\mathcal{Y}}_1 \rightarrow \mathcal{Y} \times_C C_1$ from a smooth variety $\widehat{\mathcal{Y}}_1$ such that the induced fibration $\widehat{\mathcal{Y}}_1 \rightarrow C_1$ is semi-stable. We may also assume without loss of generality that $\widehat{\mathcal{Y}}_1 \rightarrow \mathcal{Y} \times_C C_1$ induces an isomorphism over $C_1^\circ := \pi^{-1}(C^\circ)$. Write $\mathcal{X}_1 := \mathcal{X} \times_C C_1$, and consider the Stein factorization $\widehat{\mathcal{Y}}_1 \rightarrow \mathcal{Y}_1 \rightarrow \mathcal{X}_1$ of $\widehat{\mathcal{Y}}_1 \rightarrow \mathcal{X}_1$. We have the following commutative diagram:

$$\begin{array}{ccccc}
 \widehat{\mathcal{Y}}_1 & \xrightarrow{\quad} & \mathcal{Y} \times_C C_1 & \xrightarrow{\quad} & \mathcal{Y} \\
 \downarrow \text{birational} & & \searrow & & \downarrow \gamma \\
 \mathcal{Y}_1 & \xrightarrow{\gamma_1, \text{finite}} & \mathcal{X}_1 = \mathcal{X} \times_C C_1 & \xrightarrow{\quad} & \mathcal{X} \\
 \downarrow g_1 & & \downarrow f_1 & & \downarrow f \\
 C_1 & \xrightarrow{\quad} & C_1 & \xrightarrow{\pi} & C
 \end{array}$$

$\widehat{\mathcal{Y}}_1 \xrightarrow{\text{semi-stable}} C_1$ (curved arrow on the left)
 $\mathcal{Y} \xrightarrow{g} C$ (curved arrow on the right)

Note that $\gamma_1|_{g_1^{-1}(C_1^\circ)}$ is a quasi-étale cover. Since $\widehat{\mathcal{Y}}_1 \rightarrow C_1$ has reduced fibers, it follows that γ_1 is étale in codimension one.

We still have to show that fibers of g_1 are klt. Note first that \mathcal{X}_1 is normal by Fact 2.6. This implies that $K_{\mathcal{X}_1}$ is well-defined and \mathbb{Q} -Cartier since $K_{\mathcal{X}_1/C_1}$ is then the pull-back of $K_{\mathcal{X}/C}$ under the projection morphism $\mathcal{X}_1 = \mathcal{X} \times_C C_1 \rightarrow \mathcal{X}$. The claim now follows from Lemma 6.2, completing the proof of the lemma. \square

We are now in position to prove Proposition 6.1.

Proof of Proposition 6.1. We maintain notation and assumptions of Proposition 6.1. Note that the statement (2) in Proposition 6.1 is an immediate consequence of (1) together with Proposition 5.4.

By Proposition 2.14, replacing X with a quasi-étale cover, if necessary, we may assume that there exist a normal variety \mathcal{X} and a flat projective morphism with connected fibers $f: \mathcal{X} \rightarrow C$ onto a smooth connected algebraic curve C such that $X \cong f^{-1}(t_0)$ for some point t_0 on C , $f^{-1}(t)$ is smooth for $t \neq t_0$, and such that $K_{\mathcal{X}/C} \sim_{\mathbb{Z}} 0$. Set $C^\circ := C \setminus \{t_0\}$ and $\mathcal{X}^\circ := \mathcal{X} \setminus \mathcal{X}_{t_0}$.

Let K be an algebraic closure of the function field of C . Applying the Beauville-Bogomolov decomposition theorem to \mathcal{X}_K (see [Bea83]), we see that, replacing C by a finite cover of some open neighborhood of t_0 , if necessary, there exists an abelian scheme $\mathcal{B}^\circ/C^\circ$, as well as finitely many families $(\mathcal{Y}_i^\circ/C^\circ)_{1 \leq i \leq s}$ of projective manifolds, and a finite étale cover

$$\gamma^\circ: \mathcal{B}^\circ \times_{C^\circ} \mathcal{Y}_1^\circ \times_{C^\circ} \cdots \times_{C^\circ} \mathcal{Y}_s^\circ \rightarrow \mathcal{X}^\circ$$

such that the $(\mathcal{Y}_i^\circ)_K$ are irreducible and simply-connected Calabi-Yau, or symplectic manifolds. In particular, we have $\pi_1^{\text{ét}}((\mathcal{Y}_i^\circ)_K) = \{1\}$. Applying [sga03, Exposé X, Théorème 3.8], we conclude that $\pi_1^{\text{ét}}((\mathcal{Y}_i^\circ)_t) = \{1\}$

for any point t on C° . On the other hand, we have $h^1\left((\mathcal{Y}_i^\circ)_t, \mathcal{O}_{(\mathcal{Y}_i^\circ)_t}\right) = 0$ by Lemma 4.3 and $K_{(\mathcal{Y}_i^\circ)_t} \sim_{\mathbb{Z}} 0$ by the adjunction formula. It follows from [Bea83] that $(\mathcal{Y}_i^\circ)_t$ has finite fundamental group for $t \neq t_0$. Hence $(\mathcal{Y}_i^\circ)_t$ is simply-connected. Using [Bea83] and Lemma 4.3 again, we obtain that for any $t \in C^\circ$, the $(\mathcal{Y}_i^\circ)_t$ are irreducible and simply-connected Calabi-Yau, or symplectic manifolds.

Write $\mathcal{Y}^\circ := \mathcal{Y}_1^\circ \times_{C^\circ} \cdots \times_{C^\circ} \mathcal{Y}_s^\circ$, and $\mathcal{Z}^\circ := \mathcal{B}^\circ \times_{C^\circ} \mathcal{Y}^\circ$. By Lemma 6.4, replacing C by a further cover, if necessary, we may also assume that there exists a normal variety \mathcal{Z} as well as a flat morphism $g: \mathcal{Z} \rightarrow C$ whose fibers have klt singularities, and a quasi-étale cover $\gamma: \mathcal{Z} \rightarrow \mathcal{X}$ such that $g^{-1}(C^\circ) = \mathcal{Z}^\circ$ and $\gamma^\circ = \gamma|_{g^{-1}(C^\circ)}$. We may also assume that $h^\circ: \mathcal{Y}^\circ \rightarrow C^\circ$ has a section. Together with the neutral section $0^\circ: C^\circ \rightarrow \mathcal{B}^\circ$, we obtain a section of $g|_{g^{-1}(C^\circ)}$, and hence a section of g . Moreover, the projection morphism $\mathcal{Z}^\circ \rightarrow \mathcal{B}^\circ$ identifies with the natural morphism $\mathcal{Z}^\circ \rightarrow (\text{Pic}^\circ(\mathcal{Z}^\circ/C^\circ))^\vee$ induced by the universal line bundle. By Proposition 4.2, there exists an abelian scheme $\mathcal{A} \rightarrow C$, as well as a normal variety $\mathcal{Y} \subset \mathcal{Z}$, and a finite étale cover $\mathcal{Y} \times_C \mathcal{A} \rightarrow \mathcal{Z}$ such that the natural map $h: \mathcal{Y} \rightarrow C$ is a flat projective morphism with connected normal fibers, and such that $h^{-1}(C^\circ) = 0^\circ \times_{C^\circ} \mathcal{Y}^\circ \cong \mathcal{Y}^\circ$. The situation is summarized in the following diagram:

$$\begin{array}{ccccccc} \mathcal{A} \times_C \mathcal{Y} & \xrightarrow{\text{étale}} & \mathcal{Z} & \xrightarrow{\gamma} & \mathcal{X} & \longrightarrow & C \\ & & \cup & & \cup & & \cup \\ & & \mathcal{Z}^\circ \cong \mathcal{B}^\circ \times_{C^\circ} \mathcal{Y}^\circ & \longrightarrow & \mathcal{X}^\circ & \longrightarrow & C^\circ \end{array}$$

Let \mathcal{L} be a relatively ample line bundle on \mathcal{Y} . Denote by $s_i^\circ: C^\circ \rightarrow \mathcal{Y}_i^\circ$ the induced section of $\mathcal{Y}_i^\circ \rightarrow C^\circ$, and consider the embedding $\mathcal{Y}_i^\circ \subset \mathcal{Y}_1^\circ \times_{C^\circ} \cdots \times_{C^\circ} \mathcal{Y}_s^\circ \cong \mathcal{Y}^\circ$ induced by the sections s_j for $j \neq i$. Denote by $p_i^\circ: \mathcal{Y}^\circ \rightarrow \mathcal{Y}_i^\circ$ the projection morphism. Then, we have

$$\mathcal{L}|_{\mathcal{Y}^\circ} \cong (p_1^\circ)^*(\mathcal{L}|_{\mathcal{Y}_1^\circ}) \otimes \cdots \otimes (p_s^\circ)^*(\mathcal{L}|_{\mathcal{Y}_s^\circ})$$

since $h^1\left((\mathcal{Y}_i^\circ)_t, \mathcal{O}_{(\mathcal{Y}_i^\circ)_t}\right) = 0$ for any point t on C° .

Suppose now that \mathcal{L} is very ample over C , and set $N_i := h^0\left((\mathcal{Y}_i^\circ)_t, \mathcal{O}_{(\mathcal{Y}_i^\circ)_t}\right) - 1$. Then $\mathcal{L}|_{\mathcal{Y}_i^\circ}$ is very ample over C° and induces an embedding $\mathcal{Y}_i^\circ \subset \mathbb{P}^{N_i} \times C^\circ$. It follows that $\mathcal{Y}_1^\circ \times_{C^\circ} \cdots \times_{C^\circ} \mathcal{Y}_s^\circ$ embeds into $\mathbb{P}^{N_1} \times \cdots \times \mathbb{P}^{N_s} \times C^\circ$. Set $N := h^0(\mathcal{Y}^\circ, \mathcal{O}_{\mathcal{Y}^\circ}) - 1 = (N_1 + 1) \cdots (N_s + 1) - 1$. Then \mathcal{L} embeds \mathcal{Y} into $\mathbb{P}^N \times C$ and the image of \mathcal{Y}° agree with the image of $\mathcal{Y}_1^\circ \times_{C^\circ} \cdots \times_{C^\circ} \mathcal{Y}_s^\circ$ under the Segre embedding $\mathbb{P}^{N_1} \times \cdots \times \mathbb{P}^{N_s} \times C^\circ \subset \mathbb{P}^N \times C^\circ$. Let $\mathcal{Y}_i \subset \mathbb{P}^{N_i} \times C$ be the closure of \mathcal{Y}_i° . Note that $\mathcal{Y}_i \rightarrow C$ is flat. Since the scheme $\text{Hilb}(\mathbb{P}^N)$ is separated, we conclude that $(\mathcal{Y}_1)_{t_0} \times \cdots \times (\mathcal{Y}_s)_{t_0} = \mathcal{Y}_{t_0} \subset \mathbb{P}^N$. This easily implies that $(\mathcal{Y}_i)_{t_0}$ is a normal projective variety with klt singularities and trivial canonical divisor, completing the proof of Proposition 6.1. \square

Proof of Theorem B. By Proposition 2.14, we may assume without loss of generality that X admits a projective \mathbb{Q} -Gorenstein smoothing. Theorem B is now an easy consequence of Propositions 6.1 and 5.4. Indeed, the only thing to check is the assertion concerning the algebra of reflexive forms. Proposition 4.4 settles the Calabi-Yau case immediately. In the symplectic case, item (2) in Proposition 5.4 together with the Bochner principle (see Theorem 5.3) yield a reflexive 2-form σ on X , symplectic on X_{reg} , while Proposition 4.4 shows that σ generates the algebra of reflexive forms. \square

6.2. Irreducible Calabi-Yau and symplectic varieties with stable tangent sheaf. In this section, we try to analyze a bit further the factors appearing in the decomposition of X' in Theorem B. These varieties have stable tangent sheaf, but that sheaf may not be strongly stable. Such varieties are conjecturally covered by either an abelian variety or a product of copies of a single irreducible, Calabi-Yau or symplectic variety, see Conjecture 1.4. Assuming that a weak singular analogue of the Beauville-Bogomolov decomposition theorem holds, we prove this conjecture.

Proposition 6.5. *Suppose that any projective variety X with klt singularities and numerically trivial canonical class admits a quasi-étale cover $Y \rightarrow X$ that splits as a product of an abelian variety and varieties with strongly stable tangent sheaves. Then Conjecture 1.4 holds.*

Proof. We maintain notation and assumptions of Conjecture 1.4. We know that there exists a quasi-étale cover $Y \rightarrow X$ of X such that Y decomposes as a product $Y \cong Y_0 \times Y_1 \times \cdots \times Y_m$ of an abelian variety Y_0 and varieties Y_i for $1 \leq i \leq m$ with strongly stable tangent sheaves. We may assume without loss of generality

that $\dim Y_i \geq 2$ for $1 \leq i \leq m$. Let $Z \rightarrow Y$ be a quasi-étale cover such that the induced quasi-étale cover $Z \rightarrow X$ is Galois, with Galois group G .

The decomposition $Y \cong Y_0 \times Y_1 \times \cdots \times Y_m$ induces a decomposition

$$T_Z \cong \mathcal{E}_0 \oplus \mathcal{E}_1 \oplus \cdots \oplus \mathcal{E}_m$$

of T_Z , where the \mathcal{E}_i for $1 \leq i \leq m$ are strongly stable sheaves. Observe that the foliation \mathcal{E}_i is induced by the Stein factorization of the projection

$$\pi_i: Z \rightarrow B_i := Y_0 \times \cdots \times Y_{i-1} \times Y_{i+1} \times \cdots \times Y_m.$$

Note also that

$$\mathcal{E}_0 \oplus \cdots \oplus \mathcal{E}_{i-1} \oplus \mathcal{E}_{i+1} \oplus \cdots \oplus \mathcal{E}_m$$

induces a flat connection on π_i . A classical result of complex analysis then implies that π_i is a locally trivial fibration for the Euclidean topology over the smooth locus of B_i . Denote by Z_i a connected component of general fiber of π_i . It comes with a quasi-étale cover $Z_i \rightarrow Y_i$.

Suppose first that $\dim Y_0 \geq 1$. Since \mathcal{E}_i is strongly stable for $1 \leq i \leq m$, we have $h^0(Z, \mathcal{E}_i) = 0$ for each $1 \leq i \leq m$. This immediately implies that \mathcal{E}_0 is stable under G , and thus $T_Z = \mathcal{E}_0$ since T_X is stable. This shows that Y is an abelian variety.

Suppose from now on that $\dim Y_0 = 0$. We claim that $\mathcal{E}_i \not\cong \mathcal{E}_j$ if $i \neq j$. Indeed, we have $\mathcal{E}_{j|Z_i} \cong \mathcal{O}_{Z_i}^{\oplus \text{rank} \mathcal{E}_j}$ while $(\mathcal{E}_{i|Z_i})^{**} \cong T_{Z_i}$ and $h^0(Z_i, T_{Z_i}) = 0$ since T_{Z_i} is strongly stable. Therefore, the group G acts on the set $1 \leq i \leq m$ of stable summands of T_Z , and since T_X is stable, this action is transitive. Thus, for any $i \in I$, there exists $g_i \in G$ such that $\mathcal{E}_i \cong g_i^* \mathcal{E}_1 \subset g_i^* T_Z \cong T_Z$. This in turn implies that $Z_i \cong Z_1$, completing the proof of the proposition. \square

To finish this section, let us rephrase some results obtained in Proposition 5.4 with the notations of Theorem B. We start by fixing on X a singular Ricci-flat Kähler metric, provided by [EGZ09]. It can be showed (see [GGK17, Proof of Proposition 7.6]) that the induced metric on $A \times X'$ is actually a product metric, and that the metric induced on X' is also a product metric compatible with the decomposition of X' . Up to passing to a further cover and inflating the abelian part, one can assume that the factors of X' have non-trivial restricted holonomy. Finally, one can pass to an holonomy cover by [GGK17, Theorem B] to ensure that the holonomy is connected. Piecing everything together, Proposition 5.4 shows that a quasi-étale cover of X splits as

$$A \times \prod_{i \in I} Y_i \times \prod_{j \in J} Z_j$$

where A is an abelian variety, and such that the following holds. Let $y_i \in Y_i$ and $z_j \in Z_j$ be smooth points.

- (1) For every $i \in I$, there exist $n_i, r_i \in \mathbb{N}$ with $n_i r_i = \dim Y_i$ such that the holonomy acts on $T_{y_i} Y_i$ by the standard product representation $\text{SU}(n_i)^{\times r_i} \circlearrowleft \mathbb{C}^{\dim Y_i}$.
- (2) For every $j \in J$, there exist $n_j, r_j \in \mathbb{N}$ with $2n_j r_j = \dim Z_j$ such that the holonomy acts on $T_{z_j} Z_j$ by the standard product representation $\text{Sp}(n_j)^{\times r_j} \circlearrowleft \mathbb{C}^{\dim Z_j}$.

7. PROOF OF THEOREM A

In this section we prove Theorem A.

Let X be a (proper) variety, let m be a positive integer, and write $A := \mathbb{C}[[x_1, \dots, x_m]]$ and $K := \mathbb{C}((x_1, \dots, x_m))$. Let also \bar{K} be an algebraic closure of K . Recall that a *deformation of X over A* is a flat morphism of schemes $f: \mathcal{X} \rightarrow \text{Spec } A$ such that $\mathcal{X} \otimes (A/\mathfrak{m}) \cong X$, where \mathfrak{m} denotes the maximal ideal of A . We say that f is a *proper deformation of X over A* if f is a proper morphism.

If f is a smooth proper morphism with geometrically connected fibers, then there is an isomorphism of fundamental groups $\pi_1^{\text{ét}}(\mathcal{X}_{\bar{K}}) \cong \pi_1^{\text{ét}}(X)$ (see [sga03, Exposé X, Théorème 3.8]). However, it is well-known that this statement becomes wrong if f is not assumed to be proper. The following will prove to be crucial.

Theorem 7.1. *Let X be a normal proper variety with klt singularities, and assume that X is smooth in codimension two. Write $A := \mathbb{C}[[x_1, \dots, x_m]]$ and $K := \mathbb{C}((x_1, \dots, x_m))$, and let \bar{K} be an algebraic closure of K . Let \mathcal{X} be a proper deformation of X over A . Suppose that $\mathcal{X}_{\bar{K}}$ is smooth with $\pi_1^{\text{ét}}(\mathcal{X}_{\bar{K}}) = \{1\}$. Then $\pi_1^{\text{ét}}(X_{\text{reg}}) = \{1\}$.*

The following example shows that Theorem 7.1 is wrong if one relaxes the assumption on the codimension of the singular locus.

Example 7.2. Let $X \subset \mathbb{P}^3$ be a cone over a smooth plane cubic curve, and let $f: \mathcal{X} \rightarrow \mathbb{P}^1$ be flat family of cubic surfaces in \mathbb{P}^3 such that $f^{-1}(0) = X$ and $f^{-1}(t)$ is smooth for a general point t on \mathbb{P}^1 . Then $\pi_1(\mathcal{X}_{\mathbb{C}(t)}) = \{1\}$ and $\pi_1(X_{\text{reg}}) \cong \mathbb{Z} \oplus \mathbb{Z}$.

The same arguments used in the proof of Lemmas [Art76, I.9.1 and I.9.2] show that the following holds.

Lemma 7.3. *Let X be a normal variety of dimension at least two, let $X^\circ \subset X_{\text{reg}}$ be an open subset, and let A be a local artinian \mathbb{C} -algebra. Let $i: X^\circ \hookrightarrow X$ be the inclusion map.*

- (1) *Let X_A be a deformation of X over A , and suppose that $X \setminus X^\circ$ has codimension at least two. Then the restriction map induces an isomorphism $\mathcal{O}_{X_A} \cong i_*(i^{-1}\mathcal{O}_{X_A})$. Let X_A^1 and X_A^2 be deformations of X over A , and let $(X_A^1)^\circ \subset X_A^1$ and $(X_A^2)^\circ \subset X_A^2$ be the open subschemes with underlying topological space X° . Then any isomorphism $(X_A^1)^\circ \cong (X_A^2)^\circ$ of A -schemes uniquely extends to an isomorphism $X_A^1 \cong X_A^2$ of A -schemes.*
- (2) *Suppose that X is affine, that $X \setminus X^\circ$ has codimension at least three, and that $\text{depth } \mathcal{O}_{X,x} \geq 3$ for any point $x \in X \setminus X_{\text{reg}}$. Let X_A° be a deformation of X° over A , and set $X_A := \text{Spec } H^0(X_A^\circ, \mathcal{O}_{X_A^\circ})$. Then X_A is a flat deformation of X over A extending X_A° .*

Proposition 7.4. *Let $\gamma: Y \rightarrow X$ be a quasi-étale cover of normal proper varieties. Suppose that X is smooth in codimension two, and that $\text{depth } \mathcal{O}_{Y,y} \geq 3$ for any point $y \in Y \setminus Y_{\text{reg}}$. Let (A, \mathfrak{m}) be a complete local noetherian \mathbb{C} -algebra with residue field \mathbb{C} , and let \mathcal{X} be a proper deformation of X over A . Then there exists a proper deformation \mathcal{Y} of Y over A , and a finite morphism $\Gamma: \mathcal{Y} \rightarrow \mathcal{X}$ extending γ . If moreover A is regular, then Γ is a quasi-étale cover.*

Proof. For any non-negative integer m , write $A_m := A/\mathfrak{m}^{m+1}$, $S := \text{Spec } A$, $S_m := \text{Spec } A_m$, and $X_m := \mathcal{X} \times_S S_m$. We will also denote by $\widehat{\mathcal{X}}$ the formal completion of \mathcal{X} along X . Note that $\widehat{\mathcal{X}}$ is the colimit of the X_m . Moreover, it comes with a proper morphism onto $\widehat{S} := \text{Spf } A$.

Let X° be the smooth locus of X , and set $Y^\circ := \gamma^{-1}(X^\circ)$. By the Nagata-Zariski purity theorem, γ branches only over the singular set of X , and hence $\gamma|_{Y^\circ}: Y^\circ \rightarrow X^\circ$ is an étale cover. Note that $Y \setminus Y^\circ$ has codimension at least three and that $Y^\circ \subset Y_{\text{reg}}$.

Let $i_m: X_m^\circ \subset X_m$ be the open subscheme with underlying topological space X° . By [Gro67, Théorème 18.1.2], there exists a finite étale cover $\gamma_m^\circ: Y_m^\circ \rightarrow X_m^\circ$ such that $Y_m^\circ \cong Y_{m+1}^\circ \times_{S_{m+1}} S_m$ for any integer $m \geq 0$. Set $Y_m := \text{Spec}_{X_m}(i_m \circ \gamma_m^\circ)_* \mathcal{O}_{Y_m^\circ}$, and denote by $\gamma_m: Y_m \rightarrow X_m$ the natural morphism. Note that Y_m° can be identified with the open subscheme of Y_m with underlying topological space Y° . Applying Lemma 7.3 above, we see that Y_m is flat over S_m , and that $\gamma_{m+1} \times_{S_{m+1}} S_m = \gamma_m$. Moreover, by [Gro05, Exposé VIII, Corollaire 2.3], γ_m is finite. Let $\widehat{\mathcal{Y}}$ denote the colimit of the Y_m ; $\widehat{\mathcal{Y}}$ is a noetherian formal scheme, proper and flat over \widehat{S} . It comes with a finite morphism $\widehat{\Gamma}: \widehat{\mathcal{Y}} \rightarrow \widehat{\mathcal{X}}$. Now, by [Gro61, Proposition 5.4.4], $\widehat{\mathcal{Y}}$ is algebraizable. More precisely, there exists a scheme \mathcal{Y} proper over S such that $\widehat{\mathcal{Y}}$ identifies with the formal completion of \mathcal{Y} along Y . Moreover, the morphism $\widehat{\Gamma}: \widehat{\mathcal{Y}} \rightarrow \widehat{\mathcal{X}}$ of formal schemes is induced by a finite morphism of schemes $\Gamma: \mathcal{Y} \rightarrow \mathcal{X}$ (see proof of [Gro61, Proposition 5.4.4]). Note that \mathcal{Y} is flat over S by [Bou61, Chapitre III, §5, Théorème 2 et Proposition 2].

Suppose from now on that A is regular. Let $x \in \mathcal{X}$ be a codimension one point, not contained in the special fiber $\mathcal{X}_{\mathfrak{m}}$, and suppose that Γ is ramified at x . Then γ must be ramified along $\overline{\{x\}} \cap \mathcal{X}_{\mathfrak{m}}$. On the other hand, $\overline{\{x\}} \cap \mathcal{X}_{\mathfrak{m}}$ has codimension one in $\mathcal{X}_{\mathfrak{m}}$ since A is regular. This yields a contradiction, and shows that Γ is a quasi-étale cover, completing the proof of the proposition. \square

Before proving Theorem 7.1 below, we note the following consequence of Proposition 7.4.

Corollary 7.5. *Let X be a normal proper variety with klt singularities, and assume that X is smooth in codimension two. Write $A := \mathbb{C}[[x_1, \dots, x_m]]$, and let \mathcal{X} be a proper deformation of X over A . Then the natural map $\pi_1^{\text{ét}}(X_{\text{reg}}) \rightarrow \pi_1^{\text{ét}}(\mathcal{X}_{\text{reg}})$ is injective.*

Proof. Given a finite étale cover $\gamma^\circ: Y^\circ \rightarrow X_{\text{reg}}$ with Y° connected, we need to show that there exists a finite étale cover $\Gamma^\circ: \mathcal{Y}^\circ \rightarrow \mathcal{X}_{\text{reg}}$ such that Y° is a connected component of $\mathcal{Y}^\circ|_{(\Gamma^\circ)^{-1}(X_{\text{reg}})}$ and such that γ° is induced by Γ° .

Let $\gamma: Y \rightarrow X$ be the normalization of X in the function field of Y° . Note that γ is a quasi-étale cover. Applying [Kol97, Proposition 3.16] to γ , we see that Y has klt singularities. It follows that Y is Cohen-Macaulay by [KMM87, Theorem 1.3.6]. By the Nagata-Zariski purity theorem, γ branches only over the singular set of X , and hence Y is smooth in codimension two. Combining the previous two assertions, one sees that the assumption from Proposition 7.4 about the depth of points in $Y \setminus Y_{\text{reg}}$ is satisfied. Applying the aforementioned proposition then proves the corollary. \square

Proof of Theorem 7.1. By the semicontinuity theorem, we have $h^0(\mathcal{X}_{\bar{K}}, \mathcal{O}_{\mathcal{X}_{\bar{K}}}) = h^0(X, \mathcal{O}_X) = 1$, and hence $\mathcal{X}_{\bar{K}}$ is connected.

Let $\gamma^\circ: Y^\circ \rightarrow X_{\text{reg}}$ be a finite étale cover with Y° connected, and let $\gamma: Y \rightarrow X$ be the normalization of X in the function field of Y° . Note that γ is a quasi-étale cover. The same argument used in the proof of Corollary 7.5 shows that there exists a deformation \mathcal{Y} of Y over A and a quasi-étale cover $\Gamma: \mathcal{Y} \rightarrow \mathcal{X}$ over A extending γ . By the semicontinuity theorem again, we see that $\mathcal{Y}_{\bar{K}}$ is connected. Moreover, by the Nagata-Zariski purity theorem, the finite morphism $\gamma_{\bar{K}}: \mathcal{Y}_{\bar{K}} \rightarrow \mathcal{X}_{\bar{K}}$ is étale, and hence an isomorphism since $\pi_1^{\text{ét}}(\mathcal{X}_{\bar{K}}) = \{1\}$. This implies that γ is an isomorphism as well, completing the proof of the theorem. \square

We will also need the following observation. The proof follows the line of argument given in [KM92, Corollary 12.1.9].

Lemma 7.6. *Let X be a normal projective variety with klt singularities. If X is smooth in codimension two, then any projective smoothing over an algebraic curve is a \mathbb{Q} -Gorenstein smoothing.*

Proof. Let \mathcal{X} be a normal variety, and let $f: \mathcal{X} \rightarrow C$ be a flat projective morphism with connected fibers onto a smooth connected curve C such that $X \cong f^{-1}(t_0)$ for some point t_0 on C and such that $f^{-1}(t)$ is smooth for $t \neq t_0$.

By [KMM87, Theorem 1.3.6], we know that X is Cohen-Macaulay. Let m be a positive integer such that mK_X is a Cartier divisor. Set $U := \mathcal{X} \setminus (X \setminus X_{\text{reg}}) \subset \mathcal{X}_{\text{reg}}$, and denote by $j: U \hookrightarrow \mathcal{X}$ and $i: X_{\text{reg}} = \mathcal{X}_{t_0} \cap U \hookrightarrow X$ the natural inclusions. By [Har80, Proposition 1.6], we have $j_*(\mathcal{O}_X(mK_{\mathcal{X}/C})|_U) \cong \mathcal{O}_X(mK_{\mathcal{X}/C})$ and $i_*(\mathcal{O}_X(mK_{\mathcal{X}/C})|_{X_{\text{reg}}}) \cong i_*(\mathcal{O}_X(mK_X)|_{X_{\text{reg}}}) \cong \mathcal{O}_X(mK_X)$. Applying [KM92, Lemma 12.1.8], we see that $\mathcal{O}_X(mK_{\mathcal{X}/C})|_X \cong \mathcal{O}_X(mK_X)$. This implies that $\mathcal{O}_X(mK_{\mathcal{X}/C})$ is a Cartier divisor, proving the lemma. \square

We are now in position to prove our main result.

Proof of Theorem A. We maintain notation and assumptions of Theorem A.

Applying Lemma 2.11 and Lemma 7.6, we see that we may assume without loss of generality that X admits a projective \mathbb{Q} -Gorenstein smoothing $f: \mathcal{X} \rightarrow C$ over an algebraic curve C . Let t_0 be a point on C such that $X \cong f^{-1}(t_0)$. By Proposition 6.1 and Lemma 7.6 again, we may also assume that \mathcal{X}_t is an irreducible and simply-connected Calabi-Yau, or symplectic manifold for any point $t \neq t_0$, and that T_X is slope-stable with respect to any ample polarization on X .

Denote by A the completion of the local ring \mathcal{O}_{C, t_0} . Note that $A \cong \mathbb{C}[[t]]$ by a theorem of Cohen. Let K denotes the field of fractions of A , and let \bar{K} be an algebraic closure of K . Write $S := \text{Spec } A$, and $\mathcal{X} := \mathcal{X} \times_C S$.

Applying [sga03, Exposé X, Théorème 3.8], we see that $\pi_1^{\text{ét}}(\mathcal{X}_{\bar{K}}) = \{1\}$. It follows from Theorem 7.1 that $\pi_1^{\text{ét}}(X_{\text{reg}}) = \{1\}$, and hence the tangent sheaf T_X is strongly stable. The theorem now follows from Theorem 5.2. \square

Remark 7.7. There is an alternative proof of Theorem A in the case where \mathcal{X}_t is irreducible symplectic, still assuming the $X := \mathcal{X}_0$ is smooth in codimension two. Indeed, in that case a theorem of Namikawa [Nam06, Corollary 2] asserts that if $\pi: Y \rightarrow X$ is a \mathbb{Q} -factorial terminalization of X , then Y is smooth and π is a symplectic resolution. Note that \mathbb{Q} -factorial terminalizations always exist for varieties with canonical singularities by [BCHM10, Corollary 1.4.3]. By a result of Kaledin (see [Kal01, Proposition 1.2]), a symplectic resolution is a semi-small morphism; in particular, $2 \text{codim } \pi^{-1}(X_{\text{sing}}) \geq \text{codim } X_{\text{sing}}$, and therefore $\text{codim } \pi^{-1}(X_{\text{sing}}) \geq 2$. Set $Y^\circ := Y \setminus \pi^{-1}(X_{\text{sing}})$, and note that $Y^\circ \cong X_{\text{reg}}$ and $\text{codim}(Y \setminus Y^\circ) \geq 2$. In particular, we have $\pi_1(Y^\circ) \cong \pi_1(Y)$. By [Tak03, Theorem 1.1], we also have $\pi_1(Y) \cong \pi_1(X)$, and from [Kol93, Lemma 5.2.2]), we see that $\pi_1(X) = 0$. Eventually, X_{reg} is simply-connected, and as T_X is stable by Proposition 5.4, it is automatically strongly stable.

8. EXAMPLES

In this section, we first give examples of smoothable (irreducible) Calabi-Yau and symplectic varieties. We also collect examples which illustrate to what extent our results are sharp. We maintain notation of Section 5.

8.1. Examples of smoothable Calabi-Yau varieties.

Example 8.1 (Nodal hypersurfaces). Let X be a nodal degree $n+2$ hypersurface in \mathbb{P}^{n+1} with $n \geq 3$. This means that the singularities of X are isolated and locally analytically isomorphic to the germ at 0 of the quadric

$$Q := \left\{ z \in \mathbb{C}^{n+1} \mid \sum_{i=1}^{n+1} z_i^2 = 0 \right\}.$$

Then X has canonical Gorenstein singularities, and it is smooth in codimension 2. Moreover, X has trivial canonical bundle and is smoothable by irreducible Calabi-Yau manifolds. By Theorem 7.1, one has $\pi_1^{\text{ét}}(X_{\text{reg}}) = \{1\}$. Then, Proposition 4.4 shows that X is an irreducible Calabi-Yau variety.

This applies in particular to

$$X := \{[x_0 : \cdots : x_4] \in \mathbb{P}^4 \mid x_0 f + x_1 g = 0\}$$

where f and g are general homogeneous polynomials in x_0, \dots, x_4 of degree 4. Indeed, it is clear that X has 16 (isolated) nodal singularities. In that example, there is an alternative proof that bypasses Theorem 7.1. Indeed, blowing up the plane $S := (x_0 = x_1 = 0) \subset \mathbb{P}^4$ yields a small resolution $\pi: \tilde{X} \rightarrow X$ so in particular, we must have $\pi_1(X_{\text{reg}}) \cong \pi_1(\tilde{X} \setminus \pi^{-1}(S)) \cong \pi_1(\tilde{X})$ since $\text{codim } \pi^{-1}(S) \geq 2$. Applying successively [Tak03, Theorem 1.1] and the Lefschetz's hyperplane theorem for fundamental groups, we see that $\pi_1(\tilde{X}) \cong \pi_1(X) = \{1\}$, and hence $\pi_1(X_{\text{reg}}) = \{1\}$ as claimed.

Example 8.2 (Calabi-Yau threefolds with non-nodal singularities). Let $S_1 \rightarrow \mathbb{P}^1$ and $S_2 \rightarrow \mathbb{P}^1$ be rational elliptic surfaces with sections, and let $X := S_1 \times_{\mathbb{P}^1} S_2$ (see [Sch88]). In [Nam94, Example 5.9], Namikawa gives conditions on the singular fibers of each fibration under which X is a smoothable \mathbb{Q} -factorial 3-fold with trivial canonical class with one isolated terminal singularity locally analytically isomorphic to the germ at 0 of the hypersurface

$$\{z \in \mathbb{C}^4 \mid z_1^2 + z_2^2 + z_3(z_3 + z_4)(z_3 - z_4) = 0\}.$$

As in the example above, one concludes that X is an irreducible Calabi-Yau variety.

Example 8.3 (Calabi-Yau threefolds with \mathbb{Q} -factorial isolated rational hypersurface singularities). If X is a projective variety of dimension three with canonical singularities and trivial canonical bundle, then Namikawa and Steenbrink proved in [NS95, Theorem 1.3] that X admits a flat deformation to a smooth Calabi-Yau threefold provided that X has only \mathbb{Q} -factorial, isolated, rational hypersurface singularities. If X is singular, then it is an irreducible Calabi-Yau variety. The argument is as follows.

Suppose from now on that X is singular, and let $Y \rightarrow X$ be any quasi-étale cover. We claim that Y is also singular. Suppose otherwise, and let $Z \rightarrow Y$ be a quasi-étale cover such that the induced finite morphism $Z \rightarrow X$ is Galois. By the Nagata-Zariski purity theorem, $Z \rightarrow Y$ is étale, and thus Z is smooth as well. This in turn implies that X has quotient singularities. On the other hand, any isolated quotient singularity in dimension at least three is rigid by [Sch71, Theorem 2] (see also Remark 1.2), yielding a contradiction. By the Nagata-Zariski purity theorem again, we conclude that Y has isolated singularities.

Applying Proposition 6.1, we see that there exist a quasi-étale cover $Y \rightarrow X$ and a smoothing of Y into irreducible and simply-connected Calabi-Yau or symplectic manifolds. By Proposition 5.4 and Theorem 7.1, Y is an irreducible Calabi-Yau variety, and hence so is X .

If the global assumption on \mathbb{Q} -factoriality is dropped, [NS95, Theorem 2.4] shows that one can still improve the singularities of X by deforming it to a variety with only nodal singularities.

Remark 8.4. It should be noted that from our classification point of view, threefolds are completely understood as they are known to satisfy a singular analogue of the Beauville-Bogomolov decomposition theorem by [Dru16]. Therefore, Examples 8.2 and 8.3 should be thought as illustrative rather than new. For instance, [Dru16] can be used along the same lines as in Example 8.3 above to prove that if X is a smoothable projective variety of dimension three with canonical isolated singularities and trivial canonical class, then X is an irreducible Calabi-Yau variety.

8.2. Examples of smoothable symplectic varieties. Let X be a normal variety. Recall that we say that X is a *symplectic variety* if X has canonical singularities and there exists $\omega \in H^0(X_{\text{reg}}, \Omega_{X_{\text{reg}}}^2)$ everywhere non-degenerate. If $\pi: \tilde{X} \rightarrow X$ is a resolution of singularities of X , then ω extends to a holomorphic 2-form on \tilde{X} by [GKKP11, Theorem 1.4]. Let now $\pi: \tilde{X} \rightarrow X$ be a \mathbb{Q} -factorial terminalization of X . Recall that the existence of π is established in [BCHM10, Corollary 1.4.3]. Then [Nam06, Corollary 2] asserts that X is smoothable if and only if \tilde{X} is smooth. Note that ω automatically extends to a symplectic form on \tilde{X}_{reg} since π is crepant. In particular, if X is smoothable, then X admits a symplectic resolution. Conversely, if X admits a symplectic resolution, then X is smoothable by [Nam01, Theorem 2.2]. More precisely, any smoothing X_t of X by symplectic manifolds is a flat deformation of \tilde{X} .

Note that Namikawa's theorem [Nam06, Corollary 2] provides smoothings by Kähler manifolds which are not projective.

8.3. Examples of smoothable symplectic varieties with stable but not strongly stable tangent sheaf. Recall from Proposition 5.4 and Theorem 7.1 that if a normal projective variety X with klt singularities and $K_X \equiv 0$ whose tangent sheaf is not strongly stable admits a projective smoothing into simply-connected irreducible Calabi-Yau, or symplectic manifolds then $\text{codim}(X \setminus X_{\text{reg}}) = 2$.

Example 8.5 (Singular Kummer surface). In the following example, we consider a degeneration of $K3$ surfaces to a singular Kummer surface X . More precisely, let $X = A/\langle \pm 1 \rangle$ where A is a principally polarized abelian surface. Then the following holds.

- The Kummer surface X admits a \mathbb{Q} -Gorenstein projective smoothing $f: \mathcal{X} \rightarrow C$ by $K3$ surfaces.
- The tangent sheaf T_X is not strongly stable.
- Let \mathcal{L} be a relatively ample line bundle on \mathcal{X} , and denote by g the Ricci-flat Kähler metric on X_{reg} given by [EGZ09, Theorem 7.5] applied to $(\mathcal{X}_0, \mathcal{L}|_{\mathcal{X}_0})$. Given $x \in X_{\text{reg}}$, we have

$$\text{Hol}(X_{\text{reg}}, g)_x \cong \mathbb{Z}/2\mathbb{Z} \quad \text{and} \quad \text{Hol}^\circ(X_{\text{reg}}, g)_x = \{1\}.$$

- We have $\pi_1(X) = \{1\}$ and $\pi_1(X_{\text{reg}})$ is an extension of \mathbb{Z}^4 by $\mathbb{Z}/2\mathbb{Z}$.

It is well-known that X can be realized as a singular quartic surface in \mathbb{P}^3 (see [BL04, Theorem 4.8.1]). In particular, it can be seen as the fiber over some point $t_0 \in C$ of a projective flat family $f: \mathcal{X} \rightarrow C$ of quartic surfaces over a smooth algebraic curve C such that \mathcal{X}_t is smooth if $t \neq t_0$. The total space \mathcal{X} is an hypersurface in $C \times \mathbb{P}^3$, and hence \mathbb{Q} -Gorenstein. For $t \neq t_0$, \mathcal{X}_t is a smooth quartic surface, and thus it is a $K3$ surface.

The tangent sheaf T_X is not strongly stable as X admits a quasi-étale cover $\gamma: A \rightarrow A/\langle \pm 1 \rangle$ of degree two and $T_A \cong \mathcal{O}_A^{\oplus 2}$ is obviously not stable.

By [EGZ09, Theorem 7.5], there exists a unique closed positive $(1, 1)$ -current ω with bounded potentials on X such that $[\omega] = [c_1(\mathcal{L}|_X)] \in H^2(X, \mathbb{R})$ and such that the restriction of ω to X_{reg} is a smooth Kähler form with zero Ricci curvature. Note that g is the Riemannian metric associate with $\omega|_{X_{\text{reg}}}$ on X_{reg} . We claim that g is flat, or equivalently that $\text{Hol}^\circ(X_{\text{reg}}, g)_x = \{1\}$. Indeed, it is known that ω is of orbifold type, that is, $\gamma^*\omega$ defines a smooth Ricci-flat Kähler metric on A . On the other hand, it is well-known that any Ricci-flat metric on a torus is flat. Next, we show that $\text{Hol}(X_{\text{reg}}, g)_x \cong \mathbb{Z}/2\mathbb{Z}$. Note that the natural surjection $\pi_1(X_{\text{reg}}) \twoheadrightarrow \text{Hol}(X_{\text{reg}}, g)_x / \text{Hol}^\circ(X_{\text{reg}}, g)_x$ together with the fact that $(\gamma|_{\gamma^{-1}(X_{\text{reg}})})^*g$ is flat yields a surjective map $\mathbb{Z}/2\mathbb{Z} \twoheadrightarrow \text{Hol}(X_{\text{reg}}, g)_x$. Suppose that $\text{Hol}(X_{\text{reg}}, g)_x$ is trivial. Then, by Bochner principle, one would get a non-zero element $\sigma \in H^0(X, \Omega_X^{[1]})$, and therefore a non-zero global 1-form on A invariant under the involution $a \mapsto -a$, yielding a contradiction. Thus

$$\text{Hol}(X_{\text{reg}}, g)_x \cong \mathbb{Z}/2\mathbb{Z}.$$

As for the fundamental group, X is simply-connected by Lefschetz theorem. Moreover, a degree two étale cover of X_{reg} is isomorphic to the complement of 16 points in an abelian surface, therefore the fundamental group of this cover is \mathbb{Z}^4 .

Example 8.6 (Symmetric square of a $K3$ surface). This following example was already exhibited in [GKP16, Example 8.6] as a variety with stable but not strongly stable tangent sheaf. Let S be a $K3$ surface, and let $X := S \times S / \langle i \rangle$ where i is the natural involution $(s_1, s_2) \mapsto (s_2, s_1)$ of $S \times S$. Recall from [Bea83, Section

6] that the Hilbert scheme $S^{[2]}$ parametrizing length 2 zero-dimensional subschemes on S is an irreducible symplectic manifold which admits a birational crepant morphism $S^{[2]} \rightarrow X$. Then the following holds.

- The variety X admits a Kähler deformation to a smooth irreducible symplectic variety.
- The tangent sheaf X is not strongly stable.
- Let $x \in X_{\text{reg}}$. There exists a Ricci-flat Kähler metric g on X_{reg} such that

$$\text{Hol}(X_{\text{reg}}, g)_x \cong (\text{SU}(2) \times \text{SU}(2)) \rtimes \mathbb{Z}/2\mathbb{Z} \quad \text{and} \quad \text{Hol}^\circ(X_{\text{reg}}, g)_x \cong \text{SU}(2) \times \text{SU}(2).$$

- We have $\pi_1(X) = \{1\}$ and $\pi_1(X_{\text{reg}}) \cong \mathbb{Z}/2\mathbb{Z}$.

The claim on the existence of a smoothing follows from the existence of the symplectic resolution $S^{[2]} \rightarrow X$ combined with a theorem of Namikawa [Nam06, Corollary 2] (see also Section 8.2).

The tangent sheaf T_X is not strongly stable as X admits a quasi-étale cover $\gamma: S \times S \rightarrow X$ of degree two and $T_{S \times S}$ is obviously not stable.

Let us consider the two projections $p_i: S \times S \rightarrow S$ with $i = 1, 2$. Given a Ricci-flat Kähler metric g_S on S , $p_1^*g_S + p_2^*g_S$ defines a Kähler Ricci-flat metric on $S \times S$ that descends to a Ricci-flat Kähler metric g on X_{reg} . Let y be a point on $S \times S$ such that $\gamma(y) = x$. As the restricted holonomy is preserved by passing to an étale cover, we must have

$$\text{Hol}^\circ(X_{\text{reg}}, g)_x \cong \text{Hol}^\circ(S \times S \setminus \Delta, \gamma^*g)_y = \text{SU}(2) \times \text{SU}(2)$$

where $\Delta \subset S \times S$ denotes the diagonal.

As in the previous example, we have a surjective map $\mathbb{Z}/2\mathbb{Z} \rightarrow \text{Hol}(X_{\text{reg}}, g)_x / \text{Hol}^\circ(X_{\text{reg}}, g)_x$. Suppose that $\text{Hol}(X_{\text{reg}}, g)_x^\circ = \text{Hol}^\circ(X_{\text{reg}}, g)_x$. Then, by the Bochner principle, one would get two linearly independent reflexive 2-forms on X , hence two independent 2-forms on $S \times S$ invariant by the involution, yielding a contradiction. This shows that

$$\text{Hol}(X_{\text{reg}}, g)_x \cong (\text{SU}(2) \times \text{SU}(2)) \rtimes \mathbb{Z}/2\mathbb{Z}.$$

By a theorem of Armstrong (see [Arm82]), we have $\pi_1(X) = \{1\}$. As for X_{reg} , it admits a degree two cover from $S \times S \setminus \Delta$. Since $\text{codim}\Delta = 2$ and $S \times S$ is simply-connected, we conclude that $S \times S \setminus \Delta$ is simply-connected as well, and hence $\pi_1(X_{\text{reg}}) \cong \mathbb{Z}/2\mathbb{Z}$.

Remark 8.7. As already mentioned, Namikawa's theorem [Nam06, Corollary 2] provides smoothings by Kähler manifolds which are not projective. We do not know whether X admits a projective smoothing by irreducible symplectic varieties.

REFERENCES

- [Arm82] M. A. Armstrong, *Calculating the fundamental group of an orbit space*, Proc. Amer. Math. Soc. **84** (1982), no. 2, 267–271.
- [Art76] Michael Artin, *Lectures on deformations of singularities. Notes by C. S. Seshadri, Allen Tannenbaum*, Lectures on Mathematics and Physics. 54. Bombay: Tata Institute of Fundamental Research. II, 127 p. (1976), 1976.
- [BCHM10] Caucher Birkar, Paolo Cascini, Christopher D. Hacon, and James McKernan, *Existence of minimal models for varieties of log general type*, J. Amer. Math. Soc. **23** (2010), no. 2, 405–468.
- [Bea83] Arnaud Beauville, *Variétés Kähleriennes dont la première classe de Chern est nulle*, J. Differential Geom. **18** (1983), no. 4, 755–782 (1984).
- [Bes87] Arthur L. Besse, *Einstein manifolds*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 10, Springer-Verlag, Berlin, 1987.
- [BL04] Christina Birkenhake and Herbert Lange, *Complex abelian varieties. 2nd augmented ed.*, 2nd augmented ed. ed., Berlin: Springer, 2004.
- [BLR90] Siegfried Bosch, Werner Lütkebohmert, and Michel Raynaud, *Néron models*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 21, Springer-Verlag, Berlin, 1990.
- [Bou61] N. Bourbaki, *éléments de mathématique. Fascicule XXVIII. Algèbre commutative. Chapitre 3: Graduations, filtrations et topologies. Chapitre 4: Idéaux premiers associés et décomposition primaire*, Actualités Scientifiques et Industrielles, No. 1293, Hermann, Paris, 1961.
- [Bri10] Michel Brion, *Some basic results on actions of nonaffine algebraic groups*, Symmetry and spaces, Progr. Math., vol. 278, Birkhäuser Boston, Inc., Boston, MA, 2010, pp. 1–20.
- [BS76] Constantin Bănică and Octavian Stănăşilă, *Algebraic methods in the global theory of complex spaces*, Editura Academiei, Bucharest; John Wiley & Sons, London-New York-Sydney, 1976, Translated from the Romanian.
- [Con00] Brian Conrad, *Grothendieck duality and base change*, Lecture Notes in Mathematics, vol. 1750, Springer-Verlag, Berlin, 2000.

- [DG67] J. Dieudonné and A. Grothendieck, *Critères différentiels de régularité pour les localisés des algèbres analytiques*, J. Algebra **5** (1967), 305–324.
- [Dru16] Stéphane Druel, *A decomposition theorem for singular spaces with trivial canonical class of dimension at most five*, Preprint [arXiv:1606.09006v2](https://arxiv.org/abs/1606.09006v2), 2016.
- [DS14] S. Donaldson and S. Sun, *Gromov-Hausdorff limits of Kähler manifolds and algebraic geometry*, Acta Math. **213** (2014), no. 1, 63–106.
- [EGZ09] Philippe Eyssidieux, Vincent Guedj, and Ahmed Zeriahi, *Singular Kähler-Einstein metrics*, J. Am. Math. Soc. **22** (2009), no. 3, 607–639.
- [GGK17] Daniel Greb, Henri Guenancia, and Stefan Kebekus, *Klt varieties with trivial canonical class: holonomy, differential forms, and fundamental groups*, Preprint [arXiv:1704.01408](https://arxiv.org/abs/1704.01408), 2017.
- [GKKP11] Daniel Greb, Stefan Kebekus, Sándor J. Kovács, and Thomas Peternell, *Differential forms on log canonical spaces*, Publ. Math. Inst. Hautes Études Sci. (2011), no. 114, 87–169. MR 2854859
- [GKP11] Daniel Greb, Stefan Kebekus, and Thomas Peternell, *Singular spaces with trivial canonical class*, To appear in: Minimal models and extremal rays — proceedings of the conference in honor of Shigefumi Mori’s 60th birthday, Advanced Studies in Pure Mathematics, Kinokuniya Publishing House, Tokyo, 2011.
- [GKP16] ———, *Singular spaces with trivial canonical class*, Minimal Models and Extremal Rays, Kyoto, 2011, Adv. Stud. Pure Math., vol. 70, Mathematical Society of Japan, Tokyo, 2016, Preprint [arXiv:1110.5250](https://arxiv.org/abs/1110.5250), pp. 67–113.
- [GP11] Philippe Gille and Patrick Polo (eds.), *Schémas en groupes (SGA 3). Tome I. Propriétés générales des schémas en groupes*, Documents Mathématiques (Paris) [Mathematical Documents (Paris)], 7, Société Mathématique de France, Paris, 2011, Séminaire de Géométrie Algébrique du Bois Marie 1962–64. [Algebraic Geometry Seminar of Bois Marie 1962–64], A seminar directed by M. Demazure and A. Grothendieck with the collaboration of M. Artin, J.-E. Bertin, P. Gabriel, M. Raynaud and J-P. Serre, Revised and annotated edition of the 1970 French original.
- [Gro61] A. Grothendieck, *éléments de géométrie algébrique. III. étude cohomologique des faisceaux cohérents. I*, Inst. Hautes Études Sci. Publ. Math. (1961), no. 11, 167.
- [Gro65] ———, *Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. II*, Inst. Hautes Études Sci. Publ. Math. (1965), no. 24, 231.
- [Gro66] ———, *Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. III*, Inst. Hautes Études Sci. Publ. Math. (1966), no. 28, 255.
- [Gro67] ———, *éléments de géométrie algébrique. IV. étude locale des schémas et des morphismes de schémas IV*, Inst. Hautes Études Sci. Publ. Math. (1967), no. 32, 361.
- [Gro95a] Alexander Grothendieck, *Technique de descente et théorèmes d’existence en géométrie algébrique. V. Les schémas de Picard: théorèmes d’existence*, Séminaire Bourbaki, Vol. 7, Soc. Math. France, Paris, 1995, pp. Exp. No. 232, 143–161.
- [Gro95b] ———, *Techniques de construction et théorèmes d’existence en géométrie algébrique. IV. Les schémas de Hilbert*, Séminaire Bourbaki, Vol. 6, Soc. Math. France, Paris, 1995, pp. Exp. No. 221, 249–276.
- [Gro05] ———, *Cohomologie locale des faisceaux cohérents et théorèmes de Lefschetz locaux et globaux (SGA 2)*, Documents Mathématiques (Paris) [Mathematical Documents (Paris)], 4, Société Mathématique de France, Paris, 2005, Séminaire de Géométrie Algébrique du Bois Marie, 1962, Augmenté d’un exposé de Michèle Raynaud. [With an exposé by Michèle Raynaud], With a preface and edited by Yves Laszlo, Revised reprint of the 1968 French original.
- [Har77] Robin Hartshorne, *Algebraic geometry*, Springer-Verlag, New York, 1977, Graduate Texts in Mathematics, No. 52.
- [Har80] ———, *Stable reflexive sheaves*, Math. Ann. **254** (1980), no. 2, 121–176.
- [Kal01] Dmitry Kaledin, *Symplectic resolutions: deformations and birational maps*, Preprint [arXiv:0012008](https://arxiv.org/abs/0012008), 2001.
- [Kar00] Kalle Karu, *Minimal models and boundedness of stable varieties*, J. Algebraic Geom. **9** (2000), no. 1, 93–109.
- [Kaw85] Yujiro Kawamata, *Minimal models and the Kodaira dimension of algebraic fiber spaces*, J. Reine Angew. Math. **363** (1985), 1–46.
- [KKMSD73] G. Kempf, Finn Faye Knudsen, D. Mumford, and B. Saint-Donat, *Toroidal embeddings. I*, Lecture Notes in Mathematics, Vol. 339, Springer-Verlag, Berlin-New York, 1973.
- [KM92] János Kollár and Shigefumi Mori, *Classification of three-dimensional flips*, J. Amer. Math. Soc. **5** (1992), no. 3, 533–703.
- [KM98] ———, *Birational geometry of algebraic varieties*, Cambridge Tracts in Mathematics, vol. 134, Cambridge University Press, Cambridge, 1998, With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original.
- [KMM87] Yujiro Kawamata, Katsumi Matsuda, and Kenji Matsuki, *Introduction to the minimal model problem*, Algebraic geometry, Sendai, 1985, Adv. Stud. Pure Math., vol. 10, North-Holland, Amsterdam, 1987, pp. 283–360.
- [Kol86] János Kollár, *Higher direct images of dualizing sheaves. II*, Ann. of Math. (2) **124** (1986), no. 1, 171–202.
- [Kol93] ———, *Shafarevich maps and plurigenera of algebraic varieties*, Invent. Math. **113** (1993), no. 1, 177–215.
- [Kol97] ———, *Singularities of pairs*, Algebraic geometry—Santa Cruz 1995, Proc. Sympos. Pure Math., vol. 62, AMS, 1997, pp. 221–287.
- [Laz04] Robert Lazarsfeld, *Positivity in algebraic geometry. I*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 48, Springer-Verlag, Berlin, 2004, Classical setting: line bundles and linear series.

- [LWX14] Chi Li, Xiaowei Wang, and Chenyang Xu, *On proper moduli spaces of smoothable Kähler-Einstein Fano varieties*, Preprint [arXiv:1411.0761](https://arxiv.org/abs/1411.0761), 2014.
- [MFK94] D. Mumford, J. Fogarty, and F. Kirwan, *Geometric invariant theory*, third ed., *Ergebnisse der Mathematik und ihrer Grenzgebiete (2) [Results in Mathematics and Related Areas (2)]*, vol. 34, Springer-Verlag, Berlin, 1994.
- [Nak04] Noboru Nakayama, *Zariski-decomposition and abundance*, *MSJ Memoirs*, vol. 14, Mathematical Society of Japan, Tokyo, 2004.
- [Nam94] Yoshinori Namikawa, *On deformations of calabi-yau 3-folds with terminal singularities*, *Topology* **33** (1994), no. 3, 429–446.
- [Nam01] ———, *Deformation theory of singular symplectic n -folds*, *Math. Ann.* **319** (2001), no. 3, 597–623.
- [Nam06] ———, *On deformations of \mathbb{Q} -factorial symplectic varieties*, *J. Reine Angew. Math.* **599** (2006), 97–110.
- [NS95] Yoshinori Namikawa and J. H. M. Steenbrink, *Global smoothing of Calabi-Yau threefolds*, *Invent. Math.* **122** (1995), no. 2, 403–419.
- [RZ11a] Xiaochun Rong and Yuguang Zhang, *Continuity of extremal transitions and flops for Calabi-Yau manifolds*, *J. Differential Geom.* **89** (2011), no. 2, 233–269, Appendix B by Mark Gross.
- [RZ11b] Wei-Dong Ruan and Yuguang Zhang, *Convergence of Calabi-Yau manifolds*, *Adv. Math.* **228** (2011), no. 3, 1543–1589.
- [Sch71] Michael Schlessinger, *Rigidity of quotient singularities*, *Invent. Math.* **14** (1971), 17–26.
- [Sch88] Chad Schoen, *On fiber products of rational elliptic surfaces with section*, *Math. Z.* **197** (1988), no. 2, 177–199.
- [Ser01] Jean-Pierre Serre, *Exposés de séminaires (1950-1999)*, *Documents Mathématiques (Paris) [Mathematical Documents (Paris)]*, vol. 1, Société Mathématique de France, Paris, 2001.
- [sga03] *Revêtements étales et groupe fondamental (SGA 1)*, *Documents Mathématiques (Paris) [Mathematical Documents (Paris)]*, 3, Société Mathématique de France, Paris, 2003, Séminaire de géométrie algébrique du Bois Marie 1960–61. [Algebraic Geometry Seminar of Bois Marie 1960-61], Directed by A. Grothendieck, With two papers by M. Raynaud, Updated and annotated reprint of the 1971 original [Lecture Notes in Math., 224, Springer, Berlin].
- [SSY16] Cristiano Spotti, Song Sun, and Chengjian Yao, *Existence and deformations of Kähler-Einstein metrics on smoothable \mathbb{Q} -Fano varieties*, *Duke Math. J.* **165** (2016), no. 16, 3043–3083.
- [Tak03] Shigeharu Takayama, *Local simple connectedness of resolutions of log-terminal singularities*, *Internat. J. Math.* **14** (2003), no. 8, 825–836.
- [Yau78] Shing-Tung Yau, *On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation. I.*, *Commun. Pure Appl. Math.* **31** (1978), 339–411.

STÉPHANE DRUEL: INSTITUT FOURIER, UMR 5582 DU CNRS, UNIVERSITÉ GRENOBLE ALPES, CS 40700, 38058 GRENOBLE CEDEX 9, FRANCE

E-mail address: stephane.drue1@univ-grenoble-alpes.fr

HENRI GUENANCIA: DEPARTMENT OF MATHEMATICS, STONY BROOK UNIVERSITY, STONY BROOK, NY 11794-3651, UNITED STATES

E-mail address: henri.guenancia@stonybrook.edu