



LIPSCHITZ REGULARITY FOR ORTHOTROPIC FUNCTIONALS WITH NONSTANDARD GROWTH CONDITIONS

Pierre Bousquet, Lorenzo Brasco

► To cite this version:

Pierre Bousquet, Lorenzo Brasco. LIPSCHITZ REGULARITY FOR ORTHOTROPIC FUNCTIONALS WITH NONSTANDARD GROWTH CONDITIONS. 2018. hal-01886305

HAL Id: hal-01886305

<https://hal.science/hal-01886305>

Preprint submitted on 2 Oct 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

LIPSCHITZ REGULARITY FOR ORTHOTROPIC FUNCTIONALS WITH NONSTANDARD GROWTH CONDITIONS

PIERRE BOUSQUET AND LORENZO BRASCO

ABSTRACT. We consider a model convex functional with orthotropic structure and super-quadratic nonstandard growth conditions. We prove that bounded local minimizers are locally Lipschitz, with no restrictions on the ratio between the highest and the lowest growth rate.

CONTENTS

1. Introduction	1
1.1. Overview	1
1.2. Main result	3
1.3. Structure of the proof	4
1.4. Plan of the paper	8
2. Preliminaries	9
3. Caccioppoli-type inequalities	12
4. Local energy estimates for the regularized problem	13
4.1. Towards an iterative Moser's scheme	13
4.2. Towards higher integrability	16
5. A Lipschitz estimate	18
6. A recursive gain of integrability for the gradient	23
7. Proof of Theorem 1.1	29
Appendix A. Calculus lemmas	30
A.1. Tools for the Lipschitz estimate	30
A.2. Tools for the higher integrability	33
References	36

1. INTRODUCTION

1.1. Overview. We pursue our study of the gradient regularity for local minimizers of functionals from the Calculus of Variations, having a structure that we called *orthotropic*. We refer to our previous contributions [2, 3, 4, 5] and [6], for an introduction to the subject.

More precisely, we want to expand the investigation carried on in [6], by studying functionals of the form

$$\int f(\nabla u) dx, \quad f : \mathbb{R}^N \rightarrow \mathbb{R} \text{ convex},$$

which couple the following two features

orthotropic structure and *nonstandard growth conditions*.

2010 *Mathematics Subject Classification*. 35J70, 35B65, 49K20.

Key words and phrases. Nonstandard growth conditions, degenerate elliptic equations, Lipschitz regularity, orthotropic problems.

The first one means that we require

$$f(z) = \sum_{i=1}^N f_i(z_i), \quad \text{with } f_i : \mathbb{R} \rightarrow \mathbb{R} \text{ convex,}$$

while the second one means that

$$|z|^p - 1 \lesssim f(z) \lesssim |z|^q + 1, \quad \text{with } 1 < p < q.$$

As we will see, these two features give rise to one of the most challenging type of functionals, at least if one is interested in higher order regularity of local minimizers, i.e. regularity of their gradients.

Let us be more specific on the type of functionals we want to study. We take a vector $\mathbf{p} = (p_1, \dots, p_N)$ with $2 \leq p_1 \leq \dots \leq p_N$. Let $\Omega \subset \mathbb{R}^N$ be an open set. For every $u \in W_{\text{loc}}^{1,\mathbf{p}}(\Omega)$ and every $\Omega' \Subset \Omega$, we consider the *orthotropic functional with nonstandard growth*

$$\mathfrak{F}_{\mathbf{p}}(u, \Omega') = \sum_{i=1}^N \frac{1}{p_i} \int_{\Omega'} |u_{x_i}|^{p_i} dx.$$

We say that $u \in W_{\text{loc}}^{1,\mathbf{p}}(\Omega)$ is a *local minimizer* of $\mathfrak{F}_{\mathbf{p}}$ if

$$\mathfrak{F}_{\mathbf{p}}(u, \Omega') \leq \mathfrak{F}_{\mathbf{p}}(v, \Omega'), \quad \text{for every } v - u \in W_0^{1,\mathbf{p}}(\Omega') \quad \text{and every } \Omega' \Subset \Omega.$$

Here $W^{1,\mathbf{p}}$ and $W_0^{1,\mathbf{p}}$ are the classical anisotropic Sobolev spaces, defined for an open set $E \subset \mathbb{R}^N$ by

$$W^{1,\mathbf{p}}(E) = \{u \in L^1(E) : u_{x_i} \in L^{p_i}(E), i = 1, \dots, N\},$$

and

$$W_0^{1,\mathbf{p}}(E) = W^{1,\mathbf{p}}(E) \cap W_0^{1,1}(E).$$

It is easy to see that a local minimizer of $\mathfrak{F}_{\mathbf{p}}$ is a local weak solution of the following quasilinear equation with orthotropic structure

$$(1.1) \quad \sum_{i=1}^N \left(|u_{x_i}|^{p_i-2} u_{x_i} \right)_{x_i} = 0.$$

It is well-known that local minimizers of functionals like $\mathfrak{F}_{\mathbf{p}}$ above can be *unbounded* if the ratio

$$\frac{p_N}{p_1},$$

is too large, see the celebrated counter-examples by Giaquinta [21] and Marcellini [28, 29, 30] (see also Hong's paper [23]). In Western countries, the regularity theory for *non degenerate* functionals with nonstandard growth was initiated in the seminal papers [28, 29] by Marcellini. For *strongly degenerate functionals*, including the orthotropic functional with nonstandard growth $\mathfrak{F}_{\mathbf{p}}$ introduced above, the question has been addressed in the Russian literature, see for example the papers [24] by Kolodii, [25] by Koralev and [34] by Uralt'seva and Urdaletova.

However, in spite of a large number of papers and contributions on the subject, a satisfactory gradient regularity theory for these problems is still missing. Some higher integrability results for the gradient have been obtained for example in [18, Theorem 2.1] and [17, Theorem 5]. In any case, we point out that even the case of basic regularity (i.e. $C^{0,\alpha}$ estimates and Harnack inequalities) is still not completely well-understood, we refer to the recent paper [1] and the references therein.

1.2. Main result. In this paper, we are going to prove that *bounded* local minimizers of our orthotropic functional $\mathfrak{F}_{\mathbf{p}}$ are locally Lipschitz continuous. We point out that **no upper bounds on the ratio**

$$\frac{p_N}{p_1},$$

are needed for the result to hold.

Theorem 1.1. *Let $\mathbf{p} = (p_1, \dots, p_N)$ be such that $2 \leq p_1 \leq \dots \leq p_N$. Let $U \in W_{\text{loc}}^{1,\mathbf{p}}(\Omega)$ be a local minimizer of $\mathfrak{F}_{\mathbf{p}}$ such that*

$$U \in L_{\text{loc}}^{\infty}(\Omega).$$

Then $\nabla U \in L_{\text{loc}}^{\infty}(\Omega)$.

Remark 1.2 (L^{∞} assumption). Sharp conditions in order to get $U \in L_{\text{loc}}^{\infty}$ can be found in [20, Theorem 1] by Fusco and Sbordone, see also [19, Theorem 3.1] and the papers [11, 12] by Cupini, Marcellini and Mascolo for the case of more general functionals. Pioneering results are due to Kolodň, see [24]. We also mention the recent paper [16] by DiBenedetto, Gianazza and Vespi, where some precise a priori L^{∞} estimates on the solution are proved, see Section 6 there.

Remark 1.3 (Comparison with previous results). Some particular cases of our Theorem 1.1 can be traced back in the literature. We try to give a complete picture of the subject.

The first one is [34, Theorem 1] by Uralt'seva and Urdaletova, which proves local Lipschitz regularity for bounded minimizers, under the restrictions

$$p_1 \geq 4 \quad \text{and} \quad \frac{p_N}{p_1} < 2.$$

The method of proof of [34] is completely different from ours and is based on the so-called *Bernstein's technique*. We refer to [3] for a detailed description of their proof.

More recently, Theorem 1.1 has been proved in the two-dimensional case $N = 2$ by the second author in collaboration with Leone, Pisante and Verde, see [6, Theorem 1.4]. In this case as well, the proof is different from the one we give here, the former being based on a two-dimensional trick introduced in [3, Theorem A]. Still in dimension $N = 2$, Lindqvist and Ricciotti in [27, Theorem 1.2] proved C^1 regularity for solutions of (1.1), by extending to the case of nonstandard growth conditions a result of the authors, see [2, Main Theorem].

In the standard growth case, i.e. when $p_1 = \dots = p_N = p$, local Lipschitz regularity has been obtained in [4, Theorem 1.1]. As we will explain later, the result of [4] is the true ancestor of Theorem 1.1, since the latter is (partly) based on a generalization of the method of proof of the former. An alternative proof, based on viscosity methods, has been given by Demengel, see [14].

In [13], the same author extended her result to cover the case $p_1 < p_N$, under the assumptions

$$p_N < p_1 + 1.$$

The result of [13, Corollary 1.2] still requires $p_1 \geq 2$ and applies to *continuous* local minimizers.

Finally, Lipschitz regularity for solutions of (1.1) has been claimed in the abstract of [8]. However, a closer inspection of the assumptions of Theorem 1.2 there (see [8, equation (1.2)]) shows that their result does not cover the case of (1.1).

Remark 1.4 (A paper by Lieberman). The reader who is familiar with this subject may observe that *apparently* our Theorem 1.1 is already contained in Lieberman's paper [26]. Indeed, [26, Example 1, page 794] deals with exactly the same result for bounded minimizers, by even dropping the requirement $p_1 \geq 2$. However, Lieberman's proof seems to be affected by a crucial flaw. This is a delicate issue, thus we prefer to explain in a clean way the doubtful point of [26].

We first recall that the proof by Lieberman is inspired by Simon's paper [32], dealing with L^∞ gradient estimates for solutions of non-uniformly elliptic equations. One of the crucial tool used by Simon is a generalized version of the Sobolev inequality for functions on manifolds. This is a celebrated result by Michael and Simon himself [31, Theorem 2.1], which in turn generalizes the idea of the cornerstone paper [9] by Bombieri, De Giorgi and Miranda on the *minimal surface equation*.

The idea of [26] is to enlarge the space dimension and identify the set Ω with the flat N -dimensional submanifold $\mathcal{M} := \Omega \times \{(0, \dots, 0)\}$ contained in \mathbb{R}^{2N-1} . Then the author introduces:

- a suitable gradient operator

$$\varphi \mapsto \delta\varphi := \left(\sum_{j=1}^{2N-1} \gamma^{1,j} \varphi_{x_j}, \dots, \sum_{j=1}^{2N-1} \gamma^{2N-1,j} \varphi_{x_j} \right).$$

Here $\gamma = (\gamma^{i,j})$ is a measurable map with values into the set of positive definite symmetric $(2N-1) \times (2N-1)$ matrices;

- a suitable nonnegative measure μ defined on sets of the form $\mathcal{M} \cap E$ for all Borel sets $E \subset \mathbb{R}^{2N-1}$;
- a mean curvature-type operator $H = (H_1, \dots, H_N, H_{N+1}, \dots, H_{2N-1})$ defined on \mathcal{M} .

The key point of [26, Section 4] is to apply the Sobolev-type inequality of Michael and Simon in conjunction with a Caccioppoli inequality for the gradient, in order to circumvent the strong degeneracy of the equation (1.1). However, in order to apply the result by Michael and Simon, some conditions linking the three objects above are needed. Namely, the crucial condition

$$(1.2) \quad \int_{\mathcal{M}} [\delta_i \varphi + H_i \varphi] d\mu = 0, \quad \text{for every } \varphi \in C_0^\infty(U), \quad \text{for every } i = 1, \dots, 2N-1,$$

must be verified, where $\mathcal{M} \subset U \subset \mathbb{R}^{2N-1}$ is an open set. This is condition (1.2) in [31], which is stated to hold true within the framework of Lieberman's paper, see the proof of [26, Proposition 2.1]. However, with the definitions of μ, δ and H taken in [26], one can see that *this crucial condition fails to be verified*. Indeed, with the definitions of [26, Proposition 2.1], for $N+1 \leq i \leq 2N-1$, it holds

$$\gamma^{i,i} = 1 \quad \text{and} \quad \gamma^{i,j} = 0 \text{ for } j \neq i, \quad \text{thus} \quad \delta_i \varphi = \varphi_{x_i},$$

and

$$H_i = 0,$$

while μ coincides with the N -dimensional Lebesgue measure on Ω . Thus condition (1.2) for $N+1 \leq i \leq 2N-1$ becomes

$$\int_{\Omega} \varphi_{x_i}(x, 0, \dots, 0) dx = 0, \quad \text{for every } \varphi \in C_0^\infty(U),$$

which in general *is false*. Thus the proof of [26, Proposition 2.1] does not appear to be correct, leaving in doubt the whole proof of [26, Lemma 4.1], which contains the L^∞ gradient estimate.

1.3. Structure of the proof. The proof of Theorem 1.1 is quite involved, thus we prefer spending a large part of this introduction in order to neatly introduce the main ideas and novelties.

As usual when dealing with higher order regularity, the first issue to be tackled is that the minimizer U lacks the smoothness needed to perform all the necessary manipulations. However, this is a minor issue, which can be easily fixed by approximating our local minimizer U with solutions u_ε of uniformly elliptic problems, see Section 2. The solutions u_ε are as smooth as needed (basically, C^2 regularity is enough) and they converge to our original local minimizer U , as the small regularization parameter $\varepsilon > 0$ converges to 0. Thus it is sufficient to prove “good” a priori estimates on u_ε which are stable when ε goes to 0.

For this reason, in the rest of this subsection we will pretend that our minimizer U is smooth and explain how to get the needed a priori estimates.

The building blocks of Theorem 1.1 are the following two estimates:

A. a local $L^\infty - L^\gamma$ a priori estimate on the gradient, i.e. an estimate of the type

$$(1.3) \quad \|\nabla U\|_{L^\infty(B_r)} \leq C \left(\int_{B_R} |\nabla U|^\gamma dx \right)^{\frac{\Theta}{\gamma}},$$

where $\gamma \geq p_N + 2$ and $\Theta > 1$ are two suitable exponents. This is the content of Proposition 5.1;

B. a local higher integrability estimate *of arbitrary order* on the gradient, i.e. an estimate of the type

$$\int_{B_R} |\nabla U|^q dx \leq C_q,$$

where $1 < q < +\infty$ is arbitrary and $C_q > 0$ is a constant depending only on q , the data of the problem and the local L^∞ norm of U . This is proved in Proposition 6.1.

It is straightforward to see that once **A.** and **B.** are established, then our main result easily follows. We explain how to get both of them:

- in order to obtain **A.** we employ the same method that we successfully applied in [4, Theorem 1.1], for the standard growth case $p_1 = \dots = p_N = p$. This is based on a new class of Caccioppoli-type inequalities for ∇U , which have been first introduced by the two authors in [2] and then generalized and exploited in its full generality in [4].

In a nutshell, the idea is to take the equation satisfied by U

$$\sum_{i=1}^N \left(|U_{x_i}|^{p_i-2} U_{x_i} \right)_{x_i} = 0,$$

differentiate it with respect to x_j and then insert weird test functions of the form

$$\Phi(U_{x_j}) \Psi(U_{x_k}),$$

with $k, j \in \{1, \dots, N\}$. With these new Caccioppoli-type inequalities at hand, we can follow the same scheme as in [4, Proposition 5.1] and obtain (1.3).

We point out that, apart from a number of technical complications linked to the fact that $p_1 \neq p_N$, in the present setting there is a crucial difference with the case treated in [4]. Indeed, after a Moser-type iteration, there we obtained an a priori estimate of the type

$$(1.4) \quad \|\nabla U\|_{L^\infty(B_r)} \leq C \left(\int_{B_R} |\nabla U|^{p+2} dx \right)^{\frac{1}{p+2}}.$$

Apparently, in that case as well we needed the further higher integrability information $\nabla U \in L_{\text{loc}}^{p+2}$. However, thanks to the homogeneity of the estimate (1.4), one can use a standard interpolation trick (see the Step 4 of the proof of [4, Proposition 5.1]) and upgrade (1.4) to the following

$$\|\nabla U\|_{L^\infty(B_r)} \leq C \left(\int_{B_R} |\nabla U|^p dx \right)^{\frac{1}{p}}.$$

This does not require any prior integrability information on ∇U beyond the natural growth exponent. Thus in the standard growth case, we are dispensed with point **B.**, i.e. point **A.** is enough to conclude. On the contrary, in our case, the same trick does not apply to estimate (1.3), because of the presence of the exponent $\Theta > 1$. For this reason we need a higher integrability information on ∇U .

In order to have a better understanding of the proof described above, we refer the interested reader to the Introduction of [4], where the whole strategy for point **A.** is explained in details;

- part **B.** is the really involved point of the whole proof. At first, we point out that we do not have a good control on the exponent γ appearing in (1.3) (unless some restrictions on the ratio p_N/p_1 are imposed). For this reason, we need to gain as much integrability on ∇U as possible. This is a classical subject in the regularity theory for functionals with nonstandard growth conditions, i.e. integrability gain on the gradients of minimizers.

Since in general minimizers of this kind of functionals may be very irregular when p_1 and p_N are too far apart, usually one needs to impose some restrictions on the ratio p_N/p_1 to get some regularity. These restriction are typically of the type

$$\frac{p_N}{p_1} < c_N, \quad \text{for some constant } c_N > 0 \text{ such that } \lim_{N \rightarrow \infty} c_N = 1.$$

If one further supposes *local minimizers to be bounded*, then the previous restriction can be relaxed to conditions of the type

$$\frac{p_N}{p_1} < C \quad \text{or} \quad p_N < p_1 + C,$$

with a *universal* constant $C > 0$. In any case, to the best of our knowledge all the results appearing in the literature require some upper bound on the ratio p_N/p_1 . More precisely, all the results except one: in the very interesting paper [7] by Bildhauer, Fuchs and Zhong, the authors consider a functional with nonstandard growth of the type

$$(1.5) \quad (u, \Omega') \mapsto \int_{\Omega'} \left(\sum_{i=1}^{N-1} |u_{x_i}|^2 \right)^{\frac{p_1}{2}} dx + \int_{\Omega'} |u_{x_N}|^{p_2} dx, \quad \text{with } p_1 \leq p_2,$$

and prove that any local minimizer $u \in L_{\text{loc}}^\infty$ is such that $\nabla u \in L_{\text{loc}}^q$ for every $1 < q < +\infty$, *no matter how large the ratio p_2/p_1 is*, see [7, Theorem 1.1]. The idea of [7] is partially inspired from Choe's result [10, Theorem 3], which in turn seems to find its roots in DiBenedetto's paper [15] (see [15, Proposition 3.1]). It relies on a suitable integration by parts in conjunction with the Caccioppoli inequality for ∇u . For functionals as in (1.5), this leads to an iterative scheme of the type

$$\text{"gain of integrability on } u_{x_N} \text{"} \implies \text{"gain of integrability on } (u_{x_1}, \dots, u_{x_{N-1}}) \text{"}$$

and viceversa

$$\text{"gain of integrability on } (u_{x_1}, \dots, u_{x_{N-1}}) \text{"} \implies \text{"gain of integrability on } u_{x_N} \text{"}.$$

By means of a *doubly recursive scheme* which is quite difficult to handle, [7] exploits the full power of the above approach to avoid any unnecessary restriction on the exponents. In [10] instead, the gain of integrability was extremely simplified, at the price of taking the assumption

$$p_N < p_1 + 1.$$

Incidentally, we point out that this is the same assumption as in the aforementioned paper [13], which uses however different techniques.

We will try to detail the main difficulties of this method in a while. Before this, we point out that (1.5) is only concerned with two growth exponents. Moreover, the type of degeneracy of the functional (1.5) is much lighter than that of our functional \mathfrak{F}_p . For these reasons, even if our strategy is greatly inspired by that of [7], all the estimates have to be recast and the resulting iterative scheme becomes of far reaching complexity.

We now come to explain such an iterative scheme: by proceeding as in [10] and [7], we get an estimate of the type (see Proposition 4.3)

$$(1.6) \quad \int_{B_{r_0}} |U_{x_k}|^{p_k+2+\alpha} dx \leq C + C \sum_{i \neq k} \int_{B_{R_0}} |U_{x_i}|^{\frac{p_i-2}{p_k}(p_k+2+\alpha)} dx, \quad \text{for } k = 1, \dots, N,$$

where $C > 0$ depends on the data and on the local L^∞ norm of U , as well. Here the free parameter $\alpha \geq 0$ has to be carefully chosen, in order to improve the gradient summability. We observe that, technically speaking, *this scheme is not of Moser-type*. Indeed, the key point of (1.6) is that it entails estimates on a fixed component U_{x_j} , in terms of all the others.

• **First step** We start by using (1.6) as follows: we take $k = N$ in (1.6) and choose $\alpha \geq 0$ in such a way that

$$\frac{p_i - 2}{p_N} (p_N + 2 + \alpha) \leq p_i, \quad \text{for } i = 1, \dots, N-1.$$

It is possible to make such a choice *without imposing restrictions on p_N/p_1* , the optimal choice being¹

$$p_N + 2 + \alpha_N^{(0)} = p_N \min_{1 \leq i \leq N-1} \frac{p_i}{p_i - 2} =: p_N q_{N-1}.$$

This permits to upgrade the integrability of U_{x_N} to $L_{\text{loc}}^{p_N q_{N-1}}$. This is the end of the first step.

• **Second step** Once we gain this property on U_{x_N} , we shift to $U_{x_{N-1}}$: we take $k = N-1$ in (1.6), that we write in the following form

$$\begin{aligned} \int_{B_{r_0}} |U_{x_k}|^{p_k+2+\alpha} dx &\leq C + C \sum_{i=1}^{N-2} \int_{B_{R_0}} |U_{x_i}|^{\frac{p_i-2}{p_k}(p_k+2+\alpha)} dx \\ &\quad + C \int_{B_{R_0}} |U_{x_N}|^{\frac{p_N-2}{p_k}(p_k+2+\alpha)} dx. \end{aligned}$$

Then by using that

$$U_{x_i} \in L_{\text{loc}}^{p_i}, \quad \text{for } i = 1, \dots, N-2 \quad \text{and} \quad U_{x_N} \in L_{\text{loc}}^{p_N q_{N-1}},$$

we choose α in such a way that

$$\begin{cases} \frac{p_i - 2}{p_{N-1}} (p_{N-1} + 2 + \alpha) \leq p_i, & \text{for } i = 1, \dots, N-2, \\ \frac{p_N - 2}{p_{N-1}} (p_{N-1} + 2 + \alpha) \leq p_N q_{N-1} \end{cases}$$

If we set as above

$$q_i = \frac{p_i}{p_i - 2},$$

the optimal choice is now

$$p_{N-1} + 2 + \alpha_{N-1}^{(0)} = p_{N-1} \min \left\{ q_N q_{N-1}, \min_{1 \leq i \leq N-2} q_i \right\}.$$

However, *this is not the end* of the second step. Indeed, rather than applying (1.6) directly to the other components $U_{x_{N-2}}, \dots, U_{x_1}$ as above, we come back to U_{x_N} .

More precisely, we apply (1.6) to U_{x_N} taking into account the new information on $U_{x_{N-1}}$. This gives higher integrability for U_{x_N} . We next apply alternatively (1.6) to $U_{x_{N-1}}$ and U_{x_N} , taking into

¹For ease of presentation, in what follows we assume that $p_i > 2$ for every $i = 1, \dots, N$.

account the higher integrability gain at each step. After a finite number of iterations, it can be established that

$$U_{x_N} \in L_{\text{loc}}^{p_N q_{N-2}} \quad \text{and} \quad U_{x_{N-1}} \in L_{\text{loc}}^{p_{N-1} q_{N-2}}.$$

This is the end of the second step.

- **j -th step** (for $2 \leq j \leq N-1$) When we land on this step, we have iteratively acquired the following knowledge

$$U_{x_i} \in L_{\text{loc}}^{p_i}, \quad \text{for } i = 1, \dots, N-j \quad \text{and} \quad U_{x_i} \in L_{\text{loc}}^{p_N q_{N-j+1}}, \quad \text{for } i = N-j+2, \dots, N.$$

We then start to get into play the component $U_{x_{N-j+1}}$. We take $k = N-j+1$ in (1.6) and we choose α in such a way that

$$\begin{cases} \frac{p_i - 2}{p_{N-j+1}} (p_{N-j+1} + 2 + \alpha) \leq p_i, & \text{for } i = 1, \dots, N-j, \\ \frac{p_i - 2}{p_{N-j+1}} (p_{N-j+1} + 2 + \alpha) \leq p_i q_{N-j+1}, & \text{for } i = N-j+2, \dots, N. \end{cases}$$

This permits to infer that

$$U_{x_{N-j+1}} \in L_{\text{loc}}^{p_{N-j+1} + 2 + \alpha_{N-j+1}^{(0)}},$$

where

$$p_{N-j+1} + 2 + \alpha_{N-j+1}^{(0)} = p_{N-j+1} \min \left\{ \min_{i=1, \dots, N-j} q_i, q_{N-j+1} \min_{i=N-j+2, \dots, N} q_i \right\}.$$

As illustrated in the second step, we now use this information and start to cyclically use (1.6) on $U_{x_N}, U_{x_{N-1}}, \dots, U_{x_{N-j}}$, in order to improve their integrability. After a finite number of iterations of this algorithm, we obtain

$$U_{x_i} \in L_{\text{loc}}^{p_i q_{N-j+1}}, \quad \text{for } i = N-j+2, \dots, N.$$

This is the end of the j -th step.

- **Last step.** We fix $q_0 \geq 2$ arbitrary. We finally consider the last component U_{x_1} , as well. By using the starting information

$$U_{x_i} \in L_{\text{loc}}^{p_i q_1}, \quad \text{for } i = 2, \dots, N,$$

and proceeding as above, we finally get

$$U_{x_i} \in L_{\text{loc}}^{p_i q_0}, \quad \text{for } 1 \leq i \leq N.$$

This yields the desired conclusion.

The main difficulty of **B.** is to prove that this algorithm does not require any restriction on the exponents p_i , and that each step *ends up after a finite number of loops*.

1.4. Plan of the paper. In Section 2 the reader will find the approximating scheme and all the basic material needed to understand the sequel of the paper. Section 3 contains the crucial Caccioppoli-type inequalities for the gradient, needed to build up the Moser's scheme for point **A.** of the strategy presented above. Then in Section 4, we prove integral estimates for the gradient: the first one is a Caccioppoli inequality for power functions of the gradient (Proposition 4.2), while the second one is the self-improving scheme *à la* Bildhauer-Fuchs-Zhong (Proposition 4.3). With Sections 5 and 6, we enter into the core of the paper: they contain the $L^\infty - L^\gamma$ gradient estimate and the higher integrability estimate for the gradient, respectively. Then in the short Section 7, we eventually prove our main result.

Two technical appendices conclude the paper: they contain the study of all the intricate sequences of real numbers needed in this paper.

Acknowledgments. Part of this work has been done during a visit of P. B. to Bologna & Ferrara in February 2018 and during a visit of L. B. to Toulouse in June 2018. The latter has been financed by the ANR project “*Entropies, Flots, Inégalités*”, we wish to thank Max Fathi. Hosting institutions are kindly acknowledged.

2. PRELIMINARIES

We will use the same approximation scheme as in [3, Section 2] and [6, Section 5]. We recall that are interested in local minimizers of the following variational integral

$$\mathfrak{F}_{\mathbf{p}}(u; \Omega') = \sum_{i=1}^N \frac{1}{p_i} \int_{\Omega'} |u_{x_i}|^{p_i} dx, \quad u \in W_{\text{loc}}^{1,\mathbf{p}}(\Omega), \quad \Omega' \Subset \Omega,$$

where $\mathbf{p} = (p_1, \dots, p_N)$ and $2 \leq p_1 \leq \dots \leq p_N$. In the rest of the paper, we fix $U \in W_{\text{loc}}^{1,\mathbf{p}}(\Omega)$ a local minimizer of $\mathfrak{F}_{\mathbf{p}}$. We also fix a ball

$$B \Subset \Omega \quad \text{such that} \quad 2B \Subset \Omega \text{ as well.}$$

We use the usual notation λB to denote the ball concentric with B , scaled by a factor $\lambda > 0$. Since we have the continuous inclusion $W^{1,\mathbf{p}}(2B) \subset W^{1,p_1}(2B)$, by Poincaré inequality it holds

$$U \in L^{p_1}(2B).$$

For every $0 < \varepsilon \ll 1$ and every $x \in \overline{B}$, we set $U_\varepsilon(x) = U * \varrho_\varepsilon(x)$, where ϱ_ε is the usual family of Friedrichs mollifiers, supported in a ball of radius ε centered at the origin. We also set

$$(2.1) \quad g_{i,\varepsilon}(t) = \frac{1}{p_i} |t|^{p_i} + \frac{\varepsilon}{2} t^2, \quad t \in \mathbb{R}, \quad i = 1, \dots, N.$$

Finally, we define the regularized functional

$$\mathfrak{F}_{\mathbf{p},\varepsilon}(v; B) = \sum_{i=1}^N \int_B g_{i,\varepsilon}(v_{x_i}) dx.$$

The following preliminary result is standard, see [3, Lemma 2.5 and Lemma 2.8].

Lemma 2.1 (Basic energy estimate). *There exists $0 < \varepsilon_0 < 1$ such that for every $0 < \varepsilon \leq \varepsilon_0$, the problem*

$$(2.2) \quad \min \left\{ \mathfrak{F}_{\mathbf{p},\varepsilon}(v; B) : v - U_\varepsilon \in W_0^{1,\mathbf{p}}(B) \right\},$$

admits a unique solution u_ε . Moreover, the following uniform estimate holds

$$\sum_{i=1}^N \frac{1}{p_i} \int_B |(u_\varepsilon)_{x_i}|^{p_i} dx \leq \left(\sum_{i=1}^N \frac{1}{p_i} \int_{2B} |U_{x_i}|^{p_i} dx + \frac{\varepsilon_0}{2} \int_{2B} |\nabla U|^2 dx \right).$$

Finally, $u_\varepsilon \in C^2(B)$.

Proof. The only difference with respect to [3] is on the uniform energy estimate, due to the nonstandard growth conditions. We show how to obtain this: it is sufficient to test the minimality of u_ε against U_ε , this

gives

$$\begin{aligned} \sum_{i=1}^N \frac{1}{p_i} \int_B |(u_\varepsilon)_{x_i}|^{p_i} dx &\leq \sum_{i=1}^N \frac{1}{p_i} \int_B |(U * \varrho_\varepsilon)_{x_i}|^{p_i} dx + \frac{\varepsilon}{2} \int_B |\nabla(U * \varrho_\varepsilon)|^2 dx \\ &\leq \|\varrho_\varepsilon\|_{L^1(\mathbb{R}^N)} \left(\sum_{i=1}^N \frac{1}{p_i} \int_{2B} |U_{x_i}|^{p_i} dx + \frac{\varepsilon}{2} \int_{2B} |\nabla U|^2 dx \right). \end{aligned}$$

By using the scaling properties of the family of mollifiers, we get the conclusion. \square

As usual, we will also rely on the following convergence result.

Lemma 2.2 (Convergence to a minimizer). *With the same notation as above, we have*

$$(2.3) \quad \lim_{\varepsilon \rightarrow 0} \left[\|u_\varepsilon - U\|_{L^{p_1}(B)} + \sum_{i=1}^N \|(u_\varepsilon - U)_{x_i}\|_{L^{p_i}(B)} \right] = 0.$$

Proof. We observe that $u_\varepsilon - U_\varepsilon \in W_0^{1,\mathbf{p}}(B)$ and the set B is bounded in every direction. Thus by Poincaré inequality, we have

$$(2.4) \quad \int_B |u_\varepsilon - U_\varepsilon|^{p_i} \leq C_i |B|^{-\frac{p_i}{N}} \int_B |(u_\varepsilon - U_\varepsilon)_{x_i}|^{p_i} dx, \quad i = 1, \dots, N,$$

for some $C_i = C_i(N, p_i) > 0$. For $i = 1$, this in turn gives

$$\begin{aligned} \|u_\varepsilon\|_{L^{p_1}(B)} &\leq \|u_\varepsilon - U_\varepsilon\|_{L^{p_1}(B)} + \|U_\varepsilon\|_{L^{p_1}(B)} \\ &\leq C \|(u_\varepsilon - U_\varepsilon)_{x_1}\|_{L^{p_1}(B)} + \|U_\varepsilon\|_{L^{p_1}(B)} \\ &\leq C \|(u_\varepsilon)_{x_1}\|_{L^{p_1}(B)} + C \|U\|_{W^{1,p_1}(2B)}, \end{aligned}$$

for a constant $C = C(N, p_1) > 0$. By Lemma 2.1, the last term is uniformly bounded for $0 < \varepsilon \leq \varepsilon_0$. Thus the family $\{u_\varepsilon\}_{0 < \varepsilon \leq \varepsilon_0}$ is bounded in $W^{1,p_1}(B)$. We can infer the weak convergence in $W^{1,p_1}(B)$ of a subsequence $\{u_{\varepsilon_k}\}_{k \in \mathbb{N}}$ to a function $u \in W^{1,p_1}(B)$. This convergence is strong in $L^{p_1}(B)$, by the Rellich-Kondrašov Theorem.

For every $\varphi \in W_0^{1,\mathbf{p}}(B)$, we test the minimality of u_{ε_k} against $\varphi + U_{\varepsilon_k}$. Thus, by lower semicontinuity of the L^{p_i} norms on $L^{p_1}(B)$, we can infer

$$\begin{aligned} \sum_{i=1}^N \frac{1}{p_i} \int_B |u_{x_i}|^{p_i} dx &\leq \liminf_{k \rightarrow +\infty} \sum_{i=1}^N \frac{1}{p_i} \int_B |(u_{\varepsilon_k})_{x_i}|^{p_i} dx \\ (2.5) \quad &\leq \lim_{k \rightarrow +\infty} \sum_{i=1}^N \frac{1}{p_i} \int_B |(\varphi + U_{\varepsilon_k})_{x_i}|^{p_i} dx + \frac{\varepsilon_k}{2} \int_B |\nabla \varphi + \nabla U_{\varepsilon_k}|^2 dx \\ &= \sum_{i=1}^N \frac{1}{p_i} \int_B |(\varphi + U)_{x_i}|^{p_i} dx. \end{aligned}$$

This shows that $u_{x_i} \in L^{p_i}(B)$ for $i = 1, \dots, N$ and u solves

$$\min \left\{ \mathfrak{F}_{\mathbf{p}}(v; B) : v - U \in W_0^{1,\mathbf{p}}(B) \right\}.$$

By strict convexity of the functional $\mathfrak{F}_{\mathbf{p}}$, we thus obtain $u = U$.

We can now take $\varphi \equiv 0$ in (2.5). Since we know that $u = U$, we have equality everywhere in (2.5), thus in particular

$$(2.6) \quad \lim_{k \rightarrow +\infty} \sum_{i=1}^N \frac{1}{p_i} \int_B |(u_{\varepsilon_k})_{x_i}|^p dx = \sum_{i=1}^N \frac{1}{p_i} \int_B |U_{x_i}|^p dx, \quad i = 1, \dots, N.$$

We next observe that the weak convergence of $\{(u_{\varepsilon_k})_{x_i}\}_{k \in \mathbb{N}}$ to U_{x_i} and the lower semicontinuity of the L^{p_i} norm on L^{p_i} imply that

$$(2.7) \quad \liminf_{k \rightarrow +\infty} \int_B \left| \frac{(u_{\varepsilon_k})_{x_i} + U_{x_i}}{2} \right|^{p_i} dx \geq \int_B |U_{x_i}|^{p_i} dx, \quad i = 1, \dots, N.$$

Moreover, by Clarkson's inequality for $p_i \geq 2$, one has

$$\left\| \frac{(u_{\varepsilon_k})_{x_i} + U_{x_i}}{2} \right\|_{L^{p_i}(B)}^{p_i} + \left\| \frac{(u_{\varepsilon_k})_{x_i} - U_{x_i}}{2} \right\|_{L^{p_i}(B)}^{p_i} \leq \frac{1}{2} \left(\|(u_{\varepsilon_k})_{x_i}\|_{L^{p_i}(B)}^{p_i} + \|U_{x_i}\|_{L^{p_i}(B)}^{p_i} \right).$$

We divide by p_i , sum over $i = 1, \dots, N$ and rely on (2.6) and (2.7) to obtain

$$\lim_{k \rightarrow +\infty} \sum_{i=1}^N \frac{1}{p_i} \|(u_{\varepsilon_k})_{x_i} - U_{x_i}\|_{L^{p_i}(B)}^{p_i} = 0.$$

By using this into (2.4) with $i = 1$ and using the strong convergence of U_{ε_k} to U , we get

$$\lim_{k \rightarrow +\infty} \left[\|u_{\varepsilon_k} - U\|_{L^{p_1}(B)} + \sum_{i=1}^N \|(u_{\varepsilon_k})_{x_i} - U_{x_i}\|_{L^{p_i}(B)} \right] = 0.$$

Finally, we observe that we can repeat this argument with any subsequence of the original family $\{(u_{\varepsilon})_{\varepsilon > 0}\}$. Thus the above limit holds true for the whole family $\{u_{\varepsilon}\}_{0 < \varepsilon \leq \varepsilon_0}$ instead of $\{u_{\varepsilon_k}\}_{k \in \mathbb{N}}$ and (2.3) follows. \square

We recall that our main result Theorem 1.1 is valid for *bounded* local minimizers. Thus, the following simple uniform L^∞ estimate will be crucial.

Proposition 2.3 (Uniform L^∞ estimate). *With the notation above, let us further assume that $U \in L^\infty(2B)$. Then for every $0 < \varepsilon \leq \varepsilon_0$ we have*

$$\|u_\varepsilon\|_{L^\infty(B)} \leq \|U\|_{L^\infty(2B)}.$$

Proof. By the maximum principle, see for example [33, Theorem 2.1], we have

$$\max_B |u_\varepsilon| = \max_{\partial B} |u_\varepsilon| = \max_{\partial B} |U_\varepsilon| \leq \max_{\overline{B}} |U_\varepsilon|.$$

By recalling the construction of U_ε and using the hypothesis on U , we get the desired conclusion. \square

As in [4], the following standard technical result will be useful. The proof can be found in [22, Lemma 6.1], for example.

Lemma 2.4. *Let $0 < r < R$ and let $Z(t) : [r, R] \rightarrow [0, \infty)$ be a bounded function. Assume that for $r \leq t < s \leq R$ we have*

$$Z(t) \leq \frac{\mathcal{A}}{(s-t)^{\alpha_0}} + \frac{\mathcal{B}}{(s-t)^{\beta_0}} + \mathcal{C} + \vartheta Z(s),$$

with $\mathcal{A}, \mathcal{B}, \mathcal{C} \geq 0$, $\alpha_0 \geq \beta_0 > 0$ and $0 \leq \vartheta < 1$. Then we have

$$Z(r) \leq \left(\frac{1}{(1-\lambda)^{\alpha_0}} \frac{\lambda^{\alpha_0}}{\lambda^{\alpha_0} - \vartheta} \right) \left[\frac{\mathcal{A}}{(R-r)^{\alpha_0}} + \frac{\mathcal{B}}{(R-r)^{\beta_0}} + \mathcal{C} \right],$$

where λ is any number such that

$$\vartheta^{\frac{1}{\alpha_0}} < \lambda < 1.$$

3. CACCIOPPOLI-TYPE INEQUALITIES

The solution u_ε of the problem (2.2) satisfies the Euler-Lagrange equation

$$(3.1) \quad \sum_{i=1}^N \int g'_{i,\varepsilon}((u_\varepsilon)_{x_i}) \varphi_{x_i} dx = 0, \quad \text{for every } \varphi \in W_0^{1,\mathbf{p}}(B).$$

From now on we will systematically suppress the subscript ε on u_ε and *simply write* u .

In order to prove the gradient regularity, we need the equation satisfied by the gradient ∇u . Thus we insert a test function of the form $\varphi = \psi_{x_j} \in W_0^{1,\mathbf{p}}(B)$ in (3.1), compactly supported in B . After an integration by parts, we get

$$(3.2) \quad \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) u_{x_i x_j} \psi_{x_i} dx = 0,$$

for $j = 1, \dots, N$. We thus found the equation solved by u_{x_j} . Observe that we are legitimate to integrate by parts, since $u \in C^2(B)$ by Lemma 2.1.

The following Caccioppoli inequality can be proved exactly as [3, Lemma 3.2], we omit the details.

Lemma 3.1. *Let $\Phi : \mathbb{R} \rightarrow \mathbb{R}^+$ be a C^1 convex function. Then there exists a constant $C = C(\mathbf{p}) > 0$ such that for every function $\eta \in C_0^\infty(B)$ and every $j = 1, \dots, N$, we have*

$$(3.3) \quad \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) \left| (\Phi(u_{x_j}))_{x_i} \right|^2 \eta^2 dx \leq C \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) |\Phi(u_{x_j})|^2 \eta_{x_i}^2 dx.$$

Actually, we can drop the requirement that Φ has to be convex, under some circumstances. The resulting Caccioppoli inequality is of interest.

Lemma 3.2. *Let $-1 < \alpha \leq 0$. For every function $\eta \in C_0^\infty(B)$ and every $j = 1, \dots, N$, we have*

$$\sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) u_{x_i x_j}^2 |u_{x_j}|^\alpha \eta^2 dx \leq \frac{4}{(1+\alpha)^2} \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) |u_{x_j}|^{\alpha+2} |\eta_{x_i}|^2 dx.$$

When $\alpha < 0$, in the left hand side of the above inequality, the quantity $u_{x_i x_j}^2 |u_{x_j}|^\alpha$ is defined to be 0 on the set where u_{x_j} vanishes.

Proof. Let $\kappa > 0$. We take in (3.2) the test function

$$\psi = u_{x_j} (\kappa + |u_{x_j}|^2)^{\frac{\alpha}{2}} \eta^2,$$

where η is as in the statement. We get

$$\begin{aligned} & \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) u_{x_i x_j}^2 (\kappa + |u_{x_j}|^2)^{\frac{\alpha}{2}} \eta^2 dx \\ & + \alpha \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) u_{x_i x_j}^2 (\kappa + |u_{x_j}|^2)^{\frac{\alpha-2}{2}} |u_{x_j}|^2 \eta^2 dx \\ & = -2 \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) u_{x_i x_j} (\kappa + |u_{x_j}|^2)^{\frac{\alpha}{2}} u_{x_j} \eta \eta_{x_i} dx. \end{aligned}$$

We observe that

$$(\kappa + |u_{x_j}|^2)^{\frac{\alpha-2}{2}} |u_{x_j}|^2 \leq (\kappa + |u_{x_j}|^2)^{\frac{\alpha}{2}} \quad \text{and} \quad (\kappa + |u_{x_j}|^2)^{\frac{\alpha}{2}} |u_{x_j}| \leq (\kappa + |u_{x_j}|^2)^{\frac{\alpha+1}{2}}.$$

From the previous identity, we get (remember that α is non positive)

$$(1 + \alpha) \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) u_{x_i x_j}^2 (\kappa + |u_{x_j}|^2)^{\frac{\alpha}{2}} \eta^2 dx \leq 2 \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) |u_{x_i x_j}| (\kappa + |u_{x_j}|^2)^{\frac{\alpha+1}{2}} |\eta| |\eta_{x_i}| dx.$$

By using Young's inequality, we can absorb the Hessian term in the right-hand side, to get

$$\sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) u_{x_i x_j}^2 (\kappa + |u_{x_j}|^2)^{\frac{\alpha}{2}} \eta^2 dx \leq \frac{4}{(1 + \alpha)^2} \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) (\kappa + |u_{x_j}|^2)^{\frac{\alpha+2}{2}} |\eta_{x_i}|^2 dx.$$

By taking the limit as κ goes to 0 on both sides, and using Fatou Lemma on the left-hand side and the Dominated Convergence Theorem in the right-hand side, we get the conclusion. \square

As in the standard growth case $p_1 = \dots = p_N = p$, a key role is played by the following sophisticated Caccioppoli-type inequality for the gradient. The proof is the same as that of [4, Proposition 3.2] and we omit it. It is sufficient to observe that the proof in [4] does not depend on the particular form of the functions $g_{i,\varepsilon}$.

Proposition 3.3 (Weird Caccioppoli inequality). *Let $\Phi, \Psi : [0, +\infty) \rightarrow [0, +\infty)$ be two non-decreasing continuous functions. We further assume that Ψ is convex and C^1 . Let $\eta \in C_0^\infty(B)$ and $0 \leq \theta \leq 2$, then for every $k, j = 1, \dots, N$,*

$$\begin{aligned} & \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) u_{x_i x_j}^2 \Phi(u_{x_j}^2) \Psi(u_{x_k}^2) \eta^2 dx \\ & \leq C \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) u_{x_j}^2 \Phi(u_{x_j}^2) \Psi(u_{x_k}^2) |\nabla \eta|^2 dx \\ & + C \left(\sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) u_{x_i x_j}^2 u_{x_j}^2 \Phi(u_{x_j}^2)^2 \Psi'(u_{x_k}^2)^\theta \eta^2 dx \right)^{\frac{1}{2}} \\ & \times \left(\sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) |u_{x_k}|^{2\theta} \Psi(u_{x_k}^2)^{2-\theta} |\nabla \eta|^2 dx \right)^{\frac{1}{2}}. \end{aligned} \tag{3.4}$$

4. LOCAL ENERGY ESTIMATES FOR THE REGULARIZED PROBLEM

4.1. Towards an iterative Moser's scheme. We recall that

$$(4.1) \quad g''_{i,\varepsilon}(t) = (p_i - 1) |t|^{p_i-2} + \varepsilon.$$

We use Proposition 3.3 with the following choices

$$(4.2) \quad \Phi(t) = t^{s-1} \quad \text{and} \quad \Psi(t) = t^m, \quad \text{for } t \geq 0,$$

with $1 \leq s \leq m$. The proof of the following result is exactly the same as that of [4, Proposition 4.1].

Proposition 4.1 (Staircase to the full Caccioppoli). *Let $2 \leq p_1 \leq p_2 \leq \dots \leq p_N$ and let $\eta \in C_0^\infty(B)$. Then for every $k, j = 1, \dots, N$ and $1 \leq s \leq m$,*

$$(4.3) \quad \begin{aligned} \sum_{i=1}^N \int_{\Omega} g''_{i,\varepsilon}(u_{x_i}) u_{x_i x_j}^2 |u_{x_j}|^{2s-2} |u_{x_k}|^{2m} \eta^2 dx &\leq C \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) |u_{x_j}|^{2s+2m} |\nabla \eta|^2 dx \\ &+ C(m+1) \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) |u_{x_k}|^{2s+2m} |\nabla \eta|^2 dx \\ &+ \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) u_{x_i x_j}^2 |u_{x_j}|^{4s-2} |u_{x_k}|^{2m-2s} \eta^2 dx. \end{aligned}$$

By iterating a finite number of times the previous estimate, we get the following

Proposition 4.2 (Caccioppoli for power functions). *Take an exponent q of the form*

$$q = 2^{\ell_0} - 1, \quad \text{for a given } \ell_0 \in \mathbb{N} \setminus \{0\}.$$

Let $2 \leq p_1 \leq p_2 \leq \dots \leq p_N$ and let $\eta \in C_0^\infty(B)$. Then for every $k = 1, \dots, N$, we have

$$(4.4) \quad \begin{aligned} \int \left| \nabla \left(|u_{x_k}|^{q+\frac{p_k-2}{2}} u_{x_k} \right) \right|^2 \eta^2 dx &\leq C q^5 \sum_{i,j=1}^N \int g''_{i,\varepsilon}(u_{x_i}) |u_{x_j}|^{2q+2} |\nabla \eta|^2 dx \\ &+ C q^5 \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) |u_{x_k}|^{2q+2} |\nabla \eta|^2 dx, \end{aligned}$$

for some $C = C(N, p_k) > 0$.

Proof. The proof is essentially the same as that of [4, Proposition 4.2]. We define the two finite families of indices $\{s_\ell\}$ and $\{m_\ell\}$ through

$$s_\ell = 2^\ell, \quad m_\ell = q + 1 - 2^\ell, \quad \ell \in \{0, \dots, \ell_0\}.$$

By definition, we have

$$\begin{aligned} 1 \leq s_\ell \leq m_\ell, \quad \ell \in \{0, \dots, \ell_0 - 1\}, \\ s_\ell + m_\ell = q + 1, \quad \ell \in \{0, \dots, \ell_0\}, \\ 4s_\ell - 2 = 2s_{\ell+1} - 2, \quad 2m_\ell - 2s_\ell = 2m_{\ell+1}, \end{aligned}$$

and

$$s_0 = 1, \quad m_0 = q, \quad s_{\ell_0} = 2^{\ell_0}, \quad m_{\ell_0} = 0.$$

From inequality (4.3), we get for every $\ell \in \{0, \dots, \ell_0 - 1\}$,

$$\begin{aligned} \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) u_{x_i x_j}^2 |u_{x_j}|^{2s_\ell-2} |u_{x_k}|^{2m_\ell} \eta^2 dx \\ \leq C \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) |u_{x_j}|^{2q+2} |\nabla \eta|^2 dx \\ + C(m_\ell + 1) \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) |u_{x_k}|^{2q+2} |\nabla \eta|^2 dx \\ + \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) u_{x_i x_j}^2 |u_{x_j}|^{2s_{\ell+1}-2} |u_{x_k}|^{2m_{\ell+1}} \eta^2 dx, \end{aligned}$$

for some $C > 0$ universal. By starting from $\ell = 0$ and iterating the previous estimate up to $\ell = \ell_0 - 1$, we then get

$$\begin{aligned} \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) u_{x_i x_j}^2 |u_{x_k}|^{2q} \eta^2 dx &\leq C q^2 \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) |u_{x_j}|^{2q+2} |\nabla \eta|^2 dx \\ &\quad + C q^2 \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) |u_{x_k}|^{2q+2} |\nabla \eta|^2 dx \\ &\quad + \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) u_{x_i x_j}^2 |u_{x_j}|^{2q} \eta^2 dx, \end{aligned}$$

for a universal constant $C > 0$. For the last term, we apply the Caccioppoli inequality (3.3) with

$$\Phi(t) = \frac{|t|^{q+1}}{q+1}, \quad t \in \mathbb{R},$$

thus we get

$$\begin{aligned} \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) u_{x_i x_j}^2 |u_{x_k}|^{2q} \eta^2 dx &\leq C q^2 \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) |u_{x_j}|^{2q+2} |\nabla \eta|^2 dx \\ &\quad + C q^2 \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) |u_{x_k}|^{2q+2} |\nabla \eta|^2 dx \\ &\quad + \frac{C}{(q+1)^2} \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) |u_{x_j}|^{2q+2} |\nabla \eta|^2 dx; \end{aligned}$$

that is,

$$\begin{aligned} \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) u_{x_i x_j}^2 |u_{x_k}|^{2q} \eta^2 dx &\leq C q^2 \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) |u_{x_j}|^{2q+2} |\nabla \eta|^2 dx \\ &\quad + C q^2 \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) |u_{x_k}|^{2q+2} |\nabla \eta|^2 dx, \end{aligned} \tag{4.5}$$

possibly for a different universal constant $C > 0$.

We now recall (4.1), thus we get

$$\begin{aligned} \sum_{i=1}^N \int g''_{i,\varepsilon}(u_{x_i}) u_{x_i x_j}^2 |u_{x_k}|^{2q} \eta^2 dx &\geq \int |u_{x_k}|^{p_k-2} u_{x_k x_j}^2 |u_{x_k}|^{2q} \eta^2 dx \\ &= \left(\frac{2}{2q + p_k} \right)^2 \int \left| \left(|u_{x_k}|^{q + \frac{p_k-2}{2}} u_{x_k} \right)_{x_j} \right|^2 \eta^2 dx. \end{aligned}$$

We can sum over $j = 1, \dots, N$ to obtain

$$\sum_{i,j=1}^N \int g''_{i,\varepsilon}(u_{x_i}) u_{x_i x_j}^2 |u_{x_k}|^{2q} \eta^2 dx \geq \left(\frac{2}{2q + p_k} \right)^2 \int |\nabla \left(|u_{x_k}|^{q + \frac{p_k-2}{2}} u_{x_k} \right)|^2 \eta^2 dx.$$

This proves the desired inequality. \square

4.2. Towards higher integrability. In order to prove the higher integrability of the gradient, we will need the following self-improving estimate. This is analogous to the estimate at the basis of [7, Theorem 1.1], which deals with the case $p_1 = \dots = p_{N-1} < p_N$ only. As pointed out in the Introduction, our case will be much more involved.

Proposition 4.3. *For every $\alpha > -1$ and every $k = 1, \dots, N$, there exists a constant $C = C(N, p_k, \alpha) > 0$ such that for every pair of concentric balls $B_{r_0} \subset B_{R_0} \Subset B$, we have*

$$(4.6) \quad \begin{aligned} \int_{B_{r_0}} |u_{x_k}|^{p_k+2+\alpha} dx &\leq C R_0^N \left(\left(\frac{\|u\|_{L^\infty(B)}}{R_0 - r_0} \right)^{p_k+2+\alpha} + \varepsilon_0 \right) \\ &\quad + C \left(\frac{\|u\|_{L^\infty(B)}}{R_0 - r_0} \right)^{\frac{2}{p_k}(p_k+2+\alpha)} \int_{B_{R_0}} \sum_{i \neq k} |u_{x_i}|^{\frac{p_i-2}{p_k}(p_k+2+\alpha)} dx. \end{aligned}$$

Proof. We fix $k \in \{1, \dots, N\}$ and take $\eta \in C_0^\infty(B)$ a positive cut-off function. For every $\alpha > -1$, we estimate the quantity

$$\int |u_{x_k}|^{p_k+2+\alpha} \eta^2 dx = \int u_{x_k} u_{x_k} |u_{x_k}|^{p_k+\alpha} \eta^2 dx.$$

By integration by parts (recall that $u \in C^2(B)$), one gets

$$\begin{aligned} \int |u_{x_k}|^{p_k+2+\alpha} \eta^2 dx &= - \int u \left(u_{x_k} |u_{x_k}|^{p_k+\alpha} \eta^2 \right)_{x_k} dx \\ &= -(p_k + \alpha + 1) \int u u_{x_k x_k} |u_{x_k}|^{p_k+\alpha} \eta^2 dx - 2 \int u u_{x_k} |u_{x_k}|^{p_k+\alpha} \eta \eta_{x_k} dx. \end{aligned}$$

Hence, we have

$$(4.7) \quad \int |u_{x_k}|^{p_k+2+\alpha} \eta^2 dx \leq (p_k + \alpha + 1) \|u\|_{L^\infty(B)} \left(\int |u_{x_k x_k}| |u_{x_k}|^{p_k+\alpha} \eta^2 dx + \int |u_{x_k}|^{p_k+\alpha+1} \eta |\nabla \eta| dx \right).$$

We now use the Young's inequality for the two terms in the right-hand side: for every $\tau > 0$,

$$|u_{x_k x_k}| |u_{x_k}|^{p_k+\alpha} \leq \tau |u_{x_k}|^{p_k+\alpha+2} + \frac{1}{4\tau} |u_{x_k}|^{p_k+\alpha-2} |u_{x_k x_k}|^2$$

and

$$|u_{x_k}|^{p_k+\alpha+1} \eta |\nabla \eta| \leq \tau |u_{x_k}|^{p_k+\alpha+2} \eta^2 + \frac{1}{4\tau} |u_{x_k}|^{p_k+\alpha} |\nabla \eta|^2.$$

In the first inequality, when $p_k + \alpha - 2 < 0$, the quantity $|u_{x_k}|^{p_k+\alpha-2} |u_{x_k x_k}|^2$ is defined to be 0 on the set where $u_{x_k} = 0$.

Inserting these two inequalities into (4.7) and choosing

$$\tau = \frac{1}{4(p_k + \alpha + 1) \|u\|_{L^\infty(B)}},$$

we can absorb the two terms multiplied by τ in the left-hand side. This leads to

$$\int \eta^2 |u_{x_k}|^{p_k+2+\alpha} dx \leq C \|u\|_{L^\infty(B)}^2 \left(\int |u_{x_k x_k}|^2 |u_{x_k}|^{p_k+\alpha-2} \eta^2 dx + \int |u_{x_k}|^{p_k+\alpha} |\nabla \eta|^2 dx \right),$$

for a constant $C = C(p_k, \alpha) > 0$. Observe that

$$|u_{x_k x_k}|^2 |u_{x_k}|^{p_k+\alpha-2} = |u_{x_k x_k}|^2 |u_{x_k}|^{p_k-2} |u_{x_k}|^\alpha \leq g''_{k,\varepsilon}(u_{x_k}) |u_{x_k x_k}|^2 |u_{x_k}|^\alpha.$$

Thus, if $\alpha > 0$, we can apply the Caccioppoli inequality of Lemma 3.1 with the convex function $\Phi(t) = |t|^{\frac{\alpha}{2}+1}$. Otherwise, if $-1 < \alpha \leq 0$, we can apply Lemma 3.2. This gives

$$\int |u_{x_k x_k}|^2 |u_{x_k}|^{p_k+\alpha-2} \eta^2 dx \leq C \sum_{i=1}^N \int_B g''_{i,\varepsilon}(u_{x_i}) |u_{x_k}|^{\alpha+2} |\eta_{x_i}|^2 dx,$$

and thus we obtain

$$\begin{aligned} \int \eta^2 |u_{x_k}|^{p_k+2+\alpha} dx &\leq C \|u\|_{L^\infty(B)}^2 \int \left(|u_{x_k}|^{\alpha+2} \sum_{i=1}^N g''_{i,\varepsilon}(u_{x_i}) + |u_{x_k}|^{p_k+\alpha} \right) |\nabla \eta|^2 dx \\ &\leq C \|u\|_{L^\infty(B)}^2 \int \left(|u_{x_k}|^{\alpha+2} \sum_{i \neq k} |u_{x_i}|^{p_i-2} + |u_{x_k}|^{p_k+\alpha} + \varepsilon |u_{x_k}|^{\alpha+2} \right) |\nabla \eta|^2 dx, \end{aligned}$$

where in the second inequality the constant C may differ from the previous one. There we used (4.1).

We now fix a pair concentric balls $B_r \subset B_R \Subset B$. Applying the above estimate to a non negative cut-off function $\eta \in C_0^\infty(B_R)$ such that

$$\eta \equiv 1 \text{ on } B_r \quad \text{and} \quad \|\nabla \eta\|_{L^\infty(B_R)} \leq \frac{C}{R-r},$$

one gets

$$(4.8) \quad \int_{B_r} |u_{x_k}|^{p_k+2+\alpha} dx \leq \frac{C \|u\|_{L^\infty(B)}^2}{(R-r)^2} \int_{B_R} \left(|u_{x_k}|^{\alpha+2} \sum_{i \neq k} |u_{x_i}|^{p_i-2} + |u_{x_k}|^{p_k+\alpha} + \varepsilon |u_{x_k}|^{\alpha+2} \right) dx.$$

We now want to absorb all the terms containing u_{x_k} from the right-hand side. Thus, we apply again the Young's inequality. For every $\tau > 0$, there exists $C_0 > 0$ which depends only on N, p_k and α such that

$$|u_{x_k}|^{\alpha+2} \sum_{i \neq k} |u_{x_i}|^{p_i-2} \leq \tau |u_{x_k}|^{p_k+\alpha+2} + \frac{C_0}{\tau^{\frac{\alpha+2}{p_k}}} \sum_{i \neq k} |u_{x_i}|^{(p_i-2) \frac{p_k+\alpha+2}{p_k}},$$

and

$$|u_{x_k}|^{p_k+\alpha} \leq \tau |u_{x_k}|^{p_k+\alpha+2} + \frac{C_0}{\tau^{\frac{p_k+\alpha}{2}}}.$$

Moreover, we use that

$$\varepsilon |u_{x_k}|^{\alpha+2} \leq \varepsilon + |u_{x_k}|^{p_k+\alpha+2}.$$

thanks to the fact that $\varepsilon < 1$ and $p_k \geq 2$. Inserting these inequalities into (4.8) and choosing

$$\tau = \frac{(R-r)^2}{4C \|u\|_{L^\infty(B_R)}^2},$$

one obtains

$$\begin{aligned} \int_{B_r} |u_{x_k}|^{p_k+2+\alpha} dx &\leq \frac{1}{2} \int_{B_R} |u_{x_k}|^{p_k+2+\alpha} dx \\ &\quad + \frac{C \|u\|_{L^\infty(B_R)}^2}{(R-r)^2} \left(\frac{R^N}{\tau^{\frac{p_k+\alpha}{2}}} + \frac{1}{\tau^{\frac{\alpha+2}{p_k}}} \sum_{i \neq k} \int_{B_R} |u_{x_i}|^{(p_i-2) \frac{p_k+2+\alpha}{p_k}} dx + \varepsilon R^N \right). \end{aligned}$$

By recalling the choice of τ above, this is the same as

$$\begin{aligned} \int_{B_r} |u_{x_k}|^{p_k+2+\alpha} dx &\leq \frac{1}{2} \int_{B_R} |u_{x_k}|^{p_k+2+\alpha} dx + C R^N \left(\left(\frac{\|u\|_{L^\infty(B)}}{R-r} \right)^{p_k+\alpha+2} + \varepsilon_0 \right) \\ &\quad + C \left(\frac{\|u\|_{L^\infty(B)}}{R-r} \right)^{2 \frac{\alpha+2+p_k}{p_k}} \left(\int_{B_R} \sum_{i \neq k} |u_{x_i}|^{(p_i-2) \frac{p_k+2+\alpha}{p_k}} dx \right). \end{aligned}$$

We now fix $r_0 < R_0$ as in the statement and use the previous estimate for $r_0 \leq r < R \leq R_0$. By applying Lemma 2.4, one finally obtains that

$$\begin{aligned} \int_{B_{r_0}} |u_{x_k}|^{p_k+2+\alpha} dx &\leq C R_0^N \left(\left(\frac{\|u\|_{L^\infty(B)}}{R_0-r_0} \right)^{p_k+\alpha+2} + \varepsilon_0 \right) \\ &\quad + C \left(\frac{\|u\|_{L^\infty(B)}}{R_0-r_0} \right)^{2 \frac{p_k+\alpha+2}{p_k}} \left(\int_{B_{R_0}} \sum_{i \neq k} |u_{x_i}|^{(p_i-2) \frac{p_k+2+\alpha}{p_k}} dx \right). \end{aligned}$$

Here, the constant C depends on N, p_k and α . This concludes the proof. \square

5. A LIPSCHITZ ESTIMATE

Proposition 5.1. *Let $2 \leq p_1 \leq \dots \leq p_N$ and $0 < \varepsilon \leq \varepsilon_0$. There exist an exponent $\gamma \geq p_N + 2$, two exponents $\Theta, \beta > 1$ and a constant $C > 0$ such that for every $B_{r_0} \subset B_{R_0} \Subset B$ with $0 < r_0 < R_0 \leq 1$,*

$$(5.1) \quad \|\nabla u\|_{L^\infty(B_{r_0})} \leq \frac{C}{(R_0 - r_0)^\beta} \left(\int_{B_{R_0}} |\nabla u|^\gamma dx + 1 \right)^{\frac{\Theta}{\gamma}}.$$

The parameters γ, β, Θ and the constant C are independent of ε .

Proof. The proof is very similar to that of [4, Theorem 5.1], though some important technical modifications have to be taken into account. For simplicity, we assume throughout the proof that $N \geq 3$, so in this case the Sobolev exponent 2^* is finite. Observe that the case $N = 2$, which could be treated with minor modifications, is already contained in [6, Theorem 1.4] (the proof there is different).

As in [4], we divide the proof into four steps.

Step 1: a first iterative scheme. We can proceed as in [4, Proposition 5.1, Step 1] by replacing the term

$$\int |\nabla \eta|^2 |u_{x_k}|^{2q+p} dx,$$

there, with the following one

$$\int |\nabla \eta|^2 |u_{x_k}|^{2q+p_k} dx.$$

Then the relevant outcome is now

$$(5.2) \quad \left(\int \left(\sum_{k=1}^N |u_{x_k}|^{2q+p_k} \right)^{\frac{2^*}{2}} \eta^{2^*} dx \right)^{\frac{2}{2^*}} \leq C q^5 \sum_{i,k=1}^N \int g''_{i,\varepsilon}(u_{x_i}) |u_{x_k}|^{2q+2} |\nabla \eta|^2 dx \\ + C \int |\nabla \eta|^2 \sum_{k=1}^N |u_{x_k}|^{2q+p_k} dx.$$

We now introduce the function

$$\mathcal{U}(x) := \max_{k=1, \dots, N} |u_{x_k}(x)|,$$

and observe that

$$|u_{x_k}|^{2q+p_1} - 1 \leq |u_{x_k}|^{2q+p_k} \leq |u_{x_k}|^{2q+p_N} + 1, \quad \text{for } k = 1, \dots, N.$$

This in turn gives

$$\mathcal{U}^{2q+p_1} - 1 \leq \sum_{k=1}^N |u_{x_k}|^{2q+p_k} \leq N \mathcal{U}^{2q+p_N} + N.$$

Also, we have that

$$g''_{i,\varepsilon}(u_{x_i}) |u_{x_k}|^{2q+2} \leq C \mathcal{U}^{2q+p_N} + C, \quad \text{for every } 1 \leq i, k \leq N.$$

By further observing that

$$(\mathcal{U}^{2q+p_1})^{\frac{2^*}{2}} \leq \left(1 + \sum_{k=1}^N |u_{x_k}|^{2q+p_k}\right)^{\frac{2^*}{2}} \leq C \left(1 + \left(\sum_{k=1}^N |u_{x_k}|^{2q+p_k}\right)^{\frac{2^*}{2}}\right),$$

we obtain from (5.2)

$$\left(\int \mathcal{U}^{\frac{2^*}{2}(2q+p_1)} \eta^{2^*}\right)^{\frac{2}{2^*}} \leq C q^5 \int \mathcal{U}^{2q+p_N} |\nabla \eta|^2 dx + C q^5 \int |\nabla \eta|^2 dx + C q^5 \left(\int \eta^{2^*} dx\right)^{\frac{2}{2^*}},$$

for a possibly different $C = C(N, \mathbf{p}) > 1$. By using the Sobolev embedding $W_0^{1,2}(B) \hookrightarrow L^{2^*}(B)$

$$\left(\int \eta^{2^*} dx\right)^{\frac{2}{2^*}} \leq C \int |\nabla \eta|^2 dx,$$

thus the previous estimate leads to

$$(5.3) \quad \left(\int \mathcal{U}^{\frac{2^*}{2}(2q+p_1)} \eta^{2^*} dx\right)^{\frac{2}{2^*}} \leq C q^5 \int |\nabla \eta|^2 (\mathcal{U}^{2q+p_N} + 1) dx.$$

We fix two concentric balls $B_r \subset B_R \Subset B$, with $0 < r < R \leq 1$. Then for every pair of radius $r \leq t < s \leq R$ we take in (5.3) a standard cut-off function

$$(5.4) \quad \eta \in C_0^\infty(B_s), \quad \eta \equiv 1 \text{ on } B_t, \quad 0 \leq \eta \leq 1, \quad \|\nabla \eta\|_{L^\infty} \leq \frac{C}{s-t}.$$

This yields

$$(5.5) \quad \left(\int_{B_t} \mathcal{U}^{\frac{2^*}{2}(2q+p_1)} dx\right)^{\frac{2}{2^*}} \leq C \frac{q^5}{(s-t)^2} \int_{B_s} (\mathcal{U}^{2q+p_N} + 1) dx.$$

We define the sequence of exponents

$$\gamma_j = p_N + 2^{j+2} - 2, \quad j \geq 0,$$

and take in (5.5) $q = 2^{j+1} - 1$. This gives for every $j \geq 0$,

$$(5.6) \quad \left(\int_{B_t} \mathcal{U}^{\frac{2^*}{2}(\gamma_j+p_1-p_N)} dx\right)^{\frac{2}{2^*}} \leq C \frac{2^{5j}}{(s-t)^2} \int_{B_s} (\mathcal{U}^{\gamma_j} + 1) dx,$$

for a possibly different constant $C = C(N, \mathbf{p}) > 1$. Observe that we always have

$$\gamma_j + p_1 - p_N \geq 2^{j+2}, \quad j \in \mathbb{N},$$

thanks to the definition of γ_j .

Step 2: filling the gaps. By using the definition of γ_j , it is not difficult to see that

$$\gamma_j < \frac{2^*}{2} (\gamma_j + p_1 - p_N) \quad \Longleftrightarrow \quad j > \log_2 \left(\frac{N-2}{2} (p_N - 2) - \frac{N}{2} (p_1 - 2) \right) - 2.$$

Thus we introduce the starting index²

$$j_0 = \min \left\{ j \in \mathbb{N} : j > \log_2 \left(\frac{N-2}{2} (p_N - 2) - \frac{N}{2} (p_1 - 2) \right) - 2 \right\}.$$

By definition, this entails that

$$\gamma_{j-1} < \gamma_j < \frac{2^*}{2} (\gamma_j + p_1 - p_N), \quad \text{for every } j \geq j_0 + 1.$$

By interpolation in Lebesgue spaces, we obtain

$$\int_{B_t} \mathcal{U}^{\gamma_j} dx \leq \left(\int_{B_t} \mathcal{U}^{\gamma_{j-1}} dx \right)^{\frac{\tau_j \gamma_j}{\gamma_j - 1}} \left(\int_{B_t} \mathcal{U}^{\frac{2^*}{2} (\gamma_j + p_1 - p_N)} dx \right)^{\frac{2}{2^*} \frac{(1-\tau_j) \gamma_j}{\gamma_j + p_1 - p_N}},$$

where the interpolation exponent $0 < \tau_j < 1$ is given by

$$\tau_j = \frac{\gamma_{j-1}}{\gamma_j} \frac{\frac{2^*}{2} (\gamma_j + p_1 - p_N) - \gamma_j}{\frac{2^*}{2} (\gamma_j + p_1 - p_N) - \gamma_{j-1}}.$$

We now rely on (5.6) to get

$$\begin{aligned} \int_{B_t} \mathcal{U}^{\gamma_j} dx &\leq \left(\int_{B_t} \mathcal{U}^{\gamma_{j-1}} dx \right)^{\frac{\tau_j \gamma_j}{\gamma_j - 1}} \left(C \frac{2^{5j}}{(s-t)^2} \int_{B_s} (\mathcal{U}^{\gamma_j} + 1) dx \right)^{\frac{(1-\tau_j) \gamma_j}{\gamma_j + p_1 - p_N}} \\ &= \left[\left(C \frac{2^{5j}}{(s-t)^2} \right)^{\frac{1-\tau_j}{\tau_j} \frac{\gamma_j}{\gamma_j + p_1 - p_N}} \left(\int_{B_t} \mathcal{U}^{\gamma_{j-1}} dx \right)^{\frac{\gamma_j}{\gamma_j - 1}} \right]^{\tau_j} \left(\int_{B_s} (\mathcal{U}^{\gamma_j} + 1) dx \right)^{\frac{(1-\tau_j) \gamma_j}{\gamma_j + p_1 - p_N}}. \end{aligned}$$

By Young's inequality, for every $j \geq j_0 + 1$, we get

$$\begin{aligned} (5.7) \quad \int_{B_t} \mathcal{U}^{\gamma_j} dx &\leq \frac{(1-\tau_j) \gamma_j}{\gamma_j + p_1 - p_N} \int_{B_s} (\mathcal{U}^{\gamma_j} + 1) dx \\ &\quad + \frac{1}{\left(\frac{\gamma_j + p_1 - p_N}{(1-\tau_j) \gamma_j} \right)'} \left(C \frac{2^{5j}}{(s-t)^2} \right)^{\frac{(1-\tau_j) \gamma_j}{\gamma_j + p_1 - p_N} \left(\frac{\gamma_j + p_1 - p_N}{(1-\tau_j) \gamma_j} \right)'} \left(\int_{B_t} \mathcal{U}^{\gamma_{j-1}} dx \right)^{\frac{\gamma_j}{\gamma_j - 1} \tau_j \left(\frac{\gamma_j + p_1 - p_N}{(1-\tau_j) \gamma_j} \right)'}. \end{aligned}$$

We also introduce the second index

$$j_1 = \min \{ j \in \mathbb{N} : j > \log_2 ((N-2)(p_N - 2) - N(p_1 - 2)) - 2 \}.$$

²We use the convention that

$$\log t = -\infty \quad \text{for } t \leq 0.$$

Observe that $j_0 = 0$ whenever

$$\frac{N-2}{2} (p_N - 2) - \frac{N}{2} (p_1 - 2) < 4 \quad \text{i.e.} \quad p_N < 2 + \frac{N(p_1 - 2) + 8}{N-2}.$$

If we finally set

$$J = 1 + \max\{j_0, j_1\},$$

then by Lemma A.1 we know that

$$(5.8) \quad 0 < C_1 \leq \frac{(1 - \tau_j) \gamma_j}{\gamma_j + p_1 - p_N} \leq C_2 < 1, \quad \text{for every } j \geq J.$$

This in turn implies that for every $j \geq J$

$$\frac{1}{\left(\frac{\gamma_j + p_1 - p_N}{(1 - \tau_j) \gamma_j}\right)'} \leq \frac{1}{\left(\frac{1}{C_1}\right)'} = 1 - C_1.$$

Thus from (5.7) we get

$$(5.9) \quad \begin{aligned} \int_{B_t} \mathcal{U}^{\gamma_j} dx &\leq C_2 \int_{B_s} (\mathcal{U}^{\gamma_j} + 1) dx \\ &+ (1 - C_1) \left(C \frac{2^{5j}}{(s-t)^2}\right)^\beta \left(\int_{B_t} \mathcal{U}^{\gamma_{j-1}} dx\right)^{\frac{\gamma_j}{\gamma_{j-1} - \tau_j} \left(\frac{\gamma_j + p_1 - p_N}{(1 - \tau_j) \gamma_j}\right)'} \end{aligned}$$

for some $1 < \beta < \infty$, depending on N, p_1 and p_N . In the last inequality we also used that $s \leq R \leq 1$ and $C > 1$, together with (5.8). Finally we set

$$\varepsilon_j = \tau_j \left(\frac{\gamma_j + p_1 - p_N}{(1 - \tau_j) \gamma_j}\right)' - 1, \quad \text{for } j \geq J,$$

and rewrite (5.9) as

$$(5.10) \quad \begin{aligned} \int_{B_t} \mathcal{U}^{\gamma_j} dx &\leq C_2 \int_{B_s} \mathcal{U}^{\gamma_j} dx \\ &+ (1 - C_1) \left(C \frac{2^{5j}}{(s-t)^2}\right)^\beta \left(\int_{B_t} \mathcal{U}^{\gamma_{j-1}} dx\right)^{\frac{\gamma_j}{\gamma_{j-1} - \tau_j} (1 + \varepsilon_j)} + C_2 |B_R|, \end{aligned}$$

which holds for every $r \leq s < t \leq R$. By applying Lemma 2.4 with

$$Z(t) = \int_{B_t} \mathcal{U}^{\gamma_j} dx, \quad \alpha_0 = 2\beta, \quad \text{and} \quad \vartheta = C_2,$$

we finally obtain for every $j \geq J$,

$$(5.11) \quad \int_{B_r} \mathcal{U}^{\gamma_j} dx \leq C \left(2^{5j\beta} (R-r)^{-2\beta} \left(\int_{B_R} \mathcal{U}^{\gamma_{j-1}} dx\right)^{\frac{\gamma_j}{\gamma_{j-1} - \tau_j} (1 + \varepsilon_j)} + 1\right),$$

for some $C = C(N, p_1, p_N) > 1$.

Step 3: Moser's iteration. We now iterate the previous estimate on a sequence of shrinking balls. We fix two radii $0 < r < R \leq 1$ and define the sequence

$$R_j = r + \frac{R-r}{2^{j-J}}, \quad j \geq J.$$

We use (5.11) with $R_{j+1} < R_j$ in place of $r < R$. Thus we get

$$(5.12) \quad \int_{B_{R_{j+1}}} \mathcal{U}^{\gamma_j} dx \leq C \left(2^{7j\beta} (R-r)^{-2\beta} \left(\int_{B_{R_j}} \mathcal{U}^{\gamma_{j-1}} dx\right)^{\frac{\gamma_j}{\gamma_{j-1} - \tau_j} (1 + \varepsilon_j)} + 1\right)$$

where the constant $C > 1$ depends on N and p_1, p_N only.

We introduce the notation

$$Y_j = \int_{B_{R_j}} \mathcal{U}^{\gamma_j-1} dx,$$

thus (5.12) reads

$$Y_{j+1} \leq C \left(2^{7j\beta} (R-r)^{-2\beta} Y_j^{\frac{\gamma_j}{\gamma_j-1} (1+\varepsilon_j)} + 1 \right) \leq (C 2^{7\beta} (R-r)^{-2\beta})^j (Y_j + 1)^{\frac{\gamma_j}{\gamma_j-1} (1+\varepsilon_j)}.$$

Here, we have used again that $C > 1$ and $R \leq 1$, so that the term multiplying Y_j is larger than 1. By iterating the previous estimate starting from $j = J$ and using some standard manipulations, we obtain

$$\begin{aligned} Y_{n+1} &\leq (C 2^{7\beta} (R-r)^{-2\beta})^n (Y_n + 1)^{\frac{\gamma_n}{\gamma_n-1} (1+\varepsilon_n)} \\ &\leq (C 2^{7\beta} (R-r)^{-2\beta})^n \left((C 2^{7\beta} (R-r)^{-2\beta})^{n-1} (Y_{n-1} + 1)^{\frac{\gamma_{n-1}}{\gamma_{n-1}-2} (1+\varepsilon_{n-1})} + 1 \right)^{\frac{\gamma_n}{\gamma_n-1} (1+\varepsilon_n)} \\ &\leq (C 2^{7\beta} (R-r)^{-2\beta})^n \left(2 (C 2^{7\beta} (R-r)^{-2\beta})^{n-1} (Y_{n-1} + 1)^{\frac{\gamma_{n-1}}{\gamma_{n-1}-2} (1+\varepsilon_{n-1})} \right)^{\frac{\gamma_n}{\gamma_n-1} (1+\varepsilon_n)} \\ &\leq \dots \\ &\leq \left(2 C 2^{7\beta} (R-r)^{-2\beta} \right)^{\sum_{j=J}^n \left(j \frac{\gamma_n}{\gamma_j} \prod_{k=j+1}^n (1+\varepsilon_k) \right)} \left[Y_J + 1 \right]^{\frac{\gamma_n}{\gamma_J-1} \prod_{j=J}^n (1+\varepsilon_j)}, \end{aligned}$$

where we used that $C 2^{7\beta} (R-r)^{-2\beta} > 1$. We now simply write C in place of $2 C 2^{7\beta}$ and take the power $1/\gamma_n$ on both sides:

$$\begin{aligned} Y_{n+1}^{\frac{1}{\gamma_n}} &\leq \left(C (R-r)^{-2\beta} \right)^{\sum_{j=J}^n \frac{j}{\gamma_j} \prod_{k=j+1}^n (1+\varepsilon_k)} \left[Y_J + 1 \right]^{\frac{\prod_{j=J}^n (1+\varepsilon_j)}{\gamma_J-1}} \\ &\leq \left(C (R-r)^{-2\beta} \right)^{\Theta \sum_{j=J}^n \frac{j}{\gamma_j}} \left[Y_J + 1 \right]^{\frac{\Theta}{\gamma_J-1}}. \end{aligned}$$

In the previous estimate, we set

$$\Theta = \lim_{n \rightarrow \infty} \prod_{j=0}^n (1 + \varepsilon_j),$$

which is a finite number, thanks to Lemma A.2. We observe that $\gamma_j \sim 2^{j+2}$ as j goes to ∞ . This implies the convergence of the series above and we thus get

$$\|\mathcal{U}\|_{L^\infty(B_r)} = \lim_{n \rightarrow \infty} \left(\int_{B_{R_{n+1}}} \mathcal{U}^{\gamma_{n+1}} dx \right)^{\frac{1}{\gamma_{n+1}}} \leq C (R-r)^{-\beta'} \left(\int_{B_R} \mathcal{U}^{\gamma_{J-1}} dx + 1 \right)^{\frac{\Theta}{\gamma_J-1}},$$

for some $C = C(N, p_1, p_N) > 1$ and $\beta' = \beta'(N, p_1, p_N) > 0$. By recalling the definition of \mathcal{U} , we finally obtain

$$\|\nabla u\|_{L^\infty(B_r)} \leq C (R-r)^{-\beta'} \left(\int_{B_R} |\nabla u|^{\gamma_{J-1}} dx + 1 \right)^{\frac{\Theta}{\gamma_J-1}}.$$

This concludes the proof. \square

6. A RECURSIVE GAIN OF INTEGRABILITY FOR THE GRADIENT

The main outcome of estimate (5.1) is the following: assume that our local minimizer U has a gradient with a sufficiently high integrability, then one would be able to conclude that ∇U has to be bounded. We notice that since the explicit determination of the exponent γ in (5.1) is actually very intricate (unless some upper bounds on p_N/p_1 are imposed), we essentially need to prove that

$$\nabla U \in L_{\text{loc}}^q \quad \text{for every } q < \infty,$$

in order to be on the safe side. Thus, in order to infer the desired local Lipschitz regularity on U , we are going to prove a higher integrability estimate on ∇u_ε , which is uniform with respect to $0 < \varepsilon \leq \varepsilon_0$. This is the content of the result of this section. Remember that we simplify the notation u_ε and replace it by u .

Proposition 6.1. *Let $2 \leq p_1 \leq \dots \leq p_N < +\infty$ and $0 < \varepsilon \leq \varepsilon_0$. For every $2 \leq q_0 < +\infty$ and every $B_{R_0} \Subset B$, there exists a constant $C > 0$ such that*

$$\sum_{i=1}^N \int_{B_{R_0}} |u_{x_i}|^{p_i q_0} dx \leq C.$$

The constant C depends on $N, \mathbf{p}, q_0, R_0, \text{dist}(B_{R_0}, \partial B)$,

$$\|u\|_{L^\infty(B)} \quad \text{and} \quad \sum_{i=1}^N \int_B |u_{x_i}|^{p_i} dx.$$

Proof. We proceed to exploit the scheme of Proposition 4.3. In what follows, we use the convention that

$$\frac{p}{p-2} = +\infty,$$

whenever $p = 2$. We can also assume without loss of generality that

$$p_N > 2,$$

otherwise $p_1 = \dots = p_N = 2$ and in this case the regularity theory for our problem is well-established (U would be a harmonic function in such a situation).

We fix q_0 as in the statement and introduce the exponents

$$(6.1) \quad q_j = \min \left\{ \frac{p_j}{p_j - 2}, q_0 \right\} = \min \left\{ \left(\frac{p_j}{2} \right)', q_0 \right\}, \quad j = 1, \dots, N.$$

Since $p_1 \leq \dots \leq p_N$, we get that $q_1 \geq q_2 \geq \dots \geq q_N$ and thus

$$(6.2) \quad \min_{i=1, \dots, k} q_i = q_k, \quad \text{for every } k \in \{1, \dots, N\}.$$

We now prove by *downward induction* on $j = N, \dots, 1$ that the following fact holds: for every $B_R \Subset B$, one has

$$(6.3) \quad \begin{array}{l} \text{for every } j \in \{1, \dots, N\} \\ \text{and every } B_R \Subset B, \end{array} \quad \sum_{i=j}^N \int_{B_R} |u_{x_i}|^{p_i q_{j-1}} dx \leq C, \quad \text{with } C > 0 \text{ independent of } \varepsilon.$$

In particular, for $j = 1$, (6.3) implies

$$\sum_{i=1}^N \int_{B_R} |u_{x_i}|^{p_i q_0} dx \leq C,$$

for a uniform constant $C > 0$. The statement on the quality of the constant C will be clear from the computations below.

Initialization step. We start from $j = N$. We observe that in this case the right-hand side of (4.6) (written for $k = N$) is uniformly bounded with respect to $\varepsilon > 0$, provided that $\alpha > -1$ is chosen in such a way that for every $i \in \{1, \dots, N-1\}$, one has

$$(p_i - 2) \frac{p_N + 2 + \alpha}{p_N} \leq p_i.$$

If $p_i = 2$, this is automatically satisfied. Otherwise, this is equivalent to

$$p_N + 2 + \alpha \leq p_N \frac{p_i}{p_i - 2}.$$

We define α by

$$p_N + 2 + \alpha = p_N q_{N-1}.$$

By definition of q_{N-1} ,

$$p_N + 2 + \alpha = p_N \min \left\{ \frac{p_{N-1}}{p_{N-1} - 2}, q_0 \right\} \leq p_N \min_{1 \leq i \leq N-1} \frac{p_i}{p_i - 2},$$

as desired. We need to check that $\alpha > -1$, or equivalently

$$(6.4) \quad p_N \min \left\{ \frac{p_{N-1}}{p_{N-1} - 2}, q_0 \right\} > p_N + 1.$$

Since $q_0 \geq 2$, one has $p_N q_0 > p_N + 1$. Moreover, using that $p_{N-1} \leq p_N$, one gets $2p_N > p_{N-1} - 2$ which in turn is equivalent to

$$p_N \frac{p_{N-1}}{p_{N-1} - 2} > p_N + 1.$$

This proves (6.4).

Thus for every $B_R \Subset B_{R'} \Subset B$, it follows from (4.6) that

$$\begin{aligned} \int_{B_R} |u_{x_N}|^{p_N q_{N-1}} dx &\leq C (R')^N \left(\left(\frac{\|u\|_{L^\infty(B)}}{R' - R} \right)^{p_N q_{N-1}} + \varepsilon_0 \right) \\ &\quad + C \left(\frac{\|u\|_{L^\infty(B)}}{R' - R} \right)^{2q_{N-1}} \int_{B_{R'}} \sum_{i=1}^{N-1} |u_{x_i}|^{\frac{p_i-2}{p_N} (p_N q_{N-1})} dx. \end{aligned}$$

Since $q_i \geq q_{N-1}$ for $1 \leq i \leq N-1$, one has

$$\frac{p_i}{q_{N-1} (p_i - 2)} \geq 1.$$

Using Hölder's inequality for each term of the sum of the right-hand side, with the exponent $p_i/(q_{N-1}(p_i-2))$ and its conjugate, one gets

$$\begin{aligned} \int_{B_R} |u_{x_N}|^{p_N q_{N-1}} dx &\leq C (R')^N \left(\left(\frac{\|u\|_{L^\infty(B)}}{R' - R} \right)^{p_N q_{N-1}} + \varepsilon_0 \right) \\ &\quad + C \left(\frac{\|u\|_{L^\infty(B)}}{R' - R} \right)^{2q_{N-1}} \sum_{i=1}^{N-1} \left(\int_{B_{R'}} |u_{x_i}|^{p_i} dx \right)^{\frac{(p_i-2) q_{N-1}}{p_i}}. \end{aligned}$$

By using Lemma 2.1 and Proposition 2.3 in order to control the two terms on the right-hand side, we get a uniform (in ε) control on the $L^{p_N q_{N-1}}(B_R)$ norm of u_{x_N} . This finally establishes the initialization step $j = N$, i.e.

$$\text{for every } B_R \Subset B, \text{ we have } \int_{B_R} |u_{x_N}|^{p_N q_{N-1}} dx \leq C, \quad \text{with } C > 0 \text{ independent of } \varepsilon.$$

Inductive step. We then assume that the assertion (6.3) is true for some $j \in \{2, \dots, N\}$ and we prove it for $j - 1$. By the induction assumption, we thus know that

$$(6.5) \quad \begin{array}{l} \text{for some } j \in \{2, \dots, N\} \\ \text{and every } B_R \in B, \end{array} \quad \sum_{i=j}^N \int_{B_R} |u_{x_i}|^{p_i q_{j-1}} dx \leq C, \quad \text{with } C > 0 \text{ independent of } \varepsilon.$$

In the rest of the proof, we establish that (6.5) implies

$$(6.6) \quad \text{for every } B_R \in B, \quad \sum_{i=j-1}^N \int_{B_R} |u_{x_i}|^{p_i q_{j-2}} dx \leq C, \quad \text{with } C > 0 \text{ independent of } \varepsilon.$$

In order to prove this, as explained in the Introduction, we need to employ a *multiply iterative scheme* based on Proposition 4.3. More specifically, we start by relying on (4.6) with the choices

$$k = j - 1 \quad \text{and} \quad p_{j-1} + 2 + \alpha = p_{j-1} \min \left\{ \min_{0 \leq i \leq j-2} q_i, q_{j-1} \min_{j \leq i \leq N} q_i \right\}.$$

We first justify the fact that such a choice for α is feasible. Observe that

$$\min_{0 \leq i \leq j-2} q_i = q_{j-2} \quad \text{and} \quad \min_{j \leq i \leq N} q_i = q_N,$$

hence the condition on α is equivalent to

$$(6.7) \quad p_{j-1} + 2 + \alpha = p_{j-1} \min \{q_{j-2}, q_{j-1} q_N\}.$$

Since

$$q_{j-1} = \min \left\{ \frac{p_{j-1}}{p_{j-1} - 2}, q_0 \right\} \quad \text{with } q_0 \geq 2,$$

one has $p_{j-1} + 2 \leq p_{j-1} q_{j-1}$. By recalling that the exponents q_j are non increasing and larger than 1, this implies that

$$p_{j-1} + 2 \leq p_{j-1} q_{j-2} \quad \text{and} \quad p_{j-1} + 2 < p_{j-1} q_{j-1} q_N,$$

and thus

$$p_{j-1} + 2 + \alpha = p_{j-1} \min \{q_{j-2}, q_{j-1} q_N\} \geq p_{j-1} + 2.$$

This implies that $\alpha \geq 0$ as desired.

We next rely on the fact that by Lemma 2.1 and Proposition 2.3, we have

$$\|u\|_{L^\infty(B)} + \sum_{i=1}^N \int_B |u_{x_i}|^{p_i} dx \leq C,$$

with a constant $C > 0$ independent of $\varepsilon > 0$ and on the induction assumption (6.5), which gives a local uniform (in ε) control on

$$\sum_{i=j}^N \int_{B_R} |u_{x_i}|^{p_i q_{j-1}} dx,$$

for $B_R \in B$. Hence, the definition (6.7) of α ensures that the right-hand side of (4.6) is uniformly bounded. Thus from Proposition 4.3, we get that for every $B_R \in B$ we have

$$(6.8) \quad \int_{B_R} |u_{x_{j-1}}|^{\beta_{j-1}^{(0)}} dx \leq C,$$

with $C > 0$ independent of ε . Here, the exponent $\beta_{j-1}^{(0)}$ is given by

$$\beta_{j-1}^{(0)} = p_{j-1} + 2 + \alpha = p_{j-1} \min \{q_{j-2}, q_{j-1} q_N\}.$$

We can summarize the previous integrability information as the following estimate: for every $B_R \Subset B$,

$$(6.9) \quad \sum_{i=j-1}^N \int_{B_R} |u_{x_i}|^{\beta_i^{(0)}} dx \leq C,$$

with $C > 0$ independent of $\varepsilon > 0$ and

$$(6.10) \quad \begin{cases} \beta_{j-1}^{(0)} &= p_{j-1} \min \{q_{j-2}, q_{j-1} q_N\}, \\ \beta_i^{(0)} &= p_i q_{j-1}, \quad \text{for } i = j, \dots, N. \end{cases}$$

We proceed to define by induction a vector sequence

$$(\beta_{j-1}^{(\ell)}, \dots, \beta_N^{(\ell)}), \quad \ell \in \mathbb{N},$$

as follows: for $\ell = 0$ this is given by (6.10) and then we use the following *multiply recursive* scheme

$$(6.11) \quad \begin{cases} \beta_N^{(\ell+1)} &= p_N \min \left\{ q_{j-2}, \min_{j-1 \leq k \leq N-1} \frac{\beta_k^{(\ell)}}{p_k - 2} \right\} \\ \beta_{N-1}^{(\ell+1)} &= p_{N-1} \min \left\{ q_{j-2}, \min_{j-1 \leq k \leq N-2} \frac{\beta_k^{(\ell)}}{p_k - 2}, \frac{\beta_N^{(\ell+1)}}{p_N - 2} \right\} \\ \beta_{N-2}^{(\ell+1)} &= p_{N-2} \min \left\{ q_{j-2}, \min_{j-1 \leq k \leq N-3} \frac{\beta_k^{(\ell)}}{p_k - 2}, \min_{N-1 \leq k \leq N} \frac{\beta_k^{(\ell+1)}}{p_k - 2} \right\} \\ \vdots &= \vdots \\ \beta_{j-1}^{(\ell+1)} &= p_{j-1} \min \left\{ q_{j-2}, \min_{j \leq k \leq N} \frac{\beta_k^{(\ell+1)}}{p_k - 2} \right\}. \end{cases}$$

We first observe that this scheme is well-defined, since each $\beta_i^{(\ell+1)}$ is determined either by $(\beta_{j-1}^{(\ell)}, \dots, \beta_N^{(\ell)})$ or by an updated information on the $\beta_k^{(\ell+1)}$, with $k \geq i + 1$. Moreover, thanks to Lemma A.3 and Lemma A.4 below, we have that

$$\left\{ \beta_i^{(\ell)} \right\}_{\ell \in \mathbb{N}} \quad \text{is nondecreasing, for every } j-1 \leq i \leq N,$$

and there exists $\ell_0 \in \mathbb{N}$ such that

$$(6.12) \quad \text{for every } \ell \geq \ell_0, \quad \beta_i^{(\ell)} = p_i q_{j-2}, \quad \text{for } i = j-1, \dots, N.$$

With these definitions at hand, we now prove that

$$(6.13) \quad \text{for every } B_R \Subset B, \quad \sum_{i=j-1}^N \int_{B_R} |u_{x_i}|^{\beta_i^{(\ell)}} dx \leq C, \quad \text{for every } \ell \in \mathbb{N}, \quad \text{with } C > 0 \text{ independent of } \varepsilon.$$

By taking into account (6.12), this will eventually establish (6.6), thus concluding the proof.

In turn, the proof of (6.13) relies on an induction argument. The assertion (6.13) is true for $\ell = 0$, thanks to (6.9).

We now assume (6.13) to hold for some $\ell \in \mathbb{N}$ and establish the same for $\ell + 1$, i.e.

$$\text{for every } B_R \Subset B, \quad \sum_{i=j-1}^N \int_{B_R} |u_{x_i}|^{\beta_i^{(\ell+1)}} dx \leq C,$$

with C independent of ε . Actually, by a downward induction on $m = N, \dots, j-1$, we prove that

$$\sum_{i=m}^N \int_{B_R} |u_{x_i}|^{\beta_i^{(\ell+1)}} dx \leq C, \quad \text{with } C \text{ independent of } \varepsilon.$$

For the initialization step $m = N$, we apply (4.6) with $k = N$ and for the following choice of α :

$$p_N + 2 + \alpha = p_N \min \left\{ q_{j-2}, \min_{j-1 \leq k \leq N-1} \frac{\beta_k^{(\ell)}}{p_k - 2} \right\} = \beta_N^{(\ell+1)}.$$

In order to justify that α defined as such is non negative, we rely on the fact that for every $i \in \{j-1, \dots, N\}$, the sequence $\{\beta_i^{(\ell)}\}_{\ell \in \mathbb{N}}$ is nondecreasing. This implies that

$$\alpha \geq \beta_N^{(0)} - (p_N + 2) = p_N q_{j-1} - (p_N + 2) \geq p_N q_N - (p_N + 2) \geq 0.$$

Hence, such a choice of α is feasible.

We get that for every $B_R \Subset B_{R'} \Subset B$,

$$\begin{aligned} \int_{B_R} |u_{x_N}|^{\beta_N^{(\ell+1)}} dx &\leq C (R')^N \left(\left(\frac{\|u\|_{L^\infty(B)}}{R' - R} \right)^{\beta_N^{(\ell+1)}} + \varepsilon_0 \right) \\ &\quad + C \left(\frac{\|u\|_{L^\infty(B)}}{R' - R} \right)^{\frac{2}{p_N} \beta_N^{(\ell+1)}} \int_{B_{R'}} \sum_{i=1}^{j-2} |u_{x_i}|^{(p_i-2) \frac{\beta_N^{(\ell+1)}}{p_N}} dx \\ &\quad + C \left(\frac{\|u\|_{L^\infty(B)}}{R' - R} \right)^{\frac{2}{p_N} \beta_N^{(\ell+1)}} \int_{B_{R'}} \sum_{i=j-1}^{N-1} |u_{x_i}|^{\frac{p_i-2}{p_N} \beta_N^{(\ell+1)}} dx. \end{aligned}$$

For the terms in the first sum of the right hand side, we use Hölder's inequality with the exponent

$$\frac{p_i}{p_i - 2} \frac{p_N}{\beta_N^{(\ell+1)}},$$

and its conjugate³. In the second sum, we use Hölder's inequality with the exponent⁴

$$\frac{p_N}{p_i - 2} \frac{\beta_i^{(\ell)}}{\beta_N^{(\ell+1)}}.$$

³Observe that for $p_i = 2$ there is no need of Hölder's inequality. If $p_i > 2$, one can easily check that these exponents are larger than 1 by observing that

$$\beta_N^{(\ell+1)} \leq p_N q_{j-2} \leq p_N q_i, \quad \text{for } i \leq j-2.$$

⁴As before, there is no need of Hölder's inequality for $p_i = 2$. For $p_i > 2$, we rely on the fact that by definition

$$\beta_N^{(\ell+1)} \leq \beta_i^{(\ell)} \frac{p_N}{p_i - 2}, \quad \text{for } j-1 \leq i \leq N-1.$$

This justifies that the exponent is larger than 1.

One gets

$$\begin{aligned} \int_{B_R} |u_{x_N}|^{\beta_N^{(\ell+1)}} dx &\leq C (R')^N \left(\left(\frac{\|u\|_{L^\infty(B)}}{R' - R} \right)^{\beta_N^{(\ell+1)}} + \varepsilon_0 \right) \\ &\quad + C \left(\frac{\|u\|_{L^\infty(B)}}{R' - R} \right)^{2 \frac{\beta_N^{(\ell+1)}}{p_N}} \sum_{i=1}^{j-2} \left(\int_{B_{R'}} |u_{x_i}|^{p_i} dx \right)^{\frac{p_i-2}{p_i} \frac{\beta_N^{(\ell+1)}}{p_N}} \\ &\quad + C \left(\frac{\|u\|_{L^\infty(B)}}{R' - R} \right)^{2 \frac{\beta_N^{(\ell+1)}}{p_N}} \sum_{i=j-1}^{N-1} \left(\int_{B_{R'}} |u_{x_i}|^{\beta_i^{(\ell)}} dx \right)^{\frac{p_i-2}{p_N} \frac{\beta_N^{(\ell+1)}}{\beta_i^{(\ell)}}}. \end{aligned}$$

By using the induction assumption (6.13) to control the last term, Lemma 2.1 and Proposition 2.3 in order to control the other two, uniformly in ε , we get the desired estimate for u_{x_N} .

We now assume that for some $m \in \{j, \dots, N\}$, we have

$$(6.14) \quad \text{for every } B_R \Subset B, \quad \sum_{i=m}^N \int_{B_R} |u_{x_i}|^{\beta_i^{(\ell+1)}} dx \leq C, \quad \text{with } C > 0 \text{ independent of } \varepsilon,$$

and prove that this entails

$$\text{for every } B_R \Subset B, \quad \sum_{i=m-1}^N \int_{B_R} |u_{x_i}|^{\beta_i^{(\ell+1)}} dx \leq C, \quad \text{with } C > 0 \text{ independent of } \varepsilon.$$

Obviously, we only need to improve the control on the last component of the gradient, i.e. on $u_{x_{m-1}}$. We still rely on Proposition 4.3, this time with the choices

$$k = m - 1 \quad \text{and} \quad p_{m-1} + 2 + \alpha = p_{m-1} \min \left\{ q_{j-2}, \min_{j-1 \leq i \leq m-2} \frac{\beta_i^{(\ell)}}{p_i - 2}, \min_{m \leq i \leq N} \frac{\beta_i^{(\ell+1)}}{p_i - 2} \right\} = \beta_{m-1}^{(\ell+1)}.$$

The fact that $\beta_{m-1}^{(\ell+1)} \geq \beta_{m-1}^{(0)} \geq p_{m-1} + 2$ ensures that $\alpha \geq 0$. Hence, for every $B_R \Subset B_{R'} \Subset B$,

$$\begin{aligned} \int_{B_R} |u_{x_{m-1}}|^{\beta_{m-1}^{(\ell+1)}} dx &\leq C (R')^N \left(\left(\frac{\|u\|_{L^\infty(B)}}{R' - R} \right)^{\beta_{m-1}^{(\ell+1)}} + \varepsilon_0 \right) \\ &\quad + C \left(\frac{\|u\|_{L^\infty(B)}}{R' - R} \right)^{\frac{2}{p_{m-1}} \beta_{m-1}^{(\ell+1)}} \int_{B_{R'}} \sum_{i=1}^{j-2} |u_{x_i}|^{(p_i-2) \frac{\beta_{m-1}^{(\ell+1)}}{p_{m-1}}} dx \\ &\quad + C \left(\frac{\|u\|_{L^\infty(B)}}{R' - R} \right)^{\frac{2}{p_{m-1}} \beta_{m-1}^{(\ell+1)}} \int_{B_{R'}} \sum_{i=j-1}^{m-2} |u_{x_i}|^{(p_i-2) \frac{\beta_{m-1}^{(\ell+1)}}{p_{m-1}}} dx \\ &\quad + C \left(\frac{\|u\|_{L^\infty(B)}}{R' - R} \right)^{\frac{2}{p_{m-1}} \beta_{m-1}^{(\ell+1)}} \int_{B_{R'}} \sum_{i=m}^N |u_{x_i}|^{(p_i-2) \frac{\beta_{m-1}^{(\ell+1)}}{p_{m-1}}} dx. \end{aligned}$$

We now proceed as above: we control the last term by using the induction assumption (6.14) and the fact that if

$$\frac{\beta_{m-1}^{(\ell+1)}}{p_{m-1}} (p_i - 2) \leq \beta_i^{(\ell+1)}, \quad \text{if } m \leq i \leq N.$$

The third term is estimated thanks to the induction assumption (6.13) and the inequality⁵

$$\frac{\beta_{m-1}^{(\ell+1)}}{p_{m-1}}(p_i - 2) \leq \beta_i^{(\ell)}, \quad \text{if } j-1 \leq i \leq m-2.$$

Finally, on the two first terms, we use Lemma 2.1 and Proposition 2.3, and also that⁶

$$\frac{\beta_{m-1}^{(\ell+1)}}{p_{m-1}}(p_i - 2) \leq p_i, \quad \text{if } 1 \leq i \leq j-2.$$

This finally establishes that

$$\text{for every } B_R \Subset B, \quad \sum_{i=j-1}^N \int_{B_R} |u_{x_i}|^{\beta_i^{(\ell+1)}} dx \leq C,$$

with C independent of ε . As already explained, this is enough to safely conclude the proof. \square

7. PROOF OF THEOREM 1.1

The cornerstones of the proof of Theorem 1.1 are the uniform L^∞ estimate for the gradient of Proposition 5.1 and the uniform higher integrability estimate of Proposition 6.1. Indeed, by using Proposition 6.1 with the choice $q_0 = \gamma$ (i.e. the exponent in (5.1)), we get that for every $B_{r_0} \Subset B$ with $r_0 < 1$

$$\|\nabla u_\varepsilon\|_{L^\infty(B_{r_0})} \leq C,$$

with $C > 0$ independent of ε . Observe that to infer that C is independent of ε , we use Lemma 2.1 and Proposition 2.3. Once we have this uniform estimate at our disposal, the Lipschitz regularity of U follows with a standard covering argument, by taking into account that u_ε converges to U (see Lemma 2.2). We refer to the proof of [3, Theorem A] for details. \square

Once we have Theorem 1.1 at our disposal, we can prove a higher differentiability result *à la* Uhlenbeck. For the model case of the functional

$$\sum_{i=1}^N \frac{1}{p_i} \int |u_{x_i}|^{p_i} dx,$$

the following result considerably improves [6, Theorem 1.1].

Corollary 7.1. *Let $\mathbf{p} = (p_1, \dots, p_N)$ be such that $2 \leq p_1 \leq \dots \leq p_N$. Let $U \in W_{\text{loc}}^{1,\mathbf{p}}(\Omega)$ be a local minimizer of $\mathfrak{F}_{\mathbf{p}}$ such that*

$$U \in L_{\text{loc}}^\infty(\Omega).$$

Then

$$|U_{x_i}|^{\frac{p_i-2}{2}} U_{x_i} \in W_{\text{loc}}^{1,2}(\Omega), \quad \text{for } i = 1, \dots, N.$$

Proof. The proof is the same as the one in [6, Proposition 3.2]. It is based on Nirenberg's method of incremental quotients, which aims at differentiating the equation

$$\sum_{i=1}^N (|U_{x_i}|^{p_i-2} U_{x_i}) = 0,$$

⁵This part of the discussion is void when $m = j$.

⁶This part of the discussion is void when $j = 2$.

in a discrete sense. By proceeding as in [6], we get for every $j = 1, \dots, N$ and every pair of concentric balls $B_{r_0} \Subset B_{R_0} \Subset \Omega$

$$\sum_{i=1}^N \int_{B_{r_0}} \left| \frac{\delta_{h\mathbf{e}_j} \left(|U_{x_i}|^{\frac{p_i-2}{2}} U_{x_i} \right)}{|h|^{\frac{s_j+1}{2}}} \right|^2 dx \leq \frac{C}{(R_0 - r_0)^2} \sum_{i=1}^N \left(\int_{B_{R_0}} |U_{x_i}|^{p_i} dx \right)^{\frac{p_i-2}{p_i}} \left(\int_{B_{R_0}} \left| \frac{\delta_{h\mathbf{e}_j} U}{|h|^{\frac{s_j+1}{2}}} \right|^{p_i} \right)^{\frac{2}{p_i}},$$

see [6, equation (3.6)]. By using that $\nabla U \in L_{\text{loc}}^\infty$, we can choose

$$s_j = 1 \quad \text{so that} \quad \frac{s_j + 1}{2} = 1,$$

to control the last term on the right-hand side and obtain an estimate on

$$\sum_{i=1}^N \int_{B_{r_0}} \left| \frac{\delta_{h\mathbf{e}_j} \left(|U_{x_i}|^{\frac{p_i-2}{2}} U_{x_i} \right)}{|h|} \right|^2 dx, \quad j = 1, \dots, N,$$

which is uniform in $|h| \ll 1$. By appealing to the difference quotient characterization of Sobolev spaces, we get the conclusion. \square

APPENDIX A. CALCULUS LEMMAS

In this section, we separately present some proofs on the elementary facts for the sequences needed in the proof of our main result.

A.1. Tools for the Lipschitz estimate. In what follows, we denote as usual

$$2^* = \frac{2N}{N-2}, \quad \text{for } N \geq 3.$$

Lemma A.1. *Let $2 \leq p_1 \leq p_N$. We define*

$$j_0 = \min \left\{ j \in \mathbb{N} : j > \log_2 \left(\frac{N-2}{2} (p_N - 2) - \frac{N}{2} (p_1 - 2) \right) - 2 \right\},$$

$$j_1 = \min \left\{ j \in \mathbb{N} : j > \log_2 \left((N-2) (p_N - 2) - N (p_1 - 2) \right) - 2 \right\}.$$

and $J = 1 + \max\{j_0, j_1\}$. We set

$$\gamma_j = 2^{j+2} + p_N - 2,$$

and

$$\tau_j = \frac{\gamma_{j-1}}{\gamma_j} \frac{\frac{2^*}{2} (\gamma_j + p_1 - p_N) - \gamma_j}{\frac{2^*}{2} (\gamma_j + p_1 - p_N) - \gamma_{j-1}}.$$

Then there exist two constants $0 < C_1 < C_2 < 1$ depending on N, p_1 and p_N such that

$$C_1 \leq \frac{(1 - \tau_j) \gamma_j}{\gamma_j + p_1 - p_N} \leq C_2, \quad \text{for every } j \geq J.$$

Proof. It is easily seen that the sequence $\{\gamma_j\}_{j \in \mathbb{N}}$ is increasing. Moreover, by definition of j_0 , we have that

$$\gamma_j < \frac{2^*}{2} (\gamma_j + p_1 - p_N), \quad \text{for every } j \geq j_0,$$

thus τ_j is well-defined and positive for $j \geq j_0$. The definition of τ_j entails that

$$(A.1) \quad \frac{1}{\gamma_j} = \frac{\tau_j}{\gamma_{j-1}} + \frac{1 - \tau_j}{\frac{2^*}{2}(\gamma_j + p_1 - p_N)}.$$

thus the previous discussion implies that

$$0 < \tau_j < 1, \quad \text{for every } j \geq j_0.$$

Then the proof is by direct computation: we have

$$\begin{aligned} \frac{(1 - \tau_j) \gamma_j}{\gamma_j + p_1 - p_N} &= \left(1 - \frac{\gamma_{j-1}}{\gamma_j} \frac{\frac{2^*}{2}(\gamma_j + p_1 - p_N) - \gamma_j}{\frac{2^*}{2}(\gamma_j + p_1 - p_N) - \gamma_{j-1}} \right) \frac{\gamma_j}{\gamma_j + p_1 - p_N} \\ &= \frac{\gamma_j \frac{2^*}{2}(\gamma_j + p_1 - p_N) - \gamma_{j-1} \frac{2^*}{2}(\gamma_j + p_1 - p_N)}{\frac{2^*}{2}(\gamma_j + p_1 - p_N) - \gamma_{j-1}} \frac{1}{\gamma_j + p_1 - p_N} \\ &= \frac{2^*}{2} \frac{\gamma_j - \gamma_{j-1}}{\frac{2^*}{2}(\gamma_j + p_1 - p_N) - \gamma_{j-1}} \\ &= \frac{2^*}{2} \frac{2^{j+2} - 2^{j+1}}{\frac{2^*}{2}(2^{j+2} + p_1 - 2) - p_N - 2^{j+1} + 2} \\ &= \frac{2^*}{4} \frac{2^{j+2}}{\left(\frac{2^* - 1}{2}\right) 2^{j+2} + \frac{2^*}{2}(p_1 - 2) - (p_N - 2)}. \end{aligned}$$

We have to distinguish two cases: if

$$p_N < \frac{N}{N-2}(p_1 - 2) + 2,$$

then the function

$$(A.2) \quad t \mapsto \frac{t}{\left(\frac{2^* - 1}{2}\right) t + \frac{2^*}{2}(p_1 - 2) - (p_N - 2)},$$

is well-defined for every $t > 0$ and monotonically *increasing*. We have in this case

$$\begin{aligned} 0 < \frac{(1 - \tau_{j_0}) \gamma_{j_0}}{\gamma_{j_0} + p_1 - p_N} &\leq \frac{(1 - \tau_j) \gamma_j}{\gamma_j + p_1 - p_N} \\ &< \lim_{j \rightarrow \infty} \frac{2^*}{4} \frac{2^{j+2}}{\left(\frac{2^* - 1}{2}\right) 2^{j+2} + \frac{2^*}{2}(p_1 - 2) - (p_N - 2)} = \frac{1}{2} \frac{2^*}{2^* - 1} < 1, \end{aligned}$$

for every $j \geq j_0$.

On the other hand, if

$$(A.3) \quad p_N \geq \frac{N}{N-2}(p_1 - 2) + 2,$$

then the function (A.2) is well-defined and monotonically *decreasing* for

$$(A.4) \quad t > \frac{2}{2^* - 1} \left((p_N - 2) - \frac{2^*}{2} (p_1 - 2) \right).$$

Thus we now obtain

$$\frac{1}{2} \frac{2^*}{2^* - 1} \leq \frac{(1 - \tau_j) \gamma_j}{\gamma_j + p_1 - p_N} \leq \frac{2^*}{4} \frac{2^{j_1+2}}{\left(\frac{2^* - 1}{2} \right) 2^{j_1+2} + \frac{2^*}{2} (p_1 - 2) - (p_N - 2)},$$

for every $j \geq j_1$. Observe that the choice of j_1 assures firstly that $t = 2^{j+2}$ satisfies (A.4) whenever $j \geq j_1$ (here we use (A.3)) and secondly, that the right-hand side above is strictly smaller than 1. \square

Lemma A.2. *With the notation of Lemma A.1, we define the sequence $\{\varepsilon_j\}_{j \geq j_0}$ by*

$$(A.5) \quad 1 + \varepsilon_j = \tau_j \left(\frac{\gamma_j + p_1 - p_N}{(1 - \tau_j) \gamma_j} \right)', \quad \text{for } j \geq j_0.$$

Then

$$\varepsilon_j \sim \frac{N}{4} \frac{p_N - p_1}{2} \frac{1}{2^j}, \quad \text{for } j \rightarrow \infty.$$

In particular, we have

$$\lim_{n \rightarrow \infty} \prod_{i=j_0}^n (1 + \varepsilon_i) < +\infty.$$

Proof. We start by computing explicitly the conjugate exponent appearing in (A.5). We have

$$\begin{aligned} \left(\frac{\gamma_j + p_1 - p_N}{(1 - \tau_j) \gamma_j} \right)' &= \left(\frac{1}{1 - \tau_j} \left(1 + \frac{p_1 - p_N}{\gamma_j} \right) \right)' \\ &= \frac{\frac{1}{1 - \tau_j} \left(1 + \frac{p_1 - p_N}{\gamma_j} \right)}{\frac{1}{1 - \tau_j} \left(1 + \frac{p_1 - p_N}{\gamma_j} \right) - 1} = \frac{1 - \frac{p_N - p_1}{\gamma_j}}{\tau_j - \frac{p_N - p_1}{\gamma_j}}. \end{aligned}$$

Thus we have

$$\varepsilon_j = \tau_j \left(\frac{\gamma_j + p_1 - p_N}{(1 - \tau_j) \gamma_j} \right)' - 1 = \frac{\tau_j \left(1 - \frac{p_N - p_1}{\gamma_j} \right)}{\tau_j - \frac{p_N - p_1}{\gamma_j}} - 1 = \frac{p_N - p_1}{\gamma_j} \frac{1 - \tau_j}{\tau_j - \frac{p_N - p_1}{\gamma_j}}.$$

We now observe that

$$\frac{p_N - p_1}{\gamma_j} \frac{1 - \tau_j}{\tau_j - \frac{p_N - p_1}{\gamma_j}} \sim \frac{p_N - p_1}{\gamma_j} \frac{1 - \tau_j}{\tau_j}, \quad \text{for } j \rightarrow \infty.$$

Moreover, by using the definitions of γ_j and τ_j , we have

$$\frac{1 - \tau_j}{\tau_j} = \frac{1}{\tau_j} - 1 \sim \frac{N}{2}, \quad \text{for } j \rightarrow \infty,$$

which implies that

$$\varepsilon_j \sim \frac{N}{2} \frac{p_N - p_1}{\gamma_j}, \quad \text{for } j \rightarrow \infty.$$

By observing that $\gamma_j \sim 2^{j+2}$, we get the desired conclusion.

In order to prove the last part, it is enough to notice that

$$\prod_{j=j_0}^n (1 + \varepsilon_j) = \exp \left(\sum_{j=j_0}^n \log(1 + \varepsilon_j) \right) \quad \text{and} \quad \log(1 + \varepsilon_j) \sim \varepsilon_j, \quad \text{for } j \rightarrow \infty.$$

By using the first part of the proof and the definition of γ_j , we see that

$$\lim_{n \rightarrow \infty} \sum_{j=j_0}^n \varepsilon_j < +\infty.$$

This concludes the proof. \square

A.2. Tools for the higher integrability. In this section, we present some properties of the vector-valued sequence $\{(\beta_{j-1}^\ell, \dots, \beta_N^\ell)\}$ which were needed in the proof of Proposition 6.1, in order to complete the inductive step. We use the same notation as before: in particular, we fix $2 \leq q_0 < +\infty$ and set

$$q_j = \min \left\{ \left(\frac{p_j}{2} \right)', q_0 \right\}, \quad j = 1, \dots, N.$$

Then for a fixed index $j \in \{2, \dots, N\}$, we define

$$(A.6) \quad \begin{cases} \beta_{j-1}^{(0)} &= p_{j-1} \min \{q_{j-2}, q_{j-1} q_N\}, \\ \beta_i^{(0)} &= p_i q_{j-1}, \quad \text{for } i = j, \dots, N, \end{cases}$$

and by a recursive scheme

$$(A.7) \quad \begin{cases} \beta_N^{(\ell+1)} &= p_N \min \left\{ q_{j-2}, \min_{j-1 \leq k \leq N-1} \frac{\beta_k^{(\ell)}}{p_k - 2} \right\} \\ \beta_{N-1}^{(\ell+1)} &= p_{N-1} \min \left\{ q_{j-2}, \min_{j-1 \leq k \leq N-2} \frac{\beta_k^{(\ell)}}{p_k - 2}, \frac{\beta_N^{(\ell+1)}}{p_N - 2} \right\} \\ \beta_{N-2}^{(\ell+1)} &= p_{N-2} \min \left\{ q_{j-2}, \min_{j-1 \leq k \leq N-3} \frac{\beta_k^{(\ell)}}{p_k - 2}, \min_{N-1 \leq k \leq N} \frac{\beta_k^{(\ell+1)}}{p_k - 2} \right\} \\ \vdots &= \vdots \\ \beta_{j-1}^{(\ell+1)} &= p_{j-1} \min \left\{ q_{j-2}, \min_{j \leq k \leq N} \frac{\beta_k^{(\ell+1)}}{p_k - 2} \right\} \end{cases}$$

Lemma A.3. *For every $i \in \{j-1, \dots, N\}$, the sequence $\{\beta_i^{(\ell)}\}_{\ell \in \mathbb{N}}$ is nondecreasing.*

Proof. We proceed again by induction on ℓ .

Initialization step. We first need to prove that

$$(A.8) \quad \beta_i^{(1)} \geq \beta_i^{(0)}, \quad \text{for } i = j-1, \dots, N.$$

We establish this by downward induction on $i = N, \dots, j-1$. Indeed, for $i = N$, one has by definition

$$\beta_N^{(1)} = p_N \min \left\{ q_{j-2}, \min_{j-1 \leq k \leq N-1} \frac{\beta_k^{(0)}}{p_k - 2} \right\}.$$

By using the definition of $\beta_k^{(0)}$, the previous is the same as

$$\beta_N^{(1)} = p_N \min \left\{ q_{j-2}, q_{j-1} \min\{q_{j-2}, q_N q_{j-1}\}, q_{j-1} \min_{j \leq k \leq N-1} q_k \right\}.$$

By recalling that $q_{j-2} \geq \dots \geq q_N$, this gives

$$\beta_N^{(1)} = p_N \min \left\{ q_{j-2}, q_{j-1} \min\{q_{j-2}, q_N q_{j-1}\}, q_{j-1} q_{N-1} \right\} \geq p_N \min\{q_{j-2}, q_{j-1} q_N\}.$$

Hence,

$$\beta_N^{(0)} = p_N q_{j-1} \leq p_N \min\{q_{j-2}, q_{j-1} q_N\} = \beta_N^{(1)}.$$

This proves (A.8) for $i = N$.

We now assume that for some $i \in \{j-1, \dots, N\}$, property (A.8) holds for every $k \in \{i+1, \dots, N\}$. One proceeds to prove that (A.8) holds for i , as well. By definition of $\beta_i^{(1)}$,

$$\beta_i^{(1)} = p_i \min \left\{ q_{j-2}, \min_{j-1 \leq k \leq i-1} \frac{\beta_k^{(0)}}{p_k - 2}, \min_{i+1 \leq k \leq N} \frac{\beta_k^{(1)}}{p_k - 2} \right\}.$$

By the induction assumption, $\beta_k^{(1)} \geq \beta_k^{(0)}$ for $k \geq i+1$ and thus

$$\begin{aligned} \beta_i^{(1)} &\geq p_i \min \left\{ q_{j-2}, \min_{j-1 \leq k \leq i-1} \frac{\beta_k^{(0)}}{p_k - 2}, \min_{i+1 \leq k \leq N} \frac{\beta_k^{(0)}}{p_k - 2} \right\} \\ &= p_i \min \left\{ q_{j-2}, \frac{\beta_{j-1}^{(0)}}{p_{j-1} - 2}, \min_{\substack{j \leq k \leq N \\ k \neq i}} \frac{\beta_k^{(0)}}{p_k - 2} \right\}. \end{aligned}$$

By definition of $\beta_i^{(0)}$, this gives

$$\begin{aligned} \beta_i^{(1)} &\geq p_i \min \left\{ q_{j-2}, q_{j-1} \min\{q_{j-2}, q_N q_{j-1}\}, q_{j-1} \min_{\substack{j \leq k \leq N \\ k \neq i}} q_k \right\} \\ &= p_i \min\{q_{j-2}, q_{j-1} q_N\}. \end{aligned}$$

When $i \geq j$, this implies $\beta_i^{(1)} \geq p_i q_{j-1} = \beta_i^{(0)}$, while when $i = j-1$, one has

$$\beta_{j-1}^{(1)} \geq p_{j-1} \min\{q_{j-2}, q_N q_{j-1}\} = \beta_{j-1}^{(0)}.$$

We have thus proved (A.8) for i , which completes the proof.

Inductive step. We now assume that for an index $\ell \geq 1$, we have

$$(A.9) \quad \beta_i^{(\ell)} \geq \beta_i^{(\ell-1)}, \quad \text{for } i = j-1, \dots, N.$$

We need to prove that this entails

$$\beta_i^{(\ell+1)} \geq \beta_i^{(\ell)}, \quad \text{for } i = j-1, \dots, N,$$

as well.

We rely again on a downward induction on $i = N, \dots, j-1$. Indeed, for $i = N$, we can use (A.9), which gives

$$\beta_N^{(\ell+1)} = p_N \min \left\{ q_{j-2}, \min_{j-1 \leq k \leq N-1} \frac{\beta_k^{(\ell)}}{p_k - 2} \right\} \geq p_N \min \left\{ q_{j-2}, \min_{j-1 \leq k \leq N-1} \frac{\beta_k^{(\ell-1)}}{p_k - 2} \right\} = \beta_N^{(\ell)}.$$

We now assume that for some $i \geq j - 1$, one has

$$\beta_k^{(\ell+1)} \geq \beta_k^{(\ell)}, \quad \text{for every } k \in \{i + 1, \dots, N\}.$$

Then

$$\begin{aligned} \beta_i^{(\ell+1)} &= p_i \min \left\{ q_{j-2}, \min_{j-1 \leq k \leq i-1} \frac{\beta_k^{(\ell)}}{p_k - 2}, \min_{i+1 \leq k \leq N} \frac{\beta_k^{(\ell+1)}}{p_k - 2} \right\} \\ &\geq p_i \min \left\{ q_{j-2}, \min_{j-1 \leq k \leq i-1} \frac{\beta_k^{(\ell)}}{p_k - 2}, \min_{i+1 \leq k \leq N} \frac{\beta_k^{(\ell)}}{p_k - 2} \right\}. \end{aligned}$$

Relying now on the induction assumption (A.9), one gets

$$\beta_i^{(\ell+1)} \geq p_i \min \left\{ q_{j-2}, \min_{j-1 \leq k \leq i-1} \frac{\beta_k^{(\ell-1)}}{p_k - 2}, \min_{i+1 \leq k \leq N} \frac{\beta_k^{(\ell)}}{p_k - 2} \right\} = \beta_i^{(\ell)}.$$

This completes the proof. \square

Lemma A.4. *With the notation of the previous lemma, there exists $\ell_0 \in \mathbb{N}$ such that for every $\ell \geq \ell_0$, one has*

$$\beta_i^{(\ell)} = p_i q_{j-2}, \quad \text{for every } i = j - 1, \dots, N.$$

Proof. By using the monotonicity proved in the previous lemma, we get in particular for $i = j - 1, \dots, N$,

$$\begin{aligned} \beta_i^{(\ell+1)} &= p_i \min \left\{ q_{j-2}, \min_{j-1 \leq k \leq i-1} \frac{\beta_k^{(\ell)}}{p_k - 2}, \min_{i+1 \leq k \leq N} \frac{\beta_k^{(\ell+1)}}{p_k - 2} \right\} \\ (A.10) \quad &\geq p_i \min \left\{ q_{j-2}, \min_{\substack{j-1 \leq k \leq N \\ k \neq i}} \frac{\beta_k^{(\ell)}}{p_k - 2} \right\} \geq p_i \min \left\{ q_{j-2}, \min_{j-1 \leq k \leq N} \frac{\beta_k^{(\ell)}}{p_k - 2} \right\}. \end{aligned}$$

Dividing by p_i and observing that

$$\frac{\beta_k^{(\ell)}}{p_k - 2} \geq \frac{\beta_k^{(\ell)}}{p_k} q_k \geq \frac{\beta_k^{(\ell)}}{p_k} q_N,$$

one deduces that

$$\frac{\beta_i^{(\ell+1)}}{p_i} \geq \min \left\{ q_{j-2}, q_N \min_{j-1 \leq k \leq N} \frac{\beta_k^{(\ell)}}{p_k} \right\}.$$

Since this is true for every $i = j - 1, \dots, N$, this implies

$$(A.11) \quad \delta^{(\ell+1)} \geq \min \left\{ q_{j-2}, q_N \delta^{(\ell)} \right\}, \quad \text{where } \delta^{(\ell)} = \min_{j-1 \leq k \leq N} \frac{\beta_k^{(\ell)}}{p_k}.$$

The monotonicity of each sequence $\{\beta_k^{(\ell)}\}_{\ell \in \mathbb{N}}$ entails the monotonicity of $\{\delta^{(\ell)}\}_{\ell \in \mathbb{N}}$. We claim that there exists $\ell_0 \in \mathbb{N}$ such that one has

$$(A.12) \quad q_N \delta^{(\ell)} \geq q_{j-2}, \quad \text{for every } \ell \geq \ell_0.$$

Indeed, assume by contradiction that $q_N \delta^{(\ell)} < q_{j-2}$ for every $\ell \geq 0$. Then it follows from (A.11) that

$$\delta^{(\ell+1)} \geq q_N \delta^{(\ell)}, \quad \text{for every } \ell \geq 0.$$

which implies in turn that $\delta^{(\ell)} \nearrow +\infty$, as $\ell \rightarrow \infty$. This contradicts that $q_N \delta^{(\ell)} < q_{j-2}$ for every $\ell \geq 0$. Hence, the claim (A.12) is established. By (A.11) again, this implies that

$$\frac{\beta_i^{(\ell+1)}}{p_i} \geq \min_{j-1 \leq k \leq N} \frac{\beta_k^{(\ell+1)}}{p_k} = \delta^{(\ell+1)} \geq q_{j-2}, \quad \text{for } i = j-1, \dots, N, \text{ for every } \ell \geq \ell_0 - 1.$$

Since the opposite estimate on $\beta_i^{(\ell+1)}/p_i$ is a consequence of the definition of $\beta_i^{(\ell+1)}$, we obtain the desired conclusion. \square

REFERENCES

- [1] P. Baroni, A. Di Castro, G. Palatucci, Intrinsic geometry and De Giorgi classes for certain anisotropic problems, *Discrete Contin. Dyn. Syst. Ser. S*, **10** (2017), 647–659. [2](#)
- [2] P. Bousquet, L. Brasco, C^1 regularity of orthotropic p -harmonic functions in the plane, *Anal. PDE*, **11** (2018), 813–854. [1](#), [3](#), [5](#)
- [3] P. Bousquet, L. Brasco, V. Julin, Lipschitz regularity for local minimizers of some widely degenerate problems, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, **26** (2016), 1–40. [1](#), [3](#), [9](#), [12](#), [29](#)
- [4] P. Bousquet, L. Brasco, C. Leone, A. Verde, On the Lipschitz character of orthotropic p -harmonic functions, *Calc. Var. Partial Differential Equations*, **57** (2018), 57:88 [1](#), [3](#), [5](#), [6](#), [11](#), [13](#), [14](#), [18](#)
- [5] L. Brasco, G. Carlier, On certain anisotropic elliptic equations arising in congested optimal transport: local gradient bounds, *Adv. Calc. Var.*, **7** (2014), 379–407. [1](#)
- [6] L. Brasco, C. Leone, G. Pisante, A. Verde, Sobolev and Lipschitz regularity for local minimizers of widely degenerate anisotropic functionals, *Nonlinear Anal.*, **153** (2017), 169–199. [1](#), [3](#), [9](#), [18](#), [29](#), [30](#)
- [7] M. Bildahuer, M. Fuchs, X. Zhong, A regularity theory for scalar local minimizers of splitting-type variational integrals, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, **6** (2007), 385–404. [6](#), [7](#), [16](#)
- [8] V. Bögelein, F. Duzaar, P. Marcellini, Parabolic equations with p, q -growth, *J. Math. Pures Appl.*, **100** (2013), 535–563. [3](#)
- [9] E. Bombieri, E. De Giorgi, M. Miranda, Una maggiorazione a priori relativa alle ipersuperfici minimali non parametriche, *Arch. Rational Mech. Anal.*, **32** (1969), 255–267. [4](#)
- [10] H. J. Choe, Interior behaviour of minimizers for certain functionals with nonstandard growth, *Nonlinear Anal.*, **19** (1992), 933–945. [6](#), [7](#)
- [11] G. Cupini, P. Marcellini, E. Mascolo, Regularity of minimizers under limit growth conditions, *Nonlinear Analysis*, **153** (2017) 294–310. [3](#)
- [12] G. Cupini, P. Marcellini, E. Mascolo, Local boundedness of minimizers with limit growth conditions, *J. Optim. Theory Appl.*, **166** (2015), 1–22. [3](#)
- [13] F. Demengel, Regularity properties of viscosity solutions for fully nonlinear equations on the model of the anisotropic p -Laplacian, *Asymptot. Anal.*, **105** (2017), 27–43. [3](#), [6](#)
- [14] F. Demengel, Lipschitz interior regularity for the viscosity and weak solutions of the pseudo p -Laplacian equation, *Adv. Differential Equations*, **21** (2016), 373–400. [3](#)
- [15] E. DiBenedetto, $C^{1+\alpha}$ local regularity of weak solutions of degenerate elliptic equations, *Nonlinear Anal.*, **7** (1983), 827–850. [6](#)
- [16] E. DiBenedetto, U. Gianazza, V. Vespri, Remarks on Local Boundedness and Local Hölder Continuity of Local Weak Solutions to Anisotropic p -Laplacian Type Equations, *J. Elliptic Parabol. Equ.*, **2** (2016), 157–169. [3](#)
- [17] L. Esposito, F. Leonetti, G. Mingione, Sharp regularity for functionals with (p, q) growth, *J. Differential Equations*, **204** (2004), 5–55. [2](#)
- [18] L. Esposito, F. Leonetti, G. Mingione, Regularity for minimizers of functionals with $p - q$ growth, *NoDEA Nonlinear Differential Equations Appl.*, **6** (1999), 133–148. [2](#)
- [19] N. Fusco, C. Sbordone, Some remarks on the regularity of minima of anisotropic integrals, *Commun. Partial Differ. Equations*, **18** (1993), 153–167. [3](#)
- [20] N. Fusco, C. Sbordone, Local boundedness of minimizers in a limit case, *Manuscripta Math.*, **69** (1990), 19–25. [3](#)
- [21] M. Giaquinta, Growth conditions and regularity, a counterexample, *Manuscripta Math.*, **59** (1987), 245–248. [2](#)
- [22] E. Giusti, *Direct methods in the calculus of variations*. World Scientific Publishing Co., Inc., River Edge, NJ, 2003. [11](#)
- [23] Hong Min-Chun, Some remarks on the minimizers of variational integrals with non standard growth conditions, *Boll. Un. Mat. Ital. A*, **6** (1992), 91–101. [2](#)
- [24] I. M. Kolodnii, The boundedness of generalized solutions of elliptic differential equations, *Moscow Univ. Math. Bull.*, **25** (1970), 31–37. [2](#), [3](#)

- [25] A. G. Korolëv, Boundedness of generalized solutions of elliptic differential equations, Russian Math. Surveys, **38** (1983), 186–187. [2](#)
- [26] G. M. Lieberman, Gradient estimates for anisotropic elliptic equations, Adv. Differential Equations, **10** (2005), 767–812. [3](#), [4](#)
- [27] P. Lindqvist, D. Ricciotti, Regularity for an anisotropic equation in the plane, Nonlinear Analysis (2018), <https://doi.org/10.1016/j.na.2018.02.002> [3](#)
- [28] P. Marcellini, Regularity and existence of solutions of elliptic equations with p, q -growth conditions, J. Differential Equations, **90** (1991), 1–30. [2](#)
- [29] P. Marcellini, Regularity of minimizers of integrals of the Calculus of Variations under non standard growth conditions, Arch. Rational Mech. Anal., **105** (1989), 267–284. [2](#)
- [30] P. Marcellini, Un exemple de solution discontinue d'un problème variationnel dans le cas scalaire, preprint n. 11 dell'Ist. Mat. Univ. Firenze (1987). Available at <http://web.math.unifi.it/users/marcell/lavori> [2](#)
- [31] J. H. Michael, L. M. Simon, Sobolev and mean-value inequalities on generalized submanifolds of \mathbb{R}^n , Comm. Pure Appl. Math., **26** (1973), 361–379. [4](#)
- [32] L. Simon, Interior gradient bounds for non-uniformly elliptic equations, Indiana Univ. Math. J., **25** (1976), 821–855. [4](#)
- [33] G. Stampacchia, On some regular multiple integral problems in the calculus of variations, Comm. Pure Appl. Math., **16** (1963), 383–421. [11](#)
- [34] N. Uralt'seva, N. Urdaletova, The boundedness of the gradients of generalized solutions of degenerate quasilinear nonuniformly elliptic equations, Vest. Leningr. Univ. Math., **16** (1984), 263–270. [2](#), [3](#)

(P. Bousquet) INSTITUT DE MATHÉMATIQUES DE TOULOUSE, CNRS UMR 5219
 UNIVERSITÉ DE TOULOUSE
 F-31062 TOULOUSE CEDEX 9, FRANCE.
E-mail address: pierre.bousquet@math.univ-toulouse.fr

(L. Brasco) DIPARTIMENTO DI MATEMATICA E INFORMATICA
 UNIVERSITÀ DEGLI STUDI DI FERRARA
 VIA MACHIAVELLI 35, 44121 FERRARA, ITALY
E-mail address: lorenzo.brasco@unife.it