

The weakness of being cohesive, thin or free in reverse mathematics

Ludovic Patey

► **To cite this version:**

Ludovic Patey. The weakness of being cohesive, thin or free in reverse mathematics. *Israël Journal of Mathematics*, The Hebrew University Magnes Press, 2016, 216 (2), pp.905 - 955. <10.1007/s11856-016-1433-3>. <hal-01888606>

HAL Id: hal-01888606

<https://hal.archives-ouvertes.fr/hal-01888606>

Submitted on 5 Oct 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THE WEAKNESS OF BEING COHESIVE, THIN OR FREE IN REVERSE MATHEMATICS

LUDOVIC PATEY

ABSTRACT. Informally, a mathematical statement is *robust* if its strength is left unchanged under variations of the statement. In this paper, we investigate the lack of robustness of Ramsey's theorem and its consequence under the frameworks of reverse mathematics and computable reducibility. To this end, we study the degrees of unsolvability of cohesive sets for different uniformly computable sequence of sets and identify different layers of unsolvability. This analysis enables us to answer some questions of Wang about how typical sets help computing cohesive sets.

We also study the impact of the number of colors in the computable reducibility between coloring statements. In particular, we strengthen the proof by Dzhafarov that cohesiveness does not strongly reduce to stable Ramsey's theorem for pairs, revealing the combinatorial nature of this non-reducibility and prove that whenever k is greater than ℓ , stable Ramsey's theorem for n -tuples and k colors is not computably reducible to Ramsey's theorem for n -tuples and ℓ colors. In this sense, Ramsey's theorem is not robust with respect to his number of colors over computable reducibility. Finally, we separate the thin set and free set theorem from Ramsey's theorem for pairs and identify an infinite decreasing hierarchy of thin set theorems in reverse mathematics. This shows that in reverse mathematics, the strength of Ramsey's theorem is very sensitive to the number of colors in the output set. In particular, it enables us to answer several related questions asked by Cholak, Giusto, Hirst and Jockusch.

1. INTRODUCTION

Ramsey's theorem (RT_k^n) asserts that any k -coloring of $[\mathbb{N}]^n$ admits an infinite monochromatic set, where $[\mathbb{N}]^n$ stands for the n -tuples over \mathbb{N} . In this paper, we study the lack of *robustness* of Ramsey's theorem and its consequences under the frameworks of reverse mathematics and computable reducibility. Informally, a mathematical statement is *robust* within a given framework if its strength is invariant under slight variations of the statement. In reverse mathematics, robustness can be understood as equiprovability of the various statements over the base theory RCA_0 , while in computability, a Π_2^1 statement is robust if its variations are computably equivalent. We shall detail further reverse mathematics and computable reducibility in sections 1.1 and 1.2. Our investigations follow three axes.

Axis 1: We first study the degrees of unsolvability of cohesiveness. Given a sequence of sets of integers R_0, R_1, \dots , an infinite set is \vec{R} -*cohesive* if it is almost included in R_i or \overline{R}_i for each i . COH is a consequence of Ramsey's theorem for pairs which finds many practical applications in computability and reverse mathematics. Jockusch and Stephan [24] have studied the computational strength of cohesive sets for maximally difficult sequences of sets. We reveal that COH contains hierarchies of complexity of its instances by establishing a one-to-one correspondence between instances of COH and instances of König's lemma (KL). This shows that the strength of COH depends on the considered class of its instances, and is therefore not *inner robust*. This correspondance enables us to reprove the existence of a computable sequence of sets with no low cohesive set [24] and to answer several questions asked by Wang [46] about how randomness and genericity help in solving computably unsolvable instances of COH.

Axis 2: A simple color amalgamation argument shows that RT_k^n and RT_ℓ^n are provably equivalent in reverse mathematics whenever $k, \ell \geq 2$. In this sense, Ramsey's theorem is robust with respect to numbers of colors in reverse mathematics. However, the standard proof that $\text{RT}_k^n \rightarrow \text{RT}_{k+1}^n$ involves two applications of RT_k^n . Mileti [31] first wondered whether those two applications were really necessary. The question has been later formalized thanks to Weihrauch and computable reducibility and investigated by Dorais, Dzhafarov, Hirst, Mileti and Shafer [9],

Hirschfeldt and Jockusch [18], Brattka and Rakotoniaina [4], among others. We answer positively by proving that for every $n \geq 2$, RT_k^n is not computably reducible to RT_ℓ^n whenever $k > \ell \geq 2$. Therefore, Ramsey’s theorem is not robust with respect to the number of colors under computable reducibility.

Axis 3: Last, we investigate the reverse mathematics of a weakening of Ramsey’s theorem in which more colors are allowed in the resulting set. The thin set theorem (TS_k^n) asserts that for any k -coloring of $[\mathbb{N}]^n$, there is an infinite set H such that $[H]^n$ avoids at least one color. We show that the thin set theorem is not robust in reverse mathematics by proving that for every $n, m, \ell \geq 2$, TS_k^n does not imply TS_ℓ^m for sufficiently large k ’s. This is the first example of an infinite decreasing hierarchy in reverse mathematics. This enables us to answer several questions from Cholak, Giusto, Hirst and Jockusch [5], Montálban [33] and Hirschfeldt [17] about the strength of the thin set theorem and its strengthening, the free set theorem, with respect to Ramsey’s theorem for pairs.

1.1. Reverse mathematics

Reverse mathematics is a vast mathematical program whose goal is to classify ordinary theorems in terms of their provability strength. It uses the framework of subsystems of second order arithmetic, which is sufficiently rich to express many theorems in a natural way. The base system, RCA_0 standing for Recursive Comprehension Axiom, contains the basic first order Peano arithmetic together with the Δ_1^0 comprehension scheme and the Σ_1^0 induction scheme. Thanks to the equivalence between Δ_1^0 -definable sets and computable sets, RCA_0 can be considered as capturing “computable mathematics”. The proof-theoretic analysis of the theorems in reverse mathematics is therefore closely related to their computational analysis. See Simpson [43] for a formal introduction to reverse mathematics.

Early reverse mathematics have led to two main empirical observations: First, many ordinary (i.e. non set-theoretic) theorems require very weak set existence axioms. Second, most of those theorems are in fact *equivalent* to one of five main subsystems, known as the “Big Five”. However, among the theorems studied in reverse mathematics, a notable class of theorems fails to support the second observation, namely, Ramsey-type theorems. The underlying idea of Ramsey’s theory is that whenever a collection of objects is sufficiently large, we can always find an arbitrarily large sub-collection of objects satisfying some given structural property. Perhaps the most well-known statement is Ramsey’s theorem, stating that every coloring of tuples of integers with a finite number of colors admits an infinite monochromatic subset. The various consequences of Ramsey’s theorem usually fail to coincide with the main five subsystems, and slight variations of their statements lead to different subsystems. The study of Ramsey-type statements has been a very active research subject in reverse mathematics over the past few years [2, 6, 14, 19]. See Hirschfeldt [17] for a good introduction to recent reverse mathematics.

1.2. Reducibilities

Many theorems in reverse mathematics are Π_2^1 statements, i.e., of the form $(\forall X)(\exists Y)\Phi(X, Y)$ where Φ is an arithmetic formula. They can be considered as *problems* which usually come with a natural class of *instances*. Given an instance X , a set Y such that $\Phi(X, Y)$ holds is called a *solution* to X . For example, König’s lemma states that every infinite, finitely branching tree has an infinite path. In this statement, an instance is a infinite, finitely branching tree T , and a solution to T is an infinite path through T .

Thanks to the computational nature of the axioms of RCA_0 , given two Π_2^1 statements P and Q , a proof of implication $Q \rightarrow P$ consists in taking an arbitrary P -instance I and computing a solution to I in a computational process involving several applications of the Q principle. If the proof relativizes and can be formalized over RCA_0 (the main concern being the restriction to Σ_1^0 -induction), we obtain a proof of $\text{RCA}_0 \vdash Q \rightarrow P$. It is often the case that the proof of the implication $Q \rightarrow P$ involves only one application of Q given an instance of P . Such a reduction is called a *computable reduction*.

Definition 1.1 (Computable reducibility) Fix two Π_2^1 statements P and Q .

1. P is *computably reducible* to Q (written $P \leq_c Q$) if every P -instance I computes a Q -instance J such that for every solution X to J , $X \oplus I$ computes a solution to I .
2. P is *strongly computably reducible* to a Q (written $P \leq_{sc} Q$) if every P -instance I computes a Q -instance J such that every solution to J computes a solution to I .

Of course, proving that a statement P is not computably reducible to another statement Q is not sufficient for separating the statements over RCA_0 . For example, we shall see that Ramsey's theorem for pairs with $k + 1$ colors is not computably reducible to Ramsey's theorem for pairs with k colors, whereas the statements are known to be logically equivalent over RCA_0 . However, proving that $P \not\leq_c Q$ can be seen as a preliminary step towards the separation of the principles. Lerman et al. [29] have developed a framework for iterating a one-step non-reducibility into a separation over RCA_0 .

Other reducibility notions have been introduced to better understand the computational content of theorems from the point of view of reverse mathematics. Dorais et al. [9] studied the uniformity of the computable reductions $P \leq_c Q$ by requiring the construction of a Q -instance J given a P -instance I and the construction of a solution to I given a solution to J to be done with two fixed Turing functionals. They showed that this *uniform reducibility* is the restriction of the Weihrauch reduction to the second-order setting. Hirschfeldt and Jockusch [18] introduced a game-theoretic approach and defined a *generalized uniform reducibility* extending the notion of uniform reducibility to several applications of the statement Q . In this paper, we shall restrict ourselves to computable reducibility and provability over RCA_0 .

1.3. Degrees of unsolvability of cohesiveness

Cohesiveness plays a central role in reverse mathematics. It appears naturally in the standard proof of Ramsey's theorem, as a preliminary step to reduce an instance of Ramsey's theorem over $(n + 1)$ -tuples into a non-effective instance over n -tuples. An important part of current research about Ramsey-type principles in reverse mathematics consists in trying to understand whether cohesiveness is a consequence of stable Ramsey's theorem for pairs, or more generally whether it is a combinatorial consequence of the infinite pigeonhole principle [6, 12, 11, 46]. Chong et al. [8] recently showed using non-standard models that cohesiveness is not a proof-theoretic consequence of the pigeonhole principle. However it is not known whether or not cohesiveness is computably reducible to stable Ramsey's theorem for pairs.

Definition 1.2 (Cohesiveness) An infinite set C is \vec{R} -cohesive for a sequence of sets R_0, R_1, \dots if for each $i \in \omega$, $C \subseteq^* R_i$ or $C \subseteq^* \bar{R}_i$. A set C is *p-cohesive* if it is \vec{R} -cohesive where \vec{R} is an enumeration of all primitive recursive sets. COH is the statement "Every uniform sequence of sets \vec{R} has an \vec{R} -cohesive set."

Jockusch and Stephan [24] studied the degrees of unsolvability of cohesiveness and proved that COH admits a universal instance whose solutions are the p-cohesive sets. They characterized their degrees as those whose jump is PA relative to \emptyset' .

Cohesiveness is a Π_2^1 statement whose instances are sequences of sets \vec{R} and whose solutions are \vec{R} -cohesive sets. It is natural to wonder about the degrees of unsolvability of the \vec{R} -cohesive sets according to the sequence of sets \vec{R} . Mingzhong Cai asked whether whenever a uniformly computable sequence of sets R_0, R_1, \dots has no computable \vec{R} -cohesive set, there exists a non-computable set which does not compute one. In the opposite direction, one may wonder whether every unsolvable instance of COH is maximally difficult. A natural first approach in the analysis of the strength of a principle consists in looking in which way typical sets can help in computing a solution to an unsolvable instance. The notion of typical set is usually understood in two different ways: using the genericity approach and the randomness approach. Wang [46] answered Cai's question by investigating the solvability of cohesiveness by typical sets.

In this paper, we refine Wang's analysis by establishing a pointwise correspondence between sets cohesive for a sequence and sets whose jump computes a member of a $\Pi_1^{0, \emptyset'}$ class. Then, using the known interrelations between typical sets and Π_1^0 classes, we give precise genericity

and randomness bounds above which no typical set helps computing a cohesive set. We identify different layers of unsolvability and spot a class of instances sharing many properties with the universal instance. Emulating work in [34] on the pigeonhole principle and weak König's lemma (WKL_0), we show that some unsolvable instances of COH are combinatorial consequences of the pigeonhole principle.

1.4. Ramsey's theorem and computable reducibility

The strength of Ramsey-type statements is notoriously hard to tackle in the setting of reverse mathematics. The separation of Ramsey's theorem for pairs (RT_2^2) from the arithmetical comprehension axiom (ACA_0) was a long-standing open problem, until Seetapun and Slaman solved it [40] with his notion of cone avoidance. The question of the relation between RT_2^2 and weak König's lemma (WKL_0) remained open for many years before Cholak, Jockusch and Slaman [6] proved that WKL_0 does not imply RT_2^2 over RCA_0 . More than fifteen years after Seetapun, Liu [30] solved the remaining direction by proving that RT_2^2 does not imply WKL_0 over RCA_0 .

Definition 1.3 (Ramsey's theorem) A subset H of ω is *homogeneous* for a coloring $f : [\omega]^n \rightarrow k$ (or *f-homogeneous*) if each n -tuple over H is given the same color by f . A coloring $f : [\omega]^{n+1} \rightarrow k$ is *stable* if for every n -tuple $\sigma \in [\omega]^n$, $\lim_s f(\sigma, s)$ exists. RT_k^n is the statement "Every coloring $f : [\omega]^n \rightarrow k$ has an infinite f -homogeneous set". SRT_k^n is the restriction of RT_k^n to stable colorings.

Simpson [43, Theorem III.7.6] proved that whenever $n \geq 3$ and $k \geq 2$, $\text{RCA}_0 \vdash \text{RT}_k^n \leftrightarrow \text{ACA}_0$. Ramsey's theorem for pairs is probably the most famous example of statement escaping the Big Five. Seetapun [40] proved that RT_2^2 is strictly weaker than ACA_0 over RCA_0 . Because of the complexity of the related separations, RT_2^2 received a particular attention from the reverse mathematics community. Mileti [31] and Jockusch and Lempp [unpublished] proved that RT_2^2 is equivalent to $\text{SRT}_2^2 + \text{COH}$ over RCA_0 . Recently, Chong et al. [8] proved that SRT_2^2 is strictly weaker than RT_2^2 over RCA_0 . However they used non-standard models to separate the statements and the question whether SRT_2^2 and RT_2^2 coincide over ω -models remains open. Stable Ramsey's theorem for pairs can be characterized by a purely computability-theoretic statement.

Definition 1.4 For every $n, k \geq 1$, D_k^n is the statement "Every Δ_n^0 k -partition of the integers has an infinite subset of one of its parts". $\text{D}_{<\infty}^n$ is the statement $(\forall k)\text{D}_k^n$.

Cholak et al. [6] proved that D_k^2 and SRT_k^2 are computably equivalent and that the proof is formalizable over $\text{RCA}_0 + \text{B}\Sigma_2^0$. Later, Chong et al. [7] proved that D_2^2 implies $\text{B}\Sigma_2^0$ over RCA_0 , showing therefore that $\text{RCA}_0 \vdash \text{D}_k^2 \leftrightarrow \text{SRT}_\ell^2$ for every $k, \ell \geq 2$. Dzhafarov [12] proved that $\text{COH} \not\leq_{sc} \text{D}_{<\infty}^2$ by constructing a sequence of sets R_0, R_1, \dots such that for every $k \geq 2$, every instance of RT_k^1 hyperarithmetical in \vec{R} has a solution which does not compute an \vec{R} -cohesive set. In section 3.1, we strengthen this result by making \vec{R} uniformly Δ_2^0 and removing the effectiveness restriction on the instance of RT_k^1 . The proof reveals the combinatorial nature of the relations between cohesiveness and RT_k^1 and answers a question of Antonio Montálban. Recently, Dzhafarov [10] proved that $\text{COH} \not\leq_{sc} \text{SRT}_2^2$.

Another closely related subject of interest is the impact of the number of colors in the strength of Ramsey's theorem. For every $n \geq 1$ and $k, \ell \geq 2$, $\text{RCA}_0 \vdash \text{RT}_k^n \leftrightarrow \text{RT}_\ell^n$ by a simple color blindness argument. Whenever $k > \ell \geq 2$, the reduction of RT_k^n to RT_ℓ^n involves more than one application of RT_ℓ^n and therefore is not a computable reduction. Hirschfeldt and Jockusch [18] noticed that the proof of Dzhafarov [12] can be modified to obtain $\text{RT}_k^1 \not\leq_{sc} \text{RT}_\ell^1$. Dorais et al. [9] asked in which case $\text{RT}_k^n \leq_c \text{RT}_\ell^n$. In section 3.2, we answer by proving that $\text{SRT}_k^n \not\leq_c \text{RT}_\ell^n$ whenever $k > \ell \geq 2$ and $n \geq 2$.

1.5. The weakness of free set and thin set theorems

Simpson [43, Theorem III.7.6] proved that the hierarchy of Ramsey's theorem collapses at level three in reverse mathematics. One may wonder about some natural weakenings of Ramsey's

theorem over arbitrary tuples which remain strictly weaker than ACA_0 . Given a coloring $f : [\omega]^n \rightarrow k$, instead of stating the existence of an infinite f -homogeneous set H , we can simply require that f avoids at least one color over the set H . This is the notion of f -thin set.

Definition 1.5 (Thin set theorem) Given a coloring $f : [\omega]^n \rightarrow k$ (resp. $f : [\omega]^n \rightarrow \omega$), an infinite set H is *thin* for f if $|f([H]^n)| \leq k - 1$ (resp. $f([H]^n) \neq \omega$). For every $n \geq 1$ and $k \geq 2$, TS_k^n is the statement “Every coloring $f : [\omega]^n \rightarrow k$ has a thin set” and TS^n is the statement “Every coloring $f : [\omega]^n \rightarrow \omega$ has a thin set”. STS_k^n is the restriction of TS_k^n to stable colorings. TS is the statement $(\forall n) \text{TS}^n$.

The reverse mathematical analysis of the thin set theorem started with Friedman [14, 15]. It has been studied by Cholak et al. [5], Wang [48] and the author [34, 35] among others. Dorais et al. [9] proved that TS_k^1 is not uniformly reducible to TS_ℓ^1 whenever $\ell > k$. Hirschfeldt and Jockusch [18] extended the result to colorings over arbitrary tuples. We generalize the previous theorems by proving that $\text{TS}_k^n \not\leq_c \text{TS}_\ell^n$ whenever $\ell > k \geq 2$ and $n \geq 2$. In the case of colorings of singletons, we prove that $\text{TS}_k^1 \not\leq_{sc} \text{TS}_\ell^1$ whenever $\ell > k \geq 2$.

The free set theorem is a strengthening of the thin set theorem in which every member of a free set is a witness of thinness of the same set. Indeed, if H is an infinite f -free set for some function f , for every $a \in H$, $H \setminus \{a\}$ is f -thin with witness color a . See Theorem 3.2 in [5] for a formal version of this claim.

Definition 1.6 (Free set theorem) Given a coloring $f : [\omega]^n \rightarrow \omega$, an infinite set H is *free* for f if for every $\sigma \in [H]^n$, $f(\sigma) \in H \rightarrow f(\sigma) \in \sigma$. For every $n \geq 1$, FS^n is the statement “Every coloring $f : [\omega]^n \rightarrow \omega$ has a free set”. SFS^n is the restriction of FS^n to stable colorings. FS is the statement $(\forall n) \text{FS}^n$.

Cholak et al. [5] proved that $\text{RCA}_0 \vdash \text{RT}_2^n \rightarrow \text{FS}^n \rightarrow \text{TS}^n$ for every $n \geq 2$. Wang [48] proved that FS (hence TS) does not imply ACA_0 over ω -models. The author [34] proved that FS does not imply WKL_0 (and in fact weak König’s lemma) over RCA_0 .

Cholak et al. [5] and Montalbán [33] asked whether any of TS^2 , FS^2 , $\text{FS}^2 + \text{COH}$ and $\text{FS}^2 + \text{WKL}_0$ imply RT_2^2 over RCA_0 . Hirschfeldt [17] asked whether $\text{FS}^2 + \text{WKL}_0$ implies any of SRT_2^2 , the ascending descending sequence (ADS) and the chain antichain principle (CAC). We answer all these questions negatively by proving that for every $k \geq 2$, the conjunction of COH , WKL_0 , the Erdős-Moser theorem (EM), TS_{k+1}^2 , FS and TS implies neither STS_k^2 nor the stable ascending descending sequence (SADS) over RCA_0 .

1.6. Organization of the paper

In section 2, we establish an instance-wise correspondence between cohesive sets and degrees whose jump computes a member of a $\Pi_1^{0,\emptyset'}$ class. We take advantage of this correspondence to study how typical sets are useful to compute unsolvable instances of cohesiveness, and extend this analysis to Ramsey-type statements. In section 3.1, we reprove Dzhafarov’s result that cohesiveness is not strongly computably reducible to $\text{D}_{<\infty}^2$ with a more combinatorial proof using hyperimmunity. In section 3.2, we refine the forcing of the previous section to separate Ramsey’s theorem over computable reducibility according to the number of colors. In section 4.1, we separate variants of the thin set theorem for singletons over strong computable reducibility according to the number of colors using preservation of non-c.e. definitions. Finally, we separate the thin set theorem for pairs from Ramsey’s theorem for pairs over RCA_0 in section 4.2, and extend this separation to the full thin set theorem in section 4.3 and the full free set theorem in section 4.4.

1.7. Notation

String, sequence. Fix an integer $k \in \omega$. A *string* (over k) is an ordered tuple of integers a_0, \dots, a_{n-1} (such that $a_i < k$ for every $i < n$). The empty string is written ε . A *sequence* (over k) is an infinite listing of integers a_0, a_1, \dots (such that $a_i < k$ for every $i \in \omega$). Given $s \in \omega$, k^s is the set of strings of length s over k and $k^{<s}$ is the set of strings of length $< s$ over k .

Similarly, $k^{<\omega}$ is the set of finite strings over k and k^ω is the set of sequences (i.e. infinite strings) over k . If σ is a string, then $|\sigma|$ denotes its length. Given two strings $\sigma, \tau \in k^{<\omega}$, σ is a *prefix* of τ (written $\sigma \preceq \tau$) if there exists a string $\rho \in k^{<\omega}$ such that $\sigma\rho = \tau$. Given a sequence X , we write $\sigma \prec X$ if $\sigma = X \upharpoonright n$ for some $n \in \omega$, where $X \upharpoonright n$ denotes the restriction of X to its first n elements. A *binary string* (resp. *real*) is a *string* (resp. *sequence*) over 2 . We may identify a real with a set of integers by considering that the real is its characteristic function. Accordingly, we identify a string $\sigma \in 2^{<\omega}$ with the set $\text{set}(\sigma) = \{n < |\sigma| : \sigma(n) = 1\}$. Therefore $n \in \sigma$ means $n \in \text{set}(\sigma)$ and given a set $A \subseteq \omega$, we denote by $\sigma \cap A$ the string $\tau \in 2^{|\sigma|}$ such that $\tau(n) = 1$ if and only if $\sigma(n) = 1$ and $n \in A$. We also write $\sigma \subseteq A$ for $\text{set}(\sigma) \subseteq A$. Given a real $X \in 2^\omega$ and a string σ , we denote by X/σ the real obtained by replacing the $|\sigma|$ first bits of X by σ .

Tree, path. A tree $T \subseteq \omega^{<\omega}$ is a set downward closed under the prefix relation. The tree T is *finitely branching* if every node $\sigma \in T$ has finitely many immediate successors. A *binary tree* is a tree $T \subseteq 2^{<\omega}$. A set $P \subseteq \omega$ is a *path* through T if for every $\sigma \prec P$, $\sigma \in T$. A string $\sigma \in k^{<\omega}$ is a *stem* of a tree T if every $\tau \in T$ is comparable with σ . Given a tree T and a string $\sigma \in T$, we denote by $T^{[\sigma]}$ the subtree $\{\tau \in T : \tau \preceq \sigma \vee \tau \succeq \sigma\}$. We write $P \gg X$ to say that P is of PA degree relative to X .

Classes. Given a finite string $\sigma \in \omega^{<\omega}$, $[\sigma]$ is the set of sequences extending σ . Whenever it is clear from the context that we are working with binary strings, $[\sigma]$ denotes the set of *reals* extending σ . A $\Pi_1^{0,X}$ *class* is the collection of paths through an X -computable tree. The complement of a $\Pi_1^{0,X}$ class is a $\Sigma_1^{0,X}$ *class*. A k -*enum* of a class $\mathcal{C} \subseteq 2^\omega$ is a uniform sequence of finite sets of strings D_0, D_1, \dots such that D_s is a set of at most k binary strings of length s such that one of those is a prefix of a member of \mathcal{C} .

Sets, partitions. Given two sets A and B , we denote by $A < B$ the formula $(\forall x \in A)(\forall y \in B)[x < y]$ and by $A \subseteq^* B$ the formula $(\exists b)(\forall x \in A)[x \notin B \rightarrow x < b]$, meaning that A is contained in B except for at most finitely many elements. Given a set X and some integer k , a k -*cover* of X is a k -uple A_0, \dots, A_{k-1} such that $A_0 \cup \dots \cup A_{k-1} = X$. We may simply say k -*cover* when the set X is unambiguous. A k -*partition* is a k -cover whose sets are pairwise disjoint.

2. THE DEGREES OF UNSOLVABILITY OF COHESIVENESS

In this section, we study the degree of unsolvability of \vec{R} -cohesive sets according the degree of unsolvability of the sequence \vec{R} itself. Then we take advantage of this analysis to answer various questions about which theorems in reverse mathematics can solve a computably unsolvable instance of cohesiveness.

2.1. Cohesiveness and $\Pi_1^{0,\emptyset'}$ classes

Jockusch and Stephan characterized the p-cohesive degrees as those whose jump is of degree PA relative to \emptyset' . We clarify the situation by establishing an instance-wise correspondence between the degrees of the sets cohesive for a sequence, and the degrees whose jump computes a member of a non-empty $\Pi_1^{0,\emptyset'}$ class.

Definition 2.1 Let R_0, R_1, \dots be a uniformly computable sequence of sets. For every $\sigma \in 2^{<\omega}$, we define R_σ inductively as follows. First, $R_\varepsilon = \omega$ and then, if R_σ has already been defined for some string σ of length s , let $R_{\sigma 0} = R_\sigma \cap \overline{R_s}$ and $R_{\sigma 1} = R_\sigma \cap R_s$. For example, $R_{0110} = \overline{R_0} \cap R_1 \cap R_2 \cap \overline{R_3}$. Let $\mathcal{C}(\vec{R})$ be the $\Pi_1^{0,\emptyset'}$ class of binary sequences P such that for every $\sigma \prec P$, the set R_σ is infinite.

Our first lemma shows that the degrees of \vec{R} -cohesive sets can be characterized by their jumps. This lemma reveals in particular that low sets fail to solve unsolvable instances of cohesiveness.

Lemma 2.2 For every uniformly computable sequence of sets R_0, R_1, \dots , a set computes an \vec{R} -cohesive set if and only if its jump computes a member of $\mathcal{C}(\vec{R})$.

Proof. Fix an \vec{R} -cohesive set C . Let $P = \bigcup \{\sigma \in 2^{<\omega} : C \subseteq^* R_\sigma\}$. The sequence P is infinite and C' -computable as there exists exactly one string σ of each length such that $C \subseteq^* R_\sigma$. In particular, for every $\sigma \prec P$, R_σ is infinite, so P is a member of $\mathcal{C}(\vec{R})$.

Conversely, let X be a set whose jump computes a member P of $\mathcal{C}(\vec{R})$. By Schoenfield's limit lemma [41], there exists an X -computable function $f(\cdot, \cdot)$ such that for each x , $\lim_s f(x, s) = P(x)$. Define an \vec{R} -cohesive set $C = \bigcup_s C_s$ X -computably by stages $C_0 = \emptyset \subsetneq C_1 \subsetneq \dots$ as follows. At stage s , search for some string σ of length s and some integer $n \in R_\sigma$ greater than s such that $f(x, n) = \sigma(x)$ for each $x < |\sigma|$. We claim that such σ and n must exist, as there exists a threshold n_0 such that for every $n > n_0$, $f(x, n) = P(x)$ for each $x < s$. Let $\sigma \prec P$ be of length s . By definition of P , R_σ is infinite, so there must exist some $n \in R_\sigma$ which is greater than n_0 and s . Set $C_{s+1} = C_s \cup \{n\}$ and go to the next stage. We now check that $C = \bigcup_s C_s$ is \vec{R} -cohesive. For every $x \in \omega$, there exists a threshold n_1 such that for every $n > n_1$, $f(x, n) = P(x)$. By construction, for every element $n \in C \setminus C_{n_1}$, $n \in R_\sigma$ for some string σ such that $\sigma(x) = P(x)$. Therefore $C \subseteq^* R_x$ or $C \subseteq^* \bar{R}_x$. \square

Jockusch and Stephan [24] showed the existence of a uniformly computable sequence of sets R_0, R_1, \dots having no low \vec{R} -cohesive set. We prove that it suffices to consider any sequence \vec{R} with no computable \vec{R} -cohesive set to obtain this property.

Corollary 2.3 A uniformly computable sequence of sets R_0, R_1, \dots has a low \vec{R} -cohesive set if and only if it has a computable \vec{R} -cohesive set.

Proof. Let X be a low \vec{R} -cohesive set. By Lemma 2.2, the jump of X (hence \emptyset') computes a member of $\mathcal{C}(\vec{R})$. By a second application of Lemma 2.2, the existence of a computable \vec{R} -cohesive set follows. \square

One may naturally wonder about the shape of the $\Pi_1^{0, \emptyset'}$ classes $\mathcal{C}(\vec{R})$ for uniformly computable sequences R_0, R_1, \dots . We show through the following lemma that $\mathcal{C}(\vec{R})$ can be any $\Pi_1^{0, \emptyset'}$ class. Together with Lemma 2.2, it establishes an instance-wise correspondence between cohesive sets and $\Pi_1^{0, \emptyset'}$ classes.

Lemma 2.4 For every non-empty $\Pi_1^{0, \emptyset'}$ class $\mathcal{D} \subseteq 2^\omega$, there exists a uniformly computable sequence of sets R_0, R_1, \dots such that $\mathcal{C}(\vec{R}) = \mathcal{D}$.

Proof. By Schoenfield's limit lemma [41], there exists a computable function $g : 2^{<\omega} \times \omega \rightarrow 2$ whose limit exists and such that \mathcal{D} is the collection of X such that for every $\sigma \prec X$, $\lim_s g(\sigma, s) = 1$. We can furthermore assume that whenever $g(\sigma, s) = 1$, then for every $\tau \prec \sigma$, $g(\tau, s) = 1$, and that for every $s \in \omega$, the set $U_s = \{\sigma \in 2^s : g(\sigma, s) = 1\}$ is non-empty. We define a uniformly computable sequence of sets R_0, R_1, \dots such that $\mathcal{C}(\vec{R}) = \mathcal{D}$ by stages as follows.

As stage 0, $R_i = \emptyset$ for every $i \in \omega$. Suppose that we have already decided $R_i \upharpoonright n_s$ for every $i \in \omega$ and some $n_s \in \omega$. At stage $s+1$, we will add elements to R_0, \dots, R_s so that for each string σ of length $s+1$, $R_\sigma \upharpoonright [n_s, n_s+p] \neq \emptyset$ if and only if $\sigma \in U_{s+1}$. To do so, consider the set $U_{s+1} = \{\sigma_0, \dots, \sigma_p\}$ defined above and add $\{n_s + i : \sigma_i(j) = 1, i \leq p\}$ to R_j for each $j \leq s$. Set $n_{s+1} = n_s + p + 1$ and go to the next stage.

We claim that R_σ is infinite if and only if $\sigma \prec X$ for some $X \in \mathcal{D}$. Assume that R_σ is infinite. By construction, there are infinitely many s such that $R_\sigma \upharpoonright [n_s, n_s+p] \neq \emptyset$. So there are infinitely many stages s such that $\tau \in U_s$ ($g(\tau, s) = 1$) for some $\tau \succeq \sigma$. By assumption on g , there are infinitely many $\tau \succeq \sigma$ such that $g(\tau, s) = 1$ for infinitely many s . Therefore, by compactness, there exists some $X \in \mathcal{D}$ such that $\sigma \prec X$. Conversely, if $\sigma \prec X$ for some $X \in \mathcal{D}$, then there are infinitely many stages s such that $\tau \in U_s$ for some $\tau \succeq \sigma$. At each of these stages, $R_\sigma \upharpoonright [n_s, n_s+p] \supseteq R_\tau \upharpoonright [n_s, n_s+p] \neq \emptyset$. Therefore R_σ is infinite. \square

Jockusch et al. proved in [22] that for every $\Pi_1^{0, \emptyset'}$ class $\mathcal{C} \subseteq 2^\omega$, there exists a Π_1^0 class $\mathcal{D} \subseteq \omega^\omega$ such that $\deg(\mathcal{C}) = \deg(\mathcal{D})$, where $\deg(\mathcal{C})$ is the class of degrees of members of \mathcal{C} . For the reader

who is familiar with Weihrauch degrees, what we actually prove here is that König's lemma is the jump of the cohesiveness principle under Weihrauch reducibility. Bienvenu [personal communication] suggested the use of Simpson's Embedding Lemma [42, Lemma 3.3] to prove the reducibility of some unsolvable instances of cohesiveness to various statements.

Lemma 2.5 (Bienvenu) For every $\Sigma_3^{0,\emptyset'}$ class $\mathcal{E} \subseteq \omega^\omega$ with no \emptyset' -computable member, there exists a uniformly computable sequence of sets R_0, R_1, \dots with no computable \vec{R} -cohesive set but such that every member of \mathcal{E} computes an \vec{R} -cohesive set.

Proof. By a relativization of Lemma 3.3 in [42], there exists a $\Pi_1^{0,\emptyset'}$ class \mathcal{D} whose degrees (relative to \emptyset') are exactly $\text{deg}(\mathcal{E}) \cup PA[\emptyset']$. Therefore \mathcal{D} has no \emptyset' -computable member and every member of \mathcal{E} \emptyset' -computes a member of \mathcal{D} . By Lemma 2.4, there exists a uniformly computable sequence of sets R_0, R_1, \dots such that $\mathcal{C}(\vec{R}) = \mathcal{D}$. By Lemma 2.2, there exists no computable \vec{R} -cohesive set, but every member of \mathcal{D} (and in particular every member of \mathcal{E}) computes an \vec{R} -cohesive set. \square

2.2. How genericity helps solving cohesiveness

We now take advantage of the analysis of the previous section to deduce optimal bounds on how much genericity is needed to avoid solving an unsolvable instance of COH.

Definition 2.6 (Genericity) Fix a set of strings $S \subseteq 2^{<\omega}$. The set S is *dense* if every string has an extension in S . A real G *meets* S if it has some initial segment in S . A real G *avoids* S if it has an initial segment with no extension in S . Given an integer $n \in \omega$, a real is *n-generic* if it meets or avoids each Σ_n^0 set of strings. A real is *weakly n-generic* if it meets each Σ_n^0 dense set of strings.

By Friedberg's jump inversion theorem [13], there exists a 1-generic which is of high degree, and therefore computes a cohesive set for every uniformly computable sequence of sets. Wang [46] proved that whenever a uniformly computable sequence of sets R_0, R_1, \dots has no computable \vec{R} -cohesive sets, no weakly 3-generic computes an \vec{R} -cohesive set. He asked whether there exists a 2-generic computing an \vec{R} -cohesive set. We prove the optimality of Wang's bound by showing the existence of an unsolvable instance of COH which is solvable by a 2-generic real.

Lemma 2.7 There exists a 2-generic real G together with a uniformly computable sequence of sets R_0, R_1, \dots with no computable \vec{R} -cohesive set such that G computes an \vec{R} -cohesive set.

Proof. Fix any Δ_3^0 2-generic real G and consider the singleton $\mathcal{E} = \{G\}$. As no 2-generic is Δ_3^0 , the class \mathcal{E} has no \emptyset' -computable member. By Lemma 2.5, there exists a uniformly computable sequence of sets R_0, R_1, \dots with no computable \vec{R} -cohesive set, such that G computes an \vec{R} -cohesive set. \square

However, if we slightly increase the unsolvability of the sequence of sets, no 2-generic real helps computing a set cohesive for the sequence. Recall that a *1-enum* of a class $\mathcal{C} \subseteq 2^{<\omega}$ is a sequence of strings $\sigma_0, \sigma_1, \dots$ such that $|\sigma_s| = s$ and $[\sigma_s] \cap \mathcal{C} \neq \emptyset$ for each $s \in \omega$. The notion has been extensively studied in [34].

Theorem 2.8 For any uniformly computable sequence of sets R_0, R_1, \dots such that $\mathcal{C}(\vec{R})$ has no \emptyset' -computable 1-enum, no 2-generic real computes an \vec{R} -cohesive set.

Proof. By Jockusch [21], every n -generic set is GL_n and in particular, every 2-generic is GL_1 . Therefore, by Lemma 2.2, a 2-generic set G computes an \vec{R} -cohesive set if and only if there exists some functional Γ such that $\Gamma^{G \oplus \emptyset'}$ is a member of $\mathcal{C}(\vec{R})$. Fix a functional Γ such that $\Gamma^{G \oplus \emptyset'}$ is total for some 2-generic set G , and define the following $\Sigma_1^{0,\emptyset'}$ set:

$$W_{bad} = \{\sigma \in 2^{<\omega} : [\Gamma^{\sigma \oplus \emptyset'}] \cap \mathcal{C}(\vec{R}) = \emptyset\}$$

We claim G meets W_{bad} . Suppose for contradiction that G avoids W_{bad} . By 2-genericity of G , there exists a string $\sigma \prec G$ with no extension in W_{bad} . We show that there exists a \emptyset' -effective procedure which computes a 1-enum of $\mathcal{C}(\vec{R})$, contradicting our hypothesis.

On input n , \emptyset' -effectively search for a $\tau_n \succeq \sigma$ such that $\Gamma^{\tau_n \oplus \emptyset'} \upharpoonright n$ is defined. Such τ_n exists as $\sigma \prec G$ and $\Gamma^{G \oplus \emptyset'}$ is total. As $\tau_n \notin W_{bad}$, $[\Gamma^{\tau_n \oplus \emptyset'}] \cap \mathcal{C}(\vec{R}) \neq \emptyset$ and therefore $(\tau_n : n \in \omega)$ is a \emptyset' -computable 1-enum of $\mathcal{C}(\vec{R})$. \square

Note that if we assume that G is weakly 3-generic and therefore avoids the set $W_{bad} \cup W_{partial}$ where

$$W_{partial} = \{\sigma \in 2^{<\omega} : (\forall \tau \succeq \sigma) |\Gamma^{\tau \oplus \emptyset'}| < |\sigma|\}$$

then we can furthermore impose that $\tau_{n+1} \succeq \tau_n$ and \emptyset' -compute a member of $\mathcal{C}(\vec{R})$. This suffices to reprove that no weakly 3-generic helps solving an unsolvable instance of COH.

We now prove a theorem inspired by the proof of domination closure of p-cohesive degrees by Jockusch and Stephan [24].

Theorem 2.9 For any uniformly computable sequence of sets R_0, R_1, \dots such that $\mathcal{C}(\vec{R})$ has no \emptyset' -computable 1-enum, every \vec{R} -cohesive set is of hyperimmune degree.

Proof. Suppose for the contradiction that there exists some \vec{R} -cohesive set $C = \{a_0 < a_1 < \dots\}$ and a computable set $B = \{b_0 < b_1 < \dots\}$ such that $(\forall i)(a_i < b_i)$. For each $n \in \omega$, let $B_n = \{n, n+1, \dots, b_n\}$. Note that $a_n \in B_n$ for every n , and therefore for every length s , there exists a string σ_s of length s such that $(\exists b)(\forall n > b) R_{\sigma_s} \cap B_n \neq \emptyset$. Let $\sigma_0, \sigma_1, \dots$ be the \emptyset' -computable sequence of such strings. We claim that this sequence is a 1-enum of $\mathcal{C}(\vec{R})$, therefore contradicting our hypothesis. Indeed, as $(\exists b)(\forall n > b) R_{\sigma_s} \cap B_n \neq \emptyset$, the set R_{σ_s} is infinite and therefore $\mathcal{C}(\vec{R}) \cap [\sigma_s] \neq \emptyset$. \square

Of course, there exists some uniformly computable sequence of sets R_0, R_1, \dots with no computable \vec{R} -cohesive set but with an \vec{R} -cohesive set of hyperimmune-free degree. Simply apply Lemma 2.5 with $\mathcal{E} = \{X\}$ where X is a Δ_3^0 set of hyperimmune-free degree. Such a set is known to exist by Miller and Martin [32]. The class \mathcal{E} has no \emptyset' -computable member as every Δ_2^0 set is hyperimmune.

2.3. How randomness helps solving cohesiveness

We now explore the interrelations between cohesiveness and the measure-theoretic paradigm of typicality, namely, algorithmic randomness.

Definition 2.10 (Randomness) A Σ_n^0 (Martin-Löf) test is a sequence U_0, U_1, \dots of uniformly Σ_n^0 classes such that $\mu(U_i) \leq 2^{-i}$ for every $i \in \omega$. A real Z is n -random if for every Σ_n^0 test U_0, U_1, \dots , $Z \notin \bigcap_i U_i$. A real Z is weakly n -random if it is in every Σ_n^0 class of measure 1.

We shall say *Martin-Löf random* for *1-random*. Wang [46] proved that whenever a uniformly computable sequence of sets R_0, R_1, \dots has no computable \vec{R} -cohesive sets, there exists a Martin-Löf random real computing no \vec{R} -cohesive set. Thanks to Corollary 2.3, we know that it suffices to take any low Martin-Löf random real to obtain this property. Wang asked whether we can always ensure the existence of a 3-random real computing an \vec{R} -cohesive set whenever the instance is unsolvable. The next two lemmas answer this question by proving that it depends on the considered sequence of sets \vec{R} .

Lemma 2.11 There exists a uniformly computable sequence of sets R_0, R_1, \dots with no computable \vec{R} -cohesive set, but such that every 2-random real computes an \vec{R} -cohesive set.

Proof. Let \mathcal{D} be a $\Pi_1^{0, \emptyset'}$ class of positive measure with no \emptyset' -computable member. By Lemma 2.4, there exists a uniformly computable sequence of sets R_0, R_1, \dots such that $\mathcal{C}(\vec{R}) = \mathcal{D}$. By

Kautz [25, 26], every 2-random real is, up to prefix, a member of $\mathcal{C}(\vec{R})$. Therefore, by Lemma 2.2, every 2-random real computes an \vec{R} -cohesive set. \square

Lemma 2.12 For every $n \geq 3$, no (weakly) n -random real computes a p-cohesive set.

Proof. Jockusch and Stephan [24] proved that degrees of p-cohesive sets are those whose jump is PA relative to \emptyset' . By a relativization of Stephan [44], every 2-random real whose jump is of PA degree relative to \emptyset' is high. By Kautz [25], no weakly 3-random real is high. For every $n \geq 3$, every (weakly) n -random real is a weakly 3-random real. \square

Avigad et al. [1] introduced the principle n -WWKL stating that every Δ_n^0 tree of positive measure has a path. In particular, 1-WWKL is WWKL₀. Thanks to Lemma 2.12, for every $n \in \omega$, one can apply the usual constructions to build an ω -model of n -WWKL which does not contain any p-cohesive set and therefore is not a model of COH. Pick any n -random Z which does not compute any p-cohesive set and consider it as an infinite join $Z_0 \oplus Z_1 \oplus \dots$. By Van Lambalgen's theorem [45], the ω -structure whose second-order part is the Turing ideal $\{X : (\exists i)X \leq_T Z_0 \oplus \dots \oplus Z_i\}$ is a model of n -WWKL. Moreover it does not contain a p-cohesive set.

2.4. How Ramsey-type theorems help solving cohesiveness

In his paper separating Ramsey's theorem for pairs from weak König's lemma, Liu [30] proved that every (non-necessarily effective) set A has an infinite subset of either it or its complement which is not of PA degree. The absence of effectiveness conditions on A shows the combinatorial nature of the weakness of the infinite pigeonhole principle. On the other hand, the author [34] showed that this weakness depends on the choice of the instance of WKL₀, by constructing a computable tree with no computable path together with a Δ_2^0 set A such that every infinite subset of either A or \bar{A} computes a path through the tree. We answer a similar question for cohesiveness and study the weakness of the pigeonhole principle for typical partitions.

Lemma 2.13 There exists a Δ_3^0 (in fact low over \emptyset') set A and a uniformly computable sequence of sets R_0, R_1, \dots with no computable \vec{R} -cohesive set, such that every infinite subset of either A or \bar{A} computes an \vec{R} -cohesive set.

Proof. Fix a set A which is low over \emptyset' and bi-immune relative to \emptyset' . The set of the infinite, increasing sequences which form a subset of either A or \bar{A} is $\Pi_1^{0,A}$, hence $\Pi_2^{0,\emptyset'}$ in the Baire space:

$$\mathcal{E} = \{X \in \omega^\omega : (\forall s)[X(s) < X(s+1)] \wedge [(\forall s)(X(s) \in A) \vee (\forall s)(X(s) \in \bar{A})]\}$$

Moreover, \mathcal{E} has no \emptyset' -computable member by bi-immunity relative to \emptyset' of A . Apply Lemma 2.5 to complete the proof. \square

In a previous section, we constructed a uniformly computable sequence of sets R_0, R_1, \dots with no computable \vec{R} -cohesive set such that every 2-random real computes an \vec{R} -cohesive set. The following lemma strengthens this result by constructing an unsolvable instance of COH solvable by every infinite subset of any 2-random real.

Definition 2.14 (Diagonal non-computability) A function $f : \omega \rightarrow \omega$ is *diagonally non-computable* relative to X if for every $e \in \omega$, $f(e) \neq \Phi_e^X(e)$.

By Kjos-Hanssen [28] and Greenberg and Miller [16], a set computes a function d.n.c. relative to $\emptyset^{(n-1)}$ if and only if it computes an infinite subset of an n -random.

Lemma 2.15 There exists a uniformly computable sequence of sets R_0, R_1, \dots with no computable \vec{R} -cohesive set, such that every function d.n.c. relative to \emptyset' computes an \vec{R} -cohesive set.

Proof. The class of functions which are d.n.c. relative to \emptyset' is $\Pi_1^{0,\emptyset'}$ in the Baire space:

$$\mathcal{E} = \left\{ f \in \omega^\omega : (\forall e)[\Phi_e^{\emptyset'}(e) \uparrow \vee f(e) \neq \Phi_e^{\emptyset'}(e)] \right\}$$

Moreover, \mathcal{E} has no \emptyset' -computable member. Apply Lemma 2.5 to complete the proof. \square

In contrast with this lemma, if we require a bit more uncomputability in the \vec{R} -cohesive sets of the sequence R_0, R_1, \dots , we can ensure the existence of a function d.n.c. relative to \emptyset' which does not compute an \vec{R} -cohesive set.

Theorem 2.16 Fix a uniformly computable sequence of sets R_0, R_1, \dots such that $\mathcal{C}(\vec{R})$ has no \emptyset' -computable 1-enum. For every set X , there exists a function f d.n.c. relative to X whose jump does not compute a 1-enum of $\mathcal{C}(\vec{R})$. In particular, f does not compute an \vec{R} -cohesive set.

The proof of Theorem 2.16 is done by a bushy tree forcing argument. See the survey from Khan and Miller [27] for terminology and definitions. Fix a set X . We will construct a GL_1 function which is d.n.c. relative to X . Our forcing conditions are tuples (σ, B) where $\sigma \in \omega^{<\omega}$ and $B \subseteq \omega^{<\omega}$ is an upward-closed set k -small above σ for some $k \in \omega$. A sequence f satisfies a condition (σ, B) if $\sigma \prec f$ and B is small above every initial segment of f . Our initial condition is (ε, B_{DNC}^X) where

$$B_{DNC}^X = \{ \sigma \in \omega^{<\omega} : (\exists e)\sigma(e) = \Phi_e^X(e) \}$$

Therefore every infinite sequence f satisfying (ε, B_{DNC}^X) is d.n.c. relative to X . Thanks to the following lemma, we can prevent $f \oplus \emptyset'$ from computing a 1-enum of $\mathcal{C}(\vec{R})$. As the constructed function f is GL_1 , $f' \leq_T f \oplus \emptyset'$ does not compute a 1-enum of $\mathcal{C}(\vec{R})$.

Lemma 2.17 For every condition $c = (\sigma, B)$ and every Turing functional Γ , there exists an extension $d = (\tau, C)$ forcing $\Gamma^{f \oplus \emptyset'}$ to be partial or such that $\Gamma^{f \oplus \emptyset'}$ is not a 1-enum of $\mathcal{C}(\vec{R})$.

Proof. Suppose that B is k -small above σ . For every $n \in \omega$, define the $\Sigma_1^{0,\emptyset'}$ set $D_n = \{ \tau \in \omega^{<\omega} : \Gamma^{\tau \oplus \emptyset'}(n) \downarrow \in 2^n \}$. Make a \emptyset' -effective search for an $n \in \omega$ such that one of the following holds:

- (a) D_n is $k2^n$ -small above σ for some $n \in \omega$
- (b) $D_{n,\rho} = \{ \tau \in \omega^{<\omega} : \Gamma^{\tau \oplus \emptyset'}(n) \downarrow = \rho \}$ is k -big above σ for some string $\rho \in 2^n$ such that $[\rho] \cap \mathcal{C}(\vec{R}) = \emptyset$.

Such n exists, as otherwise, for every $n \in \omega$, D_n is $k2^n$ -big above σ . By the smallness additivity property, $D_{n,\rho}$ is k -big above σ for some $\rho \in 2^n$. For every such string ρ , $[\rho] \cap \mathcal{C}(\vec{R}) \neq \emptyset$. Therefore we can \emptyset' -compute a 1-enum of $\mathcal{C}(\vec{R})$ by searching on each input n for some ρ of length n such that $D_{n,\rho}$ is k -big above σ .

If we are in case (a), take $d = (\tau, C \cup D_n)$ as the desired extension. The condition d forces $\Gamma^{f \oplus \emptyset'}$ to be partial. If we are in case (b), by the concatenation property, there exists an extension $\tau \in D_{n,\rho}$ such that B is still k -small above τ . The condition $d = (\tau, B)$ is an extension forcing $\Gamma^{f \oplus \emptyset'}$ not to be a 1-enum of $\mathcal{C}(\vec{R})$ as $\Gamma^{f \oplus \emptyset'}(n) = \Gamma^{\tau \oplus \emptyset'}(n) = \rho$ and $[\rho] \cap \mathcal{C}(\vec{R}) = \emptyset$. \square

Looking at the proof of the previous lemma, we can \emptyset' -decide in which case we are, and then use the knowledge of f to see which path has been chosen in the bushy tree. The construction therefore yields a GL_1 sequence.

3. RAMSEY'S THEOREM AND COMPUTABLE REDUCIBILITY

The strength of Ramsey's theorem is known to remain the same when changing the number of colors in the setting of reverse mathematics. Indeed, given some coloring $f : [\omega]^n \rightarrow k^2$, we can define another coloring $g : [\omega]^n \rightarrow k$ by merging colors together by blocks of size k . After one application of RT_k^n to the coloring g , we obtain an infinite set H over which f uses at most k different colors. Another application of RT_k^n gives an infinite f -homogeneous set. This standard proof of $\text{RCA}_0 \vdash \text{RT}_k^n \rightarrow \text{RT}_{k^2}^n$ involves two applications of RT_k^n . In this section, we show that

in the computable reducibility setting, multiple applications are really necessary to reduce RT_k^n to RT_ℓ^n whenever $k > \ell$ and $n \geq 2$.

Note that two applications of RT_2^n are sufficient to deduce RT_k^n in the case $n \geq 4$, as Jockusch [20] proved that every computable instance of RT_k^n has a Π_n^0 solution, and that for every set X , there exists an X -computable instance of RT_2^n such that every solution computes $X^{(n-2)}$.

3.1. Cohesiveness and strong reducibility

We start our analysis with partitions of integers. Of course, every computable partition has an infinite computable homogeneous set, so we need to consider non-effective partitions and strong computable reducibility. The study of RT_k^1 over strong reducibility has close connections with cohesiveness. Dzhafarov [12] proved that $\text{COH} \not\leq_{sc} \text{D}_{<\infty}^2$ by iterating the following theorem.

Theorem 3.1 (Dzhafarov [12]) For every $k \geq 2$ and $\ell < 2^k$, there is a finite sequence R_0, \dots, R_{k-1} such that for all partitions $A_0 \cup \dots \cup A_{\ell-1} = \omega$ hyperarithmetical in \vec{R} , there is an infinite subset of some A_j that computes no \vec{R} -cohesive set.

Hirschfeldt and Jockusch noticed in [18] that the proof of Theorem 3.1 can be slightly modified to obtain a proof that $\text{RT}_k^1 \not\leq_{sc} \text{RT}_\ell^1$ whenever $k > \ell \geq 2$. Montálban asked whether the hyperarithmetic effectiveness restriction can be removed from Dzhafarov's theorem. We give a positive answer, which has been proved independently by Hirschfeldt and Jockusch [18]. Moreover, we show that \vec{R} can be chosen to be low.

Given two integers $k, \ell \geq 1$, we let $\pi(k, \ell)$ denote the unique $a \geq 1$ such that $k = a \cdot \ell - b$ for some $b \in [0, \ell)$. Informally, $\pi(k, \ell)$ is the minimal number of pigeons we can ensure in at least one pigeonhole, given k pigeons and ℓ pigeonholes. In particular, $\pi(k, \ell) \geq 2$ whenever $k > \ell \geq 1$. We prove the following theorem, from which we deduce several corollaries about cohesiveness and RT_k^1 .

Theorem 3.2 Fix some $k \geq 1$ and $\ell \geq 2$, some set I and a sequence of k I -hyperimmune sets B_0, \dots, B_{k-1} . For every ℓ -partition $A_0 \cup \dots \cup A_{\ell-1} = \omega$, there exists an infinite subset H of some A_i such that $\pi(k, \ell)$ sets among the B 's are $I \oplus H$ -hyperimmune.

We will postpone the proof of Theorem 3.2 until after Corollary 3.6. Using the existence of a low k -partition $B_0 \cup \dots \cup B_{k-1} = \omega$ such that $\overline{B_j}$ is hyperimmune for every $j < k$, we deduce the following corollary.

Corollary 3.3 For every $k > \ell \geq 2$, there is a low k -partition $B_0 \cup \dots \cup B_{k-1} = \omega$ such that for all ℓ -partitions $A_0 \cup \dots \cup A_{\ell-1} = \omega$, there is an infinite subset H of some A_i and a pair $j_0 < j_1 < k$ such that every infinite H -computable set intersects both B_{j_0} and B_{j_1} .

Proof. Fix some $k > \ell \geq 2$ and a low k -partition $B_0 \cup \dots \cup B_{k-1} = \omega$ such that $\overline{B_j}$ is hyperimmune for every $j < k$. Since $k > \ell \geq 2$, $\pi(k, \ell) \geq 2$. Therefore, by Theorem 3.2, for every ℓ -partition $A_0 \cup \dots \cup A_{\ell-1} = \omega$, there is an infinite subset H of some A_i and a pair $j_0 < j_1 < k$ such that $\overline{B_{j_0}}$ and $\overline{B_{j_1}}$ are H -hyperimmune. In particular, every infinite H -computable set intersects both B_{j_0} and B_{j_1} . \square

The positive answer to Montálban's question is an immediate consequence of the previous corollary.

Corollary 3.4 For every $k \geq 2$ and $\ell < 2^k$, there is a finite sequence of low sets R_0, \dots, R_{k-1} such that for all partitions $A_0 \cup \dots \cup A_{\ell-1} = \omega$, there is an infinite subset of some A_i that computes no \vec{R} -cohesive set.

Proof. Given $k \geq 2$ and $\ell < 2^k$, fix the low 2^k -partition $(B_\sigma : \sigma \in 2^k)$ whose existence is stated by Corollary 3.3. For each $i < k$, define $R_i = \bigcup_{\sigma(i)=1} B_\sigma$. Note that by disjointness of the B 's, $\overline{R_i} = \bigcup_{\sigma(i)=0} B_\sigma$. By choice of the B 's, for all ℓ -partitions $A_0 \cup \dots \cup A_{\ell-1} = \omega$, there is an

infinite subset H of some A_j and a pair $\sigma <_{lex} \tau \in 2^k$ such that every infinite H -computable set intersects both B_σ and B_τ . Let $i < k$ be the least bit such that $\sigma(i) \neq \tau(i)$. As $\sigma <_{lex} \tau$, $\sigma(i) = 0$ and $\tau(i) = 1$. By definition of R_i , $B_\tau \subseteq R_i$ and $B_\sigma \subseteq \overline{R_i}$. Therefore no infinite H -computable set is homogeneous for R_i . In particular no infinite H -computable set is \vec{R} -cohesive. \square

The construction of the B 's is done uniformly in k . We can therefore deduce the following corollary.

Corollary 3.5 There exists a sequence of low sets R_0, R_1, \dots such that every finite partition of ω has an infinite subset in one of its parts which does not compute an \vec{R} -cohesive set.

The effectiveness of B in the statement of Corollary 3.3 enables us to deduce computable non-reducibility results about stable Ramsey's theorem for pairs, thanks to the computable equivalence between SRT_ℓ^2 and the statement D_ℓ^2 .

Corollary 3.6 For every $k > \ell \geq 2$, $\text{SRT}_k^2 \not\leq_c \text{SRT}_\ell^2$.

Proof. Fix $k > \ell \geq 2$. By Corollary 3.3, there is a Δ_2^0 k -partition $B_0 \cup \dots \cup B_{k-1} = \omega$ such that for all ℓ -partitions $A_0, \dots, A_{\ell-1}$ of ω , there is an infinite subset H of some A_i which does not compute an infinite subset of any B_j . By Cholak et al. [6], for every stable computable function $f : [\omega]^2 \rightarrow \ell$, there exists a Δ_2^0 ℓ -partition $A_0 \cup \dots \cup A_{\ell-1} = \omega$ such that every infinite subset of a part computes an infinite f -homogeneous set. Therefore, for every such function f , there exists an infinite f -homogeneous set which does not compute an infinite subset of any B_j . By Schoenfield's limit lemma [41], the Δ_2^0 approximation $g : [\omega]^2 \rightarrow k$ of the k -partition $B_0 \cup \dots \cup B_{k-1} = \omega$ is a stable computable function and every infinite g -homogeneous set with color j is an infinite subset of B_j . \square

We now turn to the proof of Theorem 3.2. We shall prove it by induction over ℓ , using a forcing construction whose forcing conditions are Mathias conditions (F, X) where X is an infinite set such that the B 's are $X \oplus I$ -hyperimmune. The case where $\ell = 1$ trivially holds since $\pi(k, 1) = k$.

3.1.1. *Forcing limitlessness.* For every ℓ -partition $A_0 \cup \dots \cup A_{\ell-1} = \omega$, we want to satisfy the following scheme of requirements to ensure that $G \cap A_i$ is infinite for each $i < \ell$.

$$\mathcal{Q}_p : (\exists n_0, \dots, n_{\ell-1} > p)[n_0 \in G \cap A_0 \wedge \dots \wedge n_{\ell-1} \in G \cap A_{\ell-1}]$$

Of course, all requirements may not be satisfiable if some part A_i is finite. Usually, a forcing argument starts with the assumption that the instance is non-trivial, that is, does not admit a solution with the desired properties (cone avoiding, low, ...). In order to force the solution to be infinite, it suffices to ensure that the reservoirs satisfy the desired properties, and therefore cannot be a solution to a non-trivial instance.

In our case, we say that an ℓ -partition $A_0 \cup \dots \cup A_{\ell-1}$ is *non-trivial* if there is no infinite set H included in the complement of one of the A 's and such that the B 's are $H \oplus I$ -hyperimmune. The following lemma states that we can focus on non-trivial partitions without loss of generality.

Lemma 3.7 For every trivial ℓ -partition $A_0 \cup \dots \cup A_{\ell-1}$, there is an infinite set $H \subseteq A_i$ for some $i < \ell$ such that $\pi(k, \ell)$ sets among the B 's are $H \oplus I$ -hyperimmune.

Proof. Let $G = \{n_0 < n_1 < \dots\}$ be an infinite subset of $\overline{A_i}$ for some $i < \ell$ such that the B 's are $G \oplus I$ -hyperimmune. Define the $(\ell - 1)$ -partition $(C_j : j \neq i)$ by setting $C_j = \{s \in \omega : n_s \in A_j\}$ for each $j \neq i$. By induction hypothesis, there exists an infinite set $H_0 \subseteq C_j$ for some $j \neq i$ such that $\pi(k, \ell - 1)$ sets among the B 's are $H_0 \oplus G \oplus I$ -hyperimmune. Note that $\pi(k, \ell - 1) \leq \pi(k, \ell)$. The set $H = \{n_s : s \in H_0\}$ is an $H_0 \oplus G$ -computable subset of A_j and $\pi(k, \ell)$ sets among the B 's are $H \oplus J$ -hyperimmune. \square

Notice that the proof of Lemma 3.7 uses the induction hypothesis with a different context, namely, $G \oplus I$ instead of I . This is where we needed to use the relativized version of the

theorem in the proof. A condition $c = (F, X)$ forces \mathcal{Q}_p if there exists some $n_0, \dots, n_{m-1} > p$ such that $n_i \in F \cap A_j$ for each $i < \ell$. Therefore, if G satisfies c and c forces \mathcal{Q}_p , then G satisfies the requirement \mathcal{Q}_p . We now prove that the set of conditions forcing \mathcal{Q}_p is dense for each $p \in \omega$. Thus, every sufficiently generic filter will induce an infinite solution.

Lemma 3.8 For every condition c and every $p \in \omega$, there is an extension forcing \mathcal{Q}_p .

Proof. Fix some $p \in \omega$. It is sufficient to show that given a condition $c = (F, X)$ and some $i < \ell$, there exists an extension $d_0 = (E, Y)$ and some integer $n_i > p$ such that $n_i \in E \cap A_i$. By iterating the process for each $i < \ell$, we obtain the desired extension d . By definition of non-triviality, A_i is co-immune in X and therefore $X \cap A_i$ is infinite. Take any $n_i \in X \cap A_i \cap (p, +\infty)$. The condition $d_0 = (F \cup \{n_i\}, X \setminus [0, n_i])$ is the desired extension. \square

3.1.2. *Forcing non-homogeneity.* The second scheme of requirements aims at ensuring that for some $i < \ell$, at least $\pi(k, \ell)$ sets among the B 's are $(G \cap A_i) \oplus I$ -hyperimmune. The requirements are of the following form for each $j < k$ and each tuple of indices $\vec{e} = e_0, \dots, e_{\ell-1}$.

$$\mathcal{R}_{\vec{e}, j} : \mathcal{R}_{e_0}^{A_0, B_j} \vee \dots \vee \mathcal{R}_{e_{\ell-1}}^{A_{\ell-1}, B_j}$$

where $\mathcal{R}_e^{A, B}$ is the statement “ $\Phi_e^{(G \cap A) \oplus I}$ does not dominate p_B ”.

We claim that if all the requirements are satisfied, then $(G \cap A_i)$ has the desired property for some $i < \ell$. Indeed, if for some fixed $j < k$, all the requirements $\mathcal{R}_{\vec{e}, j}$ are satisfied, then by the usual pairing argument, there is some $i < \ell$ such that B_j is $(G \cap A_i) \oplus I$ -hyperimmune. So if all the requirements are satisfied, then by the pigeonhole principle, there is some $i < \ell$ such that $\pi(k, \ell)$ sets among the B 's are $(G \cap A_i) \oplus I$ -hyperimmune.

A condition forces $\mathcal{R}_{\vec{e}, j}$ if every set G satisfying this condition also satisfies the requirement $\mathcal{R}_{\vec{e}}$. The following lemma is the core of the forcing argument.

Lemma 3.9 For every condition $c = (F, X)$, every $j < k$ and every tuple of Turing indices \vec{e} , there exists an extension $d = (E, Y)$ forcing $\Phi_{e_i}^{(G \cap A_i) \oplus I}$ not to dominate p_{B_j} for some $i < \ell$.

Proof. Let f be the partial $X \oplus I$ -computable function which on input x , searches for a finite set of integers U such that for every ℓ -partition $Z_0 \cup \dots \cup Z_{\ell-1} = X$, there is some $i < \ell$ and some set $E \subseteq Z_i$ such that $\Phi_{e_i}^{((F \cap A_i) \cup E) \oplus I}(x) \downarrow \in U$. If such a set U is found, then $f(x) = \max(U) + 1$, otherwise $f(x) \uparrow$. We have two cases.

- Case 1: The function f is total. By $X \oplus I$ -hyperimmunity of B_j , $f(x) \leq p_{B_j}(x)$ for some x . Let U be the finite set witnessing $f(x) \downarrow$. Letting $Z_i = X \cap A_i$ for each $i < \ell$, there is some i and some finite set $E \subseteq X \cap A_i$ such that $\Phi_{e_i}^{((F \cap A_i) \cup E) \oplus I}(x) \downarrow \in U$. The condition $d = (F \cup E, X \setminus [0, \max(E)])$ is an extension forcing $\Phi_{e_i}^{(G \cap A_i) \oplus I}(x) < f(x) \leq p_{B_j}(x)$.
- Case 2: There is some x such that $f(x) \uparrow$. By compactness, the $\Pi_1^{0, X \oplus I}$ class \mathcal{C} of sets $Z_0 \oplus \dots \oplus Z_{\ell-1}$ such that $Z_0 \cup \dots \cup Z_{\ell-1} = X$ and for every $i < \ell$ and every set $E \subseteq Z_i$, $\Phi_{e_i}^{((F \cap A_i) \cup E) \oplus I}(x) \uparrow$ is non-empty. By the hyperimmune-free basis theorem [23], there is some ℓ -partition $Z_0 \oplus \dots \oplus Z_{\ell-1} \in \mathcal{C}$ such that all the B 's are $Z_0 \oplus \dots \oplus Z_{\ell-1} \oplus X \oplus I$ -hyperimmune. Let $i < \ell$ be such that Z_i is infinite. The condition $d = (F, Z_i)$ is an extension of c forcing $\Phi_{e_i}^{(G \cap A_i) \oplus I}(x) \uparrow$.

\square

3.1.3. *Construction.* We have all necessary ingredients to build an infinite set G such that each $G \cap A_i$ is infinite, and such that $\pi(k, \ell)$ sets among the B 's are $(G \cap A_i) \oplus I$ -hyperimmune for some $i < \ell$. Thanks to Lemma 3.8 and Lemma 3.9, define an infinite descending sequence of conditions $(\varepsilon, \omega) \geq c_0 \geq \dots$ such that for each $s \in \omega$,

- (a) c_s forces \mathcal{Q}_s
- (b) c_s forces $\mathcal{R}_{\vec{e}, j}$ if $s = \langle \vec{e}, j \rangle$

where $c_s = (F_s, X_s)$. Define the set $G = \bigcup_s F_s$. By (a), $G \cap A_i$ is infinite for every $i < \ell$, and by (b), each requirement $\mathcal{R}_{\bar{e},j}$ is satisfied. This finishes the proof of Theorem 3.2.

3.2. Reducibility to Ramsey's theorem for pairs

Dorais et al. [9] asked whether $\text{RT}_k^n \not\leq_c \text{RT}_\ell^n$ for every $n \geq 2$ and $k > \ell \geq 2$. Hirschfeldt and Jockusch [18] and Rakotoniaina [38] proved that SRT_k^n is not uniformly reducible to RT_ℓ^n whenever $k > \ell$. We extend the result to computable reducibility. In the first place, we shall focus on the case $n = 2$. For this, we will take advantage of the proof of RT_ℓ^2 that applies the cohesiveness principle to obtain a stable coloring $f : [\omega]^2 \rightarrow \ell$. This coloring can itself be considered as the Δ_2^0 approximation of a \emptyset' -computable ℓ -partition of ω , and therefore as a non-effective instance of RT_ℓ^1 . Any infinite subset of one of its parts computes an infinite set homogeneous for f .

In the previous section, we have shown how to diagonalize against every ℓ -partition, simply using the fact that the complement of the parts of the instance of RT_k^n are hyperimmune. The author proved in [36] that COH instances admit solutions preserving the hyperimmunity of a predefined collection of hyperimmune sets.

Theorem 3.10 (Patey [36]) For every sequence of hyperimmune sets A_0, A_1, \dots and every uniformly computable sequence of sets R_0, R_1, \dots , there is an infinite \vec{R} -cohesive set C such that the A 's are hyperimmune relative to C .

Note that this theorem is optimal in the sense that every p-cohesive set is hyperimmune. Using Theorem 3.2, we can deduce the following theorem.

Theorem 3.11 Fix some $k \geq 1$ and $\ell \geq 2$, some set I and a sequence of k I -hyperimmune sets B_0, \dots, B_{k-1} . Every I -computable coloring $f : [\omega]^2 \rightarrow \ell$ has an infinite f -homogeneous set H such that $\pi(k, \ell)$ sets among the B 's are $I \oplus H$ -hyperimmune.

Proof. Fix k , a sequence of I -hyperimmune sets B_0, \dots, B_{k-1} for some set I . Let $f : [\omega]^2 \rightarrow \ell$ be an I -computable coloring and consider the sequence of sets R_0, R_1, \dots defined for each $x \in \omega$ by

$$R_x = \{s : f(x, s) = 1\}$$

By Theorem 3.10, there is an infinite \vec{R} -cohesive set C such that the B 's are hyperimmune relative to $C \oplus I$. Let $\tilde{f} : \omega \rightarrow \ell$ be defined by $\tilde{f}(x) = \lim_{s \in C} f(x, s)$. By Theorem 3.2 relativized to $C \oplus Z$, there is an infinite \tilde{f} -homogeneous set H such that $\pi(k, \ell)$ among the B 's are $H \oplus C \oplus Z$ -hyperimmune. In particular, $H \oplus C \oplus I$ computes an infinite f -homogeneous set. \square

Using again the existence of a low k -partition $B_0 \cup \dots \cup B_{k-1}$ such that $\overline{B_j}$ is hyperimmune for every $j < k$, we deduce the following corollary.

Corollary 3.12 For every $k > \ell \geq 2$, there is a low k -partition $B_0 \cup \dots \cup B_{k-1} = \omega$ such that each computable coloring $f : [\omega]^2 \rightarrow \ell$ has an infinite f -homogeneous set H and a pair $j_0 < j_1 < k$ such that every infinite H -computable set intersects both B_{j_0} and B_{j_1} .

Proof. Fix some $k > \ell \geq 2$ and a low k -partition $B_0 \cup \dots \cup B_{k-1} = \omega$ such that $\overline{B_j}$ is hyperimmune for every $j < k$. Since $k > \ell \geq 2$, $\pi(k, \ell) \geq 2$. Therefore, by Theorem 3.11, for every RT_ℓ^2 -instance $f : [\omega]^2 \rightarrow \ell$, there is an infinite f -homogeneous set H and a pair $j_0 < j_1 < k$ such that $\overline{B_{j_0}}$ and $\overline{B_{j_1}}$ are H -hyperimmune. In particular, every infinite H -computable set intersects both B_{j_0} and B_{j_1} . \square

Using Corollary 3.12 in a relativized form, we can extend the result to colorings over arbitrary tuples.

Theorem 3.13 For every $n \geq 2$, and every $k > \ell \geq 2$, there is a Δ_n^0 k -partition $B_0 \cup \dots \cup B_{k-1} = \omega$ such that each computable coloring $f : [\omega]^n \rightarrow \ell$ has an infinite f -homogeneous set H and a pair $j_0 < j_1 < k$ such that every infinite H -computable set intersects both B_{j_0} and B_{j_1} .

Proof. This is proved in a relativized form by induction over n . The case $n = 2$ is proved by relativizing Corollary 3.12. Now assume it holds for some n in order to prove it for $n+1$. Let $P \gg \emptyset^{(n-1)}$ be such that $P' \leq \emptyset^{(n)}$. Such a set exists by the relativized low basis theorem [23]. Applying the induction hypothesis to P , there is a $\Delta_2^{0,P}$ (hence Δ_{n+1}^0) k -partition $B_0 \cup \dots \cup B_{k-1} = \omega$ such that each P -computable coloring $f : [\omega]^n \rightarrow \ell$ has an infinite f -homogeneous set H and a pair $j_0 < j_1 < k$ such that every infinite $H \oplus P$ -computable set intersects both B_{j_0} and B_{j_1} .

Let $f : [\omega]^{n+1} \rightarrow \ell$ be a computable coloring. By Jockusch [20, Lemma 5.4], there exists an infinite set C pre-homogeneous for f such that $C \leq_T P$. (A set C is *pre-homogeneous* if any two $(n+1)$ -element subsets of C with the same first n elements are assigned the same color by f .) Let $\tilde{f} : [C]^n \rightarrow \ell$ be the P -computable coloring defined for each $\sigma \in [C]^n$ by $\tilde{f}(\sigma) = f(\sigma, a)$, where $a \in A$, $a > \max(\sigma)$. Every \tilde{f} -homogeneous set is f -homogeneous. By definition of $B_0 \cup \dots \cup B_{k-1} = \omega$, there exists an infinite \tilde{f} -homogeneous (hence f -homogeneous) set H and a pair $j_0 < j_1 < k$ such that every infinite $H \oplus P$ -computable set intersects both B_{j_0} and B_{j_1} . \square

Using the fact that $D_k^n \leq_c \text{SRT}_k^n$ for every $n, k \geq 2$, we obtain the following corollary strengthening the result of Hirschfeldt and Jockusch [18] and Rakotoniaina [38].

Corollary 3.14 For every $n \geq 2$ and every $k > \ell \geq 2$, $\text{SRT}_k^n \not\leq_c \text{RT}_\ell^n$.

This answers in particular Question 7.1 of Dorais et al. [9]. The following corollary answers positively Question 5.5.3 of Mileti [31].

Corollary 3.15 There exists two stable computable functions $f_1 : [\omega]^2 \rightarrow 2$ and $f_2 : [\omega]^2 \rightarrow 2$ such that there is no computable $g : [\omega]^2 \rightarrow 2$ with the property that every set H_g homogeneous for g computes both a set H_{f_1} homogeneous for f_1 and a set H_{f_2} homogeneous for f_2 .

Proof. By Corollary 3.12 with $\ell = 2$ and $k = 3$, there exists a Δ_2^0 3-partition $B_0 \cup B_1 \cup B_2 = \omega$ such that each computable coloring $f : [\omega]^2 \rightarrow 2$ has an infinite f -homogeneous set H and a pair $j_0 < j_1 < 3$ such that every infinite H -computable set intersects both B_{j_0} and B_{j_1} . As in Corollary 3.4, we assume that the B 's are disjoint. By Schoenfield's limit lemma [41], there exist two stable computable colorings f_1 and f_2 such that $\lim_s f_1(\cdot, s) = B_0$ and $\lim_s f_2(\cdot, s) = B_1$. If $j_0 = 0$ (resp. $j_0 = 1$) then H does not compute an infinite set homogeneous for f_1 (resp. f_2). This completes the proof. \square

4. THE WEAKNESS OF FREE SET AND THIN SET THEOREMS

The combinatorics involved in our study of the free set and thin set theorems differ deeply from our analysis of Ramsey's theorem in the previous sections. Let $\text{RT}_{\ell,d}^n$ be the statement "Every coloring $f : [\omega]^n \rightarrow \ell$ has an infinite set H on which f uses at most d colors." An analysis of the thin set theorems in the continuity of the previous sections would consist in considering the computable reductions between $\text{RT}_{\ell,d}^n$ and $\text{RT}_{k,d}^n$ whenever $\ell < k$ for a fixed parameter d . In this section, we consider the variation of the parameter d , and show that different d 's lead to different subsystems of second-order arithmetic.

4.1. Thin set theorem and strong reducibility

We start our analysis with partitions of integers like we did with Ramsey's theorem. Every computable partition has an infinite computable set avoiding one of its parts. The natural reducibility to consider is therefore strong computable reducibility. In this section, we show that $\text{TS}_k^1 \not\leq_{sc} \text{TS}_{k+1}^1$. We could have proven this separation using the notion of hyperimmunity as we did in the previous section (and this is indeed the approach chosen by the author in [36]).

However, we want to apply preservation of definitions, emulating Wang's analysis of theorems in reverse mathematics in terms of preservation of definitions.

4.1.1. *Preservation of non-c.e. definitions.* The notion of preservation of definitions was introduced by Wang in [47], in the context of a new analysis of principles in reverse mathematics in terms of their definitional strength. Wang defined a set X to preserve properly Δ_2^0 definitions if every properly Δ_2^0 set (i.e. Δ_2^0 but neither Σ_1^0 nor Π_1^0) is properly $\Delta_2^{0,X}$. He deduced several separation results, and in particular constructed an ω -model of the conjunction of COH, WKL₀, the Erdős-Moser theorem (EM), the rainbow Ramsey theorem for pairs (RRT₂²) and the Π_1^0 -genericity principle (Π_1^0 G) which is not a model of TS². His analysis has been extended by the author in [37].

Definition 4.1 (Preservation of non-c.e. definitions)

1. A set X *preserves non-c.e. definitions* of some non-c.e. sets A_0, A_1, \dots if no A_i is X -c.e.
2. A Π_1^1 statement P *admits preservation of k non-c.e. definitions* if for each C , each sequence of non- C -c.e. sets A_0, \dots, A_{k-1} and each C -computable P -instance X , there exists a solution Y to X such that $Y \oplus C$ preserves non-c.e. definitions of A_0, \dots, A_{k-1} .

Wang proved [47] that COH, EM, WKL₀, RRT₂² and Π_1^0 G admit preservation of k non-c.e. definitions for every $k \in \omega$. By a trivial adaptation of Proposition 2.4 from [47], if some statement P admits preservation of k non-c.e. definitions and some other statement Q does not, then there exists an ω -model of P which is not a model of Q . We start with a trivial lemma showing that our preservation proofs subsume Wang's analysis of cone avoidance of Ramsey-type theorems from [48].

Lemma 4.2 If some statement P admits preservation of 1 non-c.e. definition, then it admits cone avoidance.

Proof. Fix any set C , any set $A \not\leq_T C$ and any C -computable P -instance X . As $A \not\leq_T C$, either A or \bar{A} is not C -c.e. Call this set B . As P admits preservation of 1 non-c.e. definition, there exists a solution Y of X such that B is not $Y \oplus C$ -c.e. In particular A is not $Y \oplus C$ -computable. \square

4.1.2. *Negative preservation results.* The following theorem can be proven by a direct adaptation of Theorem 4.3 proven by Wang [47]. However, we provide a simpler proof.

Theorem 4.3 For every $k \geq 2$, there exists a Δ_2^0 k -partition $B_0 \cup \dots \cup B_{k-1} = \omega$ such that for each $j < k$, \bar{B}_j is non-c.e. but is H -c.e. for every infinite set $H \subseteq \bar{B}_j$.

Proof. It suffices to construct a stable computable function $f : [\omega]^2 \rightarrow k$ with no infinite computable f -thin set, and such that for each $i < k$ and each $x < y < z \in \omega$,

$$f(x, y) \neq i \wedge f(y, z) \neq i \rightarrow f(x, z) \neq i$$

We first justify that those properties are sufficient for proving our theorem. Let $B_i = \{x : \lim_s f(x, s) = i\}$. Every infinite subset of \bar{B}_i computes an infinite set thin for f with witness i , therefore no \bar{B}_i is c.e. Moreover, \bar{B}_i is H -c.e. for every infinite set $H \subseteq \bar{B}_i$ since

$$\bar{B}_i = \{x \in \omega : (\exists y \in H) f(x, y) \neq i\}$$

The construction of the function f is done by a finite injury priority argument with a movable marker procedure. We want to satisfy the following scheme of requirements for each $e \in \omega$ and $i < k$:

$$\mathcal{R}_{e,i} : W_e \text{ infinite} \rightarrow (\exists x \in W_e) \lim_s f(x, s) = i$$

The requirements are given the usual priority ordering. We proceed by stages, maintaining k sets B_0, \dots, B_{k-1} which represent the limit of the function f . At stage 0, $B_{i,0} = \emptyset$ for each $i < k$ and f is nowhere defined. Moreover, each requirement $\mathcal{R}_{e,i}$ is given a movable marker $m_{e,i}$ initialized to 0.

A strategy for $\mathcal{R}_{e,i}$ requires attention at stage $s + 1$ if $W_{e,s} \subset \overline{B_{i,s}}$ and $W_{e,s} \cap [m_{e,i}, s] \neq \emptyset$. The strategy sets $B_{i,s+1} = B_{i,s} \cup [m_{e,i}, s]$, and $B_{j,s+1} = B_{j,s} \setminus [m_{e,i}, s]$ for every $j \neq i$. Then it is declared *satisfied* until some strategy of higher priority changes its marker. Each marker $m_{e',i'}$ of strategies of lower priorities is assigned the value $s + 1$.

At stage $s + 1$, assume that $B_{0,s} \cup \dots \cup B_{k-1,s} = [0, s)$ and that f is defined for each pair over $[0, s)$. For each $x \in [0, s)$, set $f(x, s) = i$ for the unique i such that $x \in B_{i,s}$. If some strategy requires attention at stage $s + 1$, take the least one and satisfy it. If no such requirement is found, set $B_{0,s+1} = B_{0,s} \cup \{s\}$ and $B_{i,s+1} = B_{i,s}$ for $i > 0$. Then go to the next stage. This ends the construction.

Each time a strategy acts, it changes the markers of strategies of lower priority, and is declared satisfied. Once a strategy is satisfied, only a strategy of higher priority can injury it. Therefore, each strategy acts finitely often and the markers stabilize. It follows that the B 's also stabilize and that f is a stable function.

Lemma 4.4 For every $i < k$ and every $x < y < z$, $f(x, y) \neq i \wedge f(y, z) \neq i \rightarrow f(x, z) \neq i$.

Proof. Suppose that $f(x, y) \neq i$ but $f(x, z) = i$ for some $i < k$. Let $s \leq z$ be the least stage such that $f(x, t) = i$ for every $t \in [s + 1, z]$. At stage $s + 1$, some strategy $\mathcal{R}_{e,i}$ moved to B_i the whole interval $[m_{e,i}, s]$. Since $m_{e',i'} \leq m_{e,i}$ for every strategy $\mathcal{R}_{e',i'}$ of higher priority, none of the elements in $[m_{e,i}, s]$ leave B_i before stage $z + 1$. As $f(x, y) \neq i$, $y \notin [s + 1, z]$ so $y \in [m_{e,i}, s]$. Therefore $y \in B_{i,z}$ and thus $f(y, z) = i$. \square

Lemma 4.5 For every $e \in \omega$ and $i < k$, $\mathcal{R}_{e,i}$ is satisfied.

Proof. By induction over the priority order. Let s_0 be a stage after which no strategy of higher priority will ever act. By construction, $m_{e,i}$ will not change after stage s_0 . If W_e is infinite, it will eventually enumerate some element u bigger than $m_{e,i}$, and therefore $\mathcal{R}_{e,i}$ will require attention at some stage $s \geq u$. As no strategy of higher priority ever acts after stage s_0 , $\mathcal{R}_{e,i}$ will receive attention, be satisfied and never be injured. \square

Satisfying $\mathcal{R}_{e,i}$ for every $e \in \omega$ and $i < k$ guarantees that f has no computable thin set. This last claim finishes the proof of Theorem 4.3. \square

Corollary 4.6 For every $k \geq 2$, STS_k^2 does not admit preservation of k non-c.e. definitions.

4.1.3. *Strong preservation of non-c.e. definitions.* Because every computable instance of TS_k^1 having a computable solution, TS_k^1 admits preservation of k non-c.e. definitions for every k . On the other hand, we would like to say that TS_k^1 does not *combinatorially* preserve k non-c.e. definitions since Theorem 4.3 shows the existence of a non-effective instance of TS_k^1 whose solutions do not preserve k non-c.e. definitions. This combinatorial notion of preservation is called *strong preservation*.

Definition 4.7 (Strong preservation of non-c.e. definitions) A Π_2^1 statement P admits strong preservation of k non-c.e. definitions if for each set C , each sequence of non- C -c.e. sets A_0, \dots, A_{k-1} and each (arbitrary) P -instance X , there exists a solution Y to X such that $Y \oplus C$ preserves non-c.e. definitions of A_0, \dots, A_{k-1} .

We have seen through Theorem 4.3 that for every $k \geq 2$, TS_k^1 does not admit strong preservation of k non-c.e. definitions. The following theorem shows the optimality of Theorem 4.3.

Theorem 4.8 For every $k \geq 2$, TS_{k+1}^1 admits strong preservation of k non-c.e. definitions.

The proof of Theorem 4.8 follows Corollary 4.10. Putting Theorem 4.3 and Theorem 4.8 together, we obtain the desired separation over strong computable reducibility.

Corollary 4.9 For every $\ell > k \geq 2$, $\text{TS}_k^1 \not\leq_{sc} \text{TS}_\ell^1$

Using the computable equivalence between the problem of finding a infinite set thin for an Δ_2^0 ℓ -partition and STS_ℓ^2 , we deduce the following corollary.

Corollary 4.10 For every $\ell > k \geq 2$, $\text{STS}_k^2 \not\leq_c \text{STS}_\ell^2$

Proof. Fix $\ell > k \geq 2$ and consider the Δ_2^0 k -partition $B_0 \cup \dots \cup B_{k-1} = \omega$ of Theorem 4.3. By Schoenfield's limit lemma [41], there exists a stable computable function $g : [\omega]^2 \rightarrow k$ such that $B_j = \{x : \lim_s g(x, s) = j\}$ for each $j < k$. Every infinite set thin for g is thin for the B 's. Fix any stable computable function $f : [\omega]^2 \rightarrow \ell$ and let $A_i = \{x : \lim_s f(x, s) = i\}$ for each $i < m$. By Theorem 4.8, there exists an infinite set H thin for the A 's which does not compute an infinite set thin for the B 's (hence for g). As $H \oplus f$ computes an infinite set G thin for f , f has an infinite f -thin set which does not compute an infinite set thin for g . \square

The remainder of this section is devoted to the proof of Theorem 4.8. Fix some set C preserving non-c.e. definitions of some sets B_0, \dots, B_{k-1} and fix some $(k+1)$ -partition $A_0 \cup \dots \cup A_k = \omega$. We will construct a set G such that $G \cap \overline{A_i}$ is infinite for each $i \leq k$ and none of the B 's are $(G \cap \overline{A_i}) \oplus C$ -c.e. for some $i \leq k$. Our forcing conditions are Mathias conditions (σ, X) where X is an infinite set of integers such that none of the B 's are $X \oplus C$ -c.e.

4.1.4. *Forcing limitlessness.* We want to satisfy the following scheme of requirements to ensure that $G \cap \overline{A_i}$ is infinite for each $i \leq k$:

$$\mathcal{Q}_p : (\exists m_0, \dots, m_k > p)[m_0 \in G \cap \overline{A_0} \wedge \dots \wedge m_k \in G \cap \overline{A_k}]$$

We say that an $(k+1)$ -partition $A_0 \cup \dots \cup A_k = \omega$ is *non-trivial* if there exists no infinite set H homogeneous for the A 's such that none of the B 's are $H \oplus C$ -c.e. Of course, every infinite set homogeneous for the A 's is thin for the A 's, so if the partition $A_0 \cup \dots \cup A_k = \omega$ is trivial, we succeed. Therefore we will assume from now on that the partition is non-trivial. A condition $c = (\sigma, X)$ *forces* \mathcal{Q}_p if there exist some $m_0, \dots, m_k > p$ such that $m_i \in \sigma \cap \overline{A_i}$ for each $i \leq k$. Therefore, if G satisfies c and c forces \mathcal{Q}_p , then G satisfies the requirement \mathcal{Q}_p . We now prove that the set of conditions forcing \mathcal{Q}_p is dense for each $p \in \omega$. Thus, every sufficiently generic filter will induce a set G such that $G \cap \overline{A_i}$ is infinite for each $i \leq k$.

Lemma 4.11 For every condition c and every p , there is an extension forcing \mathcal{Q}_p .

Proof. Fix some $p \in \omega$. It is sufficient to show that given a condition $c = (\sigma, X)$ and some $i \leq k$, there exist an extension $d_0 = (\tau, Y)$ and some integer $m_i > p$ that $m_i \in \tau \cap \overline{A_i}$. By iterating the process for each $i \leq k$, we obtain an extension forcing \mathcal{Q}_p . Suppose for the sake of contradiction that $X \cap \overline{A_i}$ is finite. One can then X -compute an infinite set $H \subseteq A_i$, contradicting non-triviality of the A 's. Therefore, there exists an $m_i \in X \cap \overline{A_i}$ such that $m_i > \max(\sigma, p)$. The condition $d_0 = (\sigma \frown m_i, X)$ is the desired extension. \square

4.1.5. *Forcing preservation.* The second scheme of requirements consists in ensuring that the sets B_0, \dots, B_{k-1} are all non- $(G \cap \overline{A_i}) \oplus C$ -c.e. for some $i \leq k$. The requirements are of the following form for each tuple of indices $\vec{e} = (e_i : i \leq k)$:

$$\mathcal{R}_{\vec{e}} : \bigwedge_{j < k} W_{e_0}^{(G \cap \overline{A_0}) \oplus C} \neq B_j \vee \dots \vee \bigwedge_{j < k} W_{e_k}^{(G \cap \overline{A_k}) \oplus C} \neq B_j$$

A condition *forces* $\mathcal{R}_{\vec{e}}$ if every set G satisfying this condition also satisfies requirement $\mathcal{R}_{\vec{e}}$. The following lemma is the core of the forcing argument.

Lemma 4.12 For every condition $c = (\sigma, X)$, every $i_0 < i_1 \leq k$, every $j < k$ and every vector of indices \vec{e} , there exists an extension d forcing either $W_{e_{i_0}}^{(G \cap \overline{A_{i_0}}) \oplus C} \neq B_j$ or $W_{e_{i_1}}^{(G \cap \overline{A_{i_1}}) \oplus C} \neq B_j$.

Proof. Let W be the set of all $a \in \omega$ such that for every 2-cover $Z_{i_0} \cup Z_{i_1} = X$, there is some $i \in \{i_0, i_1\}$ and some set $G_i \subseteq Z_i$ such that $a \in W_{e_i}^{(G_i/(\sigma \cap \overline{A_i})) \oplus C}$. The set W is $X \oplus C$ -c.e. Therefore $W \neq B_j$. Let $a \in W \Delta B_j$. We have two cases:

- Case 1: $a \in W \setminus B_j$. By definition of W , taking in particular the sets $Z_{i_0} = X \cap \overline{A_{i_0}}$ and $Z_{i_1} = X \cap \overline{A_{i_1}}$, there is some $i \in \{i_0, i_1\}$ and some finite set $G_i \subseteq Z_i$ such that $a \in W_{e_i}^{(G_i/(\sigma \cap \overline{A_i})) \oplus C}$. The condition $d = (G_i/\sigma, X)$ is an extension forcing $W_{e_i}^{(G \cap \overline{A_i}) \oplus C} \neq B_j$.
- Case 2: $a \in B_j \setminus W$. Let \mathcal{C} be the $\Pi_1^{0, X \oplus C}$ class of sets $Z_{i_0} \oplus Z_{i_1}$ such that $Z_{i_0} \cup Z_{i_1} = X$ and for every $i \in \{i_0, i_1\}$ and every set $G_i \subseteq Z_i$ $a \notin W_{e_i}^{(G_i/(\sigma \cap \overline{A_i})) \oplus C}$. By definition of W , \mathcal{C} is non-empty. As WKL_0 admits preservation of k non-c.e. definitions, there exists some $Z_{i_0} \oplus Z_{i_1} \in \mathcal{C}$ such that none of the B 's are $Z_{i_0} \oplus Z_{i_1} \oplus X \oplus C$ -c.e. Let $i \in \{i_0, i_1\}$ be such that Z_i is infinite. The condition $d = (\sigma, Z_i)$ is an extension of c forcing $W_{e_i}^{(G \cap \overline{A_i}) \oplus C} \neq B_j$. □

As usual, the following lemma iterates Lemma 4.12 and uses the fact that $k+1 > k$ to satisfy the requirement $\mathcal{R}_{\vec{e}}$.

Lemma 4.13 For every condition c , and every indices \vec{e} , there exists an extension d forcing $\mathcal{R}_{\vec{e}}$.

Proof. Fix a condition c , and iterate applications of Lemma 4.12 to obtain an extension d such that for each $j < k$, d forces $W_{e_i}^{(G \cap \overline{A_i}) \oplus C} \neq B_j$ for k different i 's. By the pigeonhole principle, there exists some $i \leq k$ such that d forces $W_{e_i}^{(G \cap \overline{A_i}) \oplus C} \neq B_j$ for each $j < k$. Therefore, d forces $\mathcal{R}_{\vec{e}}$. □

4.1.6. *Construction.* Thanks to Lemma 4.11 and Lemma 4.13, define an infinite descending sequence of conditions $(\varepsilon, \omega) \geq c_0 \geq \dots$ such that for each $s \in \omega$,

- (a) c_s forces \mathcal{Q}_s
- (b) c_s forces $\mathcal{R}_{\vec{e}}$ if $s = \langle \vec{e} \rangle$

where $c_s = (\sigma_s, X_s)$. Let $G = \bigcup_s \sigma_s$. By (a), $G \cap \overline{A_i}$ is infinite for every $i \leq k$ and by (b), G satisfies each requirement $\mathcal{R}_{\vec{e}}$. This finishes the proof of Theorem 4.8.

4.2. Thin set theorem for pairs and reverse mathematics

There is a fundamental difference in the way we proved that $\text{RT}_k^1 \not\leq_{sc} \text{RT}_\ell^1$ and that $\text{TS}_\ell^1 \not\leq_{sc} \text{TS}_k^1$ whenever $k > \ell$. In the former case, we have built an instance I of RT_k^1 satisfying some hyperimmunity properties, and used those properties to construct a solution X to each instance of RT_ℓ^1 which does not compute a solution to I . We did not ensure that those hyperimmunity properties are preserved relative to the solution X , which prevents us from iterating the construction. As it happens, those properties are not preserved as multiple applications of RT_ℓ^1 are sufficient to compute a solution to I . In the latter case, we proved that TS_ℓ^1 has an instance whose solutions do not preserve some definitional property, whereas each instance of TS_k^1 has a solution preserving it. This preservation enables us to iterate the applications of TS_k^1 and build ω -structures whose second-order part is made of sets preserving this property. We will take advantage of those observations to obtain new separations in reverse mathematics.

In this section, we prove that TS_{k+1}^2 does not imply TS_k^2 over RCA_0 for every $k \geq 2$. In particular, we answer several questions asked by Cholak, Giusto, Hirst and Jockusch [5] and by Montálban [33] about the relation between RT_2^2 and TS^2 . Dorais et al. [9] proved that $\text{RCA}_0 \vdash \text{TS}_k^n \rightarrow \text{ACA}_0$ for $n \geq 3$ whenever k is not large enough. Therefore we cannot hope to obtain the same separation result over RCA_0 for arbitrary tuples. However, we shall see that TS_k^n is not computably reducible to TS_{k+1}^n for $n, k \geq 2$.

Theorem 4.14 For every $k \geq 2$, let Φ be the conjunction of COH , WKL_0 , RRT_2^2 , $\Pi_1^0\text{G}$, EM , TS_{k+1}^2 . Over RCA_0 , Φ implies neither STS_k^2 nor SADS .

The proof of Theorem 4.14 follows Corollary 4.20. Cholak et al. [5] and Montálban [33] asked whether TS^2 implies RT_2^2 over RCA_0 . Thanks to Theorem 4.14, we answer negatively, noticing that TS_k^2 is the statement RT_k^2 and $\text{RCA}_0 \vdash \text{TS}_k^2 \rightarrow \text{TS}^2$ for every $k \geq 2$ (see Dorais et al. [9]).

Corollary 4.15 TS^2 does not imply RT_2^2 over RCA_0 .

Using the standard trick of prehomogeneous sets, we can generalize from computable non-reducibility over pairs to arbitrary tuples.

Corollary 4.16 For every $k, n \geq 2$ there exists a Δ_n^0 k -partition $B_0 \cup \dots \cup B_{k-1} = \omega$ such that every computable coloring $f : [\omega]^n \rightarrow k + 1$ has an infinite f -thin set computing no set thin for the B 's.

Proof. This is proved in a relativized form by induction over $n \geq 2$. The case $n = 2$ is obtained by relativizing the proof of Theorem 4.14, which shows indeed the existence of a Δ_2^0 k -partition $B_0 \cup \dots \cup B_{k-1} = \omega$ such that every computable coloring $f : [\omega]^2 \rightarrow k + 1$ has an infinite f -thin set computing no set thin for the B 's. Now assume it holds for some n in order to prove it for $n + 1$. By the relativized low basis theorem [23], let $P \gg \emptyset^{(n-1)}$ be such that $P' \leq \emptyset^{(n)}$. Applying the induction hypothesis to P , there is a $\Delta_2^{0,P}$ (hence Δ_{n+1}^0) k -partition $B_0 \cup \dots \cup B_{k-1} = \omega$ such that each P -computable coloring $f : [\omega]^n \rightarrow k + 1$ has an infinite f -homogeneous set H such that $H \oplus P$ does not compute an infinite set thin for the B 's.

Let $f : [\omega]^{n+1} \rightarrow k + 1$ be a computable coloring. By Jockusch [20, Lemma 5.4], there exists an infinite set C pre-homogeneous for f such that $C \leq_T P$. Let $\tilde{f} : [C]^n \rightarrow k + 1$ be the P -computable coloring defined for each $\sigma \in [C]^n$ by $\tilde{f}(\sigma) = f(\sigma, a)$, where $a \in A$, $a > \max(\sigma)$. Every \tilde{f} -thin set is f -thin. By definition of $B_0 \cup \dots \cup B_{k-1} = \omega$, there exists an infinite \tilde{f} -thin (hence f -thin) set H such that $H \oplus P$ does not compute an infinite set thin for the B 's. \square

Corollary 4.17 For every $k, n \geq 2$, $\text{STS}_k^n \not\leq_c \text{TS}_{k+1}^n$

We proved in section 4.1 that STS_k^2 does not admit preservation of k non-c.e. definitions. Jockusch noticed (see Hirschfeldt and Shore [19]) that **SADS** does not admit preservation of 2 non-c.e. definitions. We give the proof for the sake of completeness.

Theorem 4.18 **SADS** does not admit preservation of 2 non-c.e. definitions.

Proof. Tennenbaum (see Rosenstein [39]) constructed a computable linear order of order type $\omega + \omega^*$ with no computable infinite ascending or descending sequence. Let B_0 be the ω -part and B_1 be the ω^* part of this linear order. Every infinite subset of B_0 (resp. B_1) computes an infinite ascending (resp. descending) sequence, therefore B_0 and B_1 are non-c.e. The ω part (resp. ω^* part) is c.e. in every infinite ascending (resp. descending) sequence. \square

By Schoenfield's limit lemma [41], a stable computable coloring over $(n + 1)$ -tuples can be considered as a non-effective coloring over n -tuples. This consideration establishes a bridge between preservation properties for colorings over $(n + 1)$ -tuples and strong preservation properties for colorings over n -tuples. In particular, it enables us to prove preservation results by induction over n . The following lemma has been proven by the author in its full generality in [34]. Nevertheless we reprove it in the context of preservation of non-c.e. definitions.

Lemma 4.19 For every $k, n \geq 1$ and $\ell \geq 2$, if TS_ℓ^n admits strong preservation of k non-c.e. definitions, then TS_ℓ^{n+1} admits preservation of k non-c.e. definitions.

Proof. Fix any set C , k non- C -c.e. sets A_0, \dots, A_{k-1} and any C -computable coloring $f : [\omega]^{n+1} \rightarrow \ell$. Consider the uniformly C -computable sequence of sets \vec{R} defined for each $\sigma \in [\omega]^n$ and $i < \ell$ by

$$R_{\sigma,i} = \{s \in \omega : f(\sigma, s) = i\}$$

As COH admits preservation of k non-c.e. definitions, there exists some \vec{R} -cohesive set G such that $G \oplus C$ preserves non-c.e. definitions of the A 's. The cohesive set induces a $(G \oplus C)$ -computable coloring $\tilde{f} : [\omega]^n \rightarrow \ell$ defined by:

$$(\forall \sigma \in [\omega]^n) \tilde{f}(\sigma) = \lim_{s \in G} f(\sigma, s)$$

As TS_ℓ^n admits strong preservation of k non-c.e. definitions, there exists an infinite \tilde{f} -thin set H such that $H \oplus G \oplus C$ preserves non-c.e. definitions of the A 's. $H \oplus G \oplus C$ computes an infinite f -thin set. \square

Using Theorem 4.8 together with Lemma 4.19, we deduce the following corollary.

Corollary 4.20 For every $k \geq 2$, TS_{k+1}^2 admits preservation of k non-c.e. definitions.

We are now ready to prove Theorem 4.14.

Proof of Theorem 4.14. Fix some $k \geq 2$. Wang proved in [47] that COH, WKL_0 , RRT_2^2 , $\Pi_1^0\text{G}$ and EM admit preservation of k non-c.e. definitions. By Corollary 4.20, so does TS_{k+1}^2 . By Corollary 4.6 and Theorem 4.18, neither STS_k^2 nor SADS admit preservation of k non-c.e. definitions. The theorem follows by an application of Proposition 2.4 of Wang [47]. \square

4.3. Thin set theorem for tuples and reverse mathematics

In this section, we extend the preservation of non-c.e. definitions of the thin set theorem for pairs to arbitrary tuples, using the same construction pattern as Wang [48]. We deduce that TS_ℓ^n does not imply TS_k^n over RCA_0 whenever ℓ is large enough, which is informally the strongest result we can obtain since Proposition 5.3 in Dorais et al. [9] states that $\text{RCA}_0 \vdash \text{ACA}_0 \leftrightarrow \text{TS}_k^n$ for $n \geq 3$ whenever k is not large enough.

Theorem 4.21 For every $k, n \geq 1$, TS_ℓ^n admits strong preservation of k non-c.e. definitions for sufficiently large ℓ .

The proof of Theorem 4.21 begins below in section 4.3.1. Using the fact that $\text{RCA}_0 \vdash \text{TS}_\ell^n \rightarrow \text{TS}^n$ for every $n, \ell \geq 2$, we obtain the following preservation result for TS.

Corollary 4.22 For every $k \geq 1$, TS admits strong preservation of k non-c.e. definitions.

Thanks to the existing preservations of non-c.e. definitions and Proposition 2.4 from Wang [47], we deduce the following separations over ω -models.

Corollary 4.23 For every $k \geq 2$, let Φ be the conjunction of COH, WKL_0 , RRT_2^2 , $\Pi_1^0\text{G}$, EM, TS_{k+1}^2 and TS. Over RCA_0 , Φ implies neither STS_k^2 nor SADS.

The remainder of this section is devoted to the proof of Theorem 4.21.

4.3.1. *Proof structure.* We shall follow the proof structure of strong cone avoidance used by Wang [48]. Fix some $k \geq 1$. The induction works as follows:

- (A1) In section 4.2 we proved that TS_{k+1}^1 admits strong preservation of k non-c.e. definitions. This is the base case of our induction.
- (A2) Assuming that for each $t \in (0, n)$, $\text{TS}_{d_t+1}^t$ admits strong preservation of k non-c.e. definitions, we prove that $\text{TS}_{d_{n-1}+1}^n$ admits preservation of k non-c.e. definitions. This is done by Lemma 4.19.
- (A3) Then we prove that $\text{TS}_{d_n+1}^n$ admits strong preservation of k non-c.e. definitions where

$$d_n = d_1 d_{n-1} + \sum_{0 < t < n} d_t d_{n-t}$$

Properties (A1) and (A2) are already proven. We now prove property (A3). It is again done in several steps. Fix a coloring $f : [\omega]^n \rightarrow d_n + 1$ and a set C preserving non-c.e. definitions of k sets A_0, \dots, A_{k-1} .

- (S1) First, we construct an infinite set $D \subseteq \omega$ such that $D \oplus C$ preserves non-c.e. definitions of the A 's and a sequence $(I_\sigma : 0 < |\sigma| < n)$ such that for each $t \in (0, n)$ and each $\sigma \in [\omega]^t$
- (a) I_σ is a subset of $\{0, \dots, d_n\}$ with at most d_{n-t} many elements
 - (b) $(\exists b)(\forall \tau \in [D \cap (b, +\infty)]^{n-t}) f(\sigma, \tau) \in I_\sigma$
- (S2) Second, we construct an infinite set $E \subseteq D$ such that $E \oplus C$ preserves non-c.e. definitions of the A 's and a sequence $(I_t : 0 < t < n)$ such that for each $t \in (0, n)$
- (a) I_t is a subset of $\{0, \dots, d_n\}$ of size at most $d_t d_{n-t}$
 - (b) $(\forall \sigma \in [E]^t)(\exists b)(\forall \tau \in [E \cap (b, +\infty)]^{n-t}) f(\sigma, \tau) \in I_t$
- (S3) Third, we construct a sequence $(\xi_i \in [E]^{<\omega} : i < \omega)$ such that
- (a) The set $G = \bigcup_i \xi_i$ is infinite and $G \oplus C$ preserves non-c.e. definitions of the A 's
 - (b) $|f([\xi_i]^n)| \leq d_{n-1}$ and $\max(\xi_i) < \min(\xi_{i+1})$ for each $i < \omega$
 - (c) For each $t \in (0, n)$ and $\sigma \in [\bigcup_{j < i} \xi_j]^t$, $f(\sigma, \tau) \in I_t$ for all $\tau \in [\bigcup_{j \geq i} \xi_j]^{n-t}$
- (S4) Finally, we build an infinite set $H \subseteq G$ such that $H \oplus C$ preserves non-c.e. definitions of the A 's and $|f([H]^n)| \leq d_n$.

4.3.2. *Generalized cohesiveness.* Before proving that $\text{TS}_{d_n+1}^n$ admits strong preservation of k non-c.e. definitions, we need to prove strong preservation for a generalized notion of cohesiveness already used by the author in [34]. Cohesiveness can be seen as the problem which takes as an input a coloring of pairs $f : [\omega]^2 \rightarrow \ell$ and fixes the first parameter to obtain an infinite sequence of colorings of integers $f_x : \omega \rightarrow \ell$ for each $x \in \omega$. A solution to this problem is an infinite set G which is eventually homogeneous for each coloring f_x .

Going further in this approach, we can consider that cohesiveness is a degenerate case of the problem which takes as an input a coloring of pairs $f : [\omega]^2 \rightarrow \omega$ using infinitely many colors, and fixes again the first parameter to obtain an infinite sequence of colorings of integers $f_x : \omega \rightarrow \omega$. A solution to this problem is an infinite set G such that for each color i , either eventually the color will be avoided by f_x over G , or G will be eventually homogeneous for f_x with color i .

We can generalize the notion to colorings over tuples $f : [\omega]^n \rightarrow \omega$, seeing f as an infinite sequence of colorings over t -tuples $f_\sigma : [\omega]^t \rightarrow \omega$ for each $\sigma \in [\omega]^{n-t}$. We will create a set G such that at most d_t colors will appear for arbitrarily large pairs over G for each function f_σ . This set will be constructed by applying $\text{TS}_{d_t+1}^t$ to f_σ for each σ .

We do not need Theorem 4.24 in its full generality to complete our step (S1). However, it will be useful in a later section for proving that the free set theorem admits preservation of k non-c.e. definitions.

Theorem 4.24 Fix a coloring $f : [\omega]^n \rightarrow \omega$, some $t \leq n$ and suppose that $\text{TS}_{d_s+1}^s$ admits strong preservation of k non-c.e. definitions for each $s \in (0, t]$. For every set C preserving non-c.e. definitions of some sets A_0, \dots, A_{k-1} , there exists an infinite set G such that $G \oplus C$ preserves non-c.e. definitions of the A 's and for every $\sigma \in [\omega]^{<\omega}$ such that $n - t \leq |\sigma| < n$,

$$\left| \left\{ x : (\forall b)(\exists \tau \in [G \cap (b, +\infty)]^{n-|\sigma|}) f(\sigma, \tau) = x \right\} \right| \leq d_{n-|\sigma|}$$

Proof. Our forcing conditions are Mathias conditions (F, X) where $X \oplus C$ preserves non-c.e. definitions of the A 's. Lemma 3.16 in Wang [47] states that for every set G which is sufficiently generic for (F, X) , $G \oplus C$ preserves k non-c.e. definitions. It suffices therefore to prove the following lemma.

Lemma 4.25 For every condition (F, X) and $\sigma \in [\omega]^{<\omega}$ such that $n - t \leq |\sigma| < n$, for every finite set I such that $|I| = d_{n-|\sigma|}$, there exists an extension (F, \tilde{X}) such that

$$\{f(\sigma, \tau) : \tau \in [\tilde{X}]^{n-|\sigma|}\} \subseteq I \quad \text{or} \quad I \not\subseteq \{f(\sigma, \tau) : \tau \in [\tilde{X}]^{n-|\sigma|}\}$$

Proof. Define the function $g : [X]^{n-|\sigma|} \rightarrow I \cup \{\perp\}$ by $g(\tau) = f(\sigma, \tau)$ if $f(\sigma, \tau) \in I$ and $g(\tau) = \perp$ otherwise. By strong preservation of k non-c.e. definitions of $\text{TS}_{d_{n-|\sigma|+1}}^{n-|\sigma|}$, there exists an infinite subset $\tilde{X} \subseteq X$ such that $\tilde{X} \oplus C$ preserves non-c.e. definitions of the A 's and $|\{g(\tau) : \tau \in [\tilde{X}]^{n-|\sigma|}\}| \leq d_{n-|\sigma|}$. The condition (F, \tilde{X}) is the desired extension. \square

Using Lemma 3.16 in [47] and Lemma 4.25, one can define an infinite descending sequence of conditions $(\emptyset, \omega) \geq c_0 \geq c_1 \geq \dots$ such that for each $s \in \omega$

1. $c_s = (F_s, X_s)$ with $|F_s| \geq s$
2. c_s forces $W_e^{G \oplus C} \neq A_i$ if $s = \langle e, i \rangle$
3. $\{f(\sigma, \tau) : \tau \in [X_s]^{n-|\sigma|}\} \subseteq I$ or $I \not\subseteq \{f(\sigma, \tau) : \tau \in [X_s]^{n-|\sigma|}\}$ if $s = \langle \sigma, I \rangle$ and $|I| = d_{n-|\sigma|}$.

The set $G = \bigcup_s F_s$ is an infinite set such that $G \oplus C$ preserves non-c.e. definitions of the A 's. We claim that G satisfies the desired properties. Fix a $\sigma \in [\omega]^{<\omega}$ such that $n - t \leq |\sigma| < n$. Suppose that there exists $d_{n-|\sigma|} + 1$ elements $x_0, \dots, x_{d_{n-|\sigma|}}$ such that $(\forall b)(\exists \tau \in [G \cap (b, +\infty)]^{n-|\sigma|}) f(\sigma, \tau) = x_i$ for each $i \leq d_{n-|\sigma|}$. Let $I = \{x_0, \dots, x_{d_{n-|\sigma|-1}}\}$. By step $s = \langle \sigma, I \rangle$, G satisfies (F_s, X_s) such that $\{f(\sigma, \tau) : \tau \in [X_s]^{n-|\sigma|}\} \subseteq I$ or $I \not\subseteq \{f(\sigma, \tau) : \tau \in [X_s]^{n-|\sigma|}\}$. In the first case it contradicts the choice of $x_{d_{n-|\sigma|}}$ and in the second case it contradicts the choice of an element of I . This finishes the proof of Theorem 4.24. \square

4.3.3. Step (S1) : Construction of the set D . We start with the construction of an infinite set $D \subseteq \omega$ such that $D \oplus C$ preserves non-c.e. definitions of the A 's and a sequence $(I_\sigma : 0 < |\sigma| < n)$ such that for each $t \in (0, n)$ and each $\sigma \in [\omega]^t$

- (a) I_σ is a subset of $\{0, \dots, d_n\}$ with at most d_{n-t} many elements
- (b) $(\exists b)(\forall \tau \in [G \cap (b, +\infty)]^{n-t}) f(\sigma, \tau) \in I_\sigma$

Let D be the set constructed in Theorem 4.24 for $t = n - 1$. For each $\sigma \in [\omega]^{<\omega}$ such that $0 < |\sigma| < n$, let

$$I_\sigma = \{x \leq d_n : (\forall b)(\exists \tau \in [G \cap (b, +\infty)]^{n-|\sigma|}) f(\sigma, \tau) = x\}$$

By choice of D , the set I_σ has at most $d_{n-|\sigma|}$ many elements. Moreover, for each $y \leq d_n$ such that $y \notin I_\sigma$, there exists a bound b_y such that $(\forall \tau \in [D \cap (b_y, +\infty)]^{n-|\sigma|}) f(\sigma, \tau) \neq y$. So taking $b = \max(b_y : y \leq d_n \wedge y \notin I_\sigma)$, we obtain

$$(\forall \tau \in [D \cap (b, +\infty)]^{n-|\sigma|}) f(\sigma, \tau) \in I_\sigma$$

4.3.4. Step (S2) : Construction of the set E . We now construct an infinite set $E \subseteq D$ such that $E \oplus C$ preserves non-c.e. definitions of the A 's and a sequence $(I_t : 0 < t < n)$ such that for each $t \in (0, n)$

- (a) I_t is a subset of $\{0, \dots, d_n\}$ of size at most $d_t d_{n-t}$
- (b) $(\forall \sigma \in [E]^t)(\exists b)(\forall \tau \in [E \cap (b, +\infty)]^{n-t}) f(\sigma, \tau) \in I_t$

For each $t \in (0, n)$ and $\sigma \in [\omega]^t$, let $F_t(\sigma) = I_\sigma$. Using strong preservation of k non-c.e. definitions of $\text{TS}_{d_{t+1}}^t$, we build a finite sequence $D \supseteq E_1 \supseteq \dots \supseteq E_{n-1}$ such that for each $t \in (0, n)$

1. $E_t \oplus C$ preserves non-c.e. definitions of the A 's
2. $|F_t([E_t]^t)| \leq d_t$

Let $E = E_{n-1}$ and $I_t = \bigcup F_t([E]^t)$ for each $t \in (0, n)$. As for each $\sigma \in [E]^t$, $|F_t(\sigma)| = |I_\sigma| \leq d_{n-t}$, $|I_t| \leq d_t d_{n-t}$, so property (a) holds. We now check that property (b) is satisfied. Fix a $\sigma \in [E]^t$. By definition of I_t , $F_t(\sigma) = I_\sigma \subseteq I_t$. As $E \subseteq D$,

$$(\exists b)(\forall \tau \in [E \cap (b, +\infty)]^{n-t}) f(\sigma, \tau) \in I_\sigma \subseteq I_t$$

4.3.5. *Step (S3) : Construction of the set G.* Given the set E and the sequence of sets of colors $(I_t : 0 < t < n)$, we will construct a sequence $(\xi_i \in [E]^{<\omega} : i < \omega)$ such that

- (a) The set $G = \bigcup_i \xi_i$ is infinite and $G \oplus C$ preserves non-c.e. definitions of the A 's
- (b) $|f([\xi_i]^n)| \leq d_{n-1}$ and $\max(\xi_i) < \min(\xi_{i+1})$ for each $i < \omega$
- (c) For each $t \in (0, n)$ and $\sigma \in [\bigcup_{j < i} \xi_j]^t$, $f(\sigma, \tau) \in I_t$ for all $\tau \in [\bigcup_{j \geq i} \xi_j]^{n-t}$

We construct our set G by Mathias forcing (σ, X) where X is an infinite subset of E such that $X \oplus C$ preserves non-c.e. definitions of the A 's. Using property (b) of E , we can easily construct an infinite sequence $(\xi_i \in [E]^{<\omega} : i < \omega)$ satisfying properties (b) and (c) of step (S3). The following lemma shows how to satisfy property (a).

Lemma 4.26 Fix a condition (σ, X) , some $e \in \omega$ and some $j < k$. There exists an extension $(\sigma\xi, Y)$ with $|f([\xi]^n)| \leq d_{n-1}$, forcing $W_e^{G \oplus C} \neq A_j$.

Proof. Let W be the set of $a \in \omega$ such that for every coloring $g : [X]^n \rightarrow d_n + 1$, there is a set $\xi \in [X]^{<\omega}$ such that $|g([\xi]^n)| \leq d_{n-1}$ and $a \in W_e^{\sigma\xi \oplus C}$. The set W is $X \oplus C$ -c.e., therefore $W \neq A_j$. Let $a \in W \Delta A_j$. We have two cases:

- Case 1: $a \in W \setminus A_j$. In particular, taking $g = f$, there exists a set $\xi \in [X]^{<\omega}$ such that $|f([\xi]^n)| \leq d_{n-1}$ and $a \in W_e^{\sigma\xi \oplus C}$. Take the condition $(\sigma\xi, X)$ as the extension.
- Case 2: $a \in A_j \setminus W$. By definition of W , the collection \mathcal{C} of colorings $g : [X]^n \rightarrow d_n + 1$ such that for every set $\xi \in [X]^{<\omega}$ satisfying $|g([\xi]^n)| \leq d_{n-1}$, $a \notin W_e^{\sigma\xi \oplus C}$ is a non-empty $\Pi_1^{0, X \oplus C}$ class. As WKL_0 admits preservation of k non-c.e. definitions, there is some coloring $g \in \mathcal{C}$ such that $g \oplus X \oplus C$ preserves non-c.e. definitions of the A 's. By preservation of k non-c.e. definitions of $\text{TS}_{d_{n-1}+1}^n$, there exists an infinite subset $Y \subseteq X$ such that $Y \oplus C$ preserves non-c.e. definitions of the A 's and $|g([Y]^n)| \leq d_{n-1}$. The condition (σ, Y) forces $a \notin W_e^{G \oplus C}$.

□

Using Lemma 4.26 and property (b) of the set E , we can construct an infinite descending sequence of conditions $(\epsilon, E) \geq c_0 \geq \dots$ such that for each $s \in \omega$

- (i) $\sigma_{s+1} = \sigma_s \xi_s$ with $|\sigma_s| \geq s$ and $f([\xi_s]^n) \leq d_{n-1}$
- (ii) $f(\sigma, \tau) \in I_t$ for each $t \in (0, n)$, $\sigma \in [\sigma_s]^t$ and $\tau \in [X]^{n-t}$.
- (iii) c_s forces $W_e^{G \oplus C} \neq A_j$ if $s = \langle e, j \rangle$

where $c_s = (\sigma_s, X_s)$. The set $G = \bigcup_s \sigma_s$ satisfies the desired properties.

4.3.6. *Step (S4) : Construction of the set H.* Finally, we build an infinite set $H \subseteq G$ such that $H \oplus C$ preserves non-c.e. definitions of the A 's and $|f([H]^n)| \leq d_n$.

For each $i < \omega$, let $J_i = f([\xi_i]^n)$. By property (b) of step (S3), J_i is a subset of $\{0, \dots, d_n\}$ such that $|J_i| \leq d_{n-1}$. For each subset $J \subseteq \{0, \dots, d_n\}$ of size d_{n-1} , define the set

$$Z_J = \{x \in G : (\exists i)x \in \xi_i \wedge f([\xi_i]^n) \subseteq J\}$$

There exists finitely many such J 's, and the Z 's form a partition of G . Apply strong preservation of k non-c.e. definitions of $\text{TS}_{d_1+1}^1$ to obtain a finite set S of J 's of such that $|S| \leq d_1$ and an infinite set $H \subseteq \bigcup_{J \in S} Z_J \subseteq G$ such that $H \oplus G \oplus C$ preserves non-c.e. definitions of the A 's.

Lemma 4.27 $|f([H]^n)| \leq d_n$

Proof. As $H \subseteq G$, any $\sigma \in [H]^n$ can be decomposed into $\rho \hat{\ } \tau$ for some $\rho \in [\xi_i]^{<\omega}$ and some $\tau \in [\bigcup_{j \geq i} \xi_j]^{<\omega}$ with $|\rho| > 0$. If $|\tau| = 0$ then $f(\sigma) \in \bigcup_{J \in S} J$ by definition of H . If $|\tau| > 0$, then $f(\sigma) \in I_{|\rho|}$ by property (c) of step (S3). In any case

$$f(\sigma) \in \left(\bigcup_{J \in S} J \right) \cup \left(\bigcup_{t \in (0, n)} I_t \right)$$

Recall that $|S| \leq d_1$, $|J| = d_{n-1}$ for each $J \in S$, and $|I_t| \leq d_t d_{n-t}$ for each $t \in (0, n)$. Thus, applying the definition of d_n from (A3), $|f([H]^n)| \leq d_1 d_{n-1} + \sum_{0 < t < n} d_t d_{n-t} = d_n$, as desired. \square

This completes property (A3) and the proof of Theorem 4.21.

4.4. Free set theorem for tuples and non-c.e. definitions

Cholak et al. [5] studied the thin set theorem with infinitely many colors as a weakening of the free set theorem. The forcing notions used by Wang in [47] and by the author in [34] for constructing solutions to free set instances both involve the thin set theorem for a finite, but arbitrary number of colors. These constructions may suggest some relation between FS^n and TS_k^n for arbitrarily large k , but the exact nature of this relation is currently unclear.

In this section, we use the preservation of non-c.e. definitions of the thin set theorem to deduce similar preservation results for the free set theorem, and thereby separate FS from RT_2^2 over RCA_0 . More precisely, we prove the following preservation theorem.

Theorem 4.28 For every $k \in \omega$, FS admits strong preservation of k non-c.e. definitions.

The proof of Theorem 4.28 follows Corollary 4.30. Cholak et al. [5] asked whether any of FS^2 , $\text{FS}^2 + \text{COH}$ and $\text{FS}^2 + \text{WKL}_0$ imply RT_2^2 and Hirschfeldt [17] asked whether $\text{FS}^2 + \text{WKL}_0$ implies any of SRT_2^2 , ADS or CAC . We answer all of those questions negatively with the following corollary.

Corollary 4.29 For every $k \geq 2$, let Φ be the conjunction of COH , WKL_0 , RRT_2^2 , $\Pi_1^0\text{G}$, EM , TS_{k+1}^2 and FS . Over RCA_0 , Φ implies neither STS_k^2 nor SADS .

Corollary 4.30 FS does not imply RT_2^2 over ω -models.

The remainder of this section is devoted to the proof of Theorem 4.28. The proof is done by induction over the size of the tuples. The base case of our induction states that FS^0 admits strong preservation of k non-c.e. definitions. Consider FS^0 as a degenerate case of the free set theorem, where an instance is a constant c and a solution to c is an infinite set H which does not contain c . Indeed, a function $f : [\omega]^0 \rightarrow \omega$ can be considered as a constant c , and a set H is f -free if for every $\varepsilon \in [H]^0$, $f(\varepsilon) \in H \rightarrow f(\varepsilon) \in \varepsilon$. As $f(\varepsilon) \notin \varepsilon$, $f(\varepsilon) = c \notin H$. From now on, we will assume that FS^t admits strong preservation of k non-c.e. definitions for every $t \in [0, n)$.

We start with a lemma similar to Lemma 4.19.

Lemma 4.31 For every $n \geq 1$ and $k \geq 2$, if FS^n admits strong preservation of k non-c.e. definitions, then FS^{n+1} admits preservation of k non-c.e. definitions.

Proof. Fix any set C , k non- C -c.e. sets A_0, \dots, A_{k-1} and any C -computable coloring $f : [\omega]^{n+1} \rightarrow \omega$. Consider the uniformly C -computable sequence of sets \vec{R} defined for each $\sigma \in [\omega]^n$ and $y \in \omega$ by

$$R_{\sigma, y} = \{s \in \omega : f(\sigma, s) = y\}$$

As COH admits preservation of k non-c.e. definitions, there exists some \vec{R} -cohesive set G such that $G \oplus C$ preserves non-c.e. definitions of the A 's. The cohesive set induces a coloring $\tilde{f} : [\omega]^n \rightarrow \omega$ defined for each $\sigma \in [\omega]^n$ by

$$\tilde{f}(\sigma) = \begin{cases} \lim_{s \in G} f(\sigma, s) & \text{if it exists} \\ 0 & \text{otherwise} \end{cases}$$

As FS^n admits strong preservation of k non-c.e. definitions, there exists an infinite \tilde{f} -free set H such that $H \oplus G \oplus C$ preserves non-c.e. definitions of the A 's. In particular,

$$(\forall \sigma \in [H]^n)(\forall y \in H \setminus \sigma)(\exists b)(\forall s > b)f(\sigma, s) \neq y$$

$H \oplus G \oplus C$ computes an infinite f -free set. \square

4.4.1. *Trapped functions.* Although the notion of free set can be defined for every coloring over tuples of integers, we shall restrict ourselves to a particular kind of colorings: left trapped functions. The notion of trapped function was introduced by Wang in [48] to prove that FS does not imply ACA_0 over ω -models. It was later reused by the author in [34] to separate FS from WWKL_0 over ω -models.

Definition 4.32 A function $f : [\omega]^n \rightarrow \omega$ is *left (resp. right) trapped* if for every $\sigma \in [\omega]^n$, $f(\sigma) \leq \sigma(n-1)$ (resp. $f(\sigma) > \sigma(n-1)$).

The following lemma is a particular case of a more general statement proven by the author in [34]. It follows from the facts that FS^n for right trapped functions is strongly uniformly reducible to the diagonally non-computable principle (DNR), which itself is computably reducible to FS^n for left trapped functions.

Lemma 4.33 (Patey in [34]) For each $k, n \geq 1$, if FS^n for left trapped functions admits (strong) preservation of k non-c.e. definitions then so does FS^k .

It therefore suffices to prove strong preservation of k non-c.e. definitions for left trapped functions.

4.4.2. *Case of left trapped functions.* In this part, we will prove the following theorem which, together with Lemma 4.33 is sufficient to prove Theorem 4.28 by induction over n .

Theorem 4.34 For each $k, n \geq 1$, if FS^t admits strong preservation of k non-c.e. definitions for each $t \in [0, n)$, then so does FS^n for left trapped functions.

The proof of Theorem 4.34 begins after Lemma 4.36 and ends after Lemma 4.38. The two following lemmas will ensure that the reservoirs of our forcing conditions will have good properties, so that the conditions will be extensible.

Lemma 4.35 (Patey in [34]) Suppose that FS^t admits strong preservation of k non-c.e. definitions for each $t \in (0, n)$ for some $k \in \omega$. Fix a set C , some non- C -c.e. sets A_0, \dots, A_{k-1} , a finite set F and an infinite set X computable in C . For every function $f : [X]^n \rightarrow \omega$ there exists an infinite set $Y \subseteq X$ such that $Y \oplus C$ preserves non-c.e. definitions of the A 's and $(\forall \sigma \in [F]^t)(\forall \tau \in [Y]^{n-t})f(\sigma, \tau) \notin Y \setminus \tau$ for each $t \in (0, n)$.

Proof. Fix the finite enumeration $\sigma_0, \dots, \sigma_{m-1}$ of all elements of $[F]^t$ for all $t \in (0, n)$. We define a finite decreasing sequence of sets $X = Y_0 \supseteq Y_1 \supseteq \dots \supseteq Y_m$ such that for each $s < m$

- (a) none of the A 's are $Y_{s+1} \oplus C$ -c.e.
- (b) $\forall \tau \in [Y_{s+1}]^{n-|\sigma_s|} f(\sigma_s, \tau) \notin Y_{s+1} \setminus \tau$

Given some stage $s < m$ and some set Y_s , define the function $f_{\sigma_s} : [Y_s]^{n-|\sigma_s|} \rightarrow \omega$ by $f_{\sigma_s}(\tau) = f(\sigma_s, \tau)$. By strong preservation of k non-c.e. definitions of $\text{FS}^{n-|\sigma_s|}$, there exists an infinite set $Y_{s+1} \subseteq Y_s$ satisfying (a) and (b). We claim that Y_m satisfies the properties of the lemma. Fix some $\sigma \in [F]^t$ and some $\tau \in [Y_m]^{n-t}$ for some $t \in (0, n)$. There is a stage $s < m$ such that $\sigma = \sigma_s$. Moreover, $\tau \in [Y_{s+1}]^{n-|\sigma_s|}$, so by (b), $f(\sigma_s, \tau) \notin Y_{s+1} \setminus \tau$, therefore $f(\sigma, \tau) \notin Y_m \setminus \tau$, completing the proof. \square

Lemma 4.36 Suppose that $\text{TS}_{d_{t+1}}^t$ admits strong preservation of k non-c.e. definitions for each $t \in (0, n]$ and FS^t admits strong preservation of k non-c.e. definitions for each $t \in [0, n)$. For every function $f : [\omega]^n \rightarrow \omega$ and every set C preserving non-c.e. definitions of some sets A_0, \dots, A_{k-1} , there exists an infinite set X such that $X \oplus C$ preserves non-c.e. definitions of the A 's and for every $\sigma \in [X]^{<\omega}$ such that $0 \leq |\sigma| < n$,

$$(\forall x \in X \setminus \sigma)(\exists b)(\forall \tau \in [X \cap (b, +\infty)]^{n-|\sigma|})f(\sigma, \tau) \neq x$$

Proof. Let X be the infinite set constructed in Theorem 4.24 with $t = n$. For each $s < n$ and $i < d_{n-s}$, let $f_{s,i} : [X]^s \rightarrow \omega$ be the function such that $f_{s,i}(\sigma)$ is the i th element of $\{x : (\forall b)(\exists \tau \in [X \cap (b, +\infty)]^{n-s})f(\sigma, \tau) = x\}$ if it exists, and 0 otherwise. Define a finite sequence $X \supseteq X_0 \supseteq \dots \supseteq X_{n-1}$ such that for each $s < n$

1. $X_s \oplus C$ preserves non-c.e. definitions of the A 's
2. X_s is $f_{s,i}$ -free for each $i < d_{n-s}$

We claim that X_{n-1} is the desired set. Fix $s < n$ and take any $\sigma \in [X_{n-1}]^s$ and any $x \in X_{n-1} \setminus \sigma$. If $(\forall b)(\exists \tau \in [X \cap (b, +\infty)]^{n-s})f(\sigma, \tau) = x$, then by choice of X , there exists an $i < d_{n-s}$ such that $f_{s,i}(\sigma) = x$, contradicting $f_{s,i}$ -freeness of X_{n-1} . So $(\exists b)(\forall \tau \in [X \cap (b, +\infty)]^{n-s})f(\sigma, \tau) \neq x$. \square

Proof of Theorem 4.34. Fix $k \geq 2$, some set C , some non- C -c.e. sets A_0, \dots, A_{k-1} and a left trapped coloring $f : [\omega]^n \rightarrow \omega$. We will construct an infinite f -free set H such that none of the A 's is $H \oplus C$ -c.e. Our forcing conditions are Mathias conditions (F, X) such that

- (a) $X \oplus C$ preserves non-c.e. definitions of the A 's
- (b) $(\forall \sigma \in [F \cup X]^n)f(\sigma) \notin F \setminus \sigma$
- (c) $(\forall \sigma \in [F \cup X]^t)(\forall x \in (F \cup X) \setminus \sigma)(\exists b)(\forall \tau \in [(F \cup X) \cap (b, +\infty)]^{n-t})f(\sigma, \tau) \neq x$ for each $t \in [0, n)$.
- (d) $(\forall \sigma \in [F]^t)(\forall \tau \in [X]^{n-t})f(\sigma, \tau) \notin X \setminus \tau$ for each $t \in (0, n)$

Properties (c) and (d) will be obtained by Lemma 4.36 and Lemma 4.35 and are present to maintain the property (b) over extensions. A set G satisfies a condition (F, X) if it is f -free and satisfies the Mathias condition (F, X) . Our initial condition is (\emptyset, Y) where Y is obtained by Lemma 4.36.

Lemma 4.37 For every condition (F, X) there exists an extension (H, Y) such that $|H| > |F|$.

Proof. Choose an $x \in X$ such that $(\forall \sigma \in [F]^n)f(\sigma) \neq x$ and set $H = F \cup \{x\}$. By property (c) of (F, X) , there exists a b such that

$$(\forall \sigma \in [F]^t)(\forall \tau \in [X \cap (b, +\infty)]^{n-t})f(\sigma, \tau) \neq \{x\} \setminus \sigma$$

for each $t \in [0, n]$. By Lemma 4.35, there exists an infinite set $Y \subseteq X \setminus [0, b]$ such that $Y \oplus C$ preserves non-c.e. definitions of the A 's and property (d) is satisfied for (H, Y) . We claim that (H, Y) is a valid condition. Properties (a), (c) and (d) trivially hold. We now check property (b). By property (b) of (F, X) , we only need to check that $(\forall \sigma \in [F \cup Y]^k)f(\sigma) \neq x$. This follows from our choice of b . \square

Lemma 4.38 For every condition (F, X) , every $e \in \omega$ and $j < k$, there exists an extension (H, Y) forcing $W_e^{G \oplus C} \neq A_j$.

Proof. By removing finitely many elements of X , we can assume that $(\forall \sigma \in [F]^n)f(\sigma) \notin X$. For each $a \in \omega$, let \mathcal{C}_a be the $\Pi_1^{0, X \oplus C}$ class of left trapped functions $g : [X]^n \rightarrow \omega$ such that for every g -free set $E \subset X$, $a \notin W_e^{(F \cup E) \oplus C}$. Also define $W = \{a \in \omega : \mathcal{C}_a = \emptyset\}$. The set W is $X \oplus C$ -c.e. but A_j is not $X \oplus C$ -c.e., therefore $W \neq A_j$. Let $a \in W \Delta A_j$. We have two cases:

- Case 1: $a \in W \setminus A_j$. As $f \notin \mathcal{C}_a$, there exists a finite f -free set E such that $a \in W_e^{(F \cup E) \oplus C}$. Set $H = F \cup E$. By property (c) of (F, X) , there exists a b such that

$$(\forall \sigma \in [H]^t)(\forall x \in H)(\forall \tau \in [X \cap (b, +\infty)]^{n-t})f(\sigma, \tau) \neq \{x\} \setminus \sigma$$

for each $t \in [0, n)$. By Lemma 4.35, there exists an infinite set $Y \subseteq X \cap (b, +\infty)$ such that $Y \oplus C$ preserves non-c.e. definitions of the A 's and property (d) is satisfied for (H, Y) . We claim that (H, Y) is a valid condition.

Properties (a), (c) and (d) trivially hold. We now check property (b). By our choice of b , we only need to check that $(\forall \sigma \in [H]^n)f(\sigma) \notin H \setminus \sigma$. By property (b) of (F, X) , it suffices to check that $(\forall \sigma \in [H]^n)f(\sigma) \notin E \setminus \sigma$. By property (d) of (F, X) , and our initial assumption on X , we only need to check that $(\forall \sigma \in [E]^n)f(\sigma) \notin E \setminus \sigma$, which is exactly f -freeness of E .

- Case 2: $a \in A_j \setminus W$. By definition of W , $C_a \neq \emptyset$. As WKL_0 admits preservation of k non-c.e. definitions, there exists a left trapped functions $g \in C_a$ such that $g \oplus X \oplus C$ preserves non-c.e. definitions of the A 's. As FS^n admits preservation of k non-c.e. definitions, there exists some infinite g -free set $Y \subseteq X$ such that $Y \oplus C$ preserves non-c.e. definitions of the A 's. The condition (F, Y) forces $a \notin W_e^{G \oplus C}$ and therefore $W_e^{G \oplus C} \neq A_j$. □

Let Y be the set constructed in Lemma 4.36. Using Lemma 4.37 and Lemma 4.38, we can define an infinite decreasing sequence of conditions $(\emptyset, Y) \geq c_0 \geq \dots$ such that for every $s \in \omega$

- (i) $|F_s| \geq s$
- (ii) c_s forces $W_e^{G \oplus C} \neq A_j$ if $s = \langle e, j \rangle$

where $c_s = (F_s, X_s)$. Let $G = \bigcup_s F_s$. By (i), G is infinite and by (ii), none of the A 's are $G \oplus C$ -c.e. This completes the proof of Theorem 4.34. □

5. CONCLUSION

In this last section, we recall some existing open questions about the free set and thin set theorems, and state some new ones. Cholak et al. [5] asked the following question which remains open.

Question 5.1 Does TS^2 imply FS^2 over RCA_0 ?

We ask a related question motivated by the fact that the proof of cone avoidance of FS by Wang [48] and the preservation of k non-c.e. definitions of FS^2 in section 4.4 both use TS_k^2 for any k to construct a solution to an instance of FS^2 . We know by Corollary 4.29 that FS^2 does not imply TS_k^2 over RCA_0 for any k , but the reverse implication is still open.

Question 5.2 Does TS_3^2 imply FS^2 over RCA_0 ?

Cholak et al. [5] also asked whether $FS^2 + CAC$ implies RT_2^2 over RCA_0 . Using the equivalence between RT_2^2 and $EM + ADS$ proven by Bovykin and Weiermann [3], we ask the following related questions.

Question 5.3 Does any of FS^2 , TS^2 and TS_3^2 imply EM over RCA_0 ?

Acknowledgements. The author is thankful to his PhD advisor Laurent Bienvenu and to Wei Wang for useful comments and discussions. The author is funded by the John Templeton Foundation project *Structure and Randomness in the Theory of Computation* (Grant 48003). The opinions expressed in this publication are those of the author(s) and do not necessarily reflect the views of the John Templeton Foundation.

REFERENCES

- [1] Jeremy Avigad, Edward T. Dean, and Jason Rute. Algorithmic randomness, reverse mathematics, and the dominated convergence theorem. *Annals of Pure and Applied Logic*, 163(12):1854–1864, 2012.
- [2] Laurent Bienvenu, Ludovic Patey, and Paul Shafer. On the logical strengths of partial solutions to mathematical problems. Submitted. Available at <http://arxiv.org/abs/1411.5874>, 2015.
- [3] Andrey Bovykin and Andreas Weiermann. The strength of infinitary Ramseyan principles can be accessed by their densities. *Annals of Pure and Applied Logic*, page 4, 2005. To appear.
- [4] Vasco Brattka and Tahina Rakotoniaina. On the uniform computational content of Ramsey's theorem. Available at <http://arxiv.org/abs/1508.00471>, 2015.
- [5] Peter A. Cholak, Mariagnese Giusto, Jeffrey L. Hirst, and Carl G. Jockusch Jr. Free sets and reverse mathematics. *Reverse mathematics*, 21:104–119, 2001.
- [6] Peter A. Cholak, Carl G. Jockusch, and Theodore A. Slaman. On the strength of Ramsey's theorem for pairs. *Journal of Symbolic Logic*, 66(01):1–55, 2001.

- [7] C. Chong, Steffen Lempp, and Yue Yang. On the role of the collection principle for Σ_2^0 -formulas in second-order reverse mathematics. *Proceedings of the American Mathematical Society*, 138(3):1093–1100, 2010.
- [8] Chitat Chong, Theodore Slaman, and Yue Yang. The metamathematics of stable Ramsey’s theorem for pairs. *Journal of the American Mathematical Society*, 27(3):863–892, 2014.
- [9] François G. Dorais, Damir D. Dzhafarov, Jeffrey L. Hirst, Joseph R. Mileti, and Paul Shafer. On uniform relationships between combinatorial problems. *Trans. Amer. Math. Soc.*, 368(2):1321–1359, 2016.
- [10] Damir D. Dzhafarov. Strong reductions between combinatorial principles. In preparation.
- [11] Damir D. Dzhafarov. Cohesive avoidance and arithmetical sets. *arXiv preprint arXiv:1212.0828*, 2012.
- [12] Damir D. Dzhafarov. Cohesive avoidance and strong reductions. *Proceedings of the American Mathematical Society*, 143(2):869–876, 2014.
- [13] Richard Friedberg. A criterion for completeness of degrees of unsolvability. *J. Symb. Logic*, 22:159–160, 1957.
- [14] Harvey M. Friedman. Fom:53:free sets and reverse math and fom:54:recursion theory and dynamics. Available at <https://www.cs.nyu.edu/pipermail/fom/>.
- [15] Harvey M. Friedman. *Boolean Relation Theory and Incompleteness*. Lecture Notes in Logic, 2013. to appear. Available at <http://www.math.ohio-state.edu/>
- [16] Noam Greenberg and Joseph S. Miller. Lowness for Kurtz randomness. *Journal of Symbolic Logic*, 74(2):665–678, 2009.
- [17] Denis R. Hirschfeldt. *Slicing the truth*, volume 28 of *Lecture Notes Series. Institute for Mathematical Sciences. National University of Singapore*. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2015. On the computable and reverse mathematics of combinatorial principles, Edited and with a foreword by Chitat Chong, Qi Feng, Theodore A. Slaman, W. Hugh Woodin and Yue Yang.
- [18] Denis R. Hirschfeldt and Carl G. Jockusch. On notions of computability theoretic reduction between Π_1^1 principles. *To appear*.
- [19] Denis R. Hirschfeldt and Richard A. Shore. Combinatorial principles weaker than Ramsey’s theorem for pairs. *Journal of Symbolic Logic*, 72(1):171–206, 2007.
- [20] Carl G. Jockusch. Ramsey’s theorem and recursion theory. *Journal of Symbolic Logic*, 37(2):268–280, 1972.
- [21] Carl G. Jockusch. Degrees of generic sets. *Recursion Theory: its generalizations and applications*, pages 110–139, 1980.
- [22] Carl G. Jockusch, A Lewis, and Jeffrey B. Remmel. Π_1^0 -classes and Rado’s selection principle. *Journal of Symbolic Logic*, 56(02):684–693, 1991.
- [23] Carl G. Jockusch and Robert I. Soare. Π_1^0 classes and degrees of theories. *Transactions of the American Mathematical Society*, 173:33–56, 1972.
- [24] Carl G. Jockusch and Frank Stephan. A cohesive set which is not high. *Mathematical Logic Quarterly*, 39(1):515–530, 1993.
- [25] Steven M. Kautz. *Degrees of random sets*. PhD thesis, Citeseer, 1991.
- [26] Steven M. Kautz. An improved zero-one law for algorithmically random sequences. *Theoretical computer science*, 191(1):185–192, 1998.
- [27] Mushfeq Khan and Joseph S. Miller. Forcing with bushy trees. preprint, 2014.
- [28] Bjørn Kjos-Hanssen. Infinite subsets of random sets of integers. *Mathematics Research Letters*, 16:103–110, 2009.
- [29] Manuel Lerman, Reed Solomon, and Henry Towsner. Separating principles below Ramsey’s theorem for pairs. *Journal of Mathematical Logic*, 13(02):1350007, 2013.
- [30] Lu Liu. RT_2^2 does not imply WKL_0 . *Journal of Symbolic Logic*, 77(2):609–620, 2012.
- [31] Joseph Roy Mileti. *Partition theorems and computability theory*. ProQuest LLC, Ann Arbor, MI, 2004. Thesis (Ph.D.)—University of Illinois at Urbana-Champaign.
- [32] Webb Miller and Donald A. Martin. The degrees of hyperimmune sets. *Mathematical Logic Quarterly*, 14(7-12):159–166, 1968.
- [33] Antonio Montalbán. Open questions in reverse mathematics. *Bulletin of Symbolic Logic*, 17(03):431–454, 2011.
- [34] Ludovic Patey. Combinatorial weaknesses of Ramseyan principles. In preparation. Available at <http://ludovicpatey.com/media/research/combinatorial-weaknesses-draft.pdf>, 2015.
- [35] Ludovic Patey. Degrees bounding principles and universal instances in reverse mathematics. *Annals of Pure and Applied Logic*, 166(11):1165–1185, 2015.
- [36] Ludovic Patey. Iterative forcing and hyperimmunity in reverse mathematics. In Arnold Beckmann, Victor Mitraná, and Mariya Soskova, editors, *CiE. Evolving Computability*, volume 9136 of *Lecture Notes in Computer Science*, pages 291–301. Springer International Publishing, 2015.
- [37] Ludovic Patey. Controlling iterated jumps of solutions to combinatorial problems. *Computability*, 2016. To appear. Available at <http://arxiv.org/abs/1509.05340>.
- [38] Tahina Rakotoniaina. *The Computational Strength of Ramsey’s Theorem*. PhD thesis, University of Cape Town, 2015. to appear.
- [39] Joseph G. Rosenstein. *Linear orderings*, volume 98 of *Pure and Applied Mathematics*. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London, 1982.

- [40] David Seetapun and Theodore A. Slaman. On the strength of Ramsey's theorem. *Notre Dame Journal of Formal Logic*, 36(4):570–582, 1995.
- [41] Joseph R. Shoenfield. On degrees of unsolvability. *Annals of Mathematics*, 69(03):644–653, May 1959.
- [42] Stephen G. Simpson. An extension of the recursively enumerable Turing degrees. *Journal of the London Mathematical Society*, 75(2):287–297, 2007.
- [43] Stephen G. Simpson. *Subsystems of Second Order Arithmetic*. Cambridge University Press, 2009.
- [44] Frank Stephan. Martin-Löf random and PA-complete sets. In *Logic Colloquium '02*, volume 27 of *Lect. Notes Log.*, pages 342–348. Assoc. Symbol. Logic, La Jolla, CA, 2006.
- [45] Michiel Van Lambalgen. The axiomatization of randomness. *Journal of Symbolic Logic*, 55(03):1143–1167, 1990.
- [46] Wei Wang. Omitting cohesive sets. *arXiv preprint arXiv:1309.5428*, 2013.
- [47] Wei Wang. The definability strength of combinatorial principles, 2014. To appear. Available at <http://arxiv.org/abs/1408.1465>.
- [48] Wei Wang. Some logically weak Ramseyan theorems. *Advances in Mathematics*, 261:1–25, 2014.
E-mail address: ludovic.patey@computability.fr