**Quantity and quality of dispersed fine particles after low-energy water-dispersible
soil test: impact of the initial soil matrix potential**

Samouëlian, A.^{1*}, Quénard, L.², Cousin, I.², Cornu, S.³

1- LISAH, Univ Montpellier, INRA, IRD, Montpellier SupAgro, Montpellier, France

2- INRA, UR 0272 Science du Sol, Centre de Recherche d'Orléans, CS 40001, F-45075 Orléans
cedex 2, France

3- Aix Marseille Univ, CNRS, IRD, INRA, Coll France, CEREGE, Aix-en-Provence, France

*Corresponding author: anatja.samouelian@inra.fr

Abstract

Transfer of <2 µm soil particles is a key process in the dispersion of pollutants and the textural differentiation of soil. Dispersion tests involving low-energy water-dispersible clay (LEWDC) are commonly used to indicate the stock of particles <2 µm that can be mobilized. However, notable differences in LEWDC protocols exist. We discuss the impact of the initial soil matrix potential on the dispersion sensitivity of <2 µm particles and the nature (mineralogy and particle size) of the dispersed particles. The dispersion sensitivity was more differentiated among soils with a high initial water content, which allows better soil discrimination. The nature of dispersed <2 µm particles was closer to that of the bulk <2 µm soil when moist initial conditions are considered, whereas enrichment in coarse clay minerals (with a particle size mode around 2 µm) was observed with dry initial conditions. Performing the LEWDC test under both moist and dry initial conditions can be used to interpret soil functioning.

Abbreviations: LEWDC: low-energy water-dispersible clay

Highlights

- Moist initial condition allows better discrimination of particle dispersion among soils
- Nature of dispersed $<2\ \mu\text{m}$ particles varies with the initial soil matrix condition
- LEWDC test at two different initial conditions allows soil functioning analysis

Acknowledgments

This work was performed in the frame of the French ANR Blanc-AGRIPED project (ANR-10-Blanc 605) and of a PhD grant funded by the Région Centre and INRA-EA. The authors thank Olivier Josière and Lionel Cottenot for X-ray and laser particle size analysis, respectively; Gabriel Coelho, an internship fellow at INRA UR SOLS, for preliminary LEWDC tests; Bernard Renault, Hervé Gaillard and Christian Lelay for soil sampling; and Céline Ratié (US INFOSOL) and the GIS Sol for providing RMQS soil sites for resampling and the associated data. The authors are grateful to Arvalis for providing access to their long-term experimental site.

Supplemental file:

S1: Concentration of LEWDC dispersed $<2\ \mu\text{m}$ particles (LEWDC in g.kg^{-1}) for two the initial matrix potentials.

1 Introduction

Transfer of $<2\ \mu\text{m}$ soil particles is a key process in the dispersion of pollutants, as the fine particles act as vectors for many pollutants (McCarthy and Zachara, 1989), and in the textural differentiation of soil (Pedro et al., 1978). However, depending on the nature of the particles various types of contaminants could be adsorbed and transported, as previously demonstrated for radionuclides, metals, pesticides, and bacteria (e.g., McCarthy et al., 1998; Karathanasis, 1999; Gouy et al., 1999; Ryan et al., 1999, respectively). In addition, textural differentiation of soil is considered to result from a preferential transport of either smectites (e.g., Worrall et al., 1999) or other mineralogical clays (Cornu et al., 2014). Thus, determining the nature of dispersed particles is the first step in assessing the environmental impacts in such systems.

In parallel, laboratory dispersion tests have been developed to estimate soil aggregate stability or soil erodibility (Brubaker et al., 1992; Emerson, 1967; Le Bissonnais, 1996; Pojasok and Kay, 1990; Yoder, 1936). A large variety of dispersion protocols exist and have been extensively reviewed by Amézketa (1999). These tests were also used as indicators of the stock of $<2\ \mu\text{m}$ particles that can be mobilized from a given soil, as suggested by Kjaergaard et al. (2004) or, Dexter et al. (2011) and could thus also be used to determine the nature of the mobilized particles, i.e., their mineralogy and size. However, these dispersion tests have never been used for the latter determination to our knowledge. Kjaergaard et al. (2004) showed that the use of low-energy during the dispersion phase better represents the stock of particles less than $2\ \mu\text{m}$ that can be dispersed compared to the protocol based on a classical high mechanical energy. These authors proposed a “low-energy water-dispersible clay” (LEWDC) protocol that may mimic in situ particle mobilization from the upper soil horizon. Dexter et al. (2011) confirmed the importance of the initial matrix potential on the result of this test and underscored the benefit of

using two initial matrix potentials: a low potential pF 2.5 [-30 kPa] to mimic soil conditions at the end of a humid period and a high potential pF 6 [-100000 kPa] to mimic the dry topsoil at the surface.

In that context, our objective was to identify the nature of the particles released by a low-energy dispersion test considered as a proxy for clay transfer in soils and to determine if the initial matrix potentials of the soil act on the nature (mineralogical composition and size distribution) of the dispersed particles. For that purpose, we propose here a LEWDC adapted to address this objective.

2. Materials and Methods

2.1 Studied soils and total soil analysis

Seven arable soils were sampled from the A-horizon for 6 of them (Villamblain, Boigneville A, Boigneville B, Brosse, Cercottes, Mont-près-Chambord, and Faux-Perche) and from the E-horizon (Brosse) for the latter. Boigneville A and B were two plots of the same soil that experienced different agricultural practices, resulting in a higher organic carbon (OC) content in Boigneville B. According to the USDA texture classification, most soils had a silty loam texture ranging from silty clay loam and silt loam, except for Cercottes and Mont-près-Chambord, which had sandy loam and sandy clay textures, respectively (Table 1). These soils were chosen to represent a large span of content of $<2\ \mu\text{m}$ particles (from 12 to 37%) and organic carbon content (0.3 to 3%) with a focus on loamy soils that are considered as the most sensitive soils to clay transfer (Pedro et al., 1978). Their cation exchange capacity (CEC) ranged from 77 to 309 $\text{mmol}^+ \text{kg}^{-1}$, and their pH ranged from 5.4 to 8.3. More than 80% of their CEC was due to Ca^{2+} and less than 2.5% to Na^+ . For each soil, large undisturbed soil blocks were cut off the soil, along

a pit, with a knife and stored in box (30cm x 20cm x 15cm) at 4 °C before LEWDC test. Depth of sampling was 0-10 cm, except for the Brosse site where sampling took place between 35 and 45 cm depth.

Particle size distribution was determined on the whole soil according to the normalized protocol NF X 31-107 (by sieving and the Robinson pipette method); water pH, according to the normalized protocol NF ISO 10390; CEC and exchangeable cations, using the cobaltihexamine chloride method according to the normalized protocols NF ISO 23470 and NFX 31-130, respectively; and OC, according to the normalized protocol NF ISO 10694 with combustion at 1000°C.

2.2 Dispersion test protocol

LEWDC protocol consists of 3 steps which vary among authors. We built up our own protocol based on the most used settings.

- Pretreatment: The soil samples were equilibrated at a given matrix potential to control their water content. Two pretreatments were chosen: “pF 2.5” [-30 kPa] to obtain soil samples close to field capacity and “pF 6” [-100000 kPa] to obtain soil samples with a water content close to that of the soil surface during summer. The whole undisturbed soil samples were manually split into centimeter-size clods, saturated by capillarity on foam for 24 hours, and placed in a Richards apparatus (Richards, 1956) for 7 days for equilibration at a [-30 kPa] water potential for pF 2.5 [-30 kPa]. For pF 6 [-100000 kPa], the disturbed soil samples were oven dried at 35 °C for 5 days.
- Wetting: Two approaches exist in the literature, either immediate immersion after pretreatment (Emerson, 1967; Le Bissonnais, 1996; Seta and Karathanasis, 1996; Dexter et

al., 2011; Vendelboe et al., 2012) or slow resaturation (Kjaergaard et al., 2004). This last pretreatment being less common we chose an immediate immersion after pretreatment: the clods were placed in a glass bottle with osmosed water at a soil:water ratio of 1:8 (ratio classically used in the literature).

- Dispersion: For low energy dispersion the most classical approach consists in a ten fold manual inversion (Kjaergaard et al., 2004, Vendelboe et al., 2011). In this study the soil suspension in the glass bottle was manually inverted 10 times, over 1 minute. Then, the suspension was allowed to settle for 6 hours and 14 minutes for collection of $<2\ \mu\text{m}$ particle according to Stokes law, the upper 8cm were then collected.

2.3 Analysis of dispersed $<2\ \mu\text{m}$ particles

The particle concentrations of the LEWDC suspensions were measured by turbidimetry (Hach model 2100AN, Hach Compagny, Loveland, CO) after calibration. The uncertainty of the measurements was estimated at 5%.

Particle size was analyzed by a laser particle sizer (Malvern S) after further dispersion by NaPO_3 addition. The mineralogical composition of the $<2\ \mu\text{m}$ particles was determined by X-ray diffraction (XRD; Bruker-AXS, D8 Advance) on oriented slides by deposition of 1 mL of a $7.5\ \text{g L}^{-1}$ suspension that was air dry; samples were analyzed after the application of two treatments, following the methodology of Robert and Tessier (1974): air drying and ethylene glycol (EG) solvation.

The particle size and mineralogy were measured for (1) the $<2\ \mu\text{m}$ fraction of the bulk soils obtained by the Robison's pipette method and (2) the $<2\ \mu\text{m}$ fraction obtained after the LEWDC tests at the two initial matrix potentials. The nature of the dispersed particles at the Mont-près-Chambord and Cercottes sites was not analyzed due to limited sample quantities.

139
140 2.4 Data treatment: statistical analysis
141 For each LEWDC test replicates were performed. Results were presented with mean and standard
142 deviation except for XRD (all replicates presented).
143 For each site, an ANOVA at the 5% confidence level (Fisher pair-wise mean comparison
144 method) was used to assess the effects of soil properties and initial matrix potential on the
145 amount of LEWDC dispersed particles.

146
147
148 **3 Results**

149
150 3.1 Quantity of dispersed particles at different initial soil matrix potentials
151 The quantity of LEWDC dispersed particles was significantly higher at pF 2.5 [-30 kPa] than at
152 pF 6 [-1000000 kPa] in all the tested soils except for Mont-près-Chambord (Fisher's test at the
153 95% confidence level considering the two initial matrix potential together—Figure 1). The
154 greatest variation between the pF 6 [-1000000 kPa] and pF 2.5 [-30 kPa] pretreatments was
155 observed for Brosse. The difference in particle dispersion among the sites was greater at pF 2.5 [-
156 30 kPa], ranging from 37.2 g.kg⁻¹ to 5.7 g.kg⁻¹, than at pF 6 [-1000000 kPa], ranging from 5.8
157 g.kg⁻¹ to 1.4 g.kg⁻¹.
158 While comparing the quantity of particles LEWDC-dispersed by the different studied soils, i.e.
159 their sensitivity to dispersion, the soils did not follow the same sensitivity sequence for the two
160 initial matrix potential i.e., pF 6 [-1000000 kPa] and pF 2.5 [-30 kPa] (Fisher's test at the 95%
161 confidence level). Nevertheless, for both pretreatments, the site most sensitive to dispersion was
162 Brosse (Supplementary file, S1). Thus, the quantity of dispersed particles at a given matrix
163 potential is not proportional to that at the other matrix potentials.

3.2 Nature of the LEWDC dispersed $<2\ \mu\text{m}$ particles

The nature of the LEWDC particles, defined by their mineralogical composition and particle size distribution, was compared at the two initial matrix potentials and with the nature of the total $<2\ \mu\text{m}$ fraction of the soils for the five following sites: Brosse, Villamblain, Boigneville A, Boigneville B and Faux-Perche.

3.2.1 Mineralogical composition

The LEWDC dispersed $<2\ \mu\text{m}$ particles were richer in quartz at pF6 [-1000000 kPa] than at pF 2.5 [-30 kPa] and at pF 2.5 [-30 kPa] than in the total $<2\ \mu\text{m}$ soil fraction. Concerning clay minerals, three patterns were observed. For Brosse, the LEWDC dispersed $<2\ \mu\text{m}$ particles were poorer in smectite than was the total $<2\ \mu\text{m}$ fraction (Figure 2 a and b). This deficit was greater for the pretreatment at pF 6 [-1000000 kPa] than for that at pF 2.5 [-30 kPa]. For Boigneville A and B and Villamblain, the mineralogical composition of the LEWDC dispersed $<2\ \mu\text{m}$ particles at pF 2.5 [-30 kPa] was not different from that of the total $<2\ \mu\text{m}$ fraction; however, the LEWDC dispersed $<2\ \mu\text{m}$ particles were deficient in smectite relative to the total $<2\ \mu\text{m}$ fraction at pF 6 [-1000000 kPa] (Figure 2c and d—only Boigneville A is presented). For Faux-Perche, the LEWDC dispersed $<2\ \mu\text{m}$ particles had the same mineralogical composition as the total $<2\ \mu\text{m}$ fraction of the soil, irrespective of the initial matrix potentials (Figure 2e and f).

3.2.2 Particle size distribution

The particle size distribution of the total $<2\ \mu\text{m}$ fraction was bimodal for Brosse, Boigneville A and Villamblain, with two modes of similar proportion at approximately $0.2\ \mu\text{m}$ and $2\ \mu\text{m}$ (Figure 3a, c, and e). The mode at approximately $0.2\ \mu\text{m}$ is related to the presence of smectite,

while the second mode is due to either larger clay minerals (illite, kaolinite, chlorite, quartz, and feldspars) or microaggregates of smectites, as shown by Hubert et al. (2012).

For Faux-Perche, the particle size distribution of the $<2\text{ }\mu\text{m}$ fraction exhibited a unique mode at approximately $2\text{ }\mu\text{m}$ (Figure 3b), which is consistent with the absence of smectite in this soil (Figure 2e and f). For Boigneville B, the particle size distribution of the $<2\text{ }\mu\text{m}$ fraction also exhibited a main mode at approximately $2\text{ }\mu\text{m}$ but with a slight shouldering at approximately $0.53\text{ }\mu\text{m}$ (Figure 3d). The absence of a $0.2\text{ }\mu\text{m}$ mode in the particle size distribution may have been due to the presence of microaggregates of smectite stabilized by organic bonds. This is consistent with the higher OC content in Boigneville B than in Boigneville A (Table 1).

While comparing the particle size distribution of the LEWDC dispersed $<2\text{ }\mu\text{m}$ particles to that of the total $<2\text{ }\mu\text{m}$ fraction of the soils, four patterns were observed. For Brosse, the LEWDC dispersed $<2\text{ }\mu\text{m}$ particles presented a bimodal particle size distribution comparable to that of the total $<2\text{ }\mu\text{m}$ fraction but with slight shifts in the positions of the modes and their proportions (Figure 3a). After the LEWDC test, the proportion of fine particles decreased, while the proportion of coarse particles increased, especially at pF 6 $[-1000000\text{ kPa}]$.

Regarding Boigneville A and Villamblain, the LEWDC dispersed $<2\text{ }\mu\text{m}$ particles pretreated at pF 2.5 $[-30\text{ kPa}]$ exhibited a bimodal distribution with a smaller fine particle mode (and larger coarse mode) than that of the total $<2\text{ }\mu\text{m}$ fraction, while the LEWDC particles pretreated at pF 6 $[-1000000\text{ kPa}]$ had a monomodal distribution centered at the same position as the coarse mode of the total $<2\text{ }\mu\text{m}$ fraction (Figure 3c and e).

Concerning Faux-Perche, the LEWDC dispersed $<2\text{ }\mu\text{m}$ particles had a monomodal particle size distribution similar to that of the total $<2\text{ }\mu\text{m}$ fraction of the soil but with a slight increase in the mode at pF 6 $[-1000000\text{ kPa}]$ (Figure 3b).

For Boigneville B, the LEWDC dispersed $<2\ \mu\text{m}$ particles exhibited a monomodal distribution comparable to that of the total $<2\ \mu\text{m}$ fraction but with a smaller shouldering at pF 2.5 [-30 kPa]. At pF 6 [-1000000 kPa], this shouldering disappeared, and the mode was shifted toward coarser particles, with a large proportion of particles $>2\ \mu\text{m}$ probably due to aggregation processes after sampling (Figure 3d).

As smectites are the finest clay mineral particles, the decrease in the $0.2\ \mu\text{m}$ mode corresponds to the decrease in smectites observed in the XRD data, while the increase in the $2\ \mu\text{m}$ mode corresponds to an increase in other clay minerals, e.g., feldspars and quartz.

4. Discussion/Conclusion

Our results demonstrated that the amount of LEWDC dispersed $<2\ \mu\text{m}$ particles vary strongly with the initial matrix potential of the sample, and this variation differs in amplitude from one soil to another. As emphasized by Amézketa in 1999, we also support the need for protocol standardization. The air-drying pretreatment is probably easier to standardize; nevertheless, the sensitivity of soils to particle dispersion by the LEWDC test was more pronounced at pF 2.5 [-30 kPa], as observed by Dexter et al. (2011). Therefore, to determine the sensitivity of different soils to particle dispersion, we recommend a controlled, moist initial condition equal to pF 2.5 [-30 kPa].

We showed that the nature (mineralogical composition and particle size distribution) of dispersed $<2\ \mu\text{m}$ particles is not similar to that of the total $<2\ \mu\text{m}$ fraction and is a function of the initial matrix potential. The nature of the dispersed $<2\ \mu\text{m}$ particles was generally more similar to that of the total $<2\ \mu\text{m}$ fraction at pF 2.5 [-30 kPa] than at pF 6 [-1000000 kPa] and always accompanied by a slight enrichment in coarser minerals. Depending on the soil type, this particle segregation is more or less pronounced. Therefore, we propose applying the LEWDC test at two

235 contrasting initial matrix potentials with a systematic determination of the nature of the dispersed
236 $<2\ \mu\text{m}$ particles in order to interpret the results in terms of soil processes such as soil textural
237 differentiation or soil contamination via particle transfer.
238

4 References

- Amézketa, E. 1999. Soil aggregate stability: A review. *J. Sustain. Agric.* 14:83-151.
- Brubaker, S.C., C.S. Holzhey, and B.R. Brasher. 1992. Estimating the water-dispersible clay content of soils. *Soil Sci. Soc. Am. J.* 56:1614-1620.
- Cornu, S., L. Quénard, I. Cousin, and A. Samouëlian. 2014. Experimental approach of lessivage: Quantification and mechanisms. *Geoderma* 213:357-370.
- Dexter, A.R., G. Richard, J. Davy, M. Hardy, and O. Duval. 2011. Clay dispersion from soil as a function of antecedent water potential. *Soil Sci. Soc. Am. J.* 75:444-455.
- Emerson, W.W. 1967. A classification of soil aggregates based on their coherence in water. *Aust. J. Soil Res.* 5:47-57.
- Gouy, V., J.C. Dur, R. Calvet, R. Belamie, and V. Chaplain. 1999. Influence of adsorption-desorption phenomena on pesticide run-off from soil using simulated rainfall. *Pestic. Sci.* 55:175-182.
- Hubert, F., L. Caner, A. Meunier, and E. Ferrage. 2012. Unraveling complex <2 μm clay mineralogy from soils using X-ray diffraction profile modeling on particle-size sub-fractions: Implications for soil pedogenesis and reactivity. *Am. Mineral.* 97:384-398.
- Karathanasis, A.D. 1999. Subsurface migration of copper and zinc mediated by soil colloids. *Soil Sci. Soc. Am. J.* 63:830-838.
- Kjaergaard, C., L.W. de Jonge, P. Moldrup, and P. Schjønning. 2004. Water-dispersible colloids: Effects of measurement method, clay content, initial soil matric potential, and wetting rate. *Vadose Zone J.* 3:403–412.
- Le Bissonnais, Y. 1996. Aggregate stability and assessment of soil crustability and erodibility: I. Theory and methodology. *Eur. J. Soil Sci.* 47:425–437.

- 262 McCarthy, J.F., and J.M. Zachara. 1989. Subsurface transport of contaminants: Binding to mobile
263 and immobile phases in groundwater aquifers. *Environ. Sci. Technol.* 23:496-504.
- 264 McCarthy, J.F., W.E. Sanford, and P.L. Stafford. 1998. Lanthanide field tracers demonstrate
265 enhanced transport of transuranic radionuclides by natural organic matter. *Environ. Sci.*
266 *Technol.* 32:3901-3906.
- 267 NF X 31-107. 2003. Qualité du sol – Détermination de la distribution granulométrique des
268 particules du sol - Méthode à la pipette AFNOR.
- 269 NF ISO 10390. 2005. Qualité du sol -- Détermination du pH. AFNOR
- 270 NF ISO 23470. 2007. Qualité du sol - Détermination de la capacité d'échange cationique (CEC)
271 effective et des cations échangeables à l'aide d'une solution de trichlorure de
272 cobaltihexammine. AFNOR
- 273 NF X31-130. 1999. Qualité du sol - Détermination de la capacité d'échange cationique (CEC) et
274 des cations extractibles. AFNOR
- 275 NF ISO 10694. 1995. Qualité du sol Dosage du carbone organique et du carbone total après
276 combustion sèche (analyse élémentaire) AFNOR
- 277 Pedro, G., M. Jamagne, and J.C. Begon. 1978. Two routes in genesis of strongly differentiated
278 acid soils under humid, cool-temperate conditions. *Geoderma* 20:173–189.
- 279 Pojasok, T., and B.D. Kay. 1990. Assessment of a combination of wet sieving and turbidimetry to
280 characterize the structural stability of moist aggregates. *Can. J. Soil Sci.* 70:33-42.
- 281 Richards, L.A. 1956. Sample retainers for measuring water retention by soil. *Soil Sci. Soc. Am.*
282 *Proc.* 20:301-303.
- 283 Robert, M., and D. Tessier. 1974. Méthode de préparation des argiles des sols pour des études
284 minéralogiques. *Annale Agronomique* 25:859–882.

- 285 Ryan, J.N., M. Elimelech, R.W. Harvey, and P.R. Johnson. 1999. Bacteriophage PRD1 and silica
286 colloid transport and recovery in an iron oxide-coated sand aquifer. *Environ. Sci.*
287 *Technol.* 33:63-73.
- 288 Seta, A.K., and Karathanasis, A.D. 1996. Water dispersible colloids and factors influencing their
289 dispersibility from soil aggregates. *Geoderma.* 74:255-266.
- 290 Vendelboe, A.L., Moldrup, P., Heckrath, G., Jin, Y., and de Jong, L.W. 2011. Colloids and
291 phosphorus leaching from undisturbed soil cores sampled along a natural clay gradient.
292 *Soil Science.* 176:399-406
- 293 Vendelboe, A.L., Schjonning, P., Moldrup, P., Jin, Y., Merbach, I. and de Jong, L.W. 2012.
294 Colloid release from differently managed loess soil. *Soil Science.* 177:301-309
- 295 Worrall, F., A. Parker, J.E. Rae, and A.C. Johnson. 1999. A study of suspended and colloidal
296 matter in the leachate from lysimeters and its role in pesticide transport. *J. Environ. Qual.*
297 28:595-604.
- 298 Yoder, R.E. 1936. A direct method of aggregate analysis of soils and a study of the physical
299 nature of erosion losses. *Agron. J.* 28:337-351.

Table 1: Pedological characteristics of the total soils.

Site	Particle size (g.kg ⁻¹)			OC (g.kg ⁻¹)	pH _{H2O}	CEC	Ca ²⁺	Mg ²⁺	K ⁺	Na ⁺
	0-2	2-50	50-2000							
	μm									
Brosse	307	587	106	3.5	6.2	178	155	11	0	4
Villamblain	328	648	24	14.6	7.8	210	186	8.6	5.9	0.6
Boigneville A	295	633	72	12.1	6.9	147	134	6.9	4.3	0.4
Boigneville B	247	657	96	30	5.4	142	116	7.1	8.8	0.5
Faux Perche	125	825	50	11.1	7.9	77	95	4.4	4.3	0.2
Cercottes	197	143	660	12.7	8.2	140	140	4.6	11.9	0.19
Mont-près-Chambord	366	158	476	14.3	8.3	309	278	15.6	10.2	0.4

OC: organic carbon, CEC: cation exchange capacity

List of figures

Figure 1: Concentrations of dispersed 0-2 μm particles for the different studied soils for the two pretreatments. Circles correspond to the pF 2.5 pretreatment, and squares to the pF 6 pretreatment. Letters correspond to a Fisher's mean comparison test performed for all the samples with the two pretreatments at the 95 % confidence level.

Figure 2: XRD patterns of the 0-2 μm fractions of some of the studied soils: a and b—Brosse, c and d—Boigneville A, and e and f—Faux Perche. a, c and e—air dry treatment; b, d and f—EG solvation. For each site, we present the total 0-2 μm fraction of the soil (black) and the 0-2 μm fraction extracted in the LEWDC tests at pF 6 (red) and pF 2.5 (blue). Multiple curves of the same color represent replicates.

Figure 3: Particle size distribution of the 0-2 μm fractions of the studied soils: a) Brosse, b) Faux Perche, c) Boigneville A, d) Boigneville B and e) Villamblain. For each site, the total 0-2 μm fraction (black) and the 0-2 μm particles dispersed in the LEWDC test at pF 6 (red) and pF 2.5 (blue) are reported.

Figure 1: Concentrations of dispersed 0-2 μm particles for the different studied soils for the two pretreatments. Circles correspond to the pF 2.5 pretreatment, and squares to the pF 6 pretreatment. Letters correspond to a Fisher’s mean comparison test performed for all the samples with the two pretreatments at the 95 % confidence level.

Figure 2: XRD patterns of the 0-2 μm fractions of some of the studied soils: a and b—Brosse, c and d—Boigneville A, and e and f—Faux Perche. a, c and e—air dry treatment; b, d and f—EG solvation. For each site, we present the total 0-2 μm fraction of the soil (black) and the 0-2 μm fraction extracted in the LEWDC tests at pH 6 (red) and pH 2.5 (blue). Multiple curves of the same color represent replicates.

Figure 3: Particle size distribution of the 0-2 μm fractions of the studied soils: a) Brosse, b) Faux Perche, c) Boigneville A, d) Boigneville B and e) Villamblain. For each site, the total 0-2 μm fraction (black) and the 0-2 μm particles dispersed in the LEWDC test at pH 6 (red) and pH 2.5 (blue) are reported.