1 Recovery of Mediterranean steppe vegetation after cultivation: legacy effects on
2 plant composition, soil properties and functional traits

3
4 Juliane Helm^{1,2}, Thierry Dutoit³, Arne Saatkamp⁴, Solveig Franziska Bucher¹, Matthias Leiterer⁵ &
5 Christine Römermann^{1,6}

6
7 **Helm, J.** (Corresponding author, juliane.helm@bgc-jena.mpg.de)^{1,2}

8 **Dutoit, T.** (thierry.dutoit@imbe.fr)³

9 **Saatkamp, A.** (arne.saatkamp@imbe.fr)⁴

10 **Bucher, S. F.** (solveig.franziska.bucher@uni-jena.de)¹

11 **Leiterer, M.** (Matthias.Leiterer@tll.thueringen.de)⁵

12 **Römermann, C.** (christine.roemermann@uni-jena.de)^{1,6}

13
14 ¹Institute of Ecology and Evolution, Department of Plant Biodiversity, Friedrich-Schiller-University
15 Jena, Philosophenweg 16, 07743 Jena, Germany

16 ²Current address: Max-Planck Institute for Biogeochemistry, Department of Biogeochemical Processes,
17 Hans Knöll Str. 10, 07745 Jena, Germany

18 ³Institut Méditerranéen de Biodiversité et d'Ecologie, Avignon Univ, Aix Marseille Univ, CNRS, IRD,
19 IMBE, IUT site Agroparc, 337 Chemin des Meinajaries BP 61207, F-84911 Avignon cedex 09, France

20 ⁴Institut Méditerranéen de Biodiversité et d'Ecologie, Aix Marseille Univ, CNRS, IRD, Avignon Univ,
21 IMBE, Facultés St Jérôme case 421, F-13397 Marseille cedex 20, France

22 ⁵Thüringer Landesanstalt für Landwirtschaft, Naumburger Str. 98, 07702 Jena, Germany

23 ⁶German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, 04103 Leipzig,
24 Germany

Abstract

Questions: Are the vegetation structure and soil properties of fallows similar to adjacent species-rich Mediterranean steppe communities 35 years after abandonment? Is there a spontaneous re-development towards the original steppe vegetation after re-introduction of the traditional grazing system? Can differences in functional trait composition be identified according to different land-use types? Do traits of selected species affect the ability to re-colonize fallow land?

Location: Mediterranean steppe of La Crau (Southeastern France)

Methods: We conducted 80 vegetation surveys and soil analyses in the steppe and adjacent fallow land in six locations in La Crau in 2015; for three locations, data from 2001 was included. To compare the functional composition of steppe vs. fallow land, community-weighted means (CMW) of specific leaf area (SLA), leaf dry matter content (LDMC) and canopy height were calculated. To test whether trait variability is associated with the re-colonization success of 12 selected target species, we measured SLA, LDMC, canopy height, plant width and aboveground biomass on eight replicate individuals per site and species.

Results: Fallow land was characterized by lower species richness and elevated phosphorus and potassium content in the soil. Comparing vegetation relevés between 2001 and 2015 revealed that the fallows were developing towards steppe vegetation. CWMs of SLA were significantly lower and CWM of LDMC higher in steppe than in fallow land. There was no similarity in trait composition between steppe species that successfully re-colonized the fallow-land community, and no evidence that intraspecific trait variability is associated with re-colonization success of these species.

Conclusions: Even after decades of abandonment and re-introduction of the grazing system, the vegetation structure, soil properties and plant functional trait composition of fallows differ from those of adjacent steppe communities. These results illustrate the need for active restoration to accelerate the recovery of former arable land to the target steppe communities.

Keywords: Community trait means; Cultivation; Dry Grassland; Grazing; Intraspecific trait variability; Leaf dry matter content; Natural recovery; Restoration; Specific leaf area

Nomenclature: Jäger & Werner (2002); Pavon (2014)

Abbreviations: CV = coefficient of variation; CWM = community-weighted mean; ITV = intraspecific trait variability; LDMC = leaf dry matter content; SLA = specific leaf area

Running head: Recovery of Mediterranean steppe vegetation

Introduction

Semi-natural grasslands including the steppes of the Mediterranean basin are among Europe's most species-rich ecosystems (Willems 1983; WallisDeVries et al. 2002). Nevertheless, the 20th century's agricultural intensification caused a dramatic decrease in steppe grasslands worldwide and led to fragmentation as well as eutrophication of these habitats, leading to a decrease in plant diversity (Poschlod et al. 1998; White et al. 2000; Karlik & Poschlod 2009). To protect the species diversity and to better connect semi-natural grasslands, grassland remnants need to be conserved or, if already altered, restored. For the restoration of these species rich grasslands, several techniques have already been proposed (Kiehl et al. 2010 for review) as the natural colonization of characteristic grassland species into former intensively cultivated areas is usually very slow (Römermann et al. 2005, Buisson et al. 2006). It has already been shown that it might take several decades or centuries until the species composition and diversities of grasslands return (Forey & Dutoit 2012); accordingly, active restoration measures are typically needed.

To evaluate restoration success, traditional restoration studies assess the floristic similarity between restored and original (target) vegetation, focussing on species composition as well as diversity (Jaunatre et al., 2013). However, these indicators are highly dependent on site characteristics that determine the local species-pool (Stuble et al. 2017). A complementary way of assessing restoration success is to focus on the restoration of the functional composition (Laughlin 2014, Engst et al. 2016), which is especially relevant for those sites where original species are lost from the local species pool. The analyses of plant functional traits can provide insights into how land-use changes affect vegetation dynamics and function (e.g. Römermann et al. 2009, Breitschwerdt et al. 2015), but this trait-based approach has rarely been examined in connection with restoration success in Mediterranean dry grasslands.

Various habitat filters (e.g., environmental factors such as climate conditions, soil properties, or disturbance) allow only species with certain trait values to establish (Keddy 1992; Weiher & Keddy 1995; Diaz et al. 1998; Venn et al. 2011). The species that pass these filters are necessarily similar, which eventually leads to trait convergence (Diaz et al. 1998; Wilson 2007; Laughlin et al. 2012). But, species also need to be dissimilar enough to avoid competition, according to the limiting similarity hypothesis, which eventually leads to trait divergence (MacArthur & Levins 1967; Wilson 2007; Cornwell & Ackerly 2009; Laughlin et al. 2012). Comparing the functional composition of original vs. changed plant communities and analyzing the trait composition of target species that have already successfully re-established in the restoration sites seem promising as this approach complements the taxonomic evaluation of restoration success, therewith leading to a better understanding of the factors preventing successful restoration.

The aim of this study was to evaluate whether the taxonomic and functional composition of fallow lands can develop into the direction of original steppe communities without having established active restoration measures such as the addition of propagules.

We performed vegetation relevés to assess whether fallow lands re-developed into the direction of the target steppe vegetation with respect to the species composition and diversity. To elucidate the potential natural development of fallow land vegetation towards the original steppe vegetation we analyzed temporal trends in vegetation changes by comparing vegetation relevés and soil analyses from 2001 with results for 2015.

We explored the functional composition (community-weighted mean = CWM) of traits related to plant performance and growth rate and as indicators of competitive ability and environmental tolerance of species (Garnier 1992; Reich et al. 1997; Poorter et al. 2009; Lauterbach et al. 2013; Pérez-Harguindeguy et al. 2013). Nutrient-rich environments such as former agricultural land typically favor plants with high specific leaf area (SLA) and low leaf dry matter content (LDMC). Vice versa, species with trait values indicating nutrient

conservation strategies, slow growth rates and high longevity should dominate in nutrient poor environments (Chapin et al. 1993; Garnier 1992; Lavorel & Garnier 2002; Pérez-Harguindeguy et al. 2013; Poorter & Garnier 1999; Poorter & Remkes 1990; Römermann et al. 2009; Villar et al. 2005). Canopy height reflects the competitive ability of plant communities through better light assessment (Gaudet & Keddy 1988; Westoby et al. 2002), yet the investment in height incurs costs for construction and maintenance of the stem (Falster & Westoby 2003) and thus should be positively related to soil nutrient availability.

Intraspecific variation patterns in plant functional traits are a major determinant of the ability to respond and evolve with climate change and changing environmental conditions (Nicotra et al. 2010; Laforest-Lapointe et al. 2014; Bucher et al. 2017; Bucher et al. 2018). In some cases, trait variability can even be more important than particular trait values: species showing high functional variability may successfully re-establish because they manage to adapt to changing environmental conditions (Albert et al. 2011; Jung et al. 2010). In addition to the above mentioned traits we included additional plant width and biomass to test whether high intraspecific trait variability is related to re-establishment success. Both traits reflect competitive strength and should thus relate positively to soil nutrients (Pérez-Harguindeguy et al. 2013).

To summarize, this study combines a taxonomic with a trait-based approach to i) identify the effects of former land uses on present vegetation composition, species diversity and physical-chemical soil parameters, and ii) investigate whether re-development towards the original steppe vegetation can be detected after a further 14 years of recovery under traditional grazing conditions (2001 vs. 2015). Moreover, we iii) investigate whether differences in CWM of plant height, SLA and LDMC between fallow land and original steppe vegetation are still observable and attempt to iv) determine whether the traits of successful colonizers show higher functional variability than those of species that have not re-established on fallow lands yet.

Materials and Methods

Study sites

The plain of La Crau (Fig. 1) is the last remaining xeric steppe (Devaux et al. 1983) in southeastern France (43°33'N, 4°52'W). The area was formed by ancient river deposits during the Pleistocene period between 2 000 000 BP and 30 000 BP (Molliex et al. 2013). It has a Mediterranean climate with a frequent, strong northwesterly wind, the Mistral. Siliceous pebbles cover approximately 50% of La Crau, and plant roots are prevented from reaching the alluvial water table by an impermeable conglomerate lying 40–60 cm deep (Colomb & Roux 1978). Traditionally, the land has been used for itinerant sheep grazing, probably since early Neolithic times (Buisson & Dutoit 2006). Industry, military activities, quarries, highways and intensive agriculture led to fragmentation and reduction of the semi-natural steppe vegetation from approximately 60 000 ha to 11 500 ha in the 20th century (Buisson & Dutoit 2006; Deverre 1996; Etienne et al. 1998). During the period of melon cultivation (1965–1985), intensive fertilization, herbicides and deep ploughing changed the soil structure and degraded the conglomerate (Römermann et al. 2004, 2005). In contrast, cereals (1950–1966) were cultivated without deep ploughing and using less fertilizer (Devaux et al. 1983; Masip 1991). At the time of abandonment, no active restoration operations were realized. Grazing, representing an important factor for the maintenance of these grasslands, was re-introduced immediately when cultivation ceased (Dureau & Bonnefon 1998). The recovery of fallow land in our study area was thus not a strictly passive restoration technique. The flocks of sheep link remaining steppe patches (the target) and fallow lands and potentially facilitate the dispersal and establishment of steppe species into fallow lands (Fischer et al. 1996).

Vegetation surveys

To investigate whether type of cultivation and time since abandonment play a role in restoration success on fallow land, six locations containing a total of 16 study sites differing in

their land-use histories were examined (Table 1, Fig. 1). One long-abandoned and one recently abandoned cereal field were included, to represent different time spans.

We conducted vegetation relevés on 20 m x 20 m sites, placing five replicate 2 m x 2 m quadrats at each corner and in the middle (hereafter referred to as 'plots'). Species cover was estimated using the scale of Braun-Blanquet (1964). At each location, at least two sites consisting of one undisturbed steppe plot and one adjacent fallow land plot grazed by the same flock of sheep were studied. The locations Petit Abondoux, Poulagère and Couliès were sampled in May 2014 (Tanet 2014; Fig. 1), and the locations Peau de Meau, Grand Abondoux and Valigne in April 2015 (Fig. 1).

Percentages of vegetation cover, stone cover and bare ground were estimated on the five plots at the time of vegetation sampling. To estimate the standing crop, we cut the total aboveground biomass on three 0.5 m x 0.5 m randomly chosen quadrat within each site. The leaf area index (LAI), the ratio of foliage area to ground surface area (Watson 1947), of each of the five plots was measured using the LAI-2200C Plant Canopy Analyzer. Species richness S representing the total number of species, Shannon diversity index (H_s) and Evenness (E) were calculated for each plot (Mühlenberg 1993).

Soil sampling

To identify the effect of former land use on physical-chemical soil parameters, three replicate mixed soil samples of the upper soil layer (0–10 cm) were collected randomly within each site. After drying and sieving with a 2 mm mesh sieve, soil samples were analyzed at the INRA soil analysis laboratory (*Institut National de la Recherche Agronomique*, Arras, France) according to standardized protocols. Organic carbon (C in g kg⁻¹) and total nitrogen content (N in g kg⁻¹) were analyzed via the dry combustion method using a CHN elemental analyzer (Vario EL Elemental Analyzer, Elementar). Phosphorus content (P in g kg⁻¹) was measured by the Olsen method (Olsen 1954). Calcium (Ca in g kg⁻¹), magnesium (Mg in g kg⁻¹) and potassium (K in g kg⁻¹) were determined by the Metson method (Metson 1957). Additionally, soil pH was

measured in a 1:1 soil-water solution, after stirring once followed by a 24 h rest period. Carbon-nitrogen ratio (C:N) and organic matter (OM in g kg⁻¹) were calculated using total C and N results. The amount of clay (> 0.002 mm), fine silt (0.002–0.02 mm), coarse silt (0.02–0.05 mm), fine sand (0.05–0.2 mm) and coarse sand (0.2–2 mm) in g kg⁻¹ were determined according to the Robinson method (Robinson 1922) without previous decarbonation (Baize 2000).

Comparison of vegetation composition between 2001 and 2015

To investigate whether the vegetation composition of the fallow land had re-developed towards the vegetation of unaltered steppes over the previous 14 years, we compared old vegetation relevés from Römermann et al (2004, 2005) to recent vegetation relevés for the locations Peau de Meau (steppe, cereal fallow, melon fallow; Table 1), Petit Abondoux (steppe, cereal fallow; Table 1) and Valigne (steppe, melon fallow; Table 1). Ten replicate relevés were carried out per site in 2001 and five in 2015.

Trait data and community-weighted means (CWM)

We compiled trait data for SLA (the ratio of fresh leaf area to dry mass in g cm⁻²), LDMC (the ratio of dry mass to fresh mass in mg g⁻¹) and canopy height (cm) for all species observed in the vegetation relevés using the trait databases TRY (Kattge et al. 2011) and BROTE (Paula & Pausas 2013). This dataset was then supplemented with our own measurements for species missing from the databases. For our own measurements, five individuals per species were collected at the site where they were most abundant, following the approach described in Rosbakh et al. (2015). Plants were collected and measured from May to July 2015, following standardized protocols (Pérez-Harguindeguy et al. 2013). For each species, two healthy sun leaves were collected, wrapped in moist paper and stored in a plastic bag in a cool box until the fresh leaf weight was measured. The fresh leaf area was determined at 300 dpi on a flatbed scanner. Leaf area was measured on scanned fresh leaves with the Leaf Trait package for R

(Bernhardt-Römermann, unpubl.). Then, the leaves were dried at 70° C to constant mass and subsequently weighed. For each vegetation relevé, we then calculated the CWM for SLA, LDMC and canopy height according to:

Formula 1:

$$CWM_c = \sum_{i=1}^s p_{ic} \times T_{ic}$$

where the mean value of trait T for species i in the community c was weighted by species relative abundance p_{ic} in the community, with a total of S species (Garnier et al. 2004; Violle et al. 2007).

Species selection and trait measurements for intraspecific trait variability (ITV)

Species selection was based on vegetation relevés from 2001 (Römermann et al. 2004, 2005) and 2014 (Tanet 2014). The final selection was based on indicator species analysis conducted with the package ‘indicspecies’ (Cáceres & Legendre 2009) supplemented by expert knowledge. We identified three successful colonizers as those occurring originally in the steppe but also in the fallow land (*Carthamus lanatus* L., *Lobularia maritima* L. Desv., *Sideritis romana* L.). Five unsuccessful colonizers were limited to the steppe (*B. retusum*, *Hypochaeris glabra* L., *Linum trigynum* L., *Stipa capillata* L., *Trifolium campestre* Schreb.). Four species whose major distribution was in fallow land were added for comparison (*Crepis foetida* L., *Dactylis glomerata* L., *Galactites tomentosus* Moench., *Trifolium subterraneum* L.). We sampled a maximum of eight individuals per species and site if abundant: for details, see Table 2 and Table 3. Besides SLA, LDMC and canopy height, we expanded our trait set to include plant width (cm), aboveground biomass (g), seed mass (mg) and leaf carbon (leaf C in mg kg⁻¹), nitrogen (leaf N in mg kg⁻¹) and phosphorus content (leaf P in mg kg⁻¹; Table 3). Plant width was measured with a measuring tape at maximum diameter without extending the plant, biomass was dried at 70° C to constant mass and subsequently weighed. We determined average seed mass by weighing five batches of 20–100 ripe seeds per species, depending on the species. Seed

sampling was not possible for *B. retusum*, *H. glabra* and *S. capillata* as seed production was not sufficient in 2015. Due to insufficient plant material, leaf samples of the eight individuals were pooled per site prior to nutrients analyses. Thus, ITV could not be calculated thereof but leaf nutrients could be compared between the different land-use types (see Appendix S3).

To characterize the differences in trait composition between individuals measured in the steppe and the fallow land sites, a principle component analysis (PCA) was conducted (matrix: 854 individuals of 9 species x 9 traits; *H. glabra*, *B. retusum*, *T. subterraneum* were excluded because of data constraints – insufficient biomass and seeds available). Prior to analyses, data was standardized to zero mean and unit variance.

To characterize ITV, the coefficient of variation (CV) was calculated as the standard deviation divided by the mean for all measured traits (Pérez-Harguindeguy et al. 2013) based on the eight individuals per species and site. ITV was calculated for all species occurring in the steppe sites to assess whether species with higher trait variability were more likely to pass through the environmental filter and thus manage to re-establish in fallow land. Therefore, we included four unsuccessful colonizers three successful colonizers and two fallow land species in our CV calculation (Table 2).

Statistical analyses

Vegetation composition and soil properties

To test for the effect of land-use history on vegetation structure (S, H_s, E, standing crop, LAI, vegetation cover/stone cover/bare ground) and soil properties (clay, silt, sand, organic matter, total C, total N, C:N, pH, P, Ca, Mg and K), one-way ANOVA followed by a Tukey Honest Significant Difference post-hoc test was performed for normally distributed data with homogeneous variances. Any data not showing normal distribution and homogeneous variances were analyzed by the nonparametric Kruskal–Wallis test followed by the pairwise Wilcoxon test with *P*-value adjustment according to Holm.

To assess associations between differences in vegetation composition and environmental parameters, a detrended correspondence analysis (DCA) was conducted. We used species abundances of the five vegetation relevés per site as the main matrix (a total of 12 species x 16 sites). Then, environmental parameters were correlated with the axes using mean values calculated per site for each parameter. The environmental factors considered were five physical soil variables (clays, fine and coarse silt, fine and coarse sand) and eight chemical soil variables (total C, total N, C:N, pH, P, Ca, Mg and K) as well as percentage cover of vegetation, stones and bare ground and time since abandonment as deduced from the fallow land cultivation period (Table 1). All parameters were standardized to zero mean unit variance prior to analyses.

Comparison of vegetation composition between 2001 and 2015

As the 2001 data were obtained on 2m x 2m plots via the frequency method (Pfadenhauer et al. 1986), we harmonized the data for 2001 and 2015 into presence/ absence data to avoid any observer effects in the estimation of the frequencies and for the non-metric multidimensional scaling (NMDS). Species that occurred just once across the overall dataset were removed to avoid discriminating the dataset too strongly by their presence alone instead of discriminating the gradient by their frequencies.

Differences in trait composition: Community-weighted means (CWM) and intraspecific trait variability (ITV)

To analyze differences between CWM of the three land-use types and CVs, a one-way ANOVA followed by a Tukey Honest Significant Difference post-hoc test was performed for normally distributed data with homogeneous variances. Any data not showing normal distribution and homogeneous variances were analyzed by the nonparametric Kruskal–Wallis test followed by the pairwise Wilcoxon test with *P*-value adjustment according to Holm.

All statistical analyses were carried out in *R* (version 3.3.1, R Development Core Team 2015, Vienna, AT). Multivariate analyses (DCA, PCA, NMDS) were performed using the *R* packages ‘vegan’ (Oksanen et al. 2008) and ‘MASS’ (Venables & Ripley 2002).

Results

Vegetation and soil attributes

Vegetation structure and soil properties of fallow lands differed from those of adjacent steppe (Table). The steppe showed highest values in species richness, Shannon diversity and Evenness; cereal fallows typically had intermediate positions (Table 4). Melon fallows had significantly higher values in soil Ca, P and pH. The C/N ration was highest in steppes.

The DCA (Fig. 2) showed a trend towards very homogeneous vegetation composition in the steppe plots at different sites, whereas the fallow lands tended to be more heterogeneous in vegetation composition. The first axis separated steppe from fallow land and was negatively correlated with total soil C, vegetation cover and soil C:N ratio. Percentages of bare ground, clays, pH and P were positively correlated with fallow land.

Comparison of vegetation composition between 2001 and 2015

In 2001, steppe and fallow lands had distinct species composition trends indicated by the separated polygons of the NMDS (Fig. 3). Fourteen years later, the fallow land had developed towards the original steppe vegetation, while there had been little change in steppe vegetation composition. The steppe communities of both 2001 and 2015 were characterized by perennial species, such as the dominant steppe species *B. retusum* and *S. capillata* and the annual species *Aira cupaniana* Guss. and *Linum strictum* L.. Cereal fallows in 2001 commonly contained *Euphorbia falcata* L. and *Reseda phyteuma* L., whereas *Medicago monspeliaca* L. and *Poa bulbosa* L. were identified as late colonizers of these cereal fallows. Melon fallows in 2001 were characterized by *Bromus rubens* L., *Bromus intermedius* L., *Hordeum murinum* L. and *Rostraria cristata* (L.) Tzvelev. In 2015, these species had declined and *Plantago lagopus* L. and *Catapodium rigidum* (L.) C.E.Hubb. had become the typical melon fallow species. For details on species distribution see Fig. 3.

Differences in trait composition

As shown in Table 4, CWM of SLA was significantly lower in steppe than in fallow-land communities ($\chi^2 = 54.19$, $P < 0.001$) while CWM of LDMC was significantly higher in steppe communities ($F_{2,77} = 52.38$, $P < 0.001$). Canopy height differed among all the land-use types. The highest values were observed in the steppe and the lowest in cereal fallows ($F_{2,77} = 5.19$, $P < 0.01$). For individual species information see Appendix S1.

The PCA (Appendix S2) partly separated the species according to their trait composition measured in the steppe and the fallow land. *C. foetida*, *D. glomerata*, *L. trigynum* and *S. romana* showed a tendency towards a distinction between the steppe and fallow lands and thus changes in plant trait values due to land-use change, as indicated by the black arrow.

The CV did not reveal any differences between successful and unsuccessful colonizers (Fig. 4). Overall, significant differences between species were found for SLA ($\chi^2 = 27.44$, $P > 0.001$), LDMC ($\chi^2 = 31.75$, $P > 0.001$) and plant width ($\chi^2 = 36.91$, $P > 0.001$). *L. trigynum*, as a typical steppe species, varied more in SLA than the fallow land species *D. glomerata* and the successful colonizer *S. romana*. Regarding LDMC values, *L. trigynum* displayed the highest variability, while *B. retusum* and *D. glomerata* had significantly lower variability. *B. retusum* had a significantly lower CV in plant width than *L. trigynum* and *S. romana*. *S. romana* showed significantly higher variation in plant width than *C. lanatus* (successful colonizer).

Discussion

The recovery of fallow land in our study area implemented grazing and was thus not a strictly passive restoration technique. The fact that more than 35 years after cultivation has ceased the studied vegetation did not fully transform to steppe calls for more active intervention, like seeding and planting. Notwithstanding these results, the comparison of vegetation relevés from 2001 with results for 2015 indicated a (though slow) conversion of fallow land towards steppe vegetation. Furthermore, there was no particular similarity in trait composition between steppe

species that successfully re-colonized fallow lands and the fallow-land community, and no evidence that intraspecific trait variability is associated with re-colonization success of these species.

Effect of former cultivation on vegetation attributes and soil parameters

Former cultivation practices still influence present vegetation attributes. The species composition of recently abandoned cereal fallows can be distinguished from the species composition of the steppe. However, in the long-abandoned cereal fallow, there is a visible vegetation development towards the original steppe community.

Our results show that the dominant perennial steppe grass *B. retusum* has so far been unable to re-establish on the fallow land; this is not only directly explained by habitat change due to deep soil ploughing and high fertilization, but also indirectly by competition with the common fallow land species (Buisson et al. 2015; Coiffait et al. 2012). Moreover, dispersal limitation and the mainly vegetative reproduction of this steppe grass (Dureau & Bonnefon 1998) have prevented successful re-establishment. The decline of C:N ratios on the fallow land can be explained by organic matter mineralization and the absence of restitution from perennial grass species unable to re-colonize fallow land (Römermann et al. 2004, 2005).

The high percentage of bare ground is characteristic of fallow land, where stones were removed for cultivation purposes (Buisson & Dutoit 2006). However, stones create important micro-habitats for the typical steppe vegetation: they preserve humidity, serve as shelter and create safe sites for rhizomes of e.g., *B. retusum* (Bourrelly 1984; Caturla et al. 2000; Buisson et al. 2015).

According to Janssens et al. (1998), plant diversity is limited by high soil fertility, especially by plant-available phosphorus. The fallow land remains distinguished from the steppe by its elevated phosphorus and potassium content due to former fertilization, as also observed by Gough and Marrs (1990) on abandoned fields in the UK. Because of different cultivation regime intensities, i.e., deep ploughing, fertilization, the application of pesticides in melon-cultivated

areas (Masip 1991) versus less intensive cereal cultivation, phosphorus content was higher in melon fallows than in cereal fallows. Deep ploughing for melon cultivation destroyed the upper part of the geological conglomerate 40 cm deep in the soil, bringing some pieces of the matrix (composed of CaCO_3) to the surface. Hence, Ca content increased and higher pH was measured. The effect of ploughing also appears in the physical soil parameters. As a result of soil disturbance through ploughing, clays from the steppe subsoil layer (20–40 cm) were transported to the topsoil, leading to a higher clay content for the fallow land (10–20 cm; Duclos 1994). The opposite effect is observed for the coarse silt of the steppe, which was buried in the subsoil as a result of ploughing (Bulot et al. 2017; Chenot et al. 2017).

Öster et al. (2009) found that in semi-natural grassland communities in Sweden, species richness and similarity were still lower in fallow lands more than 50 years after cultivation. Differences in plant attributes between calcareous grassland and fallow land after 60–100 years were also shown by Fagan et al. (2008), Karlik & Poschlod (2009) and Forey and Dutoit (2012). Remnants of fertilization persist in the soil of fallow land for many years and still affect plant communities and the vegetation structure decades after abandonment (Willems et al. 1993; Austrheim & Olsson 1999; Lepš 1999).

Comparison of vegetation structure between 2001 and 2015

The resurvey of the research site in 2015 shows a trajectory of the former cultivated areas towards the original steppe community, even though differences in vegetation and soil structure still exist. Although common in 2001, ruderal species *H. murinum* and *R. cristata* were not recorded on melon fallows in 2015. Similarly, *Bromus* species- typical species in the first year of abandonment- were less common on fallow land in 2015. In cereal fallows, not only weed species but also some typical steppe species had established by 2015. The steppe vegetation did not change over time because the disturbance regime (i.e., extensive grazing) remained stable between 2001 and 2015 (Sauguet 2013). The more rapid development from cereal fallow vegetation towards steppe vegetation was probably due to the fact that less fertilizer was applied

during cultivation and that there was no deep ploughing (Römermann et al. 2005, Buisson et al. 2006).

Effect of former cultivation on plant functional traits

This study confirms the relevance of the leaf traits SLA and LDMC at community level to explain the response of ecosystems to land-use change, as proposed by previous studies (Díaz et al. 2007; de Bello et al. 2010; Vandewalle et al. 2010). The steppe community was characterized by smaller and denser leaves, typical of plants following a conservative strategy (Grime 1973; Saatkamp et al. 2010; Pérez-Harguindeguy et al. 2013). Our study demonstrates that the fallow land is still a nutrient-rich environment favoring communities with high SLA and low LDMC. This is in line with findings demonstrating the association of low SLA with efficient nutrient conservation strategies, nutrient-poor habitats, slow growth rates and high longevity, as outlined above (Poorter & Remkes 1990; Garnier 1992; Chapin et al. 1993; Poorter & Garnier 1999; Lavorel & Garnier 2002; Pérez-Harguindeguy et al. 2013). Our analyses showing that the cereal fallows occupy an intermediate position with respect to trait composition also reveal that type of cultivation plays an important role in re-development towards the unaltered steppe vegetation. Plant height was greater in steppe communities than in fallow land, contradicting our initial hypothesis. However, it is in line with findings from Australian ecosystems (Falster & Westoby 2003) that showed positive relationships between plant height and LAI. Our finding of higher plants in the steppe may be related to the higher fodder quality in nutrient-rich fallow land, which could lead to preferential grazing by herbivores, reducing plant height (Falster & Westoby 2003; Saatkamp et al. 2010). This differential selection by herbivores of recently impacted sites due to their higher fodder quality could also have implications for restoration (Yates et al. 2000). Tests of plant and soil mineral content of our data revealed a significantly positive correlation between soil phosphorus and leaf phosphorus content ($r = 0.67$, $P < 0.001$), soil nitrogen and leaf nitrogen content ($r = 0.35$, $P < 0.02$, analyses not shown).

We found no evidence of particular similarity between the trait values of successful colonizers and those of the species typically found in fallow-land communities. We thus have to reject the idea that colonizers simply “fit into” the assembly of recent fallow-land communities with initial assembly conditions of high nutrient levels and recently-moved soil. Alternatively, the data can be understood as indicating that successful colonizers exploit the diverging conditions created during longer grazing periods with decreased nutrient status, and that these colonizers may be better adapted to grazing than typical fallow-land species (Saatkamp et al. 2010). However, our species pool was quite small with differences in sample size, and further studies should consider a more extensive species set covering more families.

Furthermore, our data show higher functional variability to be highly species- and trait-specific, with no clear pattern revealed for the trait values of successful or unsuccessful colonizers. This supports the view that plants in fallow lands are assembled according to their adaptability to specific grazed fallow-land conditions, rather than reflecting a wide intraspecific trait variability that ensures their survival in both fallow-land and steppe communities. Hence, our study suggests that plants with specifically high ITV are not favored in the assembly process after abandonment and during grazing.

Conclusions and implications for management

Our study demonstrates that natural recovery of abandoned fallow land, notwithstanding its close proximity to the target vegetation, can take more than 35 years, even when the traditional grazing system is re-introduced just after cultivation abandonment. Regular monitoring of further vegetation development in the abandoned fallow land would enable the potential for natural recovery after a period of cultural land-use to be assessed.

In addition to monitoring natural recovery over the long term, new grazing systems could be introduced to hasten recovery or more active restoration implemented by seeding nurse or structuring species (e.g. *B. retusum*; Coiffait-Gombault et al., 2012; Buisson et al., 2015). Future analyses could cover an extended set of species and include further traits to better

elucidate assembly processes. In particular, they should consider traits closely related to soil water conditions, such as leaf venation traits and water-use efficiency, and traits related to soil stoichiometry, such as root traits, mycorrhizal type (Jaunatre et al. 2016) and seed nutrient content. Future projects need to identify project-specific restoration targets in order to implement a trait-based approach into their restoration protocol.

Acknowledgments

We would like to thank the *Conservatoire d'espaces naturels de Provence-Alpes-Côte d'Azur* (CEN PACA) and the *Réserve Naturelle Nationale des Coussouls de Crau* for their support during field work. Thanks to Julie Tanet and Adeline Bulot working at the Mediterranean Institute of Biodiversity and Ecology of Marseille for realizing the botanical relevés in 2014. We acknowledge valuable comments of Karen Russel, Viktoria Wagner, Anna Johnson and an anonymous referee. This study was supported by the TRY initiative on plant traits (<http://www.trydb.org>) hosted, developed and maintained by J. Kattge and G. Bönisch (Max-Planck-Institute for Biogeochemistry, Jena, Germany). TRY is/was supported by DIVERSITAS, IGBP, the Global Land Project, the UK Natural Environment Research Council (NERC) through its program QUEST (Quantifying and Understanding the Earth System), the French Foundation for Biodiversity Research (FRB), and GIS "Climat, Environnement et Société" France. CR acknowledges financial support by the German Research Foundation (project RO 3842/3-1). The authors declare no conflict of interest.

References

- Albert, C.H., Grassein, F., Schurr, F.M., Vieilledent, G. & Violle, C. 2011. When and how should intraspecific variability be considered in trait-based plant ecology? *Perspectives in Plant Ecology, Evolution and Systematics* 13: 217-225.
- Austrheim, G. & Olsson, E.G.A. 1999. How does continuity in grassland management after ploughing affect plant community patterns? *Plant Ecology* 145: 59-74.

- 474 Baize, D. 2000. *Guide des analyses en pédologie: 2e édition, revue et augmentée*. Editions Quae, FR.
- 475 Bucher, S.F., König, P., Menzel, A., Migliavacca, M., Ewald, J. & Römermann, C. 2018. Traits and
476 climate are associated with first flowering day in herbaceous species along elevational gradients.
477 *Ecology and Evolution* 8: 1147-1158.
- 478 Bourrelly, M. 1984. *Contribution à l'étude écologique de la Crau (Bouches-du-Rhône). Caractéristiques*
479 *floristiques, dynamique annuelle, production primaire de la Crau steppique*. Ph.D. thesis, Aix-
480 Marseille University, Marseille, FR.
- 481 Braun-Blanquet, J. 1964. *Pflanzensoziologie : Grundzüge der Vegetationskunde*, Springer-Verlag, Wien.
482 AT.
- 483 Breitschwerdt, E., Jandt, U. & Bruehlheide, H. 2015. Do newcomers stick to the rules of the residents?
484 Designing trait-based community assembly tests. *Journal of Vegetation Science* 26: 219-232.
- 485 Bucher, S. F., Auerswald, K., Grün-Wenzel, C., Higgins, S. I., Garcia Jorge, J. & Römermann, C. 2017.
486 Stomatal traits relate to habitat preferences of herbaceous species in a temperate climate. *Flora*
487 229: 107-115.
- 488 Buisson, E., Corcket, E. & Dutoit, T. 2015. Limiting processes for perennial plant reintroduction to
489 restore dry grasslands. *Restoration Ecology* 23: 947-954.
- 490 Buisson, E. & Dutoit, T. 2004. Colonisation by native species of abandoned farmland adjacent to a
491 remnant patch of Mediterranean steppe. *Plant Ecology* 174: 371-384.
- 492 Buisson, E. & Dutoit, T. 2006. Creation of the natural reserve of La Crau: implications for the creation
493 and management of protected areas. *Journal of Environmental Management* 80: 318-326.
- 494 Buisson, E., Dutoit, T., Torre, F., Römermann, C. & Poschlod, P. 2006. The implications of seed rain
495 and seed bank patterns for plant succession at the edges of abandoned fields in Mediterranean
496 landscapes. *Agriculture, Ecosystems & Environment* 115: 6-14.
- 497 Bulot, A., Potard, K., Bureau, F., Bérard, A. & Dutoit, T. 2017. Ecological restoration by soil transfer:
498 impacts on restored soil profiles and topsoil functions. *Restoration Ecology* 25: 354-366.
- 499 Cáceres, M.D. & Legendre, P. 2009. Associations between species and groups of sites: indices and
500 statistical inference. *Ecology* 90: 3566-3574.
- 501 Caturla, R.N., Raventós, J., Guàrdia, R. & Vallejo, V.R. 2000. Early post-fire regeneration dynamics of
502 *Brachypodium retusum* Pers.(Beauv.) in old fields of the Valencia region (eastern Spain). *Acta*
503 *Oecologica* 21: 1-12.
- 504 Chapin, F.S., Autumn, K. & Pugnaire, F. 1993. Evolution of suites of traits in response to environmental
505 stress. *American Naturalist*: S78-S92.
- 506 Chenot, J., Jaunatre, R., Buisson, E. & Dutoit, T. 2017. Long-term effects of topsoil transfer assessed
507 thirty years after rehabilitation of dry alluvial quarries in Southeastern France. *Ecological*
508 *Engineering* 99: 1-12.
- 509 Coiffait, C., Buisson, E. & Dutoit, T. 2012. Using a two-phase sowing approach in restoration – sowing
510 foundation species to restore and subordinate species to evaluate restoration success. *Applied*
511 *Vegetation Science* 15: 277-289.
- 512 Colomb, E. & Roux, R. 1978. La Crau, données nouvelles et interprétations. *Géologie méditerranéenne*
513 5: 303-324.
- 514 Cornwell, W.K. & Ackerly, D.D. 2009. Community assembly and shifts in plant trait distributions across
515 an environmental gradient in coastal California. *Ecological Monographs* 79: 109-126.

516 de Bello, F., Lavorel, S., Díaz, S., Harrington, R., Cornelissen, J.H., Bardgett, R.D., Berg, M.P., Cipriotti,
517 P., Feld, C.K. & Hering, D. 2010. Towards an assessment of multiple ecosystem processes and
518 services via functional traits. *Biodiversity and Conservation* 19: 2873-2893.

519 Devaux, J., Archiloque, A., Borel, L., Bourrelly, M. & Louis-Palluel, J. 1983. Notice de la carte phyto-
520 écologique de la Crau (Bouches du Rhône). *Biologie et écologie méditerranéenne* 10: 5-54.

521 Deverre, C. 1996. La nature mise au propre dans la steppe de Crau et la forêt du Var. *Études rurales*
522 141-142: 45-61.

523 Diaz, S., Cabido, M. & Casanoves, F. 1998. Plant functional traits and environmental filters at a regional
524 scale. *Journal of Vegetation Science* 9: 113-122.

525 Dureau, R. & Bonnefon, O. 1998. Etude des pratiques de gestion pastorale des coussouls. *Patrimoine*
526 *nature et pratiques pastorales en Crau*. Ed. CEEP Ecomusée de Crau: 61-89.

527 Dutoit, T., Buisson, E., Fadda, S., Henry, F., Coiffait-Gombault, C., Jaunatre, R., Alignan, J.F., Masson,
528 S. & Bulot, A. 2013. The pseudo-steppe of La Crau (South-Eastern France): origin, management
529 and restoration of a Mediterranean rangeland. In: Traba J. Morales M (eds.) *Steppe Ecosystems:*
530 *Biological Diversity, Management and Restoration*, pp. 287-301. Nova Science Publishers,
531 Hauppauge, New York, US.

532 Engst, K., Baasch, A., Erfmeier, A., Jandt, U., May, K., Schmiede, R., & Bruelheide, H. 2016.
533 Functional community ecology meets restoration ecology: Assessing the restoration success of
534 alluvial floodplain meadows with functional traits. *Journal of Applied Ecology* 53: 751-764.

535 Etienne, M., Aronson, J. & Le Floc'h, E. 1998. Abandoned lands and land use conflicts in southern
536 France. Piloting ecosystem trajectories and redesigning outmoded landscapes in the 21st century.
537 In: Rundel P.W., Montenegro G. and Jaksic F.M. (eds.) *Landscape disturbance and biodiversity*
538 *in Mediterranean-type ecosystems*, pp. 127-140. Ecological Studies, Springer-Verlag, Berlin,
539 DE.

540 Fagan, K.C., Pywell, R.F., Bullock, J.M. & Marrs, R.H. 2008. Do restored calcareous grasslands on
541 former arable fields resemble ancient targets? The effect of time, methods and environment on
542 outcomes. *Journal of Applied Ecology* 45: 1293-1303.

543 Falster, D.S. & Westoby, M. 2003. Plant height and evolutionary games. *Trends in Ecology & Evolution*
544 18: 337-343.

545 Fischer, S.F., Poschlod, P. & Beinlich, B. 1996. Experimental studies on the dispersal of plants and
546 animals on sheep in calcareous grasslands. *Journal of Applied Ecology*: 1206-1222.

547 Forey, E. & Dutoit, T. 2012. Vegetation, soils and seed banks of limestone grasslands are still impacted
548 by former cultivation one century after abandonment. *Community Ecology* 13: 194-202.

549 Garnier, E. 1992. Growth analysis of congeneric annual and perennial grass species. *Journal of Ecology*
550 80: 665-675.

551 Garnier, E., Cortez, J., Billès, G., Navas, M.-L., Roumet, C., Debussche, M., Laurent, G., Blanchard,
552 A., Aubry, D. & Bellmann, A. 2004. Plant functional markers capture ecosystem properties
553 during secondary succession. *Ecology* 85: 2630-2637.

554 Gaudet, C.L. & Keddy, P.A. 1988. A comparative approach to predicting competitive ability from plant
555 traits. *Nature* 334: 242-243.

556 Gough, M. & Marrs, R. 1990. A comparison of soil fertility between semi-natural and agricultural plant
557 communities: Implications for the creations of species-rich grassland on abandoned agricultural
558 land. *Biological Conservation* 51: 83-96.

- 559 Grime, J.P. 1973. Competitive exclusion in herbaceous vegetation. *Nature, UK* 242: 344-347.
- 560 Jäger, E.J. & Werner, K. 2002. Rothmaler, Exkursionsflora von Deutschland. *Gefäßpflanzen: Kritischer*
561 *Band 9*. Springer Spektrum, Berlin Heidelberg, DE.
- 562 Janssens, F., Peeters, A., Tallowin, J., Bakker, J., Bekker, R., Fillat, F. & Oomes, M. 1998. Relationship
563 between soil chemical factors and grassland diversity. *Plant and Soil* 202: 69-78.
- 564 Jaunatre, R., Fonvieille, N., Spiegelberger, T., Buisson, E., & Dutoit T. 2016. Recovery of arbuscular
565 mycorrhizal fungi root colonization after severe anthropogenic disturbance: four species
566 assessed in old-growth Mediterranean grassland. *Folia Geobotanica* 51: 319-332.
- 567 Jung, V., Violle, C., Mondy, C., Hoffmann, L. & Muller, S. 2010. Intraspecific variability and trait-based
568 community assembly. *Journal of Ecology* 98: 1134-1140.
- 569 Karlik, P. & Poschlod, P. 2009. History or abiotic filter: which is more important in determining the
570 species composition of calcareous grasslands? *Preslia* 81: 321-340.
- 571 Kattge, J., Diaz, S., Lavorel, S., Prentice, I., Leadley, P., Bönisch, G., Garnier, E., Westoby, M., Reich,
572 P.B., (...) & Wright, I. 2011. TRY—a global database of plant traits. *Global Change Biology* 17:
573 2905-2935.
- 574 Keddy, P.A. 1992. Assembly and response rules: two goals for predictive community ecology. *Journal*
575 *of Vegetation Science* 3: 157-164.
- 576 Kiehl, K., Kirmer, A., Donath, T. W., Rasran, L., & Hölzel, N. 2010. Species introduction in restoration
577 projects—Evaluation of different techniques for the establishment of semi-natural grasslands in
578 Central and Northwestern Europe. *Basic and Applied Ecology* 11: 285-299.
- 579 Kuznetsova, A., Brockhoff, P.B. & Christensen, R.H.B. 2015. *lmerTest: Tests in Linear Mixed Effects*
580 *Models*. R package version 2.0-29.
- 581 Laforest-Lapointe, I., Martínez-Vilalta, J. & Retana, J. 2014. Intraspecific variability in functional traits
582 matters: case study of Scots pine. *Oecologia* 175: 1337-1348.
- 583 Laughlin, D.C. 2014. Applying trait-based models to achieve functional targets for theory-driven
584 ecological restoration. *Ecology Letters* 17: 771-784.
- 585 Laughlin, D.C., Joshi, C., Bodegom, P.M., Bastow, Z.A. & Fulé, P.Z. 2012. A predictive model of
586 community assembly that incorporates intraspecific trait variation. *Ecology Letters* 15: 1291-
587 1299.
- 588 Lauterbach, D., Römermann, C., Jeltsch, F. & Ristow, M. 2013. Factors driving plant rarity in dry
589 grasslands on different spatial scales: a functional trait approach. *Biodiversity and Conservation*
590 22: 2337-2353.
- 591 Lavorel, S. & Garnier, E. 2002. Predicting changes in community composition and ecosystem
592 functioning from plant traits: revisiting the Holy Grail. *Functional ecology* 16: 545-556.
- 593 Lenth, R.V. 2016. *lsmeans: Least-Squares Means*. R Package version 2.22.
- 594 Lepš, J. 1999. Nutrient status, disturbance and competition: an experimental test of relationships in a
595 wet meadow. *Journal of Vegetation Science* 10: 219-230.
- 596 MacArthur, R. & Levins, R. 1967. The limiting similarity, convergence, and divergence of coexisting
597 species. *American Naturalist* 101: 377-385.
- 598 Masip, A. 1991. *Le peuplement végétal de la Réserve de Peau de Meau. Données pour la gestion*. Ph.D.
599 thesis, University of Barcelona. Barcelona, ES.

- Metson, A.J. 1957. Methods of chemical analysis for soil survey samples. *Soil Science* 83: 245.
- Molliex, S., Siame, L.L., Bourlès, D.L., Bellier, O., Braucher, R. & Clauzon, G. 2013. Quaternary evolution of a large alluvial fan in a periglacial setting (Crau Plain, SE France) constrained by terrestrial cosmogenic nuclide (^{10}Be). *Geomorphology* 195: 45-52.
- Mühlenberg, M. 1993. *Freilandökologie*. UTB für Wissenschaft, 3rd ed. Quelle & Meyer, Heidelberg, Wiesbaden, DE.
- Nicotra, A.B., Atkin, O.K., Bonser, S.P., Davidson, A.M., Finnegan, E., Mathesius, U., Poot, P., Purugganan, M.D., Richards, C. & Valladares, F. 2010. Plant phenotypic plasticity in a changing climate. *Trends in Plant Science* 15: 684-692.
- Oksanen, J., Kindt, R., Legendre, P., O'Hara, B., Stevens, M.H.H., Oksanen, M.J. & Suggests, M. 2007. *Vegan: Community ecology package*. R package version 2.0-10.
- Olsen, S.R. 1954. *Estimation of available phosphorus in soils by extraction with sodium bicarbonate*. United States Department of Agriculture, Washington, DC, US.
- Öster, M., Ask, K., Cousins, S.A. & Eriksson, O. 2009. Dispersal and establishment limitation reduces the potential for successful restoration of semi-natural grassland communities on former arable fields. *Journal of Applied Ecology* 46: 1266-1274.
- Paula, S. & Pausas, J. 2013. *BROT: a plant trait database for Mediterranean Basin species*. Version 2013.06. [http:// www.uv.es/jgpausas/brot.htm](http://www.uv.es/jgpausas/brot.htm).
- Pavon, D. 2014. Mémento pour l'aide à la détermination de la flore vasculaire du département des Bouches-du-Rhône. *Bulletin de la Société linnéenne de Provence*. Numéro spécial 18.
- Pérez-Harguindeguy, N., Díaz, S., Garnier, E., Lavorel, S., Poorter, H., Jaureguiberry, P., Bret-Harte, M., Cornwell, W., Craine, J., (...) & Cornelissen, J.H.C. 2013. New handbook for standardised measurement of plant functional traits worldwide. *Australian Journal of Botany* 61: 167-234.
- Pfadenhauer, J., Poschlod, P. & Buchwald, R. 1986. Überlegungen zu einem Konzept geobotanischer Dauerbeobachtungsflächen für Bayern. Teil 1. Methodik der Anlage und Aufnahme. *Bayerische Akademie für Naturschutz und Landschaftspflege* 10: 41-60.
- Poorter, H. & Garnier, E. 1999. Ecological significance of inherent variation in relative growth rate and its components. *Handbook of functional Plant Ecology* 20: 81-120.
- Poorter, H., Niinemets, Ü., Poorter, L., Wright, I.J. & Villar, R. 2009. Causes and consequences of variation in leaf mass per area (LMA): a meta-analysis. *New Phytologist* 182: 565-588.
- Poorter, H. & Remkes, C. 1990. Leaf area ratio and net assimilation rate of 24 wild species differing in relative growth rate. *Oecologia* 83: 553-559.
- Poschlod, P., Kiefer, S., Tränkle, U., Fischer, S. & Bonn, S. 1998. Plant species richness in calcareous grasslands as affected by dispersability in space and time. *Applied Vegetation Science* 1: 75-91.
- Reich, P.B., Walters, M.B. & Ellsworth, D.S. 1997. From tropics to tundra: global convergence in plant functioning. *Proceedings of the National Academy of Sciences* 94: 13730-13734.
- Robin, V., Nelle, O., Talon T., Poschlod, P., Schwartz, D., Bal M.C., Allée, P., Vernet, J.L. & Dutoit T. 2018. A comparative review of soil charcoal data: spatiotemporal patterns of origin and long-term dynamics of Western European nutrient-poor grasslands. *The Holocene* (in press)
- Römermann, C., Bernhardt-Römermann, M., Dutoit, T., Poschlod, P. & Rolando, C. 2004. Histoire culturelle de la Crau: potentialités de ré-établissement des espèces caractéristiques du coussous après abandon. *Ecologia Mediterranea* 30: 47-70.

- 642 Römermann, C., Dutoit, T., Poschlod, P. & Buisson, E. 2005. Influence of former cultivation on the
643 unique Mediterranean steppe of France and consequences for conservation management.
644 *Biological Conservation* 121: 21-33.
- 645 Römermann, C., Bernhardt-Römermann, M., Kleyer, M. & Poschlod, P. 2009. Substitutes for grazing
646 in semi-natural grasslands – do mowing or mulching represent valuable alternatives to maintain
647 vegetation structure? *Journal of Vegetation Science* 20: 1086-1098.
- 648 Römermann, C., Bucher, S. F., Hahn, M. & Bernhardt-Römermann, M. 2016. Plant functional traits -
649 fixed facts or variable depending on the season? *Folia Geobotanica* 51: 143-159.
- 650 Robinson, G. (1922). A new method for the mechanical analysis of soils and other dispersions. *The*
651 *Journal of Agricultural Science*, 12(3), 306-321.
- 652 Rosbakh, S., Römermann, C. & Poschlod, P. 2015. Specific leaf area correlates with temperature: new
653 evidence of trait variation at the population, species and community levels. *Alpine Botany* 125:
654 79-86.
- 655 Saatkamp, A., Römermann, C. & Dutoit, T. 2010. Plant functional traits show non-linear response to
656 grazing. *Folia Geobotanica*, 45: 239-252.
- 657 Sauguet, F. 2013. L'élevage ovin en Crau: situation économique, enjeux et perspectives. In: *Écologie et*
658 *conservation d'une steppe méditerranéenne. La plaine de Crau*, pp. 192-205. Editions Quae,
659 FR.
- 660 Tanet, J. 2014. *Les fourmis sont-elles de bons indicateurs de la régénération spontanée des pelouses*
661 *sèches méditerranéennes après dégradation culturelle?* Master Thesis. Aix-Marseille University,
662 Marseille, FR.
- 663 Vandewalle, M., De Bello, F., Berg, M.P., Bolger, T., Dolédec, S., Dubs, F., Feld, C.K., Harrington, R.,
664 Harrison, P.A., (...) & Woodcock, B. A.2010. Functional traits as indicators of biodiversity
665 response to land use changes across ecosystems and organisms. *Biodiversity and Conservation*
666 19: 2921-2947.
- 667 Venables, W. & Ripley, B. 2002. *Modern Applied Statistics with S*, 4th ed. Springer-Verlag, New York,
668 USA.
- 669 Venn, S.E., Green, K., Pickering, C.M. & Morgan, J.W. 2011. Using plant functional traits to explain
670 community composition across a strong environmental filter in Australian alpine snowpatches.
671 *Plant Ecology* 212: 1491-1499.
- 672 Villar, R., Maranon, T., Quero, J.L., Panadero, P., Arenas, F. & Lambers, H. 2005. Variation in relative
673 growth rate of 20 *Aegilops* species (Poaceae) in the field: the importance of net assimilation rate
674 or specific leaf area depends on the time scale. *Plant and Soil* 272: 11-27.
- 675 Violle, C., Navas, M.L., Vile, D., Kazakou, E., Fortunel, C., Hummel, I. & Garnier, E. 2007. Let the
676 concept of trait be functional! *Oikos* 116: 882-892.
- 677 WallisDeVries, M.F., Poschlod, P. & Willems, J.H. 2002. Challenges for the conservation of calcareous
678 grasslands in northwestern Europe: integrating the requirements of flora and fauna. *Biological*
679 *Conservation* 104: 265-273.
- 680 Watson, D.J. 1947. Comparative physiological studies on the growth of field crops: I. Variation in net
681 assimilation rate and leaf area between species and varieties, and within and between years.
682 *Annals of Botany* 11: 41-76.
- 683 Weiher, E. & Keddy, P.A. 1995. Assembly rules, null models, and trait dispersion: new questions from
684 old patterns. *Oikos* 74: 159-164.

- Weiher, E., Werf, A., Thompson, K., Roderick, M., Garnier, E. & Eriksson, O. 1999. Challenging Theophrastus: a common core list of plant traits for functional ecology. *Journal of Vegetation Science* 10: 609-620.
- Westoby, M. 1998. A leaf-height-seed (LHS) plant ecology strategy scheme. *Plant and Soil* 199: 213-227.
- Westoby, M., Falster, D.S., Moles, A.T., Vesk, P.A. & Wright, I.J. 2002. Plant ecological strategies: some leading dimensions of variation between species. *Annual review of Ecology and Systematics* 33: 125-159.
- White, R.P., Murray, S., Rohweder, M., Prince, S. & Thompson, K. 2000. *Grassland ecosystems*. World Resources Institute, Washington, DC, US.
- Willems, J. 1983. Species composition and above ground phytomass in chalk grassland with different management. *Vegetatio* 52: 171-180.
- Willems, J., Peet, R.K. & Bik, L. 1993. Changes in chalk-grassland structure and species richness resulting from selective nutrient additions. *Journal of Vegetation Science* 4: 203-212.
- Wilson, J.B. 2007. Trait-divergence assembly rules have been demonstrated: Limiting similarity lives! A reply to Grime. *Journal of Vegetation Science* 18: 451-452.
- Yates, C.J., Norton, D.A. & Hobbs, R.J. 2000. Grazing effects on plant cover, soil and microclimate in fragmented woodlands in south-western Australia: implications for restoration. *Austral Ecology* 25: 36-47.
- Yulin, L., Johnson, D.A., Yongzhong, S., Jianyuan, C. & Zhang, T. 2005. Specific leaf area and leaf dry matter content of plants growing in sand dunes. *Botanical Bulletin of Academia Sinica* 46: 127-134.

Supporting Information

Appendix S1. Trait values of specific leaf area (SLA), leaf dry matter content (LDMC) and canopy height for all species.

Appendix S2. Principal component analysis (PCA) of the species–trait matrix (9 species × 9 traits).

Appendix S3. Leaf nutrient content of carbon (C), nitrogen (N) and phosphorus (P) compared between original steppe and fallow lands.

Table 1. Overview of the land-use history of the study sites in La Crau studied in 2015. *Comparison between 2001 and 2015.

Location	Study sites	Land-use type	Abbreviation	Cultivation period
Peau de Meau	steppe	-	PM_CSa*	-
	steppe	-	PM_CSb	-
	steppe	-	PM_CSc	-
	cereal fallow	melons, cereals	PM_MCa*	1974–1977
	cereal fallow	melons, cereals	PM_MCb	1969–1970
	melon fallow	melons	PM_ME*	1965–1968, 1984
Petit Abondoux	steppe	-	PA_CS	-
	cereal fallow*	melons, cereals	PA_MC*	1979–1984
Grand Abondoux	steppe	-	GA_CS	-
	melon fallow	melons	GA_ME	1980–1981
Valigne	steppe	-	VA_CS*	-
	melon fallow	melons	VA_ME*	1978, 1984
Poulagère	steppe	-	PO_CS	-
	old cereal fallow	cereals	PO_OC	< 1900
Couliès	steppe	-	CO_CS	-
	recent cereal fallow	cereals	CO_RC	1994, 2000

*comparison between 2001 and 2015

Table 2. List of the 12 species, classified into unsuccessful colonizers, fallow-land species and successful colonizers, on which trait measurements were performed on the study sites (Table 1) in 2015.

Species group	Species	Abbreviation	Family	Individuals sampled per species
Unsuccessful colonizers	<i>Brachypodium retusum</i> (Pers.) P. Beauv.	Bra_ret	Poaceaea	64
	<i>Hypochaeris glabra</i> L.	Hyp_gla	Asteraceae	15
	<i>Linum trigynum</i> L.	Lin_tri	Linaceae	120
	<i>Stipa capillata</i> L.	Sti_cap	Poaceae	110
	<i>Trifolium campestre</i> Schreb.	Tri_cam	Fabaceae	59
Fallow-land species	<i>Crepis foetida</i> L.	Cre_foe	Asteraceae	128
	<i>Dactylis glomerata</i> L.	Dac_glo	Poaceae	128
	<i>Galactites tomentosus</i> Moench.	Gal_tom	Asteraceae	52
	<i>Trifolium subterraneum</i> L.	Tri_sub	Fabaceae	16
Successful colonizers	<i>Carthamus lanatus</i> L.	Car_lan	Asteraceae	125
	<i>Lobularia maritima</i> L. Desv.	Lob_mar	Brassicaceae	128
	<i>Sideritis romana</i> L.	Sid_rom	Lamiaceae	128

Table 3. Plant functional traits measured on 12 focal species (Table 2). *These plant traits are not part of the main study, but available in Supporting Information (Appendix S2, S3).

Organ	Plant trait	Abbreviation	Unit	Measurements per site
Whole plant	Canopy height		cm	8 individuals
	Plant width		cm	8 individuals
	Aboveground biomass		g	8 individuals
Leaf	Specific leaf area	SLA	g cm ⁻²	8 individuals
	Leaf dry matter content	LDMC	mg g ⁻¹	8 individuals
	Leaf carbon content*	Leaf C	mg g ⁻¹	pooled
	Leaf nitrogen content*	Leaf N	mg g ⁻¹	pooled
	Leaf phosphorus content*	Leaf P	mg g ⁻¹	pooled
Seed	Seed mass*		mg	5 individuals

Table 4. Summary of mean $\pm$ standard error (SE) of the measured vegetation and soil parameters and community-weighted means (CWM) for the three land-use types. The statistical test values are given. Different letters indicate significant differences between the three land-use types as extracted from pair-wise tests. *** $P < 0.001$; ** $P < 0.01$; * $P < 0.05$; n.s. $P > 0.05$

		Steppe	Cereal	Melon	Statistical test values
		Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE	
Vegetation	Species richness	35.3 $\pm$ 0.1 a	29.0 $\pm$ 1.6 b	22.7 $\pm$ 1.5 c	$\chi^2 = 27.5^{***}$
	Shannon diversity index	2.6 $\pm$ 0.06 a	2.3 $\pm$ 0.1 a	1.9 $\pm$ 0.09 b	$\chi^2 = 24.7^{***}$
	Evenness	0.7 $\pm$ 0.01 a	0.7 $\pm$ 0.02 a	0.6 $\pm$ 0.02 b	$\chi^2 = 15.9^{***}$
	Standing crop	42.8 $\pm$ 1.2 a	17.9 $\pm$ 0.7 b	18.7 $\pm$ 0.8 b	F = 289.5 ^{***}
	Leaf area index	3.0 $\pm$ 0.1 a	1.3 $\pm$ 0.2 b	1.0 $\pm$ 0.1 b	F = 57.2 ^{***}
	Vegetation cover	60.3 $\pm$ 1.9 a	50.2 $\pm$ 3.2 b	43.7 $\pm$ 3.4 b	F = 10.2 ^{***}
	Stone cover	46.5 $\pm$ 1.2 a	36.6 $\pm$ 2.5 b	45.0 $\pm$ 2.0 a	F = 10.7 ^{***}
	Bare ground	9.9 $\pm$ 0.7 b	16.2 $\pm$ 1.2 a	18.8 $\pm$ 2.5 a	F = 13.6 ^{***}
Soil	Clays	192.7 $\pm$ 2.0 c	202.2 $\pm$ 3.4 b	213.4 $\pm$ 2.6 a	F = 20.0 ^{***}
	Fine silt	192.1 $\pm$ 1.7	191.4 $\pm$ 1.1	194.3 $\pm$ 2.3	n.s.
	Coarse silt	161.5 $\pm$ 4.0 a	140.1 $\pm$ 1.2 b	144.6 $\pm$ 2.1 b	F = 12.0 ^{***}
	Fine sand	210.8 $\pm$ 4.8	203.7 $\pm$ 3.2	214.4 $\pm$ 5.4	n.s.
	Coarse sand	242.9 $\pm$ 6.2 b	262.7 $\pm$ 4.2 a	233.2 $\pm$ 7.8 b	F = 7.2 ^{**}
	Total Carbon	16.2 $\pm$ 0.3	15.6 $\pm$ 0.2	15.4 $\pm$ 0.6	n.s.
	Total Nitrogen	1.4 $\pm$ 0.03	1.5 $\pm$ 0.02	1.5 $\pm$ 0.05	n.s.
	Carbon-Nitrogen ratio	11.0 $\pm$ 0.06 a	10.6 $\pm$ 0.04 b	10.5 $\pm$ 0.09 b	F = 25.0 ^{***}
	Organic matter	28.0 $\pm$ 0.5	26.9 $\pm$ 0.4	26.59 $\pm$ 1.1	n.s.
	pH	6.9 $\pm$ 0.04 c	7.1 $\pm$ 0.03 b	7.25 $\pm$ 0.05 a	F = 20.6 ^{***}
	Phosphorus	0.009 $\pm$ 0.0003 b	0.03 $\pm$ 0.006 a	0.04 $\pm$ 0.007 a	F = 22.0 ^{***}
	Calcium	1.4 $\pm$ 0.03 b	1.5 $\pm$ 0.03 ab	1.6 $\pm$ 0.05 a	F = 7.2 ^{**}
	Magnesium	0.9 $\pm$ 0.009	0.9 $\pm$ 0.02	0.9 $\pm$ 0.03	n.s.
	Potassium	0.3 $\pm$ 0.01 c	0.4 $\pm$ 0.03 b	0.4 $\pm$ 0.03 a	$\chi^2 = 24.7^{***}$
CWM traits	SLA	15.2 $\pm$ 0.3 b	21.4 $\pm$ 0.7 a	21.4 $\pm$ 0.9 a	$\chi^2 = 54.2^{***}$
	LDMC	333.9 $\pm$ 5.7 a	252.4 $\pm$ 6.3 b	276.0 $\pm$ 5.5 b	F = 52.4 ^{***}
	Canopy height	32.6 $\pm$ 0.5 a	29.3 $\pm$ 1.2 b	29.5 $\pm$ 1.1 ab	F = 5.2 ^{**}

Figure 1. Map of La Crau area in France with the six selected locations. Modified from Sylvain Fadda, IMBE (In light grey, the area of steppe remnants). At every location, we chose at least two study sites that represented one undisturbed steppe and at least one fallow land.

Figure 2. Detrended correspondence analysis (DCA) of all plots. Main matrix: vegetation data (16 sites with 5 vegetation relevés; 136 species) and second matrix: environmental factors (data on 19 parameters for 16 sites). In total, plots from eight steppe sites (white), five cereal fallows (light grey; rectangle: long-abandoned field (since 1900); triangle: recently-abandoned field (since 2000) and three melon fallows (dark grey) are shown. Vectors were significantly related to the axes ($P < 0.05$).

Figure 3. Results of non-metric multidimensional scaling (NMDS) carried out on the presence/ absence vegetation data for 2001 and 2015. The color of polygons (convex hulls) indicates the land- use types (white: steppe, light grey: cereal fallows, dark grey: melon fallows) while the structure represents the time of assessment (plain: assessed in 2001 with $n = 10$, dotted: assessed in 2015 with $n = 5$). Vegetation relevés from Peau de Meau, Valigne and Petit Abondoux were included. Final stress = 0.19. For clarity, only 56 species are displayed (out of 130).

Figure 4. Coefficient of variation (CV) of (a) specific leaf area (SLA), (b) leaf dry matter content (LDMC), (c) canopy height, (d) plant width and (e) biomass for selected species occurring in the steppe sites. The scales of the y-axis differ among panels. In the right-hand corner, the total CV of plant traits is given, represented by a dashed line in the boxplots. Kruskal–Wallis tests were applied followed by the pairwise Wilcoxon test with P-value adjustment according to Holm (** $P < 0.001$; ** $P < 0.01$; * $P < 0.05$; n.s. $P > 0.05$). Different letters indicate significant differences among species. For abbreviations of species names, see Table 2.

Supporting information

Helm, J. et al. Recovery of Mediterranean steppe vegetation after cultivation: legacy effects on plant composition, soil properties and functional traits, Applied Vegetation Science.

Appendix S1. Trait values of specific leaf area (SLA), leaf dry matter content (LDMC) and canopy height for all species. Mean $\pm$ standard errors are given for the measured trait values per species and land-use type. Different letters indicate significant differences between steppe, cereal and melon fallow. *** $P < 0.001$; ** $P < 0.01$; * $P < 0.05$; n.s. $P > 0.05$

		SLA				LDMC				Canopy height			
Species		Steppe	Cereal	Melon	Statistical test values	Steppe	Cereal	Melon	Statistical test values	Steppe	Cereal	Melon	Statistical test values
Unsuccessful colonizers	Bra_ret	9.5±0.3	-	-	-	512.8±7.1	-	-	-	15.6±0.5	-	-	-
	Hyp_gla	22.7±1.6	-	-	-	152.3±13.5	-	-	-	2.0±0.4	-	-	-
	Lin_tri	17.8±0.7 a	16.2±0.6 ab	15.0±1.0 b	$\chi^2 = 7.4^*$	330.9±11.9	327.4±13.2	324.2±18.0	n.s.	11.9±0.4	11.3±0.8	10.7±0.7	n.s.
	Sti_cap	5.1±0.1	4.9±0.1	5.0±0.3	n.s.	569.9±7.2	575.6±16.9	558.0±9.1	n.s.	32.7±1.0 a	26.4±1.5 b	25.3±1.6 b	F = 9.8***
	Tri_cam	27.7±0.7	-	-	-	281.7±7.0	-	-	-	6.1±0.4	-	-	-
Fallow-land species	Cre_foe	22.2±0.6	19.9±0.6	20.1±0.7	n.s.	192.7±4.3 a	181.4±5.5 b	186.7±5.2 ab	$\chi^2 = 6.6^*$	4.4±0.3 ab	4.0±0.3 a	5.3±0.4 b	$\chi^2 = 9.7^{**}$
	Dac_glo	20.8±0.4	20.8±0.5	22.0±0.6	n.s.	361.0±4.7	360.8±5.5	360.5±5.1	n.s.	14.8±0.6 a	11.7±0.6 b	11.9±0.7 b	$\chi^2 = 11.8^{**}$
	Gal_tom	11.3±0.3 a	10.9±0.3 a	9.6±0.4 b	F = 5.5**	228.9±8.2	233.2±10.8	228.8±19.1	n.s.	28.3±1.9	32.2±2.0	27.9±2.2	n.s.
	Tri_sub	-	19.4±0.5	-	-	-	313.5±10.8	-	-	-	2.3±0.2	-	-
Successful colonizers	Car_lan	7.4±0.2 a	6.5±0.2 b	7.7±0.3 a	$\chi^2 = 15.9^{***}$	394.3±8.5 a	341.8±11.1 b	366.7±15.1 ab	F = 4.3*	23.0±1.1 a	15.8±1.0 b	25.0±1.5 a	$\chi^2 = 15.1^{***}$
	Lob_mar	11.3±0.3	10.3±0.4	10.2±0.5	n.s.	303.1±7.6	323.7±9.8	333.1±12.1	n.s.	8.9±0.3 a	6.3±0.4 b	6.2±0.3 b	F = 22.5***
	Sid_rom	24.0±0.5	22.5±0.6	23.1±0.6	n.s.	224.8±3.9	230.0±6.5	222.9±7.7	n.s.	7.0±0.4 a	3.3±0.3 c	5.4±0.6 b	$\chi^2 = 29.6^{***}$

Appendix S2. Principal component analysis (PCA) of the species–trait matrix (854 Individuals of 9 species $\times$ 9 traits; without *Brachypodium retusum* and *Trifolium subterraneum* due to data constraints). Symbols represent the location of individuals in the multidimensional trait space. Species groups are indicated by colors (white: steppe species; light grey: fallow land species; dark grey: common species). Origins of measurements are distinguished: steppe (triangle) and fallow lands (circle). As a conceptual summary, the black arrow indicates the direction of changes in plant trait values due to land-use change (from steppe to fallow land). For abbreviations of species, see Table 2. For abbreviations of functional traits, see Table 3.

Appendix S3. Leaf nutrient content of carbon (C), nitrogen (N) and phosphorus (P) compared between original steppe and fallow lands (eight individuals per site pooled). Mean $\pm$ standard errors (SE) are shown. P was measured with inductively-coupled plasma optical emission spectrometry (IPC-OES, Optima 3000, Fa. Perkin Elmer) at the TLL (Thüringer Landesanstalt für Landwirtschaft, Jena). Leaf N and leaf P content were analyzed with a CHN elemental analyzer (Vario EL Elemental Analyzer, Elementar). Leaf nutrient content could not be measured for *Hypochaeris glabra*, *Trifolium campestre* and *Trifolium subterraneum* due to insufficient plant material.

	Steppe	Cereal	Melon	Statistical test values
	Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE	
Leaf carbon content	43.24 $\pm$ 0.25 a	41.40 $\pm$ 0.34 b	42.13 $\pm$ 0.45 ab	$\chi^2 = 13.0^{**}$
Leaf nitrogen content	1.70 $\pm$ 0.08 b	2.15 $\pm$ 0.14 a	1.94 $\pm$ 0.18 ab	$\chi^2 = 6.7^*$
Leaf phosphorus content	1.26 $\pm$ 0.01 c	2.86 $\pm$ 0.07 b	3.39 $\pm$ 0.10 a	$\chi^2 = 58.1^{***}$