Site- and Regioselectivity of the Hetero-Diels-Alder and Cheletropic Additions of Sulfur Dioxide to a 2,3,6,7-Tetramethylidenebicyclo[3.2.1]octane Derivative

Brigitte Deguin, Pierre Vogel

► To cite this version:

Brigitte Deguin, Pierre Vogel. Site- and Regioselectivity of the Hetero-Diels-Alder and Cheletropic Additions of Sulfur Dioxide to a 2,3,6,7-Tetramethylidenebicyclo[3.2.1]octane Derivative. *Tetrahedron Letters*, 1993, 34 (39), pp.6269-6270. hal-01923579

HAL Id: hal-01923579

<https://hal.science/hal-01923579>

Submitted on 15 Nov 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Site- and Regioselectivity of the Hetero-Diels-Alder and Cheletropic Additions of Sulfur Dioxide to a 2,3,6,7-Tetramethyldienebicyclo[3.2.1]octane Derivative.

Brigitte Deguin and Pierre Vogel*

Section de Chimie de l'Université de Lausanne, 2, rue de la Barre, CH 1005 Lausanne, Switzerland

Summary: At -55°C SO_2 adds to 4-[(*tert*-butyl)dimethylsilyloxy]-1-methoxy-2,3,6,7-tetramethyldienebicyclo[3.2.1]octane reversibly and selectively onto the *s*-cis-butadiene moiety at C(2), C(3) giving a single regioisomeric sultine. At -45°C the cheletropic addition of SO_2 occurs selectively giving the monosulfolene attached at C(2), C(3).

Simple 1,3-dienes such as isoprene and (E)-piperilene undergo hetero-Diels-Alder additions below -60°C and give the corresponding 3,6-dihydro-1,3-oxathiin-2-oxides (sultines) with high regioselectivity.¹ The sultines are unstable above -50°C and undergo cycloreversion liberating the starting dienes and SO_2 that undergo cheletropic additions² above -40°C giving the corresponding sulfolenes.³ The Diels-Alder additions of the exocyclic tetraene 1 to strong dienophiles ($\text{X}=\text{X}$) such as ethylenetetracarbonitrile, dimethyl acetylenedicarboxylate and maleic anhydride were not site selective and gave mixture of the corresponding monoadducts 2 and 3 together with the bis-adducts 4.⁴ No significant regioselectivity was noticed for the cycloaddition of 1 to methyl propionate and methyl vinyl ketone. In contrast with these results, we report here that SO_2 adds to tetraene 1 with high site-selectivity for both its hetero-Diels-Alder and cheletropic additions, the *s*-cis-butadiene moiety at C(2), C(3) being preferred over that at C(6), C(7). Furthermore, in the case of the hetero-Diels-Alder addition 1 + SO_2 only one single regioisomeric sultine was observed.

When a 0.2 M solution of 1 in $\text{CD}_2\text{Cl}_2/\text{SO}_2$ 7:1 (w/w) was allowed to stand at -55°C (5 mm sealed NMR tube) the monosultine 5 was formed slowly. After 2 days at -55°C , 50% of 5 was formed together with <5% of sulfolene 6 and ca. 5% of an unknown compound the ^1H -NMR spectrum of which was not consistent with an isomeric sultine or sulfolene. After 7 days at -55°C , nearly 95% of 1 had reacted. At -45°C , sultine 5 underwent cycloreversion into 1 + SO_2 that reacted slowly to give 6 which was obtained in 85% yield after 24 h at 0°C . The cheletropic addition of a second equivalent of SO_2 to give the bis-sulfolene 7 was a very slow process at 0°C . A good yield of 7 (70-80%) was obtained when 6 was reacted with a large excess $\text{CD}_2\text{Cl}_2/\text{SO}_2$ 1:5 (w/w) at 20°C for 24 h. After prolonged standing at 0 - 20°C , 6 did not equilibrate with an isomeric sulfolene. One cannot exclude yet that both 5 and 6 are preferred for thermodynamic reasons rather than for kinetic reasons. Because 7 decomposed on heating, we could not establish whether 6

was formed or not by elimination of SO₂.

The structures of **5**,⁵ **6**⁶ and **7**⁷ were deduced from their ¹H- and ¹³C-NMR spectra and with the help of double irradiation experiments. In the case of **5** the homoallylic coupling constants ⁵J_{H,H} between protons H-C(3) on one hand, and protons H-C(6) and H-C(8), on the other hand,⁸ allowed one to distinguish between pseudo-equatorial and pseudo-axial protons at C(3) and C(6). Strong NOE's were observed between the proton pairs H-C(8)/H-C(9), H-C(8)/H_e-C(6) consistently with the conformation shown for **5** in the Fig. NOE was also observed between the olefinic protons and H-C(9) and MeO protons. The axial S O moiety of **5** was assumed.⁹

The selectivities reported here for the reaction of **1** with SO₂ are striking; they rise a number of questions to which we hope to propose answers when further reactions of SO₂ with polyenes analogous to **1** will have been examined.

Acknowledgments. We thank *F. Hoffmann-La Roche & Co., AG (Basel)*, the *Swiss National Science Foundation*, and the *Fonds Herbette (Lausanne)*.

References and Notes

1. Deguin, B.; Vogel, P. *J. Am. Chem. Soc.* **1992**, *114*, 9210.
2. Woodward, R. B.; Hoffmann, R. In *"The Conservation of Orbital Symmetry"*, Academic Press, New York, 1970, pp. 152.
3. De Bruin, G. *Proc. K. Ned. Akad. Wet.* **1914**, *17*, 585; Turk, S. D.; Cobb, R. L. In *"1,4-Cycloaddition Reactions"*, Hamer, J., Ed., Academic Press, New York, **1907**, pp 13.
4. Burnier, G.; Vogel, P. *Helv. Chim. Acta* **1990**, *73*, 985.
5. Data of **5**: ¹H-NMR (Bruker, ARX 400, CD₂Cl₂/SO₂, -55°C) δ_H: 5.5, 5.2, 5.1, 4.8 (4s, H₂C=C(10), H₂C=C(11)), 4.57 (ddd, ²J=15, ⁵J(H_a-C(6),H_a-C(3))=3, ⁵J(H_a-C(6),H_e-C(3))=2, H_a-C(6)), 4.20 (dd, ²J=15, ⁵J(H_a-C(6),H_a-C(3))=1.0, H_e-C(6)), 3.82 (dd, ³J=2, ⁵J(H_a-C(3),H-C(8))=1.0, H-C(8)), 3.53 (dddd, ²J=17, ⁵J(H_a-C(3),H_a-C(6))=3, ⁵J(H_a-C(3),H_e-C(6)) = ⁵J(H_a-C(3),H-C(8))=1.0, H_a-C(3)), 3.06 (dd, ²J=17, ⁵J(H_a-C(6),H_e-C(3))=2, H_e-C(3)), 2.85 (dd, ³J=6, 2, H-C(9)), 2.28 (d, ²J=10, H_{syn}-C(12)), 1.92 (dd, ²J=10, ³J=6, H_{anti}-C(12)), 0.90 (s, t-Bu), 0.16 & 0.15 (2s, Me₂Si); ¹³C-NMR (100.6 MHz, CD₂Cl₂, -55°C) δ_C: 148.0, 145.0 (2s, C(10), C(11)), 129.4, 127.4 (2s, C(2), C(7)), 109.2, 101.0 (2t, ¹J=160, C(10), C(11)), 83.6 (s, C(11)), 73.3 (d, ¹J=145, C(8)), 59.7 (t, ¹J=151, C(3)), 52.1 (q, ¹J=142, MeO), 46.3 (d, ¹J=140, C(9)), 44.8 (t, ¹J=140, C(6)), 31.3 (t, ¹J=135, C(12)), 25.3 (q, ¹J=125, t-Bu), 17.7 (s, SiC), -5.1, -5.2 (2q, ¹J=118, Me₂Si).
6. Data of **6**: ¹H-NMR (360 MHz, CD₂Cl₂/SO₂, 0°C) δ_H: 5.62, 5.38, 5.22, 4.95 (4s), 4.0-3.9 (m, H_a-C(3), H_a-C(5), H-C(7)), 3.8-3.72 & 3.62-3.60 (2m, H_b-C(3), H_b-C(5)), 3.33 (s, MeO), 2.98 (m, H-C(8)), 2.22 (d, ²J=10) & 1.93 (dd, ²J=10, ³J=6, H₂C(11)), 0.95 (s, t-Bu), 0.20 & 0.15 (2 s, Me₂Si); ¹³C-NMR (100.6 MHz, CD₂Cl₂/SO₂, 0°C) δ_C: 147.4, 145.3, 139.9, 128.6 (4s), 109.7 (t, 2C, ¹J=159), 84.3 (s, C(1)), 72.4 (d, ¹J=146, C(7)), 58.8 (t, ¹J=143, C(3)), 55.7 (t, ¹J=157, C(5)), 52.6 (q, ¹J=143, MeO), 47.4 (d, ¹J=141, C(8)), 31.8 (t, ¹J=135, C(11)), 25.8 (q, ¹J=125), 18.3 (s), -4.61 & -4.74 (2 q, ¹J=119).
7. Data of **7**: ¹H-NMR (360 MHz, SO₂/CD₂Cl₂, 20°C) δ_H: 4.1 (m, H-C(7)), 4.0-3.75 (m, 8H, H₂C(3), H₂C(5), H₂C(10), H₂C(12)), 3.35 (s, MeO), 3.08 (m, H-C(8)), 2.58 (dd, ²J=10, ³J=5) & 2.45 (d, ²J=10, H₂C(14)), 0.90 (s, t-Bu), 0.16, 0.15 (2s, Me₂Si); ¹³C-NMR (100.6 MHz, SO₂/CD₂Cl₂, 20°C) δ_C: 125.5 (s, C(2), C(4)), 115.6, 113.9 (2s, C(6), C(9)), 70.9 (s, C(1)), 65.3 (d, ¹J=143, C(7)), 58.3, 57.0, 56.8, 56.1 (4t, ¹J=146), 52.9 (q, ¹J=143, MeO), 47.2 (d, ¹J=144, C(8)), 40.8 (t, ¹J=138, C(14)), 26.1 (q, ¹J=125), 18.7 (s), -4.5, -4.6 (2q).
8. Barfield, M.; Sternhell, S. *J. Am. Chem. Soc.* **1972**, *94*, 1905; Mahaim, C.; Carrupt, P.-A.; Vogel, P. *Helv. Chim. Acta* **1985**, *68*, 2182.
9. Juaristi, E.; Cuevas, G. *Tetrahedron* **1992**, *48*, 5019 and references cited therein.