High PAH degradation and activity of degrading bacteria during alfalfa growth where a contrasted active community developed in comparison to unplanted soil

Amélia Bourceret, Corinne Leyval, Pierre Faure, Catherine Lorgeoux, Aurélie Cebon

► To cite this version:

Amélia Bourceret, Corinne Leyval, Pierre Faure, Catherine Lorgeoux, Aurélie Cebon. High PAH degradation and activity of degrading bacteria during alfalfa growth where a contrasted active community developed in comparison to unplanted soil. *Environmental Science and Pollution Research*, 2018, 25 (29), pp.29556 - 29571. 10.1007/s11356-018-2744-1 . hal-01930927

HAL Id: hal-01930927

<https://hal.science/hal-01930927>

Submitted on 13 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

High PAH degradation and activity of degrading bacteria during alfalfa growth where a contrasted active community developed in comparison to unplanted soil

Amélia Bourceret^{1,2, §}, Corinne Leyval^{1,2}, Pierre Faure^{1,2}, Catherine Lorgeoux³, Aurélie Cébron^{1,2 †}

¹Université de Lorraine, LIEC UMR7360, Faculté des Sciences et Technologies, Bd des Aiguillettes, BP 70239, 54506 Vandoeuvre-lès-Nancy, France.

²CNRS, LIEC UMR7360, Faculté des Sciences et Technologies, Bd des Aiguillettes, BP 70239, 54506 Vandoeuvre-lès-Nancy, France.

³Université de Lorraine, CNRS, CREGU, GeoRessourcesLab., UMR 7359, Faculté des Sciences et Technologies, Bd des Aiguillettes, BP 70239, 54506 Vandoeuvre-lès-Nancy, France.

[§]Current affiliation: Department Plant Microbe Interactions, Max Planck Institute for Plant Breeding Research, 50829 Köln, Germany.

[†] Corresponding author: aurelie.cebron@univ-lorraine.fr; phone: (+33) 383684296

Key words: historically contaminated soil, alfalfa, rhizosphere, RNA, bacterial diversity, PAH-degraders, functional community

Abstract

PAH biodegradation in plant rhizosphere has been investigated in many studies, but the timescale of degradation and degrading bacteria activity was rarely considered. We explored the impact of plants on the temporal variability of PAH degradation, microbial abundance, activity and bacterial community structure in a rhizotron experiment. A historically contaminated soil was spiked with PAHs, planted or not with alfalfa, over 22 days with sampling once a week. In both conditions, most of the spiked PAHs were dissipated during the first week, conducting to polar polycyclic aromatic compound production and to decreased richness and diversity of bacterial communities. We showed a rapid impact of the rhizosphere on PAH degradation via the increased activity of PAH-degrading bacteria. After 12 days, PAH degradation was significantly higher in the planted (100% degradation) than in unplanted (70%) soil. Gram-negative (*Proteobacteria*) PAH-dioxygenase genes and transcripts were higher in planted than unplanted soil and were correlated to the spiked PAH degradation. Conversely, Gram-positive (*Actinobacteria*) PAH-dioxygenase gene transcription was constant over time in both conditions. At 12 days, plant growth favoured the

activity of many *Gamma-Proteobacteria* (*Pseudomonadaceae*, *Stenotrophomonas* and *Acinetobacter*) while in unplanted soil *Alpha-Proteobacteria* (*Sphingomonadaceae*, *Sphingobium* and *Magnetospirillum*) and *Actinobacteria* (*Iamia*, *Geodermatophilaceae*, and *Solirubrobacterales*) were more active.

Introduction

Polycyclic Aromatic Hydrocarbons (PAHs) are organic pollutants resulting from coal-related activities (coking plants, gas plants) and derived activities (iron- and steel-production plants). These toxic compounds are persistent in the environment. Their high hydrophobicity, which leads to adsorption to the soil matrix, reduces their bioavailability to microorganisms (Bogan and Sullivan 2003). This aging process decreases PAH bioavailability in soils over time, and slows down the rate of PAH biodegradation (Allard et al. 2000). Microbial diversity also evolves over time, suggesting an adaptation to contamination (Bourceret et al. 2016). Thus, it is essential to better understand the temporal variations in microbial activity, to monitor it, and to relate it to the dynamics of PAHs in soils. To address this, many studies have used soils spiked with model PAH compounds (Allard et al. 2000; Thomas and Cébron 2016), but this makes them more bioavailable and therefore more degradable than aged PAHs in historically contaminated soils. Therefore it is appeared to us more realistic to study the biodegradation of a complex mixture of PAHs so as to investigate the activity of autochthonous microbial populations from aged contaminated soils and involved in biodegradation (Cébron et al. 2013).

Rhizodegradation is a cost-effective and environment-friendly process using microorganisms associated to roots to dissipate soil organic pollutants, such as PAHs (Haritash and Kaushik 2009). Plants release root exudates composed of sugars, amino and organic acids, among other compounds (Walker et al. 2003; Gao et al. 2015). They provide carbon growth sources for microorganisms, and biosurfactant compounds that increase PAH bioavailability and dissipation (Joner et al. 2002; Zhou et al. 2011). In PAH-contaminated soils, root exudates can increase microbial abundance (Kirk et al. 2005) and activity (Cébron et al. 2011). Several studies showed that plants enhanced PAH dissipation by modifying microbial diversity (Kawasaki et al. 2012) and activity (Yergeau et al. 2014; Peng et al. 2015), and selecting PAH degraders (Storey et al. 2014; Siciliano et al. 2003) in the rhizosphere. Consequently, most studies indicated that the combination of all these processes increased PAH degradation in the presence of plants (Liste and Alexander 2000; Tejeda-Agredano et al. 2013) and more

precisely in the rhizosphere. However, other experiments also evidenced no effect (Gartler et al. 2014), or slower PAH degradation in planted soil as compared to unplanted soil (Thomas and Cébron 2016). These contrasting results could be explained by spatio-temporal variations that occur within the rhizosphere and activate various processes during plant development. This hypothesis justifies further research on the time course of PAH-degradation processes and on the activity of the functional microbial community in soil-plant systems.

The temporal dynamics of rhizospheric processes could explain the variable efficiency of PAH degradation observed with plants. In uncontaminated soil, it has been extensively shown that the composition and quantity of root exudates can vary depending on the plant developmental stage and on root location (Hinsinger et al. 2005; Walker et al. 2003), and can thus modify the microbial community (Yang and Crowley 2000; Hannula et al. 2010; Baudoin et al. 2002), functional diversity (Haichar et al. 2008), and microbial activity (Baudoin et al. 2002). On the other hand, these rhizosphere processes have received little attention in contaminated environments where pollution is an additional factor impacting microorganism activity. It is therefore essential to assess the impact of both PAH pollution and root development on overall microbial activity and on the PAH-biodegrading function to know which factors are the most determining ones in a rhizoremediation context.

Among plant species tested for PAH rhizodegradation, the efficiency of legumes – e.g. alfalfa – is relevant, because they have extensive root systems that allow soil colonization and nutrient uptake (Muratova et al. 2003; D’Orazio et al. 2013). Alfalfa is a plant commonly used *in situ* during land-farming treatment of organic pollutants (Trapp and Karlson 2001). It is also commonly used in microcosm studies to evaluate rhizodegradation efficiency, and more precisely study the processes involved (Bourceret et al. 2015). A legume such as alfalfa is thus a good and realistic model plant to evaluate the time course of PAH rhizodegradation in aged contaminated soil.

This study aimed to evaluate 1) the dynamics of PAH dissipation in alfalfa rhizosphere as compared to unplanted soil, and 2) the variability of microbial community activity and PAH-degrading bacterial abundance and activity over three weeks of plant growth. We developed a specific 2-compartment rhizotron device to rapidly establish an active root system and put it in contact with PAH spiked soil so as to avoid the time gap between root growth from seeds germination and the degradation of bioavailable pollutants. Rhizotrons were filled with an aged PAH-contaminated soil, initially spiked with a complex mixture of PAHs (i.e. a soil

organic extract). Rhizotrons were sampled once a week to monitor concentrations of PAHs and potential metabolites (polar polycyclic aromatic compounds, pPAC). Dissolved organic carbon (DOC), pH, and humidity were measured to characterize the rhizosphere and soil conditions. Working on extracted DNA and RNA fractions, we also assessed total bacterial and fungal abundance, PAH-degrading bacterial abundance and transcriptional activity levels. Finally, we assessed active bacterial community structures by pyrosequencing of 16S rRNA transcripts.

Materials and Methods

Experimental device

We used an aged PAH contaminated soil, from a former coking plant (Neuves-Maisons (NM), Lorraine, France) to perform rhizotron experiments. Soil characteristics and spiking procedure were described previously by Bourceret et al. (2015). Briefly, the NM soil (1033 ± 148 mg per kg of dw soil of $\Sigma 13$ PAHs (fluorene, phenanthrene, anthracene, fluoranthene, pyrene, benzo(a)anthracene, chrysene, benzo(b)fluoranthene, benzo(k)fluoranthene, benzo(a)pyrene, dibenzo(a,h)anthracene, benzo(g,h,i)perylene, indeno(1,2,3-cd)pyrene) was air-dried, sieved at 2 mm and spiked with a complex organic pollutant extract (**Table S1**), rich in PAHs and obtained from the same soil using chloroform extraction (Cébron et al. 2013). After solvent evaporation, the spiked soil was mixed with NM soil not exposed to chloroform, in a one-tenth ratio. Experimental device was divided in two parts (**Fig. 1**): the top part (mini-rhizotron) was a poly-methyl methacrylate box (7 cm high x 29.8 cm long x 2.3 cm wide) without bottom, that fits into the second part: a rhizotron (30 cm high x 30 cm long x 2.5 cm wide) with a removable front face allowing spatially localized depth sampling as described by Bourceret et al. (2015). First, 8 mini-rhizotrons were filled with NM soil, closed at the bottom by bolting cloth (5 μ m mesh), and adjusted to 80% of the soil water holding capacity (WHC) (corresponding to 203 ml deionized water per kg of dw soil). Alfalfa (*Medicago sativa* var. Europe) was planted on 4 of these devices by seeding 2.4 g of seeds (c.a. 26 seeds per cm^2) and the other 4 devices were left unplanted. After germination, the 8 devices were placed in a plant growth chamber under controlled conditions (22°C/18°C day/night, 80% relative humidity, c.a. 250 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$, 16 hours photoperiod) and were maintained at 80% of the WHC by weighing the devices every two days. The previous growth of plant in NM soil allowed to avoid the delay between time needed for root development and the PAH degradation, that was already shown to be fast in spiked soil with a previous study (Bourceret et al. 2015). After 15 days, when roots reached the bottom of the mini-rhizotron devices, eight

rhizotrons were filled with spiked NM soil (1418 ± 193 mg of $\Sigma 13$ PAHs per kg of dw soil), bolting cloths of the mini-rhizotrons were removed and devices were fitted in their corresponding rhizotron to allow root penetration and growth in spiked soil. This time corresponded to T0. Initial spiked NM soil was sampled: six aliquots from each of the eight rhizotrons corresponding to 48 T0 samples were stored at -80°C for further analyses. Then the 4 planted and 4 unplanted devices were incubated in plant growth chamber in conditions described above. After 6 (T1), 12 (T2), 16 (T3) and 22 (T4) days of plant growth, two rhizotrons (one planted and one unplanted) were harvested and opened to allow core sampling using a stainless steel cylinder equipped with a PTFE piston (1.6 cm diameter, 2.5 cm long, thus enabling to collect soil cores of around 5 cm^3). At 3 cm from the interface of the two compartments, 6 samples with 3 cm between the core centers were collected per device (**Fig. 1**). Soil samples were prepared as previously described by Bourceret et al. (2015). Briefly, after 3 rinsing by sterile water, roots were spread on a black paper and photographed to estimate the size of each root fragment using ImageJ software (<http://rsbweb.nih.gov/ij/>). Roots were then air-dried at 60°C and weighed to quantify dry root biomass. Rhizosphere soil, collected from root rinsing water by centrifugation at 4400 rpm for 10 minutes, and bulk soil were weighed to calculate the ratio between the two fractions and were mixed together to obtain a unique soil sample stored at -80°C before analyses: soil characteristics (pH, DOC, organic acids, and sugar concentrations), PAH and pPAC concentrations, and DNA/RNA extractions. Soil humidity was measured by collecting 3 other samples on the side of the rhizotrons and by weighing the soil before and after drying at 60°C for 48 hours.

Soil characteristics

Soil solutions were prepared with 1 g of soil in distilled water (1:5 w/v ratio), and pH was measured using a pH-meter electrode (BioBlock scientific, pHM210 Radiometer analytical). After centrifugation (10 min, 4000 rpm, 20°C) supernatant was filtered ($0.45\text{ }\mu\text{m}$) and diluted 20 times to quantify dissolved organic carbon (DOC) with a total organic carbon analyzer (TOC-V CSH, Shimadzu). Sugars (inositol, trehalose, sucrose, glucose, fructose) and organic acids (formate, butyrate, oxalate) were quantified from non-diluted extract with an ion-exchange chromatograph ICS 5000 (CarboPac SA10 column, 40°C) and ICS 3000 (AS 11-HC column, 35°C), respectively.

Extraction and quantification of total PAHs, pPACs and bioavailable PAHs

An aqueous solution of hydroxypropyl- β -cyclodextrin (20 ml, 50 mM) (Acros organics, New Jersey, USA) was used to extract available PAH fraction from 1 g of fresh soil from 5 NM

soil samples before spiking, 6 T0 samples, and 48 rhizotron samples (6 samples from both planted and bare rhizotrons at each sampling time) in Oak Ridge Centrifuge tubes (FEP, Nalgene, USA) (Reid et al. 2000). After stirring (100 rpm) for 16 hours at 24 °C and 2 centrifugation steps (10000 rpm, 20 min), dichloromethane was added (24 ml) to the aqueous supernatant (c.a. 30 ml) to extract available PAHs. An Accelerated Solvent Extractor (ASE350 Dionex ®) was used to extract total PAHs and pPACs with dichloromethane (130°C, 100 bars) from 0.25 g of soil (stored at -80°C) from 12 NM soil samples before spiking, 48 T0 samples, and 48 rhizotron samples previously lyophilized and crushed (<500 µm). Dichloromethane extracts were split in two fractions to quantify PAHs using HPLC/UV-Fluo (High Pressure Liquid Chromatography) and pPACs using GC-MS (gas chromatography coupled to mass spectrometry). Firstly, the two dichloromethane extracts containing available and total PAHs were evaporated under nitrogen flow and diluted in 5 ml and 20 ml of acetonitrile, respectively. HPLC (Dionex Ultimate 3000) system, with a 100-mm long and 4.6-mm internal diameter separation column (SupelcoAscendis Express C10), 2.6-mm granulometry allowed quantifying PAHs from 10 µl of extract. From total and available extracts, 13 PAHs (16 US-EPA PAHs excluding the three more volatile molecules: naphthalene, acenaphthylene and acenaphthene) were quantified by using a UV (254 nm) and a fluorescence detector, respectively. Secondly, from dichloromethane extracts pPACs were quantified on a GC-6890 (Agilent Technologies) equipped with a DB5- MS column (60 × 0.25 mm id × 0.25µm film thickness) coupled to a MS-5973 Inert (Agilent Technologies) operating SIM mode (Single Ion Monitoring). The oven temperature program was as follows: 2 minutes at 70 °C, then from 70 °C to 130 °C at 15 °C min⁻¹, then from 130 °C to 315 °C at 4 °C min⁻¹ and then a 25 min hold at 315 °C. The carrier gas was helium at 1.4 ml min⁻¹ constant flow. Fifteen pPACs including 11 oxygenated pPACs (O-PACs) and 4 nitrogenated pPACs (heterocyclic PACs containing nitrogen N-PACs) were quantified by an internal calibration using deuterated internal standards (the list of the 15 quantified polar PAC is given in Hanser et al. (2015)). During a sequence, calibration controls were injected every 12 samples to check the overall calibration.

DNA/RNA extraction

Nucleic acids were extracted from 0.5 g of soil using a FastDNA SPIN Kit for soil (MP Biomedicals, France). DNA and RNA were co-extracted from the same aliquots of samples stored at -80°C. DNA was resuspended in 100 µl of DES (Dnase-free water) and quantified at 480 nm (Xenius, SAFAS) by using Quant-iTPicoGreenasDNA Assay kit (Invitrogen), and

then extracts were diluted to 2 ng/μl before amplification. RNA was collected from the filtrates at the DNA binding to the column step. After sodium acetate (0.1Vol, 3M, pH 5.2) and isopropanol precipitation for one night at 4°C and centrifugation, RNA was resuspended in 45 μl of Rnase-free water. Two steps of DNase (Thermo scientific) treatment (addition of 10 μl of DNase buffer (10X), 1 μl of DNase (1u/μl) and incubated 1 hour at 37°C) and purification using RNeasyMinElute Cleanup kit (Qiagen) allowed obtaining 30 μl of pure extract (presence of DNA contamination was PCR checked). After quantification at 480 nm (Xenius, SAFAS) by Quant-iTRiboGreen RNA Assay kit (Invitrogen), extracts were converted in cDNA using SuperScript III First-Strand kit (Invitrogen). Reverse transcription reaction (20 μl) was performed using 1 μl of random hexamer (50 ng/μl), 1 μl of hybridization buffer and 22.27 ng of RNA (5 min at 65°C and 1 min on ice), then 10 μl of reaction buffer (2X) and 2 μl of enzyme mix were added and incubated 10 min at 25°C, 50 min at 50°C and 5 min at 85°C.

Real-time PCR quantification

The primer sets Fung5F/FF390R (Lueders et al. 2004; Thion et al. 2013), 968F/1401R (Felske et al. 1998; Cébron et al. 2008) and PAH-RHD_αGN F/R, PAH-RHD_α GP F/R (Cébron et al. 2008), targeting the fungal 18SrRNA genes, the bacterial 16S rRNA genes, and the PAH-ring hydroxylating dioxygenase genes from Gram-negative and Gram-positive bacteria, respectively, were used to quantify by real-time PCR the abundance of active and total fungal, bacterial and PAH-degrading bacterial populations. Amplification reactions were performed on 20 μl of reaction mix containing 10 μl of iQ SYBR green SuperMix (Bio-Rad), 0.8 μl of each primer (10 μM), 0.4 μl of 3% BSA (bovine serum albumin) solution, 0.2 μl of DMSO (dimethyl sulfoxide), 0.08 μl of T4 bacteriophage gene 32 product (MP Biomedicals, France) and 1 μl of diluted gDNA or pure cDNA. Amplification and quantification were achieved using a CFX96 Real-Time PCR detection system (Bio-Rad) according to standard plasmid dilution series and program described by Cébron et al. (2008) and Thion et al. (2012). After soil humidity correction, data were expressed in number of gene copies per gram of dry weight soil, in percentage of transcripts relative to gene copies, and in percentages of 18S rRNA or PAH-dioxygenase relative to 16S rRNA gene or transcript copies.

Pyrosequencing and sequence analysis

The primer set 515F ((5'-GTG CCA GCM GCC GCG GTA A-3') / 907R (5'-CCG TCA ATT CMT TTR AGT TT-3')) (Turner et al. 1999) was used to amplify the V4-V5 region of bacterial 16S rDNA from gDNA and cDNA. Samples were identified with a MID (Multiplex

Identifier, Roche). PCR reactions were achieved as described in Bourceret et al. (2016). Beckman Coulter Genomics (Danvers, MA, USA) sequenced libraries using a 454/Roche GS-FLX Titanium system. Data were deposited in the NCBI Sequence Read Archive (SRA) under the BioProject ID: PRJNA324193. Denoised 16S rDNA sequenced were analysed in QIIME 1.8.0 pipeline (Caporaso et al. 2010) as previously described by Bourceret et al. (2017). Briefly, after filtering and reverse complement non-sense sequences, data from gDNA and cDNA were gathered and clustered in Operational Taxonomic Units (OTUs) at 97% of similarity and their representative sequences were affiliated at different taxonomic levels (from phyla to genus) using RDP classifier (Wang et al. 2007) and the Greengenes 13_5 database (McDonald et al. 2012). Then subsampling was performed to the lowest number of sequences counted in one of the samples (4,739 sequences per sample) and alpha-diversity indexes (richness and diversity estimates) were calculated using QIIME 1.8.0 pipeline.

Statistical analyses

Statistical analyses were performed using XLStat 2013 software (Addinsoft). One-way analysis of variance (ANOVA, with $p < 0.05$) followed by a Newman-Keuls multiple comparison test was performed to measure the time effect in each condition (planted and unplanted) between 0 and 22 days. Two-way analysis of variance (ANOVA, with $p < 0.05$) followed by a Newman-Keuls multiple comparison tests were performed to measure the time and plant effect between T1 and T4. Box plots were drawn using PAH data to compare their distribution and variability. Significant differences in relative abundance of dominant OTUs ($\geq 1\%$ in at least one sample) over time, were identified in cDNA samples with STAMP version 2.0.8 (Parks and Beiko 2010) by a Tukey Kramer ANOVA, with a Benjamini Hochberg correction of p value ($q < 0.05$ and Effect size (ES) > 0.26). Significant differences in the ratio cDNA/gDNA OTU abundance, according to the presence of plant was studied by Student-t test with a Benjamini Hochberg correction of p value with the differential expression function in XLStatOmics package. Canonical correspondence analysis (CCA) followed by a Monte Carlo permutation test (1000 permutations) was performed by using XLStat-ADA 2015 software (Addinsoft) to study, considering both conditions (alfalfa and unplanted soil) and 4 sampling times (0, 6, 12, 16 and 22 days), relationship between the relative abundance of the 41 most abundant OTUs ($\geq 1\%$ in at least one sample) and environmental variables (root biomass and length, pH, percentage of Gram-negative and Gram-positive PAH dioxygenase transcripts and genes, sugars, organic acids, and PAH content).

Results

Rhizosphere parameters

Humidity stayed constant (around 80% of the WRC) over time in both planted and unplanted soil rhizotrons but was globally significantly higher in unplanted soil (78.36 ± 20.91 and 70.85 ± 9.98 %, in unplanted and alfalfa planted soil, respectively). In alfalfa rhizotron, root biomass was significantly ($p < 0.05$) higher after 22 days of plant development than after 6 days, whereas root length significantly increased over time and was significantly different between 6 and 12 days and then between 16 and 22 days (**Fig. 2A**). Related to the plant growth, several parameters, highlighting the effect of root exudation on soil properties, increased over time. Globally sugar concentration and DOC increased significantly over time in the planted compared to unplanted soil. Indeed sugar concentration was significantly higher after 12 days in planted than in unplanted soil where no sugar was detected (**Fig. 2B**) and DOC stayed constant over time in both rhizotrons until the 16th day, and then increased significantly in planted soil on the 22nd day (**Fig. 2C**). At this time point, organic acids were detected in alfalfa planted rhizotron (0.21 ± 0.13 mg.kg⁻¹) but none in unplanted soil. Surprisingly soil pH increased significantly over time in both conditions and was stable after 12 days. However it increased more rapidly in the planted than in the unplanted soil rhizotron with a significant difference at 6 days (**Fig. 2D**).

PAHs and pPACs contents

The total aged PAH concentration ($\Sigma 13$ PAHs) of the NM soil (1033 ± 148 mg.kg⁻¹dw soil) was significantly increased ($p < 0.05$), i.e 27 %, after soil spiking with the organic pollutant extract and reached 1418 ± 193 mg.kg.dw soil⁻¹ at the beginning (T0) of the experiment (**Fig. 3**). The bioavailable PAH fraction was also increased by 41 % and reached 63.7 ± 18.0 mg.kg⁻¹dw soil, while it represented 37.3 ± 15.6 mg.kg⁻¹dw soil before spiking (data not shown). Spiking increased the concentration in total and bioavailable fractions of most of the 13 PAH compounds (**Fig. S1**). The total PAH concentration decreased significantly between 0 and 6 days and stayed constant until the 22nd day in both planted and unplanted rhizotrons (**Fig. 3**). From the 6th days, the total PAH concentration dropped down to 1200 ± 210 and 1024 ± 190 mg.kg.dw soil⁻¹ in unplanted and planted soil, respectively, corresponding to a decrease in 19 and 24% relative to the initial concentration, and to 70 and 100% of the spiked PAH fraction, respectively. The concentration in total PAHs was significantly lower in alfalfa than in unplanted soil when considering all times together (from 6 to 22 days, two-way ANOVA) and was mostly explained by a significant difference after 12 days (Student t-test) of incubation

(**Fig. 3**). The concentrations of several compounds decreased significantly from the 6th day (fluorene, phenanthrene, fluoranthene, pyrene, benzo(k)fluoranthene, benzo(a)pyrene and benzo(g,h,i)perylene only in planted condition) whereas the concentration of other PAHs remained constant over the time course (anthracene, benzo(a)anthracene, chrysene, benzo(b)fluoranthene, dibenzo(a,h)anthracene, indeno(1,2,3-cd)pyrene) or even increased significantly (benzo(g,h,i)perylene) from the 22nd day, in presence of plant (**Fig. S2**). Bioavailable PAH fraction decreased also significantly between T0 and the 6th day in both conditions, reaching 0.05 ± 0.01 and 0.08 ± 0.03 mg.kg⁻¹ dw soil, corresponding to ca.100 % decreases, for both planted and unplanted soil rhizotron. After one week, the bioavailable PAH fraction was significantly lower than in the soil before spiking. Ten oxygenated and one nitrogenous pPAC compounds were detected in initial soil (T0) and in both conditions after 6 days (**Table 1**). After 6 days the concentration of 4 oxygenated pPAC (methylantracene-9,10-dione, benzoanthracenedione, naphthacene-5,12-dione) was significantly higher than at T0, without significant differences between planted and unplanted soils. Benzo(cd)pyrenone was only significantly higher at 6 days than at T0 in the planted rhizotron. The concentration of the other pPAC stayed constant over the first week for both planted and unplanted soils.

Microbial communities

Abundance of active and total microbial communities

Bacterial abundance was higher than fungal, and bacteria were also more active over time in both planted and unplanted soil (mean value of $7.84 \times 10^8 \pm 6.64 \times 10^8$ 16S rRNA gene copies and $1.12 \times 10^{10} \pm 1.25 \times 10^{10}$ transcript copies g dw soil⁻¹) while fungi were 100 times less represented (mean value of $9.94 \times 10^6 \pm 7.04 \times 10^6$ 18S rRNA gene copies and $2.05 \times 10^8 \pm 1.48 \times 10^8$ transcript copies g dw soil⁻¹) (**Fig. 4A, B**). Abundance of 16S rRNA genes tended to increase over time in alfalfa and stayed constant in unplanted rhizotron whereas number of 16S rRNA transcripts increased significantly between T0 and the 6th day for both rhizotrons and then stayed constant. Abundance of 16S rRNA genes and transcripts was significantly higher in planted than unplanted rhizotron and plant effect was mainly detected at 12 (T2) and 22 days (T4). Abundance of 18S rRNA genes and transcripts tended to be constant in both unplanted and planted rhizotrons and no plant effect was observed.

Abundance of Gram-negative PAH-dioxygenase genes and transcripts increased significantly between T0 and the 6th day in both rhizotron types, and were higher in planted than in unplanted soils, with a significant plant effect after 12 days (**Fig. 4A and B**). Abundance of Gram-positive PAH-dioxygenase genes slightly increased over time whereas transcripts

largely increased between T0 and 6 days in both soil and further increased in planted soil till 12 days, resulting in a higher content in planted than in unplanted rhizotron, significant at each sampling time. The level of PAH-dioxygenase gene transcription was expressed by the ratio of the number of transcripts relative to corresponding gene copies (**Fig. 4C**). The level of PAH-dioxygenase gene transcription was significantly higher in alfalfa than in unplanted soil specifically after 12 days, and then decreased (for Gram-negative bacteria) and stayed constant (for Gram-positive bacteria) over time without significant differences with unplanted soil (**Fig. 4C**).

Richness, evenness and diversity of the active community

Globally, bacterial richness (Chao1) and diversity (Inverse of Simpson's) indices were significantly higher in unplanted than in alfalfa planted rhizotron (**Table 2**). This difference was significant after 16 days for richness and 12 days for diversity index (**Table 2**). No significant impact of plant was detected on the bacterial community evenness. For both studied conditions, bacterial richness, diversity and evenness indices decreased significantly during the 6 first days and after this time point increased significantly over time.

To explore the potential bacterial population involved into the rapid PAH degradation appearing at the beginning of the incubation period, we looked for bacterial community structure modification over time (**Fig. 5**). Among the active community, 12 phyla were identified and their proportion changed over time. In both rhizotrons, *Proteobacteria* was the more active phylum all over the incubation to the detriment of *Chloroflexi*, *Actinobacteria* and *Nitrospirae* that represented a lower proportion of the active community along the time course compared to the T0. Activity of *Beta*- and *Gamma-Proteobacteria* increased between 0 and 6 days and stayed constant or decreased until the end of incubation while the proportion of active *Alpha*-subdivision members further increased. An opposite trend was observed for the activity of *Planctomycetes*, *Acidobacteria* and *Delta-Proteobacteria* that was first repressed between 0 and 12 days and then favored at the end of the incubation (**Fig. 5**). The relative abundance of some active phyla evolved also differently over time according to the presence of alfalfa. In alfalfa planted rhizotron, *Bacteroidetes* and *TM6* were more and less active over time, respectively, while the activity of *Unclassified Proteobacteria* was transiently high and then low over the 22 days. In planted soil, 3 OTUs affiliated to *Beta-Proteobacteria* (*Thiobacillus*, and *Un.Oxalobacteraceae*), 1 OTU affiliated to *Alpha-Proteobacteria* (*Mycoplana*) and 1 OTU affiliated to *Firmicutes* (*Bacillus selenatarsenatis*) were significantly more abundant among the active community after 6 days of incubation than

at all other times, contrary to the unplanted rhizotron where no OTU was favored after 6 days (Fig. S3). After 12 days of incubation, three other OTUs belonging to *Alpha-* (*Sphingobium*), *Beta-* (*Janthinobacterium*) and *Gamma-* (PYR10d3) *Proteobacteria* were favored in planted soil. In unplanted rhizotron, 5 OTUs affiliated to *Alpha-* (*Mycoplana*, *Phenylobacterium*) and *Beta-* (*Thiobacillus*, *unclassified. Comamonadaceae*) *Proteobacteria* were significantly more abundant among the active community after 12 days of incubation.

Significant positive or negative impact of plants on shaping active bacterial community was observed on 7 and 18 OTUs after 6 and 12 days of incubation, respectively (Fig. 6). After 6 days, the activity ratio between 16S rRNA and rDNA for 3 OTUs belonging to *Gamma-* (*Pseudomonadaceae*), *Beta-* (*Achromobacter*) *Proteobacteria* and *Firmicutes* (*Paenibacillus*) was significantly higher in alfalfa planted than in unplanted soil. Four other OTUs, affiliated to *Alpha-* (*Un. Sphingomonadaceae*), *Beta-* (*Un. Burkholderiales*) *Proteobacteria* and *Actinobacteria* (*Un. Micrococcaceae*) were more active in unplanted soil. After 12 days, plant growth favored significantly the activity of 11 OTUs belonging to *Alpha-* (*Devosia*), *Beta-* (*Methylophilaceae*), *Gamma-* (*Pseudomonadaceae*, *Stenotrophomonas*, *Acinetobacter*) *Proteobacteria*, *Bacteroidetes* (*Dyadobacter*) and *Actinobacteria* (*Un. Micrococcaceae*). In unplanted soil the activity of 6 OTUs affiliated to *Alpha-Proteobacteria* (*Un. Sphingomonadaceae*, *Sphingobium*, *Magnetospirillum*) and *Actinobacteria* (*Un. Solirubrobacterales*, *Un. Geodermatophilaceae*) was significantly higher than in alfalfa planted soil. After 16 and 22 days of incubation, 56% and 76% of the OTUs whose activity is significantly impacted by the presence of plant, were more active in alfalfa planted rhizotron (Fig. S4).

Multivariate analysis

The two first axes of the canonical correspondence analysis (CCA) explained 62.5 % of bacterial diversity variability among samples (Fig. 7A and B). The F1 axis (explaining 41.8% of the variability) separates samples from the T0 (located on the right) to all other sampling times. The F2 axis (explaining 20.8% of the variability) separates samples according to the presence of plant and time, with 6 to 22 day samples from the bottom to the top. Alfalfa planted samples were separated from unplanted soil samples according to sugar and DOC contents, root development (biomass and length), abundance of Gram-negative PAH-dioxygenase genes and transcripts, and abundance of Gram-positive PAH-dioxygenase transcripts (Fig. 7A). These parameters are correlated to the higher activity of 11 bacterial OTUs, most belonging to *Proteobacteria* except one *Bacteroidetes* (*Dyalobacter*), in the

alfalfa planted rhizotron (**Fig. 7B**). These OTUs gathered bacteria affiliated to *Pseudomonadales* (*Acinetobacter* and *Pseudomonas*), *Sphingomonadales* (*Sphingobium*), *Burkholderiales* (*Janthinobacterium* and *Achromobacter*) and *Rhizobiales* (*Un. Methylobacteriaceae*) orders. Unplanted soil samples were mostly discriminated according to the higher PAH content ($\Sigma 13$ PAHs and 2-3, 4 and 5-6 cycles). According to the Pearson correlation matrix (**Table S2**), pH and the root biomass and length were positively and significantly correlated to the percentage of Gram-positive PAH-dioxygenase transcripts contrary to the abundance of 18S transcripts, showing a negative correlation. Moreover concentrations of 2-3, 4- rings and $\Sigma 13$ PAHs were significantly and negatively correlated to the percentage of Gram-negative PAH-dioxygenase transcripts, shown as the most active degrading community at the beginning of the monitoring.

Discussion

Rapid PAH dissipation in a spiked aged-contaminated soil

After 6 days of incubation 100 and 70% of the spiked PAH fraction was dissipated in the planted and unplanted rhizotrons, respectively. A similar rapid dissipation, of bioavailable PAHs after a few days was previously observed in simplified sand or soil microcosms spiked with phenanthrene (Louvel et al. 2011; Cébron et al. 2011), but never observed with such a complex PAH contaminant mixture. We evidenced the degradation of almost all spiked PAHs including fluorene, phenanthrene, fluoranthene, pyrene, benzo(k)fluoranthene and benzo(a)pyrene (which have from 2 to 5 aromatic cycles) in both conditions. In the same soil (NM) incubated in bioslurry, 30 to 55% of the aged PAH compounds (fluorene, phenanthrene, fluoranthene, pyrene, benzo(a)anthracene and chrysene) were degraded after 1.5 months of incubation (Biache et al. 2017), indicating the presence of a very effective microbial community for the degradation of a wide range of PAH compounds. Degradation of PAHs by microbial consortia and co-metabolism are often described as playing crucial roles in the degradation of PAH mixtures in the environment (Ghosal et al. 2016) because the presence of low-molecular-weight PAHs could have a synergistic effect on the degradation of more recalcitrant compounds (Van Herwijnen R et al. 2003; Roy et al. 2013). During the first 6 days of the experiment, spiked PAHs were probably the main available carbon sources in both the planted and unplanted rhizotrons, as no increase in DOC or sugar content was observed in either condition. As the microbial community of the aged-contaminated NM coking plant soil was well adapted to PAH pollutants (Cébron et al. 2009; Bourceret et al. 2016), the bacterial

community was rapidly activated, and the spiked PAH fraction was in turn rapidly biodegraded. The main parameter that limited the biodegradation of aged PAHs in this soil was their low bioavailability (Cébron et al. 2013) and spiking transiently led to more easily biodegradable PAHs. In our experiment, before being irreversibly adsorbed to soil mineral or organic materials, such as clay, black carbon or coal-tar substances (Ghosal et al. 2016), PAHs were rapidly degraded, proving that their availability is crucial for biodegradation. Besides PAH degradation, oxygenated polar compounds (pPAC) were produced during the first week of the experiment. This kind of bioavailable compounds, previously described in water from industrial wasteland (Rollin et al. 2005; Lundstedt et al. 2006) and in coal tar (Northcott and Jones 2001; Benhabib et al. 2010), could result from biotic PAH degradation. Lundstedt et al. (2007) indeed described the production of oxygenated compounds during microbial biodegradation, with apparent concentrations resulting from the balance between pPAC degradation and pPAC production during PAH biodegradation (Biache et al. 2017). The production of oxygenated pPAC possibly more available and more toxic than parent molecules could contribute to shape the bacterial community structure.

Together with PAH biodegradation, we observed that bacterial community abundance and activity rapidly increased during the first 6 days of incubation. Moreover the community composition was drastically modified in both planted and unplanted conditions. Most of bacterial OTUs with increasing activity were affiliated to *Beta*- and *Gamma-Proteobacteria*, at the expense of *Actinobacteria* whose proportions fell sharply. PAH amendment may have caused a shift of bacterial communities from oligotroph, K-strategist, Gram-positive bacteria (such as *Actinobacteria*) to copiotroph, r-strategist, Gram-negative bacteria (such as *Proteobacteria*) (Bevivino and Dalmastri 2017). These *Proteobacteria* are potentially Gram-negative PAH-degraders. We did find a predominance of bacteria known to be involved in PAH degradation, such as OTUs affiliated to the *Achromobacter*, *Janthinobacterium*, *Pyr10d3*, *Sphingobium*, and *Thiobacillus* genera and *Sphingomonadaceae*, *Micrococcaceae*, and *Comamonadaceae* families (Fernández-Luqueño et al. 2011; Kunihiro et al. 2013). The abundance and transcription level of PAH-dioxygenase genes increased significantly during the first 6 days in both conditions, confirming the higher potential of PAH bacterial degradation. Greater activity of Gram-negative than Gram-positive (*Actinobacteria*) PAH degraders was evidenced during these early days, indicating that *Proteobacteria* were most probably the main phylum involved in the biodegradation of the spiked PAHs. This finding was also supported by a higher activity of *Proteobacteria* as compared to other taxa in both planted and unplanted soils, and by a negative correlation between the activity of Gram-

negative PAH-degrading bacteria and PAH contents. Several studies indicated that PAH contamination induces a selection of and an enrichment in PAH degraders (Cébron et al. 2008; Zhang et al. 2013; Sawulski et al. 2014), and also up-regulates the transcription of the PAH-dioxygenase gene in pure culture (Kim et al. 2006) or complex communities (Louvel et al. 2011). A negative correlation between the transcription of PAH-dioxygenase genes of Gram-negative bacteria and PAH contents would confirm the major involvement of this population in early pollutant degradation, although we cannot rule out the involvement of Gram-positive *Actinobacteria* that could be the main active PAH-degraders after PAH spiking (de Menezes et al. 2012). Furthermore, fungi could also be involved in PAH degradation (Cerniglia and Sutherland 2010), but we did not detect any increase of fungal 18S rRNA gene or transcript copies during incubation in either condition, suggesting a fairly limited involvement of these microorganisms in our soil.

Plant effect

The presence of plants induced a shift in the active bacterial community structure and the PAH-degrading population. The timescale of the experiment (22 days) allowed us to monitor the root development of alfalfa, and compare processes between planted and unplanted soils. In addition to increasing the PAH biodegradation capacity, alfalfa plants globally increased the abundance and activity of bacteria, the abundance of Gram-negative PAH-degrading bacteria and the expression of both Gram-negative and Gram-positive PAH-dioxygenase genes. However, the presence of plants led to a decrease of the bacterial richness and diversity indexes. This finding contrasts with previous observation from the same soil (NM) spiked with phenanthrene and planted with ray-grass. The authors found that plants indeed induced an increase of bacterial species richness and slightly inhibited phenanthrene degradation, even if the abundance of PAH-dioxygenase genes increased, yet without modification of the transcription level as compared to unplanted soil (Thomas and Cébron 2016). These contrasting results can be explained by two different types of contamination, i.e. a PAH mixture *vs.* phenanthrene, and two different plant species with contrasted root systems and root exudation (Bourceret et al. 2015). Root exudates probably partly inhibit the degradation of one sole PAH compound, while they might help in co-metabolism processes when PAH contamination is complex and contains high-molecular-weight compounds (Rentz et al. 2005) as shown for PCB degradation (Vrchotová et al. 2013).

After 6 days, even if root biomass and length were low, the presence of alfalfa had already modified a few parameters. The pH, the abundance and proportion of active Gram-negative

PAH-degrading bacteria were significantly higher in the planted rhizotron than in the unplanted one. Similar findings about PAH-degrading bacterial activity were reported in batch experiments with sand spiked with phenanthrene (Louvel et al. 2011). After 6 days, plants had already shaped the bacterial community structure: several OTUs belonging to *Beta*- and *Gamma-Proteobacteria* were significantly more active in the planted rhizotron, while other OTUs belonging to *Alpha*-, *Beta-Proteobacteria* and *Actinobacteria* were favoured in the unplanted rhizotron. In our experiment, after 12 days, root length and sugar content were significantly higher than after 6 days, and root biomass and DOC tended to increase, suggesting a stronger rhizosphere effect. At this sampling time, significantly higher PAH-degradation was observed in the planted rhizotron than in the unplanted one. This could be linked to an increasing level of PAH-dioxygenase transcription in both Gram-negative and Gram-positive PAH-degrading bacterial populations that was significantly higher in the planted soil than in the unplanted one. Our results clearly show a positive and rapid impact of the rhizosphere on the PAH degradation potential *via* the increased activity of a wider diversity (both GN and GP) of PAH-degrading bacteria. Even if *Actinobacteria* were enriched in the unplanted soil, the activity of the PAH degraders belonging to this phylum was favored in the rhizosphere. The higher activity of Gram-positive PAH degraders could help in the degradation of more recalcitrant PAH compounds (Kanaly and Harayama 2000). We highlighted a time-dependent effect in the course of the process dynamics, which could explain why some authors did not observe an increase in the activity of PAH-degrading populations in certain rhizospheres (Kawasaki et al. 2015). The addition of artificial root exudates can increase the density of soil PAH degraders, with an increase (Joner et al. 2002) or no difference (Cébron et al. 2011) in the degradation of spiked PAHs. We showed the activity of growing roots in a more natural rhizosphere system. A rhizotron device is probably the most adapted laboratory system to study the dynamics of rhizosphere processes because it makes it possible to perform analyses on precisely located samples with specific characteristics (depth, root density, roots age...) instead of mean rhizosphere samples usually obtained from pot experiments. Rhizotrons can allow studying the root turnover (Leigh et al. 2002), the isothiocyanate degradation rate, microbial communities (Rumberger and Marschner 2003) or spatial heterogeneity of PAH degradation and microbial communities (Bourceret et al. 2015).

In our study, the carbon released by actively growing roots and also leached from the top part of the device presenting a higher root density may have activated PAH degradation through co-metabolism (Ambrosoli et al. 2005). This phenomenon could also be attributed to a

priming effect, whereby the release of labile carbon stimulates microbial density and activity (Haichar et al. 2014), promoting PAH degradation or biotransformation (Rentz et al. 2005). As shown by the canonical analysis the higher differentiation between the planted and unplanted conditions was observed after 6 and 12 days of incubation. Root growth seemed to activate bacterial population mostly belonging to *Proteobacteria*, such as OTUs affiliated to the *Gamma-* (*Pseudomonadaceae*, *Stenotrophomonas* and *Acinetobacter*), *Alpha-* (*Sphingobium*, *Devosia*), and *Beta-* (*Methylophilaceae*, *Janthinobacterium* and *Achromobacter*) subdivisions. In addition to their ability to degrade PAHs (Fernández-Luqueño et al. 2011; Tauler et al. 2015; Gupta et al. 2015), OTUs affiliated to *Proteobacteria* can be favoured in the presence of plants as compared to unplanted soil (Thomas and Cébron 2016; Thomas et al. 2016). On the contrary, in unplanted soil other members of the *Alpha* (*Sphingomonadaceae*, *Sphingobium*, *Sphingomonas*, *Mycoplana*, *Magnetospirillum*) and *Beta* (*Ralstonia*, *Oxalobacteraceae*, *Comamonadaceae*, *Thiobacillus*) subdivisions were favoured, as well as *Actinobacteria* (*Iamia*, *Geodermatophilaceae*, *Georgenia*, and *Micrococcaceae*). After 16 and 22 days of root development, when no further PAH dissipation was observed, the transcriptional level of the PAH-dioxygenase gene from Gram-negative and Gram-positive bacteria decreased or remained constant over time in both planted and unplanted soils. At these time points, root development seemed to mainly influence the structure of the bacterial community. The proportion of many OTUs indeed differed between the planted and unplanted soils, but organic pollutant degradation no longer did.

Conclusion

Our results suggest that the dissipation of spiked PAHs is very fast in the soil when these pollutants represent the main available carbon sources for microbial communities and PAH-degrading bacteria are present. The presence of plants, which release root exudates, could favour microbial degradation by co-metabolism and increase the PAH degradation yield. The transcription of the PAH-dioxygenase gene was higher in the presence of plants than in unplanted soil, and transcription from Gram-negative bacteria (*Proteobacteria*) was high during the first 12 days when most of the PAHs were degraded. In the planted condition, Gram-positive PAH-degrading bacteria (*Actinobacteria*) were also highly active, and probably contributed to the higher level of PAH degradation in the rhizosphere. The characterization of the bacterial community structure and activity over time indicated different dynamics between the two conditions, with *Beta-* and *Gamma-Proteobacteria* mostly active

in planted soil while *Alpha-Proteobacteria* and *Actinobacteria* were more active in unplanted soil.

Acknowledgments:

This study was part of the RhizOrg project funded by the Agence Nationale de la Recherche (ANR-13-JSV7-0007_01). The authors thank the GISFI (www.gisfi.fr) for soil providing, David Billet, Geraldine Kitzinger and Gilles Bessaque for technical assistance. Carbohydrate analyses were performed by Jacqueline Marchand belonging to the PTEF facility that is supported by the French national Research Agency through the laboratory of Excellence ARBRE (ANR-11-LABX-0002-01).

Figure captions

Table 1. Nitrogenous and oxygenated pPAC concentration (mg.kg^{-1}) over time (at T0 and 6 days) in planted (Alf) and unplanted (BS) soils (Means $\pm$ SD, $n=6$). Letters indicate significant differences in pPAC concentrations between T0 and 6 days, and depending on presence of plant (one-way ANOVA).

Table 2. Bacterial richness (Chao1) and diversity estimators (Inverse of Simpson's index and Evenness) in unplanted soil (BS) and soil planted with alfalfa (Alf) at the beginning (T0) and after 6, 12, 16 and 22 days of incubation from cDNA data (Means $\pm$ SD, $n=4$). The plant effect was tested (from 6 to 22 days) using a two-way analysis of variance (ANOVA) followed by Newman-Keuls multiple comparison test (letters represents groups with significant different values ($p<0.05$)). One-way ANOVA was used to test time effect (from T0 to 22 days), separately for each condition.

Figure 1. Experimental devices and sampling points. (A) The device was composed of a mini-rhizotron (in orange) fitting in a corresponding rhizotron (in black). According to the sampling map (in white) at the depth 3 cm, six samples (in green) were collected, 3 samples (in gray) were used for soil humidity measurements, and 4 supplemental samples collected between green locations allowed quantification of dissolved organic carbon (DOC). (B) Pictures of alfalfa rhizotron devices after 6, 12, 16 and 22 days of plant growth.

Figure 2. Soil and root colonization parameters over time. (A) Dry root biomass and length, (B) sugar content, (C) dissolved organic carbon (DOC) and (D) pH after 6, 12, 16 and 22 days in alfalfa planted (green) and unplanted (brown) soils. Means (n=6 for root biomass and length, and n=4 for pH and DOC) and standard errors. Letters indicate significant (p<0.05) combined effect of time and presence of plant evaluated by using two-way ANOVA followed by Neuman-Keuls multiple comparison test.

Figure 3. PAH concentration (sum of 13 PAH, i.e. the 16 US-EPA PAHs, except naphthalene, acenaphthylene and acenaphthene) after 6, 12, 16 and 22 days in planted (green) and unplanted (brown) soils. Box plots show data variability and means (black cross), n=24 (for T0 samples) and n=6 (from 6 to 22 day samples). Dashed line indicate mean of PAH concentration in the NM soil before spiking. Black asterisks indicate significant differences between planted and unplanted soil (Student t test, p<0.05) and letters indicate significant differences depending on time (one-way ANOVA).

Figure 4. Abundance values of genes (A) and transcripts (B) and ratio of transcripts relative to their corresponding genes (C) for fungal 18S rRNA, bacterial 16S rRNA and PAH-degrading bacterial communities after 6, 12, 16 and 22 days in alfalfa planted (green) and unplanted (brown) soils. Values are means and standard deviations of 3 replicates (T0) and 6 replicates for 6 to 22 days. The effect of time was evaluated for each condition (planted, unplanted) by using one-way ANOVA (p<0.05) and presence of plant was evaluated between 6 and 22 days by using two-way ANOVA (p<0.05) followed by Neuman Keuls multiple comparison test. In A and B, different letters indicate significant changes in microbial community abundance depending on time for each condition, black asterisks indicate plant effect. In C, letters indicate cross effect of plant and time on the ratio of transcript relative to gene copies.

Figure 5. Relative proportion (mean, n=4) of dominant active bacterial phyla ($\geq 1\%$ in at least one sample) in alfalfa planted (Alf) and unplanted soil (BS) at 0, 6, 12, 16 and 22 days of incubation. The twelve significant phyla were identified by a Tukey Kramer ANOVA with a Benjamini Hochberg correction of p-value (q<0.05, and effect size >0.26).

Figure 6. Ratio of the 16S rRNA relative to 16S rDNA for OTU (≥ 10 sequences in one sample) significantly discriminating plant effect, after 6 and 12 days of incubation. Both

conditions were compared using Student t-tests with Benjamini-Hochberg correction of p-value. Asterisks denote degree of significance (green and brown for higher ratio with alfalfa and unplanted soils, respectively; * $q < 0.05$; ** $q < 0.01$; *** $q < 0.001$). OTUs were affiliated at 97% of similarities to the deeper taxonomic level, leading to affiliations at different levels.

Figure 7. Canonical correspondence analysis (CCA) ordination including abiotic (A) and (B) biotic variables. (A) environmental variables (in red) (DOC, sugar and organic acid contents, pH, percentage of 18S rRNA, PAH-RHD_α GN and GP genes and transcripts relative to 16S rRNA), root biomass (RootB) and length (RootL), sum of 13 PAH concentrations, 2-3-ring PAHs (fluorene, phenanthrene, anthracene), 4-ring PAHs (fluoranthene, pyrene, benzo(a)anthracene, chrysene), 5-6-ring PAHs (benzo(b)fluoranthene, benzo(k)fluoranthene, benzo(a)pyrene, dibenzo(a,h)anthracene, benzo(g,h,i)perylene, indeno(1,2,3-cd)pyrene), and (B) the relative abundance of dominant active bacterial OTUs ($\geq 1\%$ in at least one sample, taxonomic affiliation at 97% of similarities to the deeper taxonomic level, leading to affiliations at different levels) (in black) were used to discriminate the 24 samples (4 replicates for each sampling time corresponding to 0, 6, 12, 16 and 22 days) collected for alfalfa planted (green) and unplanted (brown) rhizotrons. A Monte Carlo permutation test (1000 permutations) was performed and showed a significant ($p < 0.0001$) effect of environmental variables on the relative abundance of dominant OTUs. Un. Coma.: Un. Comamonadaceae; Un. Pseudo.: Un. Pseudomonadaceae; Un. Sphingo.: Un. Sphingomonadaceae; Un. Geo.: Un. Geodermatophilaceae; Un. Meth.: Un. Methylobacteriaceae; Phen.: Phenylobacterium; Phaeos. Fulvum: Phaeosporillum Fulvum; Un. Xantho.: Un. Xanthomonadaceae; Un. Oxalo.: Un. Oxalobacteraceae.

Supplemental data

Table S1. PAH concentration (mg.L^{-1}) in the organic extract used to spike NM soil. Values are means ($n=3$) and standard errors.

Table S2. Pearson's correlation matrix between soil properties and the percentage of PAH-dioxygenase transcripts and genes. Values in bold correspond to significant correlation between variables ($p < 0.05$).

Figure S1. Bioavailable (A) and total (B) PAH concentration in NM soil before (grey) and after (blue) spiking. Black asterisks indicate significant differences between both conditions

(Student t test, $p < 0.05$). Values are means and standard deviation of 12 and 48 samples (total fraction) and 8 and 6 samples (bioavailable fraction), before and after spiking respectively.

Figure S2. Total PAH concentration of 13 individual compounds (the 16 US-EPA PAHs except naphthalene, acenaphthylene and acenaphthene) after 6, 12, 16 and 22 days in alfalfa planted (green) and unplanted (brown) soils. Values are means and standard deviations of 24 replicates (0 day) and 6 replicates from 6 to 22 days. The effect of time was evaluated for each condition (planted, unplanted) independently by using one-way ANOVA ($p < 0.05$) and effect of presence of plant was evaluated between T1 and T4 by using two-way ANOVA ($p < 0.05$) followed by Neuman Keuls multiple comparison test. Different letters indicate significant differences in PAH concentration depending on time for each condition, black asterisks indicate plant effect.

Figure S3. Evolution over time of the relative proportion of dominant OTUs ($\geq 1\%$ in at least one sample), in alfalfa planted and unplanted soil, reported to the transcript 16S rRNA copy number based on cDNA analysis. Forty-seven significant OTUs were identified by a Tukey Kramer ANOVA with a Benjamini Hochberg correction of p-value ($q < 0.05$, and effect size > 0.26) and non-significant OTUs correspond to population presenting constant relative proportion over time. Asterisks denote OTUs with significant differences in relative abundance over time specifically in planted (green) or in unplanted soil (brown). OTUs were affiliated at the most precise level.

Figure S4. Ratio of the 16S rRNA relative to 16S rDNA for OTU (with at least 10 sequences in one sample) significantly discriminating plant effect, after 16 and 22 days of incubation. Both conditions were compared using Student t-tests with Benjamini-Hochberg correction of p-value. Asterisks denote degree of significance (green and brown for higher ratio with alfalfa planted and unplanted soils, respectively; * $q < 0.05$; ** $q < 0.01$; *** $q < 0.001$).

References

Allard A-S, Remberger M, Neilson AH (2000) The negative impact of aging on the loss of PAH components in a creosote-contaminated soil. *Int Biodeterior Biodegradation* 46:43–49 . doi: 10.1016/S0964-8305(00)00050-0

691 Ambrosoli R, Petruzzelli L, Minati JL, Marsan FA (2005) Anaerobic PAH degradation in soil
 692 by a mixed bacterial consortium under denitrifying conditions. *Chemosphere* 60:1231–
 693 1236 . doi: 10.1016/j.chemosphere.2005.02.030

694 Baudoin E, Benizri E, Guckert A (2002) Impact of growth stage on the bacterial community
 695 structure along maize roots, as determined by metabolic and genetic fingerprinting. *Appl*
 696 *Soil Ecol* 19:135–145 . doi: 10.1016/S0929-1393(01)00185-8

697 Bell TH, El-Din Hassan S, Lauron-Moreau A, et al (2014) Linkage between bacterial and
 698 fungal rhizosphere communities in hydrocarbon-contaminated soils is related to plant
 699 phylogeny. *Isme J* 8:331–343 . doi: 10.1038/ismej.2013.149

700 Benhabib K, Faure P, Sardin M, Simonnot MO (2010) Characteristics of a solid coal tar
 701 sampled from a contaminated soil and of the organics transferred into water. *Fuel*
 702 89:352–359 . doi: 10.1016/j.fuel.2009.06.009

703 Bevivino A, Dalmastri C (2017) Composition and Function of Microbial Communities in Soil
 704 Ecosystems. In: *Soil Biological Communities and Ecosystem Resilience*. pp 79–81

705 Biache C, Ouali S, Cébron A, et al (2017) Bioremediation of PAH-contaminated soils:
 706 Consequences on formation and degradation of polar-polycyclic aromatic compounds
 707 and microbial community abundance. *J Hazard Mater* 329:1–10 . doi:
 708 10.1016/j.jhazmat.2017.01.026

709 Bogan BW, Sullivan WR (2003) Physicochemical soil parameters affecting sequestration and
 710 mycobacterial biodegradation of polycyclic aromatic hydrocarbons in soil. *Chemosphere*
 711 52:1717–1726 . doi: 10.1016/S0045-6535(03)00455-7

712 Bourceret A, Cébron A, Tisserant E, et al (2016) The Bacterial and Fungal Diversity of an
 713 Aged PAH- and Heavy Metal-Contaminated Soil is Affected by Plant Cover and
 714 Edaphic Parameters. *Microb Ecol* 1–14 . doi: 10.1007/s00248-015-0682-8

715 Bourceret A, Leyval C, De Fouquet C, Cébron A (2015) Mapping the centimeter-scale spatial
 716 variability of PAHs and microbial populations in the rhizosphere of two plants. *PLoS*
 717 *One* 10:

718 Bourceret A, Leyval C, Thomas F, Cébron A (2017) Rhizosphere effect is stronger than PAH
 719 concentration on shaping spatial bacterial assemblages along centimetre-scale depth

720 gradients. *Can J Microbiol* 63:881–893 . doi: 10.1139/cjm-2017-0124

721 Caporaso JG, Kuczynski J, Stombaugh J, et al (2010) QIIME allows analysis of high-
 722 throughput community sequencing data. *Nat Methods* 7:335–336 . doi:
 723 10.1038/nmeth.f.303

724 Castro-Silva C, Ruíz-Valdiviezo VM, Valenzuela-Encinas C, et al (2013) The bacterial
 725 community structure in an alkaline saline soil spiked with anthracene. *Electron J*
 726 *Biotechnol* 16:1–20 . doi: 10.2225/vol16-issue5-fulltext-14

727 Cébron A, Beguiristain T, Faure P, et al (2009) Influence of vegetation on the in situ bacterial
 728 community and polycyclic aromatic hydrocarbon (PAH) degraders in aged PAH-
 729 contaminated or thermal-desorption-treated soil. *Appl Environ Microbiol* 75:6322–6330
 730 . doi: 10.1128/AEM.02862-08

731 Cébron A, Cortet J, Criquet S, et al (2011) Biological functioning of PAH-polluted and
 732 thermal desorption-treated soils assessed by fauna and microbial bioindicators. *Res*
 733 *Microbiol* 162:897–907 . doi: 10.1016/j.resmic.2011.02.011

734 Cébron A, Faure P, Lorgeoux C, et al (2013) Experimental increase in availability of a PAH
 735 complex organic contamination from an aged contaminated soil: Consequences on
 736 biodegradation. *Environ Pollut* 177:98–105 . doi: 10.1016/j.envpol.2013.01.043

737 Cébron A, Norini MP, Beguiristain T, Leyval C (2008) Real-Time PCR quantification of
 738 PAH-ring hydroxylating dioxygenase (PAH-RHD α) genes from Gram positive and
 739 Gram negative bacteria in soil and sediment samples. *J Microbiol Methods* 73:148–159 .
 740 doi: 10.1016/j.mimet.2008.01.009

741 Cerniglia CE, Sutherland JB (2010) Degradation of polycyclic aromatic hydrocarbons by
 742 fungi. In: (Berlin: Springer;) (ed) *Handbook of Hydrocarbon and Lipid Microbiology*,
 743 Timmis K. pp 2080–2110

744 D’Orazio V, Ghanem A, Senesi N (2013) Phytoremediation of pyrene contaminated soils by
 745 different plant species. *Clean - Soil, Air, Water* 41:377–382 . doi:
 746 10.1002/clen.201100653

747 de Menezes A, Clipson N, Doyle E (2012) Comparative metatranscriptomics reveals
 748 widespread community responses during phenanthrene degradation in soil. *Environ*

749 Microbiol 14:2577–2588 . doi: 10.1111/j.1462-2920.2012.02781.x

750 Felske A, Akkermans ADL, De Vos WM (1998) Quantification of 16S rRNAs in complex
 751 bacterial communities by multiple competitive reverse transcription-PCR in temperature
 752 gradient gel electrophoresis fingerprints. *Appl Environ Microbiol* 64:4581–4587

753 Fernández-Luqueño F, Valenzuela-Encinas C, Marsch R, et al (2011) Microbial communities
 754 to mitigate contamination of PAHs in soil—possibilities and challenges: a review.
 755 *Environ Sci Pollut Res* 18:12–30

756 Gartler J, Wimmer B, Soja G, Reichenauer TG (2014) Effects of Rapeseed Oil on the
 757 Rhizodegradation of Polyaromatic Hydrocarbons in Contaminated Soil. *Int J*
 758 *Phytoremediation* 16:671–683 . doi: 10.1080/15226514.2013.856841

759 Ghosal D, Ghosh S, Dutta TK, Ahn Y (2016) Current state of knowledge in microbial
 760 degradation of polycyclic aromatic hydrocarbons (PAHs): A review. *Front Microbiol* 7: .
 761 doi: 10.3389/fmicb.2016.01369

762 Gupta S, Pathak B, Fulekar MH (2015) Molecular approaches for biodegradation of
 763 polycyclic aromatic hydrocarbon compounds: a review. *Rev Environ Sci Biotechnol*
 764 14:241–269 . doi: 10.1007/s11157-014-9353-3

765 Haichar F el Z, Santaella C, Heulin T, Achouak W (2014) Root exudates mediated
 766 interactions belowground. *Soil Biol Biochem* 77:69–80 . doi:
 767 10.1016/j.soilbio.2014.06.017

768 Haichar F el Z, Marol C, Berge O, et al (2008) Plant host habitat and root exudates shape soil
 769 bacterial community structure. *ISME J* 2:1221–1230 . doi: 10.1038/ismej.2008.80

770 Hannula SE, de Boer W, van Veen JA (2010) In situ dynamics of soil fungal communities
 771 under different genotypes of potato, including a genetically modified cultivar. *Soil Biol*
 772 *Biochem* 42:2211–2223 . doi: 10.1016/j.soilbio.2010.08.020

773 Hanser O, Biache C, Boulangé M, et al (2015) Evolution of dissolved organic matter during
 774 abiotic oxidation of coal tar—comparison with contaminated soils under natural
 775 attenuation. *Environ Sci Pollut Res* 22:1431–1443

776 Haritash AK, Kaushik CP (2009) Biodegradation aspects of Polycyclic Aromatic
 777 Hydrocarbons (PAHs): A review. *J Hazard Mater* 169:1–15 . doi:

778 10.1016/j.jhazmat.2009.03.137

779 Hinsinger P, Gobran GR, Gregory PJ, Wenzel WW (2005) Rhizosphere geometry and
780 heterogeneity arising from root-mediated physical and chemical processes. *New Phytol*
781 168:293–303 . doi: 10.1111/j.1469-8137.2005.01512.x

782 Joner EJ, Corgié SC, Amellal N, Leyval C (2002) Nutritional constraints to degradation of
783 polycyclic aromatic hydrocarbons in a simulated rhizosphere. *Soil Biol Biochem*
784 34:859–864 . doi: 10.1016/S0038-0717(02)00018-4

785 Kanaly R, Harayama S (2000) Biodegradation of High-Molecular-Weight Polycyclic
786 Aromatic Hydrocarbons by Bacteria MINIREVIEW Biodegradation of High-Molecular-
787 Weight Polycyclic Aromatic Hydrocarbons by Bacteria. *J Bacteriol* 182:2059–2067 .
788 doi: 10.1128/JB.182.8.2059-2067.2000.Updated

789 Kawasaki A, Warren CR, Kertesz MA (2015) Specific influence of white clover on the
790 rhizosphere microbial community in response to polycyclic aromatic hydrocarbon (PAH)
791 contamination. *Plant Soil* 1–15

792 Kawasaki A, Watson E, Kertesz M (2012) Indirect effects of polycyclic aromatic hydrocarbon
793 contamination on microbial communities in legume and grass rhizospheres. *Plant Soil*
794 358:169–182 . doi: 10.1007/s11104-011-1089-z

795 Kim S-J, Kweon O, Freeman JP, et al (2006) Molecular Cloning and Expression of Genes
796 Encoding a Novel Dioxygenase Involved in Low- and High-Molecular-Weight
797 Polycyclic Aromatic Hydrocarbon Degradation in *Mycobacterium vanbaalenii* PYR-1.
798 *Appl Environ Microbiol* 72:1045–1054

799 Kirk JL, Klironomos JN, Lee H, Trevors JT (2005) The effects of perennial ryegrass and
800 alfalfa on microbial abundance and diversity in petroleum contaminated soil. *Environ*
801 *Pollut* 133:455–465 . doi: 10.1016/j.envpol.2004.06.002

802 Kunihiro M, Ozeki Y, Nogi Y, et al (2013) Benz[a]anthracene biotransformation and
803 production of ring fission products by *Sphingobium* sp. strain KK22. *Appl Environ*
804 *Microbiol* 79:4410–4420 . doi: 10.1128/AEM.01129-13

805 Leigh MB, Fletcher JS, Fu X, Schmitz FJ (2002) Root turnover: An important source of
806 microbial substrates in rhizosphere remediation of recalcitrant contaminants. *Environ Sci*

807 Technol 36:1579–1583 . doi: 10.1021/es015702i

808 Liste H-H, Alexander M (2000) Plant-promoted pyrene degradation in soil. *Chemosphere*
809 40:7–10 . doi: 10.1016/S0045-6535(99)00216-7

810 Louvel B, Cébron A, Leyval C (2011) Root exudates affect phenanthrene biodegradation,
811 bacterial community and functional gene expression in sand microcosms. *Int Biodeterior*
812 *Biodegrad* 65:947–953 . doi: 10.1016/j.ibiod.2011.07.003

813 Lueders T, Wagner B, Claus P, Friedrich MW (2004) Stable isotope probing of rRNA and
814 DNA reveals a dynamic methylotroph community and trophic interactions with fungi
815 and protozoa in oxic rice field soil. *Environ Microbiol* 6:60–72 . doi: 10.1046/j.1462-
816 2920.2003.00535.x

817 Lundstedt S, Persson Y, Öberg L (2006) Transformation of PAHs during ethanol-Fenton
818 treatment of an aged gasworks' soil. *Chemosphere* 65:1288–1294

819 Lundstedt S, White PA, Lemieux CL, et al (2007) Sources, fate, and toxic hazards of
820 oxygenated polycyclic aromatic hydrocarbons (PAHs) at PAH-contaminated sites.
821 *AMBIO A J Hum Environ* 36:475–485

822 McDonald D, Price MN, Goodrich J, et al (2012) An improved Greengenes taxonomy with
823 explicit ranks for ecological and evolutionary analyses of bacteria and archaea. *ISME J*
824 6:610–8 . doi: 10.1038/ismej.2011.139

825 Muratova AY, Turkovskaya O V., Hübner T, Kusch P (2003) Studies of the Efficacy of
826 Alfalfa and Reed in the Phytoremediation of Hydrocarbon-Polluted Soil. *Appl Biochem*
827 *Microbiol* 39:599–605 . doi: 10.1023/A:1026238720268

828 Northcott GL, Jones KC (2001) Partitioning, extractability, and formation of nonextractable
829 PAH residues in soil. 2. Effects on compound dissolution behavior. *Environ Sci Technol*
830 35:1111–1117

831 Parks DH, Beiko RG (2010) Identifying biologically relevant differences between
832 metagenomic communities. *Bioinformatics* 26:715–721

833 Peng A, Liu J, Ling W, et al (2015) Diversity and distribution of 16S rRNA and phenol
834 monooxygenase genes in the rhizosphere and endophytic bacteria isolated from PAH-
835 contaminated sites. *Sci Rep* 5:12173 . doi: 10.1038/srep12173

836 Reid BJ, Stokes JD, Jones KC, Semple KT (2000) Nonexhaustive cyclodextrin-based
837 extraction technique for the evaluation of PAH bioavailability. *Environ Sci Technol*
838 34:3174–3179 . doi: 10.1021/es990946c

839 Rentz JA, Alvarez PJJ, Schnoor JL (2005) Benzo[a]pyrene co-metabolism in the presence of
840 plant root extracts and exudates: Implications for phytoremediation. *Environ Pollut*
841 136:477–484 . doi: 10.1016/j.envpol.2004.12.034

842 Rollin C, Quiot F, DENYS S, BUREAU J (2005) Hydrocarbures Aromatiques Polycycliques.
843 Guide méthodologique. Acquisition des données d’entrée des modèles analytiques ou
844 numériques de transferts dans les sols et les eaux souterraines. INERIS–DRC–
845 66244/DESP–R01

846 Roy M, Khara P, Basu S, Dutta TK (2013) Catabolic Versatility of *Sphingobium* sp. Strain
847 PNB Capable of Degrading Structurally Diverse Aromatic Compounds. *J*
848 *Bioremediation Biodegrad* 4:1–6 . doi: 10.4172/2155-6199.1000173

849 Rumberger A, Marschner P (2003) 2-Phenylethylisothiocyanate concentration and microbial
850 community composition in the rhizosphere of canola. *Soil Biol Biochem* 35:445–452 .
851 doi: 10.1016/S0038-0717(02)00296-1

852 Sawulski P, Clipson N, Doyle E (2014) Effects of polycyclic aromatic hydrocarbons on
853 microbial community structure and PAH ring hydroxylating dioxygenase gene
854 abundance in soil. *Biodegradation* 25:835–847 . doi: 10.1007/s10532-014-9703-4

855 Siciliano SD, Germida JJ, Banks K, Greer CW (2003) Changes in Microbial Community
856 Composition and Function during a Polyaromatic Hydrocarbon Phytoremediation Field
857 Trial. *Appl Environ Microbiol* 69:483–489

858 Storey S, Ashaari MM, McCabe G, et al (2014) Microbial community structure during
859 fluoranthene degradation in the presence of plants. *J Appl Microbiol* 117:74–84 . doi:
860 10.1111/jam.12518

861 Tauler M, Vila J, Nieto JM, Grifoll M (2015) Key high molecular weight PAH-degrading
862 bacteria in a soil consortium enriched using a sand-in-liquid microcosm system. *Appl*
863 *Microbiol Biotechnol* 1–16

864 Tejeda-Agredano MC, Gallego S, Vila J, et al (2013) Influence of the sunflower rhizosphere

865 on the biodegradation of PAHs in soil. *Soil Biol Biochem* 57:830–840 . doi:
866 10.1016/j.soilbio.2012.08.008

867 Thion C, Cébron A, Beguiristain T, Leyval C (2013) Inoculation of PAH-degrading strains of
868 *Fusarium solani* and *Arthrobacter oxydans* in rhizospheric sand and soil microcosms:
869 microbial interactions and PAH dissipation. *Biodegradation* 24:569–581 . doi:
870 10.1007/s10532-013-9628-3

871 Thion C, Cébron A, Beguiristain T, Leyval C (2012) Long-term in situ dynamics of the fungal
872 communities in a multi-contaminated soil are mainly driven by plants. *FEMS Microbiol*
873 *Ecol* 82:169–181 . doi: 10.1111/j.1574-6941.2012.01414.x

874 Thomas F, Cébron A (2016) Short-term rhizosphere effect on available carbon sources,
875 phenanthrene degradation, and active microbiome in an aged-contaminated industrial
876 soil. *Front Microbiol* 7:1–15 . doi: 10.3389/fmicb.2016.00092

877 Thomas F, Lorgeoux C, Faure P, et al (2016) Isolation and substrate screening of polycyclic
878 aromatic hydrocarbon degrading bacteria from soil with long history of contamination.
879 *Int Biodeterior Biodegradation* 107:1–9

880 Trapp S, Karlson U (2001) Aspects of phytoremediation of organic pollutants. *J Soils*
881 *Sediments* 1:37–43 . doi: 10.1065/jss2001.04.007

882 Turner S, Pryer KM, Miao VP, Palmer JD (1999) Investigating deep phylogenetic
883 relationships among cyanobacteria and plastids by small subunit rRNA sequence
884 analysis1. *J Eukaryot Microbiol* 46:327–338

885 van Herwijnen R, Wattiau P, Bastiaens L, et al. (2003) Elucidationn of the metabolic pathway
886 of fluorene and cometabolicc pathways of phenanthrene , fluoranthene , anthracenee and
887 dibenzothiophene by *Sphingomonas* sp . *Res Microbiol* 154:199–206

888 Vrchotová B, Lovecká P, Dražková M, et al (2013) Influence of root exudates on the bacterial
889 degradation of chlorobenzoic acids. *Sci World J* 2013: . doi: 10.1155/2013/872026

890 Walker TS, Bais HP, Grotewold E, Vivanco JM (2003) Root exudation and rhizosphere
891 biology. *Plant Physiol* 132:44–51

892 Wang Q, Garrity GM, Tiedje JM, Cole JR (2007) Naïve Bayesian Classifier for Rapid
893 Assignment of rRNA Sequences into the New Bacterial Taxonomy. *Appl Environ*

894 Microbiol 73:5261–5267

895 Yang C-H, Crowley DE (2000) Rhizosphere Microbial Community Structure in Relation to
896 Root Location and Plant Iron Nutritional Status. Appl Environ Microbiol 66:345–351 .
897 doi: 10.1128/AEM.66.1.345-351.2000

898 Yergeau E, Sanschagrin S, Maynard C, et al (2014) Microbial expression profiles in the
899 rhizosphere of willows depend on soil contamination. ISME J 8:344–58 . doi:
900 10.1038/ismej.2013.163

901 Zhang Z, Zhao X, Liang Y, et al (2013) Microbial functional genes reveal selection of
902 microbial community by PAHs in polluted soils. Environ Chem Lett 11:11–17 . doi:
903 10.1007/s10311-012-0370-6

904 Zhou W, Yang J, Lou L, Zhu L (2011) Solubilization properties of polycyclic aromatic
905 hydrocarbons by saponin, a plant-derived biosurfactant. Environ Pollut 159:1198–1204

906

907

909

910

911 Figure 2.

912

913

914 Figure 3.

915

916 Figure 4.

917

918

921

922 Figure 6.

923

924

929 Figure S1.

930

931 Figure S2.

932

933 Figure S3.

934

935

936 Figure S4

937

938

939 **Table S2.** PAH concentration (mg.L⁻¹) in the organic extract used to spike NM soil. Values are means (n=3) and standard errors.

940

941

942

PAHs concentration (mg/L)												
Fluorene	Phenanthrene	Anthracene	Fluoranthene	Pyrene	Benzo(a) anthracene	Chrysene	Benzo(b) fluoranthene	Benzo(k) fluoranthene	Benzo(a) pyrene	Dibenzo(a,h) anthracene	Benzo(g,h,i) perylene	Indeno(1,2,3-cd) pyrene
24.8 ± 2.3	98.7 ± 4.2	35.9 ± 1.6	221.8 ± 16.7	200.9 ± 11.7	126.3 ± 12.2	101.1 ± 6.8	115.7 ± 10.8	70.7 ± 10.3	28.2 ± 7.6	70.3 ± 7.0	77.2 ± 7.8	101.7 ± 10.3

943

Variables	Time	2-3 ring PAH	4 ring PAH	5-6 ring PAH	Σ13 PAHs	pH	DOC	RootB	RootL
%18S_DNAc	-0.646	0.686	0.451	0.361	0.474	-0.603	-0.028	-0.360	-0.338
% PAH-RHDα GN_DNAc	0.106	-0.434	-0.391	-0.220	-0.339	0.059	-0.094	0.062	0.010
% PAH-RHDα GP_DNAc	0.733	-0.204	-0.253	0.137	-0.053	0.409	0.154	0.452	0.538
%18S_DNAg	-0.502	0.283	0.199	0.064	0.154	-0.386	-0.151	-0.363	-0.485
% PAH-RHDα GN_DNAg	-0.032	-0.512	-0.448	-0.530	-0.547	0.213	-0.029	0.349	0.052
% PAH-RHDα GP_DNAg	0.497	0.163	0.094	0.388	0.277	0.170	0.092	0.049	0.120

944

945

946

947

948

949 Table S1. Pearson's correlation matrix between soil properties and the percentage of PAH-dioxygenase transcript and genes. Values in bold correspond to
950 significant correlation between variables ($p < 0.05$).

951