The plastoquinone pool outside the thylakoid membrane serves in plant photoprotection as a reservoir of singlet oxygen scavengers

Brigitte Ksas, Bertrand Legeret, Ursula Ferretti, Anne Chevalier, Pavel Pospíšil, Jean Alric, Michel Havaux

► To cite this version:

Brigitte Ksas, Bertrand Legeret, Ursula Ferretti, Anne Chevalier, Pavel Pospíšil, et al.. The plastoquinone pool outside the thylakoid membrane serves in plant photoprotection as a reservoir of singlet oxygen scavengers. *Plant, Cell and Environment*, 2018, 41 (10), pp.2277-2287. 10.1111/pce.13202 . hal-01946443

HAL Id: hal-01946443

<https://hal.science/hal-01946443>

Submitted on 8 Jan 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Running title: Plastoquinone as an antioxidant

The plastoquinone pool outside the thylakoid membrane serves in plant photoprotection as a reservoir of singlet oxygen scavengers

Brigitte Ksas¹, Bertrand Legeret², Ursula Ferretti³, Anne Chevalier¹, Pavel Pospíšil³, Jean Alric¹, Michel Havaux¹

¹CEA Cadarache, CNRS UMR 7265 Biologie Végétale et Microbiologie Environnementales, Aix Marseille Université, Laboratoire d'Ecophysiologie Moléculaire des Plantes, 13108 Saint-Paul-lez-Durance, France

²CEA Cadarache, CNRS UMR 7265 Biologie Végétale et Microbiologie Environnementales, Aix Marseille Université, Laboratoire de Bioénergétique et Biotechnologie des Bactéries et Microalgues, 13108 Saint-Paul-lez-Durance, France

³Department of Biophysics, Centre of the Region Haná for Biotechnological and Agricultural Research, Faculty of Science, Palacký University, Šlechtitelů 27, 783 71, Olomouc, Czech Republic.

Correspondence

Michel Havaux. Email: michel.havaux@cea.fr

Abstract

The Arabidopsis *vte1* mutant is devoid of tocopherol and plastochromanol (PC-8). When exposed to excess light energy, *vte1* produced more singlet oxygen ($^1\text{O}_2$) and suffered from extensive oxidative damage compared to the wild type. Here we show that overexpressing the *SOLANESYL DIPHOSPHATE SYNTHASE 1 (SPS1)* gene in *vte1* induced a marked accumulation of plastoquinone (PQ-9) and rendered the *vte1 SPS1oex* plants tolerant to photooxidative stress, indicating that PQ-9 can replace tocopherol and PC-8 in photoprotection. High PQ-9 levels were associated with a noticeable decrease in $^1\text{O}_2$ production and with higher levels of PQ-C, a $^1\text{O}_2$ -specific PQ-9 oxidation product. The extra PQ-9 molecules in the *vte1 SPS1oex* plants were stored in the plastoglobules and the chloroplast envelopes, rather than in the thylakoid membranes, whereas PQ-C was found almost exclusively in the thylakoid membranes. Upon exposure of wild-type plants to high light, the thylakoid PQ-9 pool decreased, while the extra-thylakoid pool remained unchanged. In *vte1* and *vte1 SPS1oex* plants, the PQ-9 losses in high light were strongly amplified, affecting also the extra-thylakoid pool, and PQ-C was found in high amounts in the thylakoids. We conclude that the thylakoid PQ-9 pool acts as a $^1\text{O}_2$ scavenger and is replenished from the extra-thylakoid stock.

KEYWORDS

plastoquinone, antioxidant, reactive oxygen species, photoprotection, tocopherol

1 INTRODUCTION

Plastoquinone-9 (PQ-9) is essentially viewed as a component of the photosynthetic electron transport chain in chloroplasts, carrying electrons from photosystem II (PSII) to cytochrome *b6/f* (Amesz, 1973). It is thus essential for the photosynthetic activity of plants, and mutations inhibiting PQ-9 accumulation led to a lethal phenotype in Arabidopsis and maize (Cook & Miles, 1992; Norris, Barrette, & DellaPenna, 1995). Through its redox state, the PQ-9 pool can also play a role in the regulation of gene expressions and enzyme activities (Pfannschmidt et al., 2009; Rochaix, 2014). However, several observations obtained in recent *in vitro* and *in vivo* studies have raised the idea that the physiological role of PQ-9 can be more diverse than just electron shuttling from PSII to PSI. In particular, *in vitro* studies have revealed that PQ-9 has antioxidant properties, being able to quench singlet oxygen ($^1\text{O}_2$) and to inhibit oxidation of lipid membranes (Gruszka, Pawlak, & Kruk, 2008; Nowicka, Gruszka, & Kruk, 2013; Yadav, Kruk, Sinha, & Pospisil, 2010). In isolated thylakoids, maintenance of the PQ-9 pool in the reduced state with an electron transport inhibitor was associated with a lowering of high light-induced lipid

63 peroxidation, suggesting an antioxidant role for reduced PQ-9 (Hundal, Forsmark-Andrée, Ernster, &
64 Andersson, 1995).

65 Tocopherol and plastoquinone are synthesized through a common pathway until
66 homogentisate (HGA) where SOLANESYL DIPHOSPHATE SYNTHASE 1 (SPS1) introduces the bifurcation
67 towards PQ-9 and PC-8. Accumulation of PQ-9 in leaves by overexpression of the *SPS1* gene in *SPS1oex*
68 *Arabidopsis* plants reduced lipid peroxidation and increased the tolerance to photooxidative stress
69 (Ksas, Becuwe, Chevalier, & Havaux, 2015). However, because the *SPS1oex* plants also accumulated
70 plastochroman-8 (PC-8), a PQ-9 metabolite with known antioxidative activities (Kruk, Szymanska,
71 Cela, & Munne-Bosch, 2014), photoprotection could not be unambiguously and exclusively attributed
72 to PQ-9. Nevertheless, sudden exposure of *Arabidopsis* plants to high light stress resulted in a marked
73 loss of PQ-9 (Ksas et al., 2015) and to an accumulation of oxidized derivatives formed by the interaction
74 between PQ-9 and $^1\text{O}_2$ (Szymanska & Kruk, 2010), supporting the idea that PQ-9 can act as a $^1\text{O}_2$
75 scavenger *in vivo*. Similarly, in the green microalga *Chlamydomonas*, PQ-9 was observed to decrease
76 in high light when its re-synthesis was blocked by an inhibitor of hydroxylphenylpyruvate dioxygenase
77 (Kruk & Trebst, 2008).

78 In this context, an interesting observation was the various localizations of PQ-9 in thylakoid
79 membranes, plastoglobules (Lichtenthaler & Weinert, 1970; Zbierzak et al., 2009) and chloroplast
80 envelopes (Soll, Schultz, Joyard, Douce, & Block, 1985), suggesting that PQ-9 could play diverse roles
81 in the chloroplasts. To investigate these roles, we have crossed the *vte1* *Arabidopsis* mutant, deficient
82 in both tocopherols and PC-8 (Porfiriva, Bergmuller, Tropsch, Lemke, & Dörmann, 2002), with *SPS1oex*
83 lines previously described (Ksas et al., 2010) to generate *vte1 SPS1oex* plants that selectively
84 accumulate PQ-9 in the absence of PC-8. The responses of *vte1 SPS1oex* plants to high light together
85 with the analyses of PQ-9 localization and concentration demonstrate that PQ-9 is a $^1\text{O}_2$ -scavenging
86 antioxidant *in planta* and show the participation of the extra-thylakoid PQ-9 pool in this process
87 through delivery of PQ-9 to the thylakoids in order to replace oxidized PQ-9 molecules.

89 2 MATERIAL AND METHODS

90 2.1 Plant material and growth conditions

91 *Arabidopsis* plants (*Arabidopsis thaliana*, Columbia-0 ecotype) were grown in a phytotron under
92 controlled conditions of light ($150 \mu\text{mol photons m}^{-2} \text{s}^{-1}$, photoperiod 8 h), temperature ($20^\circ\text{C}/18^\circ\text{C}$,
93 day/night) and relative air humidity (60%). Several mutant/transformed plants were examined in this
94 work: the tocopherol cyclase mutant *vte1* deficient in tocopherols (Porfiriva et al., 2002), *SPS1*-
95 overexpressing lines described in a previous work (lines #12 and #14 in Ksas et al. (2015)) and a crossing
96 between them (*vte1 SPS1oex*). Selection of homozygous *vte1 SPS1oex* plants was achieved first by

screening the progeny for resistance to the herbicide BASTA and then by screening for total absence of tocopherol and PC-8 by HPLC analyses (see below for HPLC method). Photooxidative stress was induced by transferring plants to a growth chamber at 7°C/15°C (day/night, air temperature) under a PFD of 1500 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ and a photoperiod of 8 h, as previously explained (Ksas et al., 2015). Depending of the experiment, the duration of the treatment was 48 h (photooxidative stress experiments) or 8 h (dynamics of the PQ-9 pools), as indicated in the legend of the figures.

2.2 Prenyl lipid determinations

Leaf discs (3 discs, diameter 1 cm) were grinded in ethyl acetate. After centrifugation, the supernatant was filtered and evaporated on ice under a stream of N_2 . The residue was recovered in methanol/hexane (17/1) and analyzed by HPLC, as described elsewhere (Ksas et al., 2015; Szymanska, Nowicka, & Kruk, 2014; Kruk & Karpinski, 2006). The column was a Phenomenex Kinetex 2.6 μm , 100 x 4.6 mm, 100 Å. Separation of tocopherols, plastoquinone-9 and plastochromanol-8 was done in the isocratic mode with methanol/hexane (17/1) as solvent system and a flow rate of 0.8 ml min^{-1} . All prenyl lipids, except oxidized plastoquinone-9, were detected by their fluorescence at 330 nm with an excitation at 290 nm. Plastoquinone-9 in the oxidized state was measured by its absorbance at 255 nm. Typical chromatograms are shown in Supplemental Fig. S1. Plastochromanol-8 and plastoquinone-9 standards were a kind gift from Dr. J. Kruk. Tocopherol standards were purchased from Sigma. Unless specified otherwise, the PQ-9 data given in this article correspond to the total content (oxidized PQ-9 + reduced PQH₂-9).

Determination of the photoactive and non-active fractions of plastoquinone-9 was done following the protocol described in Kruk & Karpinski (2006) modified by Ksas et al. (2015). Leaf samples were frozen in liquid nitrogen as quickly as possible to avoid changes in the plastoquinone redox state during sample preparation. Two discs (diameter, 0.8 cm) were punched out from each leaf. To determine the pool of photoactive plastoquinone-9, we first measured the amount of reduced plastoquinone-9 (plastoquinol-9) in one disc after a dark-adaptation period of 2 h. The dark-adapted samples were frozen in liquid nitrogen after 3-s illumination with far-red light to ensure full oxidation of the photoactive pool of plastoquinone. Then, plastoquinol-9 was re-quantified in the disc that had been exposed to a high PFD of white light (2000 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$) for 15 s. Illumination was done in a mortar that was filled with liquid nitrogen after the 15-s illumination in order to freeze the samples in the light. The size of the photoactive pool was calculated as the difference between the plastoquinol-9 content in the light and the plastoquinol-9 content in the dark. The size of the non-photoactive pool of plastoquinone-9 was the sum of the amount of plastoquinol-9 in the dark (not reoxidable in the dark) and the amount of oxidized plastoquinone-9 in the light (not reducible by high light). This pool

corresponds mainly to the plastoquinone molecules present in the plastoglobules and the chloroplast envelope (Lichtenthaler, 2007) which are not involved in photosynthetic electron transport.

2.3 PQ-C determination by UPLC-MS/MS

Hydroxyplastoquinone (PQ-C) was analyzed using an UPLC-MS/MS system (UPLC ultimate RS 3000; Thermo Fisher with QTOF 5600; AB Sciex) connected to a Kinetex C18 2.1 × 150 mm column (Phenomenex). The APCI source was operated in negative mode. A binary solvent system was used, in which mobile phase A consisted of acetonitrile:water (60:40, v/v) and 10 mM ammonium formate and mobile phase B consisted of isopropanol:acetonitrile (90:10, v/v), 10 mM ammonium formate, and 0.1% formic acid (v/v). The gradient started with 27% of solution B and was increased to 97% of solution B within 20 min at a speed of 0.3 ml min⁻¹, and then maintained at 97% for 5 min. Solution B was then decreased to a 27% enrichment during 7 min for column re-equilibration. Column temperature was set at 45°C. Oxidized and reduced PQ-9 and PQ-C identification was achieved based on the comparison of retention times and accurate masses with standards. Reduced PQ-9 and PQ-C were obtained by reducing the oxidized PQ-9 and PQ-C standards with sodium borohydride. In samples, PQ-C and PQ-9 contents were measured in selected reaction monitoring (SRM) mode using the following transition: m/z 764.6 → m/z 202.095 ± 0.025 (CE = -70 eV) and m/z 748.6 → m/z 149.058 ± 0.025 (CE = -70 eV) respectively. Examples of extracted-ion chromatograms (XIC) for PQ-C are shown in Supplementary Fig. S2.

2.4 Chloroplast fractionation

Chloroplasts, isolated from leaves of 6-week-old plants, were hypotonically ruptured and subplastidial compartments were isolated by centrifugation using sucrose-density gradient following the protocol described in (Besagni, Eugeni Piller, & Bréhélin, 2011), except that plants were not pre-adapted to darkness. 32 fractions of 1 ml were collected from the centrifuged tubes (from top to bottom), and 400 µl of each fraction were diluted to obtain a final sucrose concentration of 5%. The diluted fractions were then grinded in mortar glass tube with 5 ml of cold ethyl acetate and sonicated (Digital Sonifer, Brandon, USA) for 1 min with a 50% 'duty cycle' at 40% of power in an ice bath. The extracts were centrifuged for 13 min at 9600 g at 4°C. 500 µl of supernatant was evaporated in a stream of nitrogen and dissolved in 300 µl of methanol/hexane (17/1, v/v). 100 µl were immediately analyzed by HPLC, as described above.

2.5 Chlorophyll fluorescence

The maximal quantum yield of PSII photochemistry was measured on attached leaves with a PAM-2000 fluorometer (Walz) as $F_v/F_m = (F_m - F_o)/F_m$ where F_m is the maximal fluorescence level obtained

with a 800-ms pulse of intense white light and F_o is the initial level obtained after a 2-s pulse of far-red light. Measurements were done in the dark in leaves dark-adapted for 30 min.

Chlorophyll fluorescence induction curves were measured with a JTS-10 spectrophotometer/fluorimeter (BloLogic, France), and the number of PSII 2-electron acceptors was estimated as described in Joliot & Joliot (2002) as the area over the fluorescence rise measured in dark-adapted leaves divided by the area over the curve for leaves infiltrated with DCMU (one electron transferred to Q_A in the light). Chlorophyll fluorescence imaging was done with a laboratory-built instrument described in Johnson et al. (2009).

2.6 Lipid peroxidation

Lipids were extracted from approximately 0.5 g of leaves frozen in liquid nitrogen. The leaves were grinded in an equivolume methanol/chloroform solution containing 5mM Triphenyl Phosphine, 1 mM 2,6-tert-butyl-p-cresol (BHT) (5 ml g⁻¹ fresh weight) and citric acid (2.5 ml g⁻¹ fresh weight), using an Ultraturax blender. Internal standard 15-HEDE was added to a final concentration 100 nmol g⁻¹ fresh weight, and mixed properly. After centrifugation at 700 rpm and 4°C for 5 min, the lower organic phase was carefully taken out with the help of a glass syringe into a 15 ml glass tube. The syringe was rinsed with approximately 2.5 ml chloroform and transferred back into the tube. The process was repeated and the lower layer was again collected and pooled to the first collection. The solvent was evaporated under N₂ gas at 40 °C. The residues were recovered in 1.25 ml absolute ethanol and 1.25 ml of 3.5 N NaOH and hydrolyzed at 80°C for 30 min. The ethanol was evaporated under N₂-gas at 40 °C for ~10 min. After cooling to room temperature, pH was adjusted to 4-5 with 2.1 ml citric acid. Hydroxy fatty acids were extracted with hexane/ether 50/50 (v/v). The organic phase was analyzed by straight phase HPLC-UV, as previously described (Montillet et al., 2004). The hydroxyoctadecatrienoic acid (HOTE) isomers (9-, 12-, 13- and 16-HOTE derived from the oxidation of the main fatty acid in Arabidopsis leaves, linolenic acid) were quantified based on the 15-HEDE internal standard.

Lipid peroxidation was also visualized in whole plants by autoluminescence imaging. Stressed plants were dark adapted for 2 h, and the luminescence emitted from the spontaneous decomposition of lipid peroxides was captured by a highly sensitive liquid N₂-cooled charge-coupled device (CCD) camera, as previously described (Birtic et al., 2011). The images were treated using Image J software (NIH, USA)

2.7 SOSG-EP fluorescence

Production of ¹O₂ was measured in attached leaves from the fluorescence of the ¹O₂-specific SOSG (Singlet Oxygen Sensor Green) fluorescent probe (Invitrogen), as described previously (Ramel et al.,

2012). With the help of a 1-ml syringe (without the needle), 100 μM SOSG was pressure-infiltrated into the leaves through the lower surface. Plants with SOSG-infiltrated leaves were exposed to a PFD of 1500 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ at 7 °C for 20 min. As a control treatment, plants with SOSG-infiltrated leaves were placed in the dark at room temperature for 20 min. SOSG-EP fluorescence was then measured from leaf discs punched from the SOSG-infiltrated leaves using a fiberoptics-equipped Perkin-Elmer spectrofluorometer (LS 50B) with a 475 nm excitation light. SOSG-EP fluorescence at 524 nm (F524) was normalized to chlorophyll fluorescence at 680 nm (F680) for each leaf disc. SOSG-EP fluorescence was calculated as: $([\text{F524 in the light} - \text{F524 in the dark}]/\text{F680}) \times 10$.

2.8 EPR spectroscopy

For the preparation of thylakoid membranes, 7 g of leaves (fresh weight) were grinded for 2 s in 50 ml of extraction buffer (330 mM sorbitol, 50 mM Tricine, 2 mM EDTA(Na₂), 1 mM MgCl₂, 2 mM Ascorbate, pH 7.7) with 5 mM dithiothreitol (DTT) in a Warring blender at low speed. The liquid phase was removed and set aside, and 50 ml of extraction buffer was added for a second extraction. The extracts were filtered onto 4 Miracloth layers, and the filtrate was centrifuged for 4 min at 1500 g at 4°C. The pellet was washed twice with the extraction buffer and centrifuged for 4 min at 1500 g at 4°C. The washed pellet was resuspended in 21 ml of lysis buffer pH 7.8 (10 mM Tricine, 10 mM NaCl, 10 mM MgCl₂) with 1 mM PMSF (phenyl methylsulfonyl) with occasional stirring for 15 min. The sample was centrifuged at 48400 g for 15 min. The pellet was resuspended in 1.75 ml of storage buffer (100 mM Tricine, 10 mM NaCl, 10 mM MgCl₂, 400 mM sucrose, pH 7.8) and stored at -80°C before analyses.

Singlet oxygen formation was monitored by electron paramagnetic resonance (EPR) spectroscopy using the spin probe 2,2,6,6-tetramethyl-4-piperidone (TEMPD) purified by vacuum distillation. The oxidation of TEMPD by ¹O₂ forms 2,2,6,6-tetramethyl-4-piperidone-1-oxyl (TEMPONE) detectable by EPR spectroscopy. Thylakoid membranes (25 $\mu\text{g Chl ml}^{-1}$) were illuminated in the presence of 50 mM TEMPD and 40 mM MES-NaOH buffer (pH = 6.5) for 30 min under 1000 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$. After illumination, thylakoid membranes were centrifuged at 1 000 x g for 2 min to separate sample from spin probe. The EPR spectra were recorded using an EPR spectrometer Mini Scope MS400 (Magnettech GmbH, Berlin, Germany). EPR measurement conditions were as follows: microwave power, 10 mW; modulation amplitude, 1 G; modulation frequency, 100 G; scan rate, 1.62 G s⁻¹.

3 RESULTS

3.1 Overexpression of the *SPS1* gene in the *vte1* mutant leads to a marked accumulation of PQ-9, mainly in the plastoglobules and the chloroplastic envelopes

Ksas et al. (2015) previously showed that overexpressing the *SPS1* gene in *Arabidopsis* brings about a noticeable increase in the leaf content in both PQ-9 and PC-8. This was confirmed here with one of the *SPS1oex* lines previously described, with a 50%-increase in the total PQ-9 content and a 3-fold increase in PC-8 compared to WT (Fig. 1A). Please note that throughout the text the term PQ-9 is used as a generic term for total PQ-9, i.e. oxidized PQ-9 + reduced PQH₂-9 (plastoquinol). In all genotypes, PQ-9 was highly reduced, with PQH₂-9 (plastoquinol) representing around 75% of the total pool (Supplemental Fig. S3). The tocopherol cyclase mutant *vte1* is devoid of both tocopherols and PC-8 (Zbierzak et al., 2009). The loss of those two lipid-soluble antioxidants in *vte1* leaves was accompanied by a significant decrease in the total PQ-9 levels, as previously reported (Ksas et al, 2015). When the *vte1* mutant was crossed with the *SPS1*-overexpressing line (line #12 in Ksas et al., 2015) to generate a double mutant *vte1 SPS1oex*, the total PQ-9 content of the leaves was strongly increased, reaching values above the WT levels. Compared to *vte1*, *vte1 SPS1oex* contained approximately 3 times more PQ-9. The changes in tocopherols observed in *vte1*, *SPS1oex* and *vte1 SPS1oex* had no significant effect on the growth phenotype of the plants in low light (Fig. 1B) and did not affect significantly the efficiency of photosynthetic electron transport in young and mature leaves as measured by the chlorophyll fluorescence parameter Φ_{PSII} (Fig. 1C). Compared to the *SPS1* overexpressors, a decrease in Φ_{PSII} was nevertheless observed in the older leaves of *vte1* and, to a lesser extent, of WT. Thus, high PQ-9 levels appear to be helpful to maintain the efficiency of PSII-mediated electron flow in old, pre-senescing leaves.

To determine the localization of the extra PQ-9 that accumulated in *SPS1*-overexpressing plants, chloroplast membranes were fractionated by ultracentrifugation on sucrose gradient (Besagni et al., 2011). As expected (Besagni et al., 2011; Vidi et al., 2006), 3 pigmented fractions were obtained (Fig. 2A): plastoglobules (PG) were found floating at the top of the tubes and forming a whitish layer, while an intermediate yellowish zone corresponding to the outer and inner chloroplast envelope membranes (ENV) separated from the green zone at the bottom containing chlorophyll-binding thylakoid membranes. 32 fractions were collected from top (fraction 1) to bottom (fraction 32) and analyzed for their PQ-9 content (Fig. 2B, Supplemental Fig. S4). PQ-9 was found in all fractions, with an almost equivalent partitioning between the thylakoids and the two other fractions (envelopes+plastoglobules) in WT leaves (Supplemental Fig. S4). We checked the abundance of the plastoglobule marker protein Plastoglobulin 35 (PGL35) in the different fractions by Western blotting. Compared to the plastoglobule fraction, the abundance of PGL35 in the envelope fraction was very low (Supplemental Fig. S5). Consequently, the almost similar amounts of PQ-9 found in the plastoglobules and the envelopes (Supplemental Fig. S4) are not attributable to a contamination of the envelopes by plastoglobules. Surprisingly, fractions 24-28 contain very little PQ-9 although they are green. It is possible that those fractions contain photosynthetic pigmented complexes released from

the thylakoids during preparation. In *vte1*, the loss of PQ-9 concerned exclusively the PG and envelope fractions (fractions 1-4 and 5-14, respectively), with the levels in the thylakoids (fractions 26-32) being virtually unchanged relative to WT. Similarly, the accumulation of total PQ-9 in *SPS1oex* and *vte1 SPS1oex* leaves impacted mainly the PG and envelope fractions. Thus, thylakoid membranes appear to accommodate a relatively constant amount of PQ-9 (presumably ensuring optimum electron transport) while the PG and envelope fractions function as PQ-9 storage sites that can absorb large variations in total PQ-9 in the chloroplasts. This was confirmed by the estimation of the photochemically active and non-active pools of PQ-9 using a previously described procedure based on HPLC analysis of reduced and oxidized PQ-9 in the dark and upon intense illumination (Ksas et al., 2015). For each genotype, we found that the pool sizes of the active plus non-active pools (Fig. 2C) were matching, within experimental accuracy, the total pool size (Fig. 1A; in nmol cm⁻², 0.91±0.07 in WT, 1.43 ± 0.15 in *SPS1oex*, 0.60±0.04 in *vte1* and 1.73±0.08 in *vte1 SPS1oex*). Figure 2C shows that the photoactive pool (reflecting the thylakoid PQ-9 pool involved in photosynthetic electron transport) showed relatively little variations in the different genotypes while the non-photoactive pool (reflecting the pool, located outside the thylakoid membranes, not involved in electron transport) exhibited strong variations, with *SPS1oex* and *vte1 SPS1oex* having by far the largest pools of extra-thylakoid PQ-9 (Fig. 2C), in agreement with the analyses of Fig. 2B and Supplemental Fig. S4.

It is also possible to estimate the size of the functional pool of PSII electron acceptors by *in vivo* chlorophyll fluorescence measurements (Malkin & Kok, 1966; Forbush & Kok, 1968). The chlorophyll fluorescence induction curves of Fig. 2D showed little differences between the different genotypes, confirming that the functional pool in the thylakoids was not much affected by the total PQ-9 levels. From the fluorescence induction curves of Fig. 2D and the corresponding curves obtained in the presence of DCMU, we estimated that the number of PSII two-electron acceptors is in the range 13-16 molecules (insert of Fig. 2D), which is consistent with previous estimations in various photosynthetic organisms (Forbush & Kok, 1968; Kolber & Falkowski, 1993).

3.2 Accumulation of total PQ-9 in the *vte1 SPS1oex* mutant compensated for the lack of tocopherols and PC-8 and increased tolerance to photooxidative stress

Plants grown in low light (150 μmol photons m⁻² s⁻¹) were transferred to photooxidative stress conditions (1500 μmol photons m⁻² s⁻¹, 7°C). After 2 d, this treatment led to leaf bleaching (Fig. 3A), low Fv/Fm ratio indicative of PSII photoinhibition (Fig. 3D) and enhanced lipid peroxidation (Fig. 3B and C) in WT leaves. These symptoms of photodamage were markedly attenuated in *SPS1oex* plants, as previously shown (Ksas et al., 2015). In striking contrast, *vte1* exhibited a very high sensitivity to photooxidative stress, showing dramatic effects on leaves (severe bleaching and loss of turgescence), intense lipid peroxidation and drastic inhibition of PSII activity relative to WT. This confirms the

photoprotective role of tocopherol and/or PC-8 which are absent in *vte1* (Havaux, Eymery, Porfirova, Rey & Dörmann, 2005). Interestingly, overexpression of *SPS1* suppressed the photobleaching phenotype of *vte1*. Furthermore, *vte1 SPS1oex* was more tolerant to lipid peroxidation than *vte1*. The data of Fig. 3 thus demonstrate that total PQ-9 can compensate for the lack of tocopherol and PC-8 in *vte1* leaves, leading to a markedly increased tolerance to photooxidative stress. This effect was confirmed in another *SPS1*-overexpressing *vte1* mutant and under longer stress exposure (4 d) (Supplemental Figure S6).

3.3 Plastoquinone as a singlet oxygen scavenger in planta

$^1\text{O}_2$ production by illuminated leaves was monitored by measuring the intensity of SOSG-endoperoxide (EP) fluorescence at 525 nm (Flors et al., 2006; Kim, Fujitsuka, & Majima, 2013). As shown in Fig. 4A, the photosensitivity of *vte1* leaves correlated with an increased production of $^1\text{O}_2$ compared to WT, in line with the known function of tocopherols as $^1\text{O}_2$ quenchers (Fahrenholtz, Doleiden, Trozzolo, & Lamola, 1974; Di Mascio, Devasagayam, Kaiser, & Sies, 1990; Krieger-Liszkay, Fufezan, & Trebst, 2008; Choe, 2107). It was decreased in *vte1 SPS1oex*, showing that replacement of tocopherols by total PQ-9 restores the $^1\text{O}_2$ quenching capacities. This finding was confirmed by the analysis of $^1\text{O}_2$ production by illuminated thylakoid membranes using EPR spectroscopy and the $^1\text{O}_2$ -specific spin probe TEMPONE (Fig. 4C-D). The production rate of $^1\text{O}_2$ was the highest in *vte1* thylakoid membranes, whereas accumulation of PQ-9 by *SPS1* overexpression in both WT and *vte1* backgrounds resulted in the lowest $^1\text{O}_2$ production levels in the light, emphasizing the role of total PQ-9 in the quenching of $^1\text{O}_2$ *in vivo*.

Oxidation of PQ-9 by $^1\text{O}_2$ has been shown to produce some specific derivatives, such as the hydroxy derivative PQ-C (Szymanska et al., 2014). The latter compound was shown to be a stable product of $^1\text{O}_2$ action in plants, particularly when $^1\text{O}_2$ production is elevated during short-term exposure to high light stress. Quantification of PQ-C in control Arabidopsis leaves showed the presence of this compound in all genotypes (Fig. 4B). Similarly to PQ-9, PQ-C was predominantly in the reduced state (PQH₂-OH) (Supplemental Fig. S3). PQ-C accumulation is consistent with the attribution of a $^1\text{O}_2$ scavenging function to PQ-9: the amount of this PQ-9 hydroxy derivative, formed by $^1\text{O}_2$, is the highest in *vte1 SPS1oex* where total PQ-9 is the most abundant and is the major lipid-soluble antioxidant in the absence of tocopherol and PC-8.

3.4 High light stress preferentially consumes the thylakoidal PQ-9 pools

Arabidopsis plants were exposed for 2.5 h to high PFD (1500 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$) and low temperature (7°C). The total PQ-9 concentration was decreased by around 30% in WT leaves after this treatment (Fig. 5A). Fig. 5B shows that the PQ-9 losses were particularly marked for the thylakoid-located photochemically active PQ-9 pool (-60%) while the extra-thylakoid, photochemically non-

active pool was not significantly affected. Since the photosystems in the thylakoids are the main source of ROS in the light, a preferential oxidation of this photochemically active pool is expected. This is actually confirmed by the analysis of $^1\text{O}_2$ -specific PQ-C compound in the different membrane fractions of the chloroplasts: PQ-C was predominantly found in the thylakoids, with the concentration in the plastoglobules and the envelopes being very low (Fig. 5C). In *vte1*, the photoinduced decrease in total PQ-9 content was very marked (>50%) (Fig. 5A), and the photochemically active pool was almost completely depleted after the high light treatment (Fig. 5B). However, contrary to WT, the size of the extra-thylakoid, photochemically non-active PQ-9 pool was also decreased in the mutant, although to a lesser extent (~40%) than the active pool (Fig. 5B). Fig. 5C shows that, similarly to WT, PQ-C in *vte1* chloroplasts was present mainly in the thylakoids, confirming that PQ-9 oxidation occurred at this level. As shown at the leaf level (Fig. 3), the PQ-C concentration was noticeably higher in *vte1* thylakoids compared to WT thylakoids, indicating enhanced oxidation of PQ-9 in this mutant. In *vte1 SPS1oex* leaves which rely on high levels of PQ-9 for their phototolerance, a considerable loss of total PQ-9 was observed during high light stress (Fig. 5A, 5B), with a very strong accumulation of PQ-C in the thylakoids (Fig. 5C). Thus, our data suggest that, under conditions of massive oxidative degradation of PQ-9 in the thylakoids of *vte1* and *vte1 SPS1oex* chloroplasts, there is a transfer of PQ-9 molecules from the plastoglobules and/or the envelope to the thylakoid membranes in order to compensate for PQ-9 photooxidation, hence lowering the PQ-9 levels in the storage sites outside the thylakoid membranes. Owing to their high accumulation levels of total PQ-9, *vte1 SPS1oex* leaves were able to maintain the total PQ-9 content close to the WT levels after the high light treatment (Fig. 5A), and this was associated with a high resistance to photooxidative stress (Fig. 3).

4 DISCUSSION

The essential role of PQ-9 in photosynthesis as an electron carrier has long been recognized (Amesz, 1973; Crane, 2010). However, it is clear from this study that PQ-9 also fulfills an important photoprotective function that is not directly related to photosynthetic electron transport. Actually, only a fraction of the total PQ-9 pool is located in the thylakoid membranes where it can participate in linear and cyclic electron fluxes. We estimated that, in WT *Arabidopsis* leaves under normal growth conditions, the photochemically active PQ-9 corresponds to around 15 molecules per PSII and 50% of the total PQ-9 content. The latter value is higher than previous reports (ca. 30%, Kruk & Karpinski, 2006; Szymanska & Kruk 2010), possibly due to different growth conditions. The size of the total thylakoid-associated PQ-9 pool was found to be stable in control conditions, showing relatively little variation among the different genotypes studied here or in response to changes in the total PQ-9 content of the leaves. This stability contrasts with the non-photochemically active pool, the size of which exhibited large variations. As confirmed in this study, the latter PQ-9 pool partitions between

the plastoglobules (Zbierzak et al., 2009) and the envelope membranes (Soll et al., 1985). The size of the extra-thylakoid PQ-9 pool drastically decreased in the photosensitive *vte1* mutant, whereas it was selectively increased when PQ-9 biosynthesis was stimulated. The highest PQ-9 levels were found in *vte1 SPS1oex* leaves in which approximately two-thirds of the total PQ-9 pool was not in the thylakoids. In contrast, only 40% of the total pool was outside the thylakoid membrane in the *vte1* single mutant that contains the lowest levels of total PQ-9. Somehow, the PQ-9 reservoir stored in the plastoglobules and the chloroplast envelopes functions as a buffer that can manage large variations in the total PQ-9 levels in the leaves.

The multi-location storage of PQ-9 suggests a multi-functionality for this compound. Previously, PQ-9 was reported to be involved in biosynthesis pathways, as a component of phytoene desaturase in the carotenoid biosynthesis pathway (Norris et al., 1995) and of NDC1, a type-II NAD(P)H quinone oxidoreductase involved in vitamin K1 synthesis (Eugeni Piller et al., 2011). This work shows that it is also a ROS scavenger protecting the chloroplasts against photodamage by a mechanism that could possibly be similar to the antioxidant activity of tocopherol. Indeed, overexpression of *SPS1* in the tocopherol-deficient *vte1* mutant concomitantly enhanced leaf PQ-9 levels and decreased plant photosensitivity, indicating that PQ-9 can replace tocopherols in photoprotection. Tocopherol has been shown to act as a $^1\text{O}_2$ quencher (Fahrenholtz et al., 1974; Di Mascio et al., 1990; Krieger-Liszkay et al., 2008; Choe, 2017) and a terminator of lipid peroxidation (Burton & Ingold, 1986; Ham & Liebler, 1995). PQ-9 can also do both. Indeed, Nowicka et al. (2013) showed that reduced or oxidized PQ-9 incorporated into liposomes precludes lipid oxidation, and the present work demonstrates that high total PQ-9 levels lowered $^1\text{O}_2$ accumulation levels. Moreover, high light stress leads to a loss of PQ-9 and a concomitant formation of PQ-C indicating $^1\text{O}_2$ oxidation of PQ-9. As expected, the accumulation of PQ-C was the highest in *vte1 SPS1oex* leaves which rely the most on total PQ-9 for their resistance to photooxidative stress. PQ-C was found almost exclusively in the thylakoids where ROS are produced in the light, either by excitation energy transfer ($^1\text{O}_2$) or by electron transfers (superoxide anion radical, hydrogen peroxide or hydroxyl radical) (Asada, 2006; Krieger-Liszkay et al., 2008; Li, Wakao, Fischer & Niyogi, 2009; Pospisil, 2014).

PQ-9, especially when it is in the reduced state, possesses a number of features that are likely to make this compound an excellent antioxidant. Firstly, it is present in high amounts as a diffusible molecule in thylakoids. As shown in Fig. 1, the concentration of total PQ-9 in *Arabidopsis* leaves is higher than tocopherol + plastochromanol levels. Leaf enrichment in total PQ-9 relative to tocopherols can be even amplified during long-term acclimation of plants to high light intensities (Ksas et al., 2015; Lichtenthaler, 2007). Second, the PQ-9 side chain bears several double bonds contrary to tocopherols. Double bonds are known to be preferential targets of $^1\text{O}_2$ for oxidation (Triantaphylidès & Havaux, 2009), and therefore the unsaturated side-chain of PQ-9 could provide a possibility for PQ-9 to quench

$^1\text{O}_2$ by a chemical mechanism. Nevertheless, the antioxidant activity is mainly beard by the phenolic group of reduced PQ-9 (Kim & Min, 2008), so that reduced PQ-9 is a better $^1\text{O}_2$ quencher than oxidized PQ-9. Third, in thylakoid membranes, photosynthetic electron transport relies on the interaction between PQ-9 and the PSII reaction centers which are the main $^1\text{O}_2$ generators during photosynthesis (Krieger-Liszkay et al., 2008). It was initially hypothesized that, in the thylakoid membranes, PQ-9 is located in the fluid bilayer-midplane region allowing rapid lateral movements between cytochrome b6/f and PSII complexes (Millner & Barber, 1984). However, more recent data favor a close compartmentization of PSII, PQ-9 and cytochrome b6/f complex in membrane microdomains (Lavergne & Joliot, 1991; Johnson, Vasilev, Olsen, & Hunter, 2014). Moreover, PQ-9 diffusion towards/from PSII was recently proposed to occur via several entries/exits in the PSII reaction center with an exchange cavity where PQ-9 can diffuse around (Van Eerden, Melo, Frederix, Periole, & Marrink, 2017). These characteristics of the interactions between PSII and PQ-9 are likely to augment the opportunities for PQ-9 to scavenge $^1\text{O}_2$ produced in the reaction centers. Consequently, PQ-9 could constitute a first line of defense against photosynthesis-produced ROS, supplementing the action of the β -carotene molecules bound to the PSII centers.

Chemical quenching involves oxidation of the quencher and therefore total PQ-9 is expected to be consumed during its antioxidant activity, as indeed observed in plants exposed to photooxidative stress conditions (Ksas et al., 2015; Fig. 5). In the *vte1* mutant, the total PQ-9 levels are constitutively lower than in WT, and this can be attributed to the chronic oxidation of PQ-9 in the absence of tocopherol. The increased levels of PQ-C in the *vte1* mutants compared to WT corroborate this interpretation. This phenomenon concerns mainly the extra-thylakoid PQ-9 pool, with the total thylakoid-located PQ-9 levels being virtually unchanged relative to WT levels under control conditions. Since ROS are produced in the light in the thylakoids, a dynamic exchange of PQ-9 molecules must exist between the thylakoids and their storage sites in the plastoglobules and the envelopes to replace photooxidized PQ-9 molecules in the thylakoid membranes with new PQ-9 molecules and to maintain a constant pool in the thylakoid membranes. Accordingly, plastoglobules have been shown to be physically coupled to thylakoids in a way allowing bidirectional channeling of lipid metabolites (Austin, Frost, Vidi, Kessler, & Staehelin, 2006, Bréhélin & Kessler, 2008). However, experimental data measuring the nature and the rate of this metabolite exchange are not available. When Arabidopsis plants were suddenly exposed to high light stress, total PQ-9 levels drastically decreased in the thylakoids, as expected when excess light energy leads to high $^1\text{O}_2$ production levels by the photosystems. Moreover, PQ-C was detected predominantly in the thylakoid fractions, indicating that $^1\text{O}_2$ oxidation of PQ-9 occurs at this level. This is in agreement with a previous study by Szymanska and Kruk (2010) who showed a selective decrease in the photochemically active PQ-9 in Arabidopsis leaves exposed to high light stress (Szymanska & Kruk, 2010). However, we also observed a significant

reduction of the total PQ-9 concentration, though less pronounced, in the photochemically non-active pool when the photostress was severe and PQ-9 degradation was very pronounced such as in the *vte1* background. This effect of high light stress is consistent with a transfer of PQ-9 molecules from the plastoglobules and/or envelopes to the thylakoids to compensate the oxidative modification of PQ-9 during its scavenging activity. In this context, it is worth mentioning that reduced PQ-9 showed a high mobility in lipid membranes compared to α -tocopherol (Jemiola-Rzeminska, Kruk, & Strzalka, 2003). Nevertheless, the rate of this transfer appeared to be slow compared to the rate of PQ-9 oxidation in high light since it could not maintain the pool size of photochemically active PQs to the control levels measured in low light.

Because PQ-9 somehow functions as a sacrificial antioxidant consumed during oxidative stress, it must be re-synthesized for the pool being re-filled. Accordingly, most genes of the PQ-9 biosynthesis pathway are strongly upregulated during acclimation of Arabidopsis to high light intensities (Ksas et al., 2015; Block et al., 2013). This induction of PQ-9 biosynthesis genes was accompanied by a concomitant rise in the size of the total PQ-9 pool while there was less effect on the tocopherol content (Ksas et al., 2015; Szymanska & Kruk, 2010). The selective accumulation of PQ-9 molecules during growth in high light environments was reported in other species such as beech or fig trees (Lichtenthaler, 2007), consistently with the photoprotective role described for this compound in the present work.

ACKNOWLEDGMENTS

We would like to thank the Phytotec platform (Commissariat à l'Energie Atomique et aux Energies Alternatives, Cadarache) for growing plants under normal and stress conditions. We also thank Jerzy Kruk for providing us with PQ-9 and PC-8 standards. This work was supported by the French National Research Agency (ANR project SLOSAM, 14-CE02-0010-02). The PP laboratory was supported by the Ministry of Education, Youth and Sports of the Czech Republic grant LO1204 (Sustainable development of research in the Centre of the Region Haná from the National Program of Sustainability I). The UPLC-MS/MS of the HelioBiotec platform was funded by the European Union, the Région Provence-Alpes-Côte d'Azur, the French Ministry of Research and CEA.

CONFLICT OF INTEREST

The authors claim no conflict of interest

REFERENCES

Amesz, J. (1973) The function of plastoquinone in photosynthetic electron transport. *Biochimica et Biophysica Acta*, 301, 35-51.

480 Asada, K. (2006) Production and scavenging of reactive oxygen species in chloroplasts and their
 481 functions. *Plant Physiology*, 141, 391-396.

482 Austin, J.R. 2nd, Frost, E., Vidi, P.A., Kessler, F., & Staehelin, L.A. (2006) Plastoglobules are lipoprotein
 483 subcompartments of the chloroplast that are permanently coupled to thylakoid membranes and
 484 contain biosynthetic enzymes. *Plant Cell*, 18, 1693-1703.

485 Besagni, C., Eugeni Piller, L., & Bréhélin, C. (2011) Preparation of plastoglobules from Arabidopsis
 486 plastids for proteomic analysis and other studies. *Methods in Molecular Biology*, 775, 223-239.

487 Birtic, S., Ksas, B., Genty, B., Mueller, M.J., Triantaphylides, C., & Havaux, M. (2011) Using spontaneous
 488 photon emission to image lipid oxidation patterns in plant tissues. *Plant Journal*, 67, 1103-1115.

489 Block, A., Fristedt, R., Rogers, S., Kumar, J., Barnes, B., Barnes, J., Elowsky, C.G., Wamboldt, Y.,
 490 Mackenzie, S.A., Redding, K., Merchant, S.S., & Basset, G.J. (2013) Functional modeling identifies
 491 paralogous solanesyl-diphosphate synthases that assemble the side chain of plastoquinone-9 in
 492 plastids. *Journal of Biological Chemistry*, 200, 27594-27606

493 Bréhélin, C., & Kessler, F. (2008) The plastoglobule: a bag full of lipid biochemistry tricks.
 494 *Photochemistry and Photobiology*, 84, 1388-1394.

495 Burton, G.W., & Ingold, K.U. (1986) Vitamin E: Application of the Principles of Physical Organic
 496 Chemistry to the Exploration of Its Structure and Function. *Accounts of Chemical Research*, 19, 194-
 497 201.

498 Choe, E. (2017) Effects and mechanism of minor compounds in oil on lipid oxidation. In *Food Lipids:*
 499 *Chemistry, Nutrition, and Biotechnology* (ed C.A. Koh), pp. 567-590. CRC Press, Boca Raton.

500 Cook, W.B., & Miles, D. (1992) Nuclear mutations affecting plastoquinone accumulation in maize.
 501 *Photosynthesis Research*, 31, 99-111.

502 Crane, F.L. (2010) Discovery of plastoquinone: a personal perspective. *Photosynthesis Research*, 103,
 503 195-209.

504 Di Mascio, P., Devasagayam, T.P., Kaiser, S., & Sies, H. (1990) Carotenoids, tocopherols and thiols as
 505 biological singlet molecular oxygen quenchers. *Biochemical Society Transactions*, 18, 1054-1056.

506 Eugeni Piller, L., Besagni, C., Ksas, B., Rumeau, D., Bréhélin, C., Glauser, G., Kessler, F., & Havaux, M.
 507 (2011) Chloroplast lipid droplet type II NAD(P)H quinone oxidoreductase is essential for
 508 prenylquinone metabolism and vitamin K1 accumulation. *Proceedings National Academy of*
 509 *Sciences USA*, 108;14354-14359.

510 Fahrenholtz, S.R., Doleiden, F.H., Trozzolo, A.M., & Lamola, A.A. (1974) On the quenching of singlet
 511 oxygen by alpha-tocopherol. *Photochemistry and Photobiology*, 20, 505-509.

512 Falk, J., & Munné-Bosch, S. (2010) Tocochromanol functions in plants: antioxidant and beyond. *Journal*
 513 *of Experimental Botany*, 61, 1549-1566.

514 Flors, C., Fryer, M.J., Waring, J., Reeder, B., Bechtold, U., Mullineaux, ..., Baker, N.R. (2006) Imaging the
 515 production of singlet oxygen in vivo using a new fluorescent sensor, Singlet Oxygen Sensor Green.
 516 *Journal of Experimental Botany*, 57, 1725-1734.

517 Forbush, B., & Kok, K. (1968) Reaction between primary and secondary electron acceptors of
 518 photosystem II of photosynthesis. *Biochimica et Biophysica Acta*, 162, 243-253.

519 Gruszka, J., Pawlak, A., & Kruk, J. (2008) Tocochromanols, plastoquinol, and other biological
 520 prenyllipids as singlet oxygen quenchers – determination of singlet oxygen rate constants and
 521 oxidation products. *Free Radicals in Biology and Medicine*, 45, 920-928.

522 Ham, A.-J. L., & Liebler, D.C. (1995) Vitamin E Oxidation in Rat Liver Mitochondria. *Biochemistry*, 34,
 523 5754-5761.

524 Havaux, M., Eymery, F., Porfirova, S., Rey, P., & Dörmann, P. (2005) Vitamin E protects against
 525 photoinhibition and photooxidative stress in *Arabidopsis thaliana*. *Plant Cell*, 17, 3451-3469.

526 Hernandez, V.A., Eriksson, E.K., & Edwards, K. (2015) Ubiquinone-10 alters mechanical properties and
 527 increases stability of phospholipid membranes. *Biochimica et Biophysica Acta*, 1848, 2233-2243.

528 Hundal, T., Forsmark-Andrée, P., Ernster, L., & Andersson, B. (1995) Antioxidant activity of reduced
 529 plastoquinone in chloroplast thylakoid membranes. *Archives Biochemistry Biophysics*, 324, 117-
 530 122.

531 Jemiola-Rzeminska, M., Kruk, J., and Strzalka, K. (2003) Anisotropy measurements of intrinsic
 532 fluorescence of prenyllipids reveal much higher mobility of plastoquinol than alpha-tocopherol in
 533 model membranes. *Chemistry and Physics of Lipids*, 123, 233-243.

534 Johnson, M.P., Vasilev, C., Olsen, J.D., & Hunter, C.N. (2014) Nanodomains of cytochrome b6f and
 535 photosystem II complexes in spinach grana thylakoid membranes. *Plant Cell*, 26, 3051-3061.

536 Johnson, X., Vandystadt, G., Bujaldon, S., Wollman, F.-A., Dubois, R., Roussel, P., ..., Béal, D. (2009) A
 537 new setup for in vivo fluorescence imaging of photosynthetic activity. *Photosynthesis Research*,
 538 102, 85-93.

539 Joliot, P., & Joliot, A. (2002) Cyclic electron transfer in plant leaf. *Proceedings National Academy of*
 540 *Sciences USA*, 99, 10209-10214.

541 Kim, H.J., & Min, D.B. (2008) Tocopherol stability and prooxidant mechanisms of oxidized tocopherols
 542 in lipids. In *Food Lipids: Chemistry, Nutrition, and Biotechnology* (eds C.A. Koh & Min D.B.), pp. 435-
 543 447. CRC Press, Boca Raton.

544 Kim, S., Fujitsuka, M., & Majima, T. (2013) Photochemistry of singlet oxygen sensor green. *Journal*
 545 *Physical Chemistry B*, 117, 13986-13992.

546 Kolber, Z., & Falkowski, P.G. (1993) Use of active fluorescence to estimate phytoplankton
 547 photosynthesis in situ. *Limnology and Oceanography*, 38, 1646-1665.

548 Krieger-Liszkay, A., Fufezan, C., & Trebst, A. (2008) Singlet oxygen production in photosystem II and
 549 related protection mechanism. *Photosynthesis Research*, 98, 551-561.

550 Kruk, J., & Karpinski, S. (2006) An HPLC-based method of estimation of the total redox state of
 551 plastoquinone in chloroplasts, the size of the photochemically active plastoquinone-pool and its
 552 redox state in thylakoids of Arabidopsis. *Biochimica et Biophysica Acta*, 1757, 1669-1675.

553 Kruk, J., & Trebst, A. (2008) Plastoquinol as a singlet oxygen scavenger in photosystem II. *Biochimica
 554 et Biophysica Acta*, 1777, 154-162.

555 Kruk, J., Szymanska, R., Cela, J., & Munne-Bosch, S. (2014) Plastochromanol-8: Fifty years of research.
 556 *Photochemistry*, 108, 9-16.

557 Ksas, B., Becuwe, N., Chevalier, A., & Havaux, M. (2015) Plant tolerance to excess light energy and
 558 photooxidative damage relies on plastoquinone biosynthesis. *Scientific Reports*, 5, 10919.

559 Lavergne, J., & Joliot, P. (1991) Restricted diffusion in photosynthetic membranes. *Trends in
 560 Biochemical Sciences*, 16, 129-134.

561 Li, Z., Wakao, S., Fischer, B.B., & Niyogi, K.K. (2009) Sensing and responding to excess light. *Annual
 562 Review Plant Biology*, 60, 239-260.

563 Lichtenthaler, H.K., & Weinert, H. (1970) The correlation between lipoquinone accumulation and
 564 plastoglobuli formation in the chloroplasts of *Ficus elastica* Roxb. *Zeitschrift für
 565 Naturforschung*, 25b, 619–623/266,

566 Lichtenthaler, H.K. (2007) Biosynthesis, accumulation and emission of carotenoids, alpha-tocopherol,
 567 plastoquinone, and isoprene in leaves under high photosynthetic irradiance. *Photosynthesis
 568 Research*, 92, 163-179.

569 Malkin, S., & Kok, B. (1966) Fluorescence induction studies in isolated chloroplasts. I. Number of
 570 components involved in the reaction and quantum yields. *Biochimica et Biophysica Acta*, 126, 413-
 571 432.

572 Millner, P.A., & Barber, J. (1984) Plastoquinone as a mobile redox carrier in the photosynthetic
 573 membrane. *FEBS Letters*, 169, 1-6.

574 Montillet, J.-L., Cacas, J.-L., Garnier, L., Montane, M.H., Douki, T., Bessoule, J.J., ..., Triantaphylides, C.
 575 (2004) The upstream oxylipin profile of *Arabidopsis thaliana*: A tool to scan for oxidative
 576 stresses. *Plant Journal*, 40, 439–451.

577 Norris, S.R., Barrette, T.R., & DellaPenna, D. (1995) Genetic dissection of carotenoid synthesis in
 578 arabidopsis defines plastoquinone as an essential component of phytoene desaturation. *Plant Cell*,
 579 7, 2139-2149.

580 Nowicka, B., & Kruk, J. (2010) Occurrence, biosynthesis and function of isoprenoid quinones.
 581 *Biochimica et Biophysica Acta*, 1797, 1587-1605.

582 Nowicka, B., Gruszka, J., & Kruk, J. (2013) Function of plastochromanol and other biological prenyllipids
 583 in the inhibition of lipid peroxidation- A comparative study in model systems. *Biochimica et*
 584 *Biophysica Acta*, 1828,233-240.

585 Pfannschmidt, T., Bräutigam, K., Wagner, R., Dietzel, L., Schröter, Y., Steiner, S., & Nykytenko, A. (2009)
 586 Potential regulation of gene expression in photosynthetic cells by redox and energy state:
 587 approaches towards better understanding. *Annals of Botany*, 103, 599-607.

588 Porfirova, S., Bergmüller, E., Tropsch, S., Lemke, R., & Dörmann, P. (2002) Isolation of an Arabidopsis
 589 mutant lacking vitamin E and identification of a cyclase essential for all tocopherol biosynthesis.
 590 *Proceedings National Academy Sciences USA*, 99, 12495-12500.

591 Pospíšil, P. (2014) The role of metals in production and scavenging of reactive oxygen species in
 592 photosystem II. *Plant Cell Physiology*, 55, 1224–1232.

593 Ramel, F., Birtic, S., Cuiné, S., Triantaphylidès, C., Ravanat, J.L., & Havaux, M. (2012) Chemical
 594 quenching of singlet oxygen by carotenoids in plants. *Plant Physiology*, 158, 1267-1278.

595 Rochaix, J.D. (2014) Redox regulation of thylakoid protein kinases and photosynthetic gene expression.
 596 *Antioxidants and Redox Signaling*, 18, 2184-2201.

597 Soll, J., Schultz, G., Joyard, J., Douce, R., & Block, M.A. (1985) Localization and synthesis of
 598 prenylquinones in isolated outer and inner envelope membranes from spinach chloroplasts.
 599 *Archives of Biochemistry and Biophysics*, 238, 290-299.

600 Szymańska, R., Nowicka, B., & Kruk, J. (2014) Hydroxy-plastochromanol and plastoquinone-C as singlet
 601 oxygen products during photo-oxidative stress in Arabidopsis. *Plant and Cell Environment*, 37, 1464-
 602 1473.

603 Szymanska, R., & Kruk, J. (2010) Plastoquinol is the Main Prenylipid Synthesized During Acclimation to
 604 High Light Conditions in Arabidopsis and is Converted to Plastochromanol by Tocopherol Cyclase.
 605 *Plant and Cell Physiology*, 51, 537-546

606 Triantaphylides, C., & Havaux, M. (2009) Singlet oxygen in plants: production, detoxification and
 607 signaling. *Trends in Plant Science*, 14, 219-228.

608 Van Eerden, F.J., Melo, M.N., Frederix, P.W.J.M., Periole, X., & Marrink, S.J. (2017) Exchange pathways
 609 of plastoquinone and plastoquinol in the photosystem II complex. *Nature Communications*,
 610 8,15214.

611 Vidi, P.A., Kanwischer, M., Baginsky, S., Austin, J.R., Csucs, G., Dörmann, P., ..., Bréhélin, C. (2006)
 612 Tocopherol cyclase (VTE1) localization and vitamin E accumulation in chloroplast plastoglobule
 613 lipoprotein particles. *Journal of Biological Chemistry*, 281,11225-11234.

614 Yadav, D.K., Kruk, J., Sinha, R.K., & Pospisil, P. (2010) Singlet oxygen scavenging activity of plastoquinol
 615 in photosystem II of high plants: Electron paramagnetic resonance spin-trapping study. *Biochimica*
 616 *et Biophysica Acta*, 1797, 1807-1811.

617 Zbierzak, A.M., Kanwischer, M., Wille, C., Vidi, P.A., Giavalisco, P., Lohmann, A., ..., Dörmann, P. (2009)
618 Intersection of the tocopherol and plastoquinone metabolic pathways at the plastoglobule.
619 *Biochemical Journal*, 425, 389-399.
620
621

Figure legends

FIGURE 1 Accumulation of PQ-9 in the Arabidopsis *vte1 SPS1oex* plants. A) α -tocopherol (α -Toc), PC-8 and PQ-9 concentrations in leaves of WT Arabidopsis, the single mutant *vte1*, the *SPS1* overexpressor *SPS1oex* and the double mutant *vte1 SPS1oex* (obtained by crossing *vte1* with the *SPS1oex* line #12 previously described in Ksas et al. (2015)). Data are mean values of 4 separate measurements $\pm$ SD. B) Picture of the different genotypes at the age of 4 weeks. C) Images of the efficiency of PSII-mediated electron transport Φ_{PSII} in plants illuminated with red light of PFD 230 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$. * and ***, different from WT at $P < 0.01$ and 0.001 (Student's t-test).

FIGURE 2 Localization of PQ-9 in the chloroplasts. A) Fractionation of chloroplast membranes by ultracentrifugation on sucrose gradient showing zones corresponding to the plastoglobules (PG), the envelopes and the thylakoids. The scale on the right shows the position of the different fractions collected for the PQ-9 analyses. B) Quantification of PQ-9 in 1-ml fractions from top (fraction 1) to bottom (fraction 32) for WT, *vte1*, *SPS1oex* and *vte1 SPS1oex* leaves. The same amounts of chlorophyll were loaded on the sucrose gradient. C) Estimation of the photochemically active and non-active pools of PQ-9 in Arabidopsis leaves. Data are mean values of 4 separate measurements $\pm$ SD. D) Induction of chlorophyll fluorescence in intact leaves. Insert: pool sizes of PSII 2-electron acceptors estimated from the fluorescence induction curves in the presence and absence of DCMU. ** and ***, different from WT at $P < 0.005$ and $p < 0.001$ (Student's t-test).

FIGURE 3 PQ-9 accumulation increases plant tolerance to photooxidative stress. Plants were exposed for 2 d to high light stress ($1500 \mu\text{mol photons m}^{-2} \text{s}^{-1}$) and low temperature (7°C). A) Picture of the plants after the high light treatment. B) Autoluminescence imaging of lipid peroxidation after high light treatment. The color palette shows the luminescence signal intensity from low (black) to high values (white). C) HOTE levels in leaves (in nmoles per mg fresh weight (F.W.)) before and after the high light stress. Data are mean values of 3 or 4 separate measurements $\pm$ SD. D) Maximal PSII photochemical efficiency, as measured by the Fv/Fm chlorophyll fluorescence ratio, before and after the high light stress. data are mean values of 8-to-12 separate measurements $\pm$ SD. *, ** and ***, different from WT and $P < 0.01$, 0.005 and 0.001 (Student's t-test).

FIGURE 4 $^1\text{O}_2$ scavenging by PQ-9. A) $^1\text{O}_2$ production by leaves in the light as measured by SOSG-EP fluorescence. Attached leaves infiltrated with SOSG were exposed for 20 min to light ($1500 \mu\text{mol photons m}^{-2} \text{s}^{-1}$) at 7°C . Data are mean values of minimum 4 or 5 measurements. B) Total PQ-C levels in leaves before and after high light stress ($1500 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ for 2d at 7°C). Data are mean

values (in nmoles per mg leaf fresh weight) of 3 separate measurements $\pm$ SD. C-D) $^1\text{O}_2$ production by thylakoids as measured by the spin probe TEMPONE and EPR spectroscopy. No signal was detected in the dark. C) TEMPONE EPR traces and D) quantification of the TEMPONE EPR signal. Data are mean values of 3 separate measurements $\pm$ SD. *, ** and ***, different from WT at $P < 0.01$, $P < 0.005$ and $P < 0.001$ (Student's t-test).

FIGURE 5 Dynamics of the PQ-9 pools. Effects of high light stress (HL = 2.5 h at $1500 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$) on A) the PQ-9 levels in Arabidopsis leaves and B) on the repartition of PQ-9 between the photochemically active pool (in the thylakoids) and the non-active pool (in the plastoglobules and the envelopes). Data are mean values of minimum 3 separate measurements $\pm$ SD. C) PQ-C concentration in different chloroplast fractions, plastoglobules (PG), envelopes (ENV) and thylakoids separated by ultracentrifugation on sucrose gradient (fractions 1-4, 5-10 and 30-32, respectively, see Fig. 2B). LL corresponds to the growth light conditions ($150 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$). Data are mean values of 3 experiments $\pm$ SD. *, ** and ***, different from LL at $P < 0.01$, $P < 0.005$ and $P < 0.001$ (Student's t-test).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5