Recent biomonitoring reports on phosphate ester flame retardants. A short review

Anne Marie Saillenfait¹ · Sophie Ndaw¹ · Alain Robert¹ · Jean Philippe Sabaté¹

¹ Institut National de Recherche et de Sécurité, Rue du Morvan, CS, 60027, 54519 Vandoeuvre Cedex, France

✉ Anne Marie Saillenfait

Tel.: + 33 83502039.

Email address: anne-marie.saillenfait@inrs.fr

Abstract

Organophosphate triesters (PEFRs) are used increasingly as flame retardants and plasticizers in a variety of applications, such as building materials, textiles, and electric and electronic equipment. They have been proposed as alternatives to brominated flame retardants. This updated review shows that biomonitoring has gained incrementally greater importance in evaluating human exposure to PEFRs, and it holds the advantage of taking into account the multiple potential sources and various intake pathways of PEFRs. Simultaneous and extensive internal exposure to a broad range of PEFRs has been reported worldwide. Their metabolites, mainly dialkyl or diaryl diesters, have been used as biomarkers of exposure and have been ubiquitously detected in the urine of adults and children in the general population. Concentrations and profiles of PEFR urinary metabolites are seen to be variable and are highly dependent on individual and environmental factors, including age, country regulation of flame retardants, and types and quantities of emissions in microenvironments, as well as analytical procedures. Additional large biomonitoring studies, using a broad range of urinary diesters and hydroxylated metabolites, would be useful to improve the validity of the biomarkers and to refine assessments of human exposure to PEFRs.

Keywords

Flame retardant · Organophosphates · Exposure assessment · Human biomonitoring · Health

Introduction

Organophosphorous flame retardants are one of the most common groups of flame retardants (FRs). They primarily consist of organophosphate triesters (PEFRs). The total global consumption of organophosphorous compounds used as FRs was estimated to be about 207 000 tons in 2004. In 2006, chlorinated phosphate and non-chlorinated organophosphorous FRs represented 11 and 10 % of FR consumption in Europe, respectively (NEG, 2009; Arcadis EBRC, 2011; EFRA, 2007). Since the early 2000s, brominated flame retardants such as certain polybrominated diphenyl ethers (PBDE), have been progressively phased out or seen their use restricted in many regions of the world (e.g., North America, Japan, and Europe), because of their bioaccumulation, persistence, and potential health effects. The halogen-free PEFRs are considered as possible alternatives to PBDE, and their production and use have been increasing over the past few years. PEFRs are also applied as plasticizers in polymers such as PVC (e.g., aryl phosphates), cellulosic fibers, polyurethane foams (e.g., chlorinated phosphate), and engineering plastics (e.g., polycarbonate/acrylonitrile butadiene styrene-PC/ABS, and polyphenylene oxide/high impact polystyrene-PPO/HIPS). Other applications of phosphate esters are as anti-foam agents, additives in hydraulic fluids, and lubricants. PEFRs have a broad application field and are extensively used alone or in combination with other FRs in a variety of industries, including plastics, furniture, textiles, construction, electrical engineering and electronics, transportation (e.g., road and rail vehicles), and the petroleum industry (NEG, 2009; ATSDR, 2012). Triphenyl phosphate (THPP) is also present in personal care products such as nail polish (Mendelsohn et al., 2016).

PEFRs are frequently applied as chemical additives and they are not chemically bound to the polymers (i.e. unlike reactive FRs). They can be released from treated industrial and commercial products by abrasion, leaching and/or volatilization during their lifetime. They have been detected in a wide range of environmental samples around the world (e.g., indoor dust) and concern about human exposure to PEFRs is increasing. In addition, some recent epidemiological studies have suggested that certain PEFRs may have possible health effects, such as interference with endocrine and reproduction functions. Tris(2-chloroethyl) phosphate (TCEP), tris(1,3-dichloro-2-propyl) phosphate (TDCIPP), and tris(2,3-dibromopropyl)phosphate (TDBPP) are listed as carcinogenic substances under California's proposition 65 (OEHHA, 2017). In Europe, several PEFRs have been classified as substances suspected of causing cancer (e.g., TCEP, TDCIPP, tributyl phosphate - TNBP) and/or that may damage fertility (e.g., TCEP, tri-*ortho*-cresyl phosphate-ToCP) (ECHA, 2018).

While the occurrence of PEFRs in indoor dust has been extensively described across the world, less is known about internal exposure to PEFRs in humans. Several human studies have used urinary biomarkers of PEFRs to monitor exposure to these FRs among workers and the general population.

This short review compiles the available data on PEFRs in human urine published from 2011 to May 2018. A comprehensive search was performed in the Pubmed database using the search terms "organophosphate flame retardant" and "exposure", or "flame retardant" and "urine". Only full text articles were reviewed. Studies were included if the PEFRs were measured in a biological matrix (e.g., urine, hair, milk). There were no restrictions on the size and age of the study population, geographical region, or study design (e.g., pooled data).

Environmental occurrence and routes of exposure

PEFRs have been found worldwide in diverse outdoor environments, including river water, groundwater, and wastewater, with individual concentrations ranging from several ng/l to tens of µg/l (Van der Veen et al., 2012; Wei et al., 2015; Ali et al., 2017). They have been ubiquitously detected in

floor and surface dust from various indoor environments, including private houses, vehicles, and various public and work places (e.g., offices, daycare centers, electronic equipment stores and recycling plants, and hospitals). PEFRs concentrations in floor and surface dust were in the range of 0.02 ng/g to tens of µg/g. The concentrations of PEFRs measured in indoor air were generally around tens to hundreds of ng/m³. Higher PEFR levels in indoor air or dust have occasionally been reported in occupational settings (e.g., recycling of electronics) or microenvironments (e.g., cars) (Wu et al., 2016; Ali et al., 2017; Zhou et al., 2017; Bello et al., 2018; Björnsdotter et al., 2018). Concentrations in outdoor air are approximately 1 to 4 orders of magnitude less than in indoor air (Wei et al., 2015). The ubiquity of PEFRs in the environment indicates that the general population is likely to be exposed to several of these chemicals, through multiple sources, on a daily basis.

PEFRs can enter the human body *via* several routes. Recent studies have shown that dermal absorption may contribute substantially to the total body burden of PEFRs (Abdallah et al., 2016; Mendelsohn et al., 2016; Liu et al., 2017; Bello et al., 2018; Frederiksen et al., 2018). Ingestion of dust and dermal exposure to dust and treated materials (e.g., clothes and furniture) are considered primary sources of exposure to PEFRs. For volatile or semi-volatile PEFRs (e.g., TCEP, tris(1-chloro-2-propyl) phosphate - TCIPP), air and suspended particles inhalation may be a significant intake pathway (Schreder et al., 2016; Xu et al., 2016; He et al., 2018c). Ingestion of contaminated food (e.g., by migration from plastic packaging) may contribute to oral intake of PEFRs, but its contribution appears to vary substantially between compounds, as well as between and within populations (Zhang et al., 2016; Zheng et al., 2016; Xu et al., 2017; Poma et al., 2017, 2018).

Human health effects

The toxicological profiles of halogenated and non-halogenated PEFRs have been reviewed by several (environmental) agencies, in particular to evaluate their suitability as alternatives for PBDE FRs (NEG, 2009; ATSDR, 2012; Van den Eede et al., 2012; US EPA, 2015; Ministry of Environment and Food of Denmark, 2016). Assessments almost entirely relied on experimental studies. Critical effects were found to differ from one compound to another. A few PEFRs were identified as being toxic to the male reproductive system (i.e. TCEP, TDCIPP, ToCP), potentially carcinogenic (i.e. TCEP and TDCIPP), and/or toxic to specific organs (i.e. kidney for TCEP). Because of their toxic potential, the chlorinated PEFRs TCEP and TDCIPP are presently subject to regulations in several countries (mainly Northern America, Europe, Japan) (OEHHA, 2017; Canada Safety Consumer Act, 2018; ECHA, 2018). The US EPA has established an oral reference dose of 0.01 mg/kg/day for TCIPP, TEHP and TNBP; 0.02 mg/kg/day for TmCP and TDCIPP; and 0.007 mg/kg/day for TCEP (US EPA, 2017).

Despite the increasing use of a wide range of PEFRs and the ubiquitous exposure to PEFRs among the general population, human data on the potential health effects of PEFRs are still limited, especially regarding long term exposure and the risks for children. The epidemiological studies published since 2010 mainly addressed respiratory outcomes (asthma, rhinitis), and endocrine and reproductive effects (Table 1). Results of a few studies have raised concern about the possible association between exposure to some PEFRs, and alteration of thyroid hormone regulation and male reproduction (e.g., sperm quality) (Meeker et al., 2010; Meeker et al., 2013a; Hoffman et al., 2017c; Preston et al. 2017; Soubry et al., 2017; Carignan et al., 2018a). However, at present, there is not enough consistent information from which to draw firm conclusions about the adverse health effects of PEFRs (as a class or specific) in humans.

The isomer ToCP has proven to be neurotoxic and to inhibit both cholinesterase and neuropathy target esterase (NTE) activity (NEG, 2009; ATSDR, 2012; Van den Eede et al., 2012; US EPA, 2015; Ministry of Environment and Food of Denmark, 2016). Worldwide, it has been associated with numerous cases of delayed neuropathy and paralysis of the extremities in humans (Petroianu et al., 2016). Consequently, there has been a significant reduction in the commercial use of ToCP, e.g., in aircraft engine oil.

Biomarkers of exposure

Cholinesterase activity

The neurotoxic properties of ToCP have mainly been attributed to its metabolite cresyl saligenin phosphate. This reactive intermediate binds covalently to a serine moiety of butyrylcholinesterase in blood. The resulting adducts can be determined by mass-spectrometry and have been proposed as a biomarker for measuring exposure to ToCP (Schopfer et al., 2014; Tacal et al., 2014; Johnson et al., 2015). In addition, the American Conference of Governmental Industrial Hygienists (ACGIH) recommended erythrocyte cholinesterase activity as a biological exposure index (BEI) for ToCP. However this biomarker is not specific and can be inhibited by other OPs, such as organophosphorous pesticides (NEG, 2009; ATSDR, 2012).

Urinary biomonitoring

Urinary PEFRs or their metabolites appear to be the preferred non-invasive biomarkers for identifying and quantifying human exposure to PEFRs. They provide integrated information on total body burden, covering all types of sources and exposure pathways (i.e. inhalation, dermal absorption, and oral uptake), and they can be used to quantify an individual's exposure.

Diester metabolites

Information on the metabolism of PEFRs in humans is still limited and there are differences in the information available for different compounds. A common metabolic pathway has been proposed for the three types of PEFR triesters, i.e. trialkyl, triaryl, and trihaloalkyl/aryl phosphate esters. This was mainly based on *in vivo* studies in rodents and *in vitro* studies using human hepatocytes or liver fractions (Ballesteros-Gomez et al., 2015a and b; Hou et al., 2016; Van den Eede et al., 2013a, 2015a, 2016a, b and c). The first steps in the biotransformation of these triesters lead to the rapid formation of diesters or monoesters by hydrolysis of one or two ether bonds between the phosphate group and the substituent, and to a variety of hydroxylated metabolites that undergo glucuronide and sulfate conjugation. Indeed, several dialkyl or diaryl phosphates have been detected in human urine, including bis(2-chloroethyl) phosphate (BCEP), bis(1-chloro-2-propyl) phosphate (BCIPP), bis(1,3-dichloro-2-propyl) phosphate (BDCIPP), dibutyl phosphate (BNBP), and diphenyl phosphate (DPHP) (Table 1). These diesters are expected to be important and stable metabolites of TCEP, TCIPP, TDCIPP, TNBP, and triphenyl phosphate (TPHP), respectively (Table 2). Hence, most biomonitoring studies have focused on the determination of dialkyl or diaryl phosphate metabolites in urine to quantify human exposure levels to PEFRs.

However, there have been concerns regarding the use of urinary DPHP as a biomarker of exposure levels of the parent TPHP. DPHP may lack specificity since other aryl organophosphate esters containing at least two phenyl substituents [e.g., bisphenol A bis(diphenyl phosphate) and resorcinol bis(diphenyl) phosphate] have the potential to form DPHP after being hydrolysed and may contribute to DPHP urinary levels (Ballesteros-Gomez et al., 2015b; He et al., 2018a). In addition,

DPHP itself is currently a commercially available product (e.g., as catalyst for resin manufacturing). Therefore, Van den Eede et al. (2016b) recommended using DPHP as a biomarker of aryl-PFRs rather than of TPHP only. In contrast with TPHP, the production of DPHP from 2-ethylhexyl diphenyl phosphate by human serum hydroxylase *in vitro* was found to be minor and thus it was not considered to be likely confounding factor (Van den Eede et al, 2016b).

Other metabolites

A few hydroxylated metabolites of PEFRs have recently been identified in urine samples from adults and children (Dodson et al., 2014; Van den Eede et al., 2015b; Hammel et al., 2016; Kosarac et al., 2016; Su et al., 2016; Bui et al., 2017; He et al., 2018a; Hoffman et al., 2017a, 2018; Phillips et al., 2018; Völkel et al., 2018). Urinary bis(2-butoxyethyl)-(2-hydroxyethyl) phosphate (BBOEHEP) was used to monitor exposure to tris(2-butoxyethyl) phosphate (TBOEP) (Van den Eede et al, 2015b; He et al., 2018a; Völkel et al., 2018). Hydroxylated metabolites of TPHP (i.e. 4-hydroxyphenyl diphenyl phosphate, 4-hydroxyphenyl phenyl phosphate), have been considered as potential specific urinary biomarkers of TPHP exposure (Van den Eede et al., 2013a, 2015b; Dodson et al., 2014; Su et al., 2016). However, they were only occasionally detected, and at very low levels, in human urine samples (glucuronide and sulfate conjugates, or the sum of free form and conjugates) (Van den Eede et al., 2015b, 2016b; Su et al., 2016). In several studies, the hydroxylated metabolite of TCIPP, bis(1-chloro-2-propyl) 1-hydroxy-2-propyl phosphate (BCIPHPP), appeared to be a major urinary metabolite and therefore a candidate biomarker of human exposure to this PEFR (Van den Eede et al., 2015b; Butt et al., 2016; Hammel et al., 2016; Hoffman et al., 2017a; Bello et al., 2018; He et al., 2018a; Phillips et al., 2018). Total hydroxylated metabolite (i.e. the sum of free and conjugated forms) was usually measured after enzymatic de-conjugation treatment of the urine samples with sulfatase and β -glucuronidase. The free form of BCIPHPP was reported to be barely detectable (Kosarac et al., 2016).

Unmetabolized PEFRs

The parent compounds have also been tested as potential urinary biomarkers of exposure to the OP triester FRs. Considering their notable presence in urine, monitoring of the unchanged TCPE and TEHP along with their corresponding diester metabolites was considered useful for better estimation of the actual exposure (Dodson et al., 2014; He et al., 2018a). With the exception of TCEP and TEHP, unchanged PEFRs were detected in lower frequencies and concentrations than their related diester metabolites, suggesting that they were less suitable biomarkers (Van den Eede et al., 2015b; He et al., 2018a). Furthermore, additive PEFRs can leach from treated rubber and plastic storage materials and possible background contamination of collected samples must therefore be considered.

Chemical analysis

Sensitive methods are being developed to improve the limits of detection and concurrently quantify a broad number of chlorinated and non-chlorinated diester and/or selected hydroxylated OP metabolites in human urine samples. Typical analytical techniques, including gas chromatography-tandem mass spectrometry (GC-MS/MS) (Schindler et al. 2009a,b), high or ultra-performance liquid chromatography-tandem mass spectrometry (HPLC-MS/MS or UPLC-MS/MS), with electrospray or atmospheric pressure chemical ionization (ESI or APCI) have been used successfully in numerous biomonitoring studies (Cooper et al., 2011; Reemtsma et al., 2011; Van den Eede et al., 2013b; Su et al., 2015; Kosarac et al., 2016; Petropoulou et al., 2016; Jayatilaka et al., 2017), as has high resolution mass spectrometry (UPLC-HRMS) (Cequier et al., 2014). These same sensitive methods are also being developed for use with other non-invasive matrices such as hair, nails and human milk (Sundkvist et al., 2010; Lu et al., 2014; Kucharska et al. 2014; Liu et al., 2015; Alves et al., 2017).

Occurrence of PEFRs metabolites in human urine

General population

Metabolites of PEFRs, essentially the diesters, were omnipresent in the urine samples collected from the general population across different countries, and there was simultaneous exposure to several PEFRs (Table 3). Reported occurrences and concentrations varied substantially between individual PEFR compounds.

BDCIPP and DPHP were the most commonly detected diester metabolites in the urine of children, mothers, and the general population, and were also the most frequently analyzed (Table 3). Median levels of BDCIPP and DPHP were generally in the range of $\mu\text{g/l}$ (about 0.1-3 $\mu\text{g/l}$), but values of hundreds of $\mu\text{g/l}$ were reported in urine samples of a few individuals from various geographic areas. Within each of the different studies, concentrations were highly variable between individuals and could differ by two orders of magnitude. DPHP was consistently found at high frequencies (in most cases >90 %) in the general population in Europe, the United States and China, suggesting ubiquitous exposure to DPHP or its parent compounds (e.g., TPHP or other aryl-PEFRs such as 2-ethylhexyl diphenyl phosphate) around the world.

Highly variable detection frequencies were reported for BCIPP, BBNP, and BBOEP. In general, their median levels were around, or less than, 0.3 $\mu\text{g/l}$.

Information on the occurrence of BCEP and isopropylphenyl phenyl phosphate (ipPP) is more limited. These were detected in more than half of the urine samples in the large majority of studies that monitored these metabolites. In most studies, their median concentrations were in the range of 0.2-2 $\mu\text{g/l}$.

In almost all available studies, di-*ortho*-cresyl phosphate (DoCP) and/or di-*para*-cresyl phosphate (DpCP) (determined alone or together) were detected only occasionally, and/or at relatively low levels (i.e. median levels <0.02 $\mu\text{g/l}$) in recent studies in China and USA, suggesting limited exposure to the precursors of these metabolites in these general populations (Schindler et al., 2013; Fromme et al., 2014; Kosarac et al., 2016; Lu et al., 2017; Romano et al., 2017; Chen et al., 2018; Ospina et al., 2018). However, higher frequencies were reported in some occupationally exposed populations (Jayatilaka et al., 2017; Tao et al., 2018). DpCP was more abundant than di-*meta*-cresyl phosphate (DmCP) and DoCP. The synthesis and commercial compositions of TCP have in fact changed over time. Because of its neurotoxic properties, efforts have been made to minimize the amount of the *ortho* isomer present in commercial products containing TCP (NEG, 2009; ATSDR, 2012; US EPA, 2015).

Other PEFRs metabolites were more rarely analyzed. Tert-butyl phenyl phenyl phosphate (tb-PPP) and bis(2-ethylhexyl) phosphate (BEHP) were detected infrequently (Su et al., 2015; Butt et al., 2016; Castorina et al., 2017b; Hoffman et al., 2017a; Soubri et al., 2017; Carignan et al., 2018a and b; Deziel et al., 2018; He et al., 2018a; Hoffman et al., 2018; Ingle et al., 2018; Sun et al., 2018). Dibenzyl phosphate (DBzP) was not detected in urine samples collected in the U.S.A. (Romano et al., 2017; Jayatilaka et al., 2017; Ospina et al., 2018).

Workers

In addition to the general population, urinary biomarkers have been used to assess exposure to PEFRs in a number of workplaces (Table 4). There are some indications that internal exposure may be higher than the background exposure of the general population during several types of occupational activity. For example, a recent study conducted in Australia showed that urinary levels of BCIPHP among spray polyurethane foam applicators were approximately fifty times higher than urinary levels found in the general population (Bello et al., 2018). Numerous other worker groups are expected to be more heavily exposed than the general population, especially when workers are in direct contact with large volumes of PEFRs as pure chemicals or at high concentrations in technical formulations at industrial sites and in manufacturing (e.g., at electronics dismantling facilities or electronic goods recycling areas). Measurements of PEFRs in air and dust in various occupational settings have also shown that the work environment may noticeably contribute to external exposure to PEFRs (Makinen et al., 2009; Ali et al., 2014; Wei et al., 2015; Zheng et al., 2017; Zhou et al., 2017; Muenhor et al., 2017; Bello et al., 2018; Ceballos et al., 2018; Shen et al., 2018). Nevertheless, information on the nature and extent of occupational exposures to PEFRs, especially in terms of measurements of an individual's internal exposure, is still limited and warrants further investigations (characterization, quantification, and contribution to total PEFR burden).

Possible bias, limitations and strengths of the reviewed studies

The available biomonitoring data should be analyzed in the context of several influencing factors that have already been identified in a number of studies on the evaluation of human internal exposure to PEFRs.

Concentrations of urinary PEFR metabolites varied greatly both between the populations studied and from individual to individual within cohorts (Hoffman et al., 2017a; Preston et al., 2017). Except for BDCIPP and DPHP which were typical worldwide contaminants, there was no strong common pattern for the compositional profile of urinary PEFR metabolites. This may be explained by differences in FR regulations, dietary habits, lifestyle, and use of PEFRs in household products and indoor environments (e.g., building material), between the various countries and/or study locations (Carignan et al., 2013; Butt et al., 2016; Lu et al., 2017; Chen et al., 2018; He et al., 2018b). Other factors were reported to have an impact on urinary PEFR metabolite concentrations, including timing (e.g., season of collection) (Hoffman et al., 2017a and b; Deziel et al., 2018; Ingle et al., 2018; Phillips et al., 2018), sex (e.g., women tend to have higher levels of DPHP than men, Hoffman et al., 2015b; Preston et al., 2017; He et al., 2018a; Ospina et al., 2018), behavior and activity patterns (e.g., hand washing and cleaning routines, nail painting) (Abdallah et al., 2016; Mendelsohn et al., 2016; He et al., 2018b) and age (Van den Eede et al., 2015b; Lu et al., 2017; He et al., 2018a; Ospina et al., 2018; Sun et al., 2018). Urinary concentrations of the main PEFR metabolites were generally higher in toddlers than in adults (Butt et al., 2014 and 2016; Cequier et al., 2015; Hoffman et al., 2015a; Van den Eede et al., 2015b; Chen et al., 2018; He et al., 2018a and b; Ospina et al., 2018). This is an international trend, generally attributed to the tendency of young children to crawl on the floor and to their elevated hand-to-mouth contact behavior, both of which result in increased oral and dermal contact with indoor settled dust and with products containing these chemicals (e.g., plastic toys). Differences in pharmacokinetics with age cannot be excluded.

Long-term temporal trends in the urinary levels of some PEFR metabolites have been reported among adults and/or children in the United States. Concentrations of BDCIPP in urine samples collected in 2014-2015 were 16.5 times higher than those collected in 2002-2003, while

concentrations of DPHP increased at much lower rates until 2011 (Hoffman et al., 2017b). This may be related to changes in the use of specific PEFRs to meet the more stringent regulation of certain FRs, and improvements in the fire safety standards required for finished consumer products (e.g., furniture and textiles). For example, TCEP and TDCPP have recently been restricted or banned in children's products in several states in the USA (Vermont General Assembly, 2013; US EPA, 2015; Council of Columbia, 2016, Department of Ecology State of Washington, 2016).

In addition to the studied populations and the sources of PEFR emissions, sampling strategies may affect the study results. Human observations (Carignan et al., 2016) and *in vitro* and *in vivo* rat studies suggest that PEFRs are rapidly metabolized and eliminated in urine. PEFR half-lives in humans are generally estimated to be on the order of a few hours. The use of single spot urine samples in most studies may not represent metabolite concentration over time and may contribute to the variability in the metabolite concentrations. However, a slower urinary elimination of some metabolites (i.e. BBOEP) was recently observed in volunteers following an oral administration of TBOEP (Völkel et al., 2018). Several studies collected multiple samples over one day or the course of the study to limit within-subject variability (Meeker et al., 2013b; Hoffman et al., 2015b; Cequier et al., 2015; Su et al., 2015; Carignan et al., 2016; Hammel et al., 2016; Preston et al., 2017; He et al., 2018b; Phillips et al., 2018).

Analytical treatment of the biological samples may be critical for the measurement of PEFR urinary metabolites, e.g., conditions of urine collection and storage (Petropoulou et al., 2016; Carignan et al., 2017). Differences in the detection and quantification limits of the analytical methods employed to quantify urinary metabolites may also account for the broad range of detection rates of some metabolites within and/or across studies. Van den Eede et al. (2013b) showed that improvement of the LOQ method resulted in a higher detection frequency of BCEP and BDCIPP. In several studies, the method limit of detection (MLOD) of the hydroxylated metabolite of TCIPP, BCIPHPP, was much lower than that of the diester, BCIPP (at least ten folds - Butt et al., 2016; Hammel et al., 2016; Hoffman et al., 2017a; He et al., 2018a). BCIPHPP was in fact found at a higher incidence than BCIPP in recent biomonitoring studies that measured both metabolites in urine samples from the general population (Butt et al., 2016; Hammel et al., 2016; Hoffman et al., 2017a and 2018).

The biotransformation of PEFRs has not been extensively investigated in animals and humans and their potential metabolic pathways are principally based on qualitative *in vitro* analyses. *In vivo*, the triesters may undergo very little transformation, and/or several major metabolites other than diesters may be formed (Hou et al., 2016; Völkel et al., 2018). In addition, urine may not be the sole excretion pathway for certain PEFRs. Diester metabolites were the main metabolites targeted in urine for all PEFRs. However, there may be qualitative and quantitative metabolic differences between the compounds and/or between the metabolite kinetics. If the measured metabolite was not the best urinary biomarker of exposure, this would lead to underestimation of exposure for some PEFRs. For example, the diester metabolites of TBOEP and TCIPP were not always the main metabolites formed *in vitro* by human liver preparations. Several potential hydroxylated derivatives have been considered for urinary monitoring of certain PEFRs (e.g., BBOEHEP for TBOEP and BCIPHPP for TCIPP) (Van den Eede et al., 2015b; Butt et al., 2016; Hammel et al., 2016; Hoffman et al., 2017a and 2018; Bello et al., 2018; He et al., 2018a and b; Phillips et al., 2018). In fact, BBOEP and BBOEHEP were detected in 80 % of urine samples from volunteers orally administered a single dose of TBOEP (20 µg/kg b.w.), with comparable median values (0.16 and 0.18 µg/l, respectively) (Völkel et al., 2018). However, the maximum concentration of BBOEHEP was much higher than that of BBOEP (3700 and 69 pmol/kg b.w., respectively) and was reached within 1-2 hours. In contrast, BBOEP showed some maxima within 25 hours, before a smooth decline.

Several biomonitoring studies with large cohort size provide robust information on general population exposures to PEFRs. They related to a representative sample of the U.S. general population (Ospina et al., 2018), adults in China (Lu et al., 2017) and the United States (Carignan et al., 2018a and b), children in China (Chen et al., 2018) and Germany (Fromme et al., 2014), and pregnant women in the United States (Castorina et al., 2017; Hoffman et al., 2017a).

No consistent and/or uniform correlation could be established between urinary levels of some PEFR metabolites (mainly diesters) and the concentrations of the corresponding parent compounds in hand wipes or in indoor dust samples from various microenvironments (Carignan et al., 2013; Meeker et al., 2013b; Dodson et al., 2014; Fromme et al., 2014; Cequier et al., 2015; Hoffman et al., 2015b; Hammel et al., 2016; Castorina et al., 2017b; Larsson et al., 2018; Phillips et al., 2018; Tao et al., 2018; Völkel et al., 2018). Associations were generally specific to the PEFR. Some weak or positive correlations were reported, but inconstantly, for the pairs TCEP/BCEP, TDCIPP/BCIPP, TPHP/DPHP, and/or TBOEP/BBOEP. Urinary biomarkers are indicators of integrated personal exposure. Each PEFR may have several different sources and pathways of exposure, and dust sampled from specific indoor microenvironments may not be the sole and/or the primary contributor to the body burden.

Occurrence of PEFRs in other human samples

Most human biomonitoring studies have used urine as biological matrix to evaluate exposure to PEFRs. Less is known about the possibility of using PEFR levels in segments of hair and/or nails as retrospective non-invasive biomarkers for PEFR monitoring. The main PEFRs (unchanged compounds) were detected in most of the hair samples collected in various countries (e.g., TCIPP, TDCIPP, TPHP) (Table 5). Levels measured in hair were highly variable between individuals, with concentrations ranging from ng/g to high concentrations of several µg/g within the study populations (e.g., TDCIPP and TPHP, Kurcharska et al., 2015a; Liu et al., 2016). It was suggested that PEFR levels in the hair are derived from a combination of both external exposure from air and dust and internal exposure. PEFRs in hair reflect long term exposure while the occurrence of PEFR metabolites in urine most likely corresponds to recent exposure (Kurcharska et al., 2015b; Alves et al., 2017).

A number of studies have reported the presence of PEFRs in other human tissues and body fluids. PEFRs were frequently detected in placenta (Ding et al., 2016; Zhao et al., 2017) and breast milk (Sunddkvist et al., 2010; Kim et al., 2014; He et al., 2018a). The median concentration of total PEFRs was around 10-100 ng/g of lipids in breast milk from Sweden and several Asian countries, indicating that substantial exposure occurs at a young age via breastfeeding (Sunddkvist et al., 2010; Kim et al., 2014). The parent compounds (Liu et al., 2016; Zhao et al., 2016; Li et al., 2017; Ma et al., 2017; Qiao et al. 2017) and their metabolites (Bui et al., 2017) were found in human serum and blood in a few studies. The metabolism of parent PEFRs tends to occur rapidly and the measurement of metabolites concentrations in urine is generally preferred to the invasive measurement of the non-metabolized chemicals in serum for exposure assessment.

Conclusion

This short review shows that the use of urinary levels of PEFRS metabolites for monitoring internal human exposure to these emerging pollutants is widespread and has gained increasing attention over the past few years. The biomonitoring studies confirm ubiquitous exposure of the general population to PEFRS all over the world, and potentially higher exposures in children and among a number of occupational populations. The levels and compositional patterns of urinary metabolites varied as a function of factors, such as the location and time of sampling. Further

information on the toxicokinetics of PEFs in humans and the continued development and validation of bioanalytical methods will allow refinement of the current biomarkers of exposure to these chemicals. Additional biomonitoring data on PEFs are still needed in order to reduce the uncertainty in estimating human exposure, to identify the populations at risk and any possible associations with adverse health effects, to follow exposure trends, and to evaluate governmental prevention strategies and programs.

Conflict of interest

The Authors declare that they have no conflict of interest.

References

- Abdallah MAE, Pawar G, Harrad S (2016) Human dermal absorption of chlorinated organophosphate flame retardants; implications for human exposure. *Toxicol Appl Pharmacol* 291: 28-37
- Agency for Toxic Substances and Disease Registry (ATSDR). U.S. Department of Health and Human Services. Public Health Service. (2012) Toxicological profile for phosphate ester flame retardants.
- Ali N, Shahzad K, Rashid MI, Shen H, Ismail IMI, Eqani SAMAS (2017) Currently used organophosphate and brominated flame retardants in the environment of China and other developing countries (2000-2016). *Environ Sci Pollut Res* 24:18721-18741
- Ali N, Mehdi T, Malik RN, AMAS Eqani S, Kamal A, Dirtu AC, Neels H, Covaci A (2014) Levels and profile of several classes of organic contaminants in matched indoor dust and serum samples from occupational settings of Pakista. *Environ Pollut* 193:269-276
- Alves A, Covaci A, Voospoels S (2017) Method development for assessing the human exposure to organophosphate flame retardants in hair and nails. *Chemosphere* 168:692-698
- Anderson J (2015) Comment on Schindler, BK; Weiss, T; Schütze, A; et al. Occupational exposure of air crews to tricresyl phosphate isomers and organophosphate flame retardants after fume events”, *Arch Toxicol* (2013) 87:645-648. *Arch Toxicol* 89:259-261
- Araki A, Saito I, Kanazawa A, Morimoto K, Nakayama K, Shibata E, Tanaka M, Takigawa T, Yoshimura T, Chikara H, Saijo Y, Kishi R (2014) Phosphorous flame retardants in indoor dust and their relation to asthma and allergies of inhabitants. *Indoor Air* 24:3-15
- Arcadis EBRC (2011) Evaluation of data on flame retardants in consumer products – Final report. Contract number 17.020200/09/549040. European Commission Health and Consumer Products. <https://www.ifv.nl/kennisplein/Documents/20110426-ec-arcadis-ebrc-evaluation-of-data-on-flame-retardants-in-consumer-products.pdf>
- Ballesteros-Gomez A, Erratico CA, Van den Eele N, Ionas AC, Leonards PEG, Covaci A (2015a) In vitro metabolism of 2-ethylhexyldiphenyl phosphate (EDHPP) by human liver microsomes. *Toxicol Lett* 232:203-212
- Ballesteros-Gomez A, Van den Eede N, Covaci A (2015b) In vitro human metabolism of the flame retardant resorcinol bis(diphenylphosphate) (RDP). *Environ Sci Technol* 49:3897-3904
- Bello A, Carigan CC, Xue Y, Stapleton HM, Bello D (2018) Exposure to organophosphate flame retardants in spray polyurethane foam applicators: Role of dermal exposure. *Environ Int* 113:55-65
- Bergh C, Aberg KM, Svartengren M, Emenius G, Ostman C (2011) Organophosphate and phthalate esters in indoor air: a comparison between multi-storey buildings with high and low prevalence of sick building symptoms. *J Environ Monit* 13:2001-2009
- Björnsdotter MK, Romera-Garcia E, Borrell J, de Boer J, Rubio S, Ballesteros-Gomez A (2018) Presence of diphenyl phosphate and aryl-phosphate flame retardants in indoor dust from different microenvironments in Spain and the Netherlands and estimation of human exposure. *Environ Int* 112:59-67

- Bui TT, Xu F, Van den Eede N, Cousins AP, Covaci A, Cousins IT (2017) Probing the relationship between external and internal human exposure of organophosphate flame retardants using pharmacokinetic modeling. *Environ Pollut* 230:550-560
- Butt CM, Hoffman K, Chen A, Lorenzo A, Congleton J, Stapleton HM (2016) Regional comparison of organophosphate flame retardants (PFR) urinary metabolites and tetrabromobenzoic acid (TBBA) in mother-toddler pairs from California and New Jersey. *Environ Int* 94:627-634
- Butt CM, Congleton J, Hoffman K, Fang M, Stapleton HM (2014) Metabolites of organophosphate flame retardants and 2-ethylhexyl tetrabromobenzoate in urine from paired mothers and toddlers. *Environ Sci Technol* 48:10432-10438
- Canada Safety Consumer Act. Ministry of Justice. (2018) Consolidation S.C. 2010, c. 21 Current to March 26, 2018 Last amended on December 12, 2016. <http://laws-lois.justice.gc.ca/PDF/C-1.68.pdf>
- Canbaz D, van Velzen JM, Hallner E, Zwinderman AH, Wickman M, Leonards PEG, van Ree R, van Rijt LS (2016) Exposure to organophosphate and polybrominated diphenyl ether flame retardants via indoor dust and childhood asthma. *Indoor Air* 26:403-413
- Carignan CC, Minguez-Alarcon L, Williams PL, Meeker JD, Stapleton HM, Butt CM, Toth TL, Ford JB, Hauser R, (2018a) Paternal urinary concentrations of organophosphate flame retardant metabolites, fertility measures, and pregnancy outcomes among couples undergoing in vitro fertilization. *Environ Int* 111:232-238
- Carignan CC, Minguez-Alarcon L, Butt CR, Williams PL, Meeker JD, Stapleton HM, Toth TL, Ford JB, Hauser R (2018b) Urinary concentrations of organophosphate flame retardant metabolites and pregnancy outcomes among women undergoing in vitro fertilization. *Environ Health Perspect* 125:087018-1-087018-8
- Carignan CC, Butt CM, Stapleton HM, Meeker JD, Minguez-Alarcon L, Williams PL, Hauser R (2017) Influence of storage vial material on measurement of organophosphate flame retardant metabolites in urine. *Chemosphere* 181:440-446
- Carignan CC, Fang M, Stapleton HM, Heiger-Bernays W, McClean MD, Webster TF (2016) Urinary biomarkers of flame retardant among collegiate U.S. gymnasts. *Environ Int* 94:362-368
- Carignan CC, McClean MD, Cooper EM, Watkins DJ, Fraser AJ, Heiger-Bernays W, Stapleton HM, Webster TF (2013) Predictors of tris(1,3-dichloro-2-propyl) phosphate metabolite in the urine of office workers. *Environ Int* 55:56-61
- Castorina R, Badman A, Stapleton HM, Butt C, Avery D, Harley KG, Gunier RB, Holland N, Eskenazi B (2017a) Current-use flame retardants: Maternal exposure and neurodevelopment in children of the CHAMACOS cohort. *Chemosphere* 18:574-580
- Castorina R, Butt C, Stapleton HM, Avery D, Harley KG, Holland N, Eskenazi B, Bradman A (2017b) Flame retardants and their metabolites in the homes and urine of pregnant women residing in California (the CHAMACOS cohort). *Chemosphere* 179:159-166
- Ceballos DM, Broadwater K, Page E, Croteau E, La Guardia MJ (2018) Occupational exposure to polybrominated diphenyl ether (PBDEs) and other flame retardant foam additives at gymnastics

studios: Before, during and after the replacement of pit foam with PBDE-free foams. *Environ Int* 116:1-9

Cequier E, Sakhi AK, Marcé RM, Becher G, Thomsen C (2015) Human exposure pathways to organophosphate triesters – A biomonitoring study of mother-child pairs. *Environ Int* 75:159-165

Cequier E, Marce RM, Becher G, Thomsen C (2014) A high-throughput method for determination of metabolites of organophosphate flame retardants in urine by ultra performance liquid chromatography-high resolution mass spectrometry. *Anal Chim Acta* 845: 98-104

Chen Y, Fang J, ren L, Fan R, Zhang J, Liu G, Zhou L, Chen D, Yu Y, Lu S (2018) Urinary metabolites of organophosphate esters in children in South China: Concentrations, profiles and estimated daily intake. *Environ Pollut* 235:358-364

Cooper EM, Covaci A, van Nuijs ALN, Webster TF, Stapleton HM (2011) Analysis of flame retardant metabolites bis(1,3-dichloro-2-propyl) phosphate (BDCPP) and diphenyl phosphate (DPP^o) in urine using liquid chromatography-tandem mass spectrometry. *Anal Bioanal Chem* 401:2123-2132

Council of the District of Columbia (2016) D.C. Law 21-108. Carcinogenic Flame Retardant Prohibition Amendment Act of 2016. <https://code.dccouncil.us/dc/council/laws/21-108.html>

Department of Ecology, State of Washington (2016) Flame retardants concentration limits. <https://ecology.wa.gov/Waste-Toxics/Reducing-toxic-chemicals/Childrens-Safe-Products-Act/Flame-retardant-limits>

Deziel NC, Yi H, Stapleton HM, Huang H, Zhao N, Zhang Y (2018) A case-control study of exposure to organophosphate flame retardants and risk of thyroid cancer in women. *BMC Cancer* 18:637

Ding J, Xu Z, Huang W, Feng L, Yang F (2016) Organophosphate ester flame retardants and plasticizers in human placenta in Eastern China. *Sci Total environ* 554-555: 211-217

Dirtu AC, Van den Eede N, Malarvannan G, Ionas AC, Covaci A (2012) Analytical methods for selected emerging contaminants in human matrices - A review. *Anal Bioanal Chem* 404:2555-2581

Dodson RE, Van den Eede N, Covaci A, Perovich LJ, Brody JG, Rudel RA (2014) Urinary biomonitoring of phosphate flame retardants: Levels in California adults and recommendations for further studies. *Environ Sci Technol* 48:13625-13633

Engvall K, Hult M, Corner R, Lampa E, Norbäck D, Emenius G (2010) A new multiple regression model to identify multi-family houses with a high prevalence of sick building symptoms “SBS”, within the healthy sustainable house study in Stockholm (3H). *Int Arch Occup Environ Health* 83:85-94

European Chemicals Agency (ECHA). (2018). <https://echa.europa.eu>

European Flame Retardants Association (EFRA) (2007) Flame retardants. http://www.bsef-japan.com/index/files/080909_3.pdf

Feng L, Ouyang F, Liu L, Wang X, Wang X, Li YJ, Murtha A, Shen H, Zhang J, Zhang JJ (2017) Levels of urinary metabolites of organophosphate flame retardants, TDCIPP, and TPHP, in pregnant women in Shanghai. *J Environ Public Health* 2016:9416054

- Frederiksen M, Stapleton HM, Vorkamp K, Webster TF, Jensen NM, Sorensen JA, Nielsen F, Knudsen LE, Sorensen LS, Clausen PA, Nielsen JB (2018) Dermal uptake and percutaneous penetration of organophosphate esters in a human skin ex vivo model. *Chemosphere*, in press
- Fromme H, Lahrz T, Kraft M, Fembacher L, Mach C, Dietrich S, Burkardt R, Völkel W, Göen T (2014) Organophosphate flame retardants and plasticizers in the air and dust in German daycare centers and human biomonitoring in visiting children (LUPE 3). *Environ Int* 71:158-163
- Hammel SC, Hoffman K, Webster TF, Anderson KA, Stapleton HM (2016) Measuring personal exposure to organophosphate flame retardants using silicone wristbands and hand wipes. *Environ Sci Technol* 50:4483-4491
- He C, Toms LML, Thai P, Van den Eede N, Wang X, Li Y, Baduel C, Harden FA, Heffernan AL, Hobson P, Cocaci A, Mueller JF (2018a) Urinary metabolites of organophosphate esters: Concentration and age trends in Australian children. *Environ Int* 111:124-130
- He C, English K, Baduel C, Thai P, Jagals P, Ware RS, Li Y, Wang X, Sly PD, Mueller JF (2018b) Concentrations of organophosphate flame retardants and plasticizers in urine from young children in Queensland, Australia and associations with environmental factors. *Environ Res* 164:262-270
- He C, Wang X, Thai P, Baduel C, Gallen C, Banks A, Bainton P, English K, Mueller JF (2018c) Organophosphate and brominated flame retardants in Australian indoor environments: Levels, sources, and preliminary assessment of human exposure. *Environ Pollut* 235:670-679
- Hoffman K, Stapleton HM, Lorenzo A, Butt CM, Adair L, Herring AH, Daniels JL (2018) Prenatal exposure to organophosphates and associations with birthweight and gestational length. *Environ Int* 116:248-254
- Hoffman K, Lorenzo A, Butt CM, Adair L, Herring AH, Stapleton HM, Daniels JL (2017a) Predictors of urinary flame retardant concentration among pregnant women. *Environ Int* 98:96-101
- Hoffman K, Butt CM, Webster TF, Preston EV, Hammel SC, Makey C, Lorenzo AM, Cooper EM, Carignan C, Meeker JD, Hauser R, Soubry A, Murphy SK, Price TM, Hoyo C, Mendelson E, Congleton J, Daniels JL, Stapleton HM (2017b) Temporal trends in exposure to organophosphate flame retardants in the United States. *Environ Sci Technol Lett* 4:112-117
- Hoffman K, Lorenzo A, Butt CM, Hammel SC, Henderson BB, Roman SA, Scheri RP, Stapleton HM, Sosa JA (2017c). Exposure to flame retardant chemicals and occurrence and severity of papillary thyroid cancer: A case-control study. *Environ Int* 107:235-242
- Hoffman K, Butt CM, Chen A, Limkakeng AT, Stapleton HM (2015a) High exposure to organophosphate flame retardants in infants: associations with baby products. *Environ Sci Technol* 49:14554-14559
- Hoffman K, Garantziotis S, Birnbaum LS, Stapleton HM (2015b) Monitoring indoor exposure to organophosphate flame retardants: Hand wipes and house dust. *Environ Health Perspect* 123: 160-165
- Hoffman K, Daniels JL, Stapleton HM (2014) Urinary metabolites of organophosphate flame retardants and their variability in pregnant women. *Environ Int* 63:169-172
- Hou R, Xu Y, Wang Z (2016) Review of OFPRs in animals and humans: Absorption, bioaccumulation, metabolism, and internal exposure research. *Chemosphere* 153:78-90

- Ingle ME, Minguez-Alarcon L, Carignan CC, Butt CM, Stapleton HM, Williams PL, Ford JB, Hauser R, Meeker JD (2018) The association between urinary concentrations of phosphorous-containing flame retardant metabolites and semen parameters among men from a fertility clinic. *Int J Hyg Environ Health* *in press*
- Jayatilaka NK, Restrepo P, Williams LT, Ospina M, Valentin-Blasini L, Calafat AM (2017) Quantification of three chlorinated dialkyl phosphates, diphenyl phosphate, 2,3,4,5-tetrabromobenzoic acid, and four other organophosphates in human urine by solid phase extraction-high performance liquid chromatography-tandem mass spectrometry. *Anal Bioanal Chem* 409:1323-1332
- Johnson D, Carter MD, Crow BS, Isenberg SL, Graham LA, Erol HA, Watson CM, Pantazides BG, van der Schans MJ, Langenberg JP, Noort D, Blake TA, Thomas JD, Johnson RC (2015) Quantitation of ortho-cresyl phosphate adducts to butyrylcholinesterase in human serum by immunomagnetic-UHPLC-MS/MS. *J Mass Spectrom* 50:683-692
- Kanazawa A, Saito I, Araki A, Takeda M, Ma M, Saijo Y, Kishi R (2010) Association between indoor exposure to semi-volatile organic compounds and building-related symptoms among the occupants of residential dwellings. *Indoor air* 20:72-84
- Kim JW, Isobe T, Muto M, Tue NM, Katsura K, Malarvannan G, Sudaryanto A, Chang KH, Prudente M, Viet PH, Takahashi S, Tanabe S (2014) Organophosphorus flame retardants (PFRs) in human breast milk from several Asian countries. *Chemosphere* 116:91-97
- Kosarac I, Kubwado C, Foster WG (2016) Quantitative determination of nine urinary metabolites of organophosphate flame retardants using solid phase extraction and ultra performance liquid chromatography coupled to tandem mass spectrometry (UPLC-MS/MS)
- Kucharska A, Cequier E, Thomsen C, Becher G, Covaci A, Voorspoels S (2015a) Assessment of human hair as an indicator of exposure to organophosphate flame retardants. Case study on Norwegian mother-child cohort. *Environ Int* 83:50-57
- Kucharska A, Covaci A, Vanermen G, Voorspoels S (2015b) Non-invasive biomonitoring for PFRs and PBDEs: New insights in analysis of human hair externally exposed to selected flame retardants. *Sci Total Environ* 505:1062-1071
- Kucharska A, Covaci A, Vanermen G, Voorspoels S (2014) Development of a broad spectrum method for measuring flame retardants – Overcoming the challenges of non-invasive human biomonitoring studies. *Anal Bioanal Chem* 406:6665-6675
- Li P, Jin J, Wang Y, Hu J, Xu M, Sun Y, Ma Y (2017) Concentrations of organophosphorus, polybromobenzene, and polybrominated diphenyl ether flame retardants in human serum, and relationships between concentrations and donor age. *Chemosphere* 171:654-660.
- Lipscom ST, McClelland MM, MacDonald M, Cardenas A, Anderson KA, Kile ML (2017) Cross-sectional study of social behaviors in preschool children and exposure to flame retardants. *Environ Health* 16:23
- Liu X, Yu G, Cao Z, Wang B, Huang J, Deng S, Wang Y (2017) Occurrence of organophosphorus flame retardants on skin wipes: Insight into human exposure from dermal absorption. *Environ Int* 98:113-119

- Liu LY, He K, Hites RA, Salamova A (2016) Hair and nails as noninvasive biomarkers of human exposure to brominated and organophosphate flame retardants. *Environ Sci Technol* 50:3065-3073
- Liu LY, Salamova A, He K, Hites RA (2015) Analysis of polybrominated diphenyl ethers and emerging halogenated and organophosphate flame retardants in human hair and nails. *J Chromatogr A* 1406:251-257
- Lu SY, Li YX, Zhang T, Cai D, Ruan JJ, Huang MZ, Wang L, Zhang JQ, Qiu RL (2017) Effect of E-waste recycling on urinary metabolites of organophosphate flame retardants and plasticizers and their association with oxidative stress. *Environ Sci Technol* 51:2427-2437
- Ma Y, Jin J, Li P, Xu M, Sun Y, Wang Y, Yuan H (2017) Organophosphate ester flame retardants concentrations and distribution in serum from inhabitants of Shandong, China, and changes between 2011 and 2015. *Environ Toxicol Chem* 36:414-421
- Mäkinen MS, Makinen MR, Koistinen JT, Pasanen AL, Pasanen PO, Kalliokoski PJ, et al. (2009) Respiratory and dermal exposure to organophosphorous flame retardants and tetrabromobisphenol a at five work environments. *Environ Sci Technol* 43:941-947
- Martin J, Möder M, Gaudl A, Alonso E, Reemtsma T (2015) Multi-class method for biomonitoring of hair samples using gas chromatography-mass spectrometry. *Anal Bioanal Chem* 407:8725-8734
- Meeker JD, Cooper EM, Stapleton HM, Hauser R (2013a) Exploratory analysis of urinary metabolites of phosphorous-containing flame retardants in relation to markers of male reproductive health. *Endocr Disruptors* 1:e26306
- Meeker JD, Cooper EM, Stapleton HM, Hauser R (2013b) Urinary metabolites of organophosphate flame retardants: Temporal variability and correlations with house dust concentrations. *Environ Health Perspect* 121:580-585
- Meeker JD, Stapleton HM (2010) House dust concentrations of organophosphate flame retardants in relation to hormone levels and semen quality parameters. *Environ Health Perspect* 118:318-323
- Meeker JD, Godfrey-Bailey L, Hauser R (2007) Relationships between serum hormone levels and semen quality among men from an infertility clinic. *J Androl* 28:397-406
- Mendelsohn E, Hagopian A, Hoffman K, Butt CM, Lorenzo A, Congleton J, Webster TF, Stapleton HM (2016) Nail polish as a source of exposure to triphenyl phosphate. *Environ Int* 86:45-51
- Ministry of Environment and Food of Denmark. Environmental Protection Agency (2016) Environmental and health screening profiles of phosphorous flame retardants. A LOUS follow-up project. Environmental project No. 1823, 2016.
- Muenhor D, Moon HB, Lee S, Goosey E (2017) Organophosphorus flame retardants (PFRs) and phthalates in floor and road dust from a manual e-waste dismantling facility and adjacent communities in Thailand. *J Environ Sci Health A Tox Hazard Subst Environ Eng* 53: 79-90
- Nordic Expert Group for Criteria Documentation of Health Risks from Chemicals (NEG) (2009) 143. Phosphate triesters with flame retardants properties. NR 2010;44(6). ISBN 978-91-85971-23-7
- Office of Environmental Health Hazard (OEHHA). State of California. Assessment Safe Drinking Water and Toxic Enforcement Act of 1986. Chemicals known to the State to cause cancer or

reproductive toxicity. December 29, 2017 https://oehha.ca.gov/media/downloads/proposition-65/p65122917_0.pdf

Ospina M, Jayatilaka NK, Wong LY, Restrepo P, Calafat AM (2018) Exposure to organophosphate flame retardant chemicals in the U.S. general population: Data from the 2013-2014 National Health Nutrition Examination Survey. *Environ Int* 110:32-41

Petroianu GA (2016) Neuropathic organophosphates: from Scrugham, Heim and Lorot to Jake leg paralysis. *Pharmazie* 71:738-744

Petropoulou SS, Petreas M, Park JS (2016) Analytical methodology using ion-pair liquid chromatography-tandem mass spectrometry for the determination of four di-ester metabolites or organophosphate flame retardants in California human urine. *J Chromator A* 1434:70-80

Phillips AL, Hammel SC, Hoffman K, Lorenzo K, Chen AM, Webster TF, Stapleton HM (2018) Children's residential exposure to organophosphate ester flame retardants and plasticizers: Investigating exposure pathways in the TESIE study. *Environ Int* 116:176-185

Poma G, Sales C, Bruyland B, Christia C, Goscinnny S, Van Loco J, Covaci A (2018) Occurrence of organophosphorus flame retardants and plasticizers (PFRs) in Belgian foodstuffs and estimation of dietary exposure of the adult population. *Environ Sci Technool* 52:2331-2338

Poma G, Glynn A, Malarvannan G, Covaci A, Darnelund PO (2017) Dietary intake of phosphorus flame retardants (PFRs) using Swedish food market basket estimations. *Food Chem Toxicol* 100:1-7

Preston EV, McClean MD, Henn BC, Stapleton HM, Braverman LE, Pearce EN, Makey CM, Webster TF (2017) Associations between urinary diphenyl phosphate and thyroid function. *Environ Int* 101:158-164

Qiao L, ZhengXB, Zheng J, Lei WX, Li HF, Wang MH, He CT, Chen SJ, Yuan JG, Luo XJ, Yu YJ, Yang ZY, Mai BX (2016) analysis of human hair to assess exposure to organophosphate flame retardants: Influence of hair segments and gender differences. *Environ Res* 148:177-183

Reemtsma T, Lingott J, Roegler S (2011) Determination of 14 monoalkyl phosphates, dialkyl phosphates and dialkyl thiophosphates by LC-MS/MS in human urinary samples. *Sci Total Environ* 409:1990-1993.

Romano ME, Hawley NL, Eliot M, Calafat AM, Jayatilaka NK, Kelsey K, McGarvey S, Phipps MG, Savitz DA, Werner EF, Braun JM (2017) Variability and predictors of urinary concentrations of organophosphate flame retardant metabolites among pregnant women in Rhode Island. *Environ Health* 16:40

Shen B, Whitehead TP, Gill R, Dhaliwal J, Brown FR, Petreas M, PattonS, Hammond SK (2018) Organophosphate flame retardants in dust collected from United States fire stations. *Environ Int* 112:41-48

Schindler BK, Koslitz S, Weiss T, Broding HC, Brüning T, Bünger J (2014) Exposure of aircraft maintenance technicians to organophosphates from hydraulic fluids and turbine oils: A pilot study. *Int J Hyg Environ Health* 217:34-37

- Schindler BK, Weiss T, Schütze A, Koslitz S, Broding HC, Bünger J, Brüning T (2013) Occupational exposure of air crews to tricresyl phosphate isomers and organophosphate flame retardants after fume events. *Arch Toxicol* 87:645-648
- Schindler BK, Förster K, Angerer J (2009a) Determination of human urinary organophosphate flame retardant metabolites by solid-phase extraction and gas chromatography-tandem mass spectrometry. *J Chromatogr B* 877:375-381
- Schindler BK, Förster K, Angerer J (2009b) Quantification of two urinary metabolites of organophosphorus flame retardants by solid-phase extraction and gas chromatography-tandem mass spectrometry. *Arch Toxicol* 395:1167-1171
- Schopfer LM, Masson P, Lamourette P, Simon S, Lockridge O (2014) Detection of cresyl phosphate-modified butyrylcholinesterase in human plasma for chemical exposure associated with aerotoxic syndrome. *Anal Biochem* 461:17-26
- Schreder ED, Uding N, La Guardia MJ (2016) Inhalation a significant exposure route for chlorinated organophosphate flame retardants. *Chemosphere* 150:499-504
- Shen B, Whitehead TP, Gill R, Dhaliwal J, Brown FR, Petreas M, Patton S, Hammond SK (2018) Organophosphate flame retardants in dust collected from United States fire stations. *Environ Int* 112:41-48
- Soubry A, Hoyo C, Butt CM, Fieuws S, Price TM, Murphy SK, Stapleton HM (2017) Human exposure to flame-retardants is associated with aberrant DNA methylation at imprinted genes in sperm. *Environ Epigenet* 3:dvx003
- Su G, Lechter RJ, Yu H, Gooden DM, Stapleton HM (2016) Determination of glucuronide conjugates of hydroxyl triphenyl phosphate (OH-TPHP) metabolites in human urine and its use as a biomarker of TPHP exposure. *Chemosphere* 149:314-319
- Su G, Letcher RJ, Yu H (2015) Determination of organophosphate diesters in urine samples by a high-sensitivity method based on ultra high pressure liquid chromatography-triple quadrupole-mass spectrometry. *J Chromatogr A* 1426:154-160
- Sun Y, Gong X, Lin W, Wang Y, Wu M, Kannan K, Ma J (2018) Metabolites of organophosphate ester flame retardants in urine from Shanghai, China. *Environ Res* 164:507-515
- Sundkvist AM, Olofsson U, Haglund P (2010) Organophosphorus flame retardants and plasticizers in marine and fresh water biota and in human milk. *J Environ Monit* 12:943-951
- Tacal O, Schopfer LM (2014) Healthy F-16 pilots show no evidence of exposure to tri-ortho-cresyl phosphate through the on-board oxygen generating system. *Chem Biol Interact* 215:69-74
- Tao Y, Shang Y, Li J, Feng J, He Z, Covaci A, Wang P, Luo J, Mao X, Shi B, Hu L, Luo D, Mei S (2018) Exposure to organophosphate flame retardants of hotel room attendants in Wuhan City, China. *Environ Pollut* 236:626-633
- Thomas MB, Stapleton HM, Dills RL, Violette HD, Christakis DA, Sathyanarayana S (2017) Demographic and dietary risk factors in relation to urinary metabolites of organophosphate flame retardants in toddlers. *Chemosphere* 185: 918-925

- U.S. Environmental Protection Agency (US EPA) (2015) Flame retardants used in flexible polyurethane foams: an alternatives assessment update.
https://www.epa.gov/sites/production/files/2015-08/documents/ffr_final.pdf
- U.S. Environmental Protection Agency (US EPA) (2017) Regional Screening Level (RSL) Summary Table (TR=1E-06, HQ=0.1) November 2017. <https://semspub.epa.gov/work/HQ/197027.pdf>
- Van den Eede N, Tomy G, Tao F, Halldorson T, Harrad S, Neels H, Covaci A (2016a) Kinetics of tris (1-chloro-2-propyl) phosphate (TCIPP) metabolism in human liver microsomes and serum. *Chemosphere* 144:1299-1305
- Van den Eede N, Ballesteros-Gomez A, Neels H, Covaci A (2016b) Does biotransformation of aryl phosphate flame retardants in blood cast a new perspective on their debated biomarkers. *Environ Sci Technol* 50:12439-12445
- Van den Eede N, De Meester I, Maho W, Neels H, Covaci A (2016c) Biotransformation of three flame retardants and plasticizers in primary human hepatocytes: untargeted metabolite screening and quantitative assessment. *J Appl Toxicol* 36:1401-1408
- Van den Eede N, Erratico C, Exarchou V, Maho W, Neels H, Covaci A (2015a) In vitro biotransformation of tris(2-butoxyethyl) phosphate (TBOEP) in human liver and serum. *Toxicol Appl Pharmacol* 284:246-253
- Van den Eede N, Hefferman AL, Aylward LL, Hobson P, Neels H, Mueller JF, Covaci A (2015b) Age as a determinant of phosphate flame retardant exposure of the Australian population and identification of novel urinary PFR metabolites. *Environ Int* 74:1-8
- Van den Eede N, Maho W, Erratico C, Neels H, Covaci A (2013a) First insights in the metabolism of phosphate flame retardants and plasticizers using human liver fractions. *Toxicol Lett* 223:9-15
- Van den Eede N, Neels H, Jorens PG, Covaci A (2013b) Analysis of organophosphate flame retardant diester metabolites in human urine by liquid chromatography electrospray ionization tandem mass spectrometry. *J Chromator A* 1303: 48-53
- Van der Veen I, de Boer J (2012) Phosphorous flame retardants: Properties, production, environmental occurrence, toxicity and analysis. *Chemosphere* 88:1119-1153
- Vermont General Assembly (2013) The Vermont Statutes Online. Title 9: Commerce And Trade. Chapter 80: Flame Retardants. § 2971. Repealed. 2013, No. 85, § 1 to § 2980. Department of Health rulemaking; TCPP. <https://legislature.vermont.gov/statutes/fullchapter/09/080>
- Völkel W, Fuchs V, Wöckner M, Fromme H (2018) Toxicokinetic of tris(2-butoxyethyl) phosphate (TBOEP) in humans following single oral administration. *Arch Toxicol* 92:651–660
- Wei GL, LI DQ, Zhuo MN, Liao YS, Xie ZY, Guo TL, Li JJ, Zhang SY, Liang ZQ (2015) Organophosphorus flame retardants and plasticizers: Sources, occurrence, toxicity and human exposure. *Environ Pollut* 196:29-46
- WeissT, Schindler BK, Schütze A, Koslitz S, Broding HC, Bünger J, Brüning T (2015) Reply to the letter of Anderson J entitled “Comment on Schindler, BK; Weiss, T; Schütze, A; et al. Occupational

exposure of air crews to tricresyl phosphate isomers and organophosphate flame retardants after fume events. *Arch Toxicol* (2013) 87:645-648". *Arch Toxicol* 89:263-264

Wu M, Yu G, Cao Z, Wu D, Liu K, Deng S, Huang J, Wang B, Wang Y (2016) Characterization and human exposure assessment of organophosphate flame retardants in indoor dust from several microenvironments of Beijing, China. *Chemosphere* 150:465-471

Xu F, Tay JH, Covaci A, Padilla-Sanchez JA, Papadopoulou E, Smastuen L, Neels H, Sellström U, de Wit CA (2017) Assessment of dietary exposure to organohalogen contaminants, legacy and emerging flame retardants in a Norwegian cohort. *Environ Int* 102:236-243

Xu F, Giovanoulis G, van Waes S, Padilla-Sanchez JA, Papadopoulou E, Magner J, Haug LS, Neels H, Covaci A (2016) Comprehensive study of human external exposure to organophosphate flame retardants via air, dust, and hand wipes: the importance of sampling and assessment strategy. *Environ Sci Technol* 50:7752-7760

Yan X, Zheng X, Wang M, Zheng J, Xu R, Zhuang X, Lin Y, Ren M (2018) Urinary metabolites of phosphate flame retardants in workers occupied with e-waste recycling and incineration. DOI: 10.1016/j.chemosphere.2018.02.148

Yoshida T, Yoshida J (2012) Simultaneous analytical method for urinary metabolites of organophosphorus compounds and moth repellents in general population. *J Chromatogr B* 880: 66-73

Zhao F, Chen M, Gao F, Shen H, Hu J (2017) Organophosphorus flame retardants in pregnant women and their transfer to chorionic villi. *Environ Sci Technol* 51:6489-6497

Zhao F, Wan Y, Zhao H, Hu W, Mu D, Webster TF, Hu J (2016) Levels of blood organophosphorus flame retardants an association with changes in human sphingolipid homeostasis. *Environ Sci Technol* 50:8896-8903

Zhang X, Zou W, Mu L, Chen Y, Ren C, Hu X, Zhou Q (2016) Rice ingestion is a major pathway for human exposure to organophosphate flame retardants (OPFRs) in China. *J Hazard Mater* 318:686-693

Zheng X, Qiao L, Covaci A, Sun R, Guo H, Zheng J, Luo X, Xie Q, Mai B (2017) Brominated and phosphate flame retardants (FRs) in indoor dust from different microenvironments: Implications for human exposure via dust ingestion and dermal contact. *Chemosphere* 184:185-191

Zheng X, Xu F, Luo X, Mai B, Covaci A (2016) Phosphate flame retardants and novel brominated flame retardants in home-produced eggs from e-waste recycling region in China. *Chemosphere* 150:545-550

Zhou L, Hiltcher M, Püttmann W (2017) Occurrence and human exposure assessment of organophosphate flame retardants in indoor dust from various microenvironments of Rhine/Main region, Germany. *Indoor Air* 27:1113-1127

Table 1. A summary of recent epidemiological studies on the potential health effects of PEFRs

Author, location	Subjects and study period	Sampling	Exposure assessment ^a	Health effects related to PEFRs exposure
Hormonal and reproductive effects				
Carignan et al., 2018a USA	201 couples from the Environment and Reproductive Health (EARTH) prospective cohort study ^a 2005-2015	One or two spot urine samples per <i>in vitro</i> fertilization cycle	Urinary metabolites in males: BCIP, BDCIPP, DPHP, ip-PPP, <i>tert</i> butyl-phenylphenyl phosphate (tb-PPP)	<i>Paternal exposure and partner's pregnancy outcome</i> Association between paternal preconception BDCIPP levels and reduced probability of oocyte fertilization. No association between PFR metabolites and the proportion of cycles resulting in implantation, clinical pregnancy and live birth.
Carignan et al., 2018b USA	211 women from the Environment and Reproductive Health (EARTH) prospective cohort study ^a 2005-2015	One or two spot urine samples per <i>in vitro</i> fertilization cycle	Urinary metabolites in females: BCIP, BDCIPP, DPHP, ip-PPP, tb-PPP	<i>Maternal exposure and pregnancy outcome</i> Association between the levels of two individual metabolites (i.e. DPHP and tb-PPP) and of total metabolites, and reduced probability of successful fertilization, implantation, clinical pregnancy, and live birth.
Ingle et al., 2018 USA	220 men from the Environment and Reproductive Health (EARTH) cohort study ^a 2005-2015	One to five urine and sperm samples	Urinary metabolites in males: BCIP, BDCIPP, DPHP, ip-PPP, tb-PPP	<i>Exposure in men and semen parameters</i> No consistent association between individual metabolites and semen parameters.
Preston et al., 2017 USA	26 men and 25 women as a part of the Flame Retardant Exposure Study (FlaRE) 2010-2011	Spot urine and blood samples at 1, 6 and 12 months	Urinary DPHP	<i>Adult exposure to TPHP and circulating thyroid hormones</i> No association between DPHP levels and thyroxine (free T4), triiodothyroxine (free and total T3) or thyroid stimulating hormone (TSH) concentrations in serum. Association between DPHP levels and increased total T4, especially in women.
Soubry et al., 2017 USA	67 men as a part of the Gametic Epigenetic Reprogramming (TIEGER) cross-sectional study 2012-2013	Spot urine and sperm samples on the same day	Urinary metabolites: BCIP, BDCIPP, DPHP, ip-PPP, tb-PPP	<i>Exposure in men and DNA methylation at imprinted genes in sperm</i> Association between PEFRs metabolites (i.e.BDCIPP, DPHP, ip-PPP) and small methylation differences (hyper- or hypo-methylation of different genes specific to the metabolites).
Meeker et al., 2013a USA	33 men from couples who were infertile due to a male factor, a female factor, or both	One urine, blood, and semen sample	Urinary BDCIPP and DPHP	<i>Exposure in men, and semen parameter, and reproductive and thyroid hormones</i> Association between BDCIPP levels and decreases in sperm quality parameters, and concentrations of total T3 and FSH in serum.

	(Subset of a parent study: Meeker et al., 2007) 2003-2004			Association between DPHP levels and decreased sperm concentration (no significant changes in hormones, e.g., prolactin).
Meeker et al., 2010 (Meeker et al., 2007) USA	50 men from couples who were infertile due to a male factor, a female factor, or both (Subset of a parent study: Meeker et al., 2007)	Dust collected at the homes of men who participated in the parent study in 2002-2007. One blood and semen sample (38 samples for semen)	TDCPP and TPHP in dust	<i>Exposure in men and semen parameters and reproductive and thyroid hormones</i> Association between TDCPP levels and changes in serum concentrations of hormones (decrease in free T4 and increase in prolactin). Association between TPHP and increased prolactin and decreased sperm concentration.
Effects at birth and in childhood				
Hoffman et al., 2018 USA	349 mothers and their child from the cohort of the Pregnancy Infection and Nutrition study (PIN) 2002-2005	Single spot urine samples from the mothers during late-second or early-third trimester (24-30 weeks of pregnancy)	Urinary metabolites: BCIP, BDCIPP, DPHP, ip-PPP, tb-PPP, 1-hydroxy-2-propyl phosphate (BCIPH-IPP)	<i>Maternal exposure and birth outcomes</i> Higher levels of BDCIPP and ip-PPP associated with decreased gestational duration and increased preterm births (<37 weeks gestation) among female infants.
Castorina et al., 2017a, USA	310 mothers and their 7-year-old children from a longitudinal birth cohort study (Center for the Health Assessment of Mothers and Children of Salinas-CHAMACOS) 1999-2000	Single spot urine samples from the mothers at 26 weeks of pregnancy	Urinary metabolites: BCIP, BDCIPP, DPHP, ip-PPP, tb-PP (Castorina et al., 2017b)	<i>In utero exposure and neurodevelopmental outcome (associations analyzed: total PEFR metabolites, TDCIPP, TPHP, ip-PPP)</i> Association between DPHP and total PEFR metabolites levels, and decreased cognitive function (full scale IQ and working memory). No association between prenatal BDCIPP and ip-PPP levels and neurobehavioral development.
Lipscom et al., 2017 USA	72 children (aged 3-5 years) 2012-2013	Passive wristband samplers worn continuously for 7 days	FRs in wristbands, including brominated diphenyl ethers (BDEs) and total PEFRs	<i>Child exposure and social behavior</i> Cross-sectional association between total PEFRs levels and poorer social skills in a few domains (e.g., externalizing behavior).
Respiratory outcomes and immunotoxicity				
Sun et al., 2018 China	180 participants (130 adults, 27 students, and 33 children) 2016-2017	Single spot urine sample	9 urinary metabolites including BCEP, BCIPP, BDCIPP, DNBP, DPHP, BBOEP, BEHP	<i>Indoor exposure and allergy</i> Association between DNBP levels and self-reported symptoms of allergy.
Canbaz et al., 2016	110 children who developed	Dust collected from the	PBDEs and PEFRs in	<i>FRs in mother's mattress dust and the development of childhood</i>

Sweeden	asthma at 4 or at 8 years, matched with 110 controls from a large prospective study 1994-1996	mother's mattress two months after child birth f	dust, including TECP, TCIP, TDCIPP, TPHP, TBOEP, EHDPHP, mmp-TCP	<i>asthma</i> No association between higher concentrations of PEFRs and the development of childhood asthma.
Araki et al., 2014 Japan	516 inhabitants (adults and children) in 156 different homes Cross-sectional study. 2004-2006	Indoor floor and multi-surface dust collected in each family's home in 2004, 2005, and 2006	11 PEFRS such as TBOEP, TCIPP, TDCIPP, TPHP (most frequently detected) in dust	<i>Indoor exposure and asthma and allergy</i> Association between TNBP levels and an inhabitant's medical treatment for asthma and allergic rhinitis. Association between TCIPP and TDCIPP levels and recent medical treatment of an inhabitant for atopic dermatitis.
Bergh et al., 2011 Sweden	Adults (men and women). Part of a larger study, the Healthy Sustainable Houses study in Stockholm (3H). Frequency of SHS studied in 481 multi-family buildings with 10506 dwellings (Engvall et al., 2010). 2006	Indoor air from 169 apartments in buildings with a high incidence of reported SHS (2-4 apartment/building)	Phthalates and 15 organophosphorous flame retardants	<i>Indoor exposure and sick house syndrome (SHS) (i.e. irritation of the eyes, nose, throat, skin and coughing)</i> No association between PEFR levels and reported SHS symptoms, compared to buildings with low prevalence of SHS.
Kanazawa et al., 2010 Japan	134 inhabitants (64 men and 70 women) of 41 dwellings Cross-sectional study 2006-2007	Indoor air and dust (surface, floor) from the dwelling	Semi-volatile organic compounds including 11 PEFRs	<i>Indoor exposure and sick house syndrome (SHS)</i> Association between TBNP levels (floor dust) and reported mucosal symptoms of SHS. Inverse association between TBEP concentrations (floor dust) and reported mucosal symptoms of SHS.
Others				
Deziel et al., 2018 USA	100 cases and 100 controls Cases: Patients newly diagnosed with papillary thyroid cancer (PTC) (women) 2010-2013	Single spot urine samples	6 urinary metabolites: BCIPP, BCIHPP, BDCIPP, ip-PPP, DPHP and tert-butyl phenyl phenyl phosphate (tb-PPP)	<i>Adult exposure and PTC</i> No association between urinary PEFR metabolites concentrations measured at the time of diagnosis and risk of PTC. Tb-PPP was only detected in 6% of samples and was therefore excluded for analysis.
Hoffman et al., 2017c USA	70 cases and 70 controls Cases: Patients newly diagnosed with papillary thyroid cancer (PTC) (men and women)	Dust collected at each participant's home	FRs in household dust, including BDEs, TCEP, TCIPP, TDCIPP and TPHP	<i>Adult exposure and PTC</i> Higher levels of TCEP associated with increased odds of PTC, especially larger and more aggressive tumors.

	2014-2016			
Lu et al., 2017 China	221 adults and children 2014	Single spot urine samples	8 urinary metabolites:	<i>Adult exposure and oxidative stress (8-OHdG in urine)</i> Association between PEFR metabolite levels (i.e. DCEP, DNBP, DPHP) and a higher concentration of 8-OHdG, in e-waste dismantling sites.
Zhao et al., 2016 China	154 men and 101 women 2012	One blood sample	TCIPP, TBOEP, TPHP, TEP, TNBP, EHDPP in blood	<i>Adult exposure and changes in sphingolipid homeostasis</i> Association between levels of the six PEFRs and increased sphingomyelin concentration. Negative association between EHDPP, TPHP, and TNBP levels and sphingosine 1-phosphate concentration.

^a Participants originated from couples whose infertility diagnosis was either male factor, female factor, or a combination of both.

Table 2 Parent compounds and metabolites

Parent PEFR		PEFR metabolite	
Full name (CAS number)	Abbreviation	Full name	Abbreviation
Halogenated organophosphorous compounds			
Tris(2-chloroethyl) phosphate (115-96-8)	TCEP	Bis(2-chloroethyl) phosphate	BCEP
Tris(1-chloro-2-propyl) phosphate (13674-84-5)	TCIPP	Bis(1-chloro-2-propyl) phosphate	BCIPP
		Bis(1-chloro-2-propyl) 1-hydroxy-2-propyl phosphate	BCIPHPP
Tris(1,3-dichloro-2-propyl) phosphate (or isopropyl) (13674-87-8)	TDCIPP	Bis(1,3-dichloro-2-propyl) phosphate	BDCIPP
Non-halogenated organophosphorous compounds			
Tri-n-butyl phosphate (126-73-8)	TNBP	Di-n-butyl phosphate	DNBP
Tris(2-ethylhexyl) phosphate (78-42-2)	TEHP	Bis(2-ethylhexyl) phosphate	BEHP
Mono-substituted isopropyl triphenyl phosphate (Isopropylphenyl diphenyl phosphate) (several isomers : 55864-04-5, 69515-46-4, 64532-94-1)	Mono-ITP	Isopropylphenyl phenyl phosphate	ip-PPP
Tris(2-butoxyethyl) phosphate (78-51-3)	TBOEP	Bis(2-butoxyethyl) phosphate	BBOEP
		Bis(2-butoxyethyl)-(2-hydroxyethyl) phosphate	BBOEHEP
Triphenyl phosphate (115-86-6)	TPHP	Diphenyl phosphate	DPHP
Tricresyl phosphate (1330-78-5)	TCP	Dicresyl phosphate	DCP
<i>Ortho, meta, and para isomers (78-30-8, 563-04-2, 78-32-0, respectively)</i>	ToCP, TmCP, TpCP		
2-ethylhexyl diphenyl phosphate (1241-94-7)	EHDPP	5-hydroxy-2-ethylhexyl diphenyl phosphate	5-OH-EHDPP

Chemical structures of PEFR metabolites are given in Supplementary material S1

The commercial mixtures TCP, TCIPP, and triisopropylated phenyl phosphate contain varying amounts of their isomers, e.g., the most abundant isomer in commercial products of TCIPP is generally the completely branched isomer, CAS: 13674-84-5.

Table 3. Urinary concentrations of the principal metabolites of PEFRs in general populations (µg/l).50th percentile (max) % ≥ limit of detection (LOD)

Author, location	Sampling year	N (population)	BCEP	BCIPP	BCIPHP	BDCIPP	DNBP	ip-PPP	BBOEP	DPHP
<i>Australia</i>										
He et al., 2018a	2014-2015	Children (0-5 years) Pooled urine samples (20 children/pool, 20 pools) Not adjusted for specific gravity	<0.01 ^a (0.036) 15	0.85 ^a (3.2) 100	0.43 ^a (2.1) 100	2.6 ^a (19) 100	0.18 ^a (0.55) 100	-	0.32 ^a (0.78) 100	25 ^a (58) 100
He et al., 2018b	2015-2016	51 children (3-29 months, average 13 months). Two urine samples from two consecutive days. Not adjusted for specific gravity	<0.01 (n.r.) 33	<0.68 (n.r.) 86	0.93 (n.r.) 96	3.3 (n.r.) 100			0.10 (n.r.) 75	1 (n.r.) 94
Van den Eede et al., 2015b	2010-2011	Adults and children Pooled urine samples Not adjusted for specific gravity Campaign 1: 28 pools of 7 individuals and 44 pools of 7 individuals	-	-	n.r. (9.43) 100	n.r. (8.90) 92	n.r. (2.15) 18	-	n.r. (0.53) 6	n.r. (727) 97
	2012-2013	Campaign 2: 23 pools of 100 individuals	-	-	n.r. (7.17) 100	n.r. (3.41) 96	n.r. (0.94) 4	-	n.a. 0	n.r. (225) 100
<i>Asia</i>										
Chen et al., 2018 <i>China</i>	2015	411 children (212 aged 8-12 years and 199 aged 6-14 years) First morning void Specific gravity adjusted	1.04 (86.9) 91	0.15 (3.11) 66	-	0.05 (4.73) 29	0.12 (2.67) 77	-	0.05 (0.37) 84	0.28 (6.18) 99
Sun et al., 2018 <i>China, Shanghai</i>	2016-2017	180 participants (130 adults, 27 students, and 33 children) First morning void Not adjusted for specific gravity	n.r. (22.60) 5	n.r. (8.83) 16.7		n.r. (2.09) 21.1	0.008 (1.48) 51.7		0.097 (2.19) 68.3	0.066 (4.0) 67.8
Feng et al., 2017 <i>China, Shanghai</i>	2015	23 pregnant women Spot urine Specific gravity adjusted	-	-	-	1.58 (2.2) 17	-	-	-	0.83 (7.3) 100
Lu et al., 2017 <i>China</i>	2014	221 adults and children First morning void Not adjusted for specific gravity	1.1 (57) 71	0.097(23) 56	-	0.11 (4.5) 76	0.15 (7.8) 99	-	0.071 (2.1) 93	0.53 (36) 100
Yoshida T et al., 2012 <i>Japan</i>	n.r.	5 individuals (16-48 years) Spot urine Not adjusted for specific gravity	-	-	-	-	<LOQ (<LOQ) 40	-	-	n.a. (9.8) 20
<i>Europe</i>										

Larsson et al. 2018 <i>Sweden</i>	2015	113 children (4 years) First morning void								1.8 (35) 100
Völkel et al., 2018 <i>Germany</i>	2011- 2012	Children (20-80 months) 54 urine samples Spot urine Not adjusted for specific gravity							0.16 (n.r.) 80	
Cequier et al., 2015 <i>Norway</i>	2012	48 mothers 2 to 8 samples over a period of 24h (244 samples) Specific gravity adjusted Same population as Cequier et al., 2014	-	-	-	0.08 (2.1) 52	<MDL (0.35) 8	-	<MDL (0.27) < 1	0.63 (60) 97
		54 paired children (6-12 years, median 10 years) 2 or 3 samples over a period of 24h (112 samples) Specific gravity adjusted	-	-	-	0.23 (3.3) 61	<MDL (1.0) 15	-	<MDL (1.0) 32	1.0 (129) 97
Cequier et al., 2014	n.r.	42 mothers Spot urine Specific gravity adjusted	-	-	-	<LOQ (n.r.) 57	<LOQ (n.r.) 5	-	n.a. 0	n.r. (n.r.) 100
		42 paired children Spot urine Specific gravity adjusted	-	-	-	n.r. (n.r.) 79	<LOQ (n.r.) 14	-	<LOQ (n.r.) 33	n.r. (37) 100
Fromme et al., 2014 <i>Germany</i>	2011- 2012	312 children (22-80 months, mean 54 months) Spot urine Not adjusted for specific gravity	0.2 (13.1) 65	<0.2 (8.4) 21	-	-	0.2 (6.6) 71	-	2.0 (24.9) 90	0.8 (23.2) 91
Van den Eede et al., 2013b <i>Belgium</i>	n.r.	59 adults (23 men and 36 women) Spot urine Not adjusted for specific gravity	n.r.(9.5) 27	n.r (6.2) 3	-	n.r (3.5) 25	n.r (3.5) 5	-	n.r (7.0) 31	n.r (13) 93
Reemstma et al., 2011 <i>Germany</i>	n.r.	19 urine samples from males and females (14-85 years) Spot urine Not adjusted for specific gravity	-	-	-	-	-	-	-	1.3 (n.r) n.r.
North America										
Carignan et al., 2018a <i>USA</i> <i>Massachusetts</i>	2005 to 2015	201 men (whose partners were undergoing in vitro fertilization) Spot urine (1 sample) Specific gravity adjusted	-	n.a. 0	-	0.46 (12.39) 84	-	0.21 (3.60) 76	-	0.57 (8.54) 87
Carignan et al., 2018b <i>USA</i> <i>Massachusetts</i>	2005 to 2015	211 women undergoing in vitro fertilization Spot urine (1 or 2 samples) Specific gravity adjusted	-	n.a. 0	-	0.69 (63.4) 87	-		-	0.75 (616) 94
Deziel et al., 2018 <i>USA</i> <i>Connecticut</i>	2010- 2013	200 women (100 population- based controls and 100 women diagnosed with thyroid cancer in a	-	n.r. (n.r.) 44	0.19 (n.r.) 99	0.65 (n.r.) 97	-	2.35 (n.r.) 100	-	0.82 (n.r.) 97

		case-control study) Specific gravity adjusted								
Hoffman et al., 2018 <i>North Carolina</i>	2002- 2005	349 pregnant women (24- 30 weeks) Spot urine Specific gravity adjusted		n.r. (6.1) 49	0.42 (98.0) 98	1.85 (140) 93		7.06 (69.0) 99		1.31 (112) 84
Ingle et al., 2018	2005- 2015	220 men Spot urine (1 to 5 samples/man) (255 samples) Specific gravity adjusted				0.61 (20.24) 85		<MDL. (4.08) 67		0.70 (15.55) 86
Ospina et al., 2018 <i>National Survey US population</i>	2013- 2014	2666 spot urine samples, a random 1/3 sample of participants from the NHANES 2013-2014 Not adjusted for specific Age group: 6 year old and older	0.39 (110) 89	0.16 (46.7) 61	-	0.88 (169) 92	0.25 (70.3) 81	-	-	0.82 (193) 92
		Age group: 6-11 years (n=421)	0.66 (n.r.) n.r.	0.25 (n.r.) n.r.		2.31 (n.r.) n.r.	0.34 (n.r.) n.r.	-	-	1.70 (n.r.) n.r.
		Age group: 12-19 years (n=427)	0.57 (n.r.) n.r.	0.16 (n.r.) n.r.		1.43 (n.r.) n.r.	0.27 (n.r.) n.r.	-	-	1.44 (n.r.) n.r.
		Age group: 20-59 years (n=1266)	0.37 (n.r.) n.r.	0.16 (n.r.) n.r.		0.85 (n.r.) n.r.	0.22 (n.r.) n.r.	-	-	0.73 (n.r.) n.r.
		Age group: 60 years and older (n=552)	0.30 (n.r.) n.r.	0.13 (n.r.) n.r.		0.43 (n.r.) n.r.	0.28 (n.r.) n.r.	-	-	0.65 (n.r.) n.r.
Phillips et al., 2018	2014- 2016	203 children (38-73 months) Three spot urine samples collected over a 48-h period Specific gravity adjusted		n.r. (31.9) 80	n.r. (19.2) 97	n.r. (80.7) 100		n.r. (61.5) 100		n.r. (50.9) 99
Castorina et al., 2017b <i>USA California</i>	1999- 2000	310 pregnant women (26±2.4 weeks) Spot urine Specific gravity adjusted	-	n.a. 0		0.41 (53.1) 77.7	-	0.34 (5.47) 71.6	-	0.93 (54.1) 79.4
Hoffman et al., 2017a <i>USA North Carolina</i>	2001- 2006	349 pregnant women (24-30 weeks) Spot urine Specific gravity adjusted	-	0.7 (6.1) 48.7	0.4 (98) 98.3	1.9 (140) 92.8	-	7.1 (69) 99.4	-	1.3 (112) 83.7
Preston et al. 2017 <i>USA Massachusetts</i>	2010- 2011	51 adults (office workers, 26 men and 25 women) Spot urine, three sampling rounds, interval 6 months							n.r. (0.17-142) n.r.	

		135 samples Specific gravity adjusted								
Romano et al., 2017 <i>USA Rhode Island</i>	2014	58 pregnant women (spot urine samples collected at 12, 28 and/or 35 weeks of gestation) Specific gravity adjusted	0.31 (n.r.) 74	n.r. (n.r.) 53	-	1.18 (n.r.) 93	n.r.(n.r.) 33	-	-	0.93 (n.r.) 95
Thomas et al., 2017 <i>USA Washington</i>	2012- 2014	41 children (15-18 months) Spot urine Specific gravity adjusted Lab 1 N=21	-	-	-	5.47 (64.66) 95	-	0.48 (2.68) 81	-	2.71 (16.56) 100
		Lab 2 N=20	-	-	-	2.01 (202.4) 85	-	-	-	7.72 (26.93) 100
Butt et al., 2016 <i>USA California</i>	2015	28 mothers Spot urine Specific gravity adjusted	-	n.a. (4.0) 11	2.4 (104) 100	2.8 (14.3) 100	-	2.0 (14.8) 100	-	1.2 (3.5) 100
		33 paired children (2-70 months, mean 44 months) Spot urine Specific gravity adjusted	-	n.a. (3.4) 9	2.0 (23.2) 100	7.4 (798) 100	-	2.1 (8.5) 100	-	2.5 (82.0) 100
Carignan et al., 2016 <i>Eastern United States</i>	2012	11 female gymnasts (older than 15 years in age) Several samplings on practice day Specific gravity adjusted	-	-	-	0.76 (3.99) 100	-	-	-	8.71 (58.4) 100
Hammel et al., 2016 <i>USA North Carolina</i>	2015	40 adults (15 men and 25 women) First morning void on 3 separate days Specific gravity adjusted	-	n.a.(0.57) 18	1.12 (16.99) 100	2.06 (21.21) 100	-	-	-	1.16 (26.77) 100
Kosarac et al., 2016 <i>Canada</i>	2010- 2012	20 pregnant women (second and third trimester of pregnancy) and 4 post-partum women Spot urine Not adjusted for specific gravity	0.46 (1.25) 37	0.46 (2.41) 54	n.a. (0.53) 4 (free form, without enzymatic deconjugation)	0.26 (1.77) 29	-	-	<0.08 (1.02) 17	2.94 (25.7) 92
Petropoulou et al., 2016 <i>USA California</i>	n.r.	13 adults (8 women and 5 men) Spot urine Not adjusted for specific gravity	1.3 (15.0) 100	0.3 (3.5) 100	-	2.4 (7.3) 100	-	-	-	1.5 (5.6) 100
Hoffman et al., 2015a <i>USA North Carolina</i>	2014- 2015	43 children (2-18 months, mean 7.9 months) Spot urine Specific gravity adjusted		n.r. (7.5) 19		n.r. (541) 100		n.r. (6.1) 35		n.r. (26.5) 93
Hoffman et al., 2015b <i>USA North Carolina</i>	2012	53 adults (26 men and 27 women) Spot urine Specific gravity adjusted	-	-	-	n.r. (4.46) 83	-	-	-	n.r. (9.09) 91
Su et al., 2015 <i>Canada</i>	2014	12 urine samples from 4 individuals Spot urine, 3 consecutive days Not adjusted for specific gravity	n.r. (12.33) 100	n.r. (0.68) 42	-	n.r. (1.17) 83	n.r. (<MDL or <LOQ) 42	-	<MDL	n.r. (1.29) 75

Butt et al., 2014 <i>USA New Jersey</i>	2013-2014	19 mothers Spot urine Specific gravity adjusted	-	n.r. (0.64) 14	-	n.r. (11.0) 100	-	n.r. (2.3) 100	-	n.r. (68.7) 95
		23 paired children (1-5 years) Spot urine Specific gravity adjusted	-	n.r. (0.46) 4	-	n.r. (251) 100	-	n.r. (10.1) 92	-	n.r. (140) 100
Dodson et al., 2014 <i>USA California</i>	2011	16 adults Spot urine Not adjustment for specific gravity.	0.63 (2.1) 75	n.a. (0.97) 31	-	0.09 (3.9) 94	0.11 (0.45) 56	-	n.a. (0.71) 12	0.44 (6.8) 62
Hoffman et al., 2014 <i>USA North Carolina</i>	2011-2012	8 pregnant women (18 th and 28 th week of pregnancy: 24-h urine and first morning voids. After birth of child: 1 spot urine) 39 urine samples Not adjusted for specific gravity	-	-	-	1.1 (19.9) 97	-	-	-	1.6 (37.3) 97
Meeker et al., 2013 <i>USA Massachusetts</i>	2002-2007	45 men Spot urine Presumably specific gravity adjusted	-	-	-	0.12 (25.0) 91	-	-	-	0.27 (9.84) 96
Cooper et al., 2011 <i>North America</i>	n.r.	3 adults 9 urine samples Spot urine Specific gravity adjusted	-	-	-	0.37 (3.47) 100	-	-	-	1.81 (63.8) 100

^a In most studies, the metabolite method limit of detection (MLOD) was in the range of 0.01-0.6 µg/l, depending on the method used. It was lower for BDCIPP and BBOEP (3 ng/l) in the study of He et al., 2018a) and for BCEP, BDCIPP, and DPHP in the study of Sun et al., 2018 (2-5 ng/l). The MLOD was in the range of 1.0 to 2.7 µg/l for BCIPP in the study of Hammel et al. (2016), for BCIPP and DPHP in the study of Kosarac et al. (2016), and for DNBP and DPHP in the study of Yoshida et al. (2012). The limit of quantification (LOQ) of BCEP, BCIPP, BDCIPP, and DNBP was in the range of 0.8-1.6 µg/l in the study of Chen et al. (2018). The LOQs of BCEP and BCIPP were 1.2, and 3.7 µg/l, respectively, in the study of Van den Eede et al. (2013b). The LOQ of DNBP and DPHP were 2.3, and 2.6 µg/l, respectively, in the study of Yoshida et al. (2012).

^a Mean

n.r. Not reported.

n.a. Not applicable

- Not analyzed

Table 4. Urinary concentrations of the principal metabolites of PEFRs in workers (µg/L).

50th percentile (max) % ≥ limit of detection (LOD)

Author, location	Sampling year	N (population)	BCEP	BCIPP	BCIPHP	BDCIPP	DNBP	ip-PPP	DPHP	Other
Bello et al., 2018 <i>USA</i>	n.r.	12 spray polyurethane foam applicators (construction insulation) Spot urine pre- and post-shift (24 samples) Specific gravity adjusted	-	6.2 ^a (51.4) 100	88.8 ^a (703) 100	5.3 ^a (33.4) 100	-	27.9 ^a (134) 100	6.5 ^a (36.1) 100	
Tao et al., 2018 <i>China</i>	2016-2017	26 hotel room attendants (52 samples) Morning void and post-shift urine Specific gravity adjusted	-	-	-	0.23 (2.4) 79	0.048 (1.3) 38	-	0.24 (1.8) 87	BBOEP : 0.11 (9.0) 59 DoCP&DpCP : 0.17 (1.1) 79 and 87
Yan et al., 2018 <i>China</i>	n.r.	e-waste recycling site 88 workers (men and women) First morning void Not adjusted for specific gravity	1.77 (48.3) 94	n.r. (0.31) 16	-	0.23 (31.8) 82	n.r. (0.96) 41	-	0.70 (26.6) 93	BBOEP : n.r. (21.0) 35 DBP: n.r. (0.96) 41
		Incineration plant 30 workers (men and women) First morning void Not adjusted for specific gravity	1.44 (22.5) 97	n.r. (0.97) 40	-	0.22 (3.56) 93	-	-	0.11 (3.39) 90	BBOEP : n.r. (26.2) 43 DBP: 0.30 (34.8) 100
Jayatilaka et al., 2017 <i>USA</i>	2010-2011	146 firefighters Spot urine collected within 3h after firefighting Not adjusted for specific gravity	0.86 (10) 90	0.24 (2.9) 63	-	3.4 (44) 100	0.18 (2.4) 92	-	2.9 (28) 100	DpCP : <LOD (0.31) 34
	2015	76 adults from the general population Not adjusted for specific gravity	<LOD (4.1) 10	<LOD (0.98) 5	-	0.69 (6.8) 100	<LOD (0.26) 5	-	0.89 (5.6) 100	n.a. 0

Schindler et al., 2014 <i>Germany</i>	n.r.	5 aircraft maintenance technicians Spot urine Not adjusted for specific gravity Preshift/Postshift	0.5/0.3 (1.7)/(0.5) 100	0.2/0.2 (0.3)/(0.3) 70	-	-	12.5/23.5 (37.2)/(51.6) 100	-	2.9/3.5 (7.4)/(7.9) 100	DoP, DmCP, or DpCP < LOD (i.e. 0.5)
Carignan et al., 2013 <i>USA</i>	2009	29 office workers (women and men) Spot urine during afternoon of a work day Specific gravity adjusted	-	-	-	408 ^a .(1760) 100	-	-	-	
Schindler et al., 2013, 2009a and b (Anderson, 2015; Weiss et al., 2015) <i>Germany</i>	n.r.	332 urine samples from air crews Spot urine collected within 12h after exposure Not adjusted for specific gravity	0.33 (20.3) 82	0.16 (6.87) 65	-	-	0.28 (9.72) 100	-	1.10 (302) 100	DoP: < LOD (i.e. 0.5) DmCP: <0.5b (0.62) 0.3 DpCP < LOD (0.55) 0.3
		30 individuals from the general population (11- 68 years) Spot urine Not adjusted for specific gravity	<0.1 ^b (27.5) 50	<0.25 ^b (0.85) 12	-	-	<0.25 ^b (0.26) 4	-	0.52 (5.47) 68	DoP, DmCP, or DpCP < LOD (i.e. 0.5)

^a Geometric mean

^b Limit of detection (LOD)

n.r. Not reported

- Not analyzed

Table 5 PEFR concentrations in hair and nails (ng/g dry weight)

50th percentile (range) % ≥ **limit of detection**

Author Location N (population) Sampling year		TCEP	TCIPP	TDCIPP	TNBP	TEHP	EHDPP	TPHP	TBOEP	Other
Europe										
Alves et al., 2017 <i>Belgium</i> A woman and a man Year n.r.	Hair (scalp segment, one sample)									<u>DPHP</u> Woman ^c : 0.25 Man ^c : 0.23
	Fingernails (4 or five collections over 2 months)									Woman ^c : 40002 Man ^c : 80.5
	Toenails (4 or five collections over 2 months)									Woman ^c : 6815 Man ^c : 18.5
Kurcharska et al., 2015a <i>Norway</i> 48 mothers and their 54 children (6-12 year old) 2012	Hair (scalp segment) <u>Mothers</u>	72 (<33 ^a -163) 16		30 (<9 ^a -3744) 91	22 (5-672) 100	12 (<1 ^a -53) 96	27 (5 -265) 100	52 (5-1256) 100	65 (14-1253) 100	<u>TCP^b</u> 8 (<2 ^a -134) 78
	Hair (scalp segment) <u>Children</u>	59 (<33 ^a -118) 26		31 (<9 ^a -2698) 92 <i>n.r.</i>	11 (3 -150) 100 <i>n.r.</i>	8 (<1 ^a -118) 90 <i>n.r.</i>	21 (2 ^a -346) 100 <i>n.r.</i>	63 (6-363) 100 <i>n.r.</i>	318 (34-2411) 100 <i>n.r.</i>	<u>TCP^b</u> 8 (<2 ^a -74) 62
Martin et al., 2015 <i>Germany</i> 4 women	Hair (scalp segment)		<i>n.r.</i> (0.18-1.70) 100		0			<i>n.r.</i> (0.10-0.91) 100		

Year n.r.										
Kurcharska et al., 2014 <i>Belgium</i> 20 adults Year n.r.	Hair	55 (34-404) 70		42 (10-2969) 95	32 (7-5032) 95	10 (2-322) 75	15 (5-105) 100	59 (7-237) 100	37 (7-338) 100	<u>TCP^b</u> 5 (3-73) 65
<i>China</i>										
Qiao et al., 2016 49 adults (27 man and 22 woman) 2014	Hair (two segments)	3.61 (<3.50 ^a -64.9) 57	43.9 (<6.53 ^a -141) 98	4.14 (<1.04 ^a -73.8) 86	3.30 (<0.61 ^a -25.4) 98	24.1 (<0.05 ^a -151) 98	11.8 (5.78 ^a -78) 71	20.5 (<1.43 ^a -352) 84		<u>TiPP</u> 2.43 (<0.81 ^a -12.4) 94
<i>United States</i>										
Liu et al., 2016 <i>Indiana</i> 50 adults 2014	Hair (scalp segment)	240 (60-2740) 68	450 (100-9840) 88	360 (70-10490) 90				220 (70-4710) 98		
	Fingernail	190 (93-1860) 20	220 (74-2410) 36	300 (90-1410) 66				370 (110-59800) 74		
	Toenail	150 (100-150) 8	230 (90-5150) 32	230 (75-2300) 50				1080 (54-232900) 74		
Liu et al., 2015 <i>Indiana</i> 5 adults Year n.r.	Hair (scalp segment)	<i>n.r.</i> (<75 ^a -1950) 80	<i>n.r.</i> (290-1190) 100	<i>n.r.</i> (<75 ^a -970) 80				<i>n.r.</i> (76-310) 100 <i>r.</i>		
	Fingernail	<i>n.r.</i> (<150 ^a -<150) 0	<i>n.r.</i> (<150 ^a -<150) 0	<i>n.r.</i> (280-630) 100				<i>n.r.</i> (<150 ^a -17500) 100		

^a Limit of quantification (LOQ)

^b Sum of isomers

^c Average level

n.r. Not reported

S1. Chemical structure of the main PEFR metabolites

BECP

BCIPP

BCIPHPP

BDCIPP

DBNP

BEHP

ip-PPP

BBOEP

BBOEHEP

DPHP

DoCP

5-OH-EHDPPH