

Environmental DNA provides information on sediment sources: A study in catchments affected by Fukushima radioactive fallout

O. Evrard, J. Patrick Laceby, Gentile Francesco Ficetola, Ludovic Gielly, Sylvain Huon, Irène Lefevre, Yuichi Onda, Jérôme Poulenard

▶ **To cite this version:**

O. Evrard, J. Patrick Laceby, Gentile Francesco Ficetola, Ludovic Gielly, Sylvain Huon, et al.. Environmental DNA provides information on sediment sources: A study in catchments affected by Fukushima radioactive fallout. Science of the Total Environment, 2019, 665, pp.873-881. 10.1016/j.scitotenv.2019.02.191 . hal-02049144

HAL Id: hal-02049144
https://hal.science/hal-02049144

Submitted on 12 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

*Manuscript (double-spaced and continuously LINE and PAGE numbered)-for final publication
Click here to view linked References

Environmental DNA provides information on sediment sources: A study in catchments affected by Fukushima radioactive fallout

Olivier Evrard[1,*], J. Patrick Laceby[2], Gentile Francesco Ficetola[3,4], Ludovic Gielly[3], Sylvain Huon[5], Irène Lefèvre[1], Yuichi Onda[6], Jérôme Poulenard[7]

[1]Laboratoire des Sciences du Climat et de l'Environnement (LSCE-IPSL), Unité Mixte de Recherche 8212 (CEA/ CNRS/UVSQ), Université Paris-Saclay, Gif-sur-Yvette, France.
[2]Environmental Monitoring and Science Division (EMSD), Alberta Environment and Parks (AEP), 3115 – 12 Street NE, Calgary, Alberta, Canada.
[3]Université Grenoble Alpes, CNRS, Laboratoire d'Écologie Alpine, Grenoble, France.
[4]Departement of Environmental Science and Policy, Università degli Studi di Milano, Milano, Italy.
[5]Sorbonne Universités, UPMC Univ Paris 06, Institut d'Ecologie et des Sciences de l'environnement de Paris (iEES), Paris, France.
[6]Center for Research in Isotopes and Environmental Dynamics (CRIED), University of Tsukuba, Tsukuba, Japan.
[7]Université Grenoble Alpes, Université Savoie Mont Blanc, CNRS, EDYTEM (Environnements, DYnamiques et TErritoires de la Montagne), Chambéry, France

***Correspondence to :** olivier.evrard@lsce.ipsl.fr.

Abstract

An excessive supply of sediment is observed in numerous rivers across the world where it leads to deleterious impacts. Information on the sources delivering this material to waterbodies is required to design effective management measures, and sediment tracing or fingerprinting techniques are increasingly used to quantify the amount of sediment derived from different sources. However, the current methods used to identify the land use contributions to sediment have a limited discrimination power. Here, we investigated the potential of environmental DNA (eDNA) to provide more detailed information on the plant species found in sediment source areas as a next generation fingerprint. To this end, flood sediment deposits (*n*=12) were collected in 2017 in two catchments impacted by the Fukushima radioactive fallout along differing river sections draining forests, cropland or a mix of both land uses. Conventional fingerprints (i.e. fallout radionuclides and organic matter properties) were also measured in these samples. The conventional fingerprint model results showed that most sediment samples contained a dominant proportion of subsoil material. Nevertheless, the eDNA information effectively discriminated the three above-mentioned groups of sediment, with the dominance of tree, shrub and fern species in sediment sampled in rivers draining forests versus a majority of grass, algae and cultivated plant species in sediment collected in rivers draining cropland. Based on these encouraging results, future research should examine the potential of eDNA in mixed land use catchments where the contribution of topsoil to sediment dominates and where the cultivation of land has not been abandoned in order to better characterize the memory effect of eDNA in soils and sediment.

Keywords: Sediment fingerprinting; catchment; sediment source; FDNPP accident; ancient plant DNA (aDNA).

Introduction

Soil erosion is often exacerbated in agricultural catchments where erosion rates may be an order of magnitude greater on cropland relative to grassland and forested landscapes (Cerdan et al., 2010; Montgomery, 2007). Although particle detachment processes are generally reduced in areas with dense vegetation or an abundant litter layer, soil erosion regularly occurs on these natural landscapes (Fukuyama et al., 2010; Laceby et al., 2016b). Soil erosion rates are accelerating worldwide (Amundson et al., 2015), resulting in deleterious on-site and off-site consequences (Evrard et al., 2007).

One major off-site consequence of soil erosion are elevated suspended sediment loads that reduce fish spawning habitats (Owens et al., 2005), decrease reservoir life spans (Foucher et al., 2014) and degrade water quality (Gateuille et al., 2014). Accordingly, it is important to quantify the relative contribution of the major sources supplying suspended sediment to river networks in order to develop effective best management practices that mitigate deleterious downstream effects of accelerated soil erosion.

Sediment source fingerprinting is increasingly used (Walling, 2013) to estimate the relative source contributions of detached particles through the analysis of physical or biogeochemical properties, or 'fingerprints,' in source soils and their downstream sediments (Walling et al., 1979). A variety of fingerprinting properties can be used to estimate sediment source contributions, including fallout radionuclides (e.g. ^{137}Cs, $^{210}Pb_{xs}$, ^{7}Be) (Evrard et al., 2013), element geochemistry (Douglas et al., 2003), mineral magnetics (Walden et al., 1997), colour (Legout et al., 2013) or organic parameters (TN, TOC) and their isotopes (Gourdin et al., 2015), amongst others.

There are three fundamental approaches to sediment source fingerprinting. First, researchers can investigate the dominant erosion processes (e.g. surface or subsurface erosion) contributing material to the river network (Wallbrink et al., 1998). Second, researchers can examine temporal dynamics, to verify whether sediment transiting the river network consists of recently or previously eroded

material (Evrard et al., 2016a). Third, researchers can determine the dominant spatial source areas contributing particulate material. For this latter approach, the dominant spatial areas often investigated include lithological regions (e.g. bedrock or surficial geology)(Le Gall et al., 2017), soil types (Lepage et al., 2016), and land use or land cover (Tiecher et al., 2017).

To investigate sediment derived from different land uses or land covers, researchers are increasingly tracing the vegetation signatures from sources areas with carbon and nitrogen elemental concentrations (TOC and TN) and their stable isotope ratios ($\delta^{13}C$ and $\delta^{15}N$) (Gourdin et al., 2015; Huon et al., 2013; Laceby et al., 2016c). As the discrimination power of these properties is limited, particularly in areas with a mixture of C_3 and C_4 plants (Evrard et al., 2013), researchers are expanding their toolbox, for example, by tracing the fatty acids in sediments to their source soils with Compound-Specific Stable Isotope (CSSI) analysis (Gibbs, 2007). These approaches provide detailed information regarding the vegetative source signature of the sediments, including potentially different land uses (e.g., permanent pasture, woodland, cultivated land) or even agricultural practices (e.g. maize and stubble)(Blake et al., 2012). Nevertheless, the discrimination potential of these organic tracers is typically limited to the broad land uses (Bravo-Linares et al., 2018) and debate remains regarding the conservative behavior of these organic tracers during their transport across the landscape (Koiter et al., 2013; Reiffarth et al., 2016). Therefore, other fingerprints are required that provide more specific and potentially more conservative source information.

Environmental DNA (eDNA) is the DNA obtained from environmental samples such as water and sediment. The DNA of plants and animals living in soil and above the soil is transferred to the river network by erosion processes, and was shown to be preserved in deposited particulate matter (Giguet-Covex et al., 2014; Parducci et al., 2017; Pedersen et al., 2015). The preservation of eDNA is much higher in sediment – including in sediment dating back to the Holocene or the Pleistocene (Willerslev et al., 2003) – than in freshwater ecosystems (Dejean et al., 2011; Thomsen et al., 2012). With recent advances in High Throughput Sequencing, eDNA offers a novel approach that could

provide far more detailed sediment source information than CSSI tracers. In other fields, eDNA has already provided highly detailed information regarding specific plant types and/or land management practices with the analysis of eDNA in ancient sediment in lacustrine sediment cores (Ficetola et al., 2018; Pansu et al., 2015; Sjogren et al., 2016). However, to the best of our knowledge, eDNA has never been analysed in contemporary sediments with the objective of determining the relative contribution of different sediment sources through the improved discrimination of different vegetation species.

In catchments of Northeastern Japan impacted by radioactive fallout from the Fukushima Dai-ichi Nuclear Power Plant (FDNPP) accident in March 2011 (Evrard et al., 2015), the quantification of the land use contributions to sediment transiting the rivers to the Pacific Ocean is imperative in order to prioritize remediation works. Previous fingerprinting research based on the measurement of carbon and nitrogen elemental concentrations and their stable isotope ratios showed that subsoils were the dominant sources (45±26%) of particulate matter to the rivers, followed by cultivated land (38±19%) and forests (17±10%) including evergreen and deciduous species (Laceby et al., 2016b). However, the identification of the land use contributions was provided with limited resolution so far. Although areas cultivated with rice (*Oryza sativa L.*) were shown to supply a disproportionate amount of radiocesium contaminated sediment to the region's waterbodies (Yoshimura et al., 2016), uncertainties remain regarding the specific contribution of paddies compared to that of other cultivated fields or that of forested areas to sediment transiting these rivers.

Accordingly, the objective of this study is to explore the potential of eDNA as a next generation sediment source fingerprint. As this is, to our knowledge, the first application of eDNA in a sediment source fingerprinting context, our goal is to compare the discrimination potential of the source information provided by eDNA to that achieved with carbon and nitrogen properties (TOC, TN, $\delta^{13}C$ and $\delta^{15}N$) and fallout radionuclides (^{137}Cs). Our main objective is to demonstrate the potential utility of eDNA as a sediment source fingerprint, through highlighting its higher source discrimination

potential relative to conventional sediment fingerprinting approaches. Our secondary objective is to overview the challenges and opportunities of developing eDNA as a sediment fingerprinting parameter of the future.

Materials and Methods

Study area

This research was conducted in the Niida (275 km²) and Ota (77 km²) catchments (Fig. 1), draining the main radioactive contamination plume of the Fukushima Prefecture, Northeastern Japan. This region is exposed to a very erosive climate (mean annual rainfall of ~1400 mm), with the occurrence of spring floods and typhoons being the main hydro-sedimentary events that may lead to significant erosion and sediment transfer in rivers (Chartin et al., 2017; Laceby et al., 2016a). The Niida and Ota River catchments are characterized by land uses that are representative of those found in the fallout impacted region, with a dominance of forests (75%) and cropland (21%).

The main catchment features include an upstream coastal mountain range (<900 m) and a broad, more densely inhabited, coastal plain (i.e. <100 m). Dams are installed at the transition between both zones on the Ota River (Yokokawa Dam) and on the main Niida River tributary (Takanokura Dam) in the coastal plain (i.e., Mizunashi River) to provide water for drinking, and irrigation to the local communities. These dams create sediment disconnectivities between upper and lower catchment areas, with dams storing sediment (Chartin et al., 2017). The Niida River mainstream is devoid of dam.

Sediment sampling

Sediment samples were collected in June 2017 at sites draining only forests (*n*=5), a dominance of cropland (*n*=4), or a mix of both land uses (*n*=3; Fig. 1). These samples comprise particulate material that settled on channel banks, inset benches and floodplains during the falling limb of the last significant hydro-sedimentary event. One soil sample (0 – 5 cm surface layer) was also collected in a

paddy field located near the Ota River for comparison (Table 1). For fallout radionuclide and carbon/nitrogen measurements, 5 subsamples (~5 g per subsample) of recently deposited material at each site were taken with a plastic spatula over a 5 m reach and composited into one sample. For the eDNA analyses, 5 subsamples (~ 5 g per subsample) were collected at each site with a trowel, sterilized with a blowtorch, mixed and homogenized. Approximately 15 g of material were stored in a sealed container with silica gel bags for in-situ drying.

Fallout radionuclide analysis and modelling

Prior to analyses, samples were dried at 40°C for ~48 h, sieved to 2 mm, ground to a fine powder in an agate mortar, and pressed into 15 mL polyethylene containers for analysis. Fallout radionuclide activities were determined with gamma spectrometry using coaxial N- and P-type HPGe detectors (Canberra/Ortec). ^{137}Cs activities were measured at the 662 keV emission peak. ^{134}Cs activities were calculated as the mean of the counts found at the 605 and 796 keV emission peaks. All radionuclide activities were decay-corrected to March 14, 2011, the date of the main radionuclide fallout deposition (Kinoshita et al., 2011).

Based on the ^{137}Cs analyses, the relative contributions of contaminated (high ^{137}Cs) versus depleted (low ^{137}Cs) sources of radiocesium contaminated sediment were modelled in the Niida catchment. The model was run with the source sample data from Evrard et al. (2016b) which included 118 samples of contaminated source soil and 42 samples of depleted source material sampled in the Niida catchment. A distribution modelling approach estimated the relative contribution of these two end-members (Laceby and Olley, 2015). The objective was to estimate whether the sediment samples in the Niida catchment were derived from upstream contaminated sources (e.g. forested reaches) versus cropland or other areas in the coastal plains that did not receive significant fallout. Although the source dataset has been published in Evrard et al. (2016b), the sediment dataset is original and has not been published elsewhere.

185 ***Carbon and nitrogen analysis and modelling***

186 A subsample of the material analysed for fallout radionuclides was analysed for TOC, TN, $\delta^{13}C$ and
187 $\delta^{15}N$ with a continuous flow Elementar® VarioPyro cube analyzer coupled to a Micromass® Isoprime
188 Isotope Ratio Mass Spectrometer (EA-IRMS) at the Institute of Ecology and Environmental Science
189 (iEES) in Paris. First, TOC and $\delta^{13}C$ were measured with a set of tyrosine standards (Coplen et al.,
190 1983; Girardin and Mariotti, 1991). Second, TN and $\delta^{15}N$ were analysed with sample weights
191 optimized after the carbon analyses. Oxygen for combustion was injected during 70s (30 mL min^{-1})
192 and temperatures were set at 850°C and 1120°C for the reduction and combustion furnaces,
193 respectively (Agnihotri et al., 2014).

194 The carbon and nitrogen parameters were used to trace land use/land cover sources with a source
195 data set from Laceby et al. (2016b). This included TOC, TN, $\delta^{13}C$ and $\delta^{15}N$ data for 46 forest, 28
196 cultivated and 25 subsoil source samples analysed with identical methods at iEES. A concentration
197 dependent distribution modelling approach estimated the relative contribution of these two end-
198 members (Laceby et al., 2015) and provide the foundation for comparing the eDNA analyses with
199 modeled source contributions that differentiate agricultural sources from those anticipated from
200 forests and subsoil/decontaminated sources in the Fukushima coastal region that was impacted by
201 the FDNPP accident. Although the source dataset is published in Laceby et al. (2016b), the sediment
202 dataset is original and has not been published elsewhere.

203 ***eDNA analysis and interpretation***

204 Sediment eDNA was extracted following the protocol described in Pansu et al. (2015), targeting
205 extra-cellular eDNA. For each sample, ~15 g of sediment was mixed with 15 ml of saturated
206 phosphate buffer (Na_2HPO_4; 0.12 M, pH ≈ 8) for 15 min. Two ml of the mixture was centrifuged
207 (10min at 10000 g); 400 µl of resulting supernatant was kept as starting material for extraction using
208 the NucleoSpin® Soil kit (Macherey-Nagel, Düren, Germany), skipping the cell lysis step and following

manufacturer's instructions (Taberlet et al., 2012). The extracted DNA was eluted in 100 µl of SE buffer and used as PCR template. Seven extraction controls were also conducted. The eDNA of vascular plants was amplified with the g-h primers, which amplify a short variable region on chloroplast DNA and have very limited mismatches with sequences of vascular plants (Taberlet et al., 2018). Additionally, six PCR controls containing PCR mix and no DNA template, along with six positive PCR controls were performed (Parducci et al., 2017). Each sediment sample and control was amplified in four PCR replicates (Ficetola et al., 2015). The sequencing was conducted by 2 × 125–base pair pair-end sequencing on an Illumina HiSeq 2500 platform. The sequences of DNA were filtered with OBITools software (Boyer et al., 2016). Plant sequences were assigned with a database of the vascular plants found in Japan. Sequences with a >97% match with a plant genus were kept, providing the table of the Molecular Taxonomic Units (MOTUs) to analyse. For each sample, the sum of the reads of the four replicates was calculated. Those MOTUs associated with a total number of reads lower than 1000 were removed from further analysis. The logarithm of the reads of those sequences kept for analysis +1 was then calculated as a measure of relative abundance (Yoccoz et al., 2012). A Principal Component Analysis (PCA) was conducted in order to discriminate various groups of sediment based on the significant MOTUs detected in the samples. A hierarchical classification of the retained MOTUs was also performed and, finally, a Permutational Multivariate Analysis of Variance Using Distance Matrices (PERMANOVA) was conducted to assess the amount of variation of the eDNA explained by the sources.

Results and discussion

Fallout radionuclides

Sediment in lag deposit samples draining forest soils had both significantly higher and more variable ^{137}Cs activity concentrations (mean (M): 23,436 Bq kg^{-1}; SD: 17,810 Bq kg^{-1}) compared to rivers draining mixed land uses (both forest and cropland) (M: 3238 Bq kg^{-1}; SD: 3156 Bq kg^{-1}) and cropland only (M: 2962 Bq kg^{-1}; SD: 2002 Bq kg^{-1}). The ^{137}Cs activity concentration measured in the paddy field soil was low (1384 ±12 Bq kg^{-1}), although it remained in the range of values found in sediment collected in rivers draining cultivated land. ^{134}Cs/^{137}Cs activity ratios for all three sediment sample types were in a narrow range (0.95–1.02) consistent with the ratio expected from fallout from the FDNPP accident (Kobayashi et al., 2017).

Sediment ^{137}Cs activity concentrations were modeled in the Niida catchment to investigate whether they were derived primarily from contaminated regions in the catchments or those that have been decontaminated and/or not impacted by the initial fallout. On average, 6% (SD 7%) of sediment in the Niida catchment was modeled to be derived from sources contaminated by the initial fallout (e.g. upper catchment areas and surface soils). The forest sample from the Mizunashi reach had the highest contribution of fallout-contaminated landscapes (14%, Model Error (ME) 1.2%). The paddy field sample farther downstream on this reach had a 5% (ME 1%) contribution from contaminated areas. In contrast, on the main stem, three samples had a less than 2% contribution from contaminated areas. Interestingly, the sediment sample collected the farthest downstream in the Niida catchment had an increased contaminated area contribution of again ~14%. Overall, most of the sediment sampled was derived from sources that were not contaminated with fallout from the FDNPP (e.g. subsoils, decontaminated soils, or regions near the coast that did not receive fallout).

Carbon and nitrogen composition

256 The TN content was too low for accurate measurement in 2 of the 13 samples analyzed (FNL1243 and
257 FOL1233). These samples also had the lowest TOC (<0.6%) of all analyzed samples indicating that
258 they are mainly comprised of coarser material, most likely provided by subsoil sources. The sediment
259 collected in the forested catchments had lower mean $\delta^{13}C$ (M -28.4‰, SD 0.8‰) and $\delta^{15}N$ (M 1.8‰,
260 SD 1.6‰) compared to those sampled in cultivated catchments (M -27.1‰, SD 0.9‰ and M -4.3‰,
261 SD 0.3‰ respectively). Three of the forested catchment sediment samples, and one of the cultivated
262 catchment sediment sample (FNL1245) had low TOC (<1.2%) and low TN (<0.08%) suggesting higher
263 contributions of subsoil sources that are known to be low in organic matter content. The three
264 remaining cultivated catchment sediment samples had slightly higher TOC (M 1.4%, SD 0.2%) and TN
265 (M 0.11%, SD 0.03%). The remaining forest catchment sample (FOL1239) had very high TOC (19.3%)
266 and TN (1.1%). The carbon and nitrogen parameters suggest that the sediment collected in forested
267 and cultivated catchments likely have significant subsoil source material contributing to the low TOC
268 and TN content, with the exception of slightly elevated material derived from cultivated sources, and
269 also one sample, owing to its very high organic contents, dominated by forest material.

270 Sediment in three of the mainly forested catchments were modelled to be almost entirely derived
271 from subsoil sources (M 97%, SD 3%), whereas the high TOC sample was modelled to be entirely
272 from forested sources (100%). Importantly, there was a 0% contribution of cultivated land modelled
273 for the forested sites, supporting that this model is providing relevant predictions as there are
274 essentially no rice paddy fields in these forested regions. For the cultivated catchments, 86% (SD 8%)
275 of sediment was modelled to be derived from subsoil sources, only 13% (SD 8%) from rice paddy
276 fields or other agricultural sources, and with a mean of 2% (SD 1%) from forest sources. One of the
277 two mixed catchment sediment samples (FNL1242) was modelled to be derived entirely from subsoil
278 sources (100%) whereas the other (FNL1245), was modelled to be derived from 44% cultivated land,
279 18% forest and only 38% subsoil sources, the lowest subsoil source contribution modelled. Indeed,
280 subsoil erosion processes dominate the supply of sediment across these catchments (M 77%, SD
281 33%). Furthermore, the sampling location FNL1245 behaved differently than the other samples

again. The higher ^{137}Cs and contaminated source contribution in this sample relative to the others on this reach is likely due to the higher forest and cropland source contributions, which are diluted on the main stem of the Niida in downstream direction across the coastal plain.

eDNA

After filtering the data, 67 abundant MOTUs were kept for analyses (Appendix A). When examining those sequences individually (Fig. 2), woodland species including mosses, ferns and shrubs were almost exclusively found in sediment collected in rivers draining forested areas or mixed land uses. In contrast, grasses and rice (*Oryza sativa L.*) dominated in the samples collected in rivers draining cropland and mixed land uses. In addition, species and/or families of cultivated plants were also found in these latter sediment samples, including Brassicaceae (e.g. cabbage), Fabaceae, Apiaceae (e.g, celery, carrot) or rice. These results demonstrate the potential of eDNA to provide a very detailed discrimination of those land use sources supplying sediment, with the identification of targeted crop types for instance.

To facilitate their analysis, the MOTUs were aggregated in 6 different groups: grasses, wetland grasses and cultivated plants (MOTUs 1 to 4); algae (MOTUs 5 to 14); mosses (MOTUs 15 to 29); ferns (MOTUs 30 to 35); trees and shrubs (MOTUs 36 to 50) and other plants including cultivated crops (MOTUs 51 to 67). The PCA (Fig. 3) using these 67 MOTUs showed a clear discrimination between samples corresponding to (i) the paddy field soil (group A) and (ii) those 3 classes of sediment identified during sampling: material collected in rivers draining cultivated land (group AA), in rivers draining mixed land uses (group B) and in watercourses draining forests (group C).

The hierarchical classification shows that the above-mentioned groups of MOTUs are well discriminated by the eDNA information (Fig. 4). Results from the PERMANOVA analysis showed that grouping explains a very large amount of variation of the eDNA composition ($R^2 = 0.47$), with a strongly significant effect ($p < 0.001$).

306 ***Combining the information provided by fallout radionuclides, organic matter composition and***
307 ***eDNA***

308 For logistical reasons, the current research focused on the analysis of sediment deposits collected
309 from different types of tributaries in catchments of the Fukushima region. The analysis of fallout
310 radionuclides and organic matter properties of these samples, and the associated modeling showed
311 that the majority of these samples contained significant proportions of subsoil material. Various
312 sources may supply large quantities of subsoil material to the rivers in this region (Fig. 5). These
313 include the mobilization of deep soil horizons under forests as a result of landsliding, gullying or road-
314 side erosion (Mizugaki et al., 2008). Channel bank erosion may also provide a significant quantity of
315 material to the rivers, in un-channeled reaches (Fig. 5b). Accordingly, soil particles are not only
316 transferred by overland flow, they are also generated by erosion of deeper soil horizons.
317 Nevertheless, this deeper soil material is still tagged with eDNA. A significant number of MOTUs (67)
318 were identified in these samples containing high proportions of subsoil and the analysis of this eDNA
319 provided the discrimination of the different sediment sample groups identified during fieldwork. As a
320 result, a dominance of tree, fern and moss species were found in sediment collected in rivers
321 draining forests, whereas a majority of grass and cultivated species were detected in sediment
322 collected in watercourses draining cropland.

323 ***Implications for future research and concluding remarks***

324 Although the Fukushima case study was expected to be a relatively straightforward approach for
325 discriminating land use-derived sediment sources in different rivers draining either forested areas or
326 cropland, the situation was more complex owing to the large proportion of subsoil material transiting
327 these river systems (Fig. 5). This is also likely due to the fact that our sampling approach targeted
328 sediment deposited in the river channel after flooding events, rather than suspended matter
329 transiting rivers. However, this approach was guided by the precautions required to avoid the
330 biological contamination of the samples during their collection in the field and the necessity to

331 sterilize tools to collect the lag deposit samples rather than filter large volumes of river water
332 containing lower quantities of suspended matter.

333 Furthermore, cultivated land was abandoned by farmers after the nuclear accident in March 2011.
334 During the sampling campaign conducted in 2017, agricultural activities had not restarted in the
335 study area. This situation likely explains the dominance of subsoil material transiting these rivers as a
336 dense vegetation cover rapidly covered the abandoned fields and protected them against erosion,
337 thereby limiting the contribution from the erosion of cultivated topsoils. Furthermore, this situation
338 outlines the need to further investigate the memory effect of eDNA of cultivated plants in soils after
339 6 years of abandonment, although the presence of plants cultivated in the region before the nuclear
340 accident (e.g. cabbage, rice, mulberry) was demonstrated in the current research.

341 Based on these preliminary, encouraging results, future applications should be preferentially
342 conducted in mixed land use catchments where the contribution of surface material to sediment
343 transiting rivers is dominant and where agricultural activities have not been abandoned for several
344 years. The systematic collection of both soil and sediment samples should be conducted in order to
345 examine the memory effect of eDNA in soils and the potential changes in signatures from the soil to
346 the sediment, including samples collected in rivers and in deposition areas (e.g. ponds or lakes).

347 Although the design of un-mixing models to quantify sediment source contributions to sediment is
348 the ultimate objective of this research, multiple important phases of research are required before we
349 can model eDNA effectively. This has been recently demonstrated by many reviews in the sediment
350 tracing literature (Koiter et al., 2013; Laceby et al., 2017; Owens et al., 2016). For example, the
351 conservative behavior of eDNA must be thoroughly researched to evaluate its persistence on the
352 particles during mobilization, transport and deposition processes. Indeed, prior to modelling we need
353 to understand if eDNA properties are maintained during these processes, or vary in a predicable way.
354 Furthermore, we need to investigate whether is an affinity for eDNA to be bound to different particle
355 size fractions or even different soil and or mineral types. If eDNA indeed behaves conservatively

during sediment generation and deposition processes, without significant particle size impacts, researchers will have to then investigate what is the most appropriate approach to using eDNA in end member mixing models to quantitatively apportion source contributions to material transiting river networks. The analysis of artificial mixtures – in which known proportions of the different sources are combined in the laboratory – will likely be necessary to develop the best approach to incorporate eDNA into end member mixing models. We believe that the quantitative use of eDNA in sediment source fingerprinting models will be extremely beneficial for managing the degradation of our riverine, lacustrine and potentially estuarine environments through providing a more detailed source information than many of the currently employed fingerprints.

Acknowledgements

This work has been supported by the French National Research Agency (ANR) in the framework of the AMORAD project (ANR-11-RSNR-0002). The authors are grateful to Véronique Vaury for conducting the carbon and nitrogen parameter analyses.

References

Agnihotri, R. et al., 2014. Experimental Setup and Standardization of a Continuous Flow Stable Isotope Mass Spectrometer for Measuring Stable Isotopes of Carbon, Nitrogen and Sulfur in Environmental Samples. Mapan, 29(3): 195-205.

Amundson, R. et al., 2015. Soil and human security in the 21st century. Science, 348(6235).

Boyer, F. et al., 2016. obitools: a unix-inspired software package for DNA metabarcoding. Molecular ecology resources, 16(1): 176-182.

Bravo-Linares, C. et al., 2018. First use of a compound-specific stable isotope (CSSI) technique to trace sediment transport in upland forest catchments of Chile. The Science of the total environment, 618: 1114-1124.
Cerdan, O. et al., 2010. Rates and spatial variations of soil erosion in Europe: A study based on erosion plot data. Geomorphology, 122(1-2): 167-177.
Chartin, C. et al., 2017. The impact of typhoons on sediment connectivity: lessons learnt from contaminated coastal catchments of the Fukushima Prefecture (Japan). Earth Surface Processes and Landforms, 42: 306-317.
Chartin, C. et al., 2013. Tracking the early dispersion of contaminated sediment along rivers draining the Fukushima radioactive pollution plume. Anthropocene, 1: 23-34.
Coplen, T.B., Kendall, C. and Hopple, J., 1983. Comparison of stable isotope reference samples. Nature, 302: 236-238.
Dejean, T. et al., 2011. Persistence of environmental DNA in freshwater ecosystems. PloS one, 6(8): e23398.
Douglas, G., Palmer, M. and Caitcheon, G., 2003. The provenance of sediments in Moreton Bay, Australia: a synthesis of major, trace element and Sr-Nd-Pb isotopic geochemistry, modelling and landscape analysis, The Interactions between Sediments and Water. Springer, pp. 145-152.
Evrard, O., Bielders, C.L., Vandaele, K. and van Wesemael, B., 2007. Spatial and temporal variation of muddy floods in central Belgium, off-site impacts and potential control measures. Catena, 70(3): 443-454.
Evrard, O. et al., 2016a. Combining multiple fallout radionuclides (^{137}Cs, ^{7}Be, $^{210}Pb_{xs}$) to investigate temporal sediment source dynamics in tropical, ephemeral riverine systems. Journal of Soils and Sediments, 16(3): 1130-1144.
Evrard, O. et al., 2015. Radiocesium transfer from hillslopes to the Pacific Ocean after the Fukushima Nuclear Power Plant accident: A review. Journal of environmental radioactivity, 148: 92-110.
Evrard, O. et al., 2016b. Quantifying the dilution of the radiocesium contamination in Fukushima coastal river sediment (2011–2015). Scientific Reports, 6: 1-8.
Evrard, O. et al., 2013. Tracing sediment sources in a tropical highland catchment of central Mexico by using conventional and alternative fingerprinting methods. Hydrological Processes, 27(6): 911-922.
Ficetola, G.F. et al., 2015. Replication levels, false presences and the estimation of the presence/absence from eDNA metabarcoding data. Molecular Ecology Resources, 15(3): 543-56.
Ficetola, G.F. et al., 2018. DNA from lake sediments reveals long-term ecosystem changes after a biological invasion. Science advances, 4(5): eaar4292.
Foucher, A. et al., 2014. Increase in soil erosion after agricultural intensification: Evidence from a lowland basin in France. Anthropocene, 7: 30-41.
Fukuyama, T. et al., 2010. Quantifying the impact of forest management practice on the runoff of the surface-derived suspended sediment using fallout radionuclides. Hydrological Processes, 24(5): 596-607.
Gateuille, D. et al., 2014. Mass balance and decontamination times of Polycyclic Aromatic Hydrocarbons in rural nested catchments of an early industrialized region (Seine River basin, France). The Science of the total environment, 470-471: 608-17.
Gibbs, M.M., 2007. Identifying Source Soils in Contemporary Estuarine Sediments: A New Compound-Specific Isotope Method. Estuaries and Coasts, 31(2): 344-359.
Giguet-Covex, C. et al., 2014. Long livestock farming history and human landscape shaping revealed by lake sediment DNA. Nature Communications, 5.
Girardin, C. and Mariotti, A., 1991. Analyse isotopique du ^{13}C en abondance naturelle dans le carbone organique: un système automatique avec robot préparateur. Cah ORSTOM Ser Pedofil, 26: 371-380.

435 Gourdin, E. et al., 2015. Sources and export of particle-borne organic matter during a monsoon flood
436 in a catchment of northern Laos. Biogeosciences, 12(4): 1073-1089.
437 Huon, S. et al., 2013. Long-term soil carbon loss and accumulation in a catchment following the
438 conversion of forest to arable land in northern Laos. Agriculture, Ecosystems & Environment,
439 169: 43-57.
440 Kinoshita, N. et al., 2011. Assessment of individual radionuclide distributions from the Fukushima
441 nuclear accident covering central-east Japan. Proceedings of the National Academy of
442 Sciences of the United States of America, 108(49): 19526-9.
443 Kobayashi, S. et al., 2017. Low $^{134}Cs/^{137}Cs$ ratio anomaly in the north-northwest direction from the
444 Fukushima Dai-ichi Nuclear Power Station. Journal of environmental radioactivity, 178-179:
445 84-94.
446 Koiter, A.J., Owens, P.N., Petticrew, E.L. and Lobb, D.A., 2013. The behavioural characteristics of
447 sediment properties and their implications for sediment fingerprinting as an approach for
448 identifying sediment sources in river basins. Earth-Science Reviews, 125: 24-42.
449 Laceby, J.P. et al., 2016a. Rainfall erosivity in catchments contaminated with fallout from the
450 Fukushima Daiichi nuclear power plant accident. Hydrology and Earth System Sciences, 20(6):
451 2467-2482.
452 Laceby, J.P. et al., 2017. The challenges and opportunities of addressing particle size effects in
453 sediment source fingerprinting: A review. Earth-Science Reviews, 169: 85-103.
454 Laceby, J.P., Huon, S., Onda, Y. and Evrard, O., 2016b. Do forests represent a long-term source of
455 contaminated particulate matter in the Fukushima Prefecture? Journal of Environmental
456 Management, 183: 742–753.
457 Laceby, J.P., Huon, S., Onda, Y., Vaury, V. and Evrard, O., 2016c. Do forests represent a long-term
458 source of contaminated particulate matter in the Fukushima Prefecture? Journal of
459 environmental management, 183(Pt 3): 742-753.
460 Laceby, J.P. and Olley, J., 2015. An examination of geochemical modelling approaches to tracing
461 sediment sources incorporating distribution mixing and elemental correlations. Hydrological
462 Processes, 29: 1669-1685
463 Laceby, J.P., Olley, J., Pietsch, T.J., Sheldon, F. and Bunn, S.E., 2015. Identifying subsoil sediment
464 sources with carbon and nitrogen stable isotope ratios. Hydrological Processes, 29(8): 1956-
465 1971.
466 Le Gall, M. et al., 2017. Examining suspended sediment sources and dynamics during flood events in
467 a drained catchment using radiogenic strontium isotope ratios ($^{87}Sr/^{86}Sr$). Chemical Geology,
468 449: 147-157.
469 Legout, C. et al., 2013. Quantifying suspended sediment sources during runoff events in headwater
470 catchments using spectrocolorimetry. Journal of Soils and Sediments, 13(8): 1478-1492.
471 Lepage, H. et al., 2016. Investigating the source of radiocesium contaminated sediment in two
472 Fukushima coastal catchments with sediment tracing techniques. Anthropocene, 13: 57-68.
473 Mizugaki, S. et al., 2008. Estimation of suspended sediment sources using ^{137}Cs and $^{210}Pb_{ex}$ in
474 unmanaged Japanese cypress plantation watersheds in southern Japan. Hydrological
475 Processes, 22(23): 4519-4531.
476 Montgomery, D.R., 2007. Soil erosion and agricultural sustainability. Proceedings of the National
477 Academy of Sciences, 104(33): 13268-13272.
478 Owens, P.N. et al., 2005. Fine-grained sediment in river systems: environmental significance and
479 management issues. River Research and Applications, 21(7): 693-717.
480 Owens, P.N. et al., 2016. Fingerprinting and tracing the sources of soils and sediments: Earth and
481 ocean science, geoarchaeological, forensic, and human health applications. Earth-Science
482 Reviews, 162: 1-23.
483 Pansu, J. et al., 2015. Reconstructing long-term human impacts on plant communities: an ecological
484 approach based on lake sediment DNA. Molecular Ecology, 24(7): 1485-98.
485 Parducci, L. et al., 2017. Ancient plant DNA in lake sediments. New Phytologist, 214(3): 924-942.

Pedersen, M.W. et al., 2015. Ancient and modern environmental DNA. Philos Trans R Soc Lond B Biol Sci, 370(1660): 20130383.
Reiffarth, D.G., Petticrew, E.L., Owens, P.N. and Lobb, D.A., 2016. Sources of variability in fatty acid (FA) biomarkers in the application of compound-specific stable isotopes (CSSIs) to soil and sediment fingerprinting and tracing: A review. The Science of the total environment, 565: 8-27.
Sjogren, P. et al., 2016. Lake sedimentary DNA accurately records 20th Century introductions of exotic conifers in Scotland. New Phytol.
Taberlet, P., Bonin, A., Zinger, L. and Coissac, E., 2018. Environmental DNA: For Biodiversity Research and Monitoring. Oxford University Press.
Taberlet, P. et al., 2012. Soil sampling and isolation of extracellular DNA from large amount of starting material suitable for metabarcoding studies. Mol Ecol, 21(8): 1816-20.
Thomsen, P.F. et al., 2012. Monitoring endangered freshwater biodiversity using environmental DNA. Mol Ecol, 21(11): 2565-73.
Tiecher, T. et al., 2017. Quantifying land use contributions to suspended sediment in a large cultivated catchment of Southern Brazil (Guaporé River, Rio Grande do Sul). Agriculture, Ecosystems & Environment, 237: 95-108.
Walden, J., Slattery, M.C. and Burt, T.P., 1997. Use of mineral magnetic measurements to fingerprint suspended sediment sources: approaches and techniques for data analysis. Journal of Hydrology, 202(1-4): 353-372.
Wallbrink, P.J., Murray, A.S., Olley, J. and Olive, L.J., 1998. Determining sources and transit times of suspended sediment in the Murrumbidgee River, New South Wales, Australia, using fallout ^{137}Cs and ^{210}Pb. Water Resources Research, 4: 879-887.
Walling, D., Peart, M., Oldfield, F. and Thompson, R.J.N., 1979. Suspended sediment sources identified by magnetic measurements. 281(5727): 110.
Walling, D.E., 2013. The evolution of sediment source fingerprinting investigations in fluvial systems. Journal of Soils and Sediments, 13(10): 1658-1675.
Willerslev, E. et al., 2003. Diverse plant and animal genetic records from Holocene and Pleistocene sediments. Science, 300(5620): 791-5.
Yoccoz, N.G. et al., 2012. DNA from soil mirrors plant taxonomic and growth form diversity. Molecular Ecology, 21(15): 3647-55.
Yoshimura, K., Onda, Y. and Wakahara, T., 2016. Time Dependence of the ^{137}Cs Concentration in Particles Discharged from Rice Paddies to Freshwater Bodies after the Fukushima Daiichi NPP Accident. Environmental science & technology, 50(8): 4186-4193.

Figure captions

Figure 1. A) Location of the radioactive contamination plume and the selected Niida and Ota

catchments in the Fukushima Prefecture, Japan. B) The background map corresponds to the total radiocesium initial contamination of the soils (5-cm topsoil layer) decay-corrected to the 14 March, 2011 (Chartin et al., 2013). C) Sampling design in the Niida and Ota catchment along with land use.

Figure 2. Number of reads associated with a selection of plant types identified in the analysed samples collected in rivers draining cropland [C], forests [F] or a mix of cropland and forests [M].

Figure 3. Results of the Principal Component Analysis conducted to discriminate different groups of sediment samples based on the 67 significant MOTUs detected: group A – paddy field soil (in black); group AA – sediment collected in rivers draining cultivated land (in red); group B – sediment collected in rivers draining mixed land uses (in green); group C – sediment collected in rivers draining forests (in blue).

Figure 4. Results of the hierarchical classification conducted on the 67 significant MOTUs found in the analysed soil and sediment samples. Only those variables with cos > 0.3 were shown on the graph.

Figure 5. Example of sources delivering subsoil material to the rivers in the study site: (a) subsoil erosion in forests; (b) exposed channel bank in an un-channeled river section; (c) shallow landsliding on a roadside nearby a paddy field; (d) deeper landslide on a forested hillslope.

Table 1
Click here to download Table: Table1.docx

Table 1. Location and characteristics of the sediment sampling sites, concentrations in fallout radionuclides (^{137}Cs, ^{134}Cs) and associated ^{134}Cs/^{137}Cs activity ratio, organic matter properties in the soil/sediment samples.

Sample ID	Basin	Sub-basin	Type	Y	X	Sampling date	^{137}Cs (Bq kg^{-1})	^{134}Cs (Bq kg^{-1})	^{134}Cs/^{137}Cs	TN (%)	TOC (%)	TOC/TN	δ^{13}C (‰)	δ^{15}N (‰)
FOL1233	Ota	Ota	Cultivated	37.60223	140.992928	28/06/2017	875	860	0.98	-	0.6	-	-26.3	-
FOS1235	Ota	*Soil*	Paddy	37.603681	140.955704	28/06/2017	1385	1360	0.98	0.26	3.2	12.2	-23.8	5.4
FOL1234	Ota	Ota	Cultivated	37.603668	140.953856	28/06/2017	3210	3265	1.02	0.14	1.62	11.6	-25.9	4.4
FOL1236	Ota	Ota	Cultivated	37.597836	140.925567	28/06/2017	6335	6260	0.99	0.09	1.22	13.6	-27.4	3.9
FOL1237	Ota	Ota	Forest	37.556682	140.874831	29/06/2017	17380	17080	0.98	0.05	0.75	15.0	-27.6	3.4
FOL1238	Ota	Ota	Forest	37.564072	140.872601	29/06/2017	20530	20560	1.00	0.06	0.91	15.2	-28.0	2.4
FOL1239	Ota	Ota	Forest	37.578823	140.874961	29/06/2017	48730	48855	1.00	1.10	19.29	17.5	-29.3	-0.4
FNL1240	Niida	Mizunashi	Forest	37.627235	140.875257	29/06/2017	7105	7050	0.99	0.08	1.2	15.3	-28.8	1.7
FNL1241	Niida	Mizunashi	Cultivated	37.639792	140.934335	29/06/2017	3435	3465	1.01	0.11	1.2	11.3	-28.1	4.2
FNL1242	Niida	Niida	Mix	37.651152	140.958309	29/06/2017	1895	1870	0.99	0.06	0.7	11.3	-26.1	6.3
FNL1243	Niida	Niida	Mix	37.661327	140.911026	29/06/2017	980	965	0.98	-	0.3	-	-25.1	-
FNL1244	Niida	Niida	Cultivated	37.664088	140.899675	29/06/2017	960	935	0.97	0.06	0.6	10.7	-27.0	4.5
FNL1245	Niida	Niida	Mix	37.651603	140.889993	29/06/2017	6845	6535	0.95	0.35	4.5	12.7	-26.8	3.3

Table 2
Click here to download Table: Table 2.docx

Table 2: Dilution model results for the Niida catchment including the Cs-137 activities and modeled contribution from upstream areas that were contaminated by fallout by the FDNPP accident

Sample ID	Basin	Sub-basin	Type	Cs-137 Bq kg^{-1}	Error Bq kg^{-1}	Contam. Source (%)	Error (%)
FNL1240	Niida	Mizunashi	Forest	7104	25	14.0%	1.2%
FNL1241	Niida	Mizunashi	Cultivated	3435	18	4.6%	1.0%
FNL1242	Niida	Niida	Mix	1893	15	1.4%	1.0%
FNL1243	Niida	Niida	Mix	977	9	0.0%	1.0%
FNL1244	Niida	Niida	Cultivated	957	9	0.0%	1.0%
FNL1245	Niida	Niida	Mix	6843	41	13.7%	1.2%

Table 3
Click here to download Table: Table 3.docx

Table 3: Results from the Carbon and nitrogen parameter model for the Niida and the Ota catchments including the Cs-137 activities and modeled contribution from upstream areas that were contaminated by fallout by the FDNPP accident

Sample ID	Basin	Sub-basin	Type	Forest (%)	ME (%)	Crop (%)	ME (%)	Subsoil (%)	ME (%)
FOS1235	Ota	*Soil*	Paddy	4%	1%	64%	2%	32%	3%
FOL1234	Ota	Ota	Cultivated	2%	2%	21%	3%	76%	10%
FOL1236	Ota	Ota	Cultivated	3%	2%	8%	12%	89%	20%
FOL1237	Ota	Ota	Forest	0%	1%	0%	5%	100%	15%
FOL1238	Ota	Ota	Forest	1%	3%	0%	5%	98%	17%
FOL1239	Ota	Ota	Forest	100%	1%	0%	1%	0%	1%
FNL1240	Niida	Mizunashi	Forest	6%	3%	1%	3%	93%	15%
FNL1241	Niida	Mizunashi	Cultivated	1%	2%	16%	3%	82%	16%
FNL1242	Niida	Niida	Mix	0%	1%	0%	4%	100%	9%
FNL1244	Niida	Niida	Cultivated	0%	1%	4%	2%	96%	15%
FNL1245	Niida	Niida	Mix	18%	1%	44%	3%	38%	16%

Figure 1

Click here to download high resolution image




Figure 2

Click here to download high resolution image




Figure 3

Click here to download high resolution image




Figure 4

Click here to download high resolution image





Figure 5

Click here to download high resolution image



Supplementary material for on-line publication only

Click here to download Supplementary material for on-line publication only: Supplementary-Material.docx

Supplementary material for on-line publication only

Click here to download Supplementary material for on-line publication only: AppendixA.xlsx