



HAL
open science

Reconstruction and stable recovery of source terms appearing in diffusion equations

Yavar Kian, Masahiro Yamamoto

► **To cite this version:**

Yavar Kian, Masahiro Yamamoto. Reconstruction and stable recovery of source terms appearing in diffusion equations. *Inverse Problems*, 2019, 35 (11), pp.115006. 10.1088/1361-6420/ab2d42 . hal-02049174

HAL Id: hal-02049174

<https://hal.science/hal-02049174>

Submitted on 26 Feb 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

RECONSTRUCTION AND STABLE RECOVERY OF SOURCE TERMS APPEARING IN DIFFUSION EQUATIONS

YAVAR KIAN AND MASAHIRO YAMAMOTO

ABSTRACT. We consider the inverse source problem of determining a source term depending on both time and space variable for fractional and classical diffusion equations in a cylindrical domain from boundary measurements. With suitable boundary conditions we prove that some class of source terms which are independent of one space direction, can be reconstructed from boundary measurements. Actually, we prove that this inverse problem is well-posed. We establish also some results of Lipschitz stability for the recovery of source terms which we apply to the stable recovery of time-dependent coefficients.

Keywords: Inverse source problems, fractional diffusion equation, reconstruction, well-posedness, stability estimate.

Mathematics subject classification 2010 : 35R30, 35R11.

1. INTRODUCTION

1.1. Statement. Let $d \geq 2$, $\Omega = \omega \times (-\ell, \ell)$ and $\tilde{\Omega} = \omega \times (0, \ell)$, and $\omega \subset \mathbb{R}^{d-1}$ be a bounded domain with \mathcal{C}^2 boundary. We set $Q = (0, T) \times \Omega$ and $\tilde{Q} = (0, T) \times \tilde{\Omega}$. Let $\nu = \nu(x)$ be the outward unit normal vector to $\partial\Omega$ or $\partial\tilde{\Omega}$ at x . In what follows, we define \mathcal{A} by the differential operator

$$\mathcal{A}u(x) = - \sum_{i,j=1}^d \partial_{x_i} (a_{ij}(x) \partial_{x_j} u), \quad x \in \Omega,$$

where $a_{ij} = a_{ji} \in \mathcal{C}^2(\bar{\omega} \times [-\ell, \ell])$, $1 \leq i, j \leq d$, satisfy

$$\begin{aligned} \sum_{i,j=1}^d a_{ij}(x) \xi_i \xi_j &\geq c |\xi|^2, \quad x \in \bar{\Omega}, \quad \xi = (\xi_1, \dots, \xi_d) \in \mathbb{R}^d, \\ a_{jd}(x', \pm\ell) &= 0, \quad a_{dd}(x', \pm\ell) > 0, \quad x' \in \bar{\omega}, \quad j = 1, \dots, d-1. \end{aligned} \tag{1.1}$$

For $\alpha \in (0, 2) \setminus \{1\}$, we denote by ∂_t^α the Caputo fractional derivative with respect to t given by

$$\partial_t^\alpha u(t, x) := \frac{1}{\Gamma([\alpha] + 1 - \alpha)} \int_0^t (t-s)^{[\alpha]-\alpha} \partial_s^{[\alpha]+1} u(s, x) ds, \quad (t, x) \in Q,$$

and by ∂_t^1 the usual derivative in t . We set

$$m_\alpha = \begin{cases} 0, & 0 < \alpha \leq 1, \\ 1, & 1 < \alpha < 2, \end{cases}$$

$T \in (0, +\infty)$, $\Sigma = (0, T) \times \partial\Omega$, $0 < \alpha < 2$ and we consider the following problem

$$\begin{cases} \partial_t^\alpha u + \mathcal{A}u = F(t, x), & (t, x', x_d) = (t, x) \in Q, \\ \partial_t^k u(0, x) = 0, & x \in \Omega, k = 0, \dots, m_\alpha. \end{cases} \quad (1.2)$$

We associate with this problem the following boundary conditions

$$\partial_{x_d}^m u(t, x', \pm\ell) = 0, \quad (t, x') \in (0, T) \times \omega, \quad m = 0, 1, \quad (1.3)$$

$$\partial_\nu^n u(t, x) = 0, \quad (t, x) \in (0, T) \times \partial\omega \times (-\ell, \ell), \quad n = 0, 1. \quad (1.4)$$

In the same way, we consider the problem

$$\begin{cases} \partial_t^\alpha u + \mathcal{A}u + q(t, x')u = F(t, x), & (t, x', x_d) = (t, x) \in \tilde{Q}, \\ \partial_t^k u(0, x) = 0, & x \in \tilde{\Omega}, k = 0, m_\alpha. \end{cases} \quad (1.5)$$

$$\partial_{x_d} u(t, x', 0) = u(t, x', \ell) = 0, \quad (t, x') \in (0, T) \times \omega, \quad (1.6)$$

$$u(t, x) = 0, \quad (t, x) \in (0, T) \times \partial\omega \times (0, \ell). \quad (1.7)$$

In the present paper, we treat the inverse problem of determining the source term F or the coefficient q from measurements of the solution of (1.2)-(1.4) or (1.5)-(1.7) on a subboundary of the cylindrical domain Q or \tilde{Q} .

1.2. Obstruction against the uniqueness. We recall that there is an obstruction against the recovery of general source terms F from any type of measurements of the solution of (1.2)-(1.4) (resp. (1.5)-(1.7)) on $(0, T) \times \partial\Omega$ (resp. $(0, T) \times \partial\tilde{\Omega}$). Indeed, choose $\chi \neq 0$, in $C_0^\infty(Q)$ and consider $F := \partial_t^\alpha \chi + \mathcal{A}\chi$. From the uniqueness of the weak solution of problem (1.2)-(1.4) (see Section 1.4 for more details and see Definition 1.1 below for the definition of weak solutions), we know that $u = \chi$ and, since $\chi \neq 0$, we deduce that $F \neq 0$. However, we have

$$\partial_\nu^k u(t, x) = \partial_\nu^k \chi(t, x) = 0, \quad (t, x) \in (0, T) \times \partial\Omega, \quad k = 0, 1, \dots,$$

with ν the outward unit normal vector of $\partial\Omega$.

Facing this obstruction against the uniqueness, we will consider source terms of the form

$$F(t, x', x_d) := f(t, x')R(t, x', x_d), \quad (t, x', x_d) \in (0, T) \times \omega \times (-\ell, \ell) \quad (1.8)$$

and, assuming that R is known, we will consider the problem of determining f .

1.3. Motivations. We recall that the problem (1.2)-(1.4) (resp. (1.5)-(1.7)) is associated with different models of diffusion. The non-integer value of the power α , is frequently used for describing anomalous diffusion derived from continuous-time random walk models (see e.g., [27]). In this context the recovery of the source term f can be seen as the recovery of a time evolving source of diffusion. For instance, in the case $\alpha = 1$ and $d = 3$, our problem can be associated with the recovery of a source moving in the subset ω of the plan \mathbb{R}^2 from a single measurement of the heat flux at the boundary. Such a problem can be associated with the determination of different properties of materials such as metal (see e.g., [22] for the heat equation). We refer to [11] for applications of recovery of source terms of the form (1.8) to the recovery of moving sources in the electrodynamics. For non-integer value of the power α , in the spirit of [28] (see also [16]), our inverse problem can be seen as the recovery of a moving source of diffusion of a contaminant under the ground. As unknown sources, we assume the form of (1.8), which can be interpreted for example that an unknown source $f(x', t)$ depends only on the depth variable and t in the case of $d = 2$, which corresponds to a layer structure, and on the planar locations (x_1, x_2) and t but not on the depth in the case of $d = 3$, which may be a good approximation if Ω is a very thin domain in the direction of x_3 .

1.4. Known results. Inverse source problems have received a lot of attention these last decades among the mathematical community (see [13] for an overview). For diffusion equations corresponding to the case $\alpha = 1$ with time independent source terms, several authors investigated the conditional stability (e.g. [5, 35, 36]). Following the Bukhgeim-Klibanov approach introduced in [3], [12] established Lipschitz stable recovery of the source from one Neumann boundary measurement. In [6], the authors derived a stability estimate for this problem from a single Neumann observation of the solution on an arbitrary portion of the boundary. This last result has been extended to an infinite cylindrical domain by [18]. For fractional diffusion equations corresponding to the case $\alpha \neq 1$, in spite of the physical backgrounds related to various anomalous diffusion phenomena stated above, to our best knowledge, there is no result in the mathematical literature dealing with the recovery of source terms, depending on both time and space variables, of the form (1.8). We mention also the work of [14, 15, 16, 17, 19, 23, 31] where some inverse coefficient problems and some related results have been considered.

The above mentioned results are all concerned with the determination of time-independent source terms $f(x)$. Several authors considered also the recovery of time-dependent source terms. In [7, 30] the authors proved the stable recovery of a source term $f(t)$ depending only on the time variable from measurements of solutions at one spatial point over time interval. As long as the classical partial differential equations with natural number α are concerned, some papers have also been devoted to the unique existence and the stability in finding source terms of the form (1.8) (see e.g. [2, 10, 13]). In particular, in [13, Section 6.3] the author proved the reconstruction and the unique recovery of source terms of the form (1.8) appearing in a parabolic equation on the half space.

Our main results stated below, seem the first achievements for the inverse source problem of determining $f(t, x')$ in (1.8) for fractional partial differential equations.

1.5. Preliminary properties. In the present paper, following [20, 30], we consider solutions of problem (1.2)-(1.6) (resp. (1.5)-(1.7)) in the following weak sense.

Definition 1.1. *Let $F \in L^2(Q)$ (resp. $F \in L^2((0, T) \times \tilde{\Omega})$). We say that problem (1.2)-(1.4) (resp. (1.5)-(1.7) with $q = 0$) admits a weak solution u if there exists $v \in L^2_{loc}(\mathbb{R}^+; L^2(\Omega))$ such that:*

1) $v|_Q = u$ and $\inf\{\varepsilon > 0 : e^{-\varepsilon t} v \in L^1(\mathbb{R}^+; L^2(\Omega))\} = 0$,

2) for all $p > 0$ the Laplace transform $V(p) = \int_0^{+\infty} e^{-pt} v(t, \cdot) dt$ with respect to t of v , satisfies (1.3)-(1.4) (resp. (1.6)-(1.7)) and solves

$$(\mathcal{A} + p^\alpha)V(p) = \hat{F}(p) \quad \text{in } \Omega,$$

where $\hat{F}(p) = \mathcal{L}[F(t, \cdot)\mathbf{1}_{(0, T)}(t)](p) = \int_0^T e^{-pt} F(t, \cdot) dt$ and $\mathbf{1}_{(0, T)}$ is the characteristic function of $(0, T)$.

By the results in [24, 30], we can prove that for $F \in L^2(Q)$ problem (1.2)-(1.4) (resp. (1.5)-(1.7)) admits a unique solution $u \in L^2(0, T; H^1(\Omega))$ satisfying $\partial_t^\alpha u, \mathcal{A}u \in L^2(Q)$. For sake of completeness we recall this result in the Appendix.

In (1.5), we assume that $a_{ij} \in C^3(\bar{\Omega})$, $i, j \in \{1, \dots, d\}$, satisfy the following condition

$$a_{ij}(x', -x_d) = a_{ij}(x', x_d), \quad (x', x_d) \in \bar{\Omega}, \quad i, j \in \{1, \dots, d\}. \quad (1.9)$$

Then we consider the problem

$$\left\{ \begin{array}{l} \partial_t^\alpha u + \mathcal{A}u = F(t, x), \quad (t, x', x_d) \in (0, T) \times \tilde{\Omega}, \\ u(t, x) = 0, \quad (t, x) \in (0, T) \times [\partial\omega \times (0, \ell) \cup \omega \times \{\ell\}], \\ -\partial_{x_d} u(t, x', 0) = 0 \quad (t, x') \in (0, T) \times \omega, \\ \partial_t^k u(0, x) = 0, \quad x \in \Omega, \quad k = 0, m_\alpha. \end{array} \right. \quad (1.10)$$

Here we consider weak solutions in the sense of Definition 1.1 and, following [20, 30], we can prove that there exists an operator valued function $J(t) \in \mathcal{B}(L^2(\tilde{\Omega}))$ such that the solution of (1.10) takes the form

$$u(t, \cdot) = \int_0^t J(t-s)F(s, \cdot) ds.$$

Using this definition, for $q \in L^\infty((0, T) \times \omega)$ we can define the solution of

$$\left\{ \begin{array}{l} \partial_t^\alpha u + \mathcal{A}u + q(t, x')u = F(t, x), \quad (t, x', x_n) \in (0, T) \times \tilde{\Omega}, \\ u(t, x) = 0, \quad (t, x) \in (0, T) \times [\partial\omega \times (0, \ell) \cup \omega \times \{\ell\}], \\ -\partial_{x_d} u(t, x', 0) = 0 \quad (t, x') \in (0, T) \times \omega, \\ \partial_t^k u(0, x) = 0, \quad x \in \Omega, \quad k = 0, \dots, m_\alpha, \end{array} \right. \quad (1.11)$$

in the mild sense which is defined by the integral equation

$$u(t, \cdot) = - \int_0^t J(t-s)qu(s, \cdot)ds + \int_0^t J(t-s)F(s, \cdot)ds.$$

1.6. Main results. From now on, we assume that F takes the form (1.8). For our first result we need an assumption on ω and \mathcal{A} that guarantees the elliptic regularity of the operator \mathcal{A} . Indeed, due to the fact that the domain Ω is only Lipschitz, some extra assumptions will be required for guaranteeing the elliptic regularity of \mathcal{A} with the boundary conditions (1.3)-(1.4). For this purpose, for $m, n = 0, 1$, we introduce the condition:

(Hmn) For all $v \in H^1(\Omega)$ satisfying $\mathcal{A}v \in L^2(\Omega)$ and (1.3)-(1.4) with these value of m, n , we have $v \in H^2(\Omega)$ and there exists $C > 0$ depending only on \mathcal{A} and Ω such that

$$\|v\|_{H^2(\Omega)} \leq C(\|\mathcal{A}v\|_{L^2(\Omega)} + \|v\|_{L^2(\Omega)}).$$

Note that conditions (H00) and (H11) will be fulfilled if, for instance, we assume that ω is convex. Indeed, in that case Ω will also be convex and, in virtue of [9, Theorem 3.2.1.2] and [9, Theorem 3.2.1.3], (H00) and (H11) will be fulfilled. In the same way, assuming that

$$a_{id} = 0, \quad \partial_{x_j} a_{dd} = 0, \quad \partial_{x_a} a_{ij} = 0, \quad i, j \in \{1, \dots, d-1\}, \quad (1.12)$$

we deduce from a separation of variable argument similar to [8, Lemma 2.4] that, for all $m, n = 0, 1$, (Hmn) is fulfilled.

Using the conditions (Hmn), we obtain the following.

Theorem 1.2. *Let (H00), (H10) be fulfilled and assume that $R, \partial_{x_a} R \in L^\infty(Q)$ and there exists a constant $c > 0$ such that*

$$|R(t, x', \ell)| \geq c, \quad (t, x') \in (0, T) \times \omega. \quad (1.13)$$

Assume also that the condition

$$\partial_{x_a} a_{ij} = 0, \quad i, j = 1, \dots, d-1, \quad (1.14)$$

is fulfilled. Then, for $f \in L^2((0, T) \times \omega)$, the solution u of (1.2)-(1.4) with $m = 1$ and $n = 0$ satisfies $u \in H^3(-\ell, \ell; L^2((0, T) \times \omega))$, $\partial_t^\alpha u, \mathcal{A}u \in H^1(-\ell, \ell; L^2((0, T) \times \omega))$. Therefore, we can define

$$h := (t, x') \mapsto \frac{[\partial_t^\alpha u + (\mathcal{A} + a_{dd}\partial_{x_a}^2)u](t, x', \ell)}{R(t, x', \ell)} \in L^2((0, T) \times \omega). \quad (1.15)$$

Moreover, for every $f \in L^2((0, T) \times \omega)$, we can define an operator valued function $K(t, s) \in \mathcal{B}(L^2(\omega))$, $t, s \in (0, T)$, such that f solves the integral equation

$$f(t, \cdot) = h(t, \cdot) + \int_0^t K(t, s)f(s, \cdot)ds, \quad t \in (0, T), \quad (1.16)$$

which is well-posed. Finally, for every $(h, f) \in L^2((0, T) \times \omega) \times L^2((0, T) \times \omega)$ satisfying (1.16) the solution u of (1.2)-(1.4) satisfies (1.15). In the same way, assuming that (H01) and (H11) are fulfilled, the same results hold true for the problem (1.2)-(1.4) with $m = 1$ and $n = 1$.

For problem (1.11), we consider first the following condition:

(\tilde{H}) For all $v \in H^{\max(1, s)}(\tilde{\Omega})$ satisfying $\mathcal{A}v \in H^s(\tilde{\Omega})$, $s \in [0, 2]$, and (1.6)-(1.7), we have $v \in H^{2+s}(\Omega)$ and there exists $C > 0$ depending only on \mathcal{A} , s and Ω such that

$$\|v\|_{H^{2+s}(\Omega)} \leq C(\|\mathcal{A}v\|_{H^s(\Omega)} + \|v\|_{H^s(\Omega)}).$$

Assuming that ω is of class \mathcal{C}^4 and using a separation of variable argument similar to [8, Lemma 2.4], one can check that (1.12) implies (\tilde{H}).

Using (\tilde{H}) we obtain the following well-posedness result.

Proposition 1.3. *Assume that (\tilde{H}) is fulfilled. Let $\gamma \in (0, 1)$ be such that for $\alpha \in (0, 1]$, $\gamma \in (1/2, 1)$ and for $\alpha \in (1, 2)$, $\gamma \in (1/2, 1/\alpha)$. Fix $p \in (1, +\infty)$ such that $\frac{1}{p} < \min(1 - \alpha\gamma, \alpha(1 - \gamma))$ and let $F \in W^{1,p}(0, T; L^2(\tilde{\Omega})) \cap \mathcal{C}([0, T]; H^{2\gamma}(\tilde{\Omega}))$ satisfy $F|_{t=0} = 0$. Let $q \in \mathcal{C}^1([0, T]; L^\infty(\omega)) \cap \mathcal{C}([0, T]; W^{2,\infty}(\omega))$. Then problem (1.11) admits a unique weak solution $u \in \mathcal{C}([0, T]; H^{2(1+\gamma)}(\tilde{\Omega})) \cap \mathcal{C}^1([0, T]; H^\gamma(\tilde{\Omega}))$.*

Applying this well-posedness result, we can state our second main result as follows.

Theorem 1.4. *Assume that (\tilde{H}), (H00), (1.9) and (1.14) are fulfilled, $\alpha \in (0, 1]$, ω is of \mathcal{C}^4 and $\gamma \in (3/4, 1)$. Fix $p \in (1, +\infty)$ such that $\frac{1}{p} < \min(1 - \alpha\gamma, \alpha(1 - \gamma))$. Let $q \in \mathcal{C}^1([0, T]; L^\infty(\omega)) \cap \mathcal{C}([0, T]; W^{2,\infty}(\omega))$, $f \in W^{1,p}(0, T; L^\infty(\omega)) \cap \mathcal{C}([0, T]; W^{2,\infty}(\omega))$ and $R \in W^{1,\infty}(0, T; L^2(\tilde{\Omega})) \cap \mathcal{C}([0, T]; H^{2\gamma}(\tilde{\Omega}))$ satisfy $R, \partial_{x_d} R \in L^\infty((0, T) \times \tilde{\Omega})$ and (1.13). Assume also that*

$$f(0, x) = 0, \quad x \in \Omega.$$

Then, for $\delta \in (0, 2\gamma - \frac{3}{4})$, there exists a constant C depending on $q, R, \tilde{\Omega}, T, \alpha, \mathcal{A}, \delta$, such that

$$\|f\|_{L^\infty(0, T; L^2(\omega))} \leq C(\|\partial_{x_d} u(\cdot, \cdot, \ell)\|_{L^\infty(0, T; H^{\frac{3}{2}}(\omega))} + \|\partial_{x_d} u(\cdot, \cdot, \ell)\|_{W^{1,\infty}(0, T; H^\delta(\omega))}). \quad (1.17)$$

Applying this result, we can also prove the stable recovery of the coefficient q appearing in the problem

$$\left\{ \begin{array}{l} \partial_t^\alpha v + \mathcal{A}v + q(t, x')v = 0, \quad (t, x) \in (0, T) \times \tilde{\Omega}, \\ v(t, x', \ell) = h_0(t, x') \quad (t, x') \in (0, T) \times \omega, \\ v(t, x) = h_1(t, x), \quad (t, x) \in (0, T) \times \partial\omega \times (-\ell, \ell), \\ \partial_{x_d} v(t, x', -\ell) = 0 \quad (t, x') \in (0, T) \times \omega, \\ v(0, x) = w_0, \quad x \in \tilde{\Omega} \end{array} \right. \quad (1.18)$$

with $\alpha \in (0, 1]$, h_k , $k = 0, 1$, w_0 such that there exists $H \in \mathcal{C}^1([0, T]; H^{2+\gamma}(\tilde{\Omega})) \cap W^{2,p}(0, T; H^2(\tilde{\Omega}))$ satisfying

$$\left\{ \begin{array}{l} H(t, x', \ell) = h_0(t, x') \quad (t, x') \in (0, T) \times \omega, \\ H(t, x) = h_1(t, x), \quad (t, x) \in (0, T) \times \partial\omega \times (0, \ell), \\ \partial_{x_d} H(t, x', 0) = H(t, x', 0) = 0 \quad (t, x') \in (0, T) \times \omega, \\ H(0, x) = w_0(x), \quad x \in \tilde{\Omega}, \\ \partial_t^\alpha H(0, x) = -\mathcal{A}H(0, x) - q(0, x')H(0, x), \quad x = (x', x_d) \in \tilde{\Omega}. \end{array} \right. \quad (1.19)$$

The result for the determination of q can be stated as follows.

Corollary 1.5. *Let the condition of Theorem 1.4 be fulfilled with $f = 0$, $d \leq 3$ and let h_k , $k = 0, 1$, w_0 be given by (1.19) and the condition*

$$|h_0(t, x')| \geq c, \quad (t, x') \in (0, T) \times \omega \quad (1.20)$$

be fulfilled. Fix $q_j \in \mathcal{C}^1([0, T]; L^\infty(\omega)) \cap \mathcal{C}([0, T]; W^{2,\infty}(\omega))$, $j = 1, 2$, such that

$$\|q_j\|_{W^{1,\infty}(0,T;L^\infty(\omega))} + \|q_j\|_{L^\infty(0,T;W^{2,\infty}(\omega))} \leq M,$$

$$q_1(0, x') = q_2(0, x') = q(0, x'), \quad x' \in \omega$$

and consider the solution v_j of (1.18) with $q = q_j$. Then, for $\delta \in (0, 2\gamma - \frac{3}{4})$, we have

$$\|q_1 - q_2\|_{L^\infty(0,T;L^2(\omega))} \leq C \left(\|\partial_{x_d}(v_1 - v_2)|_{x_d=\ell}\|_{L^\infty(0,T;H^{\frac{3}{2}}(\omega))} + \|\partial_{x_d}(v_1 - v_2)|_{x_d=\ell}\|_{W^{1,\infty}(0,T;H^\delta(\omega))} \right), \quad (1.21)$$

where the constant C depends on α , c , T , Ω , a , \mathcal{A} , α , M , h_0 , h_1 , w_0 , δ .

1.7. Comments about our results. To our best knowledge Theorem 1.2 and 1.4 are the first results of recovery of a source term depending on both time and space variables for fractional diffusion equations of the form (1.2) when $\alpha \neq 1$. For $\alpha = 1$, we refer to [13, Section 6.3] addressing this inverse problem with Ω corresponding to the half space and see also [2]. In contrast to [13], we state our result on a bounded cylindrical domain and we restrict our analysis to solutions lying in Sobolev spaces while [13, Section 6.3] is stated with Hölder continuous functions. Moreover our approach admits a natural extension to fractional diffusion equations ($\alpha \neq 1$).

Let us remark that Theorem 1.2 gives a reconstruction algorithm for the recovery of the source term f under consideration. It is actually stated as a well-posedness result for the pair of functions (u, f) appearing in (1.2) and (1.8). In contrast to Theorem 1.2, Theorem 1.4 provides only a stability estimate. However, Theorem 1.4 can be applied to more general boundary conditions and it can also be applied to the stable recovery of a coefficient depending on both time and space variables (see Corollary 1.5).

Applying Theorem 1.4, we prove in Corollary 1.5 the stable recovery of the coefficient of order zero q provided $\partial_{x_d} q = 0$. It seems that this result is the first result of stable recovery of a coefficient depending on both time and space variables for a fractional diffusion equation. In [20, Theorem 3.6], the authors derived a similar result for the heat equation ($\alpha = 1$) stated with stronger regularity conditions and measurements on both $\gamma \times (0, \ell)$ and $\omega \times \{\ell_1\}$, with $\gamma \subset \partial\omega$ and $\ell_1 \in (0, \ell]$. Even, for $\alpha = 1$, our result improves the one of [20, Theorem 3.6] in terms of regularity conditions and restriction of the data.

Theorem 1.4 is stated only for $\alpha \in (0, 1]$ but it can be extended to $\alpha \in (0, 2)$ without any difficulty. Indeed, one can easily extend our argumentation to the case $\alpha \in (1, 2)$. In Theorem 1.4 we have restricted our analysis to $\alpha \in (0, 1]$ in order to simplify the statement of this theorem and its proof.

1.8. Outline. This paper is organized as follows. In Section 2 we prove Theorem 1.2. Section 3 is devoted to the proof of Theorem 1.4 and in Section 4 we consider the application of Theorem 1.4 stated in Corollary 1.5. Finally, in the Appendix we recall and prove a result related to regularity of solutions of (1.2).

2. PROOF OF THEOREM 1.2

We start with considering the first part of Theorem 1.2. For this purpose, we assume that (H00), (H10), (1.13)-(1.14) are fulfilled and we will show (1.16). We denote by A (resp. \tilde{A}) the operator \mathcal{A} acting on $L^2(\Omega)$ with domain

$$D(A) := \{g \in L^2(\Omega) : \mathcal{A}g \in L^2(\Omega), u|_{\partial\Omega} = 0\}$$

$$\text{(resp. } D(\tilde{A}) := \{g \in L^2(\Omega) : \mathcal{A}g \in L^2(\Omega), u|_{\partial\omega \times (0, \ell)} = 0, \partial_{x_d} u|_{x_d = \pm \ell} = 0\}$$

Thanks to (1.1) we know that A (resp. \tilde{A}) are selfadjoint operators with a spectrum consisting of a non-decreasing sequence of non-negative eigenvalues $(\lambda_n)_{n \geq 1}$ (resp. $(\tilde{\lambda}_n)_{n \geq 1}$). Moreover, conditions (H00) and (H10) imply that $D(A)$ and $D(\tilde{A})$ embedded continuously into $H^2(\Omega)$. Let us also introduce an orthonormal basis in the Hilbert space $L^2(\Omega)$ of eigenfunctions $(\varphi_n)_{n \geq 1}$ (resp. $(\tilde{\varphi}_n)_{n \geq 1}$) of A (resp. \tilde{A}) associated to the non-decreasing sequence of eigenvalues $(\lambda_n)_{n \geq 1}$ (resp. $(\tilde{\lambda}_n)_{n \geq 1}$). We consider also the operator valued function $S(t)$ (resp. $\tilde{S}(t)$) defined by

$$S(t)h = \sum_{n=1}^{\infty} t^{\alpha-1} E_{\alpha, \alpha}(-\lambda_n t^\alpha) \langle h, \varphi_n \rangle_{L^2(\Omega)} \varphi_n, \quad h \in L^2(\Omega),$$

$$\tilde{S}(t)h = \sum_{n=1}^{\infty} t^{\alpha-1} E_{\alpha, \alpha}(-\tilde{\lambda}_n t^\alpha) \langle h, \tilde{\varphi}_n \rangle_{L^2(\Omega)} \tilde{\varphi}_n, \quad h \in L^2(\Omega),$$

where $E_{\alpha,\alpha}$ corresponds to the Mittag-Leffler function given by

$$E_{\alpha,\alpha}(z) = \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(\alpha(n+1))}, \quad z \in \mathbb{C}.$$

Following [20, 30] (see also Lemma 3.1), one can check that problem (1.2) admits a unique weak solution $u \in L^2(0, T; D(\tilde{A}))$ taking the form

$$u(t, \cdot) = \int_0^t \tilde{S}(t-s)F(s, \cdot)ds, \quad t \in (0, T).$$

Recall that the function v given by Definition 1.1 takes the form

$$v(t, \cdot) = \int_0^{\min(t, T)} \tilde{S}(t-s)F(s, \cdot)ds, \quad t \in (0, +\infty).$$

Moreover, using the fact that $D(\tilde{A})$ embedded continuously into $H^2(\Omega)$, we deduce by interpolation that for $s_1 \in (\frac{3}{4}, 1)$, $D(\tilde{A}^{s_1})$ embedded continuously into $H^{2s_1}(\Omega)$. Therefore, applying formula (1.148) of [29, Theorem 1.6], one can check that $\inf\{\varepsilon > 0 : e^{-\varepsilon t}v \in L^1(\mathbb{R}^+; H^{2s_1}(\Omega))\} = 0$. Fixing $w = \partial_{x_d}u$ we deduce that, for all $p > 0$, the Laplace transform in time $W(p)$ of the extension of w to $\mathbb{R}_+ \times \Omega$ given by $w = \partial_{x_d}v$, with v defined in Definition 1.1, is lying in $H^{2s_1-1}(\Omega)$ and it satisfies

$$\begin{cases} (\mathcal{A} + p^\alpha)W(p) = -(\partial_{x_d}\mathcal{A})V(p) + \partial_{x_d}\hat{F}(p), & \text{in } \Omega, \\ W(p) = 0, & \text{on } \partial\Omega. \end{cases}$$

Note that here we use the fact that $W(p) \in H^{2s_1-1}(\Omega)$, with $2s_1 - 1 > \frac{1}{2}$, for defining its trace on $\partial\Omega$. Moreover, applying formula (1.148) of [29, Theorem 1.6] and the fact that $D(A^{1/2}) = H_0^1(\Omega)$, we can extend $S(t)$, $t > 0$, to a bounded operator from $H^{-1}(\Omega)$ to $L^2(\Omega)$ satisfying

$$\|S(t)\|_{\mathcal{B}(H^{-1}(\Omega), L^2(\Omega))} \leq Ct^{\alpha/2-1}.$$

In addition, using the fact that $v \in L_{loc}^2(\mathbb{R}^+; D(\mathcal{A})) \subset L_{loc}^2(\mathbb{R}^+; H^2(\Omega))$, we deduce that $(\partial_{x_d}\mathcal{A})v \in L_{loc}^2(\mathbb{R}^+; L^2(\Omega))$. Thus, extending F by zero to $(0, +\infty) \times \Omega$ and using the fact that

$$\inf\{\varepsilon > 0 : e^{-\varepsilon t}(\partial_{x_d}\mathcal{A})v \in L^1(\mathbb{R}^+; H^{-1}(\Omega))\} = 0,$$

we deduce that the function

$$\begin{aligned} w_1(t, \cdot) &:= \int_0^t S(t-s)[(\partial_{x_d}\mathcal{A})v(s, \cdot) + \partial_{x_d}F(s, \cdot)]ds \\ &= \int_0^t S(t-s_1)(\partial_{x_d}\mathcal{A}) \left(\int_0^{s_1} \tilde{S}(s_1-s_2)f(s_2, \cdot)R(s_2, \cdot)ds_2 \right) ds_1 + \int_0^t S(t-s)f(s_2, \cdot)\partial_{x_d}R(s_2, \cdot)ds. \end{aligned}$$

is well defined and the Laplace transform in time of w_1 coincide with the one of w . This proves that

$$\partial_{x_d}u = - \int_0^t S(t-s_1)(\partial_{x_d}\mathcal{A}) \left(\int_0^{s_1} \tilde{S}(s_1-s_2)f(s_2, \cdot)R(s_2, \cdot)ds_2 \right) ds_1 + \int_0^t S(t-s)f(s_2, \cdot)\partial_{x_d}R(s_2, \cdot)ds. \quad (2.1)$$

With the additional assumptions (H10), (H11) and the fact that ω is C^2 one can extend these arguments to problem (1.2)-(1.4), with $m = n = 1$, by using the fact that, for any $s_1 \in (0, 1/2)$, $C_0^\infty(\Omega)$ is dense in $H^{s_1}(\Omega)$.

In view of (1.14), for $w = \partial_{x_d} u$, we have

$$-(\partial_{x_d} \mathcal{A})u = \partial_{x_d}(\partial_{x_d} a_{dd} \partial_{x_d} u) + \sum_{j=1}^{d-1} [\partial_{x_j}(\partial_{x_d} a_{jd} \partial_{x_d} u) + \partial_{x_d}(\partial_{x_d} a_{jd} \partial_{x_j} u)] = B_1 w + B_2 u, \quad (2.2)$$

where

$$B_1 w := \partial_{x_d} a_{dd} \partial_{x_d} w + 2 \sum_{j=1}^{d-1} \partial_{x_d} a_{jd} \partial_{x_j} w + \left(\sum_{j=1}^d \partial_{x_j} \partial_{x_d} a_{jd} \right) w, \quad (2.3)$$

$$B_2 u = \sum_{j=1}^d (\partial_{x_d}^2 a_{jd}) \partial_{x_j} u. \quad (2.4)$$

Then, from (2.1) and the above arguments, we deduce that w solves the integral equation

$$\begin{aligned} w(t, \cdot) &= \int_0^t S(t-s) B_1 w(s, \cdot) ds + \int_0^t S(t-s_1) B_2 \left(\int_0^{s_1} \tilde{S}(s_1-s_2) f(s_2, \cdot) R(s_2, \cdot) ds_2 \right) ds_1 \\ &\quad + \int_0^t S(t-s) f(s, \cdot) \partial_{x_d} R(s, \cdot) ds. \end{aligned} \quad (2.5)$$

Now let us consider the following.

Lemma 2.1. *The integral equation (2.5) admits a unique solution $w \in L^2(0, T; H^2(\Omega))$ satisfying*

$$\|w(t, \cdot)\|_{H^1(\Omega)} \leq C \int_0^t (t-s)^{\alpha/2-1} \|f(s, \cdot)\|_{L^2(\omega)} ds, \quad t \in (0, T), \quad (2.6)$$

with $C > 0$ depending on R, \mathcal{A}, T . Moreover, we have $\partial_t^\alpha w \in L^2(Q)$.

Proof. We introduce the maps $\mathcal{K}_1, \mathcal{M} : L^2(0, T; H^1(\Omega)) \rightarrow L^2(0, T; H^1(\Omega))$, $\mathcal{K}_2 : L^2((0, T) \times \omega) \rightarrow L^2(0, T; H^1(\Omega))$ defined by

$$\mathcal{K}_1 v(t, \cdot) := \int_0^t S(t-s) B_1 v(s, \cdot) ds,$$

$$\mathcal{K}_2 f(t, \cdot) := \int_0^t S(t-s_1) B_2 \left(\int_0^{s_1} \tilde{S}(s_1-s_2) f(s_2, \cdot) R(s_2, \cdot) ds_2 \right) ds_1 + \int_0^t S(t-s) f(s, \cdot) \partial_{x_d} R(s, \cdot) ds$$

and $\mathcal{M}v = \mathcal{K}_1 v + \mathcal{K}_2 f$. Applying formula (1.148) of [29, Theorem 1.6] and the fact that both $D(A^{\frac{1}{2}})$ and $D(\tilde{A}^{\frac{1}{2}})$ embedded continuously into $H^1(\Omega)$, we find

$$\|\mathcal{K}_1 v(t, \cdot)\|_{H^1(\Omega)} \leq C \|\mathcal{K}_1 v(t, \cdot)\|_{D(A^{\frac{1}{2}})} \leq C \int_0^t \frac{(t-s)^{\frac{\alpha}{2}-1}}{\Gamma(\alpha/2)} \|v(s, \cdot)\|_{H^1(\Omega)} ds, \quad t \in (0, T) \quad (2.7)$$

$$\begin{aligned} \|\mathcal{K}_2 f(t, \cdot)\|_{H^1(\Omega)} &\leq C \left(\int_0^t (t-s)^{\frac{\alpha}{2}-1} \|f(s, \cdot)\|_{L^2(\omega)} ds + \int_0^t \int_0^{s_1} (t-s_1)^{\frac{\alpha}{2}-1} (s_1-s_2)^{\frac{\alpha}{2}-1} \|f(s_2, \cdot)\|_{L^2(\omega)} ds_2 ds_1 \right) \\ &\leq C \int_0^t (t-s)^{\frac{\alpha}{2}-1} \|f(s, \cdot)\|_{L^2(\omega)} ds, \quad t \in (0, T). \end{aligned} \quad (2.8)$$

Following the proof of [7, Proposition 1], for $n \in \mathbb{N}$, we find by iteration

$$\|\mathcal{K}_1^n v(t, \cdot)\|_{H^1(\Omega)} \leq C^n \int_0^t \frac{(t-s)^{\frac{n\alpha}{2}-1}}{\Gamma(n\alpha/2)} \|v(s, \cdot)\|_{H^1(\Omega)} ds, \quad t \in (0, T) \quad (2.9)$$

$$\|\mathcal{K}_1^{n-1} \mathcal{K}_2 f(t, \cdot)\|_{H^1(\Omega)} \leq C^n \int_0^t \frac{(t-s)^{\frac{n\alpha}{2}-1}}{\Gamma(n\alpha/2)} \|f(s, \cdot)\|_{L^2(\omega)} ds, \quad t \in (0, T). \quad (2.10)$$

It follows that for $n \in \mathbb{N}$ sufficiently large we have

$$\|\mathcal{M}^n v_1(t, \cdot) - \mathcal{M}^n v_2(t, \cdot)\|_{H^1(\Omega)} \leq C^n \int_0^t \frac{(t-s)^{\frac{n\alpha}{2}-1}}{\Gamma(n\alpha/2)} \|v_1(s, \cdot) - v_2(s, \cdot)\|_{H^1(\Omega)} ds, \quad t \in (0, T)$$

and an application of the Young inequality for convolution product implies that

$$\begin{aligned} \|\mathcal{M}^n v_1 - \mathcal{M}^n v_2\|_{L^2(0, T; H^1(\Omega))} &\leq C^n \left(\int_0^T \frac{t^{\frac{n\alpha}{2}-1}}{\Gamma(n\alpha/2)} dt \right) \|v_1 - v_2\|_{L^2(0, T; H^1(\Omega))} \\ &\leq \frac{C^n T^{\frac{n\alpha}{2}}}{\frac{n\alpha}{2} \Gamma(n\alpha/2)} \|v_1 - v_2\|_{L^2(0, T; H^1(\Omega))} \\ &\leq \frac{C^n T^{\frac{n\alpha}{2}+1}}{\Gamma(n\alpha/2+1)} \|v_1 - v_2\|_{L^2(0, T; H^1(\Omega))}. \end{aligned}$$

Then, using the fact that

$$\lim_{n \rightarrow +\infty} \frac{C^n T^{\frac{n\alpha}{2}+1}}{\Gamma(n\alpha/2+1)} = 0,$$

we deduce that there exists $n_0 \in \mathbb{N}$ such that \mathcal{M}^{n_0} is a contraction. Moreover, conditions (2.9)-(2.10) imply

$$\begin{aligned} \|\mathcal{M}^n v\|_{L^2(0, T; H^1(\Omega))} &\leq \frac{C^n T^{\frac{n\alpha}{2}+1}}{\Gamma(n\alpha/2+1)} \|v\|_{L^2(0, T; H^1(\Omega))} + \left(\sum_{k=1}^n \frac{C^k T^{\frac{k\alpha}{2}+1}}{\Gamma(k\alpha/2+1)} \right) \|f\|_{L^2((0, T) \times \omega)} \\ &\leq \frac{C^n T^{\frac{n\alpha}{2}+1}}{\Gamma(n\alpha/2+1)} \|v\|_{L^2(0, T; H^1(\Omega))} + \left(\sum_{k=1}^{\infty} \frac{C^k T^{\frac{k\alpha}{2}+1}}{\Gamma(k\alpha/2+1)} \right) \|f\|_{L^2((0, T) \times \omega)}. \end{aligned}$$

Therefore, by eventually increasing the size of n_0 , we deduce that \mathcal{M}^{n_0} admits a unique fixed point $w \in L^2(0, T; H^1(\Omega))$ which by uniqueness of this fixed point is also a fixed point of \mathcal{M} . Moreover, in view of (2.9)-(2.10), for $n_1 \in \mathbb{N}$ satisfying $n_1 \geq \frac{4}{\alpha}$ and for a.e $t \in (0, T)$, we have

$$\begin{aligned} \|w(t, \cdot)\|_{H^1(\Omega)} &= \|\mathcal{M}^{n_1} w(t, \cdot)\|_{H^1(\Omega)} \\ &\leq C \left(\int_0^t (t-s)^{\frac{n_1\alpha}{2}-1} \|w(s, \cdot)\|_{H^1(\Omega)} ds + \int_0^t (t-s)^{\frac{\alpha}{2}-1} \|f(s, \cdot)\|_{L^2(\omega)} ds \right) \\ &\leq C \left(T^{\frac{n_1\alpha}{2}-1} \int_0^t \|w(s, \cdot)\|_{H^1(\Omega)} ds + \int_0^t (t-s)^{\frac{\alpha}{2}-1} \|f(s, \cdot)\|_{L^2(\omega)} ds \right). \end{aligned}$$

Therefore, applying Gronwall inequality for function lying in $L^2(0, T)$ (see e.g [1, Lemma 6.3]), we find

$$\|w(t, \cdot)\|_{H^1(\Omega)} \leq C \left(\int_0^t (t-s)^{\frac{\alpha}{2}-1} \|f(s, \cdot)\|_{L^2(\omega)} ds \right) e^{Ct}, \quad t \in (0, T),$$

which clearly implies (2.6). Finally, using the fact that

$$B_1 w, s_1 \mapsto B_2 \left(\int_0^{s_1} \tilde{S}(s_1 - s_2) f(s_2, \cdot) R(s_2, \cdot) ds_2 \right), \partial_{x_d} F \in L^2(Q),$$

we deduce from Lemma 3.1 and assumption (H00), that \mathcal{M} takes values in $L^2(0, T; H^2(\Omega))$ and therefore $w \in L^2(0, T; H^2(\Omega))$. In the same way, we prove that $\partial_t^\alpha w \in L^2(Q)$. \square

According to Lemma 2.1, we have $u \in L^2(0, T; H^1(\Omega))$, $\mathcal{A}u$, $\partial_t^\alpha u \in L^2(Q)$ and $w = \partial_{x_d} u \in L^2(0, T; H^2(\Omega))$, $\partial_t^\alpha \partial_{x_d} u$, $\mathcal{A} \partial_{x_d} u \in L^2(Q)$. Combining this with (2.2), we deduce that

$$x_d \mapsto u(\cdot, \cdot, x_d) \in H^3((-\ell, \ell); L^2((0, T) \times \omega)) \cap H^1(-\ell, \ell; L^2(0, T; H^2(\omega))),$$

$$x_d \mapsto \partial_t^\alpha u(\cdot, \cdot, x_d) \in H^1((-\ell, \ell); L^2((0, T) \times \omega)).$$

Then, (1.2) implies

$$\begin{aligned} a_{dd} \partial_{x_d} w(t, x', \ell) &= a_{dd} \partial_{x_d}^2 u(t, x', \ell) = [\partial_t^\alpha u + (\mathcal{A} + a_{dd} \partial_{x_d}^2) u](t, x', \ell) - R(t, x', \ell) f(t, x') \\ &= R(t, x', \ell) [h(t, x') - f(t, x')], \quad (t, x') \in (0, T) \times \omega, \end{aligned} \quad (2.11)$$

with h given by (1.15). Fixing $r \in (3/4, 1)$ and $\tau_1 : L^2(0, T; H^{2r}(\Omega)) \ni y \longrightarrow \frac{a_{dd} \partial_{x_d} y(t, x', \ell)}{R(t, x', \ell)} \in L^2((0, T) \times \omega)$, we obtain

$$f(t, \cdot) = h(t, \cdot) - \tau_1 w(t, \cdot), \quad t \in (0, T).$$

Moreover, applying (2.2), (2.5) and using the fact that $u, w \in L^2(0, T; H^1(\Omega))$, we get

$$(\partial_{x_d} \mathcal{A}) \left(\int_0^{s_1} \tilde{S}(s_1 - s_2) f(s_2, \cdot) R(s_2, \cdot) ds_2 \right) ds_1 = (\partial_{x_d} \mathcal{A}) u \in L^2(Q), \quad (2.12)$$

which combined with (H00) implies

$$\begin{aligned} f(t, \cdot) &= h(t, \cdot) - \tau_1 \left(\int_0^t S(t - s_1) (\partial_{x_d} \mathcal{A}) \left(\int_0^{s_1} \tilde{S}(s_1 - s_2) f(s_2, \cdot) R(s_2, \cdot) ds_2 \right) ds_1 \right) \\ &\quad - \tau_1 \left(\int_0^t S(t - s) f(s, \cdot) \partial_{x_d} R(s, \cdot) ds \right), \quad t \in (0, T). \end{aligned} \quad (2.13)$$

In view of (H00) by interpolation $D(A^r)$ embedded continuously into $H^{2r}(\Omega)$. Moreover, in view of formula (1.148) of [29, Theorem 1.6], we have

$$\begin{aligned} &\left\| S(t - s_1) (\partial_{x_d} \mathcal{A}) \left(\int_0^{s_1} \tilde{S}(s_1 - s_2) f(s_2, \cdot) R(s_2, \cdot) ds_2 \right) \right\|_{H^{2r}(\Omega)} \\ &\leq C \left\| S(t - s_1) (\partial_{x_d} \mathcal{A}) \left(\int_0^{s_1} \tilde{S}(s_1 - s_2) f(s_2, \cdot) R(s_2, \cdot) ds_2 \right) \right\|_{D(A^r)} \\ &\leq C (t - s_1)^{\alpha(1-r)-1} \left\| (\partial_{x_d} \mathcal{A}) \left(\int_0^{s_1} \tilde{S}(s_1 - s_2) f(s_2, \cdot) R(s_2, \cdot) ds_2 \right) \right\|_{L^2(\Omega)}, \quad s_1 \in (0, t). \end{aligned}$$

On the other hand, applying Young inequality for convolution product, we obtain

$$\begin{aligned} &\int_0^T \int_0^t \left\| S(t - s_1) (\partial_{x_d} \mathcal{A}) \left(\int_0^{s_1} \tilde{S}(s_1 - s_2) f(s_2, \cdot) R(s_2, \cdot) ds_2 \right) \right\|_{H^{2r}(\Omega)} ds_1 dt \\ &\leq C \int_0^T \int_0^t (t - s_1)^{\alpha(1-r)-1} \left\| (\partial_{x_d} \mathcal{A}) \left(\int_0^t \tilde{S}(s_1 - s_2) f(s_2, \cdot) R(s_2, \cdot) ds_2 \right) \right\|_{L^2(\Omega)} ds_1 dt \\ &\leq CT^{1-r} \left\| (\partial_{x_d} \mathcal{A}) \left(\int_0^t \tilde{S}(s_1 - s_2) f(s_2, \cdot) R(s_2, \cdot) ds_2 \right) \right\|_{L^1(0, T; L^2(\Omega))} \\ &\leq C \left\| (\partial_{x_d} \mathcal{A}) \left(\int_0^t \tilde{S}(s_1 - s_2) f(s_2, \cdot) R(s_2, \cdot) ds_2 \right) \right\|_{L^2(Q)} < \infty. \end{aligned}$$

Thus, by Fubini theorem for a.e. $t \in (0, T)$, we have

$$s_1 \mapsto S(t - s_1)(\partial_{x_d}\mathcal{A}) \left(\int_0^{s_1} \tilde{S}(s_1 - s_2)f(s_2, \cdot)R(s_2, \cdot)ds_2 \right) \in L^1(0, t; H^{2r}(\Omega)).$$

In the same way, for a.e. $t \in (0, T)$, we get

$$s \mapsto S(t - s)f(s, \cdot)\partial_{x_d}R(s, \cdot) \in L^1(0, t; H^{2r}(\Omega)).$$

Therefore, from (2.13), for a.e. $t \in (0, T)$, we find

$$\begin{aligned} f(t, \cdot) = & h(t, \cdot) - \int_0^t \tau_1 S(t - s_1)(\partial_{x_d}\mathcal{A}) \left(\int_0^{s_1} \tilde{S}(s_1 - s_2)f(s_2, \cdot)R(s_2, \cdot)ds_2 \right) ds_1 \\ & - \int_0^t \tau_1 S(t - s)f(s, \cdot)\partial_{x_d}R(s, \cdot)ds. \end{aligned} \quad (2.14)$$

This proves (1.16), let us prove that this problem is well-posed. We fix the maps $\mathcal{G}, \mathcal{H} : L^2((0, T) \times \omega) \rightarrow L^2((0, T) \times \omega)$, with

$$\begin{aligned} \mathcal{H}g(t, \cdot) := & - \int_0^t \tau_1 S(t - s_1)(\partial_{x_d}\mathcal{A}) \left(\int_0^{s_1} \tilde{S}(s_1 - s_2)g(s_2, \cdot)R(s_2, \cdot)ds_2 \right) ds_1 \\ & - \int_0^t \tau_1 S(t - s)g(s, \cdot)\partial_{x_d}R(s, \cdot)ds, \quad g \in L^2((0, T) \times \omega) \end{aligned}$$

and

$$\mathcal{G}g(t, \cdot) := h(t, \cdot) + \mathcal{H}g(t, \cdot), \quad g \in L^2((0, T) \times \omega).$$

Note that

$$\begin{aligned} \|\mathcal{H}g(t, \cdot)\|_{L^2(\omega)} & \leq \int_0^t \left\| \tau_1 S(t - s_1)(\partial_{x_d}\mathcal{A}) \left(\int_0^{s_1} \tilde{S}(s_1 - s_2)g(s_2, \cdot)R(s_2, \cdot)ds_2 \right) \right\|_{L^2(\omega)} ds_1 \\ & \quad + \int_0^t \|\tau_1 S(t - s)g(s, \cdot)\partial_{x_d}R(s, \cdot)\|_{L^2(\omega)} ds \\ & \leq C \int_0^t \left\| S(t - s_1)(\partial_{x_d}\mathcal{A}) \left(\int_0^{s_1} \tilde{S}(s_1 - s_2)g(s_2, \cdot)R(s_2, \cdot)ds_2 \right) \right\|_{H^{2r}(\Omega)} ds_1 \\ & \quad + C \int_0^t \|S(t - s)g(s, \cdot)\partial_{x_d}R(s, \cdot)\|_{H^{2r}(\Omega)} ds \\ & \leq C(t^{\alpha(1-r)-1}\mathbf{1}_{(0, +\infty)}) * \left\| (\partial_{x_d}\mathcal{A}) \left(\mathbf{1}_{(0, T)}(t) \int_0^t \tilde{S}(t - s)g(s, \cdot)R(s, \cdot)ds \right) \right\|_{L^2(\Omega)} \\ & \quad + C \|\partial_{x_d}R\|_{L^\infty(Q)} (t^{\alpha(1-r)-1}\mathbf{1}_{(0, +\infty)}) * (\|g(\cdot, \cdot)\|_{L^2(\omega)} \mathbf{1}_{(0, T)}), \quad t \in (0, T). \end{aligned} \quad (2.15)$$

On the other hand, in view of (2.2), fixing $w_1 \in L^2(0, T; H^1(\Omega))$ the solution of (2.5) with $f = g$ we obtain

$$(\partial_{x_d}\mathcal{A}) \left(\int_0^t \tilde{S}(t - s)g(s, \cdot)R(s, \cdot)ds \right) = B_1 w_1(t, \cdot) + B_2 \int_0^t S(t - s)gR(s, \cdot)ds,$$

where B_1 and B_2 are defined in formula (2.3)-(2.4). Thus, applying Lemma 2.1 and formula (1.148) of [29, Theorem 1.6], we obtain

$$\left\| (\partial_{x_d}\mathcal{A}) \left(\int_0^t \tilde{S}(t - s)g(s, \cdot)R(s, \cdot)ds \right) \right\|_{L^2(\Omega)} \leq C \int_0^t (t - s)^{\frac{\alpha}{2}-1} \|g(s, \cdot)\|_{L^2(\omega)} ds, \quad t \in (0, T).$$

Combining this with (2.15), we get

$$\|\mathcal{H}g(t, \cdot)\|_{L^2(\omega)} \leq C \int_0^t \frac{(t-s)^{\alpha(1-r)-1}}{\Gamma(\alpha(1-r))} \|g(s, \cdot)\|_{L^2(\omega)} ds, \quad t \in (0, T).$$

By iteration, for all $n \in \mathbb{N}$, we deduce that

$$\|\mathcal{H}^n g(t, \cdot)\|_{L^2(\omega)} \leq C^n \int_0^t \frac{(t-s)^{n\alpha(1-r)-1}}{\Gamma(n\alpha(1-r))} \|g(s, \cdot)\|_{L^2(\omega)} ds, \quad t \in (0, T)$$

and in a similar way to Lemma 2.1, we deduce that there exists $n_2 \in \mathbb{N}$ such that \mathcal{G}^{n_2} is a contraction and \mathcal{G} admits a unique fixed point $f \in L^2((0, T) \times \omega)$ satisfying

$$\|f(t, \cdot)\|_{L^2(\omega)} = \|\mathcal{G}f(t, \cdot)\|_{L^2(\omega)} \leq \|h(t, \cdot)\|_{L^2(\omega)} + C \int_0^t (t-s)^{\alpha(1-r)-1} \|f(s, \cdot)\|_{L^2(\omega)} ds, \quad t \in (0, T).$$

Therefore, in view of [7, Lemma 3] (see also [33, Theorem A]), we have

$$\|f(t, \cdot)\|_{L^2(\omega)} \leq C \left(\|h(t, \cdot)\|_{L^2(\omega)} + \int_0^t (t-s)^{\alpha(1-r)-1} \|h(s, \cdot)\|_{L^2(\omega)} ds \right), \quad t \in (0, T),$$

and an application of the Young inequality yields

$$\|f\|_{L^2((0, T) \times \omega)} \leq C \|h\|_{L^2((0, T) \times \omega)}.$$

This proves the well-posedness of (1.14) and the reconstruction of f from the data h . Now let us consider the proof of the last part of Theorem 1.2. For this purpose, we fix $(h_1, f) \in L^2((0, T) \times \omega) \times L^2((0, T) \times \omega)$ satisfying (1.16) with $h = h_1$ and we consider $u \in L^2(0, T; H^1(\Omega))$ solving (1.2). Following the above argumentation, we can define $h \in L^2((0, T) \times \omega)$ given by (1.15) and f solves (1.16). This implies that

$$h_1(t, \cdot) = f - \int_0^t K(t, s) f(s, \cdot) ds = h(t, \cdot), \quad t \in (0, T).$$

Therefore we have $h = h_1$ and the proof of Theorem 1.2 for (1.2)-(1.4), with $m = 1$ and $n = 0$, is completed. Using similar arguments, one can check that this result is still true for the problem (1.2)-(1.4), with $m = 1$ and $n = 1$.

3. PROOF OF THEOREM 1.4

This section is devoted to the proof of Theorem 1.4. In contrast to the preceding section, for u solving (1.11), $\partial_{x_d} u$ will not be a solution of an initial boundary value problem with homogeneous boundary condition. However, with suitable regularity conditions on F we can consider the trace of u at $\{x_d = \ell\}$. We will start by proving Proposition 1.3

Proof of Proposition 1.3. We consider first the case $q = 0$. Let A be the operator \mathcal{A} acting on $L^2(\tilde{\Omega})$ with domain

$$D(A) := \{g \in L^2(\tilde{\Omega}) : \mathcal{A}g \in L^2(\tilde{\Omega}), u|_{\partial\omega \times (0, \ell)} = 0, u|_{x_d = \ell} = 0, \partial_{x_d} u|_{x_d = 0} = 0\}.$$

The spectrum of A consists of a non-decreasing sequence of strictly positive eigenvalues $(\lambda_n)_{n \geq 1}$. Let us also introduce an orthonormal basis in the Hilbert space $L^2(\tilde{\Omega})$ of eigenfunctions $(\varphi_n)_{n \geq 1}$ of A

associated with the non-decreasing sequence of eigenvalues $(\lambda_n)_{n \geq 1}$. Then, for $n \in \mathbb{N}$, the solution of (1.11) is given by

$$u_n(t) := \langle u(t, \cdot), \varphi_n \rangle_{L^2(\tilde{\Omega})} = \int_0^t (t-s)^{\alpha-1} E_{\alpha, \alpha}(-\lambda_n(t-s)^\alpha) F_n(s) ds,$$

where $F_n(t) := \langle F(t, \cdot), \varphi_n \rangle_{L^2(\tilde{\Omega})}$. Since $F_n \in W^{1,p}(0, T)$, $F_n(0) = 0$, applying [30, Lemma 3.2] and integrating by parts we find

$$u_n(t) = -\lambda_n^{-1} F_n(t) + \lambda_n^{-1} \int_0^t E_{\alpha, 1}(-\lambda_n(t-s)^\alpha) F_n'(s) ds.$$

Thus, we have $u = v + w$ where

$$v(t, \cdot) = A^{-1} F(t, \cdot), \quad w_n(t) := \langle w(t, \cdot), \varphi_n \rangle_{L^2(\tilde{\Omega})} = \lambda_n^{-1} \int_0^t E_{\alpha, 1}(-\lambda_n(t-s)^\alpha) F_n'(s) ds.$$

Note that, for all $t \in [0, T]$, $v(t, \cdot)$ solves the boundary value problem

$$\begin{cases} \mathcal{A}v(t, \cdot) = F(t, \cdot), & \text{in } \tilde{\Omega}, \\ v(t, x) = 0, & x \in \partial\omega \times (0, \ell), \\ v(t, x', \ell) = 0, & x' \in \omega, \\ \partial_{x_d} v(t, x', 0) = 0, & x' \in \omega. \end{cases}$$

Therefore, applying assumption (\tilde{H}) and the fact that $F \in \mathcal{C}([0, T]; H^{2\gamma}(\tilde{\Omega}))$, we deduce that $v \in \mathcal{C}([0, T]; H^{2(1+\gamma)}(\tilde{\Omega}))$. Thus, in order to complete the proof, we only need to check that $w \in \mathcal{C}([0, T]; H^{2(1+\gamma)}(\tilde{\Omega}))$. For this purpose, using the fact that $F_n' \in L^p(0, T)$ and $t \mapsto t^{\alpha-1} \in L^{p'}(0, T)$, with $1/p' = 1 - 1/p$, one can check that $w_n \in \mathcal{C}([0, T])$, $n \in \mathbb{N}$. Moreover, fixing $m, n \in \mathbb{N}$, with $m < n$, and applying formula (1.148) of [29, Theorem 1.6], we find

$$\begin{aligned} & \left\| \sum_{k=m}^n \lambda_k^{1+\gamma} w_k(t) \varphi_k \right\|_{L^2(\tilde{\Omega})} \\ & \leq \int_0^t \left\| \sum_{k=m}^n \lambda_k^\gamma E_{\alpha, 1}(-\lambda_k(t-s)^\alpha) F_k'(s) \varphi_k \right\|_{L^2(\tilde{\Omega})} ds \\ & \leq C \int_0^t (t-s)^{-\alpha\gamma} \left\| \sum_{k=m}^n F_k'(s) \varphi_k \right\|_{L^2(\tilde{\Omega})} ds. \end{aligned}$$

On the other hand, since $\frac{1}{p} < 1 - \alpha\gamma$, we have $\frac{1}{p'} = 1 - \frac{1}{p} > \alpha\gamma$ and we deduce

$$\int_0^T s^{-p'\alpha\gamma} ds < \infty.$$

Therefore, applying the Hölder inequality, we obtain

$$\left\| \sum_{k=m}^n \lambda_k^{1+\gamma} w_k(t) \varphi_k \right\|_{L^2(\tilde{\Omega})} \leq C \left\| \sum_{k=m}^n F_k'(s) \varphi_k \right\|_{L^p(0, T; L^2(\tilde{\Omega}))}.$$

Combining this with the fact that $F \in L^p(0, T; L^2(\tilde{\Omega}))$, we deduce that

$$\lim_{m, n \rightarrow \infty} \left\| \sum_{k=m}^n w_k \varphi_k \right\|_{\mathcal{C}([0, T]; D(A^{1+\gamma}))} = 0.$$

Thus, the sequence

$$\sum_{n \in \mathbb{N}} w_n \varphi_n$$

is a Cauchy sequence and therefore a convergent sequence of $\mathcal{C}([0, T]; D(A^{1+\gamma}))$. Since this sequence converges to w in the sense of $\mathcal{C}([0, T]; L^2(\tilde{\Omega}))$, we deduce that $w \in \mathcal{C}([0, T]; D(A^{1+\gamma}))$ and, in view of (\tilde{H}) , we deduce that $w \in \mathcal{C}([0, T]; H^{2(1+\gamma)}(\tilde{\Omega}))$. Now let us prove that $u \in \mathcal{C}^1([0, T]; H^{2\gamma}(\tilde{\Omega}))$. Note first that $u_n \in \mathcal{C}^1([0, T])$ with

$$u'_n(t) = F_n(0)t^{\alpha-1}E_{\alpha, \alpha}(-\lambda_n t^\alpha) + \int_0^t s^{\alpha-1}E_{\alpha, \alpha}(-\lambda_n s^\alpha)F'_n(t-s)ds = \int_0^t s^{\alpha-1}E_{\alpha, \alpha}(-\lambda_n s^\alpha)F'_n(t-s)ds.$$

Here we have used the fact that $F|_{t=0} = 0$. Repeating the above arguments and using the fact that $F \in W^{1,p}(0, T; L^2(\tilde{\Omega}))$ with $\frac{1}{p} < \alpha(1-\gamma)$, we deduce that $\partial_t u \in \mathcal{C}([0, T]; D(A^\gamma)) \subset \mathcal{C}([0, T]; H^{2\gamma}(\tilde{\Omega}))$. Therefore, we have $u \in \mathcal{C}^1([0, T]; H^{2\gamma}(\tilde{\Omega}))$.

Now let us consider the case $q \neq 0$. We introduce the map

$$\mathcal{G}(v) := \int_0^t J(t-s)F(s)ds - \int_0^t J(t-s)q(s)v(s)ds$$

defined on $\mathcal{C}([0, T]; H^{2(1+\gamma)}(\tilde{\Omega})) \cap \mathcal{C}^1([0, T]; H^{2\gamma}(\tilde{\Omega}))$ with $J(t)$ given by

$$J(t)h := \sum_{n=1}^{\infty} t^{\alpha-1}E_{\alpha, \alpha}(-\lambda_n t^\alpha) \langle h, \varphi_n \rangle \varphi_n, \quad h \in L^2(\tilde{\Omega}), \quad t > 0.$$

Then, using a classical fixed point argument combined with the preceding analysis, we deduce that \mathcal{G} admits a unique fixed point lying in $\mathcal{C}([0, T]; H^{2(1+\gamma)}(\tilde{\Omega})) \cap \mathcal{C}^1([0, T]; H^{2\gamma}(\tilde{\Omega}))$ which will be the solution of (1.11). This completes the proof of the lemma. \square

From now on and in all the remaining part of this section, we assume that $\alpha \in (0, 1]$ and that the conditions of Proposition 1.3 are fulfilled. We consider \tilde{u} defined on $(0, T) \times \omega \times (-\ell, \ell)$ by

$$\tilde{u}(t, x', -x_d) := \tilde{u}(t, x', x_d) = u(t, x', x_d), \quad (t, x', x_d) \in (0, T) \times \omega \times (0, \ell).$$

Then, it is clear that $\tilde{u}|_{(0, T) \times \tilde{\Omega}} \in \mathcal{C}([0, T]; H^{2(1+\gamma)}(\tilde{\Omega})) \cap \mathcal{C}^1([0, T]; H^\gamma(\tilde{\Omega}))$ and $\tilde{u}|_{(0, T) \times \omega \times (-\ell, 0)} \in \mathcal{C}([0, T]; H^{2(1+\gamma)}(\omega \times (-\ell, 0))) \cap \mathcal{C}^1([0, T]; H^\gamma(\omega \times (-\ell, 0)))$. Moreover, we have

$$\partial_{x_d}^{2k} \tilde{u}(t, x', 0_+) = \partial_{x_d}^{2k} \tilde{u}(t, x', 0_-), \quad \partial_{x_d} \tilde{u}(t, x', 0_+) = 0 = \partial_{x_d} \tilde{u}(t, x', 0_-), \quad (t, x') \in (0, T) \times \omega, \quad k = 0, 1.$$

Therefore, we deduce that $\tilde{u} \in \mathcal{C}([0, T]; H^3(\Omega)) \cap \mathcal{C}^1([0, T]; H^{2\gamma}(\Omega))$ and, thanks to (1.9), \tilde{u} solves

$$\left\{ \begin{array}{l} \partial_t^\alpha \tilde{u} + \mathcal{A}\tilde{u} + q(t, x')\tilde{u} = f(t, x')\tilde{R}(t, x), \quad (t, x', x_n) \in (0, T) \times \omega \times (-\ell, \ell), \\ \tilde{u}(t, x) = 0, \quad (t, x) \in (0, T) \times \partial\omega \times (-\ell, \ell), \\ \tilde{u}(t, x', \pm\ell) = 0 \quad (t, x') \in (0, T) \times \omega, \\ \partial_t^k \tilde{u}(0, x) = 0, \quad x \in \omega \times (-\ell, \ell), \quad k = 0, \dots, m_\alpha, \end{array} \right.$$

where \tilde{R} is the extension of R to $(0, T) \times \omega \times (-\ell, \ell)$ given by

$$\tilde{R}(t, x', x_d) := R(t, x', -x_d), \quad (t, x', x_d) \in (0, T) \times \omega \times (-\ell, 0).$$

Fixing $v = \partial_{x_d} \tilde{u}$ and using the fact that $\tilde{u} \in \mathcal{C}([0, T]; H^3(\Omega)) \cap \mathcal{C}^1([0, T]; H^{2\gamma}(\Omega))$, we deduce that v solves the problem

$$\left\{ \begin{array}{l} \partial_t^\alpha v + \mathcal{A}v + q(t, x')v = H(t, x', x_d) + f(t, x')\partial_{x_d} \tilde{R}(t, x), \quad (t, x', x_n) \in (0, T) \times \omega \times (-\ell, \ell), \\ v(t, x) = 0, \quad (t, x) \in (0, T) \times \partial\omega \times (-\ell, \ell), \\ v(t, x', \pm\ell) = \partial_{x_d} \tilde{u}(t, x', \pm\ell) = \pm \partial_{x_d} u(t, x', \ell), \quad (t, x') \in (0, T) \times \omega, \\ \partial_t^k v(0, x) = 0, \quad x \in \omega \times (-\ell, \ell), \quad k = 0, \dots, m_\alpha, \end{array} \right. \quad (3.1)$$

with

$$H(t, x', x_d) := \sum_{i,j=1}^d \partial_{x_i} (\partial_{x_d} a_{ij}(x) \partial_{x_j} \tilde{u}).$$

Note that here since $R, \partial_{x_d} R \in L^\infty((0, T) \times \tilde{\Omega})$, we have $\tilde{R}, \partial_{x_d} \tilde{R} \in L^\infty((0, T) \times \Omega)$. We are now in position to prove Theorem 1.4.

Proof of Theorem 1.4. In all this proof C denotes a generic constant depending on $\alpha, c, T, \Omega, \mathcal{A}, q, d$. According to Lemma 1.3 the solution u of (1.11) is lying in $\mathcal{C}([0, T]; H^{2(1+\gamma)}(\tilde{\Omega})) \cap \mathcal{C}^1([0, T]; H^\gamma(\tilde{\Omega}))$ and $v = \partial_{x_d} \tilde{u}$, with \tilde{u} the even extension of u to $(0, T) \times \omega \times (-\ell, \ell)$, which solves (3.1). Note first that projecting the equation (1.11) in $(0, T) \times \omega \times \{\ell\}$ and using the fact that, for all $(t, x') \in (0, T) \times \omega$, we have $u(t, x', \ell) = 0$, we deduce that

$$f(t, x')R(t, x', \ell) = [a_{d,d}\partial_{x_d}^2 u + 2 \sum_{j=1}^{d-1} a_{jd}\partial_{x_j}\partial_{x_d} u + \sum_{j=1}^d \partial_{x_j} a_{jd}\partial_{x_d} u](t, x', \ell), \quad (t, x') \in (0, T) \times \omega.$$

Combining this with (1.13) and the fact that $v = \partial_{x_d} u$, for all $t \in (0, T)$, we deduce that

$$\begin{aligned} \|f(t, \cdot)\|_{L^2(\omega)} &\leq c^{-1}C(\|\partial_{x_d} v(t, \cdot, \ell)\|_{L^2(\omega)} + \|\partial_{x_d} u(t, \cdot, \ell)\|_{H^1(\omega)}) \\ &\leq C(\|\partial_{x_d} v(t, \cdot, \ell)\|_{L^2(\omega)} + \|\partial_{x_d} u(t, \cdot, \ell)\|_{H^{\frac{3}{2}}(\omega)}). \end{aligned} \quad (3.2)$$

Thus, the proof of (1.17) will be completed if we can derive a suitable estimate of $\|\partial_{x_d} v(t, \cdot, \ell)\|_{L^2(\omega)}$

For this purpose, we decompose v into $v = v_1 + v_2$, where v_1 solves

$$\left\{ \begin{array}{l} \partial_t^\alpha v_1 + \mathcal{A}v_1 = 0, \quad \text{in } (0, T) \times \omega \times (-\ell, \ell), \\ v_1(t, x) = 0, \quad (t, x) \in (0, T) \times \partial\omega \times (-\ell, \ell), \\ v_1(t, x', \pm\ell) = \partial_{x_d} \tilde{u}(t, x', \pm\ell) = \pm \partial_{x_d} u(t, x', \ell), \quad (t, x') \in (0, T) \times \omega, \\ v_1(0, x) = 0, \quad x \in \omega \times (-\ell, \ell), \end{array} \right. \quad (3.3)$$

and v_2 solves

$$\left\{ \begin{array}{l} \partial_t^\alpha v_2 + \mathcal{A}v_2 = -qv(t, x) + H(t, x', x_d) + f(t, x') \partial_{x_d} \tilde{R}(t, x), \quad (t, x', x_n) \in (0, T) \times \omega \times (-\ell, \ell), \\ v_2(t, x) = 0, \quad (t, x) \in (0, T) \times \partial\omega \times (-\ell, \ell), \\ v_2(t, x', \pm\ell) = \partial_{x_d} \tilde{u}(t, x', \pm\ell) = 0, \quad (t, x') \in (0, T) \times \omega, \\ v_2(0, x) = 0, \quad x \in \omega \times (-\ell, \ell). \end{array} \right. \quad (3.4)$$

Since $u \in \mathcal{C}^1([0, T]; H^{2\gamma}(\tilde{\Omega})) \cap \mathcal{C}([0, T]; H^{2(1+\gamma)}(\tilde{\Omega}))$, we have

$$(t, x') \mapsto \partial_{x_d} u(t, x', \ell) \in \mathcal{C}^1([0, T]; H^{2\gamma-\frac{3}{2}}(\omega)) \cap \mathcal{C}([0, T]; H^{\frac{1}{2}+2\gamma}(\omega))$$

and $\partial_{x_d} u(0, \cdot, \ell) = 0$. therefore, using a classical lifting argument (e.g. [24, Theorem 8.3, Chapter 1]),

we can find $G \in \mathcal{C}^1([0, T]; H^{2\gamma-1}(\Omega)) \cap \mathcal{C}([0, T]; H^{2\gamma+1}(\Omega))$ such that

$$\begin{aligned} G(t, x', \ell) &= \partial_{x_d} u(t, x', \ell), \quad (t, x') \in (0, T) \times \omega, \\ G(0, x', x_d) &= 0, \quad (x', x_d) \in \omega \times (-\ell, \ell), \\ \|G\|_{W^{1,\infty}(0,T;L^2(\omega \times (-\ell, \ell)))} &+ \|G\|_{L^\infty(0,T;H^2(\omega \times (-\ell, \ell)))} \\ &\leq C(\|\partial_{x_d} u(\cdot, \cdot, \ell)\|_{L^\infty(0,T;H^{\frac{3}{2}}(\omega))} + \|\partial_{x_d} u(\cdot, \cdot, \ell)\|_{W^{1,\infty}(0,T;H^\delta(\omega))}). \end{aligned} \quad (3.5)$$

Therefore, we can decompose v_1 into $v_1 = G + w_1$ with w_1 solving

$$\left\{ \begin{array}{l} \partial_t^\alpha w_1 + \mathcal{A}w_1 = G_1, \quad \text{in } (0, T) \times \omega \times (-\ell, \ell), \\ w_1(t, x) = 0, \quad (t, x) \in (0, T) \times \partial\omega \times (-\ell, \ell), \\ w_1(t, x', \pm\ell) = 0, \quad (t, x') \in (0, T) \times \omega, \\ w_1(0, x) = 0, \quad x \in \omega \times (-\ell, \ell), \end{array} \right.$$

where $G_1 = -\partial_t^\alpha G - \mathcal{A}G \in \mathcal{C}([0, T]; L^2(\Omega))$. Thus, we have

$$w_1(t, \cdot) = \int_0^t S(s) G_1(t-s) ds,$$

where $S(t)$ corresponds to the operator valued function defined in Theorem 1.2. Therefore, applying

(H00) and formula (1.148) of [29, Theorem 1.6], we deduce that $w_1 \in \mathcal{C}([0, T]; H^{2\gamma}(\Omega))$ with

$$\|w_1\|_{L^\infty(0,T;H^{2\gamma}(\Omega))} \leq C \left(\int_0^T s^{\alpha(1-\gamma)-1} ds \right) \|G_1\|_{L^\infty(0,T;L^2(\Omega))}.$$

It follows that

$$\begin{aligned} \|v_1\|_{L^\infty(0,T;H^{2\gamma}(\Omega))} &\leq C(\|G_1\|_{L^\infty(0,T;L^2(\Omega))} + \|G\|_{L^\infty(0,T;H^2(\Omega))}) \\ &\leq C(\|G\|_{W^{1,\infty}(0,T;L^2(\omega \times (-\ell,\ell)))} + \|G\|_{L^\infty(0,T;H^2(\omega \times (-\ell,\ell)))}) \end{aligned}$$

and combining this with (3.5), we get

$$\|v_1\|_{L^\infty(0,T;H^{2\gamma}(\omega \times (-\ell,\ell))} \leq C(\|\partial_{x_d} u(\cdot, \cdot, \ell)\|_{L^\infty(0,T;H^{\frac{3}{2}}(\omega))} + \|\partial_{x_d} u(\cdot, \cdot, \ell)\|_{W^{1,\infty}(0,T;H^\delta(\omega))}). \quad (3.6)$$

Moreover, the estimate

$$\|\partial_{x_d} v_1(\cdot, \cdot, \ell)\|_{L^\infty(0,T;L^2(\omega))} \leq C \|v_1\|_{L^\infty(0,T;H^{2\gamma}(\omega \times (-\ell,\ell))},$$

implies

$$\|\partial_{x_d} v_1(\cdot, \cdot, \ell)\|_{L^\infty(0,T;L^2(\omega))} \leq C(\|\partial_{x_d} u(\cdot, \cdot, \ell)\|_{L^\infty(0,T;H^{\frac{3}{2}}(\omega))} + \|\partial_{x_d} u(\cdot, \cdot, \ell)\|_{W^{1,\infty}(0,T;H^\delta(\omega))}). \quad (3.7)$$

On the other hand, we have

$$v_2(t, \cdot) = \int_0^t S(t-s)[-qv(s, \cdot) + H(s, \cdot) + \partial_{x_d} \tilde{F}(s, \cdot)] ds.$$

Thus, applying (H00) and repeating the arguments of Lemma 1.3, we get

$$\|v_2(t, \cdot)\|_{H^{2\gamma}(\omega \times (-\ell,\ell))} \leq C \int_0^t (t-s)^{\alpha(1-\gamma)-1} [\|f(s, \cdot)\|_{L^2(\omega)} + \|v(s, \cdot)\|_{L^2(\omega \times (-\ell,\ell))} + \|H(s, \cdot)\|_{L^2(\omega \times (-\ell,\ell))}] ds$$

which, for all $t \in (0, T]$, implies that

$$\|\partial_{x_d} v_2(t, \cdot, \ell)\|_{L^2(\omega)} \leq C \int_0^t (t-s)^{\alpha(1-\gamma)-1} [\|f(s, \cdot)\|_{L^2(\omega)} + \|v(s, \cdot)\|_{L^2(\omega \times (-\ell,\ell))} + \|H(s, \cdot)\|_{L^2(\omega \times (-\ell,\ell))}] ds.$$

In light of (1.14), we get

$$\begin{aligned} H &= \partial_{x_d} a_{dd} \partial_{x_d}^2 \tilde{u} + 2 \sum_{j=1}^{d-1} \partial_{x_d} a_{jd} \partial_{x_j} \partial_{x_d} \tilde{u} + \sum_{j=1}^d \partial_{x_j} \partial_{x_d} a_{jd} \partial_{x_d} \tilde{u} + \sum_{j=1}^{d-1} \partial_{x_d}^2 a_{jd} \partial_{x_j} \tilde{u} \\ &= \partial_{x_d} a_{dd} \partial_{x_d} v + 2 \sum_{j=1}^{d-1} \partial_{x_d} a_{jd} \partial_{x_j} v + \left(\sum_{j=1}^d \partial_{x_j} \partial_{x_d} a_{jd} \right) v + \sum_{j=1}^{d-1} \partial_{x_d}^2 a_{jd} \partial_{x_j} \tilde{u} \end{aligned}$$

and it follows that

$$\|H(t, \cdot)\|_{L^2(\Omega)} \leq C(\|v(t, \cdot)\|_{H^1(\Omega)} + \|u(t, \cdot)\|_{H^1(\Omega)}), \quad t \in (0, T].$$

In view of the equation satisfied by v and according to the above arguments as well as the arguments used in Proposition 1.3, for all $t \in (0, T]$, we get

$$\begin{aligned} &\|v(t, \cdot)\|_{H^1(\Omega)} + \|u(t, \cdot)\|_{H^1(\Omega)} \\ &\leq C(\|f\|_{L^\infty(0,t;L^2(\omega))} + \|\partial_{x_d} u(\cdot, \cdot, \ell)\|_{L^\infty(0,T;H^{\frac{3}{2}}(\omega))} + \|\partial_{x_d} u(\cdot, \cdot, \ell)\|_{W^{1,\infty}(0,T;H^\delta(\omega))}). \end{aligned}$$

Thus, we find

$$\|H(t, \cdot)\|_{L^2(\Omega)} \leq C(\|f\|_{L^\infty(0,t;L^2(\omega))} + \|\partial_{x_d} u(\cdot, \cdot, \ell)\|_{L^\infty(0,T;H^{\frac{3}{2}}(\omega))} + \|\partial_{x_d} u(\cdot, \cdot, \ell)\|_{W^{1,\infty}(0,T;H^\delta(\omega))})$$

and it follows that

$$\begin{aligned} & \|\partial_{x_d} v_2(t, \cdot, \ell)\|_{L^2(\omega)} \\ & \leq C \left(\int_0^t (t-s)^{\alpha(1-\gamma)-1} \|f\|_{L^\infty(0,s;L^2(\omega))} ds + \|\partial_{x_d} u(\cdot, \cdot, \ell)\|_{L^\infty(0,T;H^{\frac{3}{2}}(\omega))} + \|\partial_{x_d} u(\cdot, \cdot, \ell)\|_{W^{1,\infty}(0,T;H^\delta(\omega))} \right) \end{aligned}$$

Combining this with (3.2) and (3.7), we find

$$\begin{aligned} \|f(t, \cdot)\|_{L^2(\omega)} & \leq C(\|\partial_{x_d} u(\cdot, \cdot, \ell)\|_{L^\infty(0,T;H^{\frac{3}{2}}(\omega))} + \|\partial_{x_d} u(\cdot, \cdot, \ell)\|_{W^{1,\infty}(0,T;H^\delta(\omega))}) \\ & \quad + C \int_0^t (t-s)^{\alpha(1-\gamma)-1} \|f\|_{L^\infty(0,s;L^2(\omega))} ds, \quad t \in (0, T), \end{aligned}$$

which clearly implies

$$\begin{aligned} \|f\|_{L^\infty(0,t;L^2(\omega))} & \leq C(\|\partial_{x_d} u(\cdot, \cdot, \ell)\|_{L^\infty(0,T;H^{\frac{3}{2}}(\omega))} + \|\partial_{x_d} u(\cdot, \cdot, \ell)\|_{W^{1,\infty}(0,T;H^\delta(\omega))}) \\ & \quad + C \int_0^t (t-s)^{\alpha(1-\gamma)-1} \|f\|_{L^\infty(0,s;L^2(\omega))} ds, \quad t \in (0, T). \end{aligned}$$

Then, [7, Lemma 3] (see also [33, Theorem A]), implies that

$$\|f\|_{L^\infty(0,t;L^2(\omega))} \leq C(\|\partial_{x_d} u(\cdot, \cdot, \ell)\|_{L^\infty(0,T;H^{\frac{3}{2}}(\omega))} + \|\partial_{x_d} u(\cdot, \cdot, \ell)\|_{W^{1,\infty}(0,T;H^\delta(\omega))}), \quad t \in (0, T)$$

from which we deduce (1.17). \square

3.1. Application to the recovery of coefficients. Consider v the solution of the problem

$$\left\{ \begin{array}{l} \partial_t^\alpha v + \mathcal{A}v + q(t, x')v = 0, \quad (t, x) \in (0, T) \times \tilde{\Omega}, \\ v(t, x', \ell) = h_0(t, x') \quad (t, x') \in (0, T) \times \omega, \\ v(t, x) = h_1, \quad (t, x) \in (0, T) \times \partial\omega \times (-\ell, \ell), \\ \partial_{x_d} v(t, x', -\ell) = 0 \quad (t, x') \in (0, T) \times \omega, \\ v(0, x) = w_0(x), \quad x \in \tilde{\Omega}, \end{array} \right.$$

with $w_0, h_k, k = 0, 1$, given by (1.19). Then, we can write $v = H + y$ where y solves

$$\left\{ \begin{array}{l} \partial_t^\alpha y + \mathcal{A}y + q(t, x')y = H_1(t, x), \quad (t, x) \in (0, T) \times \tilde{\Omega}, \\ y(t, x) = 0, \quad (t, x) \in (0, T) \times \partial\omega \times (0, \ell), \\ y(t, x', \ell) = 0 \quad (t, x') \in (0, T) \times \omega, \\ \partial_{x_d} y(t, x', 0) = 0 \quad (t, x') \in (0, T) \times \omega, \\ y(0, x) = 0, \quad x \in \tilde{\Omega}, \end{array} \right.$$

where $H_1 = -(\partial_t^\alpha + \mathcal{A} + q)H \in \mathcal{C}([0, T]; H^{2\gamma}(\tilde{\Omega})) \cap W^{1,p}(0, T; L^2(\tilde{\Omega}))$. Therefore, using the fact that, thanks to (1.19), we have

$$H_1(0, x) = 0, \quad x \in \tilde{\Omega}$$

and repeating the arguments of the preceding section, we can prove that $v \in \mathcal{C}([0, T]; H^{2(1+\gamma)}(\tilde{\Omega}))$ and for $d \leq 3$, the Sobolev embedding theorem implies that $v, \partial_{x_d} v \in \mathcal{C}([0, T] \times \bar{\Omega})$. Applying the previous results about recovery of source terms we can complete the proof of Corollary 1.5.

Proof of Corollary 1.5. Let $u = v_1 - v_2$ and notice that u solves (1.2) with $q = q_1$, $f = (q_2 - q_1)$ and $R = v_2$. Then, using the fact that $v_2, \partial_{x_d} v_2 \in \mathcal{C}([0, T] \times \bar{\Omega})$ and the fact that, thanks to (1.13), (1.20), we are in position to apply Theorem 1.4 from which we deduce (1.20). \square

APPENDIX

We consider the following result which can be deduced from other known results (see e.g [30, Theorem 2.2] and [23, Theorem 1.3]) considered for $\alpha \in (0, 1)$ that we extend to $\alpha \in (0, 2)$.

Lemma 3.1. *Let $F \in L^2(Q)$, $\alpha \in (0, 2)$ and $m, n = 0, 1$. Then problem (1.2)-(1.4) admits a unique solution $u \in L^2(0, T; H^1(\Omega))$, satisfying $\mathcal{A}u, \partial_t^\alpha u \in L^2(Q)$ and the following estimate holds true*

$$\|\partial_t^\alpha u\|_{L^2(Q)} + \|\mathcal{A}u\|_{L^2(Q)} + \|u\|_{L^2(Q)} \leq C \|F\|_{L^2(Q)}. \quad (3.8)$$

Proof. We prove this result for sake of completeness. We fix A the operator \mathcal{A} acting on $L^2(\Omega)$ with the boundary condition (1.3)-(1.4). We fix also the non-decreasing sequence of non-negative eigenvalues $(\lambda_k)_{k \geq 1}$ and associated eigenfunctions $(\varphi_k)_{k \geq 1}$ of A . Then, we consider

$$k_0 := \min\{k \in \mathbb{N} : \lambda_k > 0\}.$$

Since u solves (1.2)-(1.4), we have

$$u_k(t) := \langle u(t, \cdot), \varphi_k \rangle_{L^2(\Omega)} = \int_0^t (t-s)^{\alpha-1} E_{\alpha, \alpha}(-\lambda_k(t-s)^\alpha) F_k(s) ds,$$

where $F_k(t) := \langle F(t, \cdot), \varphi_k \rangle_{L^2(\Omega)}$. Therefore, we have

$$u_k(t) = (G_k \mathbf{1}_{(0, T)}) * (F_k \mathbf{1}_{(0, T)})$$

with $G_k(t) = t^{\alpha-1} E_{\alpha, \alpha}(-\lambda_k t^\alpha)$. Then, we find

$$\int_0^T \sum_{k=1}^{\infty} \lambda_k^2 |u_k(t)|^2 dt = \sum_{k=1}^{\infty} \lambda_k^2 \|(G_k \mathbf{1}_{(0, T)}) * (F_k \mathbf{1}_{(0, T)})\|_{L^2(0, T)}^2$$

and an application of the Young inequality yields

$$\int_0^T \sum_{k=1}^{\infty} \lambda_k^2 |u_k(t)|^2 dt = \sum_{k=1}^{\infty} \lambda_k^2 \|G_k\|_{L^1(0, T)}^2 \|F_k\|_{L^2(0, T)}^2.$$

On the other hand, we have

$$\|G_k\|_{L^1(0, T)} = \int_0^T s^{\alpha-1} E_{\alpha, \alpha}(-\lambda_k s^\alpha) ds = \lambda_k^{-1} \int_0^{\lambda_k^{-\frac{1}{\alpha}} T} s^{\alpha-1} E_{\alpha, \alpha}(-s^\alpha) ds, \quad k \geq k_0.$$

Combining this with formula (1.148) of [29, Theorem 1.6], we get

$$\|G_k\|_{L^1(0,T)} \leq C\lambda_k^{-1} \int_0^{+\infty} \frac{s^{\alpha-1}}{1+s^{2\alpha}} ds \leq C\lambda_k^{-1}, \quad k \geq k_0 + 1.$$

It follows that

$$\begin{aligned} \int_0^T \sum_{k=1}^{\infty} (1 + \lambda_k^2) |u_k(t)|^2 dt &= C \int_0^T \sum_{k=1}^{k_0} |u_k(t)|^2 dt + \int_0^T \sum_{k=k_0+1}^{\infty} (1 + \lambda_k^2) |u_k(t)|^2 dt \\ &\leq C \sum_{k=1}^{\infty} \|F_k\|_{L^2(0,T)}^2 \leq C \|F\|_{L^2(Q)}^2 \end{aligned}$$

and we get

$$\|u\|_{L^2(0,T;D(A))} \leq C \|F\|_{L^2(Q)}.$$

In the same way, we find

$$\partial_t^\alpha u = -\mathcal{A}u + F \in L^2(Q)$$

which implies at the same time that $\partial_t^\alpha u \in L^2(Q)$ and (3.8). This proves (3.8). \square

ACKNOWLEDGEMENTS.

The work of the first author is partially supported by the French National Research Agency ANR (project MultiOnde) grant ANR-17-CE40-0029. The second author is supported by Grant-in-Aid for Scientific Research (S) 15H05740 of Japan Society for the Promotion of Science and by The National Natural Science Foundation of China (no. 11771270, 91730303), and the "RUDN University Program 5-100".

REFERENCES

- [1] G. BAO, G. HU, Y. KIAN, T. YIN, *Inverse source problems in elastodynamics*, Inverse Problems, **34** (4) (2018), 045009.
- [2] N. YA. BEZNOSHCHENKO, *Determination of coefficients of higher terms in a parabolic equation*, Siberian Math. J., **16** (1975), 360-367.
- [3] A. BUKHGEIM AND M. KLIBANOV, *Global uniqueness of a class of multidimensional inverse problem*, Sov. Math.-Dokl., **24** (1981), 244-247.
- [4] M. CHENG, J. NAKAGAWA, M. YAMAMOTO, T. YAMAZAKI, *Uniqueness in an inverse problem for a one dimensional fractional diffusion equation*, Inverse Problems, **25** (2009), 115002.
- [5] M. CHOULLI AND M. YAMAMOTO, *Some stability estimates in determining sources and coefficients*, J. Inverse Ill-Posed Probl., **12** (3) (2004), 233-243.
- [6] M. CHOULLI AND M. YAMAMOTO, *Some stability estimates in determining sources and coefficients*, J. Inverse Ill-Posed Probl., **14** (4) (2006), 355-373.
- [7] K. FUJISHIRO AND Y. KIAN, *Determination of time dependent factors of coefficients in fractional diffusion equations*, Math. Control Related Fields, **6** (2016), 251-269.
- [8] P. GAITAN AND Y. KIAN, *A stability result for a time-dependent potential in a cylindrical domain*, Inverse Problems **29** (6) (2013), 065006.

- [9] P. GRISVARD, *Elliptic problems in nonsmooth domains*, Pitman, London, 1985.
- [10] G. HU, Y. KIAN, *Uniqueness and stability for the recovery of a time-dependent source and initial conditions in elastodynamics*, preprint, arXiv:1810.09662.
- [11] G. HU, Y. KIAN, P. LI, Y. ZHAO, *Inverse moving source problems in electrodynamics*, preprint, arXiv:1812.06215.
- [12] O. YU. IMANUVILOV AND M. YAMAMOTO, *Lipschitz stability in inverse parabolic problems by the Carleman estimate*, *Inverse Problems*, **14** (1998), 1229-1245.
- [13] V. ISAKOV, *Inverse source problems*, Mathematical surveys and monographs, AMS, 1990.
- [14] D. JIANG, Z. LI, Y. LIU, M. YAMAMOTO, *Weak unique continuation property and a related inverse source problem for time-fractional diffusion-advection equations*, *Inverse Problems*, **33** (2017), 055013.
- [15] B. JIN AND W. RUNDELL, *An inverse problem for a one-dimensional time-fractional diffusion problem*, *Inverse Problems*, **28** (2012) 75010-75028 (19pp).
- [16] B. JIN AND W. RUNDELL, *A tutorial on inverse problems for anomalous diffusion processes*, *Inverse problems*, **31** (2015), 035003.
- [17] Y. KIAN, L. OKSANEN, E. SOCCORSI, AND M. YAMAMOTO, *Global uniqueness in an inverse problem for time-fractional diffusion equations*, *J. Diff. Equat.*, **264** (2018), 1146-1170.
- [18] Y. Kian, D. Sambou and E. Soccorsi, *Logarithmic stability inequality in an inverse source problem for the heat equation on a waveguide*, to appear in *Applicable Analysis*.
- [19] Y. KIAN, E. SOCCORSI, M. YAMAMOTO, *On time-fractional diffusion equations with space-dependent variable order*, *Annales Henri Poincaré*, **19** (2018), 3855-3881.
- [20] Y. KIAN, M. YAMAMOTO, *On existence and uniqueness of solutions for semilinear fractional wave equations*, *Fract. Calc. Appl. Anal.*, **20** (2017), 117-138.
- [21] A.A. KILBAS, H.M. SRIVASTAVA AND J.J. TRUJILLO, *Theory and applications of fractional differential equations*, Elsevier, Amsterdam, 2006.
- [22] C-K. KIM, *An analytical solution to heat conduction with a moving heat source*, *Journal of Mechanical Science and Technology*, **25** (4) (2011), 895-899.
- [23] Z.LI, Y. KIAN, E. SOCCORSI, *Initial-boundary value problem for distributed order time-fractional diffusion equations*, preprint, arXiv:1709.06823.
- [24] J.-L. LIONS AND E. MAGENES, *Non-homogeneous Boundary Value Problems and Applications*, Vol. I, Springer-Verlag, Berlin, 1972.
- [25] Y. LIU, W. RUNDELL AND M. YAMAMOTO, *Strong maximum principle for fractional diffusion equations and an application to an inverse source problem*, *Fract. Calc. Appl. Anal.*, **19** (2016), 888-906.
- [26] F. MAINARDI, *On the initial value problem for the fractional diffusion-wave equation*, in: S. Rionero, T. Ruggeri (Eds.), *Waves and Stability in Continuous Media*, World Scientific, Singapore, 1994, pp. 246-251.
- [27] R. METZLER AND J. KLAFTER, *The random walk's guide to anomalous diffusion: a fractional dynamics approach*, *Physics Reports*, **339** (2000) 1-77.
- [28] J. NAKAGAWA, K. SAKAMOTO, M. YAMAMOTO, *Overview to mathematical analysis for fractional diffusion equations- new mathematical aspects motivated by industrial collaboration*, *Journal of Math-for-Industry*, Vol.2 (2010A-10), 99-108.
- [29] I. PODLUBNY, *Fractional differential equations*, Academic Press, San Diego, 1999.
- [30] K. SAKAMOTO AND M. YAMAMOTO, *Initial value/boundary value problems for fractional diffusion-wave equations and applications to some inverse problems*, *J. Math. Anal. Appl.*, **382** (2011), 426-447.
- [31] W. RUNDELL, Z. ZHANG, *Recovering an unknown source in a fractional diffusion problem*, *Journal of Computational Physics Volume*, **368** (2018), 299-314.

- [32] I.M. SOKOLOV, J. KLAFTER, A. BLUMEN, *Fractional kinetics*, Physics Today, **55** (2002), 48-54.
- [33] H. YE , J. GAO, Y. DING, *A generalized Gronwall inequality and its application to a fractional differential equation*, J. Math. Anal. Appl., **328** (2007), 1075-1081.
- [34] M. Yamamoto, *Stability, reconstruction formula and regularization for an inverse source hyperbolic problem by control method*, Inverse Problems, **11** (1995), 481-496.
- [35] M. YAMAMOTO, *Conditional stability in the determination of force terms of heat equations in a rectangle*, Mathematical and Computer Modelling **18** (1993), 79-88.
- [36] M. YAMAMOTO, *Conditional stability in the determination of densities of heat sources in a bounded domain*, Estimation and Control of Distributed Parameter Systems (W. Desch, F. Kappel and K. Kunisch), Birkhauser Verlag, Basel, 1994, 359-370.

AIX MARSEILLE UNIV, UNIVERSITÉ DE TOULON, CNRS, CPT, MARSEILLE, FRANCE

E-mail address: yavar.kian@univ-amu.fr

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO, 3-8-9 KOMABA MEGURO, TOKYO 153-8914, JAPAN, PEOPLES' FRIENDSHIP UNIVERSITY OF RUSSIA (RUDN UNIVERSITY) 6 MIKLUKHO-MAKLAYA ST, MOSCOW, 117198, RUSSIAN FEDERATION

E-mail address: myama@ms.u-tokyo.ac.jp