Chloroquine and Chloroquinoline Derivatives as Models for the Design of Modulators of Amyloid Peptide Precursor Metabolism

Patricia Melnyk, Valérie Vingtdeux, Stéphane Burlet, Sabiha Eddarkaoui, Marie-Eve Grosjean, Paul-Emmanuel Larchanché, Guillaume Hochart, Christian Sergheraert, Cecilia Estrella, Mathieu Barrier, et al.

► To cite this version:

Patricia Melnyk, Valérie Vingtdeux, Stéphane Burlet, Sabiha Eddarkaoui, Marie-Eve Grosjean, et al.. Chloroquine and Chloroquinoline Derivatives as Models for the Design of Modulators of Amyloid Peptide Precursor Metabolism. ACS Chemical Neuroscience, 2015, 6 (4), pp.559-569. 10.1021/cn5003013 . hal-02060814

HAL Id: hal-02060814

<https://hal.science/hal-02060814>

Submitted on 7 Mar 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Chloroquine and chloroquinoline derivatives as models for the design of modulators of amyloid peptide precursor metabolism

Patricia Melnyk,^{1,2,3} Valérie Vingtdeux,^{1,4} Stéphane Burlet,⁵ Sabiha Eddarkaoui,^{1,4} Marie-Eve Grosjean,^{1,4} Paul-Emmanuel Larchanché,^{1,2} Guillaume Hochart,^{1,2,3} Christian Sergheraert,¹ Cecilia Estrella,⁵ Mathieu Barrier,⁵ Virginie Poix,⁵ Pauline Plancq,⁵ Cécile Lannoo,⁵ Malika Hamdane,^{1,4} André Delacourte,^{1,4} Philippe Verwaerde,⁵ Luc Buée,^{1,4} and Nicolas Sergeant^{1,4}*

¹ Université de Lille, F-59000 Lille, France

² UDSL, EA 4481, UFR Pharmacie, F-59000 Lille, France

³ CNRS UMR8161, F-59000 Lille, France

⁴ Inserm U837, Jean-Pierre Aubert Research Center, F-59000 Lille, France

⁵ AlzProtect, F-59120 Loos, France

ABSTRACT

The amyloid precursor protein (APP) plays a central role in Alzheimer's disease (AD). Preventing deregulated APP processing by inhibiting amyloidogenic processing of carboxy-terminal fragments (APP-CTFs), and reducing the toxic effect of amyloid beta (A β) peptides remain an effective therapeutic strategy. We report the design of piperazine-containing compounds derived from chloroquine structure and evaluation of their effects on APP metabolism and ability to modulate processing of APP-CTF and production of A β peptide. Compounds which retained alkaline properties and high affinity for acidic cell compartments were the most effective. The present study demonstrates that (1) the amino side chain of chloroquine can be efficiently substituted by a bis(alkylamino)piperazine chain, (2) the quinoline nucleus can be replaced by a benzyle or a benzimidazole moiety, and (3) pharmacomodulation of the chemical structure allows the redirection of APP metabolism toward a decrease in A β peptide release, and increased stability of APP-CTFs and amyloid intracellular fragment. Moreover, the benzimidazole compound **29** increases APP-CTFs *in vivo* and shows promising activity by the oral route. Together, this family of compounds retains a lysosomotropic activity which inhibits lysosome-related A β production, and is likely to be beneficial for therapeutic applications in AD.

KEY WORDS: Chloroquine, piperazine, Alzheimer's disease, Amyloid Precursor Protein

INTRODUCTION

For decades, chloroquine (CQ) has been one of the two most widely used antimalarial drugs with moderate acute toxicity. It is known to cross the BBB.¹ Following a repositioning strategy, CQ and CQ-derived compounds have already been evaluated in several biological applications such as for prion disease,²⁻⁵ HCV^{6,7} and even cancer.^{8,9} Different mechanisms have been proposed for these applications. Unfortunately, the toxicity of CQ precludes its use in long-term treatments. However, CQ-derived compounds such as hydroxychloroquine are administered, for instance, for the treatment of systemic lupus erythematosus¹⁰ or rheumatoid polyarthritis.¹¹ CQ derivatives may therefore maintain the activity with reduced side effects. For that purpose, we previously developed a library of CQ-derivatives among, from which some compounds demonstrated antimalarial properties and reduced cytotoxic effects upon MRC-5 cells (human diploid embryonic lung cell line).¹²⁻¹⁴ Based on the recent idea that Alzheimer's disease (AD) could be driven by a prion-like spread of protein aggregates in the brain,^{15,16} some compounds of this library have also been positively evaluated for their anti-prion properties.⁵ Recently, we showed that CQ had an indirect modulatory effect on APP metabolism,¹⁷ the deregulation of which is central to AD pathophysiology.¹⁸ The role of CQ in AD has also been controversially emphasized by others in the literature.^{19,20}

According to the amyloid hypothesis, APP is expected to play a key role in AD. APP metabolism leads to the production and release of amyloid-beta peptides (A β), the major component of the amyloid deposits being its 42 aa form. The proteolytic processing of APP brings into play a sequence of cleavages involving either α - or β -secretase at the first step, followed by γ -secretase (Figure 1¹⁶).²¹ The α -secretase cleaves APP within the A β sequence, producing a soluble fragment (sAPP α) and a carboxy-terminal fragment (CTF α), and inhibits A β production.²² Through the

amyloidogenic pathway, β -secretase cleaves APP at the first amino acid residue of the A β sequence, producing a soluble fragment (sAPP β) and a transmembrane carboxy-terminal fragment (CTF β). The APP intracellular domain (AICD) is released by cleavage of CTF β at the ϵ -site by γ -secretase.²³ The ϵ -processed CTF β stub is subsequently cleaved at γ -sites by γ -secretase to generate A β peptides ranging from 46 to 36 amino acids in length, following stepwise successive cleavages along multiple interactive pathways.²⁴⁻²⁶

Current drug candidates for the treatment of amyloid pathology mainly target β - or γ -secretase, in order to avoid the production of A β peptides. Most of them have failed, essentially due to low BBB permeability or severe side effects.²² Given that the mechanism involved in APP processing is now known to be shared by a growing list of type I transmembrane proteins such as Notch, and that β - and γ -secretase have multiple substrates engaged in ubiquitous cellular and tissue functions, their inhibition may have deleterious effects, and more selective inhibition of APP processing by both proteases is a potential alternative.^{27,28} One way to circumvent these side effects of highly potent β - or γ -secretase inhibitors by using dual inhibitors was recently proposed by Strömberg *et al.*²⁹ With regards to γ -secretase, several compounds have been shown (1) to repress or reduce A β production, especially A β species longer than 40 aa, (2) to maintain AICD release and (3) to spare other γ -secretase substrates. The so-called γ -secretase modulators (GSMs) are small molecules demonstrated to act directly on γ -secretase activity, resulting in a decreased level of long A β (1-40 and 1-42) and an increased level of short A β peptides (1-37 and 1-38). We previously showed that CQ inhibits A β production, whereas levels of other APP metabolites such as APP-CTFs and AICD are maintained and even increased, and γ -secretase cleavage of Notch remains unmodified.^{17,30} Thus CQ can be considered as an indirect γ -secretase modulator since the lysosome is a compartment where A β is produced.³¹

Here, we have screened our library of CQ-derivatives in order to identify a family of compounds showing an improved effect on APP processing and lower or limited toxicity.¹²⁻¹⁴ Our objective was to identify a compound that can redirect APP metabolism towards the non-amyloidogenic pathway, *i.e.* that can decrease the level of long A β peptides more efficiently than short ones, and increase the level of AICD and sAPP α , with a higher efficiency than CQ or other lysosomotropic compounds.¹⁸ For that purpose, we first focused on quinoline compounds derived from CQ as the starting point, and increased the ability of compounds to accumulate in acidic vesicles such as lysosomes by adding a bis(alkylamino)piperazine side chain. Secondly, compounds in which the quinoline moiety was replaced by various heterocyclic or aromatic substructures were studied, with an aliphatic amino chain providing the best efficiency. Finally, we evaluated the importance of the symmetry of the compounds (Figure 2). Here, we identified a series of molecules based on a CQ-derived structure, but with a chemically different structure and improved efficiency. Moreover, among those molecules one had low toxicity and provided an interesting *in vivo* proof-of-concept.

RESULTS AND DISCUSSION

The synthetic pathways of chloroquinoline-piperazine derived compounds, heterocyclic and most benzyl-piperazine derived compounds and symmetric compounds have already been detailed, and compounds **1-34**, **37**, **39**, **40**, **42-47** have been described.^{12-14,32} In addition to previously described mono- or dibenzyl derived compounds, fluorobenzyl or thiazolyl compounds **35**, **36** and **41** were easily synthesized by reductive amination according to Scheme 1.

All compounds were tested for their potential interference with APP processing in an SY5Y human neuroblastoma cell line stably expressing the neuronal isoform of human wild-type APP695

(SY5Y-APP^{wt}), a well-established cellular model for the study of APP metabolism (Figure 3).^{17,33} Thus, to analyze the effect of our compounds on APP metabolism, SY5Y-APP^{wt} were treated or not with 1 and 5 μ M of compounds compared to the control condition or to the treatment of SY5Y-APP^{wt} with 1 or 5 μ M of CQ for 6 and 24 hours (Figure 3). Small proteolytic fragments of APP catabolism were resolved by 1D Tris-Tricine gels and detected with our well-characterized APP-CterC17 antiserum^{17,33} Identity of APP fragments was previously established by 2D Western-blot.³³ As shown on Figure 3, all APP fragments including APP-CTFs derived from the α - and β -cleavages are detected as well as the AICD. The exposure of 5 min enabled to quantify the APP-CTFs whilst following 30 min of exposure AICD signal was quantified. Following treatment with 1 or 5 μ M of compound **33**, α - and β -CTFs amounts were double when compared to CQ treatment (Figure 3, upper Western-blot). AICD production increased fivefold at 5 μ M of compound **33** following 24 hours of treatment (Figure 3, AICD panel). AICD C₂ and C₁₀ indicate the concentrations of compound necessary to double (C₂) or increase by 10-fold (C₁₀) the quantity of AICDs produced compared with untreated control conditions. These were calculated based on the measure of AICD following a wider range of concentrations.

Cytotoxicity and A β ₁₋₄₂ levels, determined for all compounds, were defined as critical parameters and were measured using *in vitro* assays. All values are reported in Tables 1, 2 and 3.

Cytotoxicity is expressed as the compound concentration causing 50% cell death (CC₅₀). IC₅₀ indicates the concentration of compound capable of inhibiting secretion of A β ₁₋₄₂ by 50% (Tables 1-3). For the most potent compounds, CTF α , secreted A β ₁₋₃₈, A β ₁₋₄₀ and sAPP α concentrations were also determined (Table 4). A preliminary ADME (Absorption, Distribution, Metabolism, Excretion) evaluation was performed (Table 5), and finally, the most potent compound was selected for acute *in vivo* administration by the oral route (Figure 4).

For the reference compound CQ, the AICD C₂ and C₁₀ were 1.4 μ M and 10.8 μ M respectively (Table 1). The AICD C₂ value of quinacrine 1.0 μ M was comparable to that of CQ, but its high cytotoxicity to SY5Y-APP^{wt} cells precluded the evaluation of other parameters. With the exception of compounds **5** and **8**, the AICD C₂ values of amide derivatives were comparable to that of CQ (1-5 μ M), although AICD C₁₀ values for most compounds were higher than that of CQ (17-40 μ M). For aromatic amides, substitution was allowed, but the presence of an electron-donating group was probably unfavorable for activity, although the permeability of cells to these compounds was not measured. In the case of aliphatic amides, *tert*-butyl amide **6** displayed the same activity as CQ, whereas the others compounds were less effective. The ability of amides to decrease the amount of secreted A β ₁₋₄₂ was comparable to that of CQ, but slightly improved in the case of aromatic or bulky hydrophobic substituents.

Overall, secondary and tertiary amines (series **Ib** and **Ic**) were more effective than their amide counterparts. However, amines were also more toxic with the exception of the cyclopropyl derivative **15**. Because of their cytotoxicity to SY5Y-APP^{wt} cells, as already shown for other cell lines,¹² the activity of monobenzyl amines **9-11** (except compound **12**) and dibenzyl derivatives **17-20** could not be further tested in this assay. The ability of secondary amines to double the amount of AICD fragment was in the same range as for CQ (1-3 μ M), although the AICD C₁₀ of these compounds was often higher (except for compound **15**). We observed that the activity and cytotoxicity on SY5Y-APP^{wt} cells of amines were highly dependent on their structure. Indeed, steric hindrance might play an important role since the activity of the *tert*-butyl derivative **14** was comparable to that of CQ, whereas compound **13** was less active. In this series, benzyl or *tert*-butyl substituents favored the decrease of A β ₁₋₄₂ production. Generally, the activity of tertiary

amines (series **Ic**) was higher than that of secondary amines. In tertiary amines, steric hindrance seemed to be important, since substitution with cyclopropylmethyl (compound **23**) or isobutyl (compound **24**) increased their activity. It was also worth noting the difference between the cyclic amines **25** and **26** in terms of cytotoxicity on SY5Y-APP^{wt} cells, whereas their influence on AICD production remained comparable, even though the A β ₁₋₄₂ IC₅₀ was four times higher for the piperidine derivative (1.0 μ M for **25** and 4.0 μ M for **26**).

It is generally accepted that the increased basicity of compounds leads to better uptake by acidic compartments owing to the pH gradient between the cytosol and the compartment. Using CQ as a starting point, we evaluated derivatives with an increased ability to accumulate in acidic vesicles by adding a bis(alkylamino)piperazine side chain. In Series **I**, compounds **9-13** and **14-19** were chosen to evaluate the influence of a supplementary basic nitrogen, while compounds **1-8** could be compared to CQ. The ability of these compounds to accumulate in acidic cell compartments was estimated by calculating a lysosome accumulation ratio (LAR), similar to the antimalarial accumulation ratio in parasite food vacuoles (VAR). LAR could be determined according to Equation 1,³⁴ based on a weak base model, proceeding from a derivation of the Henderson-Hasselbach equation in previous work by Hawley *et al.*³⁵ As expected, amides **1-8** displayed an LAR similar to that of CQ (LAR = 30 - 50.10³) but amines **9-16** and **17-24** showed a 200-fold increase in accumulation (LAR = 56 - 60.10⁵). This seemed to correlate with the comparable activity of amides **1-8** with CQ, whereas tertiary amines showed a greater influence on APP metabolism. Secondary amines, with a higher LAR compared to CQ, also showed a comparable influence on APP metabolism (AICD and A β ₁₋₄₂ secretion). It is worth noting that an aliphatic substitution of tertiary amines has little influence on AICD and A β ₁₋₄₂ secretion (except compound **22**), but increases cytotoxicity by around ten times.

The *diisobutylamino* side chain was chosen for further studies. We aimed to evaluate the role of the quinoline moiety, as we have previously done for antimalarial activity,¹³ for a family of analogs in which a common N₁,N₁-*diisobutyl*-1,4-bis(3-aminopropyl) piperazine motif was linked to a variety of aromatic entities.

Two series of compounds: heteroarylamines (compounds **27-32**, series **IIa**) and benzylamines (compounds **33-37**, series **IIb**) were selected in order to simultaneously study the influence of the nature of the chemical link on cytotoxicity, the accumulation of AICDs and the secretion of A β ₁₋₄₂ (Table 2). With regard to the replacement of the quinoline moiety by a heterocycle, benzimidazole was the most efficient in order to maintain an effect on APP metabolism compared to compound **24**. An acridine nucleus was selected for comparison with quinacrine but no results could be obtained because of its cytotoxicity. In the case of the benzoxazole ring (compound **28**), the compound decreased AICD level at a concentration of 5.0 μ M. Substitution with other heterocycles such as pyrazine (compound **31**) led to lower activity, and AICD C₁₀ could not be determined. The influence of the substitution on A β ₁₋₄₂ peptide secretion could not be measured in any of these cases. In the case of the pyrimidine and purine derivatives (compounds **30** and **32**), no activity at concentrations up to 10 μ M could be detected, with regard to either AICD fragment production or A β ₁₋₄₂ secretion.

The chloroquinoline nucleus could also be replaced by a benzyl moiety, which induced a slight decrease in the production of AICDs and A β ₁₋₄₂ (Table 2). The ability of methoxy **33**, chloro **34** or fluoro **35** derivatives to double or increase 10-fold the quantity of AICD fragments generated was in the same range as CQ (C₂ = 1 - 3 μ M and C₁₀ = 10 - 16 μ M). On the other hand, the nature of the substituent on the phenyl ring modified the quantity of secreted A β ₁₋₄₂, resulting in a comparable IC₅₀ for compounds **33** and **34**, whereas compound **35** did not induce a decrease in

A β ₁₋₄₂ secretion at 5 μ M. Conversely, compound **37**, with a pyridine ring, was found to be slightly less active in terms of AICD production than the phenyl compounds, but no effect could be detected on secreted A β ₁₋₄₂ at concentrations below 5 μ M. Surprisingly, when the phenyl ring was replaced by a thiazole ring (compound **36**), no effect was observed on the secretion of A β ₁₋₄₂, and AICD production was shown to decrease at concentrations below 5 μ M, in contrast to the other compounds of this series.

Finally, we evaluated the influence of compound symmetry (Table 3) using alkyl- or benzyl-type substituents. Except in the case of derivatives **44**, **45** and **46**, all the compounds showed higher cytotoxicity on SY5Y-APP^{wt} cells than previous series, making it difficult to evaluate their activity in this cellular model. Commercially available unsubstituted diamine, **38**, had no influence on AICD levels or secreted A β ₁₋₄₂. With benzyl substituents, at the non-cytotoxic concentration of 5 μ M, an increase in A β secretion was observed without affecting AICD levels. Surprisingly, *p*-chlorobenzyl derivative **40** behaved differently and was one of the most effective compounds both with regard to the production of AICD ($C_2 = 0.5 \mu$ M) and the secretion of A β ₁₋₄₂ ($IC_{50} = 3.1 \mu$ M), as seen for Series II. For the dialkyl compounds **45** and **46**, a significant increase in A β ₁₋₄₂ secretion was measured, with little impact on AICD levels. In contrast, the tetra-*isobutyl* compound **47** showed a comparable impact to CQ on APP metabolism ($C_2 = 3.7 \mu$ M for **47**, 1.4 μ M for CQ), but no influence on the secretion of A β ₁₋₄₂ at the concentration of 5 μ M.

The influence on accumulation in acidic compartments in this series did not seem to be critical, since compounds with a comparable theoretical LAR showed widely differing effects on APP metabolism. For instance, with similar LAR values of 15.10⁶, the presence of a *p*-fluorobenzyl group (compound **35**), increased the secretion of A β ₁₋₄₂, but did not significantly modify AICD levels compared to a *p*-methoxy or *p*-chlorobenzyl group (compounds **33** and **34**). In this series,

most compounds did not modify the secretion of A β ₁₋₄₂ at concentrations below 5 μ M. In the case of symmetric compounds (Series 3), except for bis(*p*-chlorobenzyl) or bis(*diisobutyl*) compounds **40** and **47**, no influence on the quantity of AICD could be measured, but these compounds led to an increase in the secretion of A β ₁₋₄₂ at concentrations below 5 μ M. Once again, the ability to be accumulated in lysosomes/endosomes could not be considered, since the calculated LARs were comparable (LAR = 15.10⁶). Thus, LAR values do not seem to be highly relevant or even the only criteria in explaining the relative influence of these compounds on APP metabolism compared to CQ.

Given the interest of quinoline **25**, benzimidazole **29** and benzyle **33** compounds demonstrated above, further experiments were designed to evaluate their impact on the metabolism of APP in greater detail. The neurotrophic fragments sAPP α and CTF α ³⁶, as well as the shorter A β ₁₋₃₈ and A β ₁₋₄₀ peptides, were quantified (Table 4). The closest analogs of CQ **25** and benzimidazole **29** showed the greatest effects on APP metabolism, with a CTF α C₂ value of 0.2 and 0.5 μ M respectively and a CTF α C₁₀ close to 10 μ M. Substitution with a methoxybenzyle moiety decreased the production of CTF α (C₂ = 6.7 μ M and C₁₀ = 28.4 μ M). Surprisingly, the levels of CTF α and sAPP α seemed uncorrelated for compound **25**. At a concentration of 10 μ M, which increased the quantity of CTF α produced ten times, sAPP α was only increased by 30%. At this concentration, sAPP α was increased by 78% for benzimidazole **29** and 55% for methoxybenzyle compound **33**. Since the ability of the three compounds **25**, **29** and **33** to decrease the secretion of A β ₁₋₄₂ was equivalent (IC₅₀ close to 1 μ M), benzimidazole **29** appeared more effective with regard to A β ₁₋₄₀, with an IC₅₀ of 2.0 μ M, compared to **33**, with a value of 5.1 μ M and **25**, with a value of 9.1 μ M. Interestingly, the influence of these compounds on the secretion of the short A β ₁₋₃₈ peptide was

also very different, as the IC_{50} was 5.0 μM for compounds **25** and **29**, whereas compound **33** seemed to be inactive at concentrations up to 10 μM . These two compounds were also able to inhibit the secretion of $A\beta_{1-38}$ and $A\beta_{1-40}$ more effectively. The activity of these three compounds on the metabolism of APP was also confirmed by measuring the secretion of sAPP α , a fragment previously described as being neuroprotective.³⁶

The mechanism of action of these compounds had to be deciphered. Starting from CQ, the introduction of a polyamino side chain should have led to more efficient accumulation in acidic vacuoles. Interestingly, however, when evaluated as antimalarials, these compounds were shown not to accumulate in parasite food vacuoles.³⁷ With respect to their influence on the secretion of $A\beta$ peptides of different lengths, the three compounds of interest seemed to have different or complementary mechanisms of action, especially with regard to LAR values. However, a lysosomotropic activity of these compounds cannot be ruled-out. Regarding the possible effect of AICD, whether as a transcriptional gene regulator or a toxic effect, the AICD must reach the cell nucleus. The transcriptional active AICD is supposed to originate from the β -CTF.³⁸ This mechanism supposes that a γ -secretase cleavage of the β -CTF occurs rapidly along the early endosome / lysosome pathway. AICD can also be produced later in the endosome / lysosome route and being degraded by lysosome proteases³⁹ or secreted in the extracellular space via exosomes¹⁷. Consistent with this mechanism, accumulating evidences suggest that 40% of APP metabolism occurs in the late endosome / lysosome pathway.^{31,40} Cyclopamine reduces the $A\beta$ production and redirect part of the APP processing to the lysosome compartment⁴⁰ giving similar results to that obtain with our small compounds. This effect does not imply a modulation of the gamma-secretase activity. We also tested the *in vitro* γ -secretase activity as well as the γ -secretase processing of Notch (unpublished data). Both are not modified by our compounds. Interestingly, a nuclear

accumulation of AICD was not reported in these studies further supporting the hypothesis that AICD can be produced along the endosome / lysosome pathway giving rise to a transcriptionally active fragment when produced in early endosomal route or alternatively, AICD can also be produced in late endosome compartments such as multivesicular bodies.¹⁷ Absence of nuclear signal is not due to our APP-CterC17 antibody since our antibody used to visualize the nuclear localization of AICD.⁴¹ Taken together, our past and current results strongly suggest that AICD accumulation is related to a modification of the late endosome /lysosome processing of AICD and would therefore explain why a toxic effect of the accumulation of AICD is not herein observed whereas nuclear AICD signaling is reported to be toxic.⁴²⁻⁴⁴ Hence, in sharp contrast to cyclopamine, which is a highly teratogen drug, one of our lead compound has passed the preclinical phase and is currently in clinical phase I. However, further investigation should be done to decipher the precise molecular mechanism of action of our compounds.

Some preliminary ADME parameters were evaluated before an *in vivo* evaluation (Table 5). The three compounds were highly soluble in water (>200 μ M at pH 7.4) but showed different logD values, with a very low value of 0.1 for quinoline **25**, predicting a low probability of BBB penetration. In contrast, benzimidazole **29** and compound **33** displayed intermediate values of 1.8 and 1.2, respectively. Evaluation of metabolic stability using mouse and human liver microsomes showed high stability for compound **25** (100% of the compound remaining after 1 h) and benzimidazole **29** (79 and 52% of the compound remaining after 1 h). In contrast, compound **33** showed low stability (30% and 27% after 1 h). Compounds **25** and **33** were not evaluated in this study for an *in vivo* proof-of-concept.

Finally, to determine the modulatory effects of our compounds on APP metabolism *in vivo*, a single dose treatment was carried out in wild-type C57Bl6 mice (Figure 4). Physiologically, AICDs were rarely detected and assumed to be rapidly degraded. After a single administration of compound **29** by the oral route, mice were sacrificed 24 h later and quantification by western blotting highlighted a significant increase in CTF α level. A single dose of compound **29** led to an increase in CTF α in the frontal cortex from a 5 mg/kg dose, and in the hippocampus from a 12.5 mg/kg dose. A 25mg/kg dose was the most efficient and significant, with an increase of CTF α of 29% in the frontal cortex and 20% in the hippocampus. This first *in vivo* proof-of-concept should be confirmed by further long-term administration studies of compound **29**.

CONCLUSION

Previous studies have underlined the interest of CQ and alkaline drugs as AICD enhancers and compounds capable of interacting with the endosome/lysosome pathway, a mechanism that is of growing interest in neurodegenerative diseases including AD. The present study of several series of compounds showed that (1) the chemical structure of CQ could be efficiently modified by adding a bis(alkylamino)piperazine chain, (2) the quinoline nucleus could be replaced by a benzyle or benzimidazole moiety, and (3) pharmacomodulation of the chemical structure allowed the metabolism of APP to be directed toward the decrease of A β peptide secretion or the increase of APP-CTF fragments as well as the accumulation of AICDs. Among the compounds evaluated in this study, compounds **25**, **29** and **33** showed a better *in vitro* profile than CQ on the metabolism of APP. Preliminary ADME and *in vivo* evaluation underlined the possibility that compound **29** could find therapeutic applications in neurodegenerative diseases in which APP metabolism needs to be modulated. Thus benzimidazole **29** is currently under development as a potential anti-

Alzheimer drug candidate. From a more fundamental point of view, as the most potent molecules were structurally different from CQ, this raises the question of the mechanism of action as well as the potential specific biological targets of this family of molecules.

METHODS

Chemistry

Compounds **1-34**, **37-40** and **42-47** have already been described and were used at a purity of > 95%.¹²⁻¹⁴ Chemicals and solvents were obtained from commercial sources, and used without further purification unless otherwise detailed. Reactions were monitored by TLC performed on Macherey-Nagel Alugram® Sil 60/UV₂₅₄ sheets (thickness 0.2 mm). Purification of products was carried out by either column chromatography or thick layer chromatography. Column chromatography was carried out on using Macherey-Nagel silica gel (230-400 mesh). Thick layer chromatography was performed on glass plates coated with Macherey-Nagel Sil/UV₂₅₄ (thickness 2 mm), from which the pure compounds were extracted with the following solvent system: DCM/MeOH (NH₃), 90:10. ¹H and ¹³C NMR spectra were obtained using a Bruker 300 MHz spectrometer, chemical shifts (δ) were expressed in ppm relative to TMS used as an internal standard. The attributions of the carbons were deduced after 2D experiments had been performed (COSY, HSQC and HMBC). Mass spectra were recorded on a Varian triple quadrupole 1200 W mass spectrometer equipped with a non-polar C18 TSK-gel Super ODS (4.6 x 50 mm) column, using electrospray ionization and a UV detector (diode array). The purity of final compounds was verified by two types of high pressure liquid chromatography (HPLC) columns: C18 Deltapak (C18N) and C4 Interchrom UP5WC4-25QS (C4). Analytical HPLC was performed on a Shimadzu system equipped with a UV detector set at 254 nm. Compounds were dissolved in buffer B or

MeOH and injected through a 50 μ L loop. The following eluent systems were used: buffer A (H₂O/TFA, 100:0.05) and buffer B (CH₃CN/H₂O/TFA, 80:20:0.05). HPLC retention times (HPLC t_R) were obtained, at flow rates of 1 mL/min, using the following conditions: for the 10 min method: a gradient run from 100% eluent A for 30 s, then to 100% eluent B for the next 8 min; and for the 40 min method: a gradient run from 100% eluent A for 1 min, then to 100% eluent B for the next 30 min.

(3-{4-[3-(4-Fluorobenzylamino)propyl]piperazin-1-yl}propyl)diisobutylamine (35)

A 3Å molecular sieve (1 g) was added to a solution of {3-[4-(3-aminopropyl)piperazin-1-yl]propyl}diisobutylamine¹² (150 mg, 0.48 mmol) and 4-fluorobenzaldehyde (57 μ L, 0.53 mmol) in ethanol (6 mL). The reaction medium was stirred at 20°C for 5 h and cooled to 0°C. NaBH₄ (45.4 mg, 1.2 mmol) was added portionwise over 15 min, and the reaction medium stirred at 20°C for 12 h. The mixture was filtered over Celite. The solvent was removed under reduced pressure and 20 mL water added to the residue. The product was extracted with dichloromethane, and washed with aqueous HCl 1M. The combined organic fractions were washed with water and dried over sodium sulfate. The solvent was evaporated under reduced pressure. Purification by thick layer chromatography (DCM:MeOH:NH₄OH, 90:10:1) was performed and enabled collection of the product as a yellow solid (102 mg, 50%). ¹H NMR (300 MHz, CDCl₃) δ 7.20 (dd, ³*J* = 8.7 Hz, ⁴*J*_F = 5.5 Hz, 2H, H_{aro}), 6.93 (m, ³*J* = ³*J*_F = 8.7 Hz, 2H, H_{aro}), 3.67 (s, 2H, NHCH₂), 2.59 (t, ³*J* = 6.8 Hz, 2H, NH-CH₂), 2.5-2.2 (m, 14H, N-CH₂), 1.96 (d, ³*J* = 7.2 Hz, 4H, 2 N-CH₂), 1.68 (m, 2H, 2 CH), 1.5-1.4 (m, 4H, 2 CH₂), 0.78 (d, ³*J* = 6.6 Hz, 12H, 4 CH₃). ¹³C NMR (75 MHz, CD₃OD) δ 129.8 (2 CH_{aro}), 115.1 (2 CH_{aro}), 64.1 (2 NCH₂), 57.2 (NCH₂), 57.0 (2 NCH₂), 53.5 (NHCH₂), 53.4 (2 NCH₂), 48.3 (NHCH₂), 26.7 (2 CH), 24.6 (2 CH₂), 21.1 (4 CH₃). LCMS (ESI⁺): Calc. for

[M+H]⁺: 421.36; Found: 421.47. HPLC (C₄, 35 min): t_R 6.7 min, P_{HPLC} 99%; HPLC (C₁₈, 35 min): t_R 3.8 min, P_{HPLC} > 99%.

Diisobutyl-[3-(4-{3-[(thiazol-2-ylmethyl)amino]propyl}piperazin-1-yl)propyl]amine (36)

A 3 Å molecular sieve (1 g) was added to a solution of {3-[4-(3-aminopropyl)piperazin-1-yl]propyl}diisobutylamine¹² (150 mg, 0.48 mmol) and 2-thiazolecarboxaldehyde (46.4 µL, 0.53 mmol) in ethanol (6 mL). The reaction medium was stirred at 20°C for 5 h and cooled to 0°C. NaBH₄ (45.4 mg, 1.2 mmol) was added portionwise over 15 min and the reaction medium was stirred at 20°C for 12 h. The mixture was filtered over Celite. The solvent was removed under reduced pressure and 20 mL water added to the residue. The product was extracted with dichloromethane, and washed with aqueous HCl 1M. The combined organic fractions were washed with water and dried over sodium sulfate. The solvent was evaporated under reduced pressure. Purification by thick layer chromatography (AcOEt:MeOH:NH₄OH, 90:10:2) was performed and enabled collection of the product as a yellow oil (69 mg, 35%). ¹H NMR (300 MHz, CDCl₃) δ 7.66 (d, ³J = 3.4 Hz, 1H, H_{aro}), 7.20 (d, ³J = 3.4 Hz, 1H, H_{aro}), 4.07 (s, 2H, NHCH₂), 2.70 (t, ³J = 6.7 Hz, 2H, NH-CH₂), 2.5-2.3 (m, 14H, N-CH₂), 1.97 (d, ³J = 7.2 Hz, 4H, 2 N-CH₂), 1.7-1.5 (m, 6H, 2 CH, 2 CH₂), 0.79 (d, ³J = 6.5 Hz, 12H, 4 CH₃). ¹³C NMR (75 MHz, CD₃OD) δ 142.6 (CH_{aro}), 118.9 (CH_{aro}), 64.1 (2 NCH₂), 57.0 (2 NCH₂), 53.4 (3 NHCH₂), 51.2 (NHCH₂), 48.5 (NHCH₂), 27.1 (CH₂), 26.7 (2 CH), 24.6 (CH₂), 21.1 (4 CH₃). LCMS (ESI⁺): Calc. for [M+H]⁺: 410.32; Found: 410.40. HPLC (C₄, 40 min): t_R 3.8 min, P_{HPLC} 99%; HPLC (C₁₈, 35 min): t_R 3.8 min, P_{HPLC} 99%.

(4-Fluorobenzyl)-(3-{4-[3-(4-fluorobenzylamino)propyl]piperazin-1-yl}propyl)amine (41)

A 3 Å molecular sieve (5 g) was added to a solution of 1,4-bis(3-aminopropyl)piperazine (515 µL, 2.50 mmol) and 4-fluorobenzaldehyde (590 µL, 5.50 mmol) in ethanol (20 mL). The reaction medium was stirred at 20°C for 5 h and cooled to 0°C. NaBH₄ (0.473 g, 12.50 mmol) was added portionwise over 15 min and the reaction medium was stirred at 20°C for 12 h. The mixture was filtered over Celite. The solvent was removed under reduced pressure and 20 mL water added to the residue. The product was extracted with dichloromethane, washed with aqueous HCl 1M. The combined organic fractions were washed with water and dried over sodium sulfate. The solvent was evaporated under reduced pressure. Purification by thick layer chromatography (acetone:NH₄OH, 90:10) was performed and enabled collection of the product as a yellow solid (0.18 g, 17%). ¹H NMR (300 MHz, CDCl₃) δ 7.21 (dd, ³J = 8.7 Hz, ⁴J = 5.5 Hz, 4H, H_{aro}), 6.93 (m, 4H, H_{aro}), 3.68 (s, 4H, NHCH₂), 2.60 (t, ³J = 6.8 Hz, 4H, NH-CH₂), 2.4-2.3 (m, 12H, N-CH₂), 1.63 (quint, ³J = 7.0 Hz, 4H, CH₂). ¹³C NMR (75 MHz, CD₃OD) δ 129.8 (CH_{aro}), 115.5 (CH_{aro}), 57.2 (NCH₂), 53.4 (2 NCH₂), 53.3 (NHCH₂), 48.3 (NHCH₂), 26.8 (CH₂). LCMS (ESI⁺): Calc. for [M+H]⁺: 417.27; Found: 417.04. HPLC (C₄, 40 min): t_R 5.9 min, P_{HPLC} > 99%; HPLC (C₁₈, 35 min): t_R 3.2 min, P_{HPLC} 95%.

Cell culture and treatment

The human neuroblastoma cell line SKNSH-SY5Y (SY5Y) was cultured in Dulbecco's modified Eagle medium supplemented with 10% fetal calf serum (PAA), 2 mM L-glutamine (Invitrogen), 1 mM non-essential amino-acids and penicillin/streptomycin (Invitrogen), in a 5% CO₂ humidified incubator at 37°C. The human APP695 cDNA was subcloned into eukaryotic expression vector pcDNA3.1 (Invitrogen), allowing for G418 antibiotic selection of stable clones. This APP cDNA was transfected into SY5Y cells using the ethyleneimine polymer ExGen 500 (Euromedex)

according to the manufacturer's instructions. SY5Y cells stably expressing the APP695 were selected with Geneticin G418 (Invitrogen) and one clone named SY5Y-APP^{wt} was used here.

For treatment, SY5Y-APP^{wt} cells were plated onto 12-well plates (Falcon) 24 h before drug exposure, and cultured in Dulbecco's modified Eagle medium (Invitrogen) supplemented with 10% fetal calf serum (PAA), 2 mM L-glutamine (Invitrogen), 1 mM non-essential amino acids (Invitrogen), 50 units/mL penicillin/streptomycin (Invitrogen), and 200 µg Geneticin G418 (Invitrogen), under 5% CO₂ at 37°C. Cells were exposed to drugs at the indicated concentrations for 24 h. After treatment, the conditioned medium was collected, spun at 200xg to eliminate the cell debris and frozen at -80°C for Aβ₁₋₄₂, Aβ₁₋₄₀ and sAPPα quantification. Treated SY5Y-APP^{wt} cells were collected in 50 µl of Laemmli lysis buffer containing protease inhibitors (Complete Mini, Roche Molecular Biochemicals, Meylan, France), sonicated for 5 min and stored at -80°C until use. Total protein quantification of extracted samples was performed by BCATM Protein Assay Kit (Thermo Scientific) according to the manufacturer's protocol.

Cytotoxicity

SY5Y-APP^{wt} cells cultured as described previously were seeded onto 96-well plates and incubated with the compound at 0.1; 0.3; 1; 3; 10; 30 and 100 µM, or DMSO, diluted in the same culture medium as a control, at 37°C in 5% CO₂ for 24h (n=3). Cytotoxicity was determined by using colorimetric MTS assay (CellTiter 96® Aqueous One Solution Cell Proliferation Assay-MTS Promega) according to the manufacturer's protocol. Absorbance was read at 490 nm and cytotoxicity was interpreted as the compound concentration causing 50% of cell death (CC₅₀). Results are shown as a percentage of control conditions which was considered as 100%.

Western blot analysis

Samples were heated at 85°C for 2 min with Reducing Agent (Life Technologies™) and equal quantities of total proteins (20 µg/lane) were resolved in NuPAGE® Novex® 16% Tris-Tricine precast gels (Life Technologies™). After electrophoresis, the proteins were transferred onto 0.2 µM PVDF membranes (Life Technologies™) for 1 h at 20°C using the liquid transfer system (Life Technologies™). Membranes were blocked with 5% skimmed milk in TNT (15 mM Tris buffer pH 8.4, 140 mM NaCl, 0.05% Tween-20) for 1 h at 20°C. After washing three times, the membrane were incubated with APPCter-C17 rabbit antiserum diluted 1:4,000 in TNT overnight at 4°C. APP-Cter-C17 was raised against the last 17 amino acids of the human APP sequence.⁴⁵ To develop the immunoreaction, the blots were incubated with peroxidase-conjugated purified mouse monoclonal anti-goat/sheep IgG (Sigma A 9452, MAb clone GT-34), 1:10,000 in TNT-M, for 1 h at 20°C, and developed with SuperSignal West Pico Chemiluminescent Substrate (Thermo Scientific). Membranes were scanned with LAS-4000 Mini Image System. AICD fragment (8 kDa) and CTF α (12 kDa) were detected. Images were obtained with a time exposure from 10 to 320s. Each image was opened with Adobe Photo Shop CS2 (version 9.0.2) computer program, a compose containing all WB bands was created for analysis. Bands quantification was performed by using Image J 1.37v computer program. Each band was transformed in a plot and the area under the curve was calculated. Results were expressed as arbitrary units of optical density. Membranes were then rinsed for 30 min at 20°C and reprobed with a goat polyclonal antibody against α -Actin, GAPDH or Neuron Specific Enolase (1:1000; Santa Cruz Biotechnology). *In vitro* results are shown as the concentrations of compound able to double (C₂) or multiply 10-fold (C₁₀) the quantity of AICD or CTF α quantified by Image J.

Secreted A β ₁₋₃₈, A β ₁₋₄₀, A β ₁₋₄₂ and sAPP α quantification

Conditioned medium was used to determine the secreted A β ₁₋₃₈, A β ₁₋₄₀ and A β ₁₋₄₂ concentrations, using the Human A β (1–40) and A β (1-38) Assay Kits (IBL) and the INNOTEST™ beta-Amyloid (1-42) ELISA Kit (Innogenetics). For sAPP α concentrations, the human sAPP α (highly sensitive) Assay Kit (IBL) was used according to the manufacturer's instructions. Each sample was loaded in duplicate onto a 96 well plate. Experiments were done in triplicate. Results expressed in ng/ml were compared to control conditions arbitrarily given an average value of 100%. Results are presented as IC₅₀, the concentration able to decrease to 50% the basal quantity of secreted A β peptide 1-38, 1-40 and 1-42.

Bioavailability properties

Bioavailability-related profile was measured according to Lipinski *et al.*⁴⁶ for aqueous solubility (PBS pH 7.4), and according to Sangster⁴⁷ for partition coefficient (logD, n-octanol-PBS, pH 7.4) by CEREP (Paris, France).

***In vitro* metabolic stability (mouse and human liver microsomes)**

Stock solutions of compounds were diluted in 100 mM potassium phosphate buffer (KPi, 1 μ M final concentrations) pH = 7.4, and test compounds were then incubated for 60 min in an incubator shaker (Eppendorf) at 37°C and 1,400 rpm with regenerating system (NADPH) and microsomal preparation (BD, final concentration 0.3 mg/mL in KPi buffer). Reactions were stopped with cold acetonitrile and Internal Standard (IS) CQ diphosphate was then added for further quantification

(based upon Test compound/IS ratio area). Samples were mixed thoroughly and then centrifuged at 13,000 rpm for 10 min. Supernatants were evaporated under vacuum (SpeedVac) at medium drying rate for 2 h and residues reconstituted in water + 0.1 % TFA. Finally, 10 μ L volumes were injected into the LCMS system. Microsomal stability was calculated by comparison of area ratio of parent compounds at different times.

The LCMS system for the microsomal stability assay consisted of an Orbitrap Exactive instrument (Thermo) equipped with an electrospray ionization source used in positive mode ($M+H^+$). The apparatus was managed with Xcalibur software. Tune parameters were set as: sheath gas flow rate at 70 L/min, aux gas flow rate at 20 L/min, spray voltage at 3.00 kV, capillary temperature at 275°C, capillary voltage at 95 V, tube lens voltage at 165 V and skimmer voltage at 36 V. Tray temperature was set at 4°C and oven temperature at 30°C. The analytical column was a C18 Hypersil Gold Thermo 50 x 2 mm, 1.9 μ m (Thermo). The mobile phase consisted of water + 0.05% TFA (A) and acetonitrile + 0.05% TFA (B). The linear gradient elution program was as follows: 0-100% of B for 3.5 min, followed by an isocratic hold at 100% B for 1 min and 2 min of reequilibration with 100% A for a total run of 6 min at a flow rate of 400 μ L/min. Due to the basicity of polyamines, good sensitivity was achieved using TFA in the mobile phase.

***In vivo* acute treatment**

Female 4-month-old C57Bl6 mice were treated p.o. (gavage) with either carrier (water) or compound **29** as hydrochloride at 1; 3; 6; 12.5 and 25 mg/kg (minimum number of animals per group = 6). After 24 h, mice were sacrificed and the brain immediately removed to dissect the frontal cortex and hippocampus. Tissues were stored at -80°C until western blot analysis. All experiments were carried out in accordance with the European Communities Council Directive of

November 24th, 1986 (86/609/EEC) and the experimental protocols were approved by the local institutional animal research committee (Project number: CEEA-35014, agreement number 59-350208).

Statistics

Statistical analysis was performed with Graph Pad Prism® software computer program. Analysis was performed using one-way ANOVA (F value), followed by a Fisher test. $p < 0.05$, 0.01 and 0.001 were considered statistically significant (respectively indicated *, ** and ***).

SUPPORTING INFORMATION

NMR spectra of compounds **35**, **36** and **41** are available free of charge via the Internet at <http://pubs.acs.org>.

ABBREVIATIONS

aa, amino acid; AD, Alzheimer's Disease; ADME, Absorption Distribution Metabolism Excretion; AICD, APP IntraCellular Domain; APP, Amyloid Precursor Protein; BBB, Blood-Brain Barrier; CTF, Carboxy-Terminal Fragment; CQ, Chloroquine; HCV, Hepatitis C Virus; LAR, Lysosome Accumulation Ratio; MTS, 3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenol)-2-(4-sulfo phenyl)-2H-tetrazolium salt.

AUTHOR INFORMATION

Corresponding Author

* Patricia Melnyk, EA4481, UFR Pharmacie, 3 rue du Pr Laguesse, BP83, 59006 Lille. Tel: 33 (0)3 20 96 49 49 – Fax: 33 (0)3 20 96 49 13, e-mail: patricia.melnik@univ-lille2.fr

Author Contributions

Patricia Melnyk: Responsible for medicinal chemistry team, design of the study, and writing the manuscript.

Valérie Vingt-deux: Contributed to the development of the cell assay and obtained the result.

Stéphane Burlet: Choice of compounds, involved in scientific discussion.

Sabiha Eddarkaoui: Contributed to the design of the assay and obtained the result.

Marie-Eve Grosjean: Contributed to the results.

Paul-Emmanuel Larchanché: Synthesis of new compounds.

Guillaume Hochart: Metabolic stability studies.

Christian Sergheraert: Design of the chemical library, involved in scientific discussion.

Cecilia Estrella: *In vivo* and *in vitro* study, analysis, involved in scientific discussion.

Matthieu Barrier: *In vivo* study, analysis, involved in scientific discussion.

Virginie Poix: *In vivo* study and CTF α WB.

Pauline Plancq: A β ₁₋₄₂ and sAPP α ELISA.

Cécile Lannoo: CC₅₀ determination, AICD and CTF α WB.

Malika Hamdane: developed and characterized the cell model.

André Delacourte: contributed to the design of the study, involved in scientific discussions.

Philippe Verwaerde: contributed to the design of the study, involved in scientific discussions

Luc Buée: contributed to the design of the study.

Nicolas Sergeant: responsible for the design of the study, wrote the manuscript.

All authors have approved the final version of the manuscript.

Funding Sources

This work was supported by the University of Lille 2, CNRS and Inserm. Grants were obtained from FEDER, OSEO and ANR.

ACKNOWLEDGMENT

We express our thanks to Emmanuelle Boll and Alexandre Barras for their contributions to organic synthesis and analysis, and to Nathalie Duhal and Dr Mostafa Kouach, CUMA, for help with the LCMS experiment.

The 300 MHz NMR facilities were funded by the Région Nord-Pas de Calais (France), the Ministère de la Jeunesse, de l'Education Nationale et de la Recherche (MJENR) and the Fonds Européens de Développement Régional (FEDER).

REFERENCES

1. Vodicka, P., Lim, J., Williams, D.T., Kegel, K.B., Chase, K., Park, H., Marchionini, D., Wilkinson, S., Mead, T., Birch, H., Yates, D., Lyons, K., Dominguez, C., Beconi, M., Yue, Z., Aronin, N., DiFiglia, M. (2014) Assessment of chloroquine treatment for modulating autophagy flux in brain of WT and HD mice. *J. Huntingtons Dis.* 3, 159-74.
2. Korth, C., May, B. C. H., Cohen, F. E., Prusiner, S. B. (2001) Acridine and phenothiazine derivatives as pharmacotherapeutics for prion disease. *Proc. Natl. Acad. Sci. USA* 98, 9836-9841.
3. Doh-ura, K., Iwaki, T., Caughey, B. (2000) Lysosomotropic agents and cysteine protease inhibitors inhibit scrapie-associated prion protein accumulation. *J. Virol.* 74, 4894-4897.

4. Kocisko, D. A., Baron, G. S., Rubenstein, R., Chen, J., Kuizon, S. *et al.* (2003) New inhibitors of scrapie-associated prion protein formation in a library of 2000 drugs and natural products. *J. Virol.* 77, 10288-10294.
5. Klingenstein, R., Melnyk, P., Leliveld, R. S., Ryckebusch, A., Korth, C. (2006) Similar Structure Activity Relationships of Quinoline Derivatives for antiprion and antimalarial effects. *J. Med. Chem.* 49, 5300-5308.
6. Ashfaq, U.A., Javed, T., Rehman, S., Nawaz, Z., Riazuddin, S. (2011) Lysosomotropic agents as HCV entry inhibitors. *Virol. J.* 8, 163-168.
7. Blanchard, E., Belouzard, S., Goueslain, L., Wakita, T., Dubuisson, J., Wychowski, C., Rouillé, Y. (2006) Hepatitis C virus entry depends on clathrin-mediated endocytosis. *J. Virol.* 80, 6964-72.
8. Vasquez-Martin, A., Lopez-Bonetc, E., Cuti, S., Oliveras-Ferraros, C., Del Barco, S., Martin-Castillo, B., Menendez, J.A. (2011) Repositioning chloroquine and metformin to eliminate cancer stem cell traits in pre-malignant lesions. *Drug Resist. Update* 14, 212-223.
9. Mahoney, E., Maddocks, K., Flynn, J., Jones, J., Cole, S.L., Zhang, X., Byrd, J.C., Johnson, A.J. (2013) Identification of endoplasmic reticulum stress-inducing agents by antagonizing autophagy : a new potential strategy for identification of anti-cancer therapeutics in b-cell malignancies. *Leuk. Lymphoma* 54, 2685-92.
10. Costedoat-Chalumeau, N., Leroux, G., Piette, J.-P., Amoura, Z. (2010) Why all systemic lupus erythematosus patients should be given hydroxychloroquine treatment? *Joint Bone Spine* 77, 4-5.

11. Suarez-Almazor, M.E., Belseck, E., Shea, B., Homik, J., Wells, G., Tugwell, P. (2010) Antimalarials for treating rheumatoid arthritis. *Cochrane Database Syst. Rev.* 4, CD000959.
12. Ryckebusch, A., Deprez-Poulain, R., Maes, L., Debreu-Fontaine, M.-A., Mouray, E., Grellier, P., Sergheraert, C. (2003) Synthesis and in Vitro and in Vivo Antimalarial Activity of N1-(7-Chloro-4-quinolyl)-1,4-bis(3-aminopropyl)piperazine Derivatives. *J. Med. Chem.* 46, 542-557.
13. Ryckebusch, A., Deprez-Poulain, R., Debreu-Fontaine, M.-A., Vandaele, R., Mouray, E., Grellier, P., Sergheraert, C. (2003) Synthesis and Antimalarial Evaluation of New 1,4-bis(3-aminopropyl)piperazine Derivatives. *Bioorg. Med. Chem. Lett.* 13, 3783–3787.
14. Bonnet, B., Soullez, D., Girault, S., Maes, L., Landry, V., Davioud-Charvet, E., Sergheraert, C. (2000) Trypanothione Reductase Inhibition/Trypanocidal Activity Relationships in a 1,4-Bis(3-aminopropyl)piperazine Series. *Bioorg. Med. Chem.* 8, 95-103.
15. Yin, R.H., Tan, L., Jiang, T., Yu, J.T. (2014) Prion-like Mechanisms in Alzheimer's Disease. *Curr. Alzheimer Res.* 11, 755-64.
16. Vingtdeux, V., Sergeant, N., Buee, L. (2012) Potential contribution of exosomes to the prion-like propagation of lesions in Alzheimer's disease. *Front. Physiol.* 3, 229-250.
17. Vingtdeux, V., Hamdane, M., Loyens, A., Gelé, P., Drobecq, H., Bégard, S., Galas, M.-C., Delacourte, A., Beauvillain, J.-C., Buée, L., Sergeant, N. (2007) Alkalizing drugs induce accumulation of amyloid precursor protein by-products in luminal vesicles of multivesicular bodies. *J. Biol. Chem.* 282, 18197-18205.

18. Kummer, M.P., Heneka, M.T. (2014) Truncated and modified amyloid-beta species. *Alzheimers Res. Ther.* 6, 28-36.
19. Yuyama, K., Yamamoto, N., Yanagisawa, K. (2006) Chloroquine-induced endocytic pathway abnormalities: Cellular model of GM1 ganglioside-induced Abeta fibrillogenesis in Alzheimer's disease. *FEBS Lett.* 580, 6972-6.
20. Bendiske, J., Bahr, B.A. (2003) Lysosomal activation is a compensatory response against protein accumulation and associated synaptopathogenesis--an approach for slowing Alzheimer disease? *J. Neuropathol. Exp. Neurol.* 62, 451-63.
21. De Strooper, B. (2010) Proteases and proteolysis in Alzheimer disease: a multifactorial view on the disease process. *Physiol. Rev.* 90, 465-494
22. Vingtdeux, V., Marambaud, P. (2012) Identification and biology of α -secretase. *J. Neurochem.* 120 (Suppl 1), 34-45.
23. Pardossi-Piquard, R., Checler, F. (2012) The physiology of the β -amyloid precursor protein intracellular domain AICD. *J. Neurochem.* 120 (Suppl 1), 109-24.
24. Takami, M., Nagashima, Y., Sano, Y., Ishihara, S., Morishima-Kawashima, M., Funamoto, S., Ihara, Y. (2009) Gamma-Secretase: successive tripeptide and tetrapeptide release from the transmembrane domain of beta-carboxyl terminal fragment. *J. Neurosci.* 29, 13042-52.
25. Karran, E., Mercken, M., de Strooper, B.D. (2011) The amyloid cascade hypothesis for Alzheimer's Disease: an appraisal for the development of therapeutics. *Nat. Rev. Drug Discov.* 10, 698-712.

26. Matsumura, N., Takami, M., Okochi, M., Wada-Kakuda, S., Fujiwara, H., Tagami, S., Funamoto, S., Ihara, Y. Morishima-Kawashima, M. (2014) Gamma-Secretase Associated with Lipid Rafts: Multiple Interactive Pathways in The Stepwise Processing of Beta-Carboxyl-terminal Fragment. *J. Biol. Chem.* 289, 5109–21.
27. Yan, R., Vassar, R. (2014) Targeting the beta secretase BACE1 for Alzheimer's disease therapy. *Lancet Neurol.* 13, 319–29.
28. Hall, A., Patel, T.R. (2014) Gamma-Secretase modulators: current status and future directions. *Prog. Med. Chem.* 53, 101–45.
29. Strömberg, K., Eketjäll, S., Georgievska, B., Tunblad, K., Eliason, K., Olsson, F., Radesäter, A.C., Klintonberg, R., Arvidsson, P.I., von Berg, S., Fälting, J., Cowburn, R.F., Dabrowski, M. (2015) Combining an amyloid-beta (A β) cleaving enzyme inhibitor with a γ -secretase modulator results in an additive reduction of A β production. *FEBS J.* 282, 65-73.
30. Vingtdeux, V., Hamdane, M., Bégard, S., Loyens, A., Delacourte, A., Beauvillain, J.-C., Buée, L., Marambaud, P., Sergeant, N. (2007) Intracellular pH regulates amyloid precursor protein intracellular domain accumulation. *Neurobiol. Dis.* 25, 686-696.
31. Tam, J.H., Seah, C., Pasternak, S.H. (2014) The Amyloid Precursor Protein is rapidly transported from the Golgi apparatus to the lysosome and where it is processed into beta-amyloid. *Mol. Brain.* 7, 54-71.
32. Melnyk, P., Sergeant, N., Buée, L., Delacourte, A. Use of 1,4-bis(3-aminoalkyl) piperazine derivatives in the treatment of neurodegenerative diseases. WO 2006/ 051489.

33. Vingtdeux, V., Hamdane, M., Gompel, M., Bégard, S., Drobecq, H., Ghestem, A., Grosjean, M.-E., Kostanjevecki, V., Grognet, P., Vanmechelen, E., Buée, L., Delacourte, A., Sergeant, N. (2005) Phosphorylation of amyloid precursor carboxy-terminal fragments enhances their processing by a gamma-secretase-dependent mechanism. *Neurobiol. Dis.* 20, 625–637.

34. Lysosome Accumulation Ratios (LAR) was calculated *in silico* using the following equation:

$$\text{LAR} = \frac{1 + \sum_{n=1}^4 \sum_{i=1}^n 10^{\text{pK}_{\text{ai}} - \text{pH}_{\text{L}}}}{1 + \sum_{n=1}^4 \sum_{i=1}^n 10^{\text{pK}_{\text{ai}} - \text{pH}_{\text{O}}}}$$

Where: pH_L is pH inside the lysosome (assumed to be pH 5.0) and pH_O in the cytosol (assumed to be pH 7.4).

This equation is a derivation of the Henderson-Hasselbach equation, based on predicted values of drug pK_a according to previous work of Hawley *et al.*³⁵ Values of pK_a were calculated using ACD/pK_a DB software from Advanced Chemistry Development Inc., Toronto, Canada.

35. Hawley, S. R., Bray, P. G., O'Neill, P. M., Park, B. K., Ward, S. A. (1996) The role of drug accumulation in 4-aminoquinoline antimalarial potency. The influence of structural substitution and physicochemical properties. *Biochem. Pharmacol.* 52, 723-733.

36. Chasseigneaux, S., Allinquant, B. (2012) Functions of Aβ, sAPPα and sAPPβ: similarities and differences. *J. Neurochem.* 120, 99-108.

37. Ryckebusch, A., Deprez-Poulain, R., Debreu-Fontaine, M.-A., Vandaele, R., Mouray, E., Grellier, P., Sergheraert, C. (2002) Parallel Synthesis and Antimalarial Activity of a Sulfonamide Library. *Bioorg. Med. Chem. Lett.* 12, 2595-2598.

38. Belyaev, N.D., Kellett, K.A., Beckett, C., Makova, N.Z., Revett, T.J., Nalivaeva, N.N., Hooper, N.M., Turner, A.J. (2010) The Transcriptionally Active Amyloid Precursor Protein (APP) Intracellular Domain Is Preferentially Produced from the 695 Isoform of APP in a beta-Secretase-dependent Pathway. *J. Biol. Chem.* 285, 41443–41454.
39. Asai, M., Yagishita, S., Iwata, N., Saido, T.C., Ishiura, S., Maruyama, K. (2011) An alternative metabolic pathway of amyloid precursor protein C-terminal fragments via cathepsin B in a human neuroglioma model. *FASEB J.* 25, 3720–3730.
40. Vorobyeva, A.G., Lee, R., Miller, S., Longen, C., Sharoni, M., Kandelwal, P.J., Kim, F.J., Marena, D.R., Saunders, A.J. (2014) Cyclopamine Modulates γ -Secretase-mediated Cleavage of Amyloid Precursor Protein by Altering Its Subcellular Trafficking and Lysosomal Degradation. *J. Biol. Chem.* 289, 33258–33274.
41. Goodger, Z. V., Rajendran, L., Trutzel, A. (2009) Nuclear signaling by the APP intracellular domain occurs predominantly through the amyloidogenic processing pathway. *J. Cell Sci.* 122, 3703-14.
42. Ghosal, K., Vogt, D.L., Liang, M., Shen, Y., Lamb, B.T., Pimplikar, S.W. (2009) Alzheimer's disease-like pathological features in transgenic mice expressing the APP intracellular domain. *Proc. Natl. Acad. USA* 106 (43), 18367-72.
43. Ghosal, K., Pimplikar, S.W. (2011) Aging and exitotoxic stress exacerbate neural circuit reorganization in amyloid precursor protein intracellular domain transgenic mice. *Neurobiol. Aging* 32 (12), 2320, e1-9.

44. Ohkawara, T., Nagase, H., Koh, C.-S., Nakayama, K. (2011) The amyloid precursor protein intracellular domain alters gene expression and induces neuron-specific apoptosis. *Gene* 475, 1–9.
45. Sergeant, N., David, J.P., Champain, D., Ghestem, A., Wattez, A., Delacourte, A. (2002) Progressive decrease of amyloid precursor protein carboxy terminal fragments (APP-CTFs), associated with tau pathology stages, in Alzheimer's disease. *J. Neurochem.* 81 (4), 663-672.
46. Lipinski, C.A., Lombardo, F., Dominy, B.W., Feeney, P.J. (2001) Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings. *Adv. Drug Deliv. Rev.* 46, 3-26.
47. Sangster, J. (1997). Octanol-Water Partition Coefficients: Fundamentals and Physical Chemistry. *Wiley Series in Solution Chemistry*, Vol. 2, John Wiley and Sons, Chichester.

Figure 1. Amyloid Precursor Protein (APP) metabolism (adapted from Vingtdoux *et al.* ¹⁶)

Figure 2. Chloroquine (CQ) and piperazine-derived compounds (Series I, II and III)

Figure 3: Effect of CQ-derivatives on APP metabolism.

SY5Y-APP^{wt} were treated or not (Control condition : Ctrl) with 1 or 5 μM of compound **33** (the structure is represented) and compared to the effect of chloroquine (CQ) at the same doses. After 6 or 24 hours of treatment, protein cell lysates were separated on 1D tris-tricine gels and APP metabolites, including APP-CTFs and AICD were detected with the APP-Cter-C17 antisera against the last 17 aa of the APP protein sequence. Apparent molecular weights are indicated on the right and were determined using the Novex® Sharp Pre-stained Protein Standard (3.5 to 260 kDa). The 5 min exposure is used to quantify the amount of APP-CTFs whereas the 30 min exposure is used to visualize and quantify AICD. APP-CTFs released from the β- (β-CTF and β'-CTF) or α- cleavage (α-CTF) are indicated on the left. Membranes were incubated with an anti neuron specific enolase (NSE) and used as loading control. Western-blot bands were quantified and results are reported on tables.

A: Frontal Cortex

B: Hippocampus

Figure 4. *In vivo* evaluation of compound 29.

C57Bl6 females were treated with compound 29 (as hydrochloride) for 24h. Semiquantitative levels of CTF α in frontal cortex (A) and hippocampus (B) after western blot; One-way ANOVA and Fisher test * $p < 0.05$; ** $p < 0.01$ and *** $p < 0.001$, compared to control (Ctl). The number of animals per group is indicated in parentheses. Results show the mean $\pm$ SEM.

Scheme 1. Synthesis of benzyl derivatives **35**, **36** and **41**.

Table 1. *In vitro* evaluation of compounds from Series I on SY5Y-APP^{WT} cells

	Ref	SY5Y CC ₅₀ (μM) ^a	AICD C ₂ (μM) ^b	AICD C ₁₀ (μM) ^c	Aβ ₁₋₄₂ IC ₅₀ (μM) ^d	ref	SY5Y CC ₅₀ (μM) ^a	AICD C ₂ (μM) ^b	AICD C ₁₀ (μM) ^c	Aβ ₁₋₄₂ IC ₅₀ (μM) ^d	Ref	SY5Y CC ₅₀ (μM) ^a	AICD C ₂ (μM) ^b	AICD C ₁₀ (μM) ^c	Aβ ₁₋₄₂ IC ₅₀ (μM) ^d	
CQ		30	1.4	10.8	12.7											
Quinacrine		5	1.0	>5	>5											
R		Series Ia					Series Ib					Series Ic
phenyl	1	57	2.0	16.9	10.3	9	<5				17	1.0				
4-methoxyphenyl	2	64	1.3	36.0	10.8	10	1.0				18	1.0				
4-chlorophenyl	3	17	2.1	13.9	9.6	11	1.0				19	1.0				
4-nitrophenyl	4	17	1.3	>17	7.3	12	17	1.0	>17	7.1	20	3.0				
methyl	5	>100	16.8	>100	>20	13	48	1.4	19.3	>10	21	18	0.8	5.8	11.3	
<i>t</i> -butyl	6	33	2.1	9.0	11.3	14	27	0.8	10.1	8.7	22	65	0.9	>60	18.4	
cyclopropyl	7	75	5.1	39.5	>20	15	60	1.0	6.7	16.4	23	6.5	0.6	3.0	>6	
<i>i</i> -propyl	8	> 30	10.1	21.6	15.0	16	10	3.7	>10	>10	24	5.3	0.4	2.7	1.0	
pyrrolidinyl											25	60	1.2	5.4	1.0	
piperidinyl											26	8.5	0.9	6.3	4.0	

Mean values calculated on the basis of at least three independent experiments with less than 10% deviation; nd: not determined; ^a compound concentration causing 50% of cell death after 24 h treatment; ^b Concentration doubling the quantity of AICD; ^c Concentration increasing the quantity of AICDs generated 10-fold; ^d Concentration decreasing the secretion of A β by 50%.

Table 2. *In vitro* evaluation of compounds of Series II on SY5Y cells

	Ref	SY5Y	AICD	AICD	Aβ ₁₋₄₂		Ref	SY5Y	AICD	AICD	Aβ ₁₋₄₂
		CC ₅₀ (μM) ^a	C ₂ (μM) ^b	C ₁₀ (μM) ^c	IC ₅₀ (μM) ^d			CC ₅₀ (μM) ^a	C ₂ (μM) ^b	C ₁₀ (μM) ^c	IC ₅₀ (μM) ^d
CQ		30	1.4	10.8	12.7						
R		Series IIa				R		Series IIb
7-chloro-4-quinolinyl	24	5.3	0.4	2.7	1.0	4-methoxyphenyl	33	82	1.5	10.5	1.0
6-chloro-2-methoxyacridin-9-yl	27	<5	>5	nd	>5	4-chlorophenyl	34	30	1.3	16	2.4
2-benzoxazolyl	28	>100	^e	nd	>10	4-fluorophenyl	35	>30	2	9.5	^f
2-benzimidazolyl	29	30	0.8	3.0	1.5	2-thiazolyl	36	nd	^e	^e	>10
2-pyrimidinyl	30	30	>10	nd	>10	4-pyridinyl	37	>100	3	5	>10
2-pyrazinyl	31	30	4.1	nd	>10						
6-purinyl	32	>60	>10	nd	>10						

Mean values calculated on the basis of at least three independent experiments with less than 10% deviation; nd: not determined; ^a compound concentration causing 50% of cell death after 24 h treatment; ^b Concentration doubling the quantity of AICD; ^c Concentration increasing the quantity of AICDs generated 10-fold; ^d Concentration decreasing the secretion of Aβ by 50%; ^e decrease in the quantity of AICDs; ^f increase in the quantity of Aβ at 5 μM.

Table 3. *In vitro* evaluation of compounds of Series III on SY5Y cells

	Ref	SY5Y	AICD	AICD	Aβ ₁₋₄₂	Ref	SY5Y	AICD	AICD	Aβ ₁₋₄₂
		CC ₅₀ (μM) ^a	C ₂ (μM) ^b	C ₁₀ (μM) ^c	IC ₅₀ (μM) ^d		CC ₅₀ (μM) ^a	C ₂ (μM) ^b	C ₁₀ (μM) ^c	IC ₅₀ (μM) ^d
CQ		30	1.4	10.8	12.7					
R
H	38	5	>5	nd	>5					
4-methoxyphenyl	39	10	>10	nd	^e					
4-chlorophenyl	40	5	0.5	5.0	3.1					
4-fluorophenyl	41	10	1.0	>10	1.0					
3,4-dimethoxyphenyl	42	20	>20	nd	^e					
4-quinolinyl	43	15	>10	nd	^e					
4-pyridinyl	44	>30	>10	nd	^e					
cyclopropyl	45	>30	13	nd	^e					
<i>i</i> -propyl	46	>30	>10	nd	^e	47	20	3.7	>20	>10

Mean values calculated on the basis of at least three independent experiments with less than 10% deviation; nd: not determined; ^a Compound concentration causing 50% of cell death after 24 h treatment; ^b Concentration doubling the quantity of AICD; ^c Concentration increasing the quantity of AICDs generated 10-fold; ^d Concentration decreasing the secretion of A β by 50%; ^e increase in the quantity of A β at 5 μ M.

Table 4. *In vitro* impact of compounds **25**, **29** and **33** on APP metabolism (SY5Y cells)

Ref	CTF α	CTF α	A β ₁₋₃₈	A β ₁₋₄₀	sAPP α ^d
	C ₂ (μ M) ^a	C ₁₀ (μ M) ^b	IC ₅₀ (μ M) ^c	IC ₅₀ (μ M) ^c	(10 μ M)
CQ	1.0	nd	10.0	7.0	+ 28%
25	0.2	11.2	5.0	9.1	+ 30%
29	0.5	>10	5.0	2.0	+ 78%
33	6.7	28.4	>10	5.1	+ 55%

Mean values calculated on the basis of at least three independent experiments with less than 10% deviation; nd: not determined; ^a Concentration doubling the quantity of CTF α ; ^b Concentration increasing the quantity of CTF α 10-fold; ^c Concentration inhibiting A β secretion by 50%; ^d increase in sAPP α secretion at 10 μ M.

Table 5. Preliminary ADME results

Ref	Aqueous solubility PBS pH 7.4 (μ M)	logD pH 7.4	Metabolic stability mLM ^a	Metabolic stability hLM ^b
25	227	0.1	100%	100%
29	278	1.8	79%	52%
33	200	1.2	30%	27%

Mean values calculated on the basis of two independent experiments with less than 10% deviation; ^a mLM: mouse liver microsomes; ^b hLM: human liver microsomes

Graphic Table of Contents