Nonlinear fourth order Taylor expansion of lattice Boltzmann schemes

François Dubois

► To cite this version:

François Dubois. Nonlinear fourth order Taylor expansion of lattice Boltzmann schemes. *Asymptotic Analysis*, 2022, 127 (4), pp.297-337. 10.3233/ASY-211692 . hal-02081116v2

HAL Id: hal-02081116

<https://hal.science/hal-02081116v2>

Submitted on 14 Jan 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Nonlinear fourth order Taylor expansion of lattice Boltzmann schemes

François Dubois^{ab}

^a *Laboratoire de Mathématiques d'Orsay, Faculté des Sciences d'Orsay,
Université Paris-Saclay, France.*

^b *Conservatoire National des Arts et Métiers, LMSSC laboratory, Paris, France.*

19 June 2020 *

Keywords: partial differential equations, asymptotic analysis

AMS classification: 65Q05, 76N15, 82C20.

PACS numbers: 02.70.Ns, 05.20.Dd, 47.10.+g

Abstract

We propose a formal expansion of multiple relaxation times lattice Boltzmann schemes in terms of a single infinitesimal numerical variable. The result is a system of partial differential equations for the conserved moments of the lattice Boltzmann scheme. The expansion is presented in the nonlinear case up to fourth order accuracy. The asymptotic corrections of the nonconserved moments are developed in terms of equilibrium values and partial differentials of the conserved moments. Both expansions are coupled and conduct to explicit compact formulas. The new algebraic expressions are validated with previous results obtained with this framework. The example of isothermal D2Q9 lattice Boltzmann scheme illustrates the theoretical framework.

* This contribution to appear in *Asymptotic Analysis* has been first presented at the Institut Henri Poincaré (Paris, France) the 10 January 2018. Edition 14 January 2021.

1) Introduction

The lattice Boltzmann schemes in their modern form have been developed with the contributions of d’Humières, Lallemand, Succi [38, 50, 37, 43] and many others. An underlying lattice Boltzmann equation is discretised on a cartesian grid with a finite set of velocities chosen in such a way that during one time step, an exact transport is done between two vertices of the mesh. A lattice Boltzmann scheme is composed with two steps: a nonlinear local relaxation step, followed by a linear advection scheme coupling a given vertex with a given family of neighbours. The relaxation step follows in general the “BGK” approximation introduced by Bhatnagar, Gross and Krook [4]. This numerical method allows the simulation of an important number of physical phenomena as isothermal flows, compressible flows with heat transfer, non-ideal fluids, multiphase and multi-component flows, microscale gas flows, soft-matter flows,... up to quantum mechanics. For the actual status of the method and the various applications, we refer *e.g.* to the books of Guo and Shu [32], Krüger *et al.* [35] and to the prospective article of Succi [56]. The lattice Boltzmann method is founded on a mesoscopic Boltzmann model but is not able in general to solve the Boltzmann equation or associated kinetic models. It is admitted that essentially macroscopic models are approximated with the lattice Boltzmann schemes.

The link between mesoscopic and macroscopic models is never straightforward and an analysis is necessary. The usual method of formal analysis is founded on the Chapman-Enskog [8] expansion in the spirit of the book of Huang [39]. The classical approach for the Boltzmann equation and other mesoscopic models posed in a continuum space-time. The Chapman-Enskog method has been adapted to lattice Boltzmann schemes with discrete space and time equations by Chen and Doolen [9] and by Qian and Zhou [51]. It allows the derivation the equivalent partial differential equations founded on a multiscale time analysis. This classical Chapman Enskog expansion is popular in the lattice Boltzmann community, as reported in the contributions of Lallemand and Luo [43], Philippi and Hegele [48] and Shan *et al.* [53] among others. A main drawback of this approach is the fact that multiscale expansions are used without a clear mathematical signification of the various variables and associated functions.

Independently of this framework, we have proposed in [14, 15] the Taylor expansion method for isothermal fluid problems to obtain formally equivalent partial differential equations at second order accuracy. This work was inspired by the method of equivalent equations for finite difference schemes, in a spirit proposed by Lerat and Peyret [45] and Warming and Hyett [57]. In this approach, the infinitesimal variable is simply the time step (proportional to the space step with the acoustic scaling) and there is only one infinitesimal scale for the analysis. This approach has been experimentally validated with Pierre Lallemand in various contributions [21, 22, 24, 42]. It was extended to third order accuracy for thermal and fluid problems in [16], automated at an arbitrary order in the linear case in [3], extended with an external drift in [25] and to relative velocities in [18, 19]. Our point of view is *a priori* not to consider the third and fourth order terms as approximations of Burnett or super-Burnett equations (see *e.g.* [1, 55]) but simply as errors of the lattice Boltzmann scheme when solving partial differential equations of second order type. Nevertheless, progress in

the approximation of fourth order partial differential equations have been obtained in [47]. Here, we have no particular fluid hypothesis. It can be applied to scalar or to vectorial [17, 29] lattice Boltzmann schemes. This Taylor expansion method has been also revisited with the Maxwell iterations of Yong *et al.* [58].

In this contribution, we first recall in Section 2 classic results related to the Boltzmann equation with discrete velocities. The link with multiple relaxation times lattice Boltzmann schemes as initially proposed by d’Humières [37] is presented in Section 3. Then we present the Taylor expansion method in a very general way and the result is explicit with compact formulas. Our main result (Section 4) concerns the derivation of asymptotic partial differential equations in the nonlinear case, up to fourth order accuracy. The proof for first and second orders is presented in Section 5, with a validation for the isothermal fluid case in the specific case of the D2Q9 scheme. In Section 6, the expansion up to third and fourth orders is detailed, with a confrontation with our previous work [16]. The proofs of these results need specific algebraic developments, presented in this contribution. The linear case with constant coefficients is described in Section 7.

2) Boltzmann equation with discrete velocities

In the space $\mathbb{R}^d$ of dimension d , we consider a finite set of q discrete velocities $v_j \in \mathcal{V}$ with components v_j^α for $1 \leq \alpha \leq d$. The unknowns of the Boltzmann equation are the particle densities f_j . They are functions of space x , time t and discrete velocities v_j :

$$f_j = f_j(x, t), \quad x \in \mathbb{R}^d, t \geq 0, 0 \leq j < q.$$

The vector $f(x, t) \in \mathbb{R}^q$ is constructed with the numbers $f_j(x, t)$ for $0 \leq j < q$. We introduce a collision vector $\mathbb{R}^q \ni f \mapsto Q(f) \in \mathbb{R}^q$ of components $Q_j(f)$ for $0 \leq j < q$ with given regular functions Q_j . We introduce also a small parameter $\varepsilon > 0$ that can be interpreted as a Knudsen number in the context of gas dynamics (see *e.g.* [8]).

The Boltzmann model with discrete velocities takes the form

$$(1) \quad \frac{\partial f_j}{\partial t} + \sum_{1 \leq \alpha \leq d} v_j^\alpha \frac{\partial f_j}{\partial x_\alpha} = \frac{1}{\varepsilon} Q_j(f), \quad 0 \leq j < q.$$

It has been proposed by Carleman [7], Gross [31], Broadwell [6] and intensively developed by Gatignol [27] and her co-workers.

• Moments

Our framework concerns multi relaxation times: we introduce a constant invertible matrix M called “d’Humières matrix” [37] in this contribution. This matrix defines the vector of moments $m \in \mathbb{R}^q$ by a simple product:

$$(2) \quad m_k \equiv \sum_{0 \leq j < q} M_{kj} f_j.$$

We introduce the number of conservation laws N ($1 \leq N < q$) such that the N first moments of the collision kernel are equal to zero:

$$(3) \quad \sum_{0 \leq j < q} M_{kj} Q_j(f) = 0, \quad \forall f \in \mathbb{R}^q, 0 \leq k < N.$$

Then it is natural to divide the vector of moments into two families:

$$(4) \quad m \equiv \begin{pmatrix} W \\ Y \end{pmatrix}.$$

The conserved moments or macroscopic variables W constitute a linear space of dimension N and if the Boltzmann model with discrete velocities (1) is satisfied, we have N conservation laws due to the cancellation (3):

$$\frac{\partial W_k}{\partial t} + \sum_{1 \leq \alpha \leq d} \sum_{0 \leq j < q} M_{kj} v_j^\alpha \frac{\partial f_j}{\partial x_\alpha} = 0, \quad 0 \leq k < N.$$

Observe that the nonconserved moments or microscopic variables Y generate a linear space of dimension $q - N$.

• Equilibrium states

We suppose that the equilibrium states f^{eq} defined by the conditions

$$Q(f^{\text{eq}}) = 0$$

are characterized with the help of a regular nonlinear vector field $\Phi : \mathbb{R}^N \longrightarrow \mathbb{R}^{q-N}$ such that

$$(5) \quad f^{\text{eq}} = M^{-1} \begin{pmatrix} W \\ \Phi(W) \end{pmatrix}.$$

In other words, the vector field $W \longmapsto Y^{\text{eq}} \equiv \Phi(W)$ defines the set of equilibrium states.

We suppose moreover that the jacobian matrix $dQ(f^{\text{eq}})$ at equilibrium is diagonalizable with real eigenvalues and real eigenvectors. More precisely, taking into account the hypothesis (3), we suppose that there exists a diagonal matrix

$$Z = \text{diag} \left(\frac{1}{\tau_1}, \dots, \frac{1}{\tau_{q-N}} \right)$$

of order $q - N$ with strictly positive coefficients τ_ℓ such that

$$(6) \quad M dQ(f^{\text{eq}}) M^{-1} = - \begin{pmatrix} 0 & 0 \\ 0 & Z \end{pmatrix}.$$

In other words, the jacobian operator $dQ(f^{\text{eq}})$ admits the matrix M^{-1} as a matrix of right eigenvectors and the associated eigenvalues are all non positive.

• Momentum-velocity operator

For a linear space of dimension d , we introduce the momentum-velocity operator matrix Λ defined by the relation

$$(7) \quad \Lambda = M \text{diag} \left(\sum_{1 \leq \alpha \leq d} v^\alpha \partial_\alpha \right) M^{-1}.$$

It is a $q \times q$ operator matrix composed by first-order space differential operators. It is obtained by conjugation of the first order advection operator $v \cdot \nabla$ by the d'Humières matrix M . The operator matrix Λ is nothing else than the advection operator seen in the basis of moments. We introduce a block decomposition of the momentum-velocity operator matrix associated to the decomposition (4) of the moments. We define a $N \times N$ operator

matrix A , a $N \times (q - N)$ operator matrix B , a $(q - N) \times N$ operator matrix C and a $(q - N) \times (q - N)$ operator matrix D according to

$$(8) \quad \Lambda \equiv \begin{pmatrix} A & B \\ C & D \end{pmatrix}.$$

Remember that in the following, the matrices A , B , C and D are matrices composed with first order space operators.

• **Boltmann-BGK system with discrete velocities**

We approximate now the previous discrete model with a BGK [4] type hypothesis: the state f is close to equilibrium and we approach the collision kernel Q by its first order expansion around the equilibrium state:

$$Q(f) \simeq Q(f^{\text{eq}}) + dQ(f^{\text{eq}}).(f - f^{\text{eq}}).$$

Due to the condition $Q(f^{\text{eq}}) = 0$, we obtain with this approximation that we call “BGK” in this contribution, even it is not exactly the hypothesis done in the original article [4], a new system of partial differential equations:

$$(9) \quad \frac{\partial f_j}{\partial t} + \sum_{1 \leq \alpha \leq d} v_j^\alpha \frac{\partial f_j}{\partial x_\alpha} = \frac{1}{\varepsilon} dQ_j(f^{\text{eq}}).(f - f^{\text{eq}}), \quad 0 \leq j < q.$$

Observe that f^{eq} is a function of the particle distribution f through the relations (2), (4) and (5). Taking into account the hypothesis (6), the definition (7) and the notation (8), we can write the Boltmann-BGK system with discrete velocities (9) under the form of a system of two coupled partial differential equations:

$$(10) \quad \frac{\partial W}{\partial t} + A W + B Y = 0, \quad \frac{\partial Y}{\partial t} + C W + D Y = -\frac{1}{\varepsilon} Z (Y - \Phi(W)).$$

Proposition 1. Fourth order Chapman-Enskog expansion of the Boltmann-BGK system with discrete velocities

As ε tends to zero, the conserved moments W of the Boltmann-BGK system with discrete velocities (10) satisfy formally the following asymptotic system of partial differential equations:

$$(11) \quad \frac{\partial W}{\partial t} + \Gamma_1(W) + \varepsilon \Gamma_2(W) + \varepsilon^2 \Gamma_3(W) + \varepsilon^3 \Gamma_4(W) = O(\varepsilon^4).$$

The nonlinear operators are related to the expansion of the microscopic moments:

$$(12) \quad Y = \Phi(W) + \varepsilon Z^{-1} \Psi_1(W) + \varepsilon^2 Z^{-1} \Psi_2(W) + \varepsilon^3 Z^{-1} \Psi_3(W) + O(\varepsilon^4)$$

and we have the interlaced relations

$$(13) \quad \left\{ \begin{array}{l} \Gamma_1(W) = A W + B \Phi(W) \\ \Psi_1(W) = d\Phi(W).\Gamma_1(W) - (C W + D \Phi(W)) \\ \Gamma_2(W) = B Z^{-1} \Psi_1(W) \\ \Psi_2(W) = Z^{-1} d\Psi_1(W).\Gamma_1(W) + d\Phi(W).\Gamma_2(W) - D Z^{-1} \Psi_1(W) \\ \Gamma_3(W) = B Z^{-1} \Psi_2(W) \\ \Psi_3(W) = Z^{-1} d\Psi_1(W).\Gamma_2(W) + d\Phi(W).\Gamma_3(W) - D Z^{-1} \Psi_2(W) \\ \quad + Z^{-1} d\Psi_2(W).\Gamma_1(W) \\ \Gamma_4(W) = B Z^{-1} \Psi_3(W). \end{array} \right.$$

In applications to fluid dynamics, the first order term $\Gamma_1(W)$ is associated to the Euler equations of gas dynamics and the second order term $\Gamma_2(W)$ to the viscous terms of the Navier-Stokes equations. The operators $\Gamma_3(W)$ and $\Gamma_4(W)$ represent the Burnett and super-Burnett terms respectively. With this Proposition 1, we essentially reformulate the classic Chapman Enskog expansion [8] in the discrete velocity case, as done *e.g.* in the book of Gatignol [27].

- Proof of Proposition 1.

First we inject the representation (12) inside the first equation of (10). It comes

$$\frac{\partial W}{\partial t} + A W + B (\Phi(W) + \varepsilon Z^{-1} \Psi_1(W) + \varepsilon^2 Z^{-1} \Psi_2(W) + \varepsilon^3 Z^{-1} \Psi_3(W)) = O(\varepsilon^4)$$

then after rearranging the terms,

$$\frac{\partial W}{\partial t} + A W + B \Phi(W) + \varepsilon B Z^{-1} \Psi_1(W) + \varepsilon^2 B Z^{-1} \Psi_2(W) + \varepsilon^3 B Z^{-1} \Psi_3(W) = O(\varepsilon^4).$$

The confrontation of this partial differential equation with the Ansatz (11) shows that

$$\begin{cases} \Gamma_1(W) = A W + B \Phi(W) \\ \Gamma_2(W) = B Z^{-1} \Psi_1(W) \\ \Gamma_3(W) = B Z^{-1} \Psi_2(W) \\ \Gamma_4(W) = B Z^{-1} \Psi_3(W) \end{cases}$$

as proposed in (13).

Secondly, we write the second equation of (10) under the form

$$Y = \Phi - \varepsilon Z^{-1} (\partial_t Y + C W + D Y).$$

We observe that

$$\begin{aligned} -\partial_t Y &= dY \cdot (-\partial_t W) \\ &= d(\Phi(W) + \varepsilon Z^{-1} \Psi_1(W) + \varepsilon^2 Z^{-1} \Psi_2(W)) \cdot (\Gamma_1 + \varepsilon \Gamma_2(W) + \varepsilon^2 \Gamma_3(W)) + O(\varepsilon^3) \\ &= d\Phi(W) \cdot \Gamma_1 + \varepsilon (Z^{-1} d\Psi_1(W) \cdot \Gamma_1 + d\Phi(W) \cdot \Gamma_2) \\ &\quad + \varepsilon^2 (Z^{-1} d\Psi_2(W) \cdot \Gamma_1 + Z^{-1} d\Psi_1(W) \cdot \Gamma_2 + d\Phi(W) \cdot \Gamma_3) + O(\varepsilon^3). \end{aligned}$$

We report this expression inside the value of Y :

$$\begin{aligned} Y &= \Phi + \varepsilon Z^{-1} (-\partial_t Y - C W - D Y) \\ &= \Phi + \varepsilon Z^{-1} [d\Phi(W) \cdot \Gamma_1 + \varepsilon (Z^{-1} d\Psi_1(W) \cdot \Gamma_1 + d\Phi(W) \cdot \Gamma_2) \\ &\quad + \varepsilon^2 (Z^{-1} d\Psi_2(W) \cdot \Gamma_1 + Z^{-1} d\Psi_1(W) \cdot \Gamma_2 + d\Phi(W) \cdot \Gamma_3) \\ &\quad - C W - D (\Phi(W) + \varepsilon Z^{-1} \Psi_1(W) + \varepsilon^2 Z^{-1} \Psi_2(W))] + O(\varepsilon^4) \\ &= \Phi + \varepsilon Z^{-1} (d\Phi(W) \cdot \Gamma_1 - C W - D \Phi) + \varepsilon^2 Z^{-1} (Z^{-1} d\Psi_1(W) \cdot \Gamma_1 + d\Phi(W) \cdot \Gamma_2 - D Z^{-1} \Psi_1) \\ &\quad + \varepsilon^3 Z^{-1} (Z^{-1} d\Psi_2(W) \cdot \Gamma_1 + Z^{-1} d\Psi_1(W) \cdot \Gamma_2 + d\Phi(W) \cdot \Gamma_3 - D Z^{-1} \Psi_2) + O(\varepsilon^4). \end{aligned}$$

Comparatively to the relation (12), we obtain

$$\begin{cases} \Psi_1(W) = d\Phi(W) \cdot \Gamma_1 - (C W + D \Phi) \\ \Psi_2(W) = Z^{-1} d\Psi_1(W) \cdot \Gamma_1 + d\Phi(W) \cdot \Gamma_2 - D Z^{-1} \Psi_1(W) \\ \Psi_3(W) = Z^{-1} d\Psi_1(W) \cdot \Gamma_2 + d\Phi(W) \cdot \Gamma_3 - D Z^{-1} \Psi_2 + Z^{-1} d\Psi_2(W) \cdot \Gamma_1 \end{cases}$$

and the last relations of (13) are explicited. The proof is completed. $\square$

3) Multiple Relaxation Times Lattice Boltzmann schemes

We introduce now a regular cartesian lattice $\mathcal{L}$ composed by vertices x separated by distances that are simple expressions of the space step Δx . A discrete time t is supposed to be an integer multiple of a time step $\Delta t > 0$. The unknowns of a lattice Boltzmann scheme with q discrete velocities are the particle densities f_j . They are now functions of discrete space x , discrete time t and discrete velocities v_j for $0 \leq j < q$:

$$f_j = f_j(x, t), \quad x \in \mathcal{L}, t = n \Delta t, n \in \mathbb{N}, v_j \in \mathcal{V}.$$

The discrete velocity set $\mathcal{V}$ does not depend on space or time.

Multiple relaxation times lattice Boltzmann scheme considered in this contribution consists in a simple Lie-Trotter splitting applied to the Boltzmann-BGK model with discrete velocities (9). First a relaxation step $f(t) \rightarrow R(\Delta t) f(t) \equiv f^*$ for the approximation of the system of ordinary differential equations

$$\frac{\partial f}{\partial t} = \frac{1}{\varepsilon} \mathrm{d}Q(f^{\mathrm{eq}}) \cdot (f - f^{\mathrm{eq}})$$

and secondly a transport step $f^* \rightarrow T(\Delta t) f^* \equiv f(t + \Delta t)$ for the free advection

$$\frac{\partial f_j}{\partial t} + \sum_{1 \leq \alpha \leq d} v_j^\alpha \frac{\partial f_j}{\partial x_\alpha} = 0, \quad 0 \leq j < q.$$

• Relaxation step

Due to the formulation (10), the relaxation scheme is simply described in the moment representation and we have

$$\frac{\partial W}{\partial t} = 0, \quad \frac{\partial Y}{\partial t} = -\frac{1}{\varepsilon} Z(Y - \Phi(W)).$$

After relaxation, the vector of moments m is transformed into a new vector m^* :

$$(14) \quad m^* \equiv \begin{pmatrix} W^* \\ Y^* \end{pmatrix}.$$

The conserved moments are not modified during this relaxation step:

$$(15) \quad W^* = W.$$

As remarked in [15], we use an explicit first order Euler scheme for the microscopic moments:

$$\frac{Y^* - Y}{\Delta t} = -\frac{1}{\varepsilon} Z(Y - \Phi(W)),$$

id est

$$(16) \quad Y^* = Y + S(\Phi(W) - Y)$$

with

$$(17) \quad S = \frac{\Delta t}{\varepsilon} Z.$$

Usually, as pointed by Lallemand and Luo [43], the relaxation matrix S is a diagonal matrix, in coherence with the hypothesis concerning the diagonalization of the operator $\mathrm{d}Q(f^{\mathrm{eq}})$ presented in the previous section:

$$S = \mathrm{diag}(s_1, \dots, s_{q-N}).$$

Moreover, the matrix S is a fixed invertible square matrix of dimension $(q - N) \times (q - N)$. In other words, the time step Δt is of the order of the Knudsen number ε in our asymptotic study. Finally the particle distribution f^* after relaxation is given according to

$$(18) \quad f^* = M^{-1} m^*.$$

- **Necessary conditions for stability**

We deduce from the relation (16)

$$Y^* - \Phi(W) = (I - S)(Y - \Phi(W))$$

and a natural stability condition takes the form $\|I - S\| \leq 1$, *id est*

$$0 \leq s_j \leq 2, \quad 0 \leq j < q - N.$$

The case of over-relaxation $1 < s_j < 2$ is widely used when implementing lattice Boltzmann schemes to take into account very low viscosities in fluid mechanics. We will emphasize this point in the next section.

- **Advection step**

In the second step of advection, the method of characteristics is applied in the very particular case when it is exact. In one time step each particle distribution f_j is exactly transported from the node $x - v_j \Delta t$ of the lattice $\mathcal{L}$ to the vertex $x \in \mathcal{L}$. In other words, the advection step occurs with a Courant-Friedrichs-Lewy number $cfl \equiv 1$ for all discrete velocities v_j .

- **Lattice Boltzmann scheme as a Lie-Trotter splitting scheme**

The iteration of the lattice Boltzmann scheme is finally given by the relation

$$(19) \quad f(t + \Delta t) = T(\Delta t) R(\Delta t) f(t)$$

and it is useful to explicit it for each component:

$$(20) \quad f_j(x, t + \Delta t) = f_j^*(x - v_j \Delta t, t), \quad v_j \in \mathcal{V}, \quad x \in \mathcal{L}.$$

Observe that this framework is very general. The so-called “BGK version” of the lattice Boltzmann scheme [5, 38, 50] corresponds to the choice $M = I$ and $Z = \frac{1}{\tau} I$. Usual multi-relaxation lattice Boltzmann schemes [20, 21, 37, 43] suppose in general some orthogonality property for the d’Humières matrix. This paradigm permits also the introduction of two particle distributions [2, 20, 33, 40] or eventually more with the vectorial schemes developed by Graille [29] or a variant proposed in [17]. In those last cases, the numbering $j \mapsto v_j$ of the velocities is simply non injective.

It is naturally possible to think to a second order accurate splitting scheme. A natural idea is the Strang splitting [54]:

$$f(t + \Delta t) = R\left(\frac{\Delta t}{2}\right) T(\Delta t) R\left(\frac{\Delta t}{2}\right) f(t).$$

In this case, as put in evidence by Dellar [11], one can write $\tilde{f}(t) \equiv R\left(\frac{\Delta t}{2}\right)^{-1} f(t)$ and the previous Strang scheme can be written as

$$\tilde{f}(t + \Delta t) = T(\Delta t) R\left(\frac{\Delta t}{2}\right) R\left(\frac{\Delta t}{2}\right) \tilde{f}(t).$$

This scheme, up to the shift of one half relaxation step, is identical to the Lie-Trotter splitting (19), except that the product $R(\frac{\Delta t}{2}) R(\frac{\Delta t}{2})$ plays the role of $R(\Delta t)$. A first possibility is to divide the time step Δt by a factor 2 in the relations (16)(17):

$$\bar{Y} = Y + \frac{1}{2} S (\Phi - Y), \quad Y^* = \bar{Y} + \frac{1}{2} S (\Phi - \bar{Y}).$$

Then

$$Y^* = \left(1 - S + \frac{1}{4} S^2\right) Y + \left(S - \frac{1}{4} S^2\right) \Phi(W)$$

and over-relaxation is no more possible because $s_j - \frac{1}{4} s_j^2 \leq 1$ for all $s_j \in \mathbb{R}$. So such a numerical scheme is not used by the engineers at our knowledge.

A second possibility is to enforce the relation $R(\frac{\Delta t}{2}) = \sqrt{R(\Delta t)}$, *id est*

$$\bar{Y} = \sqrt{I - S} Y + (I - \sqrt{I - S}) \Phi(W).$$

But this relation has a sense only if $s_j \leq 1$ and over-relaxation is again excluded.

An other idea is to exchange the roles of advection and relaxation in the Strang scheme:

$$f(t + \Delta t) = T\left(\frac{\Delta t}{2}\right) R(\Delta t) T\left(\frac{\Delta t}{2}\right) f(t)$$

or to use more elaborated splittings as proposed by Drui *et al.* [13]. In this case, the transport step is nomore at a Courant number equal to unity and an interpolation procedure, costly in terms of numerical viscosity or complex to implement as the semi-Lagrangian method [26], is not usually chosen by experts developping lattice Boltzmann schemes.

In conclusion of this sub-section, the naive Lie-Trotter splitting scheme (19) gives very good numerical results. It is important to put in evidence precise qualities and defects of this scheme through the following formal asymptotic analysis.

Proposition 2. Exponential form of lattice Boltzmann schemes

Consider a lattice Boltzmann scheme as defined at the previous section by the relations (2), (4), (5), (7), (8), (14), (15), (16), (18) and (20). Then we have an exponential form of the discrete iteration (20):

$$(21) \quad m(x, t + \Delta t) = \exp(-\Delta t \Lambda) m^*(x, t).$$

- Proof of Proposition 2.

We have simply a long sequence of identities:

$$\begin{aligned} m_k(x, t + \Delta t) &= \sum_j M_{kj} f_j(x, t + \Delta t) && \text{due to (2)} \\ &= \sum_j M_{kj} f_j^*(x - v_j \Delta t, t) && \text{due to (20)} \\ &= \sum_{j\ell} M_{kj} (M^{-1})_{j\ell} m_\ell^*(x - v_j \Delta t, t) && \text{due to (18)} \\ &= \sum_{j\ell} M_{kj} (M^{-1})_{j\ell} \sum_{n=0}^{\infty} \frac{1}{n!} (-\Delta t \sum_{\alpha} v_j^\alpha \partial_\alpha)^n m_\ell^*(x, t) && \text{Taylor expansion} \\ &= \sum_{\ell} \sum_{n=0}^{\infty} \frac{1}{n!} \sum_j M_{kj} \left(-\Delta t \sum_{\alpha} v_j^\alpha \partial_\alpha\right)^n (M^{-1})_{j\ell} m_\ell^*(x, t) && \text{elementary algebra} \end{aligned}$$

$$\begin{aligned}
 &= \sum_{\ell} \sum_{n=0}^{\infty} \frac{1}{n!} (-\Delta t)^n \left(\left(M \left(\sum_{\alpha} v^{\alpha} \partial_{\alpha} \right) M^{-1} \right)^n \right)_{k\ell} m_{\ell}^*(x, t) && \text{because } M \text{ is a constant matrix} \\
 &= \sum_{\ell} \sum_{n=0}^{\infty} \frac{1}{n!} (-\Delta t)^n (\Lambda^n)_{k\ell} m_{\ell}^*(x, t) && \text{due to (7)} \\
 &= \sum_{\ell} \left[\sum_{n=0}^{\infty} \frac{1}{n!} (-\Delta t \Lambda)_{k\ell}^n \right] m_{\ell}^*(x, t) && \text{elementary algebra} \\
 &= \sum_{\ell} \exp(-\Delta t \Lambda)_{k\ell} m_{\ell}^*(x, t) && \text{elementary algebra} \\
 &= \left(\exp(-\Delta t \Lambda) m^*(x, t) \right)_k && \text{elementary algebra}
 \end{aligned}$$

and the proof is completed. $\square$

We can see the relation (21) as a discrete form of the Duhamel formula for ordinary differential equations. The idea of introducing a formal exponential expansion is also present in the work of Boghosian and Coveney in the BGK case [5].

• Momentum-velocity operator for a D2Q9 lattice Boltzmann scheme

We can concretize the matrix Λ introduced in (7) with the popular two-dimensional ‘‘D2Q9’’ scheme. A set of nine velocities is defined by the Figure 1.

Figure 1: D2Q9 lattice Boltzmann scheme

To fix the ideas, we use the moment matrix presented in Lallemand and Luo [43]:

$$(22) \quad M_{D2Q9} = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 0 & \lambda & 0 & -\lambda & 0 & \lambda & -\lambda & -\lambda & \lambda \\ 0 & 0 & \lambda & 0 & -\lambda & \lambda & \lambda & -\lambda & -\lambda \\ -4\lambda^2 & -\lambda^2 & -\lambda^2 & -\lambda^2 & -\lambda^2 & 2\lambda^2 & 2\lambda^2 & 2\lambda^2 & 2\lambda^2 \\ 0 & \lambda^2 & -\lambda^2 & \lambda^2 & -\lambda^2 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \lambda^2 & -\lambda^2 & \lambda^2 & -\lambda^2 \\ 0 & -2\lambda^3 & 0 & 2\lambda^3 & 0 & \lambda^3 & -\lambda^3 & -\lambda^3 & \lambda^3 \\ 0 & 0 & -2\lambda^3 & 0 & 2\lambda^3 & \lambda^3 & \lambda^3 & -\lambda^3 & -\lambda^3 \\ 4\lambda^4 & -2\lambda^4 & -2\lambda^4 & -2\lambda^4 & -2\lambda^4 & \lambda^4 & \lambda^4 & \lambda^4 & \lambda^4 \end{bmatrix}.$$

Observe that other choices are possible with the Hermite moments developed in [48, 53]. The moments are associated with the lines of the matrix (22). They are denominated (in

this order) by $\rho, J_x, J_y, \varepsilon, x_x, x_y, q_x, q_y, h$. The lines of this invertible matrix are chosen orthogonal. For isothermal flows, the conserved moments

$$(23) \quad W = (\rho, J_x, J_y)^t$$

correspond to the three first lines of the d'Humières matrix (22). The non-conserved moments complete the family:

$$(24) \quad Y = (\varepsilon, x_x, x_y, q_x, q_y, h)^t.$$

They are linked to the six last lines of the matrix defined in (22). With the velocities defined in Figure 1 and the moment matrix (22), we construct without difficulty the operator matrix Λ for the thermal D2Q9 scheme:

$$(25) \quad \Lambda_{D2Q9} = \left[\begin{array}{ccc|cccccc} 0 & \partial_x & \partial_y & 0 & 0 & 0 & 0 & 0 & 0 \\ \frac{2\lambda^2}{3} \partial_x & 0 & 0 & \frac{1}{6} \partial_x & \frac{1}{2} \partial_x & \partial_y & 0 & 0 & 0 \\ \frac{2\lambda^2}{3} \partial_y & 0 & 0 & \frac{1}{6} \partial_y & -\frac{1}{2} \partial_y & \partial_x & 0 & 0 & 0 \\ \hline 0 & \lambda^2 \partial_x & \lambda^2 \partial_y & 0 & 0 & 0 & \partial_x & \partial_y & 0 \\ 0 & \frac{\lambda^2}{3} \partial_x & -\frac{\lambda^2}{3} \partial_y & 0 & 0 & 0 & -\frac{1}{3} \partial_x & \frac{1}{3} \partial_y & 0 \\ 0 & \frac{2}{3} \lambda^2 \partial_y & \frac{2}{3} \lambda^2 \partial_x & 0 & 0 & 0 & \frac{1}{3} \partial_y & \frac{1}{3} \partial_x & 0 \\ \hline 0 & 0 & 0 & \frac{\lambda^2}{3} \partial_x & -\lambda^2 \partial_x & \lambda^2 \partial_y & 0 & 0 & \frac{1}{3} \partial_x \\ 0 & 0 & 0 & \frac{\lambda^2}{3} \partial_y & \lambda^2 \partial_y & \lambda^2 \partial_x & 0 & 0 & \frac{1}{3} \partial_y \\ 0 & 0 & 0 & 0 & 0 & 0 & \lambda^2 \partial_x & \lambda^2 \partial_y & 0 \end{array} \right].$$

In the relation (25), we have put in evidence the block decomposition (8) for the D2Q9 scheme with the conserved moments precised in (23). Even if the space differential operators commute, the matrices that compose the momentum-velocity operator matrix does not commute! We have the very important but elementary structure

$$\Lambda^2 \equiv \begin{pmatrix} A_2 & B_2 \\ C_2 & D_2 \end{pmatrix} = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \begin{pmatrix} A & B \\ C & D \end{pmatrix} = \begin{pmatrix} A^2 + B C & A B + B D \\ C A + D C & C B + D^2 \end{pmatrix}$$

and to fix the ideas

$$\Lambda^3 \equiv \begin{pmatrix} A_3 & B_3 \\ C_3 & D_3 \end{pmatrix} = \begin{pmatrix} A_2 & B_2 \\ C_2 & D_2 \end{pmatrix} \begin{pmatrix} A & B \\ C & D \end{pmatrix} = \begin{pmatrix} A_2 A + B_2 C & A_2 B + B_2 D \\ C_2 A + D_2 C & C_2 B + D_2 D \end{pmatrix}.$$

4) Taylor expansion method

In this section, we revisit the Taylor expansion method: we specify the hypotheses, express our general result and begin the proof with the order zero.

• Hypotheses for a formal expansion

The formal Taylor expansion supposes some precise hypotheses. First, we adopt the acoustic scaling: the ratio $\lambda \equiv \frac{\Delta x}{\Delta t}$ is supposed to be constant in all this work. Moreover, the ratio $\frac{\Delta t}{\varepsilon}$ remains constant in our asymptotic analysis. Then the relaxation matrix S is fixed and invertible; it is also the case for the matrix S^{-1} proportional to the matrix Z^{-1} who plays an important role in the relations (13) of the Chapman-Enskog expansion (11)(12)(13) for the continuous in time lattice Boltzmann-BGK system.

A unintuitive result has been discovered by Hénon [36]. Due to the fact that the fully discretized in time lattice Boltzmann scheme (20) is substantially different from the continuous in time Boltzmann-BGK system with discrete velocities (9), the asymptotic expansions have similarities and differences. In particular, the lattice Boltzmann scheme put in evidence what we call the Hénon matrix Σ in this contribution. Roughly speaking, it plays a role analogous to the matrix Z^{-1} in the relations (13). But it is defined by

$$(26) \quad \Sigma \equiv S^{-1} - \frac{1}{2} \mathbf{I}.$$

This matrix emerges from the very classic second order analysis presented thereafter at Proposition 5 and detailed in the relations (40)(41). For applications to fluid dynamics, this matrix is closely related to viscosities, as explained in Section 5 for the D2Q9 example in relations (50) and (51). Then some values of the Hénon matrix Σ are chosen as small as possible in order to simulate flows with high Reynolds number. In consequence, over-relaxation is a mandatory practice for lattice Boltzmann schemes applied to high Reynolds number flows. Recall that this matrix remains fixed in our analysis.

Secondly, we suppose also that we can differentiate all the expansions relative to space and/or time. This hypothesis is mathematically absolutely non trivial and expresses that all the formal expansions should take place in very regular functional spaces. Thirdly, we use the notation $d\zeta(W).\xi$ for the action of the differential of some regular function $W \mapsto \zeta(W)$ against a test vector ξ . In particular,

$$(27) \quad \gamma_j(W) \equiv d\Phi(W).\Gamma_j(W), \quad 1 \leq j \leq 3.$$

For second order derivatives, we introduce

$$(28) \quad \partial^2 \Psi.\Gamma_1 \equiv d(d\Psi(W).\Gamma_1).\Gamma_1 = d^2\Psi(W).(\Gamma_1, \Gamma_1) + d\Psi(W).d\Gamma_1(W).\Gamma_1.$$

The asymptotic analysis occurs for a time step Δt (or a space step Δx) tending to zero. Emerging partial differential equations for the conserved variables are denoted as

$$(29) \quad \partial_t W + \Gamma_1(W) + \Delta t \Gamma_2(W) + \Delta t^2 \Gamma_3(W) + \Delta t^3 \Gamma_4(W) = O(\Delta t^4).$$

The vector $\Gamma_j(W)$ belongs in $\mathbb{R}^N$. It is obtained after j space derivations of the conserved moments W and of the equilibrium vector $\Phi(W)$. For this reason, we will speak in the following of *e.g.* $\Gamma_1(W)$ as “first order term”, even if it is a term of order zero relative to the infinitesimal Δt . For non-conserved moments (or microscopic variables), we suppose

$$(30) \quad \begin{cases} Y = \Phi(W) + S^{-1} (\Delta t \Psi_1(W) + \Delta t^2 \Psi_2(W) + \Delta t^3 \Psi_3(W)) + O(\Delta t^4) \\ Y^* = \Phi(W) + (\Sigma - \frac{1}{2} \mathbf{I}) (\Delta t \Psi_1(W) + \Delta t^2 \Psi_2(W) + \Delta t^3 \Psi_3(W)) + O(\Delta t^4) \end{cases}$$

where $\mathbf{I}$ is the identity matrix of dimension $q - N$. Observe that the two lines of (30) are equivalent thanks to (16) and (26). The vectors $\Phi_j(W)$ are analogous to $\Gamma_j(W)$ but not with the same dimension: $\Gamma_j(W) \in \mathbb{R}^N$ and $\Phi_j(W) \in \mathbb{R}^{q-N}$.

• **Main result. Recurrence formulas for fourth order expansion**

We prove in this contribution that we have the following recurrence formulas for the explicitation of the vectors $\Gamma_j(W)$ and $\Phi_j(W)$ up to fourth order accuracy:

$$(31) \quad \left\{ \begin{array}{l} \Gamma_1(W) = A W + B \Phi(W) \\ \Psi_1(W) = d\Phi(W) \cdot \Gamma_1(W) - (C W + D \Phi(W)) \\ \Gamma_2(W) = B \Sigma \Psi_1(W) \\ \Psi_2(W) = \Sigma d\Psi_1(W) \cdot \Gamma_1(W) + d\Phi(W) \cdot \Gamma_2(W) - D \Sigma \Psi_1(W) \\ \Gamma_3(W) = B \Sigma \Psi_2(W) + \frac{1}{12} B_2 \Psi_1(W) - \frac{1}{6} B d\Psi_1(W) \cdot \Gamma_1(W) \\ \Psi_3(W) = \Sigma d\Psi_1(W) \cdot \Gamma_2(W) + d\Phi(W) \cdot \Gamma_3(W) - D \Sigma \Psi_2(W) + \Sigma d\Psi_2(W) \cdot \Gamma_1(W) \\ \quad + \frac{1}{6} D d\Psi_1(W) \cdot \Gamma_1(W) - \frac{1}{12} D_2 \Psi_1(W) - \frac{1}{12} \partial^2 \Psi_1(W) \cdot \Gamma_1(W) \\ \Gamma_4(W) = B \Sigma \Psi_3(W) + \frac{1}{4} B_2 \Psi_2(W) + \frac{1}{6} B D_2 \Sigma \Psi_1(W) - \frac{1}{6} A B \Psi_2(W) \\ \quad - \frac{1}{6} B (d\gamma_1(W) \cdot \Gamma_2(W) + d\gamma_2(W) \cdot \Gamma_1(W)) - \frac{1}{6} B \Sigma \partial^2 \Psi_1(W) \cdot \Gamma_1(W) . \end{array} \right.$$

Observe that the order of the relations in (31) is mandatory. Each step has to be explicitated before the evaluation of the next expression. At second order accuracy, the three first lines of the relations (31) summarizes what is needed for a lot of applications with only first and second order partial differential equations. Observe that the compact form $\Gamma_2 = B \Sigma \Psi_1$ for the second order term put in evidence the important choice of a Hénon matrix “as small as possible”. Observe that the relations (31) have similarities with the ones obtained in (13) for the Chapman-Enskog expansion. But essentially the matrix Σ has replaced the matrix Z^{-1} and new terms appear due to the post-processing with Taylor expansions.

Proposition 3. Equilibrium state and zero order expansion

When Δt tends to zero, the microscopic moments are close to their equilibrium value:

$$(32) \quad Y = \Phi(W) + O(\Delta t), \quad Y^* = \Phi(W) + O(\Delta t).$$

• **Proof of Proposition 3.**

We expand one iteration of the scheme (21) at order zero:

$$(33) \quad m + O(\Delta t) = m^* + O(\Delta t^3).$$

Then we deduce from (33):

$$Y - Y^* = O(\Delta t).$$

Then due to the iteration (16) and the fact that the matrix S is supposed fixed, we have

$$(34) \quad Y = \Phi(W) + O(\Delta t).$$

The first relation of (32) is satisfied. It is then immediate to deduce from the scheme iteration (16) the second relation of (32). The proposition is established. $\square$

In the following, we detail all the ingredients that conduct to the main result (29) to (31).

5) Taylor expansion method at first and second order accuracy

In this section, we prove the two first orders of the expansion (29)(30)(31). We make the link with the Taylor expansion as presented in [14, 15] and we apply the result for the isothermal D2Q9 scheme.

Proposition 4. First order expansion

When Δt tends to zero, the macroscopic moments W satisfy asymptotically the following system of N first order equations:

$$(35) \quad \partial_t W + \Gamma_1(W) = O(\Delta t).$$

Moreover, the vector $\Gamma_1(W)$ for the first order dynamics satisfy

$$(36) \quad \Gamma_1(W) = A W + B \Phi(W).$$

- Proof of Proposition 4.

We expand one iteration of the scheme (21) at first order:

$$(37) \quad m + \Delta t \partial_t m + O(\Delta t^2) = m^* - \Delta t \Lambda m^* + O(\Delta t^2).$$

We can replace the vector m by its two components W and Y . We have for the first line

$$W + \Delta t \partial_t W + O(\Delta t^2) = W - \Delta t (A W + B Y^*) + O(\Delta t^2).$$

We simplify by the constant W term, divide by Δt and take into account the expansion (32):

$$\partial_t W + O(\Delta t) = -(A W + B \Phi(W)) + O(\Delta t).$$

This relation is exactly the expansion (35) with the first order dynamics $\Gamma_1(W)$ evaluated according to the relation (36).

- **Example of the D2Q9 fluid scheme**

The results presented here are essentially a reformulation of the classic work of Lallemand and Luo [43]. With the moments introduced in (22), (23) and (24), we denote by Φ_ε , Φ_{xx} and Φ_{xy} the equilibrium values of the three first non conserved moments (24). The dynamics at first order is given by (36). We have in this D2Q9 case,

$$\Gamma_1 = \begin{pmatrix} \partial_x J_x + \partial_y J_y \\ \frac{2}{3} \lambda^2 \partial_x \rho + \frac{1}{6} \partial_x \Phi_\varepsilon + \frac{1}{2} \partial_x \Phi_{xx} + \partial_y \Phi_{xy} \\ \frac{2}{3} \lambda^2 \partial_y \rho + \frac{1}{6} \partial_y \Phi_\varepsilon - \frac{1}{2} \partial_y \Phi_{xx} + \partial_x \Phi_{xy} \end{pmatrix}.$$

These relations can be easily fitted with the Euler equations of gas dynamics in two space dimensions

$$\begin{cases} \partial_t \rho + \partial_x J_x + \partial_y J_y = 0 \\ \partial_t J_x + \partial_x \left(\frac{1}{\rho} J_x^2 + p \right) + \partial_y \left(\frac{1}{\rho} J_x J_y \right) = 0 \\ \partial_t J_y + \partial_x \left(\frac{1}{\rho} J_x J_y \right) + \partial_y \left(\frac{1}{\rho} J_y^2 + p \right) = 0. \end{cases}$$

We identify the two expressions with $J_x \equiv \rho u$ and $J_y \equiv \rho v$:

$$\frac{2}{3} \lambda^2 \rho + \frac{1}{6} \Phi_\varepsilon + \frac{1}{2} \Phi_{xx} = \rho u^2 + p, \quad \Phi_{xy} = \rho u v, \quad \frac{2}{3} \lambda^2 \rho + \frac{1}{6} \Phi_\varepsilon - \frac{1}{2} \Phi_{xx} = \rho v^2 + p,$$

and we deduce the expression for the three first nonconserved moments:

$$\Phi_\varepsilon = 6p - 4\lambda^2 \rho + 3\rho(u^2 + v^2), \quad \Phi_{xx} = \rho(u^2 - v^2), \quad \Phi_{xy} = \rho u v.$$

The other nonequilibrium moments q_x , q_y and h does not play any role in the first order partial equivalent equations.

Proposition 5. Taylor expansion method at second order accuracy

An essential result for the applications is the second order asymptotic analysis for a time step Δt (or a space step Δx) tending to zero. The expansion of the microscopic variables takes the form

$$(38) \quad Y = \Phi(W) + \Delta t S^{-1} \Psi_1(W) + O(\Delta t^2)$$

and the vector $\Psi_1(W)$ satisfies the relation

$$(39) \quad \Psi_1(W) = d\Phi(W) \cdot \Gamma_1(W) - (C W + D \Phi(W)).$$

A set of second order partial differential equations emerges

$$(40) \quad \partial_t W + \Gamma_1(W) + \Delta t \Gamma_2(W) = O(\Delta t^2).$$

The vector $\Gamma_1(W)$ has been precised in (36) the vector $\Gamma_2(W)$ is obtained after two derivations of the conserved moments W and the equilibrium vector $\Phi(W)$:

$$(41) \quad \Gamma_2(W) = B \Sigma \Psi_1(W).$$

The expression of the second order term puts in evidence the importance of taking over-relaxation in the applications where Σ should also be as small as possible.

• Proof of Proposition 5.

We first consider the first order expansion of microscopic moments. From the expansion (37) of one iteration of the scheme at first order, we can extract relations for the second component. We have for the microscopic moments

$$Y + \Delta t \partial_t Y + O(\Delta t^2) = Y^* - \Delta t (C W + D Y^*) + O(\Delta t^2).$$

Then

$$Y - Y^* = -\Delta t \left(\partial_t Y + (C W + D \Phi(W)) \right) + O(\Delta t^2).$$

The explicitation of $\partial_t Y$ is given by the chain rule from the expansion (34):

$$\begin{aligned} \partial_t Y &= \partial_t (\Phi(W) + O(\Delta t)) \\ &= d\Phi(W) \cdot \partial_t W + O(\Delta t) \\ &= d\Phi(W) \cdot (-\Gamma_1) + O(\Delta t). \end{aligned}$$

Then

$$S (Y - \Phi(W)) = \Delta t (d\Phi(W) \cdot \Gamma_1 - (C W + D \Phi)) + O(\Delta t^2).$$

The relation (38) is established, and the expression of $\Psi_1(W)$ is obtained by the relation (39). Before using the expansion (38), we must expand the microscopic moments Y^* after relaxation up to first order accuracy. We have the following calculus

$$\begin{aligned} Y^* &= Y + S (\Phi(W) - Y) \\ &= Y - (\Delta t \Psi_1 + O(\Delta t^2)) && \text{due to (38)} \\ &= \Phi(W) + \left(\Sigma + \frac{1}{2} I \right) (\Delta t \Psi_1(W) + O(\Delta t^2)) - (\Delta t \Psi_1 + O(\Delta t^2)) && \text{due to (26) and (38)} \\ &= \Phi(W) + \left(\Sigma - \frac{1}{2} I \right) \Delta t \Psi_1(W) + O(\Delta t^2) \end{aligned}$$

and we have the relation

$$(42) \quad Y^* = \Phi(W) + \left(\Sigma - \frac{1}{2} I \right) \Delta t \Psi_1(W) + O(\Delta t^2).$$

The two relations (38) and (42) are nothing else than the first order terms of the general expansion (30).

- Second order partial differential equations

We expand now one iteration of the scheme (21) at second order accuracy:

$$(43) \quad m + \Delta t \partial_t m + \frac{1}{2} \Delta t^2 \partial_t^2 m + O(\Delta t^3) = m^* - \Delta t \Lambda m^* + \frac{1}{2} \Delta t^2 \Lambda^2 m^* + O(\Delta t^3).$$

We replace the vector m by its two components W and Y . We use the decomposition (8) of the momentum-velocity operator matrix Λ . We have for the first component

$$(44) \quad \begin{cases} W + \Delta t \partial_t W + \frac{1}{2} \Delta t^2 \partial_t^2 W + O(\Delta t^3) = \\ W - \Delta t (A W + B Y^*) + \frac{1}{2} \Delta t^2 (A_2 W + B_2 Y^*) + O(\Delta t^3). \end{cases}$$

We explicit some terms present in the relation (44). We have

$$\partial_t W = -\Gamma_1(W) + O(\Delta t) = -(A W + B \Phi(W)) + O(\Delta t)$$

Then, using the formal rule stated at the beginning of Section 4, we keep the order of accuracy after derivation:

$$\begin{aligned} \partial_t^2 W &= -\partial_t (\Gamma_1 + O(\Delta t)) \\ &= -d\Gamma_1 \cdot \partial_t W + O(\Delta t) \\ &= d\Gamma_1 \cdot \Gamma_1 + O(\Delta t) \\ &= A \Gamma_1 + B d\Phi \cdot \Gamma_1 + O(\Delta t). \end{aligned}$$

We deduce from (44):

$$\partial_t W = -\frac{1}{2} \Delta t \partial_t^2 W - (A W + B Y^*) + \frac{1}{2} \Delta t (A_2 W + B_2 Y^*) + O(\Delta t^2).$$

We order the various terms by powers of Δt :

$$\begin{aligned} \partial_t W &= -A W - B \left(\Phi + \left(\Sigma - \frac{1}{2} I \right) \Delta t \Psi_1 \right) - \frac{1}{2} \Delta t (A \Gamma_1 + B d\Phi \cdot \Gamma_1) \\ &\quad + \frac{1}{2} \Delta t ((A^2 + B C) W + (A B + B D) \Phi) + O(\Delta t^2) \\ &= -A W - B \Phi + \Delta t \left[-B \Sigma \Psi_1 + \frac{1}{2} B (d\Phi \cdot \Gamma_1 - C W - D \Phi) - \frac{1}{2} A (A W + B \Phi) \right. \\ &\quad \left. - \frac{1}{2} B d\Phi \cdot \Gamma_1 + \frac{1}{2} (A^2 + B C) W + \frac{1}{2} (A B + B D) \Phi \right] + O(\Delta t^2) \\ &= -\Gamma_1(W) - \Delta t B \Sigma \Psi_1(W) + O(\Delta t^2) \end{aligned}$$

and the relation (40) is proven with $\Gamma_2(W) = B \Sigma \Psi_1$ as proposed in (41). $\square$

- **Link with the original Taylor expansion method**

In [14, 15], the moments at equilibrium are denoted m_ℓ^{eq} :

$$m_\ell^{\text{eq}}(W) = \begin{cases} W_\ell & \text{if } 0 \leq \ell < N \\ (\Phi(W))_{\ell-N} & \text{if } \ell \geq N, \end{cases}$$

the coefficients $\tilde{\Lambda}_{i\ell}^\beta$ are given according to

$$(45) \quad \tilde{\Lambda}_{i\ell}^\beta = \sum_j M_{ij} v_j^\beta (M^{-1})_{j\ell}, \quad 1 \leq \beta \leq d, 0 \leq i, \ell < q,$$

are closely related to the operator matrix Λ introduced in (7). The coefficients of the Hnon's matrix (26) have a shifted numbering

$$(46) \quad \tilde{\sigma}_\ell = \sigma_{\ell-N} \equiv \frac{1}{s_{\ell-N}} - \frac{1}{2}, \quad \ell \geq N$$

and the defect of conservation $\tilde{\theta}_k$ is defined according to

$$(47) \quad \tilde{\theta}_k(W) \equiv \partial_t m_k^{\text{eq}} + \sum_{0 \leq \ell < q} \sum_{1 \leq \beta \leq d} \tilde{\Lambda}_{k\ell}^\beta \partial_\beta m_\ell^{\text{eq}}(W), \quad k \geq N.$$

Proposition 6. Equivalence at second order accuracy

The second order expansion (40), (38), (36), (39) and (41) is equivalent to the expansion proposed in [14, 15]:

$$(48) \quad \partial_t W_i + \tilde{\Lambda}_{i\ell}^\beta \partial_\beta m_\ell^{\text{eq}}(W) = \Delta t \sum_{k \geq N} \tilde{\Lambda}_{ik}^\beta \tilde{\sigma}_k \partial_\beta \tilde{\theta}_k(W) + O(\Delta t^2).$$

- Proof of Proposition 6.

Remark first that

$$\Lambda_{ik} = \sum_{j=0}^{q-1} M_{ij} \left(\sum_{1 \leq \beta \leq d} v_j^\beta \partial_\beta \right) (M^{-1})_{jk} = \sum_{1 \leq \beta \leq d} \left(\sum_{j=0}^{q-1} M_{ij} v_j^\beta (M^{-1})_{jk} \right) \partial_\beta = \sum_{1 \leq \beta \leq d} \tilde{\Lambda}_{ik}^\beta \partial_\beta.$$

Then on one hand, the component number i of the first order term $(AW + B\Phi(W))$ can be expanded in the following way:

$$\begin{aligned} (AW + B\Phi(W))_i &= \sum_{k < N} \tilde{\Lambda}_{ik}^\beta \partial_\beta W_k + \sum_{k \geq N} \tilde{\Lambda}_{ik}^\beta \partial_\beta \Phi_k \\ &= \sum_{k < N} \tilde{\Lambda}_{ik}^\beta \partial_\beta m_k^{\text{eq}} + \sum_{k \geq N} \tilde{\Lambda}_{ik}^\beta \partial_\beta m_k^{\text{eq}} \\ &= \sum_{0 \leq k < q} \tilde{\Lambda}_{ik}^\beta \partial_\beta m_k^{\text{eq}} \end{aligned}$$

and the first order terms of (40) and (48) are identical. Secondly, due to (47), we have for $k \geq N$

$$\begin{aligned} \tilde{\theta}_k &= \partial_t m_k^{\text{eq}} + \sum_{\ell \beta} \tilde{\Lambda}_{k\ell}^\beta \partial_\beta m_\ell^{\text{eq}} \\ &= \partial_t \Phi_{k-N} + (\Lambda \cdot m^{\text{eq}})_k \\ &= (d\Phi \cdot \partial_t W)_{k-N} + (\Lambda \cdot m^{\text{eq}})_k \\ &= (d\Phi \cdot (\Gamma_1 + O(\Delta t)))_{k-N} - (CW + D\Phi)_{k-N} \end{aligned}$$

and due to (39),

$$\tilde{\theta}_k = -(\Psi_1)_{k-N} + O(\Delta t), \quad k \geq N.$$

We deduce that for second order terms with $0 \leq i < N$:

$$\begin{aligned} (B \tilde{\sigma} \Psi_1)_i &= \sum_{0 \leq \ell < q-N} B_{i\ell} \tilde{\sigma}_{\ell+N} (\Psi_1(W))_\ell \\ &= \sum_{k \geq N} \sum_{0 \leq \beta \leq d} \tilde{\Lambda}_{ik}^\beta \partial_\beta \left(\sigma_k (-\tilde{\theta}_k + O(\Delta t)) \right) \end{aligned} \quad \text{with } k = \ell + N$$

$$= - \sum_{k \geq N} \sum_{0 \leq \beta \leq d} \tilde{\Lambda}_{ik}^{\beta} \tilde{\sigma}_k (\partial_{\beta} \tilde{\theta}_k) + O(\Delta t)$$

because the coefficients $\tilde{\sigma}_k$ are supposed constant. Then the relation (48) is established. $\square$

• **Fluid D2Q9 diffusive tensor**

If we compute the second order term Γ_2 for the D2Q9 scheme introduced previously, the holy grail would be to recover the viscous terms of the compressible Navier Stokes equations in two space dimensions. More precisely, with a given shear viscosity μ and a given bulk viscosity ζ , we write the viscous terms of the lattice Boltzmann expansion as

$$(49) \quad -\Delta t \Gamma_2 = \begin{pmatrix} 0 \\ \partial_j \tau_{xj} \equiv \partial_x (2\mu \partial_x u + (\zeta - \mu)(\partial_x u + \partial_y v)) + \partial_y (\mu(\partial_x v + \partial_y u)) \\ \partial_j \tau_{yj} \equiv \partial_x (\mu(\partial_x v + \partial_y u)) + \partial_y ((\zeta - \mu)(\partial_x u + \partial_y v) + 2\mu \partial_y v) \end{pmatrix} + \begin{pmatrix} 0 \\ \Psi_x \\ \Psi_y \end{pmatrix},$$

with the notation Ψ_x, Ψ_y for the error for the dissipation of momentum.

The problem to enforce $\Psi_x = \Psi_y = 0$ is ill-posed, as recognized by Dellar [10, 12], HÁZI and KÁVRÁN [34] and Prasianakis *at al.* [49] among others. Nevertheless, it can be proven that the unphysical terms involving second order space derivatives involving the density can be removed with a pressure law of the type

$$p(\rho) = \frac{\lambda^2}{3} \rho,$$

id est a sound velocity equal to $\frac{1}{\sqrt{3}}$ in the velocity unit of the lattice Boltzmann scheme. Nevertheless, with the notation Φ_{qx} and Φ_{qy} for the equilibrium functions of the moments q_x and q_y and the coefficients of the Hénon matrix Σ fixed as

$$(50) \quad \Sigma = \text{diag}(\sigma_e, \sigma_x, \sigma_x, \sigma_q, \sigma_q, \sigma_h),$$

the choice

$$\begin{cases} \Phi_{qx} = -\rho \lambda^2 u + 3\rho (u^2 + v^2) u \\ \Phi_{qy} = -\rho \lambda^2 v + 3\rho (u^2 + v^2) v, \end{cases}$$

conducts to the relation (49) with associated viscosities fixed in a very classical way:

$$(51) \quad \mu = \frac{\lambda}{3} \rho \sigma_x \Delta x, \quad \zeta = \frac{\lambda}{3} \rho \sigma_e \Delta x.$$

The equilibrium value Φ_h of the last fourth order moment h (see the relation (24)) has no incidence on the value of the diffusive term Γ_2 . The error is then reduced to third order terms relative to the velocity and we have precisely

$$\begin{cases} \Psi_x = \sigma_x \Delta t \left[\partial_x (u^3 \partial_x \rho - v^3 \partial_y \rho + 3\rho (u^2 \partial_x u - v^2 \partial_y v)) \right. \\ \quad \left. + \partial_y (-v^3 \partial_x \rho - u^3 \partial_y \rho - 3\rho (u^2 \partial_y u + v^2 \partial_x v)) \right] \\ \Psi_y = \sigma_x \Delta t \left[\partial_x (-v^3 \partial_x \rho - u^3 \partial_y \rho - 3\rho (u^2 \partial_y u + v^2 \partial_x v)) \right. \\ \quad \left. + \partial_y (-u^3 \partial_x \rho + v^3 \partial_y \rho + 3\rho (-u^2 \partial_x u + v^2 \partial_y v)) \right]. \end{cases}$$

6) Taylor expansion method at third and fourth order accuracy

The two following orders of the Taylor expansion does not set any theoretical difficulty, except the care of algebra with not so short formal expressions...

Proposition 7. Nonlinear third-order expansion

When the time step Δt has an infinitesimal value, the expansion of the microscopic variables takes the form

$$(52) \quad Y = \Phi(W) + \Delta t S^{-1} \Psi_1(W) + \Delta t^2 S^{-1} \Psi_2(W) + O(\Delta t^3).$$

The vector $\Psi_1(W)$ is evaluated at the relation (39) and $\Psi_2(W)$ satisfies the relation

$$(53) \quad \Psi_2(W) = \Sigma d\Psi_1(W). \Gamma_1(W) + d\Phi(W). \Gamma_2(W) - D \Sigma \Psi_1(W).$$

It is possible to precise a system of third order partial differential equations

$$(54) \quad \partial_t W + \Gamma_1(W) + \Delta t \Gamma_2(W) + \Delta t^2 \Gamma_3(W) = O(\Delta t^3).$$

The vectors $\Gamma_1(W)$ and $\Gamma_2(W)$ have been precised at the relations (36) and (41) respectively. We have moreover

$$(55) \quad \Gamma_3(W) = B \Sigma \Psi_2(W) + \frac{1}{12} B_2 \Psi_1(W) - \frac{1}{6} B d\Psi_1(W). \Gamma_1(W).$$

- Proof of Proposition 7.

We first establish the relations (52) and (53). We start by the expansion of the scheme (43) at order 2. For the second component of nonconserved moments, we have

$$Y + \Delta t \partial_t Y + \frac{1}{2} \Delta t^2 \partial_t^2 Y + O(\Delta t^3) = Y^* - \Delta t (C W + D Y^*) + \frac{1}{2} \Delta t^2 (C_2 W + D_2 Y^*) + O(\Delta t^3)$$

and, due to the relaxation process $Y - Y^* \equiv S(Y - \Phi(W))$, we have

$$(56) \quad S(Y - \Phi(W)) = -\Delta t (C W + D Y^* + \partial_t Y) + \frac{1}{2} \Delta t^2 (C_2 W + D_2 Y^* - \partial_t^2 Y) + O(\Delta t^3).$$

We can precise the two first time derivatives of the non-conserved moments. The relation (38) can also be written as

$$Y = \Phi(W) + \left(\Sigma + \frac{1}{2} I \right) \Delta t \Psi_1(W) + O(\Delta t^2).$$

We differentiate this previous relation relative to time:

$$\begin{aligned} \partial_t Y &= d\Phi. \partial_t W + \left(\Sigma + \frac{1}{2} I \right) \Delta t d\Psi_1. \partial_t W + O(\Delta t^2) \\ &= -d\Phi. (\Gamma_1(W) + \Delta t \Gamma_2(W) + O(\Delta t^2)) + \Delta t \left(\Sigma + \frac{1}{2} I \right) d\Psi_1. (-\Gamma_1 + O(\Delta t)) + O(\Delta t^2) \end{aligned}$$

and

$$(57) \quad \partial_t Y = -d\Phi(W). \Gamma_1(W) - \Delta t \left[d\Phi(W). \Gamma_2(W) + \left(\Sigma + \frac{1}{2} I \right) d\Psi_1(W). \Gamma_1(W) \right] + O(\Delta t^2).$$

Because we need the second time derivative $\partial_t^2 Y$ in the right hand side of (56), we differentiate the relation (57) relative to time. Using the notation (27), we deduce

$$(58) \quad \partial_t^2 Y = d\gamma_1. \Gamma_1 + O(\Delta t).$$

We precise now the expansions of $C W + D Y^*$ and $C_2 W + D_2 Y^*$. We have

$$C W + D Y^* = C W + D \left[\Phi(W) + \left(\Sigma - \frac{1}{2} I \right) \Delta t \Psi_1 \right] + O(\Delta t^2)$$

and

$$(59) \quad C W + D Y^* = \gamma_1(W) - \Psi_1(W) + \Delta t D \left(\Sigma - \frac{1}{2} I \right) \Psi_1(W) + O(\Delta t^2).$$

With a new set of operator matrices,

$$\begin{aligned}
 C_2 W + D_2 Y^* &= C_2 W + D_2 (\Phi(W) + O(\Delta t)) \\
 &= (C A + D C) W + (C B + D^2) \Phi(W) + O(\Delta t) \\
 &= C \Gamma_1 + D (d\Phi \cdot \Gamma_1 - \Psi_1) + O(\Delta t)
 \end{aligned}$$

and because $d\Phi \cdot \Gamma_1 = d\gamma_1 \cdot \Gamma_1 - (C \Gamma_1 + D d\Phi \cdot \Gamma_1)$,

$$(60) \quad C_2 W + D_2 Y^* = d(\gamma_1 - \Psi_1) \cdot \Gamma_1(W) - D \Psi_1(W) + O(\Delta t).$$

Then due to (56), (57), (58), (59) and (60), we have

$$\begin{aligned}
 S(Y - \Phi(W)) &= -\Delta t (C W + D Y^* + \partial_t Y) + \frac{1}{2} \Delta t^2 (C_2 W + D_2 Y^* - \partial_t^2 Y) + O(\Delta t^3) \\
 &= -\Delta t (\gamma_1 - \Psi_1 + \Delta t D (\Sigma - \frac{1}{2} I) \Psi_1) - \Delta t (-d\Phi \cdot \Gamma_1 - \Delta t [d\Phi \cdot \Gamma_2 + (\Sigma + \frac{1}{2} I) d\Psi_1 \cdot \Gamma_1]) \\
 &\quad + \frac{1}{2} \Delta t^2 (d(\gamma_1 - \Psi_1) \cdot \Gamma_1 - D \Psi_1) - \frac{1}{2} \Delta t^2 d\gamma_1 \cdot \Gamma_1 + O(\Delta t^3) \\
 &= \Delta t \Psi_1 + \Delta t^2 (-D \Sigma \Psi_1 + \frac{1}{2} D \Psi_1 + d\Phi \cdot \Gamma_2 + \Sigma d\Psi_1 \cdot \Gamma_1 - \frac{1}{2} D \Psi_1) + O(\Delta t^3) \\
 &= \Delta t \Psi_1 + \Delta t^2 (-D \Sigma \Psi_1 + d\Phi \cdot \Gamma_2 + \Sigma d\Psi_1 \cdot \Gamma_1) + O(\Delta t^3)
 \end{aligned}$$

and the relations (52) and (53) are established.

- Third order partial differential equations

We consider now the Taylor expansion of the relation (21) at third order:

$$(61) \quad \begin{cases} m + \Delta t \partial_t m + \frac{1}{2} \Delta t^2 \partial_t^2 m + \frac{1}{6} \Delta t^3 \partial_t^3 m + O(\Delta t^4) = \\ m^* - \Delta t \Lambda m^* + \frac{1}{2} \Delta t^2 \Lambda^2 m^* - \frac{1}{6} \Delta t^3 \Lambda^3 m^* + O(\Delta t^4). \end{cases}$$

We consider the first conserved components of the moments:

$$\begin{cases} W + \Delta t \partial_t W + \frac{1}{2} \Delta t^2 \partial_t^2 W + \frac{1}{6} \Delta t^3 \partial_t^3 W + O(\Delta t^4) = W - \Delta t (A W + B Y^*) \\ \quad + \frac{1}{2} \Delta t^2 (A_2 W + B_2 Y^*) - \frac{1}{6} \Delta t^3 (A_3 W + B_3 Y^*) + O(\Delta t^4). \end{cases}$$

We simplify the constant term W and divide by Δt . We deduce

$$(62) \quad \begin{cases} \partial_t W = -A W - B Y^* + \frac{1}{2} \Delta t (A_2 W + B_2 Y^* - \partial_t^2 W) \\ \quad - \frac{1}{6} \Delta t^2 (A_3 W + B_3 Y^* + \partial_t^3 W) + O(\Delta t^3). \end{cases}$$

We explicit the partial derivatives $\partial_t^2 W$ and $\partial_t^3 W$ at orders one and zero respectively. From the relation

$$\partial_t W + \Gamma_1 + \Delta t \Gamma_2 = O(\Delta t^2)$$

we have

$$\begin{aligned}
 \partial_t^2 W &= \partial_t (-\Gamma_1 - \Delta t \Gamma_2 + O(\Delta t^2)) \\
 &= d(-\Gamma_1 - \Delta t \Gamma_2 + O(\Delta t^2)) \cdot (\partial_t W) \\
 &= d(-\Gamma_1 - \Delta t \Gamma_2 + O(\Delta t^2)) \cdot (-\Gamma_1 - \Delta t \Gamma_2 + O(\Delta t^2))
 \end{aligned}$$

and

$$(63) \quad \partial_t^2 W = d\Gamma_1 \cdot \Gamma_1 + \Delta t (d\Gamma_1 \cdot \Gamma_2 + d\Gamma_2 \cdot \Gamma_1) + O(\Delta t^2).$$

Then

$$\begin{aligned}
 \partial_t^3 W &= \partial_t (\partial_t^2 W) \\
 &= \partial_t (d\Gamma_1 \cdot \Gamma_1) + O(\Delta t) \\
 &= d(d\Gamma_1 \cdot \Gamma_1) \cdot \partial_t W + O(\Delta t) \\
 &= -d(d\Gamma_1 \cdot \Gamma_1) \cdot \Gamma_1 + O(\Delta t)
 \end{aligned}$$

$$(64) \quad \partial_t^3 W = -\partial^2 \Gamma_1 \cdot \Gamma_1 + O(\Delta t).$$

In order to compute the coefficient Γ_3 , we precise now the expansions of the quantities $A_j W + B_j Y^*$ for $j = 1, 2$ and 3 . We have

$$\begin{aligned}
 A W + B Y^* &= A W + B (\Phi(W) + (\Sigma - \tfrac{1}{2} I) (\Delta t \Psi_1 + \Delta t^2 \Psi_2) + O(\Delta t^3)) \\
 &= A W + B \Phi + \Delta t B (\Sigma - \tfrac{1}{2} I) \Psi_1 + \Delta t^2 B (\Sigma - \tfrac{1}{2} I) \Psi_2 + O(\Delta t^3) \\
 A_2 W + B_2 Y^* &= A_2 W + B_2 (\Phi(W) + (\Sigma - \tfrac{1}{2} I) \Delta t \Psi_1) + O(\Delta t^2) \\
 &= (A^2 + B C) W + (A B + B D) \Phi + \Delta t B_2 (\Sigma - \tfrac{1}{2} I) \Psi_1 + O(\Delta t^2) \\
 &= A \Gamma_1 + B (\gamma_1 - \Psi_1) + \Delta t B_2 (\Sigma - \tfrac{1}{2} I) \Psi_1 + O(\Delta t^2) \\
 &\quad \text{because } C W + D \Psi_1 = \gamma_1 - \Psi_1 \\
 &= d\Gamma_1 \cdot \Gamma_1 - B \Psi_1 + \Delta t B_2 (\Sigma - \tfrac{1}{2} I) \Psi_1 + O(\Delta t^2) \\
 &\quad \text{because } d\Gamma_1 \cdot \Gamma_1 = A \Gamma_1 + B \gamma_1, \\
 A_3 W + B_3 Y^* &= A_3 W + B_3 \Phi(W) + O(\Delta t) \\
 &= (A_2 A + B_2 C) W + (A_2 B + B_2 D) \Phi + O(\Delta t) \\
 &= A_2 \Gamma_1 + B_2 (\gamma_1 - \Psi_1) + O(\Delta t) \\
 &= (A^2 + B C) \Gamma_1 + (A B + B D) \gamma_1 - B_2 \Psi_1 + O(\Delta t) \\
 &= A (A \Gamma_1 + B \gamma_1) + B (C \Gamma_1 + D \gamma_1) - B_2 \Psi_1 + O(\Delta t) \\
 &= A d\Gamma_1 \cdot \Gamma_1 + B (C \Gamma_1 + D \gamma_1) - B_2 \Psi_1 + O(\Delta t) \\
 &\quad \text{because } d\Gamma_1 \cdot \Gamma_1 = A \Gamma_1 + B \gamma_1 \\
 &= A d\Gamma_1 \cdot \Gamma_1 + B d(\gamma_1 - \Psi_1) \cdot \Gamma_1 - B_2 \Psi_1 + O(\Delta t) \\
 &\quad \text{because } C \Gamma_1 + D \gamma_1 = d(\gamma_1 - \Psi_1) \cdot \Gamma_1 \\
 &= \partial^2 \Gamma_1 \cdot \Gamma_1 - B d\Psi_1 \cdot \Gamma_1 - B_2 \Psi_1 + O(\Delta t) \\
 &\quad \text{because } \partial^2 \Gamma_1 \cdot \Gamma_1 \equiv d(d\Gamma_1 \cdot \Gamma_1) \cdot \Gamma_1 = A d\Gamma_1 \cdot \Gamma_1 + B d\gamma_1 \cdot \Gamma_1.
 \end{aligned}$$

We deduce now from (62), (63), (64) and the previous expressions:

$$\begin{aligned}
 \partial_t W &= -A W - B Y^* + \tfrac{1}{2} \Delta t (A_2 W + B_2 Y^* - \partial_t^2 W) - \tfrac{1}{6} \Delta t^2 (A_3 W + B_3 Y^* + \partial_t^3 W) + O(\Delta t^3) \\
 &= -(A W + B \Phi + \Delta t B (\Sigma - \tfrac{1}{2} I) \Psi_1 + \Delta t^2 B (\Sigma - \tfrac{1}{2} I) \Psi_2) \\
 &\quad + \tfrac{1}{2} \Delta t (d\Gamma_1 \cdot \Gamma_1 - B \Psi_1 + \Delta t B_2 (\Sigma - \tfrac{1}{2} I) \Psi_1 - d\Gamma_1 \cdot \Gamma_1 - \Delta t (d\Gamma_1 \cdot \Gamma_2 + d\Gamma_2 \cdot \Gamma_1)) \\
 &\quad - \tfrac{1}{6} \Delta t^2 (\partial^2 \Gamma_1 \cdot \Gamma_1 - B d\Psi_1 \cdot \Gamma_1 - B_2 \Psi_1 - \partial^2 \Gamma_1 \cdot \Gamma_1) + O(\Delta t^3) \\
 &= -\Gamma_1 - \Delta t B \Sigma \Psi_1 + \Delta t^2 (-B \Sigma \Psi_2 + \tfrac{1}{2} B \Psi_2 + \tfrac{1}{2} B_2 \Sigma \Psi_1 - \tfrac{1}{4} B_2 \Psi_1 - \tfrac{1}{2} (d\Gamma_1 \cdot \Gamma_2 + d\Gamma_2 \cdot \Gamma_1) \\
 &\quad + \tfrac{1}{6} B d\Psi_1 \cdot \Gamma_1 + \tfrac{1}{6} B_2 \Psi_1) + O(\Delta t^3) \quad \text{because } \partial^2 \Gamma_1 \cdot \Gamma_1 = A d\Gamma_1 \cdot \Gamma_1 + B d\gamma_1 \cdot \Gamma_1 \\
 &= -\Gamma_1 - \Delta t B \Sigma \Psi_1 + \Delta t^2 (-B \Sigma \Psi_2 + \tfrac{1}{2} B (\Sigma d\Psi_1 \cdot \Gamma_1 + \gamma_2 - D \Sigma \Psi_1) + \tfrac{1}{2} (A B + B D) \Sigma \Psi_1 \\
 &\quad - \tfrac{1}{12} B_2 \Psi_1 - \tfrac{1}{2} (A \Gamma_2 + B \gamma_2) - \tfrac{1}{2} B \Sigma d\Psi_1 \cdot \Gamma_1 + \tfrac{1}{6} B d\Psi_1 \cdot \Gamma_1) + O(\Delta t^3) \\
 &= -\Gamma_1 - \Delta t \Gamma_2 - \Delta t^2 (B \Sigma \Psi_2 + \tfrac{1}{12} B_2 \Psi_1 - \tfrac{1}{6} B d\Psi_1 \cdot \Gamma_1) + O(\Delta t^3)
 \end{aligned}$$

and the relations (54) and (55) are established. $\square$

In [16], we have proposed a third order expansion of nonlinear lattice Boltzmann schemes for two specific applications: scalar advection diffusion ($N = 1$) and fluid system ($N = d + 1$) with mass and momentum conservation. In the following proposition, we show that our previous result at third order accuracy can be reformulated with the compact formulation (54) (55) of Proposition 6.

Proposition 8. Third-order formal expansion for thermics and fluids

With the notations (45), (46) and (47) introduced at Proposition 5, we suppose moreover

$$M_{0j} = 1, \quad M_{\alpha j} = v_j^\alpha, \quad 0 \leq j < q, \quad 1 \leq \alpha \leq d.$$

The first moments are denoted by $\rho \equiv \sum_j f_j$ and $J_\alpha \equiv \sum_j v_j^\alpha f_j$. For the scalar advection diffusion ($N = 1$), the third order equivalent partial differential equation (54) (55) can be written

$$(65) \quad \partial_t \rho + \partial_\alpha J_\alpha^{\text{eq}} - \Delta t \tilde{\sigma}_\alpha \partial_\alpha \tilde{\theta}_\alpha + \Delta t^2 \left[\Lambda_{\beta\ell}^\gamma \left(\tilde{\sigma}_\beta \tilde{\sigma}_\ell - \frac{1}{12} \right) \partial_\beta \partial_\gamma \tilde{\theta}_\ell + \left(\tilde{\sigma}_\beta^2 - \frac{1}{6} \right) \partial_t (\partial_\beta \tilde{\theta}_\beta) \right] = O(\Delta t^3)$$

with $J_\alpha^{\text{eq}} \equiv \Phi_\alpha(\rho)$ and the defects of conservation $\tilde{\theta}_k(W)$ defined in (47). For the fluid system ($N = d + 1$), we have

$$(66) \quad \begin{cases} \partial_t \rho + \partial_\alpha J_\alpha - \frac{1}{12} \Delta t^2 \Lambda_{\beta\ell}^\gamma \partial_\beta \partial_\gamma \tilde{\theta}_\ell = O(\Delta t^3) \\ \partial_t J_\alpha + \Lambda_{\alpha k}^\beta \partial_\beta m_k^{\text{eq}} - \Delta t \Lambda_{\alpha k}^\beta \tilde{\sigma}_k \partial_\beta \tilde{\theta}_k \\ \quad + \Delta t^2 \left[\Lambda_{\alpha k}^\beta \Lambda_{k\ell}^\gamma \left(\tilde{\sigma}_k \tilde{\sigma}_\ell - \frac{1}{12} \right) \partial_\beta \partial_\gamma \tilde{\theta}_\ell + \Lambda_{\alpha k}^\beta \left(\tilde{\sigma}_k^2 - \frac{1}{6} \right) \partial_\beta \partial_t \tilde{\theta}_k \right] = O(\Delta t^3). \end{cases}$$

The relations (65) and (66) are exactly the ones proposed with numbers (35), (40) and (41) in the reference [16].

- Proof of Proposition 8.

We first consider the conservation defect $\tilde{\theta}_k(W) \equiv \partial_t m_k^{\text{eq}} + \sum_{\ell\beta} \tilde{\Lambda}_{k\ell}^\beta \partial_\beta m_\ell^{\text{eq}}$ introduced in (47). We define a vector conservation defect θ of dimension $q - N$ just obtained by a shift of component numbering:

$$\theta_k = \tilde{\theta}_{k+N}(W), \quad 0 \leq k < q - N.$$

We have for $0 \leq k < q - N$:

$$\begin{aligned} \theta_k &= (\text{d}\Phi \cdot \partial_t W)_k + (\tilde{\Lambda} \cdot m^{\text{eq}})_{k+N} \\ &= (\text{d}\Phi \cdot \partial_t W)_k + (C W + D \Phi(W))_k \\ &= (\text{d}\Phi \cdot (-\Gamma_1 - \Delta t \Gamma_2))_k + (C W + D \Phi(W))_k + O(\Delta t^2) && \text{due to the expansion (29)} \\ &= -(\Psi_1 + \Delta t \text{d}\Phi \cdot \Gamma_2)_k + O(\Delta t^2) && \text{due to the relation (31)} \end{aligned}$$

and

$$\theta = -\Psi_1 - \Delta t \text{d}\Phi \cdot \Gamma_2 + O(\Delta t^2).$$

In particular, $\partial_t \theta = -\partial_t \Psi_1 + O(\Delta t) = -\text{d}\Psi_1 \cdot (-\Gamma_1) + O(\Delta t)$ and

$$\partial_t \theta = \text{d}\Psi_1 \cdot \Gamma_1 + O(\Delta t).$$

Due to (41) and (55), we have the following calculus:

$$\begin{aligned} \Gamma_2 + \Delta t \Gamma_3 &= B \Sigma \Psi_1 + \Delta t (B \Sigma \Psi_2 - \frac{1}{12} B_2 \Psi_1 + \frac{1}{6} B \text{d}\Psi_1 \cdot \Gamma_1) \\ &= B \Sigma (-\theta + \Delta t \text{d}\Phi \cdot \Gamma_2 + O(\Delta t^2)) + \Delta t B \Sigma (D \Sigma \Psi_1 - \text{d}\Phi \cdot \Gamma_2 - \Sigma \text{d}\Psi_1 \cdot \Gamma_1) \\ &\quad - \frac{1}{12} \Delta t B_2 \Psi_1 + \frac{1}{6} \Delta t B \text{d}\Psi_1 \cdot \Gamma_1 \\ &= -B \Sigma \theta + \Delta t \left[(B \Sigma D \Sigma - \frac{1}{12} B_2) \Psi_1 + B \left(\frac{1}{6} - \Sigma^2 \right) \text{d}\Psi_1 \cdot \Gamma_1 \right] + O(\Delta t^2) \\ &= -B \Sigma \theta + \Delta t \left[- (B \Sigma D \Sigma - \frac{1}{12} B_2) \theta + B (\Sigma^2 - \frac{1}{6}) \partial_t \theta \right] + O(\Delta t^2). \end{aligned}$$

The third-order partial equivalent equations (54) can be written under the form

$$\partial_t W + \tilde{\Lambda} m^{\text{eq}} + \Delta t B \Sigma \theta + \Delta t^2 \left[\left(B \Sigma D \Sigma - \frac{1}{12} B_2 \right) \theta - B \left(\Sigma^2 - \frac{1}{6} \right) \partial_t \theta \right] = O(\Delta t^3).$$

- We explicit now the following cartesian components:

$$\begin{aligned}
 (B \Sigma D \Sigma \theta)_i &= -\tilde{\Lambda}_{ik}^\beta \tilde{\sigma}_k (-\tilde{\Lambda}_{k\ell}^\gamma \tilde{\sigma}_\ell \partial_\beta \partial_\gamma \tilde{\theta}_\ell = \tilde{\Lambda}_{ik}^\beta \tilde{\sigma}_k \tilde{\Lambda}_{k\ell}^\gamma \tilde{\sigma}_\ell \partial_\beta \partial_\gamma \tilde{\theta}_\ell, \\
 (B_2 \theta)_i &= (\tilde{\Lambda}_{ik}^\beta \partial_\beta) (\tilde{\Lambda}_{k\ell}^\gamma \partial_\gamma) \tilde{\theta}_\ell = \tilde{\Lambda}_{ik}^\beta \tilde{\Lambda}_{k\ell}^\gamma \partial_\beta \partial_\gamma \tilde{\theta}_\ell, \\
 ((B \Sigma D \Sigma - \frac{1}{12} B_2) \tilde{\theta})_i &= \tilde{\Lambda}_{ik}^\beta \tilde{\Lambda}_{k\ell}^\gamma (\tilde{\sigma}_k \tilde{\sigma}_\ell - \frac{1}{12}) \partial_\beta \partial_\gamma \tilde{\theta}_\ell \\
 \text{and } -(B (\Sigma^2 - \frac{1}{6}) \partial_t \tilde{\theta})_i &= \tilde{\Lambda}_{ik}^\beta (\tilde{\sigma}_k^2 - \frac{1}{6}) \partial_\beta \partial_t \tilde{\theta}_k.
 \end{aligned}$$

We deduce a new form of the third-order partial equivalent equations for $0 \leq i < N$:

$$(67) \quad \left\{ \begin{aligned} &\partial_t W_i + \tilde{\Lambda}_{ik}^\alpha \partial_\alpha m_k^{\text{eq}} - \Delta t \tilde{\Lambda}_{ik}^\beta \tilde{\sigma}_k \partial_\beta \tilde{\theta}_k \\ &+ \Delta t^2 [\tilde{\Lambda}_{ik}^\beta \tilde{\Lambda}_{k\ell}^\gamma (\tilde{\sigma}_k \tilde{\sigma}_\ell - \frac{1}{12}) \partial_\beta \partial_\gamma \tilde{\theta}_\ell + \tilde{\Lambda}_{ik}^\beta (\tilde{\sigma}_k^2 - \frac{1}{6}) \partial_\beta \partial_t \tilde{\theta}_k] = O(\Delta t^3). \end{aligned} \right.$$

- We focus on the mass conservation ($i = 0$). We have

$$\tilde{\Lambda}_{0\ell}^\alpha = M_{0j} v_j^\alpha (M^{-1})_{j\ell} = M_{\alpha j} (M^{-1})_{j\ell} = \delta_{\alpha\ell}.$$

Then

$$\tilde{\Lambda}_{0k}^\beta \tilde{\sigma}_k \partial_\beta \tilde{\theta}_k = \delta_{\beta k} \tilde{\sigma}_k \partial_\beta \tilde{\theta}_k = \tilde{\sigma}_\beta \partial_\beta \tilde{\theta}_\beta.$$

In the thermal case, we have only one conservation law and $\tilde{\sigma}_\beta \neq 0$. Thus we have enlightened the second order term of the left hand side of the relation (65). In the fluid case, the momentum (*id est* moments numbered from 1 to d) are conserved. In consequence, $\tilde{\sigma}_\beta = 0$ and we have no second order term in the mass conservation (66).

Now for the first term relative to third order term of (67), we have

$$\tilde{\Lambda}_{0k}^\beta \tilde{\Lambda}_{k\ell}^\gamma (\tilde{\sigma}_k \tilde{\sigma}_\ell - \frac{1}{12}) \partial_\beta \partial_\gamma \tilde{\theta}_\ell = \delta_{\beta k} \tilde{\Lambda}_{k\ell}^\gamma (\tilde{\sigma}_k \tilde{\sigma}_\ell - \frac{1}{12}) \partial_\beta \partial_\gamma \tilde{\theta}_\ell = \tilde{\Lambda}_{\beta\ell}^\gamma (\tilde{\sigma}_\beta \tilde{\sigma}_\ell - \frac{1}{12}) \partial_\beta \partial_\gamma \tilde{\theta}_\ell.$$

In the thermal case, $\tilde{\sigma}_\beta \neq 0$ and this term is exactly the first of the two third order terms of the relation (65). In the fluid case, $\tilde{\sigma}_\beta = 0$ and this term has a more compact expression.

Finally,

$$(68) \quad \tilde{\Lambda}_{0k}^\beta \tilde{\Lambda}_{k\ell}^\gamma (\tilde{\sigma}_k \tilde{\sigma}_\ell - \frac{1}{12}) \partial_\beta \partial_\gamma \tilde{\theta}_\ell = \begin{cases} \tilde{\Lambda}_{\beta\ell}^\gamma (\tilde{\sigma}_\beta \tilde{\sigma}_\ell - \frac{1}{12}) \partial_\beta \partial_\gamma \tilde{\theta}_\ell & \text{thermics} \\ -\frac{1}{12} \tilde{\Lambda}_{\beta\ell}^\gamma \partial_\beta \partial_\gamma \tilde{\theta}_\ell & \text{fluid.} \end{cases}$$

For the second term relative to third order, we have

$$\tilde{\Lambda}_{0k}^\beta (\tilde{\sigma}_k^2 - \frac{1}{6}) \partial_\beta \partial_t \tilde{\theta}_k = (\tilde{\sigma}_\beta^2 - \frac{1}{6}) \partial_t (\partial_\beta \tilde{\theta}_\beta)$$

In the thermal case, $\tilde{\sigma}_\beta \neq 0$ and $\tilde{\Lambda}_{0k}^\beta (\tilde{\sigma}_k^2 - \frac{1}{6}) \partial_\beta \partial_t \tilde{\theta}_k = (\tilde{\sigma}_\beta^2 - \frac{1}{6}) \partial_t (\partial_\beta \tilde{\theta}_\beta)$. In the fluid case, the momentum is conserved and $\tilde{\theta}_\beta = O(\Delta t)$ and $\tilde{\Lambda}_{0k}^\beta (\tilde{\sigma}_k^2 - \frac{1}{6}) \partial_\beta \partial_t \tilde{\theta}_k = O(\Delta t)$. In a synthetic way,

$$(69) \quad \tilde{\Lambda}_{0k}^\beta (\tilde{\sigma}_k^2 - \frac{1}{6}) \partial_\beta \partial_t \tilde{\theta}_k = \begin{cases} (\tilde{\sigma}_\beta^2 - \frac{1}{6}) \partial_t (\partial_\beta \tilde{\theta}_\beta) & \text{thermics} \\ O(\Delta t) & \text{fluid.} \end{cases}$$

We deduce from (68) and (69) the expansion (65) in the thermal case and the first relation of (66) in the fluid case.

- For the momentum equation of the fluid case, $1 \leq i = \alpha \leq d$ and the relation (67) can be rewritten as

$$\left\{ \begin{aligned} &\partial_t J_\alpha + \tilde{\Lambda}_{\alpha k}^\alpha \partial_\alpha m_k^{\text{eq}} - \Delta t \tilde{\Lambda}_{\alpha k}^\beta \tilde{\sigma}_k \partial_\beta \tilde{\theta}_k \\ &+ \Delta t^2 [\tilde{\Lambda}_{\alpha k}^\beta \tilde{\Lambda}_{k\ell}^\gamma (\tilde{\sigma}_k \tilde{\sigma}_\ell - \frac{1}{12}) \partial_\beta \partial_\gamma \tilde{\theta}_\ell + \tilde{\Lambda}_{\alpha k}^\beta (\tilde{\sigma}_k^2 - \frac{1}{6}) \partial_\beta \partial_t \tilde{\theta}_k] = O(\Delta t^3). \end{aligned} \right.$$

This relation is exactly the second relation of (66). $\square$

Proposition 9. Nonlinear expansion at fourth order

When the time step Δt has an infinitesimal value, the expansion of the microscopic variables takes the form

$$(70) \quad Y = \Phi(W) + \Delta t S^{-1} \Psi_1(W) + \Delta t^2 S^{-1} \Psi_2(W) + \Delta t^3 S^{-1} \Psi_3(W) + O(\Delta t^4).$$

The vectors $\Psi_1(W)$ and $\Psi_2(W)$ have been evaluated at the relations (39) and (53). We have

$$(71) \quad \begin{cases} \Psi_3(W) = d\Phi(W). \Gamma_3(W) + \Sigma d\Psi_1(W). \Gamma_2(W) + \Sigma d\Psi_2(W). \Gamma_1(W) - D \Sigma \Psi_2(W) \\ \quad + \frac{1}{6} D d\Psi_1(W). \Gamma_1(W) - \frac{1}{12} D_2 \Psi_1(W) - \frac{1}{12} \partial^2 \Psi_1(W). \Gamma_1(W). \end{cases}$$

A system of fourth order partial differential equations is asymptotically satisfied by the lattice Boltzmann scheme:

$$(72) \quad \partial_t W + \Gamma_1(W) + \Delta t \Gamma_2(W) + \Delta t^2 \Gamma_3(W) + \Delta t^3 \Gamma_4(W) = O(\Delta t^4).$$

The vectors $\Gamma_1(W)$, $\Gamma_2(W)$ and $\Gamma_3(W)$ have been precised at the relations (36), (41) and (55) respectively. We have now

$$(73) \quad \begin{cases} \Gamma_4(W) = B \Sigma \Psi_3(W) + \frac{1}{4} B_2 \Psi_2(W) + \frac{1}{6} B D_2 \Sigma \Psi_1 - \frac{1}{6} A B \Psi_2(W) \\ \quad - \frac{1}{6} B d\gamma_1(W). \Gamma_2(W) - \frac{1}{6} B d\gamma_2(W). \Gamma_1(W) - \frac{1}{6} B \Sigma \partial^2 \Psi_1(W). \Gamma_1(W). \end{cases}$$

- Proof of Proposition 9.

We first establish the relations (70) and (71). We start by the expansion of the scheme (61) at order 3. For the second component of nonconserved moments, we have

$$\begin{cases} Y + \Delta t \partial_t Y + \frac{1}{2} \Delta t^2 \partial_t^2 Y + \frac{1}{6} \Delta t^3 \partial_t^3 Y + O(\Delta t^4) = Y^* - \Delta t (C W + D Y^*) \\ \quad + \frac{1}{2} \Delta t^2 (C_2 W + D_2 Y^*) - \frac{1}{6} \Delta t^3 (C_3 W + D_3 Y^*) + O(\Delta t^4). \end{cases}$$

The relaxation process can be written $Y - Y^* \equiv S(Y - \Phi(W))$. We deduce

$$(74) \quad \begin{cases} S(Y - \Phi(W)) = -\Delta t (C W + D Y^* + \partial_t Y) + \frac{1}{2} \Delta t^2 (C_2 W + D_2 Y^* - \partial_t^2 Y) \\ \quad - \frac{1}{6} \Delta t^3 (C_3 W + D_3 Y^* + \partial_t^3 Y) + O(\Delta t^4). \end{cases}$$

We have to precise the three first time derivatives of the non-conserved moments. The relation (70) can also be written at second order accuracy:

$$Y = \Phi(W) + \left(\Sigma + \frac{1}{2} I \right) \Delta t \Psi_1(W) + \left(\Sigma + \frac{1}{2} I \right) \Delta t^2 \Psi_2(W) + O(\Delta t^3).$$

We differentiate this previous relation relative to time:

$$\begin{aligned} \partial_t Y &= d\Phi. \partial_t W + \left(\Sigma + \frac{1}{2} I \right) \Delta t d\Psi_1. \partial_t W + \left(\Sigma + \frac{1}{2} I \right) \Delta t^2 d\Psi_2. \partial_t W + O(\Delta t^3) \\ &= -d\Phi. (\Gamma_1(W) + \Delta t \Gamma_2(W) + \Delta t^2 \Gamma_3(W) + O(\Delta t^3)) \\ &\quad - \Delta t \left(\Sigma + \frac{1}{2} I \right) d\Psi_1. (\Gamma_1 + \Delta t \Gamma_2 + O(\Delta t^2)) + \Delta t^2 \left(\Sigma + \frac{1}{2} I \right) d\Psi_2. (-\Gamma_1 + O(\Delta t)) + O(\Delta t^3) \end{aligned}$$

and

$$(75) \quad \begin{cases} \partial_t Y = -d\Phi(W). \Gamma_1(W) - \Delta t [d\Phi(W). \Gamma_2(W) + \left(\Sigma + \frac{1}{2} I \right) d\Psi_1(W). \Gamma_1(W)] \\ \quad - \Delta t^2 [d\Phi(W). \Gamma_3(W) + \left(\Sigma + \frac{1}{2} I \right) d\Psi_1(W). \Gamma_2(W) \\ \quad + \left(\Sigma + \frac{1}{2} I \right) d\Psi_2(W). \Gamma_1(W)] + O(\Delta t^3). \end{cases}$$

We need the second time derivative $\partial_t^2 Y$ and third order time derivative $\partial_t^3 Y$ in the right hand side of (74). We differentiate the relation (75) relative to time. Using the notation (27), we deduce

$$\partial_t^2 Y = -d\gamma_1.(-\Gamma_1 - \Delta t \Gamma_2) - \Delta t \left[-d\gamma_2.\Gamma_1 - \left(\Sigma + \frac{1}{2} I\right) \partial^2 \Phi(W).\Gamma_1(W) \right] + O(\Delta t^2)$$

and

$$(76) \quad \partial_t^2 Y = d\gamma_1.\Gamma_1 + \Delta t \left[d\gamma_1.\Gamma_2 + d\gamma_2.\Gamma_1 + \left(\Sigma + \frac{1}{2} I\right) \partial^2 \Psi_1(W).\Gamma_1(W) \right] + O(\Delta t^2).$$

Finally,

$$(77) \quad \partial_t^3 Y = -\partial^2 \gamma_1(W).\Gamma_1(W) + O(\Delta t).$$

We precise now the expansions of $C W + D Y^*$, $C_2 W + D_2 Y^*$ and $C_3 W + D_3 Y^*$. We have

$$C W + D Y^* = C W + D \left[\Phi(W) + \left(\Sigma - \frac{1}{2} I\right) \Delta t \Psi_1 + \left(\Sigma - \frac{1}{2} I\right) \Delta t^2 \Psi_2 \right] + O(\Delta t^2)$$

and

$$(78) \quad \begin{cases} C W + D Y^* = \gamma_1(W) - \Psi_1(W) + \Delta t D \left(\Sigma - \frac{1}{2} I\right) \Psi_1(W) \\ \quad \quad \quad + \Delta t^2 D \left(\Sigma - \frac{1}{2} I\right) \Psi_2(W) + O(\Delta t^3). \end{cases}$$

For the second order term,

$$C_2 W + D_2 Y^* = C_2 W + D_2 \left[\Phi(W) + \Delta t \left(\Sigma - \frac{1}{2} I\right) \Psi_1 \right] + O(\Delta t^2)$$

and due to (60),

$$(79) \quad C_2 W + D_2 Y^* = d(\gamma_1 - \Psi_1).\Gamma_1(W) - D \Psi_1(W) + \Delta t D_2 \left(\Sigma - \frac{1}{2} I\right) \Psi_1 + O(\Delta t^2).$$

We have finally for the third order terms

$$\begin{aligned} C_3 W + D_3 Y^* &= C_3 W + D_3 \Phi(W) + O(\Delta t) \\ &= (C_2 A + D_2 C) W + (C_2 B + D_2 D) \Phi(W) + O(\Delta t) \\ &= C_2 \Gamma_1 + D_2 (\gamma_1 - \Psi_1) + O(\Delta t) \\ &= (C A + D C) \Gamma_1 + (C B + D^2) \gamma_1 - D_2 \Psi_1 + O(\Delta t) \\ &= C d\gamma_1.\Gamma_1 + D d(\gamma_1 - \Psi_1).\Gamma_1 - D_2 \Psi_1 + O(\Delta t) \\ &= C d\gamma_1.\Gamma_1 + D d\gamma_1.\Gamma_1 - D d\Psi_1.\Gamma_1 - D_2 \Psi_1 + O(\Delta t) \end{aligned}$$

and

$$(80) \quad C_3 W + D_3 Y^* = \partial^2(\gamma_1 - \Psi_1).\Gamma_1 - D d\Psi_1.\Gamma_1 - D_2 \Psi_1 + O(\Delta t)$$

Then due to (74), (75), (76), (77), (78), (79) and (80), we have

$$\begin{aligned} S(Y - \Phi(W)) &= -\Delta t (C W + D Y^* + \partial_t Y) + \frac{1}{2} \Delta t^2 (C_2 W + D_2 Y^* - \partial_t^2 Y) \\ &\quad - \frac{1}{6} \Delta t^3 (C_3 W + D_3 Y^* + \partial_t^3 Y) + O(\Delta t^4) \\ &= -\Delta t \left(\gamma_1 - \Psi_1 + \Delta t D \left(\Sigma - \frac{1}{2} I\right) \Psi_1 + \Delta t^2 D \left(\Sigma - \frac{1}{2} I\right) \Psi_2 \right) \\ &\quad - \Delta t \left[-d\Phi.\Gamma_1 - \Delta t (d\Phi.\Gamma_2 + \left(\Sigma + \frac{1}{2} I\right) d\Psi_1.\Gamma_1) \right. \\ &\quad \quad \left. - \Delta t^2 (d\Phi.\Gamma_3 + \left(\Sigma + \frac{1}{2} I\right) d\Psi_1.\Gamma_2 + \left(\Sigma + \frac{1}{2} I\right) d\Psi_2.\Gamma_1) \right] \\ &\quad + \frac{1}{2} \Delta t^2 (d(\gamma_1 - \Psi_1).\Gamma_1 - D \Psi_1 + \Delta t D_2 \left(\Sigma - \frac{1}{2} I\right) \Psi_1) \\ &\quad - \frac{1}{2} \Delta t^2 (d\gamma_1.\Gamma_1 + \Delta t [d\gamma_1.\Gamma_2 + d\gamma_2.\Gamma_1 + \left(\Sigma + \frac{1}{2} I\right) \partial^2 \Psi_1.\Gamma_1]) \\ &\quad - \frac{1}{6} \Delta t^3 (\partial^2(\gamma_1 - \Psi_1).\Gamma_1 - D d\Psi_1.\Gamma_1 - D_2 \Psi_1) - \frac{1}{6} \Delta t^3 (-\partial^2 \gamma_1.\Gamma_1) + O(\Delta t^4) \end{aligned}$$

$$\begin{aligned}
 &= \Delta t \Psi_1 + \Delta t^2 \Psi_2 + \Delta t^3 \left[-D(\Sigma - \tfrac{1}{2}I) \Psi_2 + d\Phi \cdot \Gamma_3 + (\Sigma + \tfrac{1}{2}I) d\Psi_1 \cdot \Gamma_2 + (\Sigma + \tfrac{1}{2}I) d\Psi_2 \cdot \Gamma_1 \right. \\
 &\quad \left. + \tfrac{1}{2} D_2 (\Sigma - \tfrac{1}{2}I) \Psi_1 - \tfrac{1}{2} d\gamma_1 \cdot \Gamma_2 - \tfrac{1}{2} d\gamma_2 \cdot \Gamma_1 - \tfrac{1}{2} (\Sigma + \tfrac{1}{2}I) \partial^2 \Psi_1 \cdot \Gamma_1 + \tfrac{1}{6} \partial^2 \Psi_1 \cdot \Gamma_1 + \tfrac{1}{6} D d\Psi_1 \cdot \Gamma_1 \right. \\
 &\quad \left. + \tfrac{1}{6} D_2 \Psi_1 \right] + O(\Delta t^4) \\
 &= \Delta t \Psi_1 + \Delta t^2 \Psi_2 + \Delta t^3 \left[-D \Sigma \Psi_2 + \tfrac{1}{2} D (\Sigma d\Psi_1 \cdot \Gamma_1 + \gamma_2 - D \Sigma \Psi_1) + d\Phi \cdot \Gamma_3 + \Sigma d\Psi_1 \cdot \Gamma_2 \right. \\
 &\quad \left. + \tfrac{1}{2} d\Psi_1 \cdot \Gamma_2 + \Sigma d\Psi_2 \cdot \Gamma_1 + \tfrac{1}{2} (\Sigma \partial^2 \Psi_1 \cdot \Gamma_1 + d\gamma_2 \cdot \Gamma_1 - D \Sigma d\Psi_1 \cdot \Gamma_1) + \tfrac{1}{2} (C B + D^2) \Sigma \Psi_1 \right. \\
 &\quad \left. - \tfrac{1}{4} D_2 \Psi_1 - \tfrac{1}{2} d\gamma_1 \cdot \Gamma_2 - \tfrac{1}{2} d\gamma_2 \cdot \Gamma_1 - \tfrac{1}{2} \Sigma \partial^2 \Psi_1 \cdot \Gamma_1 - \tfrac{1}{12} \partial^2 \Psi_1 \cdot \Gamma_1 + \tfrac{1}{6} D d\Psi_1 \cdot \Gamma_1 + \tfrac{1}{6} D_2 \Psi_1 \right] \\
 &\quad + O(\Delta t^4) \\
 &= \Delta t \Psi_1 + \Delta t^2 \Psi_2 + \Delta t^3 \left[d\Phi \cdot \Gamma_3 - D \Sigma \Psi_2 + \tfrac{1}{2} D \gamma_2 + \Sigma d\Psi_1 \cdot \Gamma_2 + \tfrac{1}{2} d\Psi_1 \cdot \Gamma_2 + \Sigma d\Psi_2 \cdot \Gamma_1 \right. \\
 &\quad \left. + \tfrac{1}{2} C B \Sigma \Psi_1 - \tfrac{1}{12} D_2 \Psi_1 - \tfrac{1}{2} d\gamma_1 \cdot \Gamma_2 - \tfrac{1}{12} \partial^2 \Psi_1 \cdot \Gamma_1 + \tfrac{1}{6} D d\Psi_1 \cdot \Gamma_1 \right] + O(\Delta t^4) \\
 &= \Delta t \Psi_1 + \Delta t^2 \Psi_2 + \Delta t^3 \left[d\Phi \cdot \Gamma_3 - D \Sigma \Psi_2 + \tfrac{1}{2} D \gamma_2 + \Sigma d\Psi_1 \cdot \Gamma_2 + \tfrac{1}{2} (-C \Gamma_2 - D \gamma_2 + d\gamma_1 \cdot \Gamma_2) \right. \\
 &\quad \left. + \Sigma d\Psi_2 \cdot \Gamma_1 + \tfrac{1}{2} C B \Sigma \Psi_1 - \tfrac{1}{12} D_2 \Psi_1 - \tfrac{1}{2} d\gamma_1 \cdot \Gamma_2 - \tfrac{1}{12} \partial^2 \Psi_1 \cdot \Gamma_1 + \tfrac{1}{6} D d\Psi_1 \cdot \Gamma_1 \right] + O(\Delta t^4) \\
 &= \Delta t \Psi_1 + \Delta t^2 \Psi_2 + \Delta t^3 \left[d\Phi \cdot \Gamma_3 - D \Sigma \Psi_2 + \Sigma d\Psi_1 \cdot \Gamma_2 + \Sigma d\Psi_2 \cdot \Gamma_1 - \tfrac{1}{12} D_2 \Psi_1 - \tfrac{1}{12} \partial^2 \Psi_1 \cdot \Gamma_1 \right. \\
 &\quad \left. + \tfrac{1}{6} D d\Psi_1 \cdot \Gamma_1 \right] + O(\Delta t^4).
 \end{aligned}$$

The relations (70) and (71) are established.

- Fourth order partial differential equations

We consider now the Taylor expansion of the relation (21) at fourth order:

$$\begin{aligned}
 m + \Delta t \partial_t m + \tfrac{1}{2} \Delta t^2 \partial_t^2 m + \tfrac{1}{6} \Delta t^3 \partial_t^3 m + \tfrac{1}{24} \Delta t^4 \partial_t^4 m + O(\Delta t^5) \\
 = m^* - \Delta t \Lambda m^* + \tfrac{1}{2} \Delta t^2 \Lambda^2 m^* - \tfrac{1}{6} \Delta t^3 \Lambda^3 m^* + \tfrac{1}{24} \Delta t^4 \Lambda^4 m^* + O(\Delta t^5).
 \end{aligned}$$

For the first components (conserved moments):

$$\begin{aligned}
 W + \Delta t \partial_t W + \tfrac{1}{2} \Delta t^2 \partial_t^2 W + \tfrac{1}{6} \Delta t^3 \partial_t^3 W + \tfrac{1}{24} \Delta t^4 \partial_t^4 W + O(\Delta t^5) = W - \Delta t (A W + B Y^*) \\
 + \tfrac{1}{2} \Delta t^2 (A_2 W + B_2 Y^*) - \tfrac{1}{6} \Delta t^3 (A_3 W + B_3 Y^*) + \tfrac{1}{24} \Delta t^4 (A_4 W + B_4 Y^*) + O(\Delta t^5).
 \end{aligned}$$

We simplify the constant term W and divide by Δt . We deduce

$$(81) \quad \begin{cases} \partial_t W = -(A W + B Y^*) + \tfrac{1}{2} \Delta t (A_2 W + B_2 Y^* - \partial_t^2 W) \\ \quad - \tfrac{1}{6} \Delta t^2 (A_3 W + B_3 Y^* + \partial_t^3 W) + \tfrac{1}{24} \Delta t^3 (A_4 W + B_4 Y^* - \partial_t^4 W) + O(\Delta t^4). \end{cases}$$

We explicit the partial derivatives $\partial_t^2 W$, $\partial_t^3 W$ and $\partial_t^4 W$ at orders two, one and zero respectively. We start with the relation

$$\partial_t W + \Gamma_1 + \Delta t \Gamma_2 + \Delta t^2 \Gamma_3 = O(\Delta t^3)$$

where Γ_1 , Γ_2 and Γ_3 have been evaluated previously. We have

$$\begin{aligned}
 \partial_t^2 W &= \partial_t (-\Gamma_1 - \Delta t \Gamma_2 - \Delta t^2 \Gamma_3 + O(\Delta t^3)) \\
 &= d(-\Gamma_1 - \Delta t \Gamma_2 - \Delta t^2 \Gamma_3 + O(\Delta t^3)) \cdot (\partial_t W) \\
 &= d(-\Gamma_1 - \Delta t \Gamma_2 - \Delta t^2 \Gamma_3 + O(\Delta t^2)) \cdot (-\Gamma_1 - \Delta t \Gamma_2 - \Delta t^2 \Gamma_3) + O(\Delta t^3)
 \end{aligned}$$

and

$$(82) \quad \partial_t^2 W = d\Gamma_1 \cdot \Gamma_1 + \Delta t (d\Gamma_1 \cdot \Gamma_2 + d\Gamma_2 \cdot \Gamma_1) + \Delta t^2 (d\Gamma_1 \cdot \Gamma_3 + d\Gamma_2 \cdot \Gamma_2 + d\Gamma_3 \cdot \Gamma_1) + O(\Delta t^2).$$

Then

$$\begin{aligned}
 \partial_t^3 W &= \partial_t (\partial_t^2 W) \\
 &= \partial_t (A \Gamma_1 + B d\Phi \cdot \Gamma_1 + \Delta t \partial_t (d\Gamma_2 \cdot \Gamma_1 + d\Gamma_1 \cdot \Gamma_2)) + O(\Delta t^2)
 \end{aligned}$$

$$\begin{aligned}
 &= (A d\Gamma_1 + B d(d\Phi.\Gamma_1)).(-\Gamma_1 - \Delta t \Gamma_2 + O(\Delta t^2)) \\
 &\quad + \Delta t \partial_t (B \Sigma d\Psi_1.\Gamma_1 + A \Gamma_2 + B d\Phi.\Gamma_2) + O(\Delta t^2) \\
 &= \partial_t (d\Gamma_1.\Gamma_1 + \Delta t (d\Gamma_1.\Gamma_2 + d\Gamma_2.\Gamma_1)) + O(\Delta t^2) \\
 &= d(d\Gamma_1.\Gamma_1 + \Delta t (d\Gamma_1.\Gamma_2 + d\Gamma_2.\Gamma_1)).\partial_t W + O(\Delta t^2) \\
 &= d(d\Gamma_1.\Gamma_1 + \Delta t (d\Gamma_1.\Gamma_2 + d\Gamma_2.\Gamma_1)).(-\Gamma_1 - \Delta t \Gamma_2) + O(\Delta t^2) \\
 &= -d(d\Gamma_1.\Gamma_1).\Gamma_1 - \Delta t [d(A \Gamma_2 + B \gamma_2 + B \Sigma d\Psi_1.\Gamma_1).\Gamma_1 + d(A \Gamma_1 + B \gamma_1).\Gamma_2] + O(\Delta t^2) \\
 &= -\partial^2 \Gamma_1.\Gamma_1 - \Delta t [A d\Gamma_2.\Gamma_1 + B d\gamma_2.\Gamma_1 + B \Sigma \partial^2 \Psi_1.\Gamma_1 + A d\Gamma_1.\Gamma_2 + B d\gamma_1.\Gamma_2] + O(\Delta t^2)
 \end{aligned}$$

and we have at first order for this third order derivative

$$(83) \quad \begin{cases} \partial_t^3 W = -\partial^2 \Gamma_1.\Gamma_1 - \Delta t [A (d\Gamma_2.\Gamma_1 + d\Gamma_1.\Gamma_2) \\ \quad + B (d\gamma_2.\Gamma_1 + d\gamma_1.\Gamma_2 + \Sigma \partial^2 \Psi_1.\Gamma_1)] + O(\Delta t^2). \end{cases}$$

Last but not least, we have

$$\begin{aligned}
 \partial_t^4 W &= \partial_t (-\partial^2 \Gamma_1.\Gamma_1 + O(\Delta t)) \\
 &= -\partial_t (A d\Gamma_1.\Gamma_1 + B d\gamma_1.\Gamma_1) + O(\Delta t) \\
 &= d(A d\Gamma_1.\Gamma_1 + B d\gamma_1.\Gamma_1).(\Gamma_1 + O(\Delta t))
 \end{aligned}$$

and finally

$$(84) \quad \partial_t^4 W = A \partial^2 \Gamma_1.\Gamma_1 + B \partial^2 \gamma_1.\Gamma_1 + O(\Delta t).$$

For the determination of the coefficient Γ_4 , we precise the expansions of the quantities $A_j W + B_j Y^*$ for $j = 1$ to $j = 4$. We use the previous evaluations done in the proof of the previous propositions:

$$\begin{aligned}
 A W + B Y^* &= A W + B [\Phi(W) + (\Sigma - \tfrac{1}{2} I) (\Delta t \Psi_1 + \Delta t^2 \Psi_2 + \Delta t^3 \Psi_3)] + O(\Delta t^4) \\
 &= A W + B \Phi + \Delta t B (\Sigma - \tfrac{1}{2} I) \Psi_1 + \Delta t^2 B (\Sigma - \tfrac{1}{2} I) \Psi_2 + \Delta t^3 B (\Sigma - \tfrac{1}{2} I) \Psi_3 + O(\Delta t^4) \\
 A_2 W + B_2 Y^* &= A_2 W + B_2 [\Phi(W) + (\Sigma - \tfrac{1}{2} I) (\Delta t \Psi_1 + \Delta t^2 \Psi_2)] + O(\Delta t^3) \\
 &= (A^2 + B C) W + (A B + B D) \Phi + \Delta t B_2 (\Sigma - \tfrac{1}{2} I) \Psi_1 + \Delta t^2 B_2 (\Sigma - \tfrac{1}{2} I) \Psi_2 + O(\Delta t^3) \\
 &= d\Gamma_1.\Gamma_1 - B \Psi_1 + \Delta t B_2 (\Sigma - \tfrac{1}{2} I) \Psi_1 + \Delta t^2 B_2 (\Sigma - \tfrac{1}{2} I) \Psi_2 + O(\Delta t^3) \\
 A_3 W + B_3 Y^* &= A_3 W + B_3 [\Phi(W) + \Delta t (\Sigma - \tfrac{1}{2} I) \Psi_1] + O(\Delta t^2) \\
 &= (A_2 A + B_2 C) W + (A_2 B + B_2 D) \Phi + \Delta t B_3 (\Sigma - \tfrac{1}{2} I) \Psi_1 + O(\Delta t^2) \\
 &= \partial^2 \Gamma_1.\Gamma_1 - B d\Psi_1.\Gamma_1 - B_2 \Psi_1 + \Delta t B_3 (\Sigma - \tfrac{1}{2} I) \Psi_1 + O(\Delta t^2) \\
 A_4 W + B_4 Y^* &= A_4 W + B_4 \Phi(W) + O(\Delta t) \\
 &= (A_3 A + B_3 C) W + (A_3 B + B_3 D) \Phi + O(\Delta t) \\
 &= A_3 \Gamma_1 + B_3 (\gamma_1 - \Psi_1) + O(\Delta t) \\
 &= (A_2 A + B_2 C) \Gamma_1 + (A_2 B + B_2 D) \gamma_1 - B_3 \Psi_1 + O(\Delta t) \\
 &= A_2 d\Gamma_1.\Gamma_1 + B_2 d(\gamma_1 - \Psi_1).\Gamma_1 - B_3 \Psi_1 + O(\Delta t) \\
 &= (A^2 + B C) d\Gamma_1.\Gamma_1 + (A B + B D) d\gamma_1.\Gamma_1 - B_2 d\Psi_1.\Gamma_1 - B_3 \Psi_1 + O(\Delta t) \\
 &= A \partial^2 \Gamma_1.\Gamma_1 + B \partial^2 (\gamma_1 - \Psi_1).\Gamma_1 - B_2 d\Psi_1.\Gamma_1 - B_3 \Psi_1 + O(\Delta t) \\
 &= A \partial^2 \Gamma_1.\Gamma_1 + B \partial^2 \gamma_1.\Gamma_1 - B \partial^2 \Psi_1.\Gamma_1 - B_2 d\Psi_1.\Gamma_1 - B_3 \Psi_1 + O(\Delta t).
 \end{aligned}$$

We deduce now from (81), (82), (83), (84) and the previous expressions:

$$\begin{aligned}
 \partial_t W &= -(A W + B Y^*) + \frac{1}{2} \Delta t (A_2 W + B_2 Y^* - \partial_t^2 W) - \frac{1}{6} \Delta t^2 (A_3 W + B_3 Y^* + \partial_t^3 W) \\
 &\quad + \frac{1}{24} \Delta t^3 (A_4 W + B_4 Y^* - \partial_t^4 W) + O(\Delta t^4) \\
 &= -(A W + B \Phi + \Delta t B (\Sigma - \frac{1}{2} I) \Psi_1 + \Delta t^2 B (\Sigma - \frac{1}{2} I) \Psi_2 + \Delta t^3 B (\Sigma - \frac{1}{2} I) \Psi_3) \\
 &\quad + \frac{1}{2} \Delta t [d\Gamma_1 \cdot \Gamma_1 - B \Psi_1 + \Delta t B_2 (\Sigma - \frac{1}{2} I) \Psi_1 + \Delta t^2 B_2 (\Sigma - \frac{1}{2} I) \Psi_2 \\
 &\quad - (d\Gamma_1 \cdot \Gamma_1 + \Delta t (d\Gamma_1 \cdot \Gamma_2 + d\Gamma_2 \cdot \Gamma_1) + \Delta t^2 (d\Gamma_1 \cdot \Gamma_3 + d\Gamma_2 \cdot \Gamma_2 + d\Gamma_3 \cdot \Gamma_1))] \\
 &\quad - \frac{1}{6} \Delta t^2 [\partial^2 \Gamma_1 \cdot \Gamma_1 - B d\Psi_1 \cdot \Gamma_1 - B_2 \Psi_1 + \Delta t B_3 (\Sigma - \frac{1}{2} I) \Psi_1 \\
 &\quad - \partial^2 \Gamma_1 \cdot \Gamma_1 - \Delta t (A (d\Gamma_2 \cdot \Gamma_1 + d\Gamma_1 \cdot \Gamma_2) + B (d\gamma_2 \cdot \Gamma_1 + d\gamma_1 \cdot \Gamma_2 + \Sigma \partial^2 \Psi_1 \cdot \Gamma_1))] \\
 &\quad + \frac{1}{24} \Delta t^3 [A \partial^2 \Gamma_1 \cdot \Gamma_1 + B \partial^2 \gamma_1 \cdot \Gamma_1 - B \partial^2 \Psi_1 \cdot \Gamma_1 - B_2 d\Psi_1 \cdot \Gamma_1 - B_3 \Psi_1 \\
 &\quad - (A \partial^2 \Gamma_1 \cdot \Gamma_1 + B \partial^2 \gamma_1 \cdot \Gamma_1)] + O(\Delta t^4) \\
 &= -\Gamma_1 - \Delta t \Gamma_2 - \Delta t^2 \Gamma_3 + \Delta t^3 [-B (\Sigma - \frac{1}{2} I) \Psi_3 + \frac{1}{2} B_2 (\Sigma - \frac{1}{2} I) \Psi_2 \\
 &\quad - \frac{1}{2} (d\Gamma_1 \cdot \Gamma_3 + d\Gamma_2 \cdot \Gamma_2 + d\Gamma_3 \cdot \Gamma_1) + \frac{1}{6} (-B_3 (\Sigma - \frac{1}{2} I) \Psi_1 + A (d\Gamma_2 \cdot \Gamma_1 + d\Gamma_1 \cdot \Gamma_2) \\
 &\quad + B (d\gamma_2 \cdot \Gamma_1 + d\gamma_1 \cdot \Gamma_2 + \Sigma \partial^2 \Psi_1 \cdot \Gamma_1)) + \frac{1}{24} (A \partial^2 \Gamma_1 \cdot \Gamma_1 + B \partial^2 \gamma_1 \cdot \Gamma_1 \\
 &\quad - B \partial^2 \Psi_1 \cdot \Gamma_1 - B_2 d\Psi_1 \cdot \Gamma_1 - B_3 \Psi_1 - A \partial^2 \Gamma_1 \cdot \Gamma_1 - B \partial^2 \gamma_1 \cdot \Gamma_1)] + O(\Delta t^4) \\
 &= -\Gamma_1 - \Delta t \Gamma_2 - \Delta t^2 \Gamma_3 - \Delta t^3 [B \Sigma \Psi_3 - \frac{1}{2} B (\gamma_3 + \Sigma d\Psi_1 \cdot \Gamma_2 + \Sigma d\Psi_2 \cdot \Gamma_1 - D \Sigma \Psi_2 \\
 &\quad + \frac{1}{6} D d\Psi_1 \cdot \Gamma_1 - \frac{1}{12} D_2 \Psi_1 - \frac{1}{12} \partial^2 \Psi_1 \cdot \Gamma_1) - \frac{1}{2} (A B + B D) \Sigma \Psi_2 + \frac{1}{4} B_2 \Psi_2 \\
 &\quad + \frac{1}{2} (A \Gamma_3 + B \gamma_3 + B \Sigma d\Psi_1 \cdot \Gamma_1 + d\Gamma_3 \cdot \Gamma_1) + \frac{1}{6} B_3 \Sigma \Psi_1 - \frac{1}{12} B_3 \Psi_1 - \frac{1}{6} A B \Sigma d\Psi_1 \cdot \Gamma_1 \\
 &\quad - \frac{1}{6} A (A \Gamma_2 + B \gamma_2) - \frac{1}{6} B (d\gamma_2 \cdot \Gamma_1 + d\gamma_1 \cdot \Gamma_2 + \Sigma \partial^2 \Psi_1 \cdot \Gamma_1) + \frac{1}{24} B \partial^2 \Psi_1 \cdot \Gamma_1 + \frac{1}{24} B_2 d\Psi_1 \cdot \Gamma_1 \\
 &\quad + \frac{1}{24} B_3 \Psi_1] + O(\Delta t^4) \\
 &= -\Gamma_1 - \Delta t \Gamma_2 - \Delta t^2 \Gamma_3 - \Delta t^3 [B \Sigma \Psi_3 - \frac{1}{2} B \Sigma d\Psi_2 \cdot \Gamma_1 - \frac{1}{12} B D d\Psi_1 \cdot \Gamma_1 + \frac{1}{24} B D_2 \Psi_1 \\
 &\quad + \frac{1}{24} B \partial^2 \Psi_1 \cdot \Gamma_1 - \frac{1}{2} A B \Sigma \Psi_2 + \frac{1}{4} B_2 \Psi_2 + \frac{1}{2} A (B \Sigma \Psi_2 - \frac{1}{6} B d\Psi_1 \cdot \Gamma_1 + \frac{1}{12} B_2 \Psi_1) \\
 &\quad + \frac{1}{2} (B \Sigma d\Psi_2 \cdot \Gamma_1 - \frac{1}{6} B \partial^2 \Psi_1 \cdot \Gamma_1 + \frac{1}{12} B_2 d\Psi_1 \cdot \Gamma_1) + \frac{1}{6} B_3 \Sigma \Psi_1 - \frac{1}{24} B_3 \Psi_1 - \frac{1}{6} A B \Sigma d\Psi_1 \cdot \Gamma_1 \\
 &\quad - \frac{1}{6} A^2 \Gamma_2 - \frac{1}{6} A B \gamma_2 - \frac{1}{6} B (d\gamma_2 \cdot \Gamma_1 + d\gamma_1 \cdot \Gamma_2 + \Sigma \partial^2 \Psi_1 \cdot \Gamma_1) + \frac{1}{24} B \partial^2 \Psi_1 \cdot \Gamma_1 \\
 &\quad + \frac{1}{24} B_2 d\Psi_1 \cdot \Gamma_1] + O(\Delta t^4) \\
 &= -\Gamma_1 - \Delta t \Gamma_2 - \Delta t^2 \Gamma_3 - \Delta t^3 [B \Sigma \Psi_3 - \frac{1}{12} B_2 d\Psi_1 \cdot \Gamma_1 + \frac{1}{24} B D_2 \Psi_1 + \frac{1}{4} B_2 \Psi_2 + \frac{1}{24} A B_2 \Psi_1 \\
 &\quad + \frac{1}{6} B_3 \Sigma \Psi_1 - \frac{1}{24} B_3 \Psi_1 - \frac{1}{6} A B \Sigma d\Psi_1 \cdot \Gamma_1 - \frac{1}{6} A^2 B \Sigma \Psi_1 - \frac{1}{6} A B \gamma_2 \\
 &\quad - \frac{1}{6} B (d\gamma_2 \cdot \Gamma_1 + d\gamma_1 \cdot \Gamma_2 + \Sigma \partial^2 \Psi_1 \cdot \Gamma_1) + \frac{1}{12} B_2 d\Psi_1 \cdot \Gamma_1] + O(\Delta t^4) \\
 &= -\Gamma_1 - \Delta t \Gamma_2 - \Delta t^2 \Gamma_3 - \Delta t^3 [B \Sigma \Psi_3 + \frac{1}{4} B_2 \Psi_2 + \frac{1}{6} B_3 \Sigma \Psi_1 - \frac{1}{6} A B \Sigma d\Psi_1 \cdot \Gamma_1 \\
 &\quad - \frac{1}{6} A^2 B \Sigma \Psi_1 - \frac{1}{6} A B (\Psi_2 - \Sigma d\Psi_1 \cdot \Gamma_1 + D \Sigma \Psi_1) - \frac{1}{6} B (d\gamma_2 \cdot \Gamma_1 + d\gamma_1 \cdot \Gamma_2 + \Sigma \partial^2 \Psi_1 \cdot \Gamma_1)] \\
 &\quad + O(\Delta t^4) \quad \text{because } A B_2 + B D_2 = B_3 \\
 &= -\Gamma_1 - \Delta t \Gamma_2 - \Delta t^2 \Gamma_3 - \Delta t^3 [B \Sigma \Psi_3 + \frac{1}{4} B_2 \Psi_2 + \frac{1}{6} B D_2 \Sigma \Psi_1 - \frac{1}{6} A B \Psi_2 \\
 &\quad - \frac{1}{6} B (d\gamma_2 \cdot \Gamma_1 + d\gamma_1 \cdot \Gamma_2 + \Sigma \partial^2 \Psi_1 \cdot \Gamma_1)] + O(\Delta t^4)
 \end{aligned}$$

because $B_3 - A(A B + B D) = B D_2$. The relations (72) and (73) are established and the proposition is proven. $\square$

7) Revisiting the “Berlin algorithm” in the linear case

In [3], we have presented the Berlin explicit algorithm in order to determine, with the help of formal calculus, the equivalent partial differential equations of a linear lattice Boltzmann scheme. We do not enter here into the details of our previous contribution because the linearized version of the present algorithm is much more simple.

Proposition 10. Linearized general expansion at fourth order

We suppose that the equilibrium value of the nonconserved moments is a linear function:

$$\Phi(W) = E W$$

with a given fixed $(q - N) \times N$ rectangle matrix E . Then the nonequilibrium moments can be expanded as

$$Y = [E + S^{-1}(\Delta t \beta_1 + \Delta t^2 \beta_2 + \Delta t^3 \beta_3)] W + O(\Delta t^4).$$

The $(q - N) \times N$ matrices β_j are linear operators of order j . The equivalent partial equivalent equations are linear and can be written as

$$\partial_t W + [\alpha_1 + \Delta t \alpha_2 + \Delta t^2 \alpha_3 + \Delta t^3 \alpha_4] W + O(\Delta t^4),$$

and α_j is a $N \times N$ differential matrix of order j . In other terms, $\Phi_j(W) = \beta_j W$ and $\Gamma_j(W) = \alpha_j W$ are linear functions of the conserved moments. The coefficient matrices α_j and β_j are computed according to a linearized version of (31):

$$(85) \quad \begin{cases} \alpha_1 = A + B E \\ \beta_1 = E \alpha_1 - C - D E \\ \alpha_2 = B \Sigma \beta_1 \\ \beta_2 = \Sigma \beta_1 \alpha_1 + E \alpha_2 - D \Sigma \beta_1 \\ \alpha_3 = B \Sigma \beta_2 + \frac{1}{12} B_2 \beta_1 - \frac{1}{6} B \beta_1 \alpha_1 \\ \beta_3 = \Sigma \beta_1 \alpha_2 + E \alpha_3 - D \Sigma \beta_2 + \Sigma \beta_2 \alpha_1 + \frac{1}{6} D \beta_1 \alpha_1 - \frac{1}{12} D_2 \beta_1 - \frac{1}{12} \beta_1 \alpha_1^2 \\ \alpha_4 = B \Sigma \beta_3 + \frac{1}{4} B_2 \beta_2 + \frac{1}{6} B D_2 \Sigma \beta_1 - \frac{1}{6} A B \beta_2 \\ \quad - \frac{1}{6} B E \alpha_1 \alpha_2 - \frac{1}{6} B E \alpha_2 \alpha_1 - \frac{1}{6} B \Sigma \beta_1 \alpha_1^2. \end{cases}$$

- Proof of Proposition 10.

First observe that $d\Phi \cdot \xi = E \xi$ for any test vector ξ . Due to the relation (36), we have

$\Gamma_1 = A W + B \Phi(W) = A W + B E W = (A + B E) W$ and $\alpha_1 = A + B E$. Due to (39), we have moreover

$$\begin{aligned} \Psi_1 &= d\Phi(W) \cdot \Gamma_1(W) - (C W + D \Phi(W)) \\ &= E \alpha_1 W - C W - D E W \\ &= (E \alpha_1 - C - D E) W \end{aligned}$$

and the second relation of (85) is established. Due to (41),

$$\begin{aligned} \Gamma_2(W) &= B \Sigma \Psi_1(W) \\ &= B \Sigma \beta_1 W. \end{aligned}$$

Then $\alpha_2 = B \Sigma \beta_1$ and the third line of (85) is proven. We have now from (53),

$$\Psi_2(W) = \Sigma d\Psi_1(W) \cdot \Gamma_1(W) + d\Phi(W) \cdot \Gamma_2(W) - D \Sigma \Psi_1(W)$$

$$\begin{aligned}
 &= \Sigma \beta_1 \alpha_1 W + E \alpha_2 W - D \Sigma \beta_1 W \\
 &= (\Sigma \beta_1 \alpha_1 + E \alpha_2 - D \Sigma \beta_1) W
 \end{aligned}$$

and $\beta_2 = \Sigma \beta_1 \alpha_1 + E \alpha_2 - D \Sigma \beta_1$ as suggested in the fourth line of (85). From the relation (55), we have now

$$\begin{aligned}
 \Gamma_3(W) &= B \Sigma \Psi_2(W) + \frac{1}{12} B_2 \Psi_1(W) - \frac{1}{6} B d\Psi_1(W). \Gamma_1(W) \\
 &= B \Sigma \beta_2 W + \frac{1}{12} B_2 \beta_1 W - \frac{1}{6} B \beta_1 \cdot \alpha_1 W \\
 &= (B \Sigma \beta_2 + \frac{1}{12} B_2 \beta_1 - \frac{1}{6} B \beta_1 \cdot \alpha_1) W.
 \end{aligned}$$

Then $\alpha_3 = B \Sigma \beta_2 + \frac{1}{12} B_2 \beta_1 - \frac{1}{6} B \beta_1 \cdot \alpha_1$ as proposed in the fifth line of (85). Observe now that due to (28), we have

$$\begin{aligned}
 \partial^2 \Psi_1. \Gamma_1(W) &= d^2 \Psi_1(W).(\Gamma_1, \Gamma_1) + d\Psi_1(W).d\Gamma_1(W). \Gamma_1 \\
 &= d\Psi_1(W).d\Gamma_1(W). \Gamma_1 && \text{because } \Psi_1 \text{ is linear} \\
 &= \beta_1 \alpha_1^2 W.
 \end{aligned}$$

Then due to (71), we have in this linear case,

$$\begin{aligned}
 \Psi_3(W) &= d\Phi(W). \Gamma_3(W) + \Sigma d\Psi_1(W). \Gamma_2(W) + \Sigma d\Psi_2(W). \Gamma_1(W) - D \Sigma \Psi_2(W) \\
 &\quad + \frac{1}{6} D d\Psi_1(W). \Gamma_1(W) - \frac{1}{12} D_2 \Psi_1(W) - \frac{1}{12} \partial^2 \Psi_1(W). \Gamma_1(W) \\
 &= E \alpha_3 W + \Sigma \beta_1 \alpha_2 W + \Sigma \beta_2 \alpha_1 W - D \Sigma \beta_2 W + \frac{1}{6} D \beta_1 \alpha_1 W \\
 &\quad - \frac{1}{12} D_2 \beta_1 W - \frac{1}{12} \beta_1 \alpha_1^2 W \\
 &= (E \alpha_3 + \Sigma \beta_1 \alpha_2 + \Sigma \beta_2 \alpha_1 - D \Sigma \beta_2 + \frac{1}{6} D \beta_1 \alpha_1 - \frac{1}{12} D_2 \beta_1 - \frac{1}{12} \beta_1 \alpha_1^2) W
 \end{aligned}$$

and $\beta_3 = E \alpha_3 + \Sigma \beta_1 \alpha_2 + \Sigma \beta_2 \alpha_1 - D \Sigma \beta_2 + \frac{1}{6} D \beta_1 \alpha_1 - \frac{1}{12} D_2 \beta_1 - \frac{1}{12} \beta_1 \alpha_1^2$. We have finally, due to (73),

$$\begin{aligned}
 \Gamma_4(W) &= B \Sigma \Psi_3(W) + \frac{1}{4} B_2 \Psi_2(W) + \frac{1}{6} B D_2 \Sigma \Psi_1 - \frac{1}{6} A B \Psi_2(W) - \frac{1}{6} B d\gamma_1(W). \Gamma_2(W) \\
 &\quad - \frac{1}{6} B d\gamma_2(W). \Gamma_1(W) - \frac{1}{6} B \Sigma \partial^2 \Psi_1(W). \Gamma_1(W) \\
 &= B \Sigma \beta_3 W + \frac{1}{4} B_2 \beta_2 W + \frac{1}{6} B D_2 \Sigma \beta_1 W - \frac{1}{6} A B \beta_2 W - \frac{1}{6} B E \alpha_2 W \\
 &\quad - \frac{1}{6} B E \alpha_1 W - \frac{1}{6} B \Sigma \beta_1 \alpha_1^2 W \\
 &= (B \Sigma \beta_3 + \frac{1}{4} B_2 \beta_2 + \frac{1}{6} B D_2 \Sigma \beta_1 - \frac{1}{6} A B \beta_2 - \frac{1}{6} B E \alpha_2 - \frac{1}{6} B E \alpha_1 - \frac{1}{6} B \Sigma \beta_1 \alpha_1^2) W
 \end{aligned}$$

and the proposition is established. $\square$

8) Conclusion

In this contribution, we have extended the Taylor expansion method of a multiple relaxation times lattice Boltzmann scheme up to fourth order accuracy. With this expansion, nonlinear partial differential equations for the conserved variables and various differential expressions for the nonconserved moments emerge naturally. This expansion has been validated with previous works on the subject.

Observe that the main result is implemented with this framework in the “pyLBM” software [30]. The next step is the introduction of source terms, in the spirit proposed in [25], and other geometries like triangles, in a way suggested in [23]. The adaptation of the Taylor expansion method to other variants of lattice Boltzmann schemes as recentered schemes

[28, 19], regularized lattice Boltzmann [44, 48, 53] or entropic lattice Boltzmann schemes [41] are also into study. After a first step in [47], the link between the single scale Taylor expansion and the multiple scale Chapman-Enskog is also a project for the near future.

Acknowledgments

The author thanks Bruce Boghosian, Filipa Caetano, Loïc Gouarin, Benjamin Graille, Pierre Lallemand and Paulo Philippi for impressive discussions. The scientific exchanges with the referee greatly improved the matter of this contribution. All the formal computations have been implemented with SageMath [52].

References

- [1] R. Agarwal, K.Y. Yun, R. Balakrishnan. “Beyond Navier Stokes - Burnett equations for flow simulations in continuum-transition regime” *Physics of Fluids* vol. **13**, p. 3061-3085, 2001.
- [2] F. Alexander, S. Chen and J. Sterling. “Lattice Boltzmann thermohydrodynamic”, *Physical Review E*, vol. **47**, R2249-R2252, 1993.
- [3] A. Augier, F. Dubois, B. Graille, P. Lallemand. “On rotational invariance of Lattice Boltzmann schemes”, *Computers and Mathematics with Applications*, vol. **67**, p. 239-255, 2014.
- [4] P. Bhatnagar, E. Gross, M. Krook. “A Model for Collision Processes in Gases I. Small Amplitude Processes in Charged and Neutral One-Component Systems”, *Physical Review*, vol. **94**, p. 511-525, 1954.
- [5] B. Boghosian, P. Coveney. “Inverse Chapman-Enskog Derivation of the Thermohydrodynamic Lattice-BGK Model for the Ideal Gas”, *International Journal of Modern Physics C*, vol. **9**, p. 1231-1245, 1998.
- [6] J. Broadwell. “Study of rarefied shear flow by the discrete velocity method”, *Journal of Fluid Mechanics*, vol. **19**, p. 401-414, 1964.
- [7] T. Carleman. *Problèmes mathématiques dans la théorie cinétique des gaz*, Publications Scientifiques de l’Institut Mittag-Leffler, Almqvist & Wiksells Boktryckeri AB, Uppsala University, 1957.
- [8] S. Chapman, T. G. Cowling. *The mathematical theory of non-uniform gases*, Cambridge University Press, 1939.
- [9] S. Chen, G. D. Doolen. “Lattice Boltzmann Method for Fluid Flows”, *Annual Review of Fluid Mechanics*, vol. **30**, p. 329-364, 1998.
- [10] P. Dellar. “Non-hydrodynamic modes and a priori construction of shallow water lattice Boltzmann equations”, *Physical Review E*, vol. **65**, 036309, 2002.
- [11] P. Dellar. “An interpretation and derivation of the lattice Boltzmann method using Strang splitting”, *Computers & Mathematics with Applications*, vol. **65**, p. 129-141, 2013.
- [12] P. Dellar. “Lattice Boltzmann algorithms without cubic defects in Galilean invariance on standard lattices”, *Journal of Computational Physics*, vol. **259**, p. 270-283, 2014.

- [13] F. Drui, E. Franck, P. Helluy, L. Navoret. “An analysis of over-relaxation in a kinetic approximation of systems of conservation laws”, *Comptes Rendus Mécanique*, vol. **347**, p. 259-269, 2019.
- [14] F. Dubois. “Une introduction au schéma de Boltzmann sur réseau”, *ESAIM Proceedings and Surveys*, vol. **18**, p. 181-215, 2007.
- [15] F. Dubois. “Equivalent partial differential equations of a Boltzmann scheme”, *Computers and mathematics with applications*, vol. **55**, p. 1441-1449, 2008.
- [16] F. Dubois. “Third order equivalent equation of lattice Boltzmann scheme”, *Discrete and Continuous Dynamical Systems-Series A*, vol. **23**, issue 1/2, p. 221-248, 2009.
- [17] F. Dubois. “Simulation of strong nonlinear waves with vectorial lattice Boltzmann schemes”, *International Journal of Modern Physics C*, vol. **25**, issue 12, 1441014, 2014.
- [18] F. Dubois, T. Février, B. Graille. “Third order equivalent equation for the relative velocity lattice Boltzmann schemes with one conservation law”, arxiv.org/abs/1502.02143, 7 February 2015.
- [19] F. Dubois, T. Février, B. Graille. “Lattice Boltzmann schemes with relative velocities” *Communications in Computational Physics*, vol. **17**, issue 4, p. 1088-1112, 2015
- [20] F. Dubois, B. Graille, P. Lallemand. “Recovering the full Navier Stokes equations with lattice Boltzmann schemes”, *30th International Symposium on Rarefied Gas Dynamics*, Jul 2016, Victoria, BC, Canada. American Institute of Physics Proceedings, vol. **1786**, (040003), pp.40003 - 40003, 2016.
- [21] F. Dubois, P. Lallemand. “Towards higher order lattice Boltzmann schemes ”, *Journal of Statistical Mechanics: Theory and Experiment*, P06006, 2009.
- [22] F. Dubois, P. Lallemand. “Quartic Parameters for Acoustic Applications of Lattice Boltzmann Scheme”, *Computers and Mathematics with Applications*, vol. **61**, p. 3404-3416, 2011.
- [23] F. Dubois, P. Lallemand. “On triangular lattice Boltzmann schemes for scalar problems”, *Communications in Computational Physics*, vol. **13**, p. 649-670, 2013.
- [24] F. Dubois, P. Lallemand. “Comparison of Simulations of Convective Flows”, *Communications in Computational Physics*, vol. **17**, p. 1169-1184, 2015.
- [25] F. Dubois, P. Lallemand, M.M. Tekitek. “Taylor expansion method for linear lattice Boltzmann schemes with an external drift. Application to boundary conditions”, High Order Nonlinear Numerical Schemes for Evolutionary PDEs, *Springer Lecture Notes in Computational Science and Engineering*, vol. **99**, R. Abgrall *et al.* Editors, p. 89-107, 2014.
- [26] S. Fletcher. *Semi-Lagrangian Advection Methods and Their Applications in Geoscience*, Elsevier Science Publishing Co, United States, 2019.
- [27] R. Gatignol. *Théorie cinétique des gaz à répartition discrète de vitesses*, Springer Lecture Notes in Physics, vol. **36**, 1975.
- [28] M.C. Geier, A. Greiner, J.G. Korvink. “Cascaded digital lattice Boltzmann automata for high Reynolds number flow”, *Physical Review E*, vol. **73**, p. 066705, 2006.

- [29] B. Graille. “Approximation of mono-dimensional hyperbolic systems: a lattice Boltzmann scheme as a relaxation method”, *Journal of Computational Physics*, vol. **266**, p. 74-88, 2014.
- [30] B. Graille, L. Gouarin. “pyLBM”, see www.imo.universite-paris-saclay.fr/~graille/pylbn.php and pypi.org/project/pylbn, 2017.
- [31] E. Gross. “Recent investigations of the Boltzmann equation”, *Rarefied Gas Dynamics*, vol. **3**, Proceedings of the First International Symposium held in Nice, F.M. Devienne Editor, Pergamon Press, London, 1960.
- [32] Z. Guo, C. Shu. *Lattice Boltzmann Method and Its Applications in Engineering*, World Scientific, 2013.
- [33] Z. Guo, C. Zheng, B. Shi. “Thermal lattice Boltzmann equation for low Mach number flows: decoupling model”, *Physical Review E*, vol. **75**, 036704, 2007.
- [34] G. Házi, P. Kávrán. “On the cubic velocity deviations in lattice Boltzmann methods”, *Journal of Physics A: Mathematical and General*, vol. **39**, Issue 12, p. 3127-3136, 2006.
- [35] T. Krüger, H. Kusumaatmaja, A. Kuzmin, O. Shardt, G. Silva, E.M. Viggen. *The Lattice Boltzmann Method: Principles and Practice*, Springer Verlag, 2017.
- [36] M. Hénon. “Viscosity of a Lattice Gas”, *Complex Systems*, vol. **1**, p. 763-789, 1987.
- [37] D. d’Humières. “Generalized Lattice-Boltzmann Equations”, in *Rarefied Gas Dynamics: Theory and Simulations*, vol. **159** of *AIAA Progress in Astronautics and Astronautics*, p. 450-458, 1992.
- [38] F. J. Higuera, S. Succi, R. Benzi. *Lattice gas dynamics with enhanced collisions*, Europhysics Letters, vol. **9**, issue 4, p. 345-349, 1991.
- [39] K. Huang. *Statistical Mechanics*, John Wiley & Sons Inc, New York, 1963.
- [40] I. V. Karlin, D. Sichau and S. S. Chikatamarla. “Consistent two-population lattice Boltzmann model for thermal flows”, *Physical Review E*, vol. **88**, 063310, 2013.
- [41] I. V. Karlin, A. N. Gorban, S. Succi, V. Boffi. “Maximum Entropy Principle for Lattice Kinetic Equations”, *Physical Review Letters*, vol. **81**, p. 6-9, 1998.
- [42] P. Lallemand, F. Dubois. “Comparison of Simulations of Convective Flows”, *Communications in Computational Physics*, vol. **17**, p. 1169-1184, 2015.
- [43] P. Lallemand, L.-S. Luo. “Theory of the lattice Boltzmann method: Dispersion, dissipation, isotropy, Galilean invariance, and stability”, *Physical Review E*, vol. **61**, p. 6546-6562, 2000.
- [44] J. Latt, B. Chopard. “Lattice Boltzmann method with regularized pre-collision distribution functions”, *Mathematics and Computers in Simulation*, vol. **72**, p. 165-168, 2006.
- [45] A. Lerat, R. Peyret. “Noncentered Schemes and Shock Propagation Problems”, *Computers and Fluids*, vol. **2**, p. 35-52, 1974.
- [46] H. Otomo, B.M. Boghosian, F. Dubois. “Two complementary lattice-Boltzmann-based analyses for nonlinear systems”, *Physica A: Statistical Mechanics and its Applications*, vol. **486**, p. 1000-1011, 2017.

- [47] H. Otomo, B.M. Boghosian, F. Dubois. “Efficient lattice Boltzmann models for the Kuramoto-Sivashinsky equation”, *Computers and Fluids*, vol. **172**, p. 683-688, 2018.
- [48] P. Philippi, L. Hegele. “From the continuous to the lattice Boltzmann equation: The discretization problem and thermal models”, *Physical Review E*, vol. **73**, Issue 5, 056702, 2006.
- [49] N.I. Prasianakis, I. Karlin, I. Mantzaras, K. Boulouchos. “Lattice Boltzmann method with restored Galilean invariance”, *Physical Review E*, vol. **79**, 066702, 2009.
- [50] Y.-H. Qian, D. d’Humières, P. Lallemand. *Lattice BGK models for Navier-Stokes equation*, Europhysics Letters, vol. **17**, issue 6, p. 479-484, 1992.
- [51] Y.H. Qian, Y. Zhou. “Higher-order dynamics in lattice-based models using the Chapman-Enskog method”, *Physical Review E*, vol. **61**, Issue 2, p. 2103-2106, 2000.
- [52] SageMath, the Sage Mathematics Software System (Version 7.5.1), The Sage Developers, <http://www.sagemath.org>, 2017.
- [53] X. Shan, X. F. Yuan, H. Chen H. “Kinetic theory representation of hydrodynamics: a way beyond the Navier-Stokes equation”, *Journal of Fluid Mechanics*, vol. **550**, p. 413-441, 2006.
- [54] G. Strang. “On the construction and comparison of difference schemes”, *SIAM Journal on Numerical Analysis*, vol. **5**, p. 506-517, 1968.
- [55] H. Struchtrup. “Failures of the Burnett and super-Burnett equations in steady state processes”, *Continuum Mechanics and Thermodynamics*, vol. **17**, p. 43-50, 2005.
- [56] S. Succi. “Lattice Boltzmann 2038”, *Europhysics Letters*, vol. **109**, 50001, 2015.
- [57] R.F. Warming, B.J. Hyett. “The modified equation approach to the stability and accuracy analysis of finite difference methods”, *Journal of Computational Physics*, vol. **14**, p. 159-179, 1974.
- [58] W.-A. Yong, W. Zhao, L.-S. Luo. “Theory of the Lattice Boltzmann method: Derivation of macroscopic equations via the Maxwell iteration” *Physical Review E*, vol. **93**, 033310, 2016.