Syntheses of Natural and Biologically Relevant Products through Asymmetric Metal-Catalyzed Domino Reactions. A Review

Hélène Pellissier

► To cite this version:

Hélène Pellissier. Syntheses of Natural and Biologically Relevant Products through Asymmetric Metal-Catalyzed Domino Reactions. A Review. Organic Preparations and Procedures International: The New Journal for Organic Synthesis, 2019, pp.311-344. 10.1080/00304948.2019.1590681 . hal-02106065

HAL Id: hal-02106065

<https://hal.science/hal-02106065>

Submitted on 22 Apr 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Syntheses of Natural and Biologically Relevant Products through Asymmetric Metal-Catalyzed Domino Reactions. A Review

Hélène Pellissier

Aix Marseille Univ, CNRS, Centrale Marseille, iSm2,
Marseille, France

Introduction	2
I. Syntheses Based on Palladium-Catalyzed Domino Reactions	2
1. <i>Heck-Initiated Domino Reactions as Key Steps</i>	2
2. <i>Wacker-Initiated Domino Reactions as Key Steps</i>	6
3. <i>Other Domino Reactions as Key Steps</i>	12
II. Syntheses Based on Copper-Catalyzed Domino Reactions	12
1. <i>Michael-Initiated Domino Reactions as Key Steps</i>	12
2. <i>Other Domino Reactions as Key Steps</i>	17
III. Syntheses Based on Rhodium-Catalyzed Domino Reactions	19
IV. Syntheses Based on Aluminum-Catalyzed Domino Reactions	23
V. Syntheses Based on Domino Reactions Catalyzed by Other Metals	25
Conclusion	30
Abbreviations	31
References	32

Syntheses of Natural and Biologically Relevant Products through Asymmetric Metal-Catalyzed Domino Reactions. A Review

Hélène Pellissier

Aix Marseille Univ, CNRS, Centrale Marseille, iSm2, Marseille, France

Introduction

A challenging goal in synthetic chemistry is the synthesis of complex products from simple starting materials by using one-step processes such as domino reactions.^{1–16} A domino reaction was defined by Tietze in 1993 as a reaction involving two or more bond-forming transformations, occurring under strictly the same reaction conditions without adding additional reagents and catalysts.^{17–20} These economic processes avoid costly protecting groups and time-consuming purification procedures after each step. In the last two decades, an impressive number of these powerful reactions have been developed, allowing the construction of very complex molecules in single steps from simple materials often with very high stereocontrol and yields. In most cases, remarkable enantioselectivities have been reported in reactions promoted by either chiral metal catalysts or organocatalysts. Especially, the considerable impact of the advent of asymmetric transition-metal catalysis has allowed the immense development of many highly efficient enantioselective metal-catalyzed domino reactions. The wide variety of these processes well reflects the variety of metals employed to promote them. This review collects for the first time the total syntheses of natural and biologically important products employing an enantioselective metal-catalyzed domino reaction as a key step, covering the years from 1996 to the present.

I. Syntheses Based on Palladium-Catalyzed Domino Reactions

1. Heck-Initiated Domino Reactions as Key Steps

Many syntheses of natural and bioactive products include as key steps asymmetric domino processes^{21–27} initiated by a Heck reaction.^{28–30} As an early example, the first total synthesis of the naturally occurring pentacyclic polyketide (+)-xestoquinone was developed by Keay *et al.*, in 1996.³¹ This compound exhibits important biological activities, such as anti-malarial, antiproliferative and cardiotonic properties. The key step of this unprecedented synthesis was an asymmetric palladium-catalyzed domino intramolecular Heck/cyclization reaction of a functionalized naphthyl triflate. As shown in *Scheme 1*, the reaction was promoted in toluene at 110°C by a combination of Pd₂(dba)₃ as precatalyst and (*S*)-BINAP as chiral ligand in the presence of pentamethylpiperidine (PMP). It afforded the corresponding key pentacyclic chiral intermediate in good yield (82%); however, the enantioselectivity

Scheme 1

was only moderate (68% *ee*). The latter was subsequently submitted to a catalytic hydrogenation on Pd/C followed by oxidation with ceric ammonium nitrate (CAN) to give the final natural product (+)-xestoquinone (*Scheme 1*).

A possible mechanism for the key enantioselective palladium-catalyzed domino intramolecular Heck/cyclization reaction of the total synthesis of (+)-xestoquinone (*Scheme 1*) is proposed in *Scheme 2*. It involves a *syn*-addition insertion of a palladium complex on the *Re*-face of the vinyl group to give a pentacyclic palladium intermediate which is further submitted to a *syn*- β -elimination to afford the final domino product.

The same year, Shibasaki *et al.* applied a related methodology to synthesize the marine natural product (+)-halenaquinone, exhibiting antibiotic, cardiogenic and protein tyrosine kinase inhibitory activities.^{32,33} In this case, the key step of the synthesis was the enantioselective palladium-catalyzed domino intramolecular Heck/cyclization reaction of a bicyclic triflate performed at 60°C in THF as solvent in the presence of a chiral palladium catalyst generated *in situ* from Pd(OAc)₂ and the chiral ligand (*S*)-BINAP mentioned above. It afforded the corresponding domino product in both high yield (78%) and enantioselectivity (87% *ee*), as shown in *Scheme 3*. The latter was further converted into the expected (+)-halenaquinone through fifteen supplementary steps which are detailed in *Scheme 3*. First, the domino product was converted into the corresponding alcohol through a two-step sequence. The further transformation of this alcohol into the corresponding triflate followed by reaction with a cyanohydrin triisopropylsilyl ether and then treatment with NaF provided the corresponding α,β -acetylenic ketone. After protection of the carbonyl function of this compound as an acetal and that of the ethynyl functionality with a triisopropylsilyl group, the resulting compound was submitted to benzylic oxidation with DDQ to give the corresponding ketone. Then, successive treatments with O₂ in the presence of *tert*-BuOK, NaI and TsOH afforded an α -iodo enol. Subsequently, the latter underwent cyclization in the presence of Pd₂(dba)₃ to give the corresponding pentacyclic compound which finally provided the anticipated (+)-halenaquinone through successive desilylation by treatment with tetrabutylammonium fluoride (TBAF) and CAN oxidation.

Scheme 2

Later in 2007, Zhu *et al.* developed another type of enantioselective palladium-catalyzed domino sequence initiated by a Heck reaction to be applied in a total synthesis of the naturally occurring butyryl-cholinesterase inhibitor (-)-physostigmine, which is employed for the treatment of glaucoma and Alzheimer's diseases.³⁴ As shown in *Scheme 4*, *ortho*-iodoanilidines reacted at 120°C in DMF as solvent with $\text{K}_4[\text{Fe}(\text{CN})_6]$ through a domino intramolecular Heck/cyanation reaction in the presence of $\text{Pd}(\text{dba})_2$ as precatalyst and (*S*)-DIFLUORPHOS as chiral ligand. This one-pot process led to the corresponding chiral oxindoles in moderate to good yields (54–78%) and enantioselectivities (61–79% *ee*). The authors demonstrated that one of these products ($\text{R}^1 = \text{OMe}$, $\text{R}^2 = \text{H}$) could be used as key intermediate in a total synthesis of (-)-physostigmine. As illustrated in *Scheme 4*, the reductive cyclization of this domino product with LiAlH_4 led to the corresponding hexahydropyrroloindole which was subsequently submitted to *N*-methylation under reductive amination conditions to afford esermethole. Then, cleavage of the methoxy group of esermethole with HBr led to the corresponding phenol which was finally submitted to NaH in the presence of *N*-succinimidyl-*N*-methylcarbamate to give the expected (-)-physostigmine.

In 2015, the same authors also developed a total synthesis of the natural powerful inhibitor of acetyl-cholinesterase (+)-esermethole.³⁵ It involved a chiral oxindole/azole bis-heterocycle as key intermediate which was generated through the first example of an asymmetric domino Heck/intermolecular arylation reaction. As shown in *Scheme 5*, the domino process between oxadiazoles and *N*-aryl acrylamides was promoted in acetonitrile at 80°C by a chiral palladium catalyst *in situ* generated from $\text{PdCl}_2(\text{MeCN})_2$ and a chiral Phox-type ligand in the presence of *N,N,N,N*-tetramethylguanidine (TMG) as base. The formed chiral domino bicyclic products bearing a quaternary stereocenter

Scheme 3

were obtained in moderate to excellent yields (55-94%) and high enantioselectivities (85-99% *ee*). One of these compounds ($\text{R}^1 = \text{R}^2 = \text{Me}$, $\text{R}^3 = \text{H}$, $\text{R}^4 = \text{Ph}$) obtained with 71% yield and 94% *ee* was converted into (+)-esermethole through five supplementary steps. The first step dealt with the reduction of this oxindole with LiAlH_4 , providing the corresponding pyrroloindoline without erosion of the enantiomeric purity. The latter product was successively submitted to reductive cleavage of the *N-N* bond of the hydrazine by treatment with SmI_2 , followed by *N*-methylation with formaldehyde and $\text{NaBH}(\text{OAc})_3$, which led to the corresponding *N*-methylated product. The latter was further converted into (+)-esermethole through successive regioselective bromination with NBS and CuI-catalyzed methoxylation of the resulting bromide.

In 2015, Lete *et al.* reported a novel access to the tetracyclic core of lycorane alkaloids, such as (+)-lycorine, based on an enantioselective intramolecular domino double Heck reaction of *N*-benzyl-2,3-dialkenylpyrroles.³⁶ As shown in Scheme 6, this process was promoted by a chiral palladium catalyst generated *in situ* from $\text{Pd}(\text{OAc})_2$ and (*R*)-BINAP as chiral ligand in the presence of 1,2,2,6,6-pentamethylpiperidine (PMP) as base, providing the corresponding chiral pyrrolophenanthridines in moderate to good yields (38-68%) and low to moderate enantioselectivities (7-72% *ee*).

Scheme 4

2. Wacker-Initiated Domino Reactions as Key Steps

Another important process based on palladium catalysis is the Wacker reaction.^{37,38} In 2005, Tietze *et al.* showed that it was possible to combine this process with a Heck reaction in an enantioselective domino Wacker/Heck reaction.³⁹ Later in 2006, this novel methodology was applied by these authors as key step in a novel total synthesis of vitamin E.⁴⁰ As depicted in [Scheme 7](#), this key reaction consisted in the enantioselective palladium-catalyzed domino Wacker/Heck reaction of a benzyl-protected phenol with methyl vinyl ketone performed at room temperature in dichloromethane as solvent. It was promoted by a chiral palladium catalyst generated *in situ* from Pd(TFA) $_2$ and (*S,S*)-Bn-BOXAX as chiral ligand in the presence of *p*-benzoquinone as reoxidant, leading to the corresponding chiral chroman in good yield (84%) and excellent enantioselectivity (97% ee), as shown in [Scheme 7](#).

$\text{R}^1 = \text{Me, Bn}$
 $\text{R}^2 = \text{Me, } n\text{-Hex}$
 $\text{R}^3 = \text{H, Br, 6-Cl, 7-Me}$
 $\text{R}^4 = \text{Ph, } p\text{-Tol, } p\text{-PhC}_6\text{H}_4, m\text{-MeOC}_6\text{H}_4, p\text{-ClC}_6\text{H}_4, \text{Bn}$

Synthesis of (+)-esermethole:

Scheme 5

This product was further converted into vitamin E through six supplementary steps, beginning with its hydrogenation on PtO_2 . Then, the boron enolate of the resulting compound was generated by treatment with diisopropylethylamine (DIPEA) as base and $c\text{-Hex}_2\text{BCl}$, and then submitted to an aldol reaction with a chiral aldehyde to give an α,β -unsaturated ketone after treatment with p -toluenesulfonic acid. The latter was subsequently submitted to

the addition of methyllithium to its carbonyl moiety to give the corresponding tertiary alcohol, which was converted into the corresponding diene over two steps by means of an elimination catalyzed by *p*-toluenesulfonic acid. Finally, this diene was hydrogenated over Pd/C with simultaneous deprotection of the benzylated phenolic hydroxyl group to give vitamin E.

In 2007, Tietze *et al.* reported another type of enantioselective palladium-catalyzed domino reaction initiated by a Wacker reaction, namely a three-component domino Wacker/carbonylation reaction.⁴¹ The process occurred between unsaturated phenols, alcohols and carbon dioxide to give the corresponding chiral chromans/benzodioxins with both high yields and enantioselectivities of up to 99% and 99% ee, respectively. The reaction was catalyzed by a palladium complex prepared *in situ* from $\text{Pd}(\text{TFA})_2$ and (*S,S*)-Bn-BOXAX as chiral ligand in the presence of *p*-benzoquinone as reoxidant under carbon dioxide atmosphere at ambient pressure. In 2008, the utility of this methodology was demonstrated by the same authors in a total synthesis of the natural

Scheme 7

product 4-dehydroxydiversonol.⁴² As shown in *Scheme 8*, the key step of this synthesis was the asymmetric domino Wacker/carbonylation reaction of an alkenyl phenol with methanol and carbon dioxide to give at room temperature the corresponding chiral ester in high yield (80%) and excellent enantioselectivity (96% *ee*). The latter was further converted into 4-dehydroxydiversonol through eight subsequent steps. The two first steps involved a reduction by treatment with diisobutylaluminum hydride (DIBAL-H) followed by a Wittig – Horner reaction of the thus-formed aldehyde with trimethyl phosphonoacetate, providing the corresponding α,β -unsaturated ester. The latter was submitted to the hydrogenation of its double bond on Pd/C followed by the oxidation of the chroman moiety by treatment with *t*-BuOOH to give the corresponding chromanone. Then, the intramolecular acylation at the α -position of the keto functionality by the ester

Scheme 8

moiety of this chromanone performed in the presence of TiCl_4 and TEA led to the corresponding tetrahydroxanthone. This compound was then oxidized by treatment with dimethyl dioxirane (DMDO) to afford the corresponding *trans*- α -hydroxy diketone as a single diastereomer. Finally, the reduction of the latter with NaBH_4 gave exclusively a *trans*-diol which was further deprotected by treatment with BBr_3 to afford the final 4-dehydroxydiversonol (*Scheme 8*).

In 2016, the same methodology was also applied by these authors to develop a formal total synthesis of the natural and biologically active pentacyclic chroman (-)-siccanin.⁴³ As shown in *Scheme 9*, the key step of this synthesis was the treatment of the same alkenyl phenol by a chiral palladium catalyst generated *in situ* from $\text{Pd}(\text{TFA})_2$ and enantiomeric ligand (*R,R*)-Bn-BOXAX under the same reaction conditions, affording the corresponding chiral chroman in good yield (71%) and high enantioselectivity (93% *ee*). This key intermediate was converted into (-)-siccanin through eight steps, beginning with its reduction with diisobutylaluminum hydride to give the corresponding

Scheme 9

aldehyde. The latter was submitted to an aldol reaction with a trimethylsilyl enol ether in the presence of BF₃(Et₂O) to provide the corresponding aldol product. Then, this compound was treated with Burgess reagent to give the related α,β -unsaturated ketone as a major diastereomer which was further separated and hydrogenated on Pd/C into the corresponding cyclohexanone. The latter was then submitted to a Peterson olefination

Scheme 10

to provide the corresponding methylene cyclohexane. A subsequent Sharpless dihydroxylation led to the corresponding diol. This compound constituted a late intermediate in the synthesis of (-)-siccanin earlier reported by Trost *et al.*, in 2003.⁴⁴

A possible mechanism for the key asymmetric domino Wacker/carbonylation reaction of the precedent synthesis is depicted in *Scheme 10*. It involves the addition of a palladium complex on the *Si*-face of the external double bond to give an intermediate palladium derivative. The latter was then submitted to a CO insertion followed by methoxylation to afford the final domino product.

3. Other Domino Reactions as Key Steps

In 2009, Trost *et al.* developed a total synthesis of the naturally occurring anticancer agent (+)-agelastatin A which was based on an enantioselective palladium-catalyzed domino double allylic alkylation reaction.⁴⁵ As illustrated in *Scheme 11*, the domino reaction of a *bis*-allylic carbonate with a diamine led to a key intermediate piperazinone in high yield (82%) and excellent enantioselectivity (>97% *ee*) when catalyzed by a chiral palladium complex *in situ* generated from $[\text{Pd}_2(\text{dba})_3](\text{CHCl}_3)$ and Trost ligand (*R,R*)-L_{ST} in the presence of acetic acid. This key piperazinone was further converted into (+)-agelastatin A through four further steps, the first one of which was the copper-catalyzed aziridination of the domino product into the corresponding tetracyclic aziridine. The treatment of the latter with $\text{In}(\text{OTf})_3$ afforded a tricyclic α -amino ketone. Then, the required installation of the D-ring urea of (+)-agelastatin A was achieved by treating this α -amino ketone with methyl isocyanate in the presence of Cs_2CO_3 to give the corresponding tetracyclic compound. Finally, removing the *N*-Ts and *N*-OMe groups was achieved through SmI_2 -mediated radical reduction, leading to (+)-agelastatin A.

II. Syntheses Based on Copper-Catalyzed Domino Reactions

1. Michael-Initiated Domino Reactions as Key Steps

Related to the growing interest in green chemistry, domino reactions promoted by copper catalysts represent a challenge in organic chemistry. Among them, a number of asymmetric domino reactions are initiated by a Michael addition. For example, an enantioselective three-component copper-catalyzed domino Michael/aldol reaction

Scheme 11

constituted the key step in a total synthesis of (–)-prostaglandin E₁ methyl ester reported by Feringa *et al.*, in 2001.⁴⁶ As shown in [Scheme 12](#), this key reaction occurred between a dioxane acetal, a dialkylzinc reagent and an aldehyde in the presence of a copper catalyst *in situ* generated from Cu(OTf)₂ and a chiral phosphoramidite as ligand, giving the corresponding Michael/aldol domino product in 60% yield as an inseparable 83:17 mixture of diastereomers. The reduction of the ketone moiety in this product proceeded stereoselectively using Zn(BH₄)₂ to give the corresponding diol as a major diastereomer with 94% ee and 63% yield. In the next step, the silyl group was removed by treatment with TBAF and then the resulting compound was acetylated with Ac₂O in pyridine to give the corresponding diacetate. Then, the latter was submitted to a 1,3-allylic transposition in the presence of Pd(MeCN)₂Cl₂ to give the corresponding allylic acetate. After deacetylation with K₂CO₃ in methanol, a chiral 1,5-diol was

Scheme 12

obtained. The final step of the synthesis dealt with the deprotection of the ketone functionality of this compound by using $(\text{NH}_4)_2\text{Ce}(\text{NO}_3)_6$, leading to (–)-prostaglandin E₁ methyl ester, as detailed in [Scheme 12](#).

Scheme 13

In 2010, the same authors reported another enantioselective copper-catalyzed domino reaction initiated by a Michael addition which constituted the key step of formal syntheses of natural products cascarillic acid and grenadamide.⁴⁷ This asymmetric domino Michael/intramolecular alkylation reaction occurred between Grignard reagents and 4-chloro- α,β -unsaturated esters/thioesters/ketones in the presence of a combination of CuI and (R)-Tol-BINAP as chiral ligand, as shown in [Scheme 13](#). It afforded the corresponding chiral *trans*-1-alkyl-2-substituted cyclopropane esters/thioesters/ketones in moderate to excellent yields (56->95%) and high enantioselectivities (84-98% ee). These various products constituted key intermediates for formal syntheses of cascarillic acid and grenadamide previously reported by Cheeseman and Bull.⁴⁸

In 2012, Riant *et al.* described the preparation of a key intermediate for a total synthesis of the natural diterpene marrubiin exhibiting antinociceptive and expectorant effects. This was based on an enantioselective copper-catalyzed domino reductive Michael/aldol cyclization reaction.⁴⁹ As shown in [Scheme 14](#), this domino process began with the reductive Michael reaction of a diketooester with phenylsilane performed in the presence of a chiral copper catalyst derived from a Taniaphos ligand, leading to

formal synthesis of marrubiin:

Scheme 14

the corresponding bicyclic chiral domino product as a single *cis*-diastereomer in good yield (80%) and excellent enantioselectivity (95% *ee*). This product was further submitted to dehydration by successive treatments with TFA, TMSCH₂N₂ and SOCl₂ to give the corresponding non-conjugated cyclohexenone. The latter was then methylated in the presence of LDA and methyl iodide to provide the corresponding bicyclic ester after required protection of the ketone group as dioxolane. This compound constituted a crucial intermediate in the synthesis of marrubiin previously reported.⁵⁰

2. Other Domino Reactions as Key Steps

In 2011, Hajra and Bar developed an enantioselective copper-catalyzed domino aziridination/Friedel – Crafts cyclization reaction as key step in a total synthesis of the dopamine D1 agonist A-86929.⁵¹ As depicted in *Scheme 15*, the reaction of a functionalized styrene with PhINNs was promoted by a copper catalyst *in situ* generated from Cu(OTf)₂ and a chiral bisoxazoline ligand, leading to the corresponding chiral bicyclic domino product in good yield (82%), complete *trans*-diastereoselectivity (>98% *de*) and excellent enantioselectivity (95% *ee*). This product was converted into A-86929 through four additional steps, beginning with the reaction of its sodium salt generated by treatment with NaH with MOMCl, providing the corresponding methoxy methyl ether. The Pictet – Spengler-type cyclization of this compound using TMSOTf afforded a tetracyclic product which was subsequently deprotected by treatment with *p*-methoxythiophenol and K₂CO₃ to give the corresponding secondary amine. Finally, the demethylation of the latter performed with BBr₃ accomplished the synthesis of A-86929.

Earlier in 2006, Knochel *et al.* had developed a synthesis of the highly toxic alkaloid (*S*)-(+)-coniine, which is known to induce curare-type paralysis.⁵² The key step of this synthesis was an asymmetric copper-catalyzed three-component reaction among butyraldehyde, trimethylsilylacetylene and dibenzylamine catalyzed by a chiral complex derived from CuBr and (*R*)-QUINAP as ligand (*Scheme 16*). This domino process afforded the corresponding chiral propargylamine in both excellent yield (90%) and enantioselectivity (90% *ee*). This product was further converted into (*S*)-(+)-coniine through six steps. First, the domino compound was desilylated by treatment with TBAF to give the corresponding alkyne which was subsequently deprotonated with *n*-BuLi and then alkylated with ethylene oxide. After silylation of the resulting alcohol with TIPSCl, the corresponding TIPS ether was obtained. Then, the latter was successively submitted to hydrogenation on Pd/C, desilylation with Bu₄NF and intramolecular Mitsunobu reaction to finally afford (*S*)-(+)-coniine.

In 2013, Watanabe and Shibasaki reported a formal total synthesis of antiviral oseltamivir phosphate (Tamiflu) which employed as crucial step another enantioselective copper-catalyzed three-component reaction.⁵³ As shown in *Scheme 17*, the reaction occurred among bis(2-phenylallyl)amine, ethyl propiolate and an aldehyde and was promoted by a chiral copper catalyst *in situ* generated from CuBr and a (*R,R*)-O-PINAP ligand. It afforded the corresponding chiral domino product in good yield (84%) and moderate enantioselectivity (76% *ee*). This compound was further converted through a seven-step sequence into Corey's intermediate for the total synthesis of oseltamivir phosphate. The first step of the sequence dealt with the poisoned palladium-catalyzed hydrogenation of the domino product into the corresponding triene followed by a Dieckmann condensation in the presence of LiHMDS to give the corresponding six-membered compound. The ketone functionality of the latter was further reduced with NaBH₄ to give the corresponding alcohol. Then, the two 2-phenylallyl groups of this alcohol were removed by allylic substitution in the presence of Pd(PPh₃)₄ and *N,N*-dimethylbarbituric acid as nucleophile, which was followed by introduction of the Boc group to give the corresponding *N*-Boc-protected amine. Finally, mesylation of the latter followed by subsequent β-elimination afforded Corey's intermediate for the total synthesis of oseltamivir phosphate.

synthesis of A-86929:

Scheme 15

Scheme 16

III. Syntheses Based on Rhodium-Catalyzed Domino Reactions

Domino carbonyl ylide formation/1,3-dipolar cycloaddition reactions allow a direct access to many complex heterocycles.^{54–56} In 2010, Hashimoto *et al.* developed an asymmetric version of this methodology as a significant step in a total synthesis of a natural product.⁵⁷ The key intermediate of this synthesis was formed through the enantioselective rhodium-catalyzed domino reaction of the formyl-derived carbonyl ylide generated from a diazo compound with 4-hydroxy-3-methoxyphenylacetylene. When the reaction was catalyzed by $\text{Rh}_2(\text{S-TCPTTL})_4$, this cycloadduct was obtained in good yield (73%) and excellent enantioselectivity (95% ee), as illustrated in [Scheme 18](#). This product was further converted into the expected natural product having blood circulation properties through eight steps. In the first one, the phenolic ester group of the domino product was protected as *tert*-butyldiphenylsilyl ether. Then, treatment of this ether by NaHMDS followed by addition of PhNTf_2 and subsequent palladium-catalyzed reduction of the resulting enol triflate led to the corresponding 1,4-diene. Then, the reduction of the latter with LiAlH_4 and subsequent silylation provided the corresponding diether which was further submitted to allylic oxidation with SeO_2 to afford the corresponding allylic alcohol as a single diastereomer. The latter was oxidized with MnO_2 to give the corresponding α,β -unsaturated ketone which was finally converted into the

Scheme 17

desired natural product through removing the two TBDPS protecting groups by treatment with TBAF.

The same authors also described the synthesis of another natural and biologically active product, descurainin, on the basis of the same domino reaction albeit catalyzed with the enantiomeric catalyst $\text{Rh}_2(\text{R-TCPTTL})_4$.⁵⁸ As shown in [Scheme 19](#), by using this catalyst the corresponding enantiomeric cycloadduct was obtained in good yield (77%) and excellent enantioselectivity (95% *ee*). A sequence of twelve further

Scheme 18

Scheme 19

supplementary steps allowed expected descurainin to be achieved from this domino product. In the first step, hydrogenation of the latter on Pd/C led to the corresponding saturated bicyclic compound as a single diastereomer, which was then treated with $(\text{KSO}_3)_2\text{NO}$ in the presence of KH_2PO_4 to give the corresponding *o*-quinone. The latter was directly converted into a catechol by treatment with $\text{Na}_2\text{S}_2\text{O}_4$. Subsequent treatment of this catechol with methyl iodide and K_2CO_3 led to the corresponding per-methylated compound. Then, treatment of the latter with NaHMDS followed by addition of PhNTf_2 and subsequent palladium-catalyzed reduction of the resulting enol triflate gave the corresponding alkene. The further reduction of the latter with LiAlH_4 led to the corresponding alcohol. Subsequent treatment of this alcohol with NbCl_5 allowed regioselective demethylation to give the corresponding phenol as a sole product. Then, protection of the two hydroxyl groups of this phenol with TBDPSCl provided the corresponding diether which was further submitted to an allylic oxidation with SeO_2 followed by an oxidation with MnO_2 to give the corresponding α,β -unsaturated ketone. Finally, the removal of the two TBDPS protecting groups of this enone with TBAF completed the total synthesis of descurainin (*Scheme 19*).

IV. Syntheses Based on Aluminum-Catalyzed Domino Reactions

In 1996, Shibasaki *et al.* reported the first catalytic asymmetric domino Michael/aldol reaction.⁵⁹ Promoted by a heterobimetallic chiral complex, such as $\text{AlLibis}[(R)\text{-binaphthoxide}]$, this three-component reaction between cyclopentenones, aldehydes and methyl-substituted dibenzyl malonates led to the corresponding multifunctionalized chiral cyclopentanones with enantioselectivities of up to 91% *ee*. In 1998, this methodology was applied by the same authors to develop a total synthesis of 11-deoxy-PGF_{1 α} .⁶⁰ As shown in *Scheme 20*, the three-component domino Michael/aldol reaction of cyclopentenone, an aldehyde and dibenzyl methylmalonate performed in the presence of enantiomeric $\text{AlLibis}[(S)\text{-binaphthoxide}]$ provided the corresponding chiral product in 84% yield and up to 88% *de*. This domino product was further converted through fourteen steps into 11-deoxy-PGF_{1 α} . The sequence began with the conversion of the domino product into the corresponding α,β -unsaturated ketone through successive mesylation and elimination. Then, hydrosilylation of this ketone catalyzed by chlorotris(triphenylphosphine)rhodium gave the corresponding *trans*-ketone which was subsequently reduced with L-Selectride into the corresponding alcohol as a single product. Then, this alcohol was protected with *tert*-butyldimethylsilyl trifluoromethanesulfonate to give the corresponding TBS ether. The latter was further transformed into the corresponding methyl ketone through successive treatments with hydrogen on Pd/C, $\text{Pb}(\text{OAc})_4$ and K_2CO_3 . Then, the aldol reaction of this methyl ketone with hexanal followed by elimination provided the corresponding α,β -unsaturated ketone. The subsequent treatment of this ketone with La-(*S*)-3-(hydroxymethyl)binaphthol complex led to the corresponding epoxide which was further subjected to Wharton reaction conditions to give the corresponding allylic alcohol. Finally, the deprotection of this alcohol by treatment with HF led to 11-deoxy-PGF_{1 α} .

Another asymmetric aluminum-catalyzed three-component reaction was developed by Shibasaki *et al.*, involving a pyridine derivative, TMSCN and a carbonyl chloride as substrates.⁶¹ Promoted by a chiral aluminum catalyst *in situ* generated from Et_2AlCl and a BINOL-derived sulfoxide chiral ligand, it afforded the corresponding chiral cyanide in quantitative yield (98%) and high enantioselectivity (91% *ee*), as shown in

Scheme 20

Scheme 21

Scheme 21. The synthetic utility of this methodology was demonstrated by its application to the formal synthesis of the dopamine D_4 receptor-selective antagonist CP-293,019.⁶² Indeed, the domino product was converted through nine steps into an important primary alcohol intermediate in the synthesis of CP-293,019. The synthetic route began with a two-step hydrogenation of the domino product followed by a protection/deprotection sequence, leading to the corresponding tetrahydropyridine. Reduction of the latter with NaBH_3CN *via* an iminium cation proceeded in a 4:1 ratio, and the isolated major diastereomer was annulated to give a bicyclic diamine. Then, the known key primary alcohol intermediate⁶³ was achieved from this bicyclic diamine in three supplementary steps.

V. Syntheses Based on Domino Reactions Catalyzed by Other Metals

In 1999, Ihara and Nemoto reported a total synthesis of the natural product (+)-equile-nin based on an asymmetric manganese-catalyzed domino epoxidation/ring-expansion

Scheme 22

reaction of a cyclopropylidene.^{64,65} This domino process was catalyzed by a chiral pre-formed salen manganese complex in the presence of sodium hypochlorite as oxidant, leading through successive epoxidation and ring-expansion to the corresponding chiral cyclobutanone in moderate yield (55%) and good enantioselectivity (78% *ee*), as

Scheme 23

illustrated in [Scheme 22](#). The latter was converted into the steroid (+)-equilenin through five supplementary steps. This sequence began with the stereoselective conversion of the domino product into the corresponding isopropenylcyclobutanol by Grignard reaction with isopropenylmagnesium bromide in the presence of cerium trichloride. Then, a palladium-mediated cascade ring expansion/insertion reaction occurred to give the corresponding 11-methylene steroid as a diastereomeric mixture which was subsequently treated with osmium tetroxide and sodium periodate to give the corresponding diketone. The latter was further submitted to hydrogenolysis on Pt/C in the presence of PdCl_2 to afford equilenin methyl ether. A subsequent deprotection performed in the presence of BBr_3 led finally to (+)-equilenin.

With the aim of synthesizing precursors to GABA analogues, Sudalai and Paraskar developed in 2006 a novel enantioselective cobalt-catalyzed domino reductive cyclization reaction of substituted γ -azido- α,β -unsaturated esters to afford the corresponding γ -lactams with enantioselectivities of up to 98% ee.⁶⁶ The process was promoted by a cobalt catalyst *in situ* generated from CoCl_2 and a chiral oxazoline ligand in the

Scheme 24

presence of NaBH_4 as reducing agent. This methodology was applied to the syntheses of (*R*)-rolipram and (*R*)-baclofen which are respectively a potent inhibitor of the cardiac cyclic AMP phosphodiesterase and a GABA agonist. As shown in [Scheme 23](#) (first equation), (*R*)-rolipram was directly obtained with both high yield (92%) and enantioselectivity (92% *ee*) from the domino reaction of a γ -azido olefinic ester prepared in six steps from 2-methoxyphenol. The synthesis of (*R*)-baclofen was achieved through hydrolysis with 6M HCl of another domino product ([Scheme 23](#), second equation) also obtained with both high yield (82%) and enantioselectivity (89% *ee*).

In 2007, Kobayashi *et al.* reported the total synthesis of the natural cytotoxic C_2 -symmetric cyclic decapeptide onchidin.⁶⁷ The key step of this synthesis was an enantioselective Mannich-type reaction between 2-amino-*m*-cresol, 5-hexynal and a ketene silyl acetal derived from phenyl propionate which was promoted by a chiral zirconium catalyst derived from a fluorinated (*S*)-BINOL as ligand. As shown in [Scheme 24](#), this three-component process led to the formation of the corresponding chiral

mechanism of the domino aldol/cyclization:

Scheme 25

Mannich *anti*-product in both good yield (85%) and diastereoselectivity (82% *de*) combined with excellent enantioselectivity (96% *ee*). Then, this domino product was converted into the corresponding *N*-Boc-protected dipeptide in four steps. The latter was further coupled with the hydrochloride salt of a tridepsipeptide to give the corresponding pentadepsipeptide. This compound was coupled with another pentadepsipeptide to give the corresponding linear decadepsipeptide. Cleavage of the TMS group followed by *N*-Boc deprotection and final macrocyclization afforded onchidin.

In 2016, Dixon *et al.* reported a novel total synthesis of the antibiotic (-)-chloramphenicol based on an enantioselective silver-catalyzed domino aldol/cyclization reaction.⁶⁸ In this key step, *p*-nitrobenzaldehyde reacted with an isocyano acetate to give the corresponding *trans*-oxazoline in moderate yield (72%) and both good diastereoselectivity (82% *de*) and enantioselectivity (89% *ee*), as illustrated in *Scheme 25*. This process was promoted at room temperature with a chiral silver catalyst *in situ* generated from Ag₂O and a cinchona alkaloid as ligand. The formed chiral domino product was further converted into (-)-chloramphenicol through four supplementary steps, which are detailed in *Scheme 25*. The ring-opening of the domino product using thionyl chloride provided the corresponding amino alcohol which was further acetylated with dichloroacetyl chloride to give the corresponding dichloroacetamide. Finally, the chemoselective reduction of the ester group in this compound with sodium borohydride delivered (-)-chloramphenicol.

Conclusion

This review collects total asymmetric syntheses of natural and/or biologically important products using enantioselective metal-catalyzed domino reactions as key steps. It shows that these economic one-pot processes have already allowed many important complex molecules to be achieved from simple starting materials. The increasing diversity of these fascinating reactions well reflects that of the metals employed to promote them. For example, various types of chiral palladium catalysts have been applied to promote asymmetric Heck-initiated domino reactions with enantioselectivities of up to 94% *ee* which were employed as key steps in total syntheses of different natural and/or bioactive products, such as (-)-xestoquinone, (+)-halenaquinone, (-)-physostigmine and (+)-esermethole. Other chiral palladium complexes were also used to promote asymmetric Wacker-initiated domino reactions achieved with up to 96% *ee* used as key steps in total syntheses of vitamin E, 4-dehydroxydiversonol and (-)-siccanin. Moreover, another type of palladium-catalyzed domino process, including a domino double allylic alkylation reaction, was employed in the total synthesis of (+)-agelastatin, providing enantioselectivities of up to >97% *ee*. Related to the growing interest in green chemistry, domino reactions promoted by green copper catalysts represent a challenge in organic chemistry. In this context, many types of asymmetric domino reactions catalyzed by a variety of chiral copper complexes have been applied as key steps in total syntheses. For example, copper-catalyzed Michael-initiated domino processes achieved with enantioselectivities of 94-98% *ee* have allowed the synthesis of important products, such as PGE₁ methyl ester, grenadamide and marrubiin. Moreover, total syntheses of A-86929, (*S*)-(+)-coniine and Tamiflu have been based on other enantioselective domino reactions including three-component ones with enantioselectivities of up to 95% *ee*. Along with palladium and copper chiral complexes, rhodium chiral catalysts have been applied in different types of domino reactions with 88-91% *ee*, constituting the key steps in the total syntheses of 11-deoxy-PG_{1α} and CP-293,019. Moreover, the

use of chiral complexes derived from other metals, such as manganese, cobalt, zirconium and silver, also provided excellent enantioselectivities of up to 96% *ee* in different types of domino reactions in the total syntheses of (+)-equilenin, (*R*)-baclofen, onchidin and (-)-chloramphenicol. Undoubtedly, many other enantioselective metal-promoted asymmetric domino reactions will be discovered in the near future and further applied as key economic steps in total synthesis.

Abbreviations

BINAP	2,2'-bis(diphenylphosphino)-1,1'-binaphthyl
BINOL	1,1'-bi-2-naphthol
Bn	benzyl
Boc	<i>tert</i> -butoxycarbonyl
BOXAX	2,2'-bis(oxazolyl)-1,1'-binaphthyl
Bt	benzotriazol
CAN	ceric ammonium nitrate
dba	(<i>E,E</i>)-dibenzylideneacetone
DBU	1,8-diazabicyclo[5.4.0]undec-7-ene
DDQ	2,3-dichloro-5,6-dicyano- <i>p</i> -benzoquinone
de	diastereomeric excess
DEAD	diethyl azodicarboxylate
DIBAL (DIBAL-H)	diisobutylaluminium hydride
DIFLUORPHOS	5,5'-bis(diphenylphosphino)-2,2,2',2'-tetrafluoro-4,4'-bi-1,3-benzodioxole
DIPEA	diisopropylethylamine
DMAP	4-(dimethylamino)pyridine
DMDO	dimethyl dioxirane
DMF	<i>N,N</i> -dimethylformamide
EDC	1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride
ee	enantiomeric excess
GABA	γ -aminobutyric acid
Hex	hexyl
HMDS	hexamethyldisilazide
LDA	lithium diisopropylamide
MOM	methoxymethyl
NBS	<i>N</i> -bromosuccinimide
NMI	<i>N</i> -methylimidazole
Ms	mesyl
M.S.	molecular sieves
NBS	<i>N</i> -bromosuccinimide
NMI	<i>N</i> -methylimidazole
Ns	nosyl (4-nitrobenzene sulfonyl)
Phox	phosphinooxazoline
PINAP	4-[2-(diphenylphosphino)-1-naphthalenyl]- <i>N</i> -[1-phenylethyl]-1-phthalazinamine
PMP	1,2,2,6,6-pentamethylpiperidine
QUINAP	1-(2-diphenylphosphino-1-naphthyl)isoquinoline
r.t.	room temperature
SES	β -trimethylsilylethanesulfonyl

Taniaphos	[2-diphenylphosphinoferrocenyl](N,N-dimethylamino)(2-diphenylphosphinophenyl)methane
TBAF	tetrabutylammonium fluoride
TBDPS	<i>tert</i> -butyldiphenylsilyl
TBHP	<i>tert</i> -butylhydroperoxide
TBS	<i>tert</i> -butyldimethylsilyl
TCPTTL	N-tetrachlorophthaloyl- <i>tert</i> -leucinate
TEA	triethylamine
Tf	trifluoromethanesulfonyl
TFA	trifluoroacetic acid
THF	tetrahydrofuran
TIPS	triisopropylsilyl
TMG	1,1,3,3-tetramethylguanidine
TMS	trimethylsilyl
Tol	tolyl
Ts	4-toluenesulfonyl (tosyl)
Val	valine

1. J. Zhu and H. Bienaymé, *Multicomponent Reactions*, p. 484, Wiley-VCH, Weinheim, 2005.
2. H. Pellissier, *Tetrahedron*, **62**, 1619 (2006).
3. D. Enders, C. Grondal, and M. R. M. Hüttl, *Angew. Chem. Int. Ed.*, **46**, 1570 (2007).
4. G. Guillena, D. J. Ramon, and M. Yus, *Tetrahedron: Asymmetry*, **18**, 693 (2007).
5. C. J. Chapman and C. G. Frost, *Synthesis*, 1 (2007).
6. A.-N. Alba, X. Companyo, M. Viciano, and R. Rios, *Curr. Org. Chem.*, **13**, 1432 (2009).
7. B. B. Touré and D. G. Hall, *Chem. Rev.*, **109**, 4439 (2009).
8. C. Grondal, M. Jeanty, and D. Enders, *Nature Chemistry*, **2**, 167 (2010).
9. L. Albrecht, H. Jiang, and K. A. Jørgensen, *Angew. Chem. Int. Ed.*, **50**, 8492 (2011).
10. H. Pellissier, *Adv. Synth. Catal.*, **354**, 237 (2012).
11. C. De Graaff, E. Ruijter, and R. V. A. Orru, *Chem. Soc. Rev.*, **41**, 3969 (2012).
12. H. Clavier and H. Pellissier, *Adv. Synth. Catal.*, **354**, 3347 (2012).
13. H. Pellissier, *Chem. Rev.*, **113**, 442 (2013).
14. H. Pellissier, *Adv. Synth. Catal.*, **358**, 2194 (2016).
15. S. A. Snyder, *Science of Synthesis. Applications of Domino Transformations in Organic Synthesis*, p. 1284, Thieme Verlag, Stuttgart, 2016.
16. T. Chanda and J. C.-G. Zhao, *Adv. Synth. Catal.*, **360**, 2 (2018).
17. L. F. Tietze and U. Beifuss, *Angew. Chem. Int. Ed. Engl.*, **32**, 131 (1993).
18. L. F. Tietze, *Chem. Rev.*, **96**, 115 (1996).
19. L. F. Tietze, G. Brasche, and K. Gericke, *Domino Reactions in Organic Synthesis*, p. 631, Wiley-VCH, Weinheim, 2006.

20. L. F. Tietze, *Domino Reactions - Concepts for Efficient Organic Synthesis*, p. 648, Wiley-VCH, Weinheim, 2014.
21. O. Reiser, M. Weber, and A. de Meijere, *Angew. Chem. Int. Ed. Engl.*, **59**, 74 (1989).
22. K. Albrecht, O. Reiser, M. Weber, and A. de Meijere, *Angew. Chem. Int. Ed. Engl.*, **28**, 1037 (1989).
23. S. P. Watson, G. R. Knox, and N. M. Heron, *Tetrahedron Lett.*, **35**, 9763 (1994).
24. R. Grigg, P. Kennewell, A. Teasdale, and V. Sridharan, *Tetrahedron Lett.*, **34**, 153 (1993).
25. R. Grigg and V. Sridharan, *Tetrahedron Lett.*, **33**, 7965 (1992).
26. G. D. Harris, R. J. Herr, and S. M. Weinreb, *J. Org. Chem.*, **58**, 5452 (1993).
27. F. E. Meyer, K. H. Ang, A. G. Steinig, and A. de Meijere, *Synlett*, 191 (1994).
28. R. F. Heck, *J. Am. Chem. Soc.*, **90**, 5518 (1968).
29. M. Oestreich, *The Mizoroki – Heck Reaction*, p. 587, Wiley, Chichester, 2009.
30. D. Mc Cartney and P. J. Guiry, *Chem. Soc. Rev.*, **40**, 5122 (2011).
31. S. P. Maddaford, N. G. Andersen, W. A. Cristofoli, and B. A. Keay, *J. Am. Chem. Soc.*, **118**, 10766 (1996).
32. A. Kojima, T. Takemoto, M. Sodeoka, and M. Shibasaki, *J. Org. Chem.*, **61**, 4876 (1996).
33. A. Kojima, T. Takemoto, M. Sodeoka, and M. Shibasaki, *Synthesis*, **581** (1998).
34. A. Pinto, Y. Jia, and L. Neuville, J. Zhu, *Chem. Eur. J.*, **13**, 961 (2007).
35. W. Kong, Q. Wang, and J. Zhu, *J. Am. Chem. Soc.*, **137**, 16028 (2015).
36. E. Coya, N. Sotomayor, and E. Lete, *Adv. Synth. Catal.*, **357**, 3206 (2015).
37. J. Reinhard, *Angew. Chem. Int. Ed.*, **48**, 9034 (2009).
38. R. I. McDonald, G. Liu, and S. S. Stahl, *Chem. Rev.*, **111**, 2981 (2011).
39. L. F. Tietze, K. M. Sommer, J. Zinngrebe, and F. Stecker, *Angew. Chem. Int. Ed.*, **44**, 257 (2005).
40. L. F. Tietze, F. Stecker, J. Zinngrebe, and K. M. Sommer, *Chem. Eur. J.*, **12**, 8770 (2006).
41. L. F. Tietze, J. Zinngrebe, D. A. Spiegl, and F. Stecker, *Heterocycles*, **74**, 473 (2007).
42. L. F. Tietze, D. A. Spiegl, F. Stecker, J. Major, C. Raith, and C. Grobe, *Chem. Eur. J.*, **14**, 8956 (2008).
43. L. F. Tietze, S. Jackenroll, D. Ganapathy, and J. R. Reiner, *Synlett*, **27**, 96 (2016).
44. B. M. Trost, H. C. Shen, and J. P. Surivet, *Angew. Chem. Int. Ed.*, **42**, 3943 (2003).
45. B. M. Trost, G. Dong, *Chem. Eur. J.*, **15**, 6910 (2009).
46. L. A. Arnold, R. Naasz, A. J. Minnaard, and B. L. Feringa, *J. Am. Chem. Soc.*, **123**, 5841 (2001). L. A. Arnold, R. Naasz, A. J. Minnaard, and B. L. Feringa, *J. Org. Chem.*, **67**, 7244 (2002).
47. T. Den Hartog, A. Rudolph, B. Macia, A. J. Minnaard, and B. L. Feringa, *J. Am. Chem. Soc.*, **132**, 14349 (2010).
48. M. Cheeseman and S. D. Bull, *Synlett*, 1119 (2006).

49. J. Deschamp, T. Hermant, and O. Riant, *Tetrahedron*, **68**, 3457 (2012).
50. L. Mangoni, M. Adinolfi, G. Laonigro, and R. Caputo, *Tetrahedron*, **28**, 611 (1972).
51. S. Hajra and S. Bar, *Chem. Commun.*, 3981 (2011).
52. N. Gommermann and P. Knochel, *Chem. Eur. J.*, **12**, 4380 (2006).
53. K. Alagiri, M. Furutachi, K. Yamatsugu, N. Kumagai, T. Watanabe, and M. Shibasaki, *J. Org. Chem.*, **78**, 4019 (2013).
54. K. V. Gothelf and K. A. Jorgensen, *Chem. Rev.*, **98**, 863 (1998).
55. A. Padwa and S. F. Hornbuckle, *Chem. Rev.*, **91**, 263 (1991).
56. A. Padwa, *Chem. Soc. Rev.*, **38**, 3072 (2009).
57. N. Shimada, T. Hanari, Y. Kurosaki, K. Takeda, M. Anada, H. Nambu, M. Shiro, and S. Hashimoto, *J. Org. Chem.*, **75**, 6039 (2010).
58. N. Shimada, T. Hanari, Y. Kurosaki, M. Anada, H. Nambu, and S. Hashimoto, *Tetrahedron Lett.*, **51**, 6572 (2010).
59. T. Arai, H. Sasai, K.-i. Aoe, K. Okamura, T. Date, and M. Shibasaki, *Angew. Chem. Int. Ed. Engl.*, **35**, 104 (1996).
60. K.-i. Yamada, T. Arai, H. Sasai, and M. Shibasaki, *J. Org. Chem.*, **63**, 3666 (1998).
61. K. Funabashi, H. Ratni, M. Kanai, and M. Shibasaki, *J. Am. Chem. Soc.*, **123**, 10784 (2001).
62. E. Ichikawa, M. Suzuki, K. Yabu, M. Albert, M. Kanai, and M. Shibasaki, *J. Am. Chem. Soc.*, **126**, 11808 (2004).
63. M. A. Sanner, T. A. Chappie, A. R. Dunaiskis, A. F. Fliri, K. A. Desai, S. H. Zorn, E. R. Jackson, C. G. Johnson, J. M. Morrone, P. A. Seymour, M. J. Majchrzak, W. S. Faraci, J. L. Collins, D. B. Duignan, C. C. Prete, J. S. Lee, and A. Trozzi, *Bioorg. Med. Chem. Lett.*, **8**, 725 (1998).
64. H. Nemoto, M. Yoshida, K. Fukumoto, and M. Ihara, *Tetrahedron Lett.*, **40**, 907 (1999).
65. M. Yoshida, M. A.-H. Ismail, H. Nemoto, and M. Ihara, *J. Chem. Soc., Perkin Trans. I*, 2629 (2000).
66. A. S. Paraskar and A. Sudalai, *Tetrahedron*, **62**, 4907 (2006).
67. S. Kobayashi, J. Kobayashi, R. Yazaki, and M. Ueno, *Chem. Asian J.*, **2**, 135 (2007).
68. A. Franchino, P. Jakubec, and D. J. Dixon, *Org. Biomol. Chem.*, **14**, 93 (2016).