Concentration of osmoregulated periplasmic glucans (OPGs) modulates the activation level of the RcsCD RcsB phosphorelay in the phytopathogen bacteria *Dickeya dadantii*

Sébastien Bontemps-Gallo, Edwige Madec, Jacqueline Dondeyne, Brigitte Delrue, Catherine Robbe-Masselot, Olivier Vidal, Anne-France Prouvost, Gilles Boussemart, Jean-Pierre Bohin, Jean-Marie Lacroix

► To cite this version:

Sébastien Bontemps-Gallo, Edwige Madec, Jacqueline Dondeyne, Brigitte Delrue, Catherine Robbe-Masselot, et al.. Concentration of osmoregulated periplasmic glucans (OPGs) modulates the activation level of the RcsCD RcsB phosphorelay in the phytopathogen bacteria *Dickeya dadantii*. *Environmental Microbiology*, 2013, *Plant-Microbe Interactions*, 15 (3), pp.881-894. 10.1111/1462-2920.12054 . hal-02133974

HAL Id: hal-02133974

<https://hal.science/hal-02133974>

Submitted on 11 Feb 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Concentration of Osmoregulated Periplasmic Glucans (OPGs) modulates the activation level of the RcsCD-RcsB phosphorelay in the phytopathogen bacteria *Dickeya dadantii*.

Sébastien Bontemps-Gallo^{1,2}, Edwige Madec^{1,2}, Jacqueline Dondeyne^{1,2}, Brigitte Delrue², Catherine Robbe-Masselot^{1,2}, Olivier Vidal^{1,2}, Anne-France Prouvost^{1,2,3}, Gilles Boussemart^{1,2,4}, Jean-Pierre Bohin^{1,2}, and Jean-Marie Lacroix^{*1,2}

¹*Unité de Glycobiologie Structurale et Fonctionnelle, UMR CNRS 8576, IFR 147, Université des Sciences et Technologies de Lille, 59655 Villeneuve d'Ascq, France*

²*Université Lille Nord de France, Lille, France*

³*Present address: VitamFero, 31 avenue Monge, 37200 Tours, France*

⁴*Present address: INRA Dijon, UMR 1229, 17 rue Sully, BP 86510, 21065 Dijon, France*

*corresponding author. Mailing address: Université des Sciences et Technologies de Lille, Unité de Glycobiologie Structurale et Fonctionnelle, UMR CNRS 8576 IFR147, 59655 Villeneuve d'Ascq Cedex, France. Fax: +33 3 20 43 65 55. Phone: +33 3 20 43 65 92. E-mail: jean-marie.lacroix@univ-lille1.fr.

Summary

Osmoregulated periplasmic glucans (OPGs) are general constituents of many Proteobacteria. Synthesis of these oligosaccharides is repressed by increased osmolarity of the medium. OPGs are important factors required for full virulence in many zoo- or phytopathogens including *Dickeya dadantii*. The phytopathogen enterobacterium *D. dadantii* causes soft-rot disease on a wide range of plant species. The total loss of virulence of *opg*-negative strains of *D. dadantii* is linked to the constitutive activation of the RcsCD RcsB phosphorelay highlighting relationship between this phosphorelay and OPGs. Here we show that OPGs control the RcsCD RcsB activation in a concentration dependent manner, are required for proper activation of this phosphorelay by medium osmolarity, and a high concentration of OPGs *in planta* is maintained to achieve the low level of activation of the RcsCD RcsB phosphorelay required for full virulence in *D. dadantii*.

1 **Introduction**

2 Osmoregulated periplasmic glucans (OPGs) are a family of periplasmic oligosaccharides
3 found in the envelope of most Proteobacteria. Their two common features are that glucose is
4 the sole constitutive sugar and their abundance in the periplasm increases as the osmolarity of
5 the medium decreases (Bohin *et al.*, 2006). Numerous studies indicate that these glucans
6 belong to the common virulence factors for zoopathogens and phytopathogens Proteobacteria,
7 since mutants devoid of OPGs exhibit a reduced or a non-virulent phenotype (Loubens *et al.*,
8 1993; Mahajan-Miklos *et al.*, 1999; Page *et al.*, 2001; Bhagwat *et al.*, 2009). In Enterobacteria,
9 the glucose backbone synthesis of OPGs is catalyzed by the *opgGH* operon products, and
10 *opgG* or *opgH* mutant strains are completely devoid of OPGs. OpgG and OpgH together
11 catalyze the synthesis of linear glucans containing 5 to 12 glucose units joined by β ,1-2
12 linkages and branched by β ,1-6 linkages (Bohin *et al.*, 2006).

13 *Dickeya dadantii*, an Enterobacterium, causes soft rot disease in a wide range of plant species,
14 including many economically important crops (Perombelon *et al.*, 2002). This opportunistic
15 pathogen devastates crops in storage facilities or in growing plants. In *D. dadantii*, *opgG* or
16 *opgH* mutant strains, completely devoid of OPGs, show a pleiotropic phenotype including
17 increased synthesis of exopolysaccharides, loss of motility, induction of a general stress
18 response and complete loss of virulence on potato tubers and chicory leaves. These
19 phenotypes were confirmed by a proteomic analysis and suggest that strains devoid of OPGs
20 are impaired in the perception of their environment (Page *et al.*, 2001; Bouchart *et al.*, 2007).
21 The isolation of revertant mutant strains in the *rscCDB* locus encoding a phosphorelay system,
22 restoring several of the wild-type phenotypes, indicated that OPGs are required in the
23 perception of environmental changes and that these glucans are needed to achieve a low level
24 of activation of the RcsCD RcsB signaling pathway (Bouchart *et al.*, 2010).

Two component systems (also called phosphorelay) are the main systems used by bacteria to sense and respond to variations of the environment. Under stimuli essentially unknown, a sensor histidine kinase autophosphorylates and transfers its phosphate group to its cognate response regulator which in turn regulates the expression of different target genes (Buelow *et al.*, 2010).

The RcsCD RcsB phosphorelay is restricted to enterobacteria. RcsC is the transmembrane kinase, RcsB the phosphorylated regulator, and phosphorylation occurs via the intermediate transmembrane protein RcsD. In several enterobacteria, it was shown that activation of the RcsCD RcsB system enhanced exopolysaccharides synthesis but also cell division, by increasing expression of the *ftsAZ* operon, (Carballès *et al.*, 1999; Bouchart *et al.*, 2010) and decreased virulence but also motility by decreasing expression of the *flhDC* operon (Francez-Charlot *et al.*, 2003; Garcia-Calderon *et al.*, 2005; Majdalani *et al.*, 2005; Bouchart *et al.*, 2010). The RcsCD RcsB system is induced by different stimuli including high osmolarity (Sledjeski *et al.*, 1996; Hagiwara *et al.*, 2003; Mouslim *et al.*, 2003). In addition, the RcsCD RcsB phosphorelay activity is enhanced by the outer membrane lipoprotein RcsF in *E. coli* (Majdalani *et al.*, 2005) and repressed by the transmembrane IgaA protein in *S. enterica* (Cano *et al.*, 2002).

Data presented in this paper indicate that OPGs concentration modulate activity of the RcsCD RcsB phosphorelay. OPGs are present in bacteria throughout the virulence process *in planta* and are required at a minimal concentration for full virulence.

Results

Periplasmic OPGs concentration depends on arabinose concentration in the NFB3835 strain

Previously, we showed that OPGs control the RcsCD RcsB phosphorelay in *D. dadantii* by preventing its activation and that total loss of OPGs (*opgG* strains) leads to a constitutive activation of this phosphorelay (Bouchart *et al.*, 2010). Thus, it was tempting to speculate that

variations of OPGs concentration affect activation of the RcsCD RcsB phosphorelay. To unravel the relationship between OPGs and the RcsCD RcsB phosphorelay, we decided to test the impact of variation of OPGs concentration on the activation of the RcsCD RcsB phosphorelay.. In *Escherichia coli*, the *araBAD* operon is known to be tightly regulated by the concentration of L-arabinose in the medium by the AraC protein (Guzman *et al.*, 1995). OPGs concentration synthesized by the NFB3835 strain (*araC-P_{BAD}opgGH*, see experimental procedures) was measured in M63 medium containing various L-arabinose concentrations ranging from 0 g l⁻¹ to 1 g l⁻¹ (Fig. 1). No OPGs could be detected without L-arabinose. OPGs concentration increased in cells as the L-arabinose concentration increased in the medium to reach a maximal concentration of 20 µg mg⁻¹ of proteins when 1 g l⁻¹ arabinose was added to the medium. OPGs concentration remained the same when higher L-arabinose concentrations (2 g l⁻¹ or 5 g l⁻¹) were added to the medium (data not shown). This strain can be used to test relationship between OPGs and the RcsCD RcsB phosphorelay, since OPGs concentration of this strain depends on L-arabinose concentration of the medium up to 1 g l⁻¹.

Increasing concentrations of OPGs decrease the activation of the RcsCD RcsB phosphorelay

Relationship between OPGs and the RcsCD RcsB phosphorelay was followed by expression of transcriptional fusions derived from two operons known to be regulated in an opposite way by the RcsCD RcsB phosphorelay in *D. dadantii* (Bouchart *et al.*, 2010). The RcsCD RcsB system regulates motility by down-regulation of the *flhDC* master operon encoding activators required for expression of the flagellar apparatus genes. The *ftsAZ* operon is needed for cell division. One of the promoters regulating this operon is up-regulated by the RcsCD RcsB phosphorelay and it was cloned in the *ftsA-uidA* fusion (Bouchart *et al.*, 2010). The *flhD-uidA* and the *ftsA-uidA* fusions were transduced into the NFB3835 strain and expressions of both fusions were measured after growth in M63 medium with increasing concentrations of L-arabinose. Motility was evaluated in the NFB3835 strain on M63 medium swimming plates

1 with increasing concentrations of L-arabinose. As a control, the respective motility levels of
2 the EC3937 (wild-type) and the NFB3723 (*opgG*) strains were measured in M63 medium
3 with or without various concentrations of L-arabinose. Motility levels remained constant
4 whatever the concentration of L-arabinose for both strains and were 2.2 fold higher for the
5 wild-type strain as compared to the *opgG* mutant strain (Fig. 2A). Motility of the NFB3835
6 strain increased as the concentration of L-arabinose increased (Fig. 2A). The lower motility
7 level was observed in the NFB3835 strain grown in M63 medium and was similar to the
8 motility level observed in the NFB3723 (*opgG*) strain grown in the same medium. The higher
9 motility level was observed in the NFB3835 strain grown in M63 medium with 1 g l⁻¹
10 arabinose added and was similar to the motility level observed in the EC3937 wild-type strain
11 grown in the same medium (Fig. 2A). Expression of the *flhD-uidA* fusion was measured after
12 growth of the NFB7300 strain in M63 medium containing the same increasing concentrations
13 of L-arabinose (Fig. 2B). As a control, the respective β -glucuronidase activity levels of the
14 NFB3800 (wild-type) and the NFB7301 (*opgG*) strains were measured in M63 medium with
15 or without various concentrations of L-arabinose. Glucuronidase activity remained constant
16 for both strains whatever the concentration of L-arabinose and was 2.3 fold higher in the wild-
17 type strain as compared to the *opgG* mutant strain (Fig. 2B). The lower level of β -
18 glucuronidase expression was measured after growth of the NFB7300 in M63 medium
19 without L-arabinose and was similar to the expression level measured in the *opgG* strain
20 (NFB7301) harboring the same fusion. The highest level of expression was measured after
21 growth in M63 medium containing 1 g l⁻¹ of L-arabinose and was similar to the expression
22 level observed in the wild-type strain harboring the same fusion (NFB3800). Increase in the
23 motility was correlated with an increased expression of the *flhD-uidA* fusion. These data
24 indicate that the concentration of OPGs modulates motility in *D. dadantii*. Expression of the
25 *ftsA-uidA* fusion was measured in the same conditions (Fig. 2C). As a control, the respective

1 glucuronidase activities levels of the NFB3809 (wild-type) and the NFB7274 (*opgG*) strains
2 were measured in M63 medium with or without various concentrations of L-arabinose.
3 Glucuronidase activity remained constant for both strains whatever the concentration of L-
4 arabinose and was 6 fold higher in the *opgG* strain as compared to the wild-type strain. The
5 lower level of expression was measured after growth of the NFB3837 strain in M63 medium
6 containing 1 g l⁻¹ L-arabinose and was similar to the expression level measured in the wild-
7 type strain (NFB3809) harboring the same fusion. The highest level was measured after
8 growth in M63 medium without L-arabinose and was similar to the expression level observed
9 in the *opgG* strain harboring the same fusion (NFB7274). These data indicate that
10 concentration of OPGs modulates *ftsA* expression in *D. dadantii*. To ensure that the
11 regulation observed occurs through the RcsCD RcsB phosphorelay, an *rscB* or an *rscC* null
12 mutations were introduced into NFB3835, NFB3837 and NFB7300 (NFB7368, NFB7371 and
13 NFB7369 respectively and NFB7400, NFB7406, NFB7405, respectively). Whatever the *rsc*
14 genetic context, motility and glucuronidase activities of both fusions remained constant
15 whatever the L-arabinose concentration and similar to the one observed for the wild-type
16 strain harboring the same fusions, respectively (Fig 2A, B and C). A recent study revealed
17 that the unphosphorylated form of RcsB is required for proper regulation of biofilm synthesis
18 (Latasa et al. 2012). Thus, the observed phenotypes in the *rscB* null mutant strains could be
19 the result of the lack of the RcsB protein itself rather than the loss of phosphorylation. An
20 *rscCDB*_{D56N} locus encoding an unphosphorylatable form of *rscB* was constructed (see
21 experimental procedures) and introduced into the *rscB* derivative strains NFB7368, NFB7369
22 and NFB7371 (NFB7401, NFB7403, NFB7402, respectively). Results were similar to those
23 observed with *rscB* null mutant strains for motility and glucuronidase activities (Fig. 2A, B
24 and C). To ensure that the RcsB cytoplasmic concentration was independent of the L-
25 arabinose concentration, a western blot analysis was performed in M63 medium with various

1 concentrations of L-arabinose added in the EC3937, NFB3723, NFB3835, NFB7112 and
2 NFB3682 strains. Similar amount of the RcsB protein was observed whatever the strain and
3 L-arabinose concentration tested (data not shown) indicating that regulation occurs through
4 variation of the phosphorylation level of the RcsB transcriptional regulator. Taken together,
5 these results indicate that OPGs control the RcsCD RcsB phosphorelay activity in a
6 concentration dependent manner.

7 *A minimal concentration of OPGs is required for D. dadantii throughout the virulence*
8 *process*

9 The RcsCD RcsB phosphorelay activity is severely repressed during virulence (Garcia-
10 Calderon *et al.*, 2005) and OPGs are required for this repression (Bouchart *et al.*, 2010). Thus,
11 OPGs concentration must affect virulence. To test the impact of OPGs concentration on
12 virulence of *D. dadantii* during pathogenesis, NFB3835 was grown in M63 medium with
13 various concentrations of L-arabinose prior to inoculation. Bacteria were centrifuged and
14 washed twice with M63 medium without a carbon source to remove arabinose to avoid OPG
15 synthesis *in planta*. Bacteria (10^7) were inoculated on chicory leaves and virulence was
16 monitored 24 h and 48 h later (Fig. 3). When no arabinose was provided to NFB3835
17 bacterial cells before inoculation (fig. 3), no maceration of chicory leaves was detected. This
18 indicates that no free arabinose is available *in planta*. A moderate maceration of chicory
19 leaves was observed at a concentration of 0.01 g l^{-1} of L-arabinose while maceration similar to
20 the wild-type strain was observed at an L-arabinose concentration of 1 g l^{-1} . Thus, because no
21 OPG could be synthesized *in planta*, severity of the disease depends on OPGs concentration
22 at time 0 and stops when OPGs concentration in bacteria falls down below the lower
23 concentration required for virulence because of dilution of OPGs during growth of bacteria.
24 This concentration could be estimated around 0.01 g l^{-1} of arabinose added (Fig. 3). This L-
25 arabinose concentration corresponds to $10 \text{ }\mu\text{g}$ of OPGs per mg of protein (Fig. 1). When

1 arabinose (1 g l^{-1}) was added to the bacterial inoculum on chicory leaves or 24 h after
2 inoculation, no virulence was observed when bacteria were grown in 63 medium before
3 inoculation while full virulence was observed whatever the arabinose concentration provided
4 in the 63 medium before inoculation (data not shown). These data indicate that a minimal
5 level of OPGs is required during all the pathogenesis process for full virulence of *D. dadantii*
6 and suggest that *D. dadantii* is able to adjust its OPG concentration *in planta*.

7 *D. dadantii* synthesizes a high concentration of OPGs in planta

8 To test this hypothesis, the concentration of OPGs was determined after growth of the wild-
9 type strain *in planta*. The EC3937 strain was grown in M63 medium and 10^7 bacteria were
10 inoculated on several chicory leaves. OPGs were extracted and purified from bacteria isolated
11 from culture in M63 medium before inoculation (0 h), and 24 h and 48 h after growth *in*
12 *planta*. For each time, CFU were determined. After 24 h of maceration, OPGs concentration
13 increased more than twofold as compared to the concentration observed at 0 h and remained
14 constant after 48 h (Fig. 4A). Because OPGs were purified on P4 gel filtration column, one
15 can argue that OPGs extracted from bacteria grown *in planta* were contaminated by
16 oligosaccharides of similar weight resulting from degradation of plant polysaccharides by
17 hydrolytic enzymes secreted by *D. dadantii* during infection. MALDI TOF mass spectrometry
18 was used to determine the sizes of the OPG backbones. Spectrum of OPGs extracted from
19 bacteria grown in M63 medium was used as a control and compared with spectra of OPGs
20 extracted from bacteria grown *in planta* after 24 h and 48 h of infection. As shown in Fig. 4B,
21 no difference in the sizes of OPGs was observed, whatever growth conditions. Each of them
22 was composed of 4 to 11 residues of glucose. This was clearly seen in mass spectra with the 8
23 $[M+Na]^+$ molecular ions at m/z 689, 851, 1013, 1175, 1337, 1499, 1661 and 1823 recovered
24 in each spectrum. For several molecular ions, an increment of 42Da was observed and
25 corresponds to the substitution of OPGs by acetyl residues, a characteristic feature of OPGs

1 isolated from *D. dadantii* (Cogez *et al.*, 2001). Thus, oligosaccharides extracted from bacteria
2 grown *in planta* were only constituted of OPGs. When bacteria were grown in media of
3 higher (0.1 or 0.2M NaCl added) osmolarity or in medium of lower osmolarity (half-strength
4 M63), no difference in the severity of the disease nor in the apparition of the symptoms were
5 observed (data not shown). Taken together, these results indicate that *in planta*, *D. dadantii*
6 adjust its OPG concentration to a higher level than the minimal concentration required to
7 reach a sufficient repression level of the RcsCD RcsB phosphorelay needed for full virulence
8 of *D. dadantii*.

9 *The RcsCD RcsB phosphorelay exhibits a low activity level during virulence*

10 This high concentration of OPGs must be related to a low activity level of the RcsCD RcsB
11 phosphorelay required for virulence. Expression of the *flhD-uidA* and *ftsA-uidA* fusions was
12 measured just before inoculation (0 h) and 24 h and 48 h after inoculation *in planta* in wild-
13 type and *rscB* genetic background (Fig. 4C). The *ftsA-uidA* fusion was expressed at a similar
14 and very low level while the *flhD-uidA* fusion was expressed at a similar and high level
15 during all the virulence process, in both genetic backgrounds and at each time tested. A
16 western blot analysis was performed with bacteria grown *in planta*. Similar amount of the
17 RcsB protein was observed after 0 h, 24 h and 48 h of maceration (data not shown). Thus,
18 these results indicate that the RcsCD RcsB phosphorelay activity is severely repressed during
19 the virulence process in *D. dadantii*.

20 *OPGs are one of the actors required for modulation of the RcsCD RcsB phosphorelay activity* 21 *by medium osmolarity*

22 OPGs synthesis and RcsCD RcsB activation are regulated by medium osmolarity in an
23 opposite way: OPG concentration decrease (Cogez *et al.*, 2001) while the RcsCD RcsB
24 phosphorelay activation occurs as the medium osmolarity increases. Thus, the RcsCD RcsB
25 activation could be modulated by osmolarity via variation of OPGs concentration. To test this

1 hypothesis, motility was evaluated on swimming plates and the *flhD-uidA* and the *ftsA-uidA*
 2 transcriptional fusion activities were determined in different genetic backgrounds at various
 3 osmolarities in M63 medium containing as a carbon source. In the wild-type strain (EC3937),
 4 motility remained constant in half strength M63 medium and in M63 medium then decreased
 5 as the osmolarity of the medium increased (Fig 5A). In the different mutant strains, motility
 6 levels remained unaffected whatever the osmolarity: as expected, a loss of motility was
 7 observed for the *opgG* strain (NFB3723) and motility similar to the wild-type strain was
 8 observed for the *rscB* (NFB7279), the *rscB_{D56N}* (NFB7112) and the *rscC* (NFB3682) strains
 9 (Fig. 5A). The expression of the *flhD-uidA* fusion harbored by the wild-type strain (NFB3800)
 10 remained constant in half strength M63 medium and in M63 medium then decreased as the
 11 osmolarity of the medium increased (Fig. 6B). As expected, expression of the *flhD-uidA*
 12 fusion in the *opgG* (NFB7301) strain remained low and constant whatever the osmolarity
 13 while the expression level of the *flhD-uidA* fusion harbored by the *rscB* (NFB7202), the
 14 *rscB_{D56N}* (NFB7399) and the *rscC* (NFB7128) strains was similar to the wild-type strain
 15 grown in M63 medium and remained unaffected when osmolarity was increased (Fig. 5B).
 16 These results indicate that repression of motility by medium osmolarity depends on variation
 17 of concentration of OPGs via activation of the RcsCD RcsB phosphorelay. For the *ftsA-uidA*
 18 transcriptional fusion activity (Fig. 7A), a low level of expression in the *rscB* (NFB7205), the
 19 *rscB_{D56N}* (NFB7408) and the *rscC* (NFB7127) strains was observed, whatever the osmolarity.
 20 Surprisingly, no increase in the expression of the *ftsA-uidA* fusion was observed in the wild-
 21 type strain (NFB3809). In the *opgG* strain (NFB7274), expression of the *ftsA-uidA* fusion was
 22 strongly increased as compared to the wild type strain and a 4 fold decrease was observed
 23 between the medium of lower osmolarity (half strength M63 medium) as compared to the
 24 medium of higher osmolarity (M63 full strength medium with 0.2 M NaCl added). This
 25 suggests that in addition to OPGs, additional factors are required for regulation of the *ftsAZ*

operon by the RcsCD RcsB phosphorelay. As a control, OPGs concentrations were determined (Fig. 6B). As previously shown (Cogez *et al.*, 2001), OPGs concentration decreased as the osmolarity increased from 36 µg of OPGs per mg of protein in half strength M63 medium to 6 µg of OPGs per mg of protein in M63 medium with 0.2 M NaCl added. A western blot analysis was performed in M63 medium with (0.2 M) or without NaCl added in an *opgG*, *rscC*, *rscB_{D56N}* and in the wild-type strains (Fig. 3C). Similar amounts of the RcsB protein were observed whatever the strain and the osmolarity tested (data not shown). These results indicate that in *D. dadantii*, variation of OPGs concentration is required for proper activation of the RcsCD RcsB phosphorelay by osmolarity, but osmoregulation is a complex multifactor phenomenon (Wood, 1999) where OPGs are needed but are only one of the actors required for regulation of the RcsCD RcsB regulon.

Discussion

OPGs are known to be an important virulence factor for many Gram-negative bacteria (Bohin, 2006) and we show in a previous paper (Bouchart *et al.*, 2010) that OPGs control the RcsCD RcsB phosphorelay. In this paper we demonstrate that in addition concentration of OPGs in the periplasm modulates activation level of the RcsCD RcsB phosphorelay in *D. dadantii*.

In *D. dadantii*, motility and the *flhDC* operon are repressed by the activation of the RcsCD RcsB phosphorelay while the *ftsA* gene is induced (Bouchart *et al.*, 2010) and reflect the RcsCD RcsB phosphorelay activation level. When OPG concentration depends on L-arabinose concentration of the medium, we showed that motility and expression of both fusions genes tested depends on L-arabinose in an RcsB dependent manner and that activation of the RcsCD RcsB phosphorelay increase as the concentration of OPGs decrease. OPGs control the RcsCD RcsB activity in a concentration dependent manner. One of the stimuli activating the RcsCD RcsB phosphorelay is osmolarity increased (Majdalani *et al.*, 2005) while in the same conditions, OPGs concentration decreased (Cogez *et al.*, 2001; Bohin *et al.*,

2006). Decreased motility and expression of the *flhD-uidA* gene fusion observed when osmolarity of the medium increased are in agreement with an increased activation of the RcsCD RcsB phosphorelay and this activation depends both on a decreased OPG concentration and on RcsB (Fig. 5 and Fig. 6). No increase in the *ftsA-uidA* expression was observed when osmolarity increase while expression of this gene is known to be enhanced under activation of the RcsCD RcsB phosphorelay. One explanation is the presence, in *D. dadantii*, of a still unknown co-regulator of RcsB affecting interaction between RcsB and its DNA regulatory sequences. This hypothesis was strengthened by the decrease expression of the *ftsA-uidA* fusion gene in the *opgG* mutant strain since without additional regulatory component, the constitutive activation of the RcsCD RcsB phosphorelay would lead to a constant and high level of expression of this fusion. In *E. coli*, RcsA, GadE or BglJ form heterodimer with RcsB and co-regulate different subset of genes of the RcsCD RcsB regulon (Majdalani *et al.*, 2005; Krin *et al.*, 2010; Stratmann *et al.*, 2012) but we did not find ortholog of these genes in the *D. dadantii* genome.

Variation in motility and in expression of target fusion genes is directly related to the modulation of the RcsB phosphorylation level itself by RcsC since i/ no variation in the expression of both target genes was observed whatever the *rscB* or *rscC* mutant strains mutant strains and ii/ no difference in RcsB amount was observed in western blot analysis (see Fig. 3) whatever the growth conditions (data not shown). No difference in expression of the *rscB* gene was also reported in *Erwinia amylovora* grown either in minimal medium or *in planta* (Wang *et al.*, 2012). Thus, the activation level of the RcsC sensor and subsequently the phosphorylation level of RcsB depend on periplasmic concentration of OPG.

A minimal concentration of OPGs is required for initiating virulence (Fig. 3). This concentration could be estimated at 10 µg of OPGs per mg of protein (Fig. 1) and was measured in M63 medium containing 0.01 g L⁻¹ arabinose corresponding to the amount found

1 in M63 medium supplemented with 0.15 M NaCl (fig. 6). This concentration of OPGs may
2 correspond to the higher level of phosphorylated RcsB allowing virulence. Full virulence was
3 observed until 20 µg per mg of proteins and was measured in M63 medium containing 1 g L⁻¹
4 arabinose or in M63 medium (fig. 6). No difference was observed in virulence (timing or
5 severity of the disease) whatever the osmolarity of the medium before inoculation *in planta*.
6 Our data indicate that *D. dadantii* may adjust its concentration of OPGs within the 24 first
7 hours of infection. This concentration of OPGs was two times higher than the minimum
8 required to ensure a low level of phosphorylated RcsB (Fig. 4). This high OPGs concentration
9 may be explained by the low osmolarity of the plant apoplast even if this low osmolarity
10 varies depending on the plant tissue (Boyer *et al.*, 2001). Another explanation is that high
11 OPGs are required for other cellular processes (Delangle *et al.*, 2007) during virulence and
12 that additional regulation enhances OPGs synthesis. In *E. coli*, the *opgGH* operon is part of
13 the envelope stress sigma E factor regulon (Dartigalongue *et al.*, 2001). In *Yersinia*
14 *enterocolitica*, it was reported that the *opgGH* operon was induced within the host (Young
15 and Miller, 1997). Thus, because OPGs are needed for virulence in several mammalian
16 pathogenic enterobacteria, it would be of interest to measure OPG concentration in these
17 bacterial species since in the gut high osmolarity environmental condition, high concentration
18 of OPGs cannot be achieved by osmoregulation.

20 **Experimental procedures**

21 *Bacterial strains, media, antibiotics and growth condition*

22 Bacterial strains and plasmids are described in Table 1. Bacteria were grown at 30°C (*D.*
23 *dadantii*) or 37°C (*E. coli*) in Lysogeny broth (LB) (Bertani, 2004), or in minimal medium
24 M63 supplemented with glycerol as a carbon source at a concentration of 2 g l⁻¹ (Miller, 1992).
25 Solid media were obtained by adding agar at 15 g l⁻¹. Osmolarity of the M63 medium

increased by adding 0.1 M or 0.2 M NaCl or decreased by a two-fold dilution. When necessary L(+)-arabinose (Calbiochem) was added in the M63 medium at indicated concentrations. X-Gluc (5-bromo-4-chloro-3-indolyl- β -D-glucuronide) was used in solid media at a concentration of 40 $\mu\text{g ml}^{-1}$. Motility was tested on M63 agar plates containing 4 g l^{-1} agar and swim diameters were measured after 30 h of incubation.

Antibiotics were used at the following concentrations: gentamicin at 3 $\mu\text{g ml}^{-1}$ (*E. coli*) or 2 $\mu\text{g.ml}^{-1}$ (*D. dadantii*), chloramphenicol and tetracycline at 25 $\mu\text{g ml}^{-1}$ (*E. coli*) or 12.5 $\mu\text{g ml}^{-1}$ (*D. dadantii*), ampicillin and kanamycin at 50 $\mu\text{g ml}^{-1}$ (*E. coli*) or 25 $\mu\text{g ml}^{-1}$ (*D. dadantii*), streptomycin and spectinomycin at 100 $\mu\text{g ml}^{-1}$ (*E. coli*) or 50 $\mu\text{g ml}^{-1}$ (*D. dadantii*).

Cloning of the opgGH operon under control of the araBAD promoter

Plasmids and primers designed for PCR are listed in Table 1. The beginning of the operon *opgGH* (without the *opgGH* promoter) was amplified by PCR (*opgGH*nhelF and *opgGH*kpnIr primers) from pNFW32 (Page *et al.*, 2001). This PCR product was digested by NheI and KpnI and cloned into pBAD18 digested by the same enzymes (pNFW208). pNFW32 was digested by EcoRI and KpnI and the DNA fragment containing the remaining of the *opgGH* operon was cloned into pNFW208 digested by the same enzymes. The resulting plasmid (pNFW212) contains the *opgGH* operon controlled by the *araBAD* promoter ($P_{BADopgGH}$). Plasmid pNFW212 was digested by AclI and Bstz17I and the DNA fragment containing the *araC* gene and $P_{BADopgGH}$ operon was inserted in the SmaI site of pUC18Not. The resulting plasmid (pNFW219) was digested by NotI and the DNA fragment containing the *araC* gene and $P_{BADopgGH}$ operon was inserted into the NotI site of pUTminiTn5-Spe. The resulting plasmid (pNFW233) was introduced in *D. dadantii* by conjugation to allow integration, after transposition of the miniTn5, of a single ectopic copy of the *araC* gene and $P_{BADopgGH}$ operon.

Construction of the mutations

The *rcsB::Gm* allele was constructed by excision of the *Cml^r* cassette from the pNFW397 (Bouchart *et al.*, 2010) digested by *HpaI* and replaced by the *Gm^r* cassette digested by the same restriction enzyme after PCR amplification (*GmhpalF* and *GmhpalR*, Table 1). The resulting plasmid (pNFW402) was introduced in *D. dadantii* cells by electroporation.

The *rcsB_{D56N}* allele was constructed by SOE (splicing by overlap extension) PCR. The upstream region was amplified using primers *RcsBDN3* and *RcsBDNsoe1* (table 1), whereas the downstream region was amplified using primers *RcsBDNsoe2* and *RcsBDN4* (Table 1). The two PCR products were spliced together using the flanking primers *RcsBDN3* and *RcsBDN4*. The mutagenesis PCR product was purified, digested with *HpaI* and *AscI*, cloned into pNFW257 digested by the same enzymes (pNFW374) and sequenced to confirm mutation. The *NotI* fragment from pNFW374, containing the *rcsCB_{D56N}D* locus, was subcloned into pUTmini-Tn5-Tet or pUTmini-Tn5-Spe digested by *NotI*. The resulting plasmids (pNFW379 and pNFW380 respectively) were introduced into *D. dadantii* cells by conjugation.

Transduction, conjugation and transformation.

Transformation of *E. coli* cells was carried out by the rubidium chloride technique (Miller, 1992). Plasmids were introduced in *D. dadantii* by electroporation (Sambrook *et al.*, 1989) or conjugation (Miller, 1992). The insertions were integrated into the *D. dadantii* chromosome by marker exchange recombination in the presence of the appropriate antibiotic, after successive cultures in low phosphate medium (Roeder *et al.*, 1985). Transduction with phage ϕ EC2 was carried out according to Resibois *et al.* (Resibois *et al.*, 1984).

Transposon mutagenesis

To allow integration, after transposition of the miniTn5, of a single ectopic copy of the *araC* gene and *P_{BAD}opgGH* operon or the *rcsCDB_{D56N}* locus, transposon mutagenesis by mini-Tn5 was performed as described in (Bouchart *et al.*, 2010). Briefly, after conjugation between an *E.*

coli strain harboring the plasmid carrying the appropriate mini-Tn5 and a *D. dadantii* strain, mutants were selected on M63 plates containing sucrose as a unique carbon source and the required antibiotic.

Determination of enzyme activities

β -glucuronidase assays were performed on crude extracts obtained from bacteria disrupted by sonication 2x 20 s (Sonifier cell disruptor B-30, Branson, 70% duty cycle, 7 microtip limit, Hold time, continuous, appropriate probe). β -glucuronidase activity was determined by spectrometric monitoring the hydrolysis of PNPU (4-nitrophenyl- β -D-glucuronide) at 405 nm. The protein concentration was determined by the Bradford assay with bovine serum albumin as standard (Bradford, 1976).

OPGs analysis

Bacteria were grown *in planta* or in M63 medium (200 ml) supplemented either with various L-arabinose concentrations or with various concentrations of NaCl until mid-log phase. Bacteria were collected by centrifugation at 4°C for 15 min at 8,000 $\times$ g. Cell pellets were resuspended in 20 ml of distilled water and extracted with 5% trichloroacetic acid (TCA). The TCA extracts were neutralized with ammonium hydroxide 10% and concentrated by rotary evaporation. The resulting material (2 ml) was then fractionated by gel filtration on a Bio-Gel P-4 (Bio-Rad) column (55 by 1.6 cm) equilibrated with 0.5% acetic acid. The column was eluted in the same buffer at a flow rate of 15 ml h⁻¹, and fractions of 1.5 ml were collected. Presence of sugar in each fraction was determined colorimetrically by the anthrone reagent procedure. Fractions containing OPGs were pooled and content was determined by the same procedure (Spiro, 1966). For mass spectrometry analysis, OPGs were subsequently desalted in water on a Bio-Gel P-2 (Bio-Rad) column (70 by 1.6 cm) and fractions containing OPGs were lyophilized.

Preparation of polyclonal antibodies against RcsB

1 A DNA fragment encoding the *rcsB* gene of *D. dadantii* was amplified by PCR using the
2 primers *rscBacf* and *rscBacr*. The PCR product was cloned into a His₆ tag expression vector,
3 pET100/D-Topo® (Invitrogen Life Technologies) to give pNFW410. The resulting His-
4 tagged RcsB was expressed in *E. coli* BL21(DE3) and the protein was purified by affinity
5 chromatography according to the manufacturer's procedure (Ni-nitrilotriacetic acid [NTA]
6 agarose; Qiagen). The purified RcsB was used to immunize rabbit.

7 *Western blot hybridization*

8 Strains of *D. dadantii* were grown at 30°C to mid-log phase in M63 or after various time of
9 infection in planta. Bacteria were collected by centrifugation and immediately lysed with
10 SDS-PAGE buffer (2% SDS, 2.5% DTT, 25 mM Tris-HCl pH 7.5, 15% glycerol), heated to
11 100°C for 5 min and loaded onto a 10% SDS-polyacrylamide gel (34). Proteins from the gel
12 were transferred onto a nitrocellulose membrane (Whatman). Western blotting was performed
13 using standard protocols with the rabbit anti-RcsB polyclonal antibodies at a dilution of
14 1:1000 and anti-rabbit secondary antibody coupled to horseradish peroxidase (dilution
15 1:10,000). Blots were imaged by chemiluminescent detection (ECL kit, GE healthcare
16 Pharmacia Biotech).

17 *Pathogenicity test*

18 Chicory leaves were inoculated as previously described (Delangle *et al.*, 2007; Bouchart *et al.*,
19 2010).

20 *Bacteria isolation from infected chicory leaves*

21 Infected chicory leaves were introduced into tubes (one leave per tube) containing 20 mL of
22 minimal medium 63 without carbon source added. Bacteria were suspended in the medium by
23 vigorous shaking of the tubes during 15 min. Leaves were then carefully removed from the
24 tube. The medium was filtered through a paper filter (previously sterilized) to separate few
25 macerated tissue release during the shaking step from bacteria (no more than 6% bacteria

were lost during this step). Bacteria were then collected by centrifugation at 4°C for 15 min at 8,000 × g. Cell pellets were resuspended in 20 mL of minimal medium 63 for OPGs analysis or in 2 mL of the same medium for β-glucuronidase assays. Before analysis or assays, an aliquot of 0.1 mL of each bacterial suspension was taken up, diluted and counted. A total of 10⁸ to 10¹⁰ CFU were recovered per chicory leave depending on the incubation time.

Matrix-assisted laser desorption/ionization mass spectrometry (MALDI-MS)

All mass spectra were acquired on a Voyager Elite (DE-STR) reflectron time-of-flight (TOF) mass spectrometer (Perseptive Biosystems, Framingham, MA), equipped with a pulsed nitrogen laser (337 nm) and a gridless delayed extraction ion source. Oligosaccharide samples were analyzed in delayed extraction mode using an accelerating voltage of 20 kV, a pulse delay time of 200 ns and a grid voltage of 66%. Detector bias gating was used to reduce the ion current for masses below 500 Da. Between 100 and 200 scans were averaged for each mass spectrum. Oligosaccharide alditols were co-crystallized with 2,5-dihydroxybenzoic acid (DHB) as matrix [10 mg ml⁻¹ of DHB in methanol/water (50/50) containing 0.1% Trifluoro Acetic acid (TFA)]. For all measurements, the "dried droplet" preparation technique was used. Typically, 1 µl of the matrix was mixed on-target with 1 µl of water-dissolved oligosaccharides and allowed to dry under an air stream. They were analyzed in positive ion mode.

Acknowledgements

This work was supported by grants from the Centre National de la Recherche Scientifique (CNRS) and from the Ministère de l'Éducation Nationale et de la Recherche.

References

Bertani, G. (2004) Lysogeny at Mid-Twentieth Century: P1, P2, and Other Experimental Systems. J Bacteriol 186: 595-600.

1 Bhagwat, A.A., Jun, W., Liu, L., Kannan, P., Dharne, M., Pheh, B., et al. (2009)
2 Osmoregulated periplasmic glucans of *Salmonella enterica* serovar Typhimurium are required
3 for optimal virulence in mice. *Microbiology* 155: 229-237.

4 Bohin, J.P., and Lacroix, J.M. (2006) Osmoregulation in the periplasm. In *The periplasm*.
5 Ehrmann, M. (ed) Washington, DC: American Society for Microbiology, pp 325-341.

6 Bouchart, F., Delangle, A., Lemoine, J., Bohin, J.P., and J.M. (2007) Proteomic analysis of a
7 non virulent mutant of the phytopathogenic bacterium *Erwinia chrysanthemi* deficient in
8 osmoregulated periplasmic glucans: change in protein expression is not restricted to the
9 envelope, but affects general metabolism. *Microbiology* 153: 760-767.

10 Bouchart, F., Boussemart G., Prouvost A.F., Cogez V., Madec E., Vidal O., et al. (2010) The
11 virulence of a *Dickeya dadantii* 3937 mutant devoid of osmoregulated periplasmic glucans
12 (OPGs) is restored by inactivation of the RcsCD RcsB phosphorelay. *J Bacteriol* 192: 3484–
13 3490.

14 Boyer, J.S. (2001) Growth-induced water potentials originate from wall yielding during
15 growth. *J Exp Bot* 52: 1483-1488.

16 Bradford, M.M. (1976) A rapid and sensitive method for the quantitation of microgram
17 quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72: 248-254.

18 Buelow, D.R., and Raivio, R. (2010) Three (and more) component regulatory systems –
19 auxiliary regulators of bacterial histidine kinases. *Mol Microbiol* 75: 547-566.

20 Cano, D.A., Dominguez-Bernal, G., Tierrez, A., Garcia-Del Portillo, F., and Casadesus, J.
21 (2002) Regulation of capsule synthesis and cell motility in *Salmonella enterica* by the
22 essential gene *igaA*. *Genetics* 162: 1513-1523.

23 Carballès, F., Bertrand, C., Bouché, J.P., and Cam, K. (1999) Regulation of *Escherichia coli*
24 cell division genes *ftsA* and *ftsZ* by the two-component system RcsC-RcsB. *Mol Microbiol* 34:
25 442-450.

26 Cogez, V., Talaga, P., Lemoine, J., and Bohin, J.P. (2001) Osmoregulated periplasmic
27 glucans of *Erwinia chrysanthemi*. *J bacteriol* 183: 3127-3133.

28 Dartigalongue, C., Missiakas, D., and Raina, S. (2001) Characterization of the *Escherichia*
29 *coli* sigma E regulon. *J biol chem* 276: 20866-20875.

30 De Lorenzo, V., and Timmis, K.N. (1994) Analysis and construction of stable phenotypes in
31 Gram-negative bacteria with Tn5- and Tn10-derived minitransposons. *Methods Enzymol* 235:
32 386-405.

De Lorenzo, V., Herrero, M., Jakubzik, U., and Timmis, K.N. (1990) Mini-Tn5 transposon derivatives for insertion mutagenesis, promoter probing, and chromosomal insertion of cloned DNA in gram-negative eubacteria. *J Bacteriol* 172: 6568-6572.

Delangle, A., Prouvost, A.F., Cogez, V., Bohin, J.P., Lacroix, J.M., and Hugouvieux-Cotte-Pattat, N. (2007) Characterization of the *Erwinia chrysanthemi* gan locus involved in galactan catabolism. *J Bacteriol* 189: 7053-7061.

Francez-Charlot, A., Laugel, B., Van Gemert, A., Dubarry, N., Wiorowski, F., Castanié-Cornet, M.P., et al. (2003) RcsCDB His-Asp phosphorelay system negatively regulates the *flhDC* operon in *Escherichia coli*. *Mol Microbiol* 49: 823-832.

Garcia-Calderon, C.B., Garcia-Quintanilla, M., Casadesus, J., and Ramos-Morales, F. (2005) Virulence attenuation in *Salmonella enterica rcsC* mutants with constitutive activation of the Rcs system. *Microbiology* 151: 579-588.

Guzman, L.M., Belin, D., Carson, M.J., and Beckwith, J. (1995) Tight regulation, modulation, and high-level expression by vectors containing the arabinose PBAD promoter. *J Bacteriol* 177: 4121-4130.

Hagiwara, D., Sugiura, M., Oshima, T., Mori, H., Aiba, H., Yamashino, T., and Mizuno, T. (2003) Genome-wide analyses revealing a signaling network of the RcsC-YojN-RcsB phosphorelay system in *Escherichia coli*. *J bacteriol* 185: 5735-5746.

Krin, E., Danchin, A., and Soutourina, O. (2010) RcsB plays a central role in H-NS-dependent regulation of motility and acid stress resistance in *Escherichia coli*. *Res microbiol* 161: 363-371.

Latasa, C., Garcia, B., Echeverz, M., Toledo-Arana, A., Valle, J., Campoy, S., et al. (2012) *Salmonella* biofilm development depends on the phosphorylation status of RcsB. *J Bacteriol* 194: 3708-3722.

Loubens, I., Debarbieux, L., Bohin, A., Lacroix, J.M., and Bohin, J.P. (1993) Homology between a genetic locus (*mdoA*) involved in the osmoregulated biosynthesis of periplasmic glucans in *Escherichia coli* and a genetic locus (*hrpM*) controlling pathogenicity of *Pseudomonas syringae*. *Mol Microbiol* 10: 329-340.

Mahajan-Miklos, S., Tan, M.W., Rahme, L.G., and Ausubel, F.M. (1999) Molecular mechanisms of bacterial virulence elucidated using a *Pseudomonas aeruginosa*-*Caenorhabditis elegans* pathogenesis model. *Cell* 96: 47-56.

Majdalani, N., and Gottesman, S. (2005) The Rcs phosphorelay: a complex signal transduction system. *Annu Rev Microbiol* 59: 379-405.

1 Miller, J.H. (1992) A short course in bacterial genetics: a laboratory manual and handbook for
2 *Escherichia coli* and related bacteria. New York USA Cold Spring Harbor Laboratory Press.

3 Mouslim, C., Latifi, T., and Groisman, E.A. (2003) Signal-dependent requirement for the co-
4 activator protein RcsA in transcription of the RcsB-regulated *ugd* gene. J of biol chem 278:
5 50588-50595.

6 Page, F., Altabe, S., Hugouvieux-Cotte-Pattat, N., Lacroix, J.M., Robert-Baudouy, J., and
7 Bohin, J.P. (2001) Osmoregulated periplasmic glucan synthesis is required for *Erwinia*
8 *chrysanthemi* pathogenicity. J Bacteriol 183: 3134-3141.

9 Perombelon, M.C.M. (2002) Potato diseases caused by soft-rot erwinias : an overview of
10 pathogenesis. Plant Pathol 51: 1-12.

11 Resibois, A., Colet, M., Faellen, M., Schoonejans, T., and Toussaint, A. (1984) Phi-EC2, a
12 new generalized transducing phage of *Erwinia chrysanthemi*. Virology 137: 102-112.

13 Roeder, D.L., and Collmer, A. (1985) Marker-exchange mutagenesis of a pectate lyase
14 isozyme gene in *Erwinia chrysanthemi*. J Bacteriol 164: 51-56.

15 Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989) Molecular cloning: a laboratory manual
16 2nd edn. New York USA, Cold Spring Harbor Laboratory press.

17 Sledjeski, D.D., and Gottesman, S. (1996) Osmotic shock induction of capsule synthesis in
18 *Escherichia coli* K-12. J bacteriol 178: 1204-1206.

19 Spiro, R.G. (1966) Analysis of sugars found in glycoproteins. Methods Enzymol 8: 3-27.

20 Stratmann, T., Pul, U., Wurm, R., Wagner, R., and Schnetz, K. (2012) RcsB-BglJ activates
21 the *Escherichia coli* *leuO* gene, encoding an H-NS antagonist and pleiotropic regulator of
22 virulence determinants. Mol microbiol 83: 1109-1123.

23 Vieira, J., and Messing, J. (1982) The pUC plasmids, an M13mp7-derived system for
24 insertion mutagenesis and sequencing with synthetic universal primers. Gene 19: 259-268.

25 Wang, D., Qi, M., Calla, B., Korban, S.S., Clough, S.J., Cock, P.J. et al. (2012) Genome-wide
26 identification of genes regulated by the Rcs phosphorelay system in *Erwinia amylovora*. Mol
27 plant-microbe interact 25: 6-17.

28 Wood, J.M. (1999) Osmosensing by bacteria: signals and membrane-based sensors. Microbiol
29 Mol biol rev 63: 230-262.

30 Yanisch-Perron, C., Vieira, J., and Messing, J. (1985) Improved M13 phage cloning vectors
31 and host strains: nucleotide sequences of the M13mp18 and pUC19 vectors. Gene 33: 103-
32 119.

33 Young, G.M., and Miller, V.L. (1997) Identification of novel chromosomal loci affecting
34 *Yersinia enterocolitica* pathogenesis. Mol microbiol 25: 319-328.

Table

Table 1. Bacterial Strains, plasmids and primers

Strain, plasmids or primer	Relevant genotype and/or phenotype ^{a,b}	Source or reference
<i>D. dadantii</i>		
EC3937	Wild type	Laboratory collection
NFB3723	<i>opgG</i> ::Cml	Laboratory collection
NFB3682	<i>rscC</i> ::Cml	Bouchart <i>et al.</i> , 2010
NFB3800	<i>flhD-uidA</i> -Kan ^r	Bouchart <i>et al.</i> , 2010
NFB3809	<i>ftsA-uidA</i> -Kan ^r	Bouchart <i>et al.</i> , 2010
NFB3835	<i>opgG</i> ::Cml, P _{BAD} <i>opgGH</i> -Spe ^r	This study
NFB3837	<i>opgG</i> ::Cml, P _{BAD} <i>opgGH</i> -Spe ^r , <i>ftsA-uidA</i> -Kan ^r	This study
NFB7112	<i>rscB</i> ::Cml, <i>rscCB</i> _{D56N} D-Spe ^r	This study
NFB7127	<i>rscC</i> ::Cml, <i>ftsA-uidA</i> -Kan ^r	Bouchart <i>et al.</i> , 2010
NFB7128	<i>rscC</i> ::Cml, <i>flhD-uidA</i> -Kan ^r	Bouchart <i>et al.</i> , 2010
NFB7202	<i>rscB</i> ::Cml, <i>flhD-uidA</i> -Kan ^r	Bouchart <i>et al.</i> , 2010
NFB7205	<i>rscB</i> ::Cml, <i>ftsA-uidA</i> -Kan ^r	Bouchart <i>et al.</i> , 2010
NFB7274	<i>opgG</i> ::Cml, <i>ftsA-uidA</i> -Kan ^r	This study
NFB7279	<i>rscB</i> ::Gm	This study
NFB7300	<i>opgG</i> ::Cml, P _{BAD} <i>opgGH</i> -Spe ^r , <i>flhD-uidA</i> -Kan ^r	This study
NFB7301	<i>opgG</i> ::Cml, <i>flhD-uidA</i> -Kan ^r	This study
NFB7368	<i>opgG</i> ::Cml, <i>rscB</i> ::Gm, P _{BAD} <i>opgGH</i> -Spe ^r	This study
NFB7369	<i>opgG</i> ::Cml, <i>rscB</i> ::Gm, P _{BAD} <i>opgGH</i> -Spe ^r , <i>flhD-uidA</i> -Kan ^r	This study
NFB7371	<i>opgG</i> ::Cml, <i>rscB</i> ::Gm, P _{BAD} <i>opgGH</i> -Spe ^r , <i>ftsA-uidA</i> -Kan ^r	This study
NFB7399	<i>rscB</i> ::Cml, <i>rscCB</i> _{D56N} D-Spe ^r , <i>flhD-uidA</i> -Kan ^r	This study
NFB7400	<i>opgG</i> ::FRT, <i>rscC</i> ::Cml, P _{BAD} <i>opgGH</i> -Spe ^r	This study
NFB7401	<i>opgG</i> ::Cml, <i>rscB</i> ::Gm, P _{BAD} <i>opgGH</i> -Spe ^r , <i>rscCB</i> _{D56N} D-Tet ^r	This study

NFB7402	<i>opgG::Cml, rcsB::Gm, P_{BAD}opgGH-Spe^r, rcsCB_{D56N}D-Tet^r, ftsA-uidA-Kan^r</i>	This study
NFB7403	<i>opgG::Cml, rcsB::Gm, P_{BAD}opgGH-Spe^r, rcsCB_{D56N}D-Tet^r, flhD-uidA-Kan^r</i>	This study
NFB7405	<i>opgG::FRT, rcsC::Cml, P_{BAD}opgGH-Spe^r, ftsA-uidA-Kan^r</i>	This study
NFB7406	<i>opgG::FRT, rcsC::Cml, P_{BAD}opgGH-Spe^r, ftsA-uidA-Kan^r</i>	This study
NFB7408	<i>rcsB::Cml, rcsCB_{D56N}D-Spe^r, ftsA-uidA-Kan^r</i>	This study
<i>E. coli</i>		
BL21 (DE3)	<i>ompT, hsdSB, gal, dcm</i>	Invitrogen
JM83	<i>ara Δ(lac-proAB), rpsL, thi, (φ80ΔlacZ-M15)</i>	Vieira <i>et al.</i> , 1982
S17-λpir	<i>recA1, thi, pro, hsdR-M+, RP4:2-Tc::Mu-Kan::Tn7, λpir</i>	De Lorenzo <i>et al.</i> , 1994
NFB3276	S17-λpir, pNFW233	This study
Plasmids		
pUC18Not	Amp ^r	Yanisch-Perron <i>et al.</i> , 1985
pUTminiTn5-Spe	mini-Tn5Spe, <i>oriR6K</i> , Spe ^r , Amp ^r	De Lorenzo <i>et al.</i> , 1990
pUTminiTn5-Tet	mini-Tn5tet, <i>oriR6K</i> , Tet ^r , Amp ^r	De Lorenzo <i>et al.</i> , 1990
pBAD18	Amp ^r , <i>araC</i> , promoter P _{BAD}	Guzman <i>et al.</i> , 1995
pNFW32	Amp ^r , <i>opgGH</i>	Page <i>et al.</i> , 2001
pNFW208	pBAD18, <i>opgG'</i>	This study
pNFW212	pBAD18, <i>opgGH</i>	This study
pNFW219	pUC18Not, <i>araC</i> , P _{BAD} <i>opgGH</i>	This study
pNFW257	pUC18Not, <i>rcsCBD</i>	Bouchart <i>et al.</i> , 2010
pNFW233	pUTminiTn5-Spe, <i>araC</i> , P _{BAD} <i>opgGH</i>	This study
pNFW374	pUC18Not, <i>rcsCB_{D56N}D</i>	This study
pNFW379	pUTminiTn5-Tet, pUC18Not, <i>rcsCB_{D56N}D</i>	This study
pNFW380	pUTminiTn5-Spe, pUC18Not, <i>rcsCB_{D56N}D</i>	This study
pNFW397	pUC18Not, <i>rcsCB::CmlD</i>	Bouchart <i>et al.</i> , 2010
pNFW402	pNFW397, Gm instead of Cml	This study

pNFW410	pET100/D-topo <i>rcsB</i>	This study
Primers	Sequence^{c,d}	
opgGHnhelf	GGGT <u>GCTAGCT</u> GATTTAGGCACACATACAGGGGG	
opgGHkpnIr	AACT <u>GGTACCAGAC</u> GGCAACGCCGTGTCGATAGCC	
gmhpalf	GTGT <u>GTTAAC</u> CCCCATCCCCCTGTTGAC	
gmhpalr	GTGT <u>GTTAAC</u> CGCAAGCTAGCTTGGCTGC	
rcsBacf	CACCATGAGCAATCTAAACGTAATTATTGCAG	
rcsBacr	CGCCGGCAAGAGCATTACTC	
rcsBDN3	CTGAAAGGTGTCTTTGCCATGCTGAATCTTCATCCC	
rcsBDNsoe1	GCATGGATAGATTGGTGATCAGGAC	
rcsBDNsoe2	GTCCTGATCACCAATCTATCCATGC	
rcsBDN4	TTCCAGCACTTTGCTGACGCTTTCC	

^a-*uidA* fusions are carried by a Kan^r mini-Tn5, *P_{BAD}opgGH* fusion is carried by a Spe^r mini-Tn5, *rcsCB_{D56N}D* is carried by a Tet^r mini-Tn5.

^bCml^r, chloramphenicol resistance; Amp^r, ampicillin resistance, Kan^r, kanamycin resistance, Spe^r, spectinomycin resistance, Str^r, streptomycin resistance, Gm^r, gentamicin resistance, Tet^r, tetracycline.

^cRestriction sites are underlined. ^dThe mutation is indicated in bold.

Figure legends

Fig. 1. OPGs concentration of the NFB3835 (*P_{BAD}opgGH*) at various concentrations of L-arabinose. Bacteria were grown until mid-log phase in M63 medium supplemented with indicated concentrations of L-arabinose. OPGs were extracted, purified and quantified as described in procedures. Results reported are the average of 3 independents experiments.

Fig. 2. Effect of the arabinose concentration on motility (A) and expression of the *flhD-uidA* (B) or the *ftsA-uidA* (C) gene fusions in various strains. Motility was measured in M63 semisolid plates supplemented with indicated L-arabinose concentrations. Swim diameters were measured after 30 h of incubation at 30°C. For fusion expression measurements, bacteria

were grown until mid log phase in the same media and broken by passing through a French pressure cell. The β -glucuronidase activities were measured with PNPU as a substrate. Specific activity is expressed as the change in optical density at 410 nm per min and per mg of protein. Results reported are the average of 3 independents experiments.

Fig. 3 Effect of arabinose concentration on pathogenicity of the NFB3835 ($P_{BAD}OpG$) strain. Bacteria were grown until mid log phase in M63 medium supplemented with indicated L-arabinose concentrations, washed twice and 10^7 bacteria were inoculated into scarified chicory leaves. Evolution of the disease symptoms were observed after 24 and 48h.

Fig. 4. Effect of OPG concentration *in planta* on RcsCD RcsB activation. For determination of OPG concentration, mass spectra or fusion activities, bacteria were collected after growth to mid log phase in 63 medium (0 h), or after growth *in planta* during 24 h, or after growth *in planta* during 48 h. For each time, CFU were determined. A/ concentration of OPGs purified from the wild-type strain (EC3937) as described in experimental procedures. B/ MALDI mass spectra acquired in positive ion mode on purified OPGs extracted from the wild-type strain. The $[M+Na]^+$ molecular ions are indicated on the top of each peak. Their respective $[M+K]^+$ molecular ions are identified by an asterisk and have an increment of 16 Da in comparison to the corresponding $[M+Na]^+$ ions. C/ Expression of the *flhD-uidA* or the *ftsA-uidA* fusion genes in the wild-type (NFB3800 and NFB3809 respectively) or the *rscB* (NFB7202 and NFB7205 respectively) strains. The β -glucuronidase activities were measured with PNPU as a substrate. Specific activity is expressed as the change in optical density at 410 nm per min and per 10^9 bacteria. Results reported are the average of 3 independents experiments.

Fig. 5 Effect of medium osmolarity on motility (A) and expression of the *flhD-uidA* (B) gene fusion in various strains. Motility was measured in half strength M63, M63, M63 supplemented with 0.1 M or M63 supplemented with 0.2M NaCl semisolid plates. For fusion expression measurements, bacteria were grown until mid log phase in the same media and broken by passing through a French pressure cell. The β -glucuronidase activities were measured with PNPU as a substrate. Specific activity is expressed as the change in optical density at 410 nm per min and per mg of protein. Results reported are the average of 3 independents experiments.

Fig. 6 A/ Effect of medium osmolarity on expression of the *ftsD-uidA* (A) gene fusion in various strains. For fusion expression measurements, bacteria were grown in half strength M63, M63, M63 supplemented with 0.1 M or M63 supplemented with 0.2M NaCl until mid-log phase and broken by passing through a French pressure cell. The β -glucuronidase activities were measured with PNPU as a substrate. Specific activity is expressed as the change in optical density at 410 nm per min and per mg of protein. Results reported are the average of 3 independents experiments. B/ OPGs concentration of the wild-type strain (EC3937) at various osmolarities. Bacteria were grown until mid log phase in the same media. OPGs were extracted, purified and quantified as described in experimental procedures. Results reported are the average of 3 independents experiments.