Ectomycorrhizal symbiosis helps plants to challenge salt stress conditions

Carmen Guerrero-Galán, Monica Calvo-Polanco, Sabine Dagmar Zimmermann

► To cite this version:

Carmen Guerrero-Galán, Monica Calvo-Polanco, Sabine Dagmar Zimmermann. Ectomycorrhizal symbiosis helps plants to challenge salt stress conditions. *Mycorrhiza*, 2019, 29 (4), pp.291-301. 10.1007/s00572-019-00894-2 . hal-02139380

HAL Id: hal-02139380

<https://hal.science/hal-02139380>

Submitted on 9 Feb 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

Ectomycorrhizal symbiosis helps plants to challenge salt stress conditions

Carmen Guerrero-Galán^{1,2}, Monica Calvo-Polanco¹ and Sabine Dagmar Zimmermann^{1*}

¹BPMP, Univ Montpellier, CNRS, INRA, SupAgro, Montpellier, France.

²Present address: Centro de Biotecnología y Genómica de Plantas, Instituto Nacional de Investigación y Tecnología Agraria y Alimentación (INIA), Universidad Politécnica de Madrid (UPM), 28223 Pozuelo de Alarcón, Spain.

*Author for correspondence: tel + 33 4 99 61 27 18; e-mail sabine.zimmermann@cnrs.fr
ORCID 0000-0002-5020-1447

Acknowledgments

CGG acknowledges the Ecole doctorale "GAIA" (University of Montpellier) for PhD fellowships financed by the French Ministry of Higher Education and Research.

Abstract

Soil salinity is an environmental constraint for plants that is currently increasing worldwide. The most common salt ions within soils, sodium and chloride, are considered as beneficial for plant development in small amounts, although their excess induces stress in most plants. Sodium is usually considered the most toxic ion in the couple, mainly because it causes an imbalance in the electrochemical gradients and competes with potassium for membrane transport systems and binding sites within cells. In addition, the sudden exposition of plants to salinity induces osmotic stress that imbalances root water and nutrient absorption and transport. Within the different strategies that plants use to cope with salinity, their association with beneficial soil microorganisms can play an important role in the adaptation to adverse environments. Among them, ectomycorrhizal fungi, specialized in colonizing the roots of woody plants, seem to have a protective effect on plants exposed to high amounts of salt. Although no precise mechanisms for this enhanced plant salt tolerance have been described yet, the improvement of plant water and mineral nutrition, and the restriction to sodium entrance and extrusion of toxic ions from the photosynthetic organs are proposed as possible strategies. This review aims at summarizing the current knowledge accumulated on the role of ectomycorrhizal symbiosis in plant tolerance to salt stress.

Keywords

Ectomycorrhizal symbiosis, sodium, salt tolerance, trees, nutrient and water transport, membrane transporters.

Introduction

Salinity, understood as the excess of sodium (Na^+) salts in the soil, is a detrimental condition for plant performance that is increasing nowadays and challenges both agricultural crops and natural ecosystems (Daliakopoulos et al. 2016). Salt stress induces both an osmotic and a toxic effect in plants which disturbs major plant metabolic processes and hampers plant water and nutrient balance. Exposure of plant roots to increasing NaCl will lead to competition of Na^+ ions with potassium (K^+) for its transporters so that Na^+ enters the plant cells through either non-specific or selective transport systems (Wu 2018). Thus, Na^+ concentrations will build up in the cytosol where it alters enzyme activity by displacement of the Ca^{2+} and K^+ ions of their binding sites (Cramer et al. 1985; Rodríguez-Navarro 2000). The abundance of Na^+ ions also changes the electrochemical gradient between both sides of the plasma membrane, affecting the energization of soil nutrient uptake (Tavakkoli et al. 2010). Palliating these negative effects diverts the energetic resources to detoxification, depleting other processes like nutrition, growth or reproduction (Munns and Tester 2008).

Maintaining K^+ contents in the cell cytoplasm and its movements in the vascular tissues is essential for salt-stressed plants to overcome the challenging environment (Shabala and Cuin 2008). Under salt stress, an initial massive influx of Na^+ in the root triggers a membrane depolarization that provokes a K^+ efflux from the cytosol (Shabala et al. 2006), accompanied by an increase in K^+ storage in the vacuole (Jiang et al. 2010). The drop in the cytosolic K^+ level induces high affinity transporters to boost the uptake from the roots (Shabala 2017). In order to prevent Na^+ from reaching the photosynthetic organs, plants try to minimize its accumulation in the tissues through changes in gene expression and protein function of membrane transporters that aim at reducing the Na^+ uptake and at limiting its allocation to the leaves, by extruding it out from root cells or into the vacuoles.

One of the first responses of plants to salinity will be the closure of stomata, the reduction of transpiration, and the adjustment of the root water uptake. It is commonly accepted that water is transferred from the soil to the root through two main paths, the cell wall continuum (apoplastic path) and through the cell-to-cell path (symplastic path) which includes the transfer of water through plasmodesmata and plasma membranes using specialized water channel proteins called aquaporins. Aquaporins are membrane intrinsic proteins (MIPs) present in all species, including fungi and plants (Nehls and Dietz 2014; Verdoucq and Maurel 2018). Under different stresses including salinity, water uptake has been mainly attributed to the tight regulation of aquaporins (Boursiac et al. 2005; Afzal et al. 2016). Plant cells also accumulate compatible osmolytes or osmoprotectants. Although the role of compatible osmolytes has not

86 been fully unravelled, it has been observed that they protect against oxidative stress inside the
87 cells (Singh et al. 2015). Moreover, several studies have demonstrated that the overexpression
88 of genes leading to increased accumulation of these solutes can confer on plants the ability to
89 tolerate salinity (Parihar et al. 2015), and also, compatible solutes will alter how water is
90 uptaken by the roots (Aroca et al. 2012; Calvo-Polanco et al. 2014).

91 Taking all this into account, we observe that plants are not devoid of defence
92 mechanisms against salinity, at least in the short term. However, in the long run, these responses
93 may not be enough against a sustained stress and require further physiological, ecological and
94 evolutionary adaptations. This is the case of root symbioses, such as those established with
95 mycorrhizal fungi, in which the microorganism improves plant water uptake and mineral
96 nutrition. The extension of the exchange surface provided by the mycelium is an important
97 factor in the enhanced absorption of water and nutrients of mycorrhizal plants as the hyphae
98 are able to penetrate narrower pores (Bogeat-Triboulot et al. 2004; Lehto and Zwiazek 2011).
99 Another beneficial effect is the improvement of soil structure that facilitates plant rooting
100 (Rillig and Mummey 2006). All this translates in a stronger growth and higher biomass
101 accumulation in mycorrhizal plants, which have a better fitness and consequently adapt better
102 to challenging environments than in non-mycorrhizal conditions (Smith and Read 2008). So
103 far, most information on the beneficial effects of mycorrhizal plant-fungal interactions in plant
104 tolerance to salinity has been gathered from arbuscular mycorrhizal symbiosis revealing several
105 mechanisms involved (Evelin et al. 2009; Lenoir et al. 2016). Other symbioses, as the
106 ectomycorrhizal interactions specific to ligneous species, have evolved in many fungal and
107 plant lineages independently, contrarily to arbuscular mycorrhiza, which is considered a
108 plesiomorphy of plants (Field et al. 2015). This evolutionary diversity implies that the
109 association with ectomycorrhizal fungi can have multiple functional roles in the physiology of
110 the host. Little is known about ectomycorrhizal fungi concerning the mechanisms that
111 contribute to alleviation of the effects of salinity on their host plants. In this review we present
112 the research carried out thus far in this field and the main conclusions we can draw from them.

114 **Ectomycorrhizal fungi and the tolerance of plants to salt stress**

115 *Soil fungi differ greatly in their behaviour in saline environments*

116 The tolerance of fungi to salt stress has been often screened by measuring their capability to
117 adapt and grow in a culture medium supplemented with NaCl. Some extremophiles can grow
118 at saturating concentrations of NaCl in their natural environments (Gunde-Cimerman et al.
119 2009), whereas others are extremely sensitive. Regarding salt tolerance of ectomycorrhizal

120 fungal species, differences have been described between the two major phyla (Tresner and
121 Hayes 1971): as a general trend, Basidiomycota are less adapted to saline conditions than
122 Ascomycota, but some species are able to support relatively high NaCl concentrations (Zalar et
123 al. 2007). As a matter of fact, there are only 21 marine species of filamentous Basidiomycota,
124 very scarce compared to the 805 marine species of filamentous Ascomycota (Jones et al. 2015).
125 Overall, the salinity tolerance trait has been most studied in Basidiomycota. It is not necessarily
126 linked to specific clades but is found frequently in members of the same species or even the
127 same genus.

128 Ectomycorrhizal fungi are not usually found in environments of extreme salinity, such
129 as salt ponds, and it is unlikely that they play a role in hypersaline environments, since
130 halophytic plants are mostly herbaceous and non-mycorrhizal (Brundrett 2009, 2017),
131 However, they have been shown to be able to adapt to saline environments, and to adapt and
132 grow in a culture medium supplemented with NaCl by *in vitro* tests. These tests on salinity
133 tolerance of different fungal strains (Table 1) were achieved at NaCl concentrations of 120-800
134 mM of *e.g.* *Hebeloma* sp. or *Laccaria* sp. (Hutchison 1990; Kernaghan et al. 2002), and of
135 isolates of the same species as *Paxillus involutus* (Zhang et al. 2008), *Suillus* spp. (Hutchison
136 1990; Dixon et al. 1993; Bois et al. 2006a; Tang et al. 2009; Obase et al. 2010), and *Pisolithus*
137 *sp.* (Matsuda et al. 2006). In all cases, ion toxicity seems to be responsible for most of the
138 negative effects observed on fungal growth, compared to the osmotic component due to the
139 abundance of solutes in the environment (Dixon et al. 1993; Chen et al. 2001; Kernaghan et al.
140 2002; Bois et al. 2006a).

141 Also, some ectomycorrhizal fungi seem to have evolved towards salt tolerance and their
142 growth can even be stimulated by the application of moderate NaCl concentrations in the range
143 of 25-50 mM (Dixon 1993; Bois et al. 2006a; Matsuda et al. 2006) or higher (Zhang et al. 2008;
144 Obase et al. 2010). Recent studies using genomic techniques, have detected how populations
145 of the ectomycorrhizal fungus *Suillus brevipes* in North America are able to adapt to stress
146 associated with climate regimes and abiotic environments (Branco et al. 2015, 2017).

148 *Ectomycorrhizal fungi improve the tolerance of plants to salt stress with different strategies*

149 Thus far, few studies have shown improved plant salt stress tolerance by ectomycorrhizal fungi
150 compared to arbuscular mycorrhizal symbiosis. The effect of the presence of ectomycorrhizal
151 fungi in alleviation of stress symptoms seems to be fungus- and/or host-dependent. Dixon et al.
152 (1993) examined ectomycorrhizal *Pinus taeda* seedlings and found different responses
153 depending on the fungal inoculum. Whereas Na⁺ salts inhibited mycorrhization by some of the

154 fungal species tested, root colonization by *Pisolithus tinctorius* and *Laccaria laccata* was not
155 affected under 80 mM. Also, a positive effect on biomass accumulation was found at mild salt
156 stress (20-40 mM NaCl) on the seedlings mycorrhized with relatively salt-tolerant fungi. Other
157 studies have tried to look into further physiological details, particularly the nutritional status of
158 the plant to understand the specific processes for adaptation (see below). Complementarily to
159 mineral and water nutrition, other processes, such as the activation of stress response pathways
160 (metabolites, hormones) by ectomycorrhizal colonization can also play a role in the positive
161 effect against environmental stresses (Luo et al. 2009; Ma et al. 2014).

162 The mechanisms responsible for the enhancement of plant tolerance to salt stress by
163 ectomycorrhizal fungi are still unclear and several strategies could be involved. First, there is
164 an undeniable redistribution of Na⁺ ions in salt-stressed mycorrhizal plants compared to their
165 non-inoculated counterparts. It seems crucial for survival that Na⁺ does not reach the
166 photosynthetic tissues, and this protective effect has been found in seedlings inoculated with
167 salt-tolerant fungi. As a general rule, ectomycorrhizal seedlings subjected to salt stress were
168 able to load less Na⁺ to the shoots when inoculated with *Hebeloma spp.* (Muhsin and Zwiazek
169 2002a; Nguyen et al. 2006; Calvo-Polanco et al. 2008a, 2008b), *Laccaria bicolor* (Bois et al.
170 2006b; Nguyen et al. 2006; Calvo-Polanco et al. 2008b) and *Suillus tomentosus* (Calvo-Polanco
171 et al. 2008a). This effect could be linked to the strong extrusion Na⁺ by the fungus observed in
172 *Populus x canescens* - *P. involutus* symbioses (Li et al. 2012). In order to limit translocation of
173 toxic Na⁺ to the aerial parts, a successful strategy of plants is the storage of Na⁺ in the root
174 apoplasm, rather than in the vacuoles of root cells, avoiding its entrance into the symplastic
175 pathway (Ottow et al. 2005). It is very likely that the Na⁺ accumulated in the root tissues is
176 retained by the fungus, in order to protect the photosynthetic apparatus of the plant to keep its
177 production of carbon compounds.

178 These findings indicate that some ectomycorrhizal fungi function as a sort of “buffer”
179 or physical barrier between the external medium and the root in order to avoid the transfer of
180 toxic Na⁺ ions into the root cortex. Following this idea, Langenfeld-Heyser et al. (2007)
181 analysed *P. x canescens* inoculated with *P. involutus* and found that the fungus did not change
182 the Na⁺ and chloride (Cl⁻) contents in the root cell walls, particularly in those of the xylem
183 vessels, but somehow the fungus delayed the transfer of Na⁺ to the shoot. An intermediate result
184 was obtained by Ma et al. (2014) with two strains of *P. involutus*, one symbiotic and one non-
185 symbiotic. The inoculation of *P. x canescens* with each of the *P. involutus* lines impaired Na⁺
186 translocation to the shoot, even though one of the lines does not form functional mycorrhizae.
187 Although these results seem somehow contradictory, these studies have been done with

different ectomycorrhizal fungi on distant plant species, gymnosperms and angiosperms, that do not necessarily have the same root and mycorrhizal structure and therefore it is expected that the mechanisms of salt storage and transfer may differ.

Ectomycorrhizal fungi improve plant mineral nutrition to alleviate salt stress

The exclusion of Na^+ from sensitive tissues is commonly accepted to be tightly related to K^+ contents. Both ions are physico-chemically similar (Benito et al. 2014) and can be transported partly by the same membrane proteins, but their roles and effects in plant cells are completely different. Potassium is an essential macronutrient for plants, in which it is kept at high concentrations (100-200 mM) to play many cellular and physiological roles, from membrane energization to photosynthesis (Ahmad and Maathuis 2014; Dreyer et al. 2017). On the other hand, Na^+ is only beneficial in small amounts (in the nano- or micromolar range) and its excess is harmful for plant tissues, except in some particular cases (Hasegawa et al. 2000). In salt stress conditions, plants can suffer from K^+ deficiency due to the competition with Na^+ for uptake systems at the plasma membrane (Botella et al. 1997). It has been demonstrated that ectomycorrhizal fungi contribute to K^+ nutrition of plants (Jentschke et al. 2001) especially under conditions of K^+ starvation (Garcia et al. 2014). Some species are able to improve the K^+ content in the leaves of their host, especially in unfavourable conditions. This has been observed in the symbiotic models *P. involutus* – *P. x canescens* (Langenfeld-Heyser et al. 2007; Luo et al. 2009), *Scleroderma bermudense* – *Coccoloba uvifera* (Bandou et al. 2006) and *Hebeloma crustuliniforme* – *Picea glauca* (Muhsin and Zwiazek 2002a). Li et al. (2012) demonstrated by flux measurements that K^+ homeostasis was maintained by *P. involutus* in salt stressed poplar, thanks to the reduction of the salt-induced leakage of K^+ and to the contribution of the fungus to the host's absorption. Thus, an adequate K^+ nutrition enhanced by the ectomycorrhizal symbiosis seems to be crucial for a better adaptation of plants to environmental stresses, and in particular to salinity (Munns and Tester 2008; Wang et al. 2013).

Another strategy against the detrimental effects of high levels of salts is given by the advantageous effects of the presence of ectomycorrhizal fungi in the improvement of the general mineral nutrition of the plant. Indeed, some ectomycorrhizal fungi are able to keep the seedlings in a healthier state, for example by increasing the absorption of Ca^{2+} , which is related as well to the K^+/Na^+ ratio in the tissues (Ma et al. 2014), or by maintaining high phosphorus and nitrogen contents in the seedlings (Muhsin and Zwiazek 2002a). Luo et al. (2011) also found evidence that ectomycorrhizal poplar (*P. involutus*) conserved a better nutritional status, which led to a reduced stress response and higher photosynthetic activity. A relative effect has

also been observed in plants that are adapted to saline environments, such as the tropical seagrass *C. uvifera* (Bandou et al. 2006), whose ectomycorrhizal roots efficiently excluded Na^+ and Cl^- ions and enhanced the absorption of phosphorus. However, no changes in the phosphorus content of salt-stressed mycorrhizal plants have been also reported (Langenfeld-Heyser et al. 2007).

Ectomycorrhizal fungi can regulate plant water transport under salt stress

In addition to the effects of salinity in ion homeostasis, NaCl is a known inhibitor of aquaporin-mediated water transport in roots (Boursiac et al. 2005; Lee et al. 2010) and is also related to the reduction of plant transpiration. The inoculation of seedlings with ectomycorrhizal fungi keeps them in an overall healthier transpiration rate (Muhsin and Zwiazek 2002b; Bois et al. 2006b; Calvo-Polanco et al. 2008b). Different studies in AM fungi (Barzana et al. 2015; Calvo-Polanco et al. 2016) and ectomycorrhizal fungi (Marjanović et al. 2005; Hacquard et al. 2013; Navarro-Ródenas et al. 2013; Xu et al. 2015; Peter et al. 2016) have already pointed out the role of mycorrhizal fungi in modifying the expression and abundance of plant aquaporins under different soil environmental conditions. The up-regulation of these traits has been usually linked to an increase in the water transport capacity of plants (Xu et al. 2015), specially under stress (Barzana et al. 2015; Calvo-Polanco et al. 2016). However, in some cases, ectomycorrhizal symbiosis had no or even a negative effect on the root hydraulic conductivity (Calvo-Polanco et al. 2011; Xu et al., 2016), but the specific processes responsible are still unknown. Taken together, the role of fungal aquaporins as modulators of water transport in stress conditions cannot be dismissed, as it might be one of the key factors for salt stress tolerance in combination with activation of pathways that involves hormones and stress signal molecules and that will need future studies.

Membrane transport systems involved in symbiotic tolerance to salt stress

Molecular mechanisms responsible for plant salinity tolerance mediated by ectomycorrhizal symbiosis include membrane transport systems both at the fungal and at the plant root membranes. Such membrane transporters would be critical for all the three adaptation strategies shown by ectomycorrhizal symbioses, for Na^+ exclusion, K^+ nutrition and water transport (Fig. 1).

Regarding the barrier function of the fungus, it would be particularly interesting to characterize the membrane transport systems responsible for Na^+ fluxes. The ENA1 ATPase (Rodríguez-Navarro and Benito 2010) and the Na^+/H^+ antiporters Nha-like (Bañuelos et al.

1998; Proft and Struhl 2004), both involved in tolerance to NaCl in fungi, are promising candidates for Na⁺ efflux from the fungus to the external medium. The first evidence for this has been found in *S. brevipes* from a salinity area that had adapted to tolerate salt stress by developing differential genomic regions including a gene encoding a membrane Na⁺/H⁺ antiporter (Branco et al. 2015). These results are also supported by experiments by Li et al. (2012), in which the pharmacological inhibition of Na⁺/H⁺ antiporters of *P. involutus* had an effect on its salt-induced extrusion of Na⁺. These results are also supported by experiments by Li et al. (2012), in which the pharmacological inhibition of Na⁺/H⁺ antiporters of *P. involutus* had an effect on its salt-induced extrusion of Na⁺. It is also possible that the excess of Na⁺ in the fungus is stored in vacuoles or endosomes rather than in the cytosol, by members of the VNX and NHX transporter families (Nass et al. 1997; Cagnac et al. 2007).

Potassium transporters and channels of the fungal and plant partners of the symbiosis could cause a differential distribution of this ion within the mycorrhizal plant (Garcia and Zimmermann, 2014). Selective fungal uptake transport systems, belonging to the Trk or HAK families, could fight the excess of external Na⁺ with a higher selective absorption than plant transporters. A further filter would be applied by the K⁺-selective channels that are thought to release K⁺ into the plant apoplasm (Guerrero-Galán et al. 2018a, b). It is also likely that fungi face the K⁺/Na⁺ imbalance through storage of K⁺ in the vacuole or Golgi and the Kha1 transporters could play this role. It is also noteworthy that, due to their similarity, Na⁺ and K⁺ can be permeated through the same transport systems with different affinities (Ariño et al. 2010).

In parallel to the K⁺ nutrition, fungal and plant aquaporins mediating water transport (Fig. 1) and their regulation are involved in salt stress responses of ectomycorrhizal plants as summarized above. These molecular players are well described in plants (Verdoucq and Maurel 2018) and have been identified in fungal genomes and partially characterized (Dietz et al. 2011; Xu et al. 2015; Peter et al. 2016). For now, seven aquaporins have been found in the fungus *L. bicolor*, of which *LbAQP1* is necessary for the development of functional ectomycorrhizae in trembling aspen (Navarro-Ródenas et al. 2015). Interestingly, in transcriptomic analysis of the symbiotic couple *Cenococcum geophilum*-*Pinus sylvestris* cultured under drought, one aquaporin (Cenge3:604158) was up-regulated, suggesting that the fungus may help the plant to cope with decreased water absorption or compensate the inhibition of plant aquaporins, although no physiological differences with non-inoculation controls was reported (Peter et al. 2016). In the future, discoveries in the area of fungal genomics will open new horizons in the

study of plant-fungal symbiosis and are expected to be critical for the improvement of plant adaptation to the presence of salt within the soil.

The role of ectomycorrhizal associations in salt stress tolerance and their use for reforestation

In controlled *in vitro* or greenhouse experiments it is relatively easy to apply inocula and to calculate the degree of root colonization. The fixed environmental conditions can be radically different from those found in the real environment, where the climate and the existing rhizospheric biodiversity can greatly change the result of the experiment (Weissenhorn et al. 2002; Onwuchekwa et al. 2014). It is possible that fungi selected and grown in laboratory conditions are less competitive than the pre-existing microorganisms, and disappear from the roots in a short time span (Onwuchekwa et al. 2014). In addition, although the effect of ectomycorrhizae in growing seedlings is evident, it is still unknown to what extent ectomycorrhizal fungi can improve salt tolerance of adult trees.

However, some research aimed to apply ectomycorrhizal symbiosis on plant seedlings intended for ecological restoration of degraded areas, for example in the saline-alkaline soils contaminated by composite tailing residues of bitumen extraction. In these areas, plants are severely affected by accumulation of Na⁺, displaying reduced uptake of essential nutrients, such as K⁺ and phosphorus, as well as leaf dieback (Renault et al. 1999). The first trials involved direct testing of salts or composite tailing water on axenic cultures of ectomycorrhizal fungi to assess which species and isolates were suitable for application in the field (Kernaghan et al. 2002; Bois et al. 2005). In parallel, several lots of autochthonous tree species were analyzed (Khasa et al. 2002; Bois et al. 2006a,b; Nguyen et al. 2006; Yi et al. 2008; Calvo-Polanco et al. 2008b, 2011). The optimal combinations of plants and fungi have been tested in experiments on oil sands reclamation sites with encouraging results with the prospect of a large-scale application. Inoculated seedlings had higher survival rates and biomass accumulation than the non-mycorrhizal plants, although the survival of the fungi in the root was not ensured (Onwuchekwa et al. 2014; Hankin et al. 2015). Research efforts are still in progress to select new pioneer symbiotic couples for land reforestation (Beaudoin-Nadeau et al. 2016).

To sum up, perspectives in this rather unexplored field of the use of ectomycorrhizal associations to struggle with salt stress could range from molecular approaches to application of ectomycorrhizal fungi in nurseries or forest cultures, offering exciting opportunities to fundamental and applied research.

323

324 References

- 325 Afzal Z, Howton TC, Sun Y, Mukhtar MS (2016) The roles of aquaporins in plant stress
326 responses. *J Dev Biol* 4:9. <https://doi.org/10.3390/jdb4010009>
- 327 Ahmad I, Maathuis FJM (2014) Cellular and tissue distribution of potassium: Physiological
328 relevance, mechanisms and regulation. *J Plant Physiol* 171:708-714.
329 <https://doi.org/10.1016/j.jplph.2013.10.016>
- 330 Ariño J, Ramos J, Sychorvá H (2010) Alkali metal cation transport and homeostasis in yeast.
331 *Microbiol Mol Biol Rev* 74:95-120. <http://dx.doi.org/10.1128/MMBR.00042-09>
- 332 Aroca R, Porcel R, Ruiz-Lozano JM (2012) Regulation of root water uptake under abiotic stress
333 conditions. *J Exp Bot* 63:43–57. <https://doi.org/10.1093/jxb/err266>
- 334 Bandou E, Lebailly F, Muller F, Dulormne M, Toribio A, Chabrol J, et al. (2006) The
335 ectomycorrhizal fungus *Scleroderma bermudense* alleviates salt stress in seagrape
336 (*Coccoloba uvifera* L.) seedlings. *Mycorrhiza* 16:559–565. [https://doi.org/10.1007/s00572-](https://doi.org/10.1007/s00572-006-0073-6)
337 006-0073-6
- 338 Bañuelos MA, Sychrová H, Bleykasten-Grosshans C, Souciet JL, Potier S (1998) The Nha1
339 antiporter of *Saccharomyces cerevisiae* mediates sodium and potassium efflux. *Microbiol*
340 144:2749–2758. <https://dx.doi.org/10.1099/00221287-144-10-2749>
- 341 Bárzana G, Aroca R, Bienert GP, Chaumont F, Ruiz-Lozano JM (2014) New insights into the
342 regulation of aquaporins by the arbuscular mycorrhizal symbiosis in maize plants under
343 drought stress and possible implications for plant performance. *Mol Plant Microbe Interact*
344 27: 349-363. <http://doi.org/10.1094/MPMI-09-13-0268-R>
- 345 Beaudoin-Nadeau M, Gagné A, Bissonnette C, Bélanger P, Fortin J, Roy S, et al. (2016)
346 Performance of ectomycorrhizal alders exposed to specific Canadian oil sands tailing
347 stressors under *in vivo* bipartite symbiotic conditions. *Can J Microbiol* 7:1-7.
348 <https://doi.org/10.1139/cjm-2015-0703>
- 349 Benito B, Haro R, Amtmann A, Cuin TA, Dreyer I (2014) The twins K⁺ and Na⁺ in plants. *J*
350 *Plant Physiol* 171:723–731. <https://doi.org/10.1016/j.jplph.2013.10.014>
- 351 Bogeat-Triboulot MB, Bartoli F, Garbaye J, Marmeisse R, Tagu D (2004) Fungal
352 ectomycorrhizal community and drought affect root hydraulic properties and soil adherence
353 to roots of *Pinus pinaster* seedlings. *Plant Soil* 267:213–223.
354 <https://doi.org/10.1007/s11104-005-5349-7>
- 355 Bois G, Piché Y, Fung M, Khasa D (2005) Mycorrhizal inoculum potentials of pure reclamation
356 materials and revegetated tailing sands from the Canadian oil sand industry. *Mycorrhiza*
357 15:149-158. <https://doi.org/10.1007/s00572-004-0315-4>
- 358 Bois G, Bertrand A, Piché Y, Fung M, Khasa D (2006a) Growth, compatible solute and salt
359 accumulation of five mycorrhizal fungal species grown over a range of NaCl concentrations.
360 *Mycorrhiza* 16:99-109. <https://doi.org/10.1007/s00572-005-0020-y>
- 361 Bois G, Bigras F, Bertrand A, Piché Y, Fung M, Khasa D (2006b) Ectomycorrhizal fungi affect
362 *Picea glauca* and *Pinus banksiana* in response to NaCl gradient. *Tree Physiol* 26:1185-1196.
363 <https://doi.org/10.1093/treephys/26.9.1185>
- 364 Botella MA, Martínez V, Pardines J, Cerdá A (1997) Salinity induced potassium deficiency in
365 maize plants. *J Plant Physiol* 150:200–205. [https://doi.org/10.1016/S0176-1617\(97\)80203-](https://doi.org/10.1016/S0176-1617(97)80203-9)
366 9
- 367 Boursiac Y, Chen S, Luu DT, Sorieul M, van den Dries N, Maurel C (2005) Early effects of

368 salinity on water transport in *Arabidopsis* roots. Molecular and cellular features of aquaporin
369 expression. *Plant Physiol* 139:790–805. <https://doi.org/10.1104/pp.105.065029>

370 Branco S, Gladieux P, Ellison CE, Kuo A, LaButti K, Lipzen A, et al. (2015) Genetic isolation
371 between two recently diverged populations of a symbiotic fungus. *Mol Ecol* 24: 2747–2758.
372 <http://doi.org/10.1111/mec.13132>

373 Branco S, Bi K, Liao H-L, Gladieux P, Badouin H, Ellison CE, et al. (2017) Continental-level
374 population differentiation and environmental adaptation in the mushroom *Suillus brevipes*.
375 *Mol Ecol* 26: 2063–2076. <https://doi.org/10.1111/mec.13892>

376 Brundrett MC (2009) Mycorrhizal associations and other means of nutrition of vascular plants:
377 Understanding the global diversity of host plants by resolving conflicting information and
378 developing reliable means of diagnosis. *Plant Soil* 320:37–77.
379 <https://doi.org/10.1007/s11104-008-9877-9>

380 Brundrett MC (2017) Global diversity and importance of mycorrhizal and nonmycorrhizal
381 plants. *Ecol Stud* 230:533–556. https://doi.org/10.1007/978-3-319-56363-3_21

382 Cagnac O, Leterrier M, Yeager M, Blumwald E (2007) Identification and characterization of
383 Vnx1p, a novel type of vacuolar monovalent cation/H⁺ antiporter of *Saccharomyces*
384 *cerevisiae*. *J Biol Chem* 282:24284–24293. <https://doi.org/10.1074/jbc.M703116200>

385 Calvo Polanco M, Zwiazek J, Voicu M (2008a) Responses of ectomycorrhizal American elm
386 (*Ulmus americana*) seedlings to salinity and soil compaction. *Plant Soil* 308:189–200.
387 <https://doi.org/10.1007/s11104-008-9619-z>

388 Calvo Polanco M, Zwiazek J, Jones M, MacKinnon M (2008b) Responses of mycorrhizal jack
389 pine (*Pinus banksiana*) seedlings to NaCl and boron. *Trees* 22:825–834.
390 <https://doi.org/10.1007/s00468-008-0243-6>

391 Calvo-Polanco M, Zwiazek JJ (2011) Role of osmotic stress in ion accumulation and
392 physiological responses of mycorrhizal white spruce (*Picea glauca*) and jack pine (*Pinus*
393 *banksiana*) to soil fluoride and NaCl. *Acta Physiol Plant* 33:1365–1373.
394 <http://doi.org/10.1007/s11738-010-0670-z>

395 Calvo-Polanco M, Sanchez-Romera B, Aroca R (2014) Mild salt stress conditions induce
396 different responses in root hydraulic conductivity of *Phaseolus vulgaris* over-time. *PloS One*
397 9:e90631. <https://doi.org/10.1371/journal.pone.0090631>

398 Calvo-Polanco M, Sánchez-Castro I, Cantos M, García JL, Azcón R, Ruiz-Lozano JM, et al.
399 (2016) Effects of different arbuscular mycorrhizal fungal backgrounds and soils on olive
400 plants growth and water relation properties under well-watered and drought conditions. *Plant*
401 *Cell Environ* 39:2498–2514. <https://doi.org/10.1111/pce.12807>

402 Chen DM, Ellul S, Herdman K, Cairney JWG (2001) Influence of salinity on biomass
403 production by Australian *Pisolithus* spp. isolates. *Mycorrhiza* 11:231–236.
404 <https://doi.org/10.1007/s005720100126>

405 Cramer GR, Läuchli A, Polito VS (1985) Displacement of Ca²⁺ by Na⁺ from the plasmalemma
406 of root cells: A primary response to salt stress? *Plant Physiol* 79:207–211.
407 <https://doi.org/10.1104/pp.79.1.207>

408 Daliakopoulos I, Tsanis I, Koutroulis A, Kourgialas N, Varouchakis A, Karatzas G, et al. (2016)
409 The threat of soil salinity: A European scale review. *Sci Total Environ* 573:727–739.
410 <https://doi.org/10.1016/j.scitotenv.2016.08.177>

411 Dietz S, von Bülow J, Beitz E, Nehls U (2011) The aquaporin gene family of the
412 ectomycorrhizal fungus *Laccaria bicolor*: Lessons for symbiotic functions. *New Phytol* 190,
413 927–940. <http://doi.org/10.1111/j.1469-8137.2011.03651.x>

- Dixon RK, Rao MV, Garg VK (1993) Salt stress affects *in vitro* growth and *in situ* symbioses of ectomycorrhizal fungi. *Mycorrhiza* 3:63–68. <https://doi.org/10.1007/BF00210694>
- Dreyer I, Gomez-Porras JL, Riedelsberger J (2017) The potassium battery: A mobile energy source for transport processes in plant vascular tissues. *New Phytol* 216:1049–1053. <https://doi.org/10.1111/nph.14667>
- Evelin H, Kapoor R, Giri B (2009) Arbuscular mycorrhizal fungi in alleviation of salt stress: A review. *Ann Bot* 104:1263–1280. <https://doi.org/10.1093/aob/mcp251>
- Field KJ, Pressel S, Duckett JG, Rimington WR, Bidartondo MI (2015) Symbiotic options for the conquest of land. *Trends Ecol Evol* 30:477–486. <https://doi.org/10.1016/j.tree.2015.05.007>
- Garcia K, Zimmermann SD (2014) The role of mycorrhizal associations in plant potassium nutrition. *Front Plant Sci* 5: 337. <http://doi.org/10.3389/fpls.2014.00337>
- Garcia K, Delteil A, Conéjéro G, Becquer A, Plassard C, Sentenac H, et al. (2014) Potassium nutrition of ectomycorrhizal *Pinus pinaster*: Overexpression of the *Hebeloma cylindrosporum* *HcTrk1* transporter affects the translocation of both K⁺ and phosphorus in the host plant. *New Phytol* 201:951–960. <http://doi.org/10.1111/nph.12603>
- Guerrero-Galán C, Delteil A, Garcia K, Houdinet G, Conéjéro G, Gaillard I, et al. (2018a) Plant potassium nutrition in ectomycorrhizal symbiosis: Properties and roles of the three fungal TOK potassium channels in *Hebeloma cylindrosporum*. *Environ Microbiol* 20:1873–1887. <http://doi.org/10.1111/1462-2920.14122>
- Guerrero-Galán C, Garcia K, Houdinet G, Zimmermann SD (2018b) *HcTOK1* participates in the maintenance of K⁺ homeostasis in the ectomycorrhizal fungus *Hebeloma cylindrosporum*, which is essential for the symbiotic K⁺ nutrition of *Pinus pinaster*. *Plant Signal Behav* 13:e1480845. <http://doi.org/10.1111/pce.13359>
- Gunde-Cimerman N, Ramos J, Plemenitaš A (2009) Halotolerant and halophilic fungi. *Mycol Res* 113:1231–1241. <https://doi.org/10.1016/j.mycres.2009.09.002>
- Hacquard S, Tisserant E, Brun A, Legué V, Martin F, Kohler A (2013) Laser microdissection and microarray analysis of *Tuber melanosporum* ectomycorrhizas reveal functional heterogeneity between mantle and Hartig net compartments. *Environ Microbiol* 5:1853–1869. <http://doi.org/10.1111/1462-2920.12080>
- Hankin S, Karst J, Landhäusser SM (2015) Influence of tree species and salvaged soils on the recovery of ectomycorrhizal fungi in upland boreal forest restoration after surface mining. *Botany* 93:267–277. <http://doi.org/10.1139/cjb-2014-0132>
- Hasegawa PM, Bressan RA, Zhu JK, Bohnert HJ (2000) Plant cellular and molecular responses to high salinity. *Annu Rev Plant Physiol Plant Mol Biol* 51:463–499. <http://doi.org/10.1146/annurev.arplant.51.1.463>
- Hutchison LJ (1990) Studies on the systematics of ectomycorrhizal fungi in axenic culture. IV. The effect of some selected fungitoxic compounds upon linear growth. *Can J Bot* 68:2172–2178. <https://doi.org/10.1139/b90-283>
- Jentschke G, Brandes B, Kuhn AJ, Schröder WH, Godbold DL (2001) Interdependence of phosphorus, nitrogen, potassium and magnesium translocation by the ectomycorrhizal fungus *Paxillus involutus*. *New Phytol* 149:327–337. <http://doi.org/10.1046/j.1469-8137.2001.00014.x>
- Jiang X, Leidi EO, Pardo JM (2010) How do vacuolar NHX exchangers function in plant salt tolerance? *Plant Signal Behav* 5:792–795. <https://dx.doi.org/10.4161/psb.5.7.11767>
- Jones EBG, Suetrong S, Sakayaroj J, Bahkali AH, Abdel-Wahab MA, Boekhout T, et al. (2015)

- Classification of marine Ascomycota, Basidiomycota, Blastocladiomycota and Chytridiomycota. *Fungal Divers* 73:1–72. <https://doi.org/10.1007/s13225-015-0339-4>
- Kernaghan G, Hambling B, Fung M, Khasa D (2002) *In vitro* selection boreal ectomycorrhizal fungi for use in reclamation of saline-alkaline habitats. *Restoration Ecol* 10:43–51. <https://doi.org/10.1046/j.1526-100X.2002.10105.x>
- Khasa P, Hambling B, Kernaghan G, Fung M, Ngimbi E (2002) Genetic variability in salt tolerance of selected boreal woody seedlings. *For Ecol Manage* 165:257–269. [https://doi.org/10.1016/S0378-1127\(01\)00623-5](https://doi.org/10.1016/S0378-1127(01)00623-5)
- Langenfeld-Heyser R, Gao J, Ducic T, Tachd P, Lu C, Fritz E, et al. (2007) *Paxillus involutus* mycorrhiza attenuate NaCl-stress responses in the salt-sensitive hybrid poplar *Populus x canescens*. *Mycorrhiza* 17:121–131. <https://doi.org/10.1007/s00572-006-0084-3>
- Lee S, Calvo-Polanco M, Chung G, Zwiazek J (2010) Role of aquaporins in root water transport of ectomycorrhizal jack pine (*Pinus banksiana*) seedlings exposed to NaCl and fluoride. *Plant Cell Environ* 33:769–780 <http://doi.org/10.1111/j.1365-3040.2009.02103.x>
- Lehto T, Zwiazek JJ (2011) Ectomycorrhizas and water relations of trees: A review. *Mycorrhiza* 21:71–90 <http://doi.org/10.1007/s00572-010-0348-9>
- Lenoir I, Fontaine J, Lounès-Hadj Sahraoui A (2016) Arbuscular mycorrhizal fungal responses to abiotic stresses: A review. *Phytochem* 123:4–15. <https://doi.org/10.1016/j.phytochem.2016.01.002>
- Li J, Bao S, Zhang Y, Ma X, Mishra-Knyrim M, Sun J, et al. (2012) *Paxillus involutus* strains MAJ and NAU mediate K⁺/Na⁺ homeostasis in ectomycorrhizal *Populus × canescens* under sodium chloride stress. *Plant Physiol* 159:1771–1786. <https://doi.org/10.1104/pp.112.195370>
- Luo Z, Janz D, Jiang X, Gö C, Wildhagen H, Tan Y, et al. (2009) Upgrading root physiology for stress tolerance by ectomycorrhizas: Insights from metabolite and transcriptional profiling into reprogramming for stress anticipation. *Plant Physiol* 151:1902–1917. <https://doi.org/10.1104/pp.109.143735>
- Luo Z, Li K, Gai Y, Göbel C, Wildhagen H, Jiang X, et al. (2011) The ectomycorrhizal fungus (*Paxillus involutus*) modulates leaf physiology of poplar towards improved salt tolerance. *Environ Exper Bot* 72:304–311. <https://doi.org/10.1016/j.envexpbot.2011.04.008>
- Ma X, Sun M, Sa G, Zhang Y, Li J, Sun J, et al. (2014) Ion fluxes in *Paxillus involutus*-inoculated roots of *Populus canescens* under saline stress. *Environ Exper Bot* 108:99–108. <https://doi.org/10.1016/j.envexpbot.2013.11.016>
- Marjanović Ž, Uehlein N, Kaldenhoff R, Zwiazek JJ, Weiss M, Hampp R, Nehls U (2005) Aquaporins in poplar: What a difference a symbiont makes! *Planta* 222:258–268. <http://doi.org/10.1007/s00425-005-1539-z>
- Matsuda Y, Sugiyama F, Nakanishi K, Ito S (2006) Effects of sodium chloride on growth of ectomycorrhizal fungal isolates in culture. *Mycoscience* 47:212–217. <https://doi.org/10.1007/S10267-006-0298-4>
- Muhsin TM, Zwiazek JJ (2002a) Colonization with *Hebeloma crustuliniforme* increases water conductance and limits shoot sodium uptake in white spruce (*Picea glauca*) seedlings. *Plant Soil* 238:217–225. <https://doi.org/10.1023/A:1014435407735>
- Muhsin TM, Zwiazek JJ (2002b) Ectomycorrhizas increase apoplastic water transport and hydraulic conductivity in *Ulmus americana* seedlings. *New Phytol* 153:153–158. <http://doi.org/10.1046/j.0028-646X.2001.00297.x>
- Munns R, Tester M (2008) Mechanisms of salinity tolerance. *Annu Rev Plant Biol* 59:651–

506 681. <https://doi.org/10.1146/annurev.arplant.59.032607.092911>

507 Nass R, Cunningham KW, Rao R (1997) Intracellular sequestration of sodium by a novel
508 Na^+/H^+ exchanger in yeast is enhanced by mutations in the plasma membrane H^+ -ATPase.
509 Insights into mechanisms of sodium tolerance. J Biol Chem 272:26145–26152.
510 <https://doi.org/10.1074/jbc.272.42.26145>

511 Navarro-Ródenas A, Bárzana G, Nicolás E, Carra A, Schubert A, Morte A (2013) Expression
512 analysis of aquaporins from desert truffle mycorrhizal symbiosis reveals a fine-tuned
513 regulation under drought. Mol Plant Microbe Interac 26:1068–1078.
514 <https://doi.org/10.1094/MPMI-07-12-0178-R>.

515 Navarro-Ródenas A, Xu H, Kemppainen M, Pardo AG, Zwiazek JJ (2015) *Laccaria bicolor*
516 aquaporin *LbAQP1* is required for Hartig net development in trembling aspen (*Populus*
517 *tremuloides*). Plant Cell Environ 38:2475–2486. <http://doi.org/10.1111/pce.12552>

518 Nehls U, Dietz S (2014) Fungal aquaporins: cellular functions and ecophysiological
519 perspectives. Appl Microbiol Biotechnol 98:8835–8851. [https://doi.org/10.1007/s00253-](https://doi.org/10.1007/s00253-014-6049-0)
520 014-6049-0

521 Nguyen H, Calvo-Polanco M, Zwiazek JJ (2006) Gas exchange and growth responses of
522 ectomycorrhizal *Picea mariana*, *Picea glauca*, and *Pinus banksiana* seedlings to NaCl and
523 Na_2SO_4 . Plant Biol 8:646–652. <https://doi.org/10.1055/s-2006-924106>

524 Obase K, Lee JK, Lee SK, Lee SY, Chun KW (2010) Variation in sodium chloride resistance
525 of *Cenococcum geophilum* and *Suillus granulatus* isolates in liquid culture. Mycobiol
526 38:225–228. <https://doi.org/10.4489/MYCO.2010.38.3.225>

527 Onwuchekwa N, Zwiazek J, Quoreshi A, Khasa D (2014) Growth of mycorrhizal jack pine
528 (*Pinus banksiana*) and white spruce (*Picea glauca*) seedlings planted in oil sands reclaimed
529 areas. Mycorrhiza 24:431–441. <https://doi.org/10.1007/s00572-014-0555-x>

530 Ottow E, Brinker M, Teichmann T, Fritz E, Kaiser W, Brosché M, et al. (2005) *Populus*
531 *euphratica* displays apoplastic sodium accumulation, osmotic adjustment by decreases in
532 calcium and soluble carbohydrates, and develops leaf succulence under salt stress. Plant
533 Physiol 139:1762–1772. <https://doi.org/10.1104/pp.105.069971>

534 Parihar P, Singh S, Singh R, Singh VP, Prasad SM (2015) Effect of salinity stress on plants and
535 its tolerance strategies: a review. Environ Sci Pollut Res 22:4056–4075.
536 <https://doi.org/10.1007/s11356-014-3739-1>

537 Peter M, Kohler A, Ohm RA, Kuo A, Krützmann J, Morin E, et al. (2016) Ectomycorrhizal
538 ecology is imprinted in the genome of the dominant symbiotic fungus *Cenococcum*
539 *geophilum*. Nature Comm 7:12662. <http://dx.doi.org/10.1038/ncomms12662>

540 Porcel R, Aroca R, Ruiz-Lozano J (2012) Salinity stress alleviation using arbuscular
541 mycorrhizal fungi. A review. Agron Sustain Dev 32:181–200.
542 <http://dx.doi.org/10.1007/s13593-011-0029-x>

543 Proft M, Struhl K (2004) MAP kinase-mediated stress relief that precedes and regulates the
544 timing of transcriptional induction. Cell 118:351–361.
545 <https://doi.org/10.1016/j.cell.2004.07.016>

546 Renault S, Paton E, Nilsson G, Zwiazek JJ, MacKinnon MD (1999) Response of boreal plants
547 to high salinity oil sands tailings water. J Environ Qual 28:1957–1962.
548 <https://doi.org/10.2134/jeq1999.00472425002800060035x>

549 Rillig M, Mummey DL (2006) Mycorrhizas and soil structure. New Phytol 171:41–53.
550 <http://doi.org/10.1111/j.1469-8137.2006.01750.x>

551 Rodriguez-Navarro A (2000) Potassium transport in fungi and plants. Biochim Biophys Acta

1469:1–30. [http://doi.org/https://doi.org/10.1016/S0304-4157\(99\)00013-1](http://doi.org/https://doi.org/10.1016/S0304-4157(99)00013-1)

Rodriguez-Navarro A, Benito B (2010) Sodium or potassium efflux ATPase a fungal, bryophyte, and protozoal ATPase. *Biochim Biophys Acta* 1798:1841–1853. <https://doi.org/10.1016/j.bbamem.2010.07.009>

Shabala S (2017) Signalling by potassium: another second messenger to add to the list? *J Exp Bot* 68:4003–4007. <https://doi.org/10.1093/jxb/erx238>

Shabala S, Demidchik V, Shabala L, Cuin TA, Smith SJ, Miller AJ, et al. (2006) Extracellular Ca^{2+} ameliorates NaCl-induced K^{+} loss from *Arabidopsis* root and leaf cells by controlling plasma membrane K^{+} permeable channels. *Plant Physiol* 141:1653–1665. <https://doi.org/10.1104/pp.106.082388>

Shabala S, Cuin TA (2008) Potassium transport and plant salt tolerance. *Physiol Plant* 133:651–669. <https://doi.org/10.1111/j.1399-3054.2007.01008.x>

Singh M, Kumar J, Singh S, Singh VP, Prasad SM (2015) Roles of osmoprotectants in improving salinity and drought tolerance in plants: a review. *Rev Environ Sci Biotechnol* 14:407–426. <https://doi.org/10.1007/s11157-015-9372-8>

Smith SE, Read D (2008) Mycorrhizal symbiosis (3rd ed.). New York: Academic Press. <http://doi.org/http://dx.doi.org/10.1016/B978-012370526-6.50017-9>

Tang M, Sheng M, Chen H, Zhang F (2009) *In vitro* salinity resistance of three ectomycorrhizal fungi. *Soil Biol Biochem* 41:948–953. <https://doi.org/10.1016/j.soilbio.2008.12.007>

Tavakkoli E, Rengasamy P, McDonald GK (2010) High concentrations of Na^{+} and Cl^{-} ions in soil solution have simultaneous detrimental effects on growth of faba bean under salinity stress. *J Exp Bot* 61:4449–4459. <https://doi.org/10.1093/jxb/erq251>

Tresner HD, Hayes JA (1971) Sodium chloride tolerance of terrestrial fungi. *Appl Microbiol* 22:210–213

Verdoucq L, Maurel C (2018) Plant Aquaporins. *Adv Bot Res* 87:25–56. <https://doi.org/10.1016/bs.abr.2018.09.011>

Wang M, Zheng Q, Shen Q, Guo S (2013) The critical role of potassium in plant stress response. *Int J Mol Sci* 14:7370–7390. <https://doi.org/10.3390/ijms14047370>

Weissenhorn I (2002) Mycorrhiza and salt tolerance of trees. EU-project MYCOREM (QLK3-1999-00097) the use of mycorrhizal fungi in phytoremediation. Final Report of Partner 9. Projects January 2000 – December 2002. <https://www.servaplant.nl/new/wp-content/uploads/2015/10/MycoremReport.pdf>

Wu H (2018) Plant salt tolerance and Na^{+} sensing and transport. *Crop J* 6:215–225. <https://doi.org/10.1016/j.cj.2018.01.003>

Xu H, Kemppainen M, El Kayal W, Lee SH, Pardo AG, Cooke JEK, Zwiazek JJ (2015) Overexpression of *Laccaria bicolor* aquaporin JQ585595 alters root water transport properties in ectomycorrhizal white spruce (*Picea glauca*) seedlings. *New Phytol* 205: 757–770. <http://doi.org/10.1111/nph.13098>

Xu H, Cooke JEK, Kemppainen M, Pardo AG, Zwiazek JJ (2016) Hydraulic conductivity and aquaporin transcription in roots of trembling aspen (*Populus tremuloides*) seedlings colonized by *Laccaria bicolor*. *Mycorrhiza* 26:441–451. <https://doi.org/10.1007/s00572-016-0681-8>

Yi H, Calvo-Polanco M, MacKinnon MD, Zwiazek JJ (2008) Responses of ectomycorrhizal *Populus tremuloides* and *Betula papyrifera* seedlings to salinity. *Environ Exp Bot* 62:357–363. <https://doi.org/10.1016/j.envexpbot.2007.10.008>

- Zalar P, Hoog GS de, Schroers H-J, Crous PW, Groenewald JZ, Gunde-Cimerman N (2007) Phylogeny and ecology of the ubiquitous saprobe *Cladosporium sphaerospermum*, with descriptions of seven new species from hypersaline environments. *Stud Mycol* 58:157-183. <https://dx.doi.org/10.3114/sim.2007.58.06>
- Zhang H, Li J, Chen S, Lu C, Wang R, Dai S (2008) Effect of NaCl on growth and ion relations in two salt-tolerant strains of *Paxillus involutus*. *For Stud China* 10:95–100. <https://doi.org/10.1007/s11632-008-0025-7>

Legends of Figure & Table

Figure 1. Fungal membrane transport systems are probably involved in plant protection against salt stress through three different strategies. Three advantages of ectomycorrhizal symbiosis seem to be involved in a better stress tolerance of plants and membrane transport systems would play a role in all of them. First, Na⁺ exclusion from the plant would depend on the extrusion of Na⁺ from the extraradical mycelium to the soil through ENA ATPases and Nha transporters. The storage of the excess of Na⁺ within vacuoles by Vnx transporters could also reduce the amount of Na⁺ that reaches the plant tissues. The second strategy is the improvement of K⁺ nutrition by the fungus, thanks to Trk and HAK uptake transporters. This ion would be transferred to the plant through TOK and/or SKC channels. Further uptake through ACU ATPases or storage in vacuoles or vesicles through Nhx and Kha transporters are also likely to occur. Finally, ectomycorrhizal fungi would help maintaining aquaporin-mediated water transport, resulting in an improved water status of the symbiotic plant.

Table 1. Summary of the most studied salt-tolerant genera of ectomycorrhizal fungi: *Hebeloma*, *Laccaria*, *Paxillus*, *Pisolithus*, *Rhizopogon* and *Suillus* (Basidiomycota). *means that the studied isolate was not salt resistant.