Generation of a superconducting vortex via Néel skyrmions

J. Baumard, J. Cayssol, F. S Bergeret, Alexandre I. Buzdin

► To cite this version:

J. Baumard, J. Cayssol, F. S Bergeret, Alexandre I. Buzdin. Generation of a superconducting vortex via Néel skyrmions. *Physical Review B: Condensed Matter and Materials Physics* (1998-2015), 2019, 99 (1), pp.014511. 10.1103/PhysRevB.99.014511 . hal-02140463

HAL Id: hal-02140463

<https://hal.science/hal-02140463>

Submitted on 27 May 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Generation of a superconducting vortex via Néel skyrmions

J. Baumard,^{1,2} J. Cayssol,¹ F. S. Bergeret,^{2,3} and A. Buzdin^{1,4}

¹*Université de Bordeaux, CNRS, LOMA, UMR 5798, F-33405 Talence, France*

²*Donostia International Physics Center (DIPC), Manuel de Lardizabal 5, E-20018 San Sebastián, Spain*

³*Centro de Física de Materiales (CFM-MPC), Centro Mixto CSIC-UPV/EHU, Manuel de Lardizabal 4, E-20018 San Sebastián, Spain*

⁴*Sechenov First Moscow State Medical University, Moscow, 119991, Russia*

(Received 22 October 2018; revised manuscript received 20 December 2018; published 14 January 2019)

We consider a type-II superconducting thin film in contact with a Néel skyrmion. The skyrmion induces spontaneous currents in the superconducting layer, which under the right condition generates a superconducting vortex in the absence of an external magnetic field. We compute the magnetic field and current distributions in the superconducting layer in the presence of Néel skyrmion.

DOI: [10.1103/PhysRevB.99.014511](https://doi.org/10.1103/PhysRevB.99.014511)

I. INTRODUCTION

Superconductor-ferromagnet heterostructures [1–4] in the presence of spin-orbit and exchange interactions are attracting great interest due to the possible realization of topological qubits based on Majorana fermions [5–11] and the fact that such systems display unconventional magnetoelectric effects [12–24]. In particular, the interplay between spin-orbit coupling and a homogeneous Zeeman or exchange field may lead to spontaneous supercurrents in bulk superconductors and hybrid structures. From a SU(2) covariant formulation of spin-dependent fields, a spin-orbit coupling and homogeneous Zeeman field is equivalent to an inhomogeneous magnetic texture that, in combination with superconducting correlations, may support spontaneous currents under certain symmetry conditions [16,25,26].

Among inhomogeneous magnetic textures, skyrmions [27–29] have attracted interest because of their nanoscale dimension (1–100 nm), topological robustness, and the low current density needed to move them, which makes them good candidates as information carriers in future memory devices [30–34]. It has been shown that a skyrmion can be stabilized when proximity coupled to an *s*-wave superconductor [35,36]. In addition, such systems can induce spontaneous currents [37], Majorana bound states [38,39], Weyl points [40], or Yu-Shiba-Rusinov-like states [41]. Moreover, Hals *et al.* [42] studied the interaction between a skyrmion and a vortex by assuming that they are stabilized in the magnetic and superconducting layers.

In this paper, we investigate the formation of a composite topological excitation between a magnetic skyrmion and a superconducting vortex in a ferromagnet-superconductor (F-S) bilayer with Rashba spin-orbit coupling. In contrast to Ref. [42], the superconducting vortex is initially absent. We show that the generation of a vortex is via the magnetoelectric effect induced by the skyrmion in the presence of a sufficiently strong spin-orbit coupling. By evaluating the free energy of the F-S system, we derive the conditions required for the creation of this vortex, and compute the current and magnetic field distributions in the superconductor.

The paper is organized as follows. In Sec. II, we introduce the free energy describing the system. In Sec. III, we derive the vortex nucleation condition. The magnetic field and current distributions are provided in Sec. IV. We finally conclude and give some perspectives implied by our work in Sec. V.

II. SETUP AND FREE ENERGY

We consider a type-II superconducting thin film of thickness d_S , characterized by the coherence length ξ and the London penetration length λ . The superconductor is in contact with a ferromagnet of thickness d_F hosting a Néel skyrmion (Fig. 1). We assume that a two-dimensional spin-orbit interaction is present in the ferromagnetic layer and described by the Rashba constant α_R . The Néel skyrmion is characterized by the following spin profile [43]:

$$\vec{S}(\vec{r}) = \eta \sin \Theta(r) \vec{e}_r + \cos \Theta(r) \vec{e}_z, \quad (1)$$

where $\vec{e}_r$ is the radial unit vector and $\vec{e}_z$ the unit vector normal to the F and S layers. The profile function $\Theta(r)$ must obey the boundary conditions $\Theta(0) = \pi$ and $\Theta(\infty) = 0$. For the analytical calculations below, we assume that $\Theta(r) = \pi(1 - r/R)$ for $r < R$, and otherwise 0. Here R denotes the radius of the skyrmion. The constant $\eta = \pm 1$ describes the skyrmion winding. The sign of η , combined with the Rashba constant α_R determines the vortex polarity. In the following we consider $\alpha_R > 0$ and $\eta = -1$.

In principle both the direct electromagnetic coupling between the skyrmion and the superconductor [44], and the magnetic proximity effect may result in the nucleation of a vortex. In this paper we only focus on the proximity effect by assuming that the exchange field and spin-orbit interaction penetrate the superconductor over the atomic thickness a , where $a \ll d_S$. For a uniform ferromagnetic layer, if the magnetization is smaller than the first critical field, $\mu_0 M \ll H_{c1}$, the standard electromagnetic interaction cannot nucleate a vortex. Even if M exceeds H_{c1} , it is possible to avoid vortex formation by designing the F and S layers such that $\mu_0 M \ll H_{c1} \frac{d_S}{d_F}$, which may be easily fulfilled if the ferromagnetic layer is much thinner than the superconductor one.

FIG. 1. A thin superconducting film proximity coupled to a ferromagnetic layer hosting a Néel skyrmion. The F layer has thickness d_F and the S layer d_S .

Let us consider temperatures for which the superconductivity is well developed, i.e., $T \ll T_c$. The free energy of the F-S bilayer can be written as

$$F = F_0 + F_{sc} + F_L + F_{mag}, \quad (2)$$

where F_0 is the free energy in the absence of superconductivity and magnetic texture, and F_{sc} is the kinetic term related to the superconducting current energy. To derive the expression of the free energy, we use the London approach, which assumes that the generated current does not modify the modulus of the superconducting order parameter. The criterion of applicability of the London approach is well known (see for example Ref. [45]): the current density should be much smaller than the critical current density $j_c \propto \frac{\Phi_0}{\mu_0 \lambda^2 \xi}$, where $\Phi_0 = \frac{h}{2e}$ (with $e > 0$) is the superconducting quantum of flux. This is always the case for Abrikosov vortices, except the narrow core region. The computation of the current (see Sec. IV B) shows that this approach is completely justified to describe the vortex generation by the skyrmion while $R \gg \xi$. Moreover, since we assume that d_S is smaller than λ , the density of superconducting current energy is nearly constant through the width d_S so F_{sc} reads:

$$F_{sc} = \int \frac{1}{2 \mu_0 \lambda_{eff}} (\vec{\phi}(\vec{r}) - \vec{A}(\vec{r}))^2 d^2 \vec{r}, \quad (3)$$

where $\lambda_{eff} = \lambda^2/d_S$ is the effective screening length for the superconductor, $\vec{\phi}$ is the gradient of the local superconducting phase (multiplied by $\hbar/2e$), and $\vec{A}$ is the vector potential. Detailed calculations are provided in Appendix A. In the presence of a vortex, the expression for the vector $\vec{\phi}$ can be obtained from the London equation as [45]:

$$\vec{\phi}(r) = \frac{\Phi_0}{2 \pi r} \vec{e}_\theta, \quad (4)$$

where $\vec{e}_\theta$ is the unit orthoradial vector.

The third contribution to the free energy, F_L in Eq. (2), corresponds to the coupling energy between the superconductor and the magnetic order induced by the skyrmion. By proximity effect, the interplay between the exchange field and the Rashba spin-orbit interaction in the ferromagnetic layer

induces a spin polarization in the superconducting film. This may give rise, for example, to a spontaneous current in the bulk superconductor near the interface to F, in the absence of an external magnetic field [22,46]. For T close to T_c , such an interaction is described by the Lifshitz invariant [47–49]. At low temperatures and for $d_S \ll \lambda$, one can consider that the spin-orbit interaction is averaged over d_S . In this case the energy F_L can be written as:

$$F_L = \int \vec{\alpha}(r) \cdot (\vec{\phi}(\vec{r}) - \vec{A}(\vec{r})) d^2 \vec{r}, \quad (5)$$

where $\vec{\alpha}(r) = \alpha(r) \vec{e}_\theta = -\alpha_0 \sin \Theta(r) \vec{e}_\theta$ (see Appendix A). The constant α_0 incorporates the Rashba constant α_R , the exchange energy h_{ex} , the thickness of the superconducting film d_S and the proximity length a :

$$\alpha_0 \approx \frac{1}{4 \mu_0 e \lambda_{eff}} \frac{a}{d_S} \frac{\alpha_R h_{ex}}{v_F^2}. \quad (6)$$

The last component F_{mag} of the free energy Eq. (2):

$$F_{mag} = \int \frac{\vec{B}^2(\vec{r})}{2 \mu_0} d^3 \vec{r}, \quad (7)$$

represents the energy of the magnetic field $\vec{B} = \vec{\nabla} \times \vec{A}$.

The current density for $d_S \ll \lambda$ in the plane $z = 0$ is given by $\vec{j} = -\frac{\partial f}{\partial \vec{A}} \delta(z)$, where f is the free-energy density in the film: $F = \int f d^2 \vec{r}$. From the Maxwell-Ampere equation $\vec{\nabla} \times \vec{B} = \mu_0 \vec{j}$, we obtain a differential equation for $\vec{A}$, which can be solved in Fourier space [45]. The solution $\vec{A}_q$ of this equation, where $\vec{A}_q$ is the two-dimensional Fourier transform of $\vec{A}$ in the layer, is given by:

$$\vec{A}_q = \frac{1}{1 + 2 q \lambda_{eff}} (\vec{\phi}_q + \mu_0 \lambda_{eff} \vec{\alpha}_q), \quad (8)$$

where $\vec{\phi}_q$, $\vec{\alpha}_q$ are the two-dimensional Fourier transforms of $\vec{\phi}(r)$ and $\vec{\alpha}(r)$, respectively. This calculation is provided in Appendix B 1.

III. CREATION OF A SUPERCONDUCTING VORTEX

A. Magnetic field induced by the skyrmion

Because of the spontaneous current generated by the skyrmion in the superconducting film, a magnetic field is created perpendicular to the layer. We first consider that there is no vortex. In this case, the term proportional to $\vec{\phi}_q$ in Eq. (8) disappears and we can derive the expression of the magnetic field distribution $\vec{B}_s(r) = B_s(r) \vec{e}_z$ in the superconducting layer. Considering that the skyrmion is small compared to λ_{eff} and focusing on small distances r from the center of the skyrmion ($r \ll \lambda_{eff}$) one can write:

$$B_s(r) = -\frac{1}{2} \mu_0 \alpha_0 \int q \Gamma(q) J_0(q r) dq, \quad (9)$$

where $\Gamma(q) = \int_0^R r \sin(\pi \frac{r}{R}) J_1(q r) dr$ and $J_0(q r)$, $J_1(q r)$ are Bessel functions of first kind. This field distribution is represented by the blue line in Fig. 2. As expected, outside of the skyrmion B_s decreases and vanishes very fast. Moreover, one can check that the magnetic flux associated to B_s is equal to zero.

FIG. 2. Magnetic field distribution in the superconducting layer. The vortex nucleation condition for a vortex with vorticity 1 is fulfilled: $R = 50 \xi$, $\alpha_R = 0.1 v_F$, and $h_{\text{eff}} = 20 k_B T_c$. The blue line corresponds to the magnetic field distribution without vortex, whereas the orange one is in the presence of the vortex.

B. Vortex nucleation condition

The condition for the superconducting vortex creation can be derived by comparing the free energy of the system with and without a vortex. We replace $\vec{A}$ by its expression, [Eq. (8)], into the contributions Eq. (3), (5), (7) to the free energy. The resulting F can be written as a sum of three terms (see Appendix B 2):

$$F = F_v + F_s + F_{\text{int}}. \quad (10)$$

The first one, proportional to Φ_0^2 , describes the self-energy of the vortex. The second term, proportional to α_R^2 , describes the energy of the current induced by the skyrmion, whereas the third term, proportional to $\Phi_0 \alpha_R$, corresponds to the interaction energy between the vortex and such current.

Since $\xi \ll r \ll \lambda_{\text{eff}}$, we can write the self-energy of the vortex in the following way:

$$F_v = \frac{1}{\pi \mu_0 \lambda_{\text{eff}}} \left(\frac{\Phi_0}{2} \right)^2 \ln \left(2 \frac{\lambda_{\text{eff}}}{\xi} \right). \quad (11)$$

By assuming $R \ll \lambda_{\text{eff}}$, we can write the current energy and the interaction term as

$$F_s = -\frac{\mu_0}{8\pi} \int \alpha_q^2 dq; \quad (12)$$

$$F_{\text{int}} = -\frac{\Phi_0}{2\pi \mu_0 e \lambda_{\text{eff}}} \frac{a}{d_s} \frac{\alpha_R}{v_F} R. \quad (13)$$

The difference of free energy, $\Delta F = F - F_s$, between the states with and without vortex reads:

$$\Delta F = \frac{\Phi_0^2}{2\pi^2 \mu_0 \lambda_{\text{eff}}} \left[\frac{\pi}{2} \ln \left(2 \frac{\lambda_{\text{eff}}}{\xi} \right) - 0.180 \frac{h_{\text{ex}}}{k_B T_c} \frac{a}{d_s} \frac{\alpha_R}{v_F} \frac{R}{\xi} \right], \quad (14)$$

where k_B is the Boltzmann constant, T_c is the critical temperature of the superconductor and the coherence length is given by $\xi = 0.180 \frac{\hbar v_F}{k_B T_c}$.

The condition for the vortex nucleation is determined by the condition $\Delta F < 0$, which gives:

$$\frac{h_{\text{eff}}}{k_B T_c} \frac{\alpha_R}{v_F} \frac{R}{\xi} > \frac{\pi}{0.36} \ln \left(2 \frac{\lambda_{\text{eff}}}{\xi} \right), \quad (15)$$

where $h_{\text{eff}} = h_{\text{ex}} \frac{a}{d_s}$ is the average effective exchange energy, with $a \ll d$.

The condition Eq. (15) gives the features of the ferromagnetic layer required to induce a vortex inside the superconducting film without any external magnetic field. Qualitatively, this result shows that if α_R , h_{eff} , or R increase, so does the magnetic field B_s [Eq. (9)], thereby favoring the appearance of the vortex.

C. Multiquanta vortices

We now discuss the possibility of nucleating a vortex carrying n superconducting flux quanta $\Phi_0 = h/2e$, with $n > 1$. So far we only considered the case $n = 1$. Let F_n be the free energy in presence of a n -quanta vortex.

$$F_n = n^2 F_v + F_s + n F_{\text{int}}. \quad (16)$$

The optimal value of n can be estimated by minimizing F_n with respect to n :

$$n_{\text{op}} \approx -\frac{F_{\text{int}}}{2 F_v} = \frac{\pi}{0.72 \ln \left(2 \frac{\lambda_{\text{eff}}}{\xi} \right)} \frac{h_{\text{eff}}}{k_B T_c} \frac{\alpha_R}{v_F} \frac{R}{\xi}. \quad (17)$$

Therefore, upon raising the Rashba coupling and/or the exchange field, it is possible to stabilize a multiquanta vortex carrying the integer value of n_{op} superconducting flux quanta. However for simplicity in what follows we assume that the spin-orbit interaction is too weak to have a vortex with vorticity larger than 1.

IV. MAGNETIC FIELD AND CURRENT DISTRIBUTIONS

A. Magnetic field

The presence of the vortex modifies the magnetic field distribution. In addition to the component B_s , stemming from the current induced by the skyrmion in the superconducting layer, there is a term originating from the vortex itself. The total magnetic field distribution can thus be written as

$$B_z(r) = B_s(r) + B_v(r). \quad (18)$$

The term $B_v(r)$ is obtained from the first term of Eq. (8):

$$B_v(r) = \frac{\Phi_0}{4\pi \lambda_{\text{eff}} r}, \quad (19)$$

for $\xi \ll r \ll \lambda_{\text{eff}}$.

The magnetic field distribution $B_z(r)$ is shown in Fig. 2 (orange line). It is assumed that the condition for appearance of a vortex is fulfilled. As expected, both B_z and B_s follow the spin direction of the skyrmion, with a sinusoidal-like shape: it is negative near the center, and positive for $r \gtrsim 0.65 R$. At $r = R$, the amplitude of the magnetic field decreases away from the skyrmion. The component B_s tends to zero very fast, whereas B_z vanishes far from the center. It decreases slowly because of the presence of the vortex, whose component B_v is proportional to $1/r$.

FIG. 3. (a) Current lines in the superconducting layer. The vortex nucleation condition for a vortex with vorticity 1 is fulfilled: $R = 50 \xi$, $\alpha_R = 0.1 v_F$, and $h_{\text{eff}} = 20 k_B T_c$. The dashed black lines represent the changes in the rotation direction of the current loops. The thickness of the lines represents the amplitude of the current. (b) Distribution of the current J in the superconducting layer.

B. Charge current

The current $\vec{J}$ in the superconducting layer is obtained from $\vec{J} = -\frac{\partial f}{\partial \vec{A}}$ (see Appendix C 2). As with the magnetic field distribution, it can be written as the sum of two contributions: one induced directly by the skyrmion, and a second stemming from the vortex.

$$\vec{J}(r) = \vec{J}_s(r) + \vec{J}_v(r). \quad (20)$$

Under the same assumptions as before, $\xi \ll r \ll \lambda_{\text{eff}}$ and $R \ll \lambda_{\text{eff}}$, one obtains

$$\vec{J}_s(r) = -\alpha_0 \int q \Gamma(q) J_1(qr) dq \vec{e}_\theta; \quad (21)$$

$$\vec{J}_v(r) = \frac{\Phi_0}{2\pi \mu_0 \lambda_{\text{eff}} r} \vec{e}_\theta. \quad (22)$$

The current lines in the film are shown in Fig. 3(a). We use the same parameters as in (Fig. 2).

Around the vortex ($r < 0.20 R$), the current is dominated by the contribution from the vortex and flows anticlockwise. As it can be seen Fig. 3(b) in this region the current is positive and decreases like $1/r$. For larger values of r within the skyrmion ($0.20 R < r < 0.95 R$), the current distribution has a sinusoidal shape, and is dominated by the contribution from the skyrmion. In this region, the current loops are clockwise. Finally, for $r > 0.95 R$, the current is again dominated by the contribution from the vortex. It decreases slowly with distance, and tends to zero far from the skyrmion.

V. CONCLUSION

We have shown that the spontaneous current induced by the skyrmion in the superconducting thin film gives rise to a magnetic field perpendicular to the layer. If the Rashba coupling exceeds a threshold value [given by Eq. (15)], the skyrmion can nucleate a superconducting vortex by magnetoelectric proximity effect in the absence of an applied external field. The vorticity is determined by the sign of the spin-orbit interaction and the skyrmion winding. Finally we outline some perspectives implied by our calculations. Even if the Rashba coupling threshold condition is not reached, it is possible to nucleate vortices in S simply by applying an external magnetic field (larger than H_{c1}), or in the case when vortices are created directly via the electromagnetic coupling with the ferromagnetic layer. In the former situation (external magnetic field), our free-energy calculations demonstrate an attractive coupling, which will pin vortices to the skyrmion for one orientation of the magnetic field. For the opposite orientation, the vortices should be pushed away by the skyrmion. Such decoration/antidecoration of the skyrmion by vortices can be, in principle, detected experimentally. Finally, we also stress that the inverse effect, namely the nucleation of a skyrmion via the proximity of a superconducting vortex, is also suggested by our results, a strong effect that could in principle be observed experimentally via magnetic force microscopy or topological Hall effect in systems such as Nb/Co/Pt [50].

ACKNOWLEDGMENTS

The authors gratefully acknowledge J. W. A. Robinson for his useful remarks and suggestions. This work was supported by EU Network COST CA16218 (NANOCOBYBRI) and the French ANR project SUPERTRONICS and OPTOFLUXONICS (A.B. and J.C.). J.B. and F.S.B. acknowledge funding by the Spanish Ministerio de Economía y Competitividad (MINECO) (Projects No. FIS2014-55987-P and No. FIS2017-82804-P). J.B. acknowledges the financial support from the Initiative d'Excellence (IDEX) of the Université de Bordeaux.

APPENDIX A: DERIVATION OF THE MAGNETOELECTRIC ENERGY F_L

We derive the expression of the coupling energy between the superconductor and magnetic order induced by the

skyrmion. We start from the Ginzburg-Landau free energy

$$F^{\text{GL}} = F_0^{\text{GL}} + \int \frac{1}{4m} |\hat{D}\Psi|^2 d^3\vec{r} + F_L^{\text{GL}}, \quad (\text{A1})$$

where $\hat{D} = (-i\hbar \vec{\nabla} + 2e\vec{A})$ is the gauge-invariant momentum operator and $\Psi = |\Psi|e^{i\vec{q}\cdot\vec{r}}$. The term F_0^{GL} contains all the terms without derivative of Ψ . The Lifshitz invariant F_L^{GL} reads [47–49]:

$$F_L^{\text{GL}} = \int \varepsilon(r)(\vec{e}_z \times \vec{S}) \cdot [\Psi^* \hat{D}\Psi + \text{H.c.}] d^3\vec{r}. \quad (\text{A2})$$

One can derive an estimate of $\varepsilon(r)$, which is constant in the region where the spin-orbit interaction is present and null elsewhere.

To obtain the expression of the wave vector $\vec{q}$, we must minimize F^{GL} with respect to $\vec{q}$ for $\vec{A} = \vec{0}$. We get:

$$q = -\frac{4m}{\hbar} \varepsilon. \quad (\text{A3})$$

From Ref. [51], we have an estimate of q :

$$q = \frac{\alpha_R h_{\text{ex}}}{\hbar v_F^2}. \quad (\text{A4})$$

By comparing the expressions (A3) and (A4), one obtains an estimate of ε :

$$\varepsilon = -\frac{\alpha_R h_{\text{ex}}}{4m v_F^2}. \quad (\text{A5})$$

For $T \ll T_c$, superconductivity is well developed. We then rewrite the free energy (A1) in the London approach by noticing that $\Psi = |\Psi|e^{i\varphi}$, where $|\Psi|$ is constant and such that $|\Psi|^2 = \frac{1}{2}n_s$ where n_s is the density of superconducting electrons. Thus, the free energy becomes:

$$F = F_0^{\text{L}} + \int \frac{e^2 n_s}{2m} (\vec{\phi} - \vec{A})^2 d^3\vec{r} - e n_s \varepsilon \int (\vec{e}_z \times \vec{S}) \cdot (\vec{\phi} - \vec{A}) d^3\vec{r}, \quad (\text{A6})$$

where $\vec{\phi} = -\frac{\Phi_0}{2\pi} \vec{\nabla}\varphi$ and $F_0^{\text{L}} = F_0 + F_{\text{mag}}$. In what follows, F_0 will be omitted.

We introduce the London coherence length:

$$\lambda^2 = \frac{m}{\mu_0 n_s e^2}. \quad (\text{A7})$$

Considering that $d_s \ll \lambda$, the quantity $\vec{\phi} - \vec{A}$ is almost constant over d_s . We emphasize that the spin-orbit interaction and the exchange field penetrate the superconducting layer over a distance a , corresponding to the atomic thickness. We also assume that the magnetization in the ferromagnetic layer is weak, thus the Zeeman field is negligible compared to the exchange field. Then we can compute the integrals of Eq. (A6) over the z direction:

$$F = F_{\text{mag}} + \frac{d_s}{2\mu_0 \lambda^2} \int (\vec{\phi} - \vec{A})^2 d^2\vec{r} - e n_s \varepsilon a \int (\vec{e}_z \times \vec{S}) \cdot (\vec{\phi} - \vec{A}) d^2\vec{r}. \quad (\text{A8})$$

Introducing the effective screening length λ_{eff} , one can notice that the second term of the free energy (A8) is exactly the

superconducting current energy F_{sc} [Eq. (3)], whereas the third term corresponds to the magnetoelectric energy F_L [Eq. (5)].

APPENDIX B: FINAL EXPRESSION OF THE FREE ENERGY F

1. Derivation of the vector potential $\vec{A}$

In this section, we derive the expression of the vector potential $\vec{A}$. Let f be the free-energy density per unit surface.

$$f = \frac{1}{2\mu_0 \lambda_{\text{eff}}} (\vec{\phi} - \vec{A})^2 + \vec{\alpha}(r) \cdot (\vec{\phi} - \vec{A}) + \int \frac{B^2}{2\mu_0} dz. \quad (\text{B1})$$

The current density for $d_s \ll \lambda$ in the plane $z = 0$ is given by

$$\vec{j} = -\frac{\partial f}{\partial \vec{A}} \delta(z) = \frac{1}{\mu_0 \lambda_{\text{eff}}} (\vec{\phi} - \vec{A}) \delta(z) + \vec{\alpha}(r) \delta(z). \quad (\text{B2})$$

We recall the Maxwell-Ampere equation in the London gauge:

$$\mu_0 \vec{j} = \vec{\nabla} \times \vec{B} = -\Delta \vec{A}. \quad (\text{B3})$$

By replacing the expression of the current density (B2) into the Maxwell-Ampere equation [Eq. (B3)], one gets:

$$-\Delta \vec{A} + \frac{1}{\lambda_{\text{eff}}} \vec{A} \delta(z) = \frac{1}{\lambda_{\text{eff}}} \vec{\phi} \delta(z) + \mu_0 \vec{\alpha}(r) \delta(z). \quad (\text{B4})$$

We introduce the following three- and two-dimensional Fourier transforms:

$$\vec{A}_{qk} = \int \vec{A}(\vec{r}, z) e^{i(\vec{q}\cdot\vec{r} + kz)} d^2\vec{r} dz; \quad (\text{B5})$$

$$\vec{A}_q = \frac{1}{2\pi} \int \vec{A}_{qk} dk = \int \vec{A}(\vec{r}) e^{i\vec{q}\cdot\vec{r}} d^2\vec{r}; \quad (\text{B6})$$

$$\vec{\phi}_q = \int \vec{\phi}(r) e^{i\vec{q}\cdot\vec{r}} d^2\vec{r} = i \frac{\Phi_0}{q} \vec{e}_\perp; \quad (\text{B7})$$

$$\vec{\alpha}_q = \int \vec{\alpha}(r) e^{i\vec{q}\cdot\vec{r}} d^2\vec{r} = i \alpha_q \vec{e}_\perp, \quad (\text{B8})$$

with $\alpha_q = 2\pi \int_0^\infty r \alpha(r) J_1(qr) dr$ where $J_1(qr)$ is a Bessel function of first kind. The unit vector $\vec{e}_\perp$ is represented in Fig. 4.

FIG. 4. The sets of coordinates $(\vec{e}_r, \vec{e}_\theta)$ and $(\vec{e}_q, \vec{e}_\perp)$, respectively, corresponding to the real space and the Fourier space.

By taking the Fourier transform of Eq. (B4), one obtains the following equation:

$$\vec{A}_{qk} = \frac{1}{q^2 + k^2} \left[\frac{1}{\lambda_{\text{eff}}} (\vec{\phi}_q - \vec{A}_q) + \mu_0 \vec{\alpha}_q \right]. \quad (\text{B9})$$

After integration of Eq. (B9) over k , one recovers the expression of $\vec{A}_q$ [Eq. (8)].

2. Free energy F

In this section, we explain how we obtain the expression (10) for the free energy F and how we determine the value of η depending on the sign of α_R .

Each term of F [Eq. (2)] can be written in terms of $\vec{A}_q$, $\vec{\phi}_q$, and $\vec{\alpha}_q$:

$$F_{\text{sc}} = \frac{1}{(2\pi)^2} \frac{1}{2\mu_0 \lambda_{\text{eff}}} \int |\vec{\phi}_q - \vec{A}_q|^2 d^2 \vec{q}; \quad (\text{B10})$$

$$F_{\text{L}} = \frac{1}{(2\pi)^2} \int \vec{\alpha}_q^* \cdot (\vec{\phi}_q - \vec{A}_q) d^2 \vec{q}; \quad (\text{B11})$$

$$F_{\text{mag}} = \frac{1}{(2\pi)^3} \int \frac{|\vec{B}_{qk}|^2}{2\mu_0} d^2 \vec{q} dk, \quad (\text{B12})$$

where $\vec{B}_{qk}$ is the Fourier transform of $\vec{B}$, and is such that

$$|\vec{B}_{qk}|^2 = k^2 |\vec{A}_{qk}|^2 + |(\vec{q} \times \vec{A}_{qk}) \cdot \vec{e}_z|^2. \quad (\text{B13})$$

We replace $\vec{A}_q$ [Eq. (B6)] and $\vec{A}_{qk}$ [Eq. (B5)] by their expression in Eqs. (B10)–(B12). After integration, one obtains:

$$F_{\text{sc}} = \frac{\lambda_{\text{eff}}}{4\pi\mu_0} \int \frac{q}{(1 + 2q\lambda_{\text{eff}})^2} (2\Phi_0 - \mu_0 \alpha_q)^2 dq; \quad (\text{B14})$$

$$F_{\text{L}} = \frac{1}{2\pi} \int \frac{\lambda_{\text{eff}} q}{1 + 2q\lambda_{\text{eff}}} (2\Phi_0 - \mu_0 \alpha_q) \alpha_q dq; \quad (\text{B15})$$

$$F_{\text{mag}} = \frac{1}{2\pi\mu_0} \int \frac{(\Phi_0 + \mu_0 \lambda_{\text{eff}} q \alpha_q)^2}{(1 + 2q\lambda_{\text{eff}})^2} dq. \quad (\text{B16})$$

Using Eq. (B14)–(B16), one can rewrite F as a sum of three terms. The first one, proportional to Φ_0^2 , is called F_v [Eq. (11)]. The second term is proportional to α_q^2 , and corresponds to F_s [Eq. (12)], and the third one, depending on the product $\Phi_0 \alpha_q$, is called F_{int} [Eq. (13)].

In order to nucleate a vortex in the superconducting layer, the difference of energy $\Delta F = F - F_s$ must be negative:

$$\Delta F = \frac{\Phi_0^2}{2\pi^2 \mu_0 \lambda_{\text{eff}}} \left[\frac{\pi}{2} \ln \left(2 \frac{\lambda_{\text{eff}}}{\xi} \right) + 0.180 \eta \frac{h_{\text{ex}}}{k_B T_c} \frac{a}{d_S} \frac{\alpha_R}{v_F} \frac{R}{\xi} \right]. \quad (\text{B17})$$

The condition $\Delta F < 0$ requires that $\eta \alpha_R < 0$: the polarity of the vortex is determined by the spin-orbit interaction and the skyrmion winding. We thus consider $\alpha_R > 0$, which implies $\eta = -1$.

APPENDIX C: MAGNETIC FIELD AND CURRENT DISTRIBUTIONS

1. Perpendicular magnetic field distribution $B(r)$

We compute the normal component $B_z(r)$ of the magnetic field distribution, which is given by $B_z(r) = (\vec{\nabla} \times \vec{A}) \cdot \vec{e}_z$. In the Fourier space, this relation becomes

$$B_{qk}^z = -i(\vec{q} \times \vec{A}_{qk}) \cdot \vec{e}_z, \quad (\text{C1})$$

where $\vec{A}_{qk}$ is obtained from Eq. (B9) after replacing $\vec{A}_q$ by its expression [Eq. (8)]. Thus

$$B_{qk}^z = \frac{1}{q^2 + k^2} \frac{2q}{1 + 2q\lambda_{\text{eff}}} (\Phi_0 + q\lambda_{\text{eff}} \mu_0 \alpha_q). \quad (\text{C2})$$

After integration over k , we get the component B_q^z , which is the Fourier transform of $B_z(r)$:

$$B_q^z = \frac{1}{1 + 2q\lambda_{\text{eff}}} (\Phi_0 + q\lambda_{\text{eff}} \mu_0 \alpha_q). \quad (\text{C3})$$

After taking the inverse Fourier transform of Eq. (C3), in the approximation $q \gg \lambda_{\text{eff}}^{-1}$ one obtains the expression (18). Notice that Eq. (19) was previously obtained in Ref. [45].

2. Current in the superconducting layer

In this section, we derive the expression of the current $\vec{J}(r)$ in the superconducting layer, obtained from the free-energy density [Eq. (B1)]:

$$\vec{J}(r) = -\frac{\partial f}{\partial \vec{A}} = \frac{1}{\mu_0 \lambda_{\text{eff}}} (\vec{\phi} - \vec{A}) + \vec{\alpha}(r). \quad (\text{C4})$$

In the Fourier space and after replacing $\vec{A}_q$ by its expression [Eq. (8)], the current becomes:

$$\vec{J}_q = \frac{2i}{1 + 2q\lambda_{\text{eff}}} \left(\frac{\Phi_0}{\mu_0} + q\lambda_{\text{eff}} \alpha_q \right) \vec{e}_\perp. \quad (\text{C5})$$

Taking into account that $\vec{e}_\perp = -\sin \theta \vec{e}_r + \cos \theta \vec{e}_\theta$ (see Fig. 4), one can perform the inverse Fourier transform of (C5). In the approximation $q \gg \lambda_{\text{eff}}^{-1}$, we finally obtain Eq. (20).

- [1] A. I. Buzdin, *Rev. Mod. Phys.* **77**, 935 (2005).
- [2] F. S. Bergeret, A. F. Volkov, and K. B. Efetov, *Rev. Mod. Phys.* **77**, 1321 (2005).
- [3] J. Linder and J. W. A. Robinson, *Nat. Phys.* **11**, 307 (2015).
- [4] M. G. Blamire and J. W. A. Robinson, *J. Phys. Condens. Matter* **26**, 453201 (2014).

- [5] A. Y. Kitaev, *Phys. Usp.* **44**, 131 (2001).
- [6] C. Nayak, S. H. Simon, A. Stern, M. Freedman, and S. Das Sarma, *Rev. Mod. Phys.* **80**, 1083 (2008).
- [7] Y. Oreg, G. Refael, and F. von Oppen, *Phys. Rev. Lett.* **105**, 177002 (2010).
- [8] R. M. Lutchyn, J. D. Sau, and S. Das Sarma, *Phys. Rev. Lett.* **105**, 077001 (2010).

- [9] C.-T. Wu, B. M. Anderson, W.-H. Hsiao, and K. Levin, *Phys. Rev. B* **95**, 014519 (2017).
- [10] A. M. Black-Schaffer and J. Linder, *Phys. Rev. B* **84**, 180509 (2011).
- [11] J. Alicea, *Rep. Prog. Phys.* **75**, 076501 (2012).
- [12] A. Buzdin, *Phys. Rev. Lett.* **101**, 107005 (2008); F. Konschelle and A. Buzdin, *ibid.* **102**, 017001 (2009).
- [13] T. Ojanen, *Phys. Rev. Lett.* **109**, 226804 (2012).
- [14] S. S. Pershoguba, K. Björnson, A. M. Black-Schaffer, and A. V. Balatsky, *Phys. Rev. Lett.* **115**, 116602 (2015).
- [15] E. M. Chudnovsky, *Phys. Rev. B* **95**, 100503 (2017).
- [16] F. Konschelle, I. V. Tokatly, and F. S. Bergeret, *Phys. Rev. B* **92**, 125443 (2015); *Europhys. Lett.* **110**, 57005 (2015).
- [17] Y. V. Fominov, A. F. Volkov, and K. B. Efetov, *Phys. Rev. B* **75**, 104509 (2007).
- [18] I. Kulagina and J. Linder, *Phys. Rev. B* **90**, 054504 (2014).
- [19] F. Konschelle, I. V. Tokatly, and F. S. Bergeret, *Phys. Rev. B* **94**, 014515 (2016).
- [20] K. Halterman, O. T. Valls, and P. H. Barsic, *Phys. Rev. B* **77**, 174511 (2008).
- [21] V. Braude and Y. V. Nazarov, *Phys. Rev. Lett.* **98**, 077003 (2007).
- [22] S. Mironov and A. Buzdin, *Phys. Rev. Lett.* **118**, 077001 (2017).
- [23] M. A. Silaev, I. V. Tokatly, and F. S. Bergeret, *Phys. Rev. B* **95**, 184508 (2017).
- [24] J. W. A. Robinson, J. D. S. Witt, and M. G. Blamire, *Science* **329**, 59 (2010).
- [25] F. S. Bergeret and I. V. Tokatly, *Phys. Rev. B* **89**, 134517 (2014).
- [26] A. S. Mel'nikov, A. V. Samokhvalov, S. M. Kuznetsova, and A. I. Buzdin, *Phys. Rev. Lett.* **109**, 237006 (2012).
- [27] A. N. Bogdanov and D. A. Yablonskii, *Zh. Eksp. Teor. Fiz.* **95**, 178 (1989) [*Sov. Phys. JETP* **68**, 101 (1989)].
- [28] U. K. Rößler, A. A. Leonov, and A. N. Bogdanov, *J. Phys.: Conf. Ser.* **303**, 012105 (2011).
- [29] A. O. Leonov, T. L. Monchesky, N. Romming, A. Kubetzka, A. N. Bogdanov, and R. Wiesendanger, *New J. Phys.* **18**, 065003 (2016).
- [30] F. Jonietz, S. Mühlbauer, C. Pfleiderer, A. Neubauer, W. Münzer, A. Bauer, T. Adams, R. Georgii, P. Böni, R. A. Duine, K. Everschor, M. Garst, and A. Rosch, *Science* **330**, 1648 (2010).
- [31] X. Z. Yu, N. Kanazawa, W. Z. Zhang, T. Nagai, T. Hara, K. Kimoto, Y. Matsui, Y. Onose, and Y. Tokura, *Nat. Commun.* **3**, 988 (2012).
- [32] N. S. Kiselev, A. N. Bogdanov, R. Schäfer, and U. K. Rößler, *J. Phys. D: Appl. Phys.* **44**, 392001 (2011).
- [33] J. Iwasaki, M. Mochizuki, and N. Nagaosa, *Nat. Nanotech.* **8**, 742 (2013).
- [34] A. Hrabec, J. Sampaio, M. Belmeguenai, I. Gross, R. Weil, S. M. Chérif, A. Stashkevich, V. Jacques, A. Thiaville, and S. Rohart, *Nat. Commun.* **8**, 15765 (2017).
- [35] A. A. Fraerman, I. R. Karetnikova, I. M. Nefedov, I. A. Shereshevskii, and M. A. Silaev, *Phys. Rev. B* **71**, 094416 (2005).
- [36] V. L. Vadimov, M. V. Sapozhnikov, and A. S. Mel'nikov, *Appl. Phys. Lett.* **113**, 032402 (2018).
- [37] D. S. Rabinovich, I. V. Bobkova, A. M. Bobkov, and M. A. Silaev, *Phys. Rev. B* **98**, 184511 (2018).
- [38] G. Yang, P. Stano, J. Klinovaja, and D. Loss, *Phys. Rev. B* **93**, 224505 (2016).
- [39] U. Güngördü, S. Sandhoefner, and A. A. Kovalev, *Phys. Rev. B* **97**, 115136 (2018).
- [40] R. Takashima and S. Fujimoto, *Phys. Rev. B* **94**, 235117 (2016).
- [41] S. S. Pershoguba, S. Nakosai, and A. V. Balatsky, *Phys. Rev. B* **94**, 064513 (2016).
- [42] K. M. D. Hals, M. Schechter, and M. S. Rudner, *Phys. Rev. Lett.* **117**, 017001 (2016).
- [43] N. Nagaosa and Y. Tokura, *Nat. Nanotech.* **8**, 899 (2013).
- [44] I. F. Lyuksyutov and V. L. Pokrovsky, *Adv. Phys.* **54**, 67 (2005).
- [45] P. G. D. Gennes, *Superconductivity of Metals and Alloys* (Benjamin, New York, 1966).
- [46] V. M. Edelstein, *Phys. Rev. Lett.* **75**, 2004 (1995).
- [47] V. M. Edelstein, *J. Phys. Condens. Matter* **8**, 339 (1996).
- [48] K. V. Samokhin, *Phys. Rev. B* **70**, 104521 (2004).
- [49] R. P. Kaur, D. F. Agterberg, and M. Sigrist, *Phys. Rev. Lett.* **94**, 137002 (2005).
- [50] A. Hrabec, N. A. Porter, A. Wells, M. J. Benitez, G. Burnell, S. McVitie, D. McGrouther, T. A. Moore, and C. H. Marrows, *Phys. Rev. B* **90**, 020402 (2014).
- [51] O. Dimitrova and M. V. Feigel'man, *Phys. Rev. B* **76**, 014522 (2007).