



**HAL**  
open science

## Malate utilization by *Schizosaccharomyces pombe*

Patricia Taillandier, Jean-Pierre Riba, Pierre Strehaiano

► **To cite this version:**

Patricia Taillandier, Jean-Pierre Riba, Pierre Strehaiano. Malate utilization by *Schizosaccharomyces pombe*. *Biotechnology Letters*, 1988, 10 (7), pp.469-472. 10.1007/BF01027058 . hal-02143216

**HAL Id: hal-02143216**

**<https://hal.science/hal-02143216>**

Submitted on 29 May 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## Open Archive Toulouse Archive Ouverte (OATAO)

OATAO is an open access repository that collects the work of some Toulouse researchers and makes it freely available over the web where possible.

This is an author's version published in: <https://oatao.univ-toulouse.fr/22868>

**Official URL** : <https://doi.org/10.1007/BF01027058>

**To cite this version :**

Taillandier, Patricia  and Riba, Jean-Pierre  and Strehaiano, Pierre  *Malate utilization by Schizosaccharomyces pombe*. (1988) *Biotechnology Letters*, 10 (7). 469-472. ISSN 0141-5492

Any correspondence concerning this service should be sent to the repository administrator:  
[tech-oatao@listes-diff.inp-toulouse.fr](mailto:tech-oatao@listes-diff.inp-toulouse.fr)

# MALATE UTILIZATION BY *SCHIZOSACCHAROMYCES POMBE*

P. TAILLANDIER, J.P. RIBA, P. STREHAIANO (\*)  
ENSIGC. CHEMIN DE LA LOGE.31078 TOULOUSE CEDEX.

## SUMMARY

Malic acid degradation by *Schizosaccharomyces pombe* was studied in synthetic and natural media. Using different concentrations from 1 to 29 g/l, malic acid was degraded. Initial deacidification rate increased with initial malate concentration. During the stationary phase, *S. pombe* was able to decompose malate added to the medium before or after sugar exhaustion.

## INTRODUCTION

*Schizosaccharomyces* yeasts present the unusual property of being able to convert malic acid into ethanol with a great efficiency during fermentation.

Malic acid is one of the main organic acids in grapes. Generally, it is necessary to remove it from wines in order to improve their organoleptic qualities and to ensure their biological stability. One means of reducing grape musts acidity might be the use of *Schizosaccharomyces*. Some attempts have been made to deacidify grape musts ( GALLANDER, 1977 ; SNOW and GALLANDER, 1979; CHARPENTIER and al. , 1985 ) or wines ( BENDA, 1973 ) with *S. pombe*. Results were variable because of the broad variety of *S. pombe* strains and because of a lack of data about the yeast kinetic and physiologic behaviour.

In order to develop a suitable biological method for grape musts deacidification using *S. pombe* we had to improve our understanding of growth and malate degradation kinetics as well as malate metabolism.

The present study deals with the linkage between malate metabolism, cellular growth and sugar assimilation.

## MATERIALS AND METHODS

### Microorganism

The yeast strain used was the *Schizosaccharomyces pombe* strain isolated by Institut Coopératif du Vin of Montpellier ( ICV ).

### Culture medium

Two media were used:

- grape must ( Gros Manseng variety ) containing 200 g/l sugars (approximately 100 g/l glucose and 100 g/l fructose ) and 5 g/l malate.
- synthetic medium consisted of the following in g/l water: glucose 100; malate 1 to 29; yeast extract 4 ;  $\text{KH}_2\text{PO}_4$  5 ;  $\text{MgSO}_4$  0.4; pH 3.

### Culture conditions

Batch experiments were performed in 2 liters fermentors (SETRIC GENIE INDUSTRIEL. Toulouse). Temperature was regulated at 30° C. PH was not regulated. Aeration rate was 0.15 VVM. Agitation rate was 280 r.p.m.

### Analytical determinations

Biomass was estimated gravimetrically. Cell viability was determined by methylene blue staining. Glucose, fructose and malic acid were measured using enzymatic procedures of

Boehringer Mannheim. Ethanol was determined by gas chromatography.

## RESULTS AND DISCUSSION

Although malate utilization has been studied for some time, the metabolic pathway for malate degradation in *S. pombe* remains unknown. Energetic processes controlling malic acid transport and break down are still not elucidated. According to most authors, malate catabolism in *Schizosaccharomyces* is strictly linked with yeast cellular growth (DITTRICH, 1963; BIDAN, 1974), and with sugar consumption (PEYNAUD and al., 1964; BRUGIRARD and ROQUES, 1972; OSOTHSILP and SUBDEN, 1986).

### Malate tolerance

To investigate malate uptake in *S. pombe* (saturation kinetics or substrate inhibition), batch cultures were performed using synthetic media containing 18-29 g/l malic acid; fermentations with synthetic medium containing 1 to 16 g/l malate have been presented elsewhere (AURIOL and al., 1987).

Concentration of malic acid (1 to 29 g/l) did not affect growth, sugar assimilation or ethanol production; 100% of malate was degraded in each experiment. No substrate inhibition of deacidification could be found. Initial deacidification rate was increasing with malate concentration. Malate uptake in *S. pombe* did not follow saturation kinetics that would have indicated a carrier mediated malate transport.

Further experiments were conducted on media containing about 7 g/l malate that is the level occurring in acid grape musts.

### Relation between deacidification and cellular growth

As shown in previous work (AURIOL and al., 1987), when acid level was high, the entire amount of malate could not be metabolized during the growth phase. Therefore, during the stationary phase resting cells continued to break-down malic acid.

Likewise, when malate was added to a culture of *S. pombe* during the stationary phase, it was totally degraded at a high rate (figure 1).




Figure 1: Malic acid degradation during the stationary phase.

Grape musts deacidification method using *S. pombe* was tested in a "Jurançon" winery (Cave Coopérative des Producteurs de Jurançon). Grape musts were sulfited before inoculation with *S. pombe* ( $1 \cdot 10^6$  cells / ml) in order to delay the growth of natural *Saccharomyces cerevisiae* flora (figure 2).




Figure 2: Grape musts deacidification. Sugars consumption. Evolution of *Saccharomyces* and *Schizosaccharomyces* populations.

After about 50 hours, deacidification of musts started before sugar up-take. *Saccharomyces* yeasts started to multiply when the sugar concentration was decreasing. *S. pombe* concentration was never greatly increasing but its viability was always about 90%. In fact, it seemed that there was little or no *S. pombe* growth.

THIBAUT (1982) noted that malate degradation started before sugar assimilation during fermentation of grape musts with pure *S. pombe*.

These three kinds of experiments clearly demonstrated that with *S. pombe* deacidification and cellular growth are strictly no-linked.

#### Relation between deacidification and sugar consumption

Grape musts deacidification attempts made at GAN winery suggested that malate degradation and sugar utilization with *S. pombe* were not so strictly linked as it was previously thought. To verify this, malate was added repeatedly to a culture of *S. pombe* after all sugar has been consumed (figure 3).

The decomposition of malate by resting cells was observed before and after sugar exhaustion with a declining deacidification rate. At the same time, the viability was decreasing rapidly from sugar exhaustion, so the decline in deacidification rate may be due to a loss of yeast activity.

In a reference culture where no malate was added after sugar exhaustion, viability remained constant, equal to 95%. Consequently, it seemed that malic acid was responsible

for the decrease of viability when no sugar was present in the culture medium.

No extracellular deacidification activity was found in the supernatant of a *S. pombe* culture in synthetic medium. Despite their loss of viability, *S. pombe* cells were able to degrade malic acid efficiently without sugar consumption ( 20 g/l malate consumed during 150 hours after sugar exhaustion ).




Figure 3: Malic acid degradation during the stationary phase before and after sugar exhaustion

## CONCLUSION

The results presented in this communication clearly show that malate degradation by *S. pombe* is neither linked with growth nor with sugar assimilation as it was previously thought. When sugar is no longer present in the culture medium, viability of yeasts decreases whereas the deacidification activity remains. It seems that malic acid is responsible for the loss of viability in the absence of sugar. Since malate is not a source of energy and cannot be the only source of carbon (KUNKEE, 1967), we do not know why it is metabolized in a medium lacking sugar.

This about malic acid utilization by *S. pombe* can help us to determine the optimal conditions for continuous biological deacidification of grape musts.

## REFERENCES

- Auriol P., Tulasi S., Goma G., Strehaiano P. (1987). R.F.Oe. 108, 37-42.  
 Benda I. (1973). Vignes et Vins. N° spécial, 31-36.  
 Bidan P., Meyer J.P., Schaeffer A. (1974). Bull. O.I.V. 523, 682-706.  
 Brugirard A., Roques J. (1972). R.F.Oe. 13, 26-36.  
 Charpentier C., Feuillat M., Gerbaux V., Auther R. (1985). C.R. Acad. Agri.France. 71, 425-432.  
 Dittrich H.H. (1963). Wein Win. 18, 392-405.  
 Gallander J.F. (1978). Am. J. Enol. Vitic. 28, 2, 65-69.  
 Kunkee R.E. (1967). Adv. Appl. Microbiol. 9, 235-300.  
 Osothsilp C., Subden R.E. (1986). J. of Bacteriol. 168, 1439-1443.  
 Peynaud E., Domerg S., Boidron A.M., Lafon-Lafourcade S., Guimberteau G. (1964). Arch. Mikrobiol. 48, 150-165.  
 Snow P.G., Gallander J.F. (1979). Am. J. Enol. Vitic. 30, 1, 45-48.  
 Thibault P.A. (1982). Thèse INSA. Toulouse.