Root systems and quotients of deformations of simple singularities

Antoine Caradot

► To cite this version:

Antoine Caradot. Root systems and quotients of deformations of simple singularities. Journal of Algebra, 2019, 526, pp.382-422. 10.1016/j.jalgebra.2019.02.020 . hal-02162828

HAL Id: hal-02162828

<https://hal.science/hal-02162828>

Submitted on 22 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution - NonCommercial| 4.0 International License

Root Systems and Quotients of Deformations of Simple Singularities

Antoine Caradot

*Institut Camille Jordan, Université Claude Bernard Lyon 1,
 43 boulevard du 11 novembre 1918, F-69622 Villeurbanne Cedex, France*

Abstract

In this article we study quotients of deformations of simple singularities, and attempt to characterise them using subsystems of simple root systems. The quotient of a semiuniversal deformation of a simple singularity of inhomogeneous type B_r ($r \geq 2$), C_r ($r \geq 3$), F_4 or G_2 by the natural symmetry of the associated Dynkin diagram is a deformation of a simple singularity of homogeneous type $X = D_s$, E_6 or E_7 , but not semiuniversal anymore. Therefore not all subdiagrams of X appear as singular configurations of the fibers of the deformation. We propose a conjecture for the types of singular configurations in terms of sub-root systems of a root system of type X and prove it for types B_2, B_3, C_3, F_4 and G_2 .

Keywords: deformations of simple singularities, simple root systems, simple singularities of inhomogeneous types, singular configurations.

2010 MSC: 14B07, 17B22, 20G05.

Table of Contents

Introduction	2
1 Deformations of simple singularities	2
1.1 Simple singularities	2
1.2 Deformations of homogeneous simple singularities	4
1.3 Deformations of inhomogeneous simple singularities	5
2 Quotients of semiuniversal deformations of inhomogeneous simple singularities	5
3 Singular configurations and root systems	11
4 Proof of Theorem 3.2	13
4.1 Strategy of the proof	13
4.2 Case $A_3 - B_2 - D_4$	14
4.3 Case $A_5 - B_3 - D_5$	16
4.4 Case $D_4 - C_3 - D_6$	18
4.5 Case $D_4 - G_2 - E_6$	22
4.6 Case $D_4 - G_2 - E_7$	24
4.7 Case $E_6 - F_4 - E_7$	27
Conclusion	34
Acknowledgements	34

Email address: caradot@math.univ-lyon1.fr (Antoine Caradot)

Preprint submitted to Journal of Algebra

February 22, 2019

Introduction

Simple singularities have first been studied by F. Klein in [9], where he classified them as quotients of the complex plane by the action of a finite subgroup Γ of SU_2 . It was then showed by P. Du Val ([6]) that the exceptional divisors of the minimal resolution of the isolated singularity of such a quotient form an arrangement of projective lines whose dual graph is a simply-laced Dynkin diagram $\Delta(\Gamma)$, and so the quotient $\mathbb{C}^2/\Gamma$ is called a simple singularity of type $\Delta(\Gamma)$. P. Slodowy extended in 1978 (and subsequently in [15] Section 6.2) the definition of a simple singularity to the non simply-laced types by adding a second finite subgroup Γ' of SU_2 containing Γ as normal subgroup. Then $\Omega = \Gamma'/\Gamma$ acts on $\mathbb{C}^2/\Gamma$ and this action can be lifted to the minimal resolution of the singularity and induces an action on the exceptional divisors which corresponds to a group of automorphisms of the Dynkin diagram of $\mathbb{C}^2/\Gamma$.

Let $\alpha : X_\Gamma \rightarrow \mathfrak{h}/W$ be the semiuniversal deformation of a simple singularity of type $\Delta(\Gamma) = A_{2r-1}$ ($r \geq 2$), D_r ($r \geq 4$) or E_6 obtained by the construction of H. Cassens and P. Slodowy in [5], with $\mathfrak{h}$ and W being the Cartan subalgebra and the associated Weyl group of the simple Lie algebra $\mathfrak{g}$ of the same type. It was shown in [4] that the automorphism group Ω of the Dynkin diagram of $\mathfrak{g}$ acts on X_Γ and $\mathfrak{h}/W$ such that α is Ω -equivariant. A result of P. Slodowy then implies that taking the restriction α^Ω of α over the Ω -fixed points of $\mathfrak{h}/W$ leads to a semiuniversal deformation of a simple singularity which is inhomogeneous and of the same type as the folding of the root lattice of $\mathfrak{g}$ by Ω . As α is Ω -equivariant, there is an action of Ω on every fiber of α^Ω , and the quotient leads to a new morphism $\overline{\alpha}^\Omega$, which was shown to be a non-semiuniversal deformation of a simple singularity of homogeneous type $\Delta(\Gamma')$ (cf. [4]).

The aim of this article is to propose a characterisation of the types of singularities which appear in the fibers of $\overline{\alpha}^\Omega$ in terms of sub-root systems of a root system of type $\Delta(\Gamma')$. The special fiber $(\overline{\alpha}^\Omega)^{-1}(\overline{0})$ is a simple singularity of type $\Delta(\Gamma')$. But as $\overline{\alpha}^\Omega$ is not semiuniversal, all subdiagrams of the Dynkin diagram of type $\Delta(\Gamma')$ cannot be found as singular configurations of the fibers of $\overline{\alpha}^\Omega$. We conjecture that there exists a subset Θ of simple roots of the root system of type $\Delta(\Gamma')$ such that the Dynkin diagram associated to the singular configuration of any fiber of $\overline{\alpha}^\Omega$ is a subdiagram of the Dynkin diagram of type $\Delta(\Gamma')$ containing the vertices associated to Θ . Furthermore, all such subdiagrams are realised as singular configurations of some fibers of $\overline{\alpha}^\Omega$.

In the first section we will present the construction of semiuniversal deformations of the simple homogeneous ([5]) and inhomogeneous ([4]) singularities. In the second section we study the regularity of the fibers of $\overline{\alpha}^\Omega$ and prove that they are always all singular. The third and fourth sections are devoted to the main conjecture as well as its proof for types B_2 , B_3 , C_3 , F_4 and G_2 .

Throughout this article the base field is the complex number field $\mathbb{C}$. Furthermore, a Dynkin diagram and its type will often be designated by the same symbol Δ .

1. Deformations of simple singularities

1.1. Simple singularities

1.1.1. Simple singularities of type A_r , D_r , E_6 , E_7 and E_8

Let Γ be a finite subgroup of SU_2 . F. Klein showed in [9] that Γ is isomorphic to either the cyclic group C_n of order n , the binary dihedral group $\mathcal{D}_n$ of order $4n$, the binary tetrahedral group $\mathcal{T}$ of order 24, the binary octahedral group $\mathcal{O}$ of order 48, or the binary icosahedral group $\mathcal{I}$ of order 120, and that the quotient $\mathbb{C}^2/\Gamma$ is a surface which injects into $\mathbb{C}^3$ as the zero set of a polynomial $R_\Gamma \in \mathbb{C}[X, Y, Z]$. Furthermore this surface presents a unique isolated singularity and is called a *simple singularity*. P. Du Val then proved ([6]) that if $s \in \mathbb{C}^2/\Gamma$ is the singular point and $\pi : \tilde{S} \rightarrow \mathbb{C}^2/\Gamma$ is the minimal resolution of $\mathbb{C}^2/\Gamma$, then the preimage of s is a union of projective lines whose intersection matrix is the additive inverse of a Cartan matrix of type $\Delta(\Gamma) = A_r$, D_r or E_r .

Γ	R_Γ	Type of $\Delta(\Gamma)$
$\mathcal{C}_n$	$X^n + YZ$	A_{n-1}
$\mathcal{D}_n$	$X(Y^2 - X^n) + Z^2$	D_{n+2}
$\mathcal{T}$	$X^4 + Y^3 + Z^2$	E_6
$\mathcal{O}$	$X^3 + XY^3 + Z^2$	E_7
$\mathcal{I}$	$X^5 + Y^3 + Z^2$	E_8

Table 1: Equations and types of homogeneous simple singularities

1.1.2. Simple singularities of type B_r , C_r , F_4 and G_2

The definition of the simple singularities of inhomogeneous types is due to P. Slodowy (cf. [15] Section 6.2).

Definition 1.1. A simple singularity of type B_r ($r \geq 2$), C_r ($r \geq 3$), F_4 or G_2 is a pair (X_0, Ω) with X_0 a simple singularity (in the sense of Section 1.1.1) and Ω a group of automorphisms of X_0 according to the following list:

Type of (X_0, Ω)	Type of X_0	Γ	Γ'	Ω
B_r ($r \geq 2$)	A_{2r-1}	$\mathcal{C}_{2r}$	$\mathcal{D}_r$	$\mathbb{Z}/2\mathbb{Z}$
C_r ($r \geq 3$)	D_{r+1}	$\mathcal{D}_{r-1}$	$\mathcal{D}_{2(r-1)}$	$\mathbb{Z}/2\mathbb{Z}$
F_4	E_6	$\mathcal{T}$	$\mathcal{O}$	$\mathbb{Z}/2\mathbb{Z}$
G_2	D_4	$\mathcal{D}_2$	$\mathcal{O}$	$\mathfrak{S}_3$

Table 2: Definition of inhomogeneous simple singularities

In each row of this table, Γ and Γ' are finite subgroups of SU_2 such that X_0 is a simple singularity of type $\Delta(\Gamma)$ (i.e. $X_0 \cong \mathbb{C}^2/\Gamma$) and $\Gamma \triangleleft \Gamma'$. Furthermore there is a natural action of $\Omega = \Gamma'/\Gamma$ on the singularity X_0 . This action lifts in a unique way to the minimal resolution $\tilde{X}_0$ of X_0 . As Ω fixes the singular point of X_0 , it will stabilise the exceptional locus in $\tilde{X}_0$. Hence we obtain an action of Ω on the dual diagram $\Delta(\Gamma)$ of the exceptional divisors. It turns out that this action agrees with the automorphism group of the Dynkin diagram.

The inhomogeneous type associated to the simple singularity (X_0, Ω) will be referred as $\Delta(\Gamma, \Gamma')$.

A simple singularity of inhomogeneous type is then a simple homogeneous singularity with a symmetry of the Dynkin diagram. One can notice that the type of (X_0, Ω) is the same as the type of the folding by Ω of a root lattice of the same type as X_0 (cf. [4] Section 1.2 for details).

Remark 1.2. The type $(A_{2r}, \mathbb{Z}/2\mathbb{Z})$ is the only case that appears in Table 1 but not in Table 2. This is because the action of the symmetry group of the Dynkin diagram of type A_{2r} fails to lift to the exceptional locus of the minimal resolution of the associated simple singularity (cf. [4] Section 1.4.2.2).

The notion of symmetry has been added to the simple singularities, therefore it is necessary to include this symmetry in the definition of the deformations of singularities of type B_r , C_r , F_4 and G_2 ([15] Section 2.6):

Definition 1.3. A deformation of a simple singularity (X_0, Ω) is an Ω -equivariant deformation of the singularity X_0 with a trivial action of Ω on the base space.

Set $\pi : X \rightarrow Y$ a deformation of X_0 . A deformation $\psi : X' \rightarrow Y'$ of X_0 is said to be induced from π by a morphism $\varphi : Y' \rightarrow Y$ if there exists a morphism $\Phi : X' \rightarrow X$ such that the following diagram is Cartesian:

$$\begin{array}{ccc} X' & \xrightarrow{\Phi} & X \\ \psi \downarrow & & \downarrow \pi \\ Y' & \xrightarrow{\varphi} & Y \end{array}$$

and

$$\begin{array}{ccc} & X_0 & \\ \swarrow & & \searrow \\ X' & \xrightarrow{\Phi} & X \end{array}$$

commutes. The condition of the first diagram being Cartesian means that X' is isomorphic to the fiber product $X \times_Y Y'$. A semiuniversal deformation $\pi_0 : X \rightarrow Y$ of a simple singularity (X_0, Ω) is a deformation of (X_0, Ω) such that any other deformation $\psi : X' \rightarrow Y'$ of (X_0, Ω) is induced from π_0 by an Ω -equivariant morphism $\varphi : Y' \rightarrow Y$ with a uniquely determined differential $d_{Y'}\varphi : T_{Y'}Y' \rightarrow T_Y Y$.

Semiuniversal deformations of simple homogeneous and inhomogeneous singularities are unique up to isomorphism, which makes them ideal objects for the study of their associated singularities.

1.2. Deformations of homogeneous simple singularities

In this section we present a construction due to H. Cassens and P. Slodowy of the pullback of a semiuniversal deformation of a simple singularity of homogeneous type. For further details on proofs and references of the statements, we invite the reader to look at [5].

Let Γ be a finite subgroup of SU_2 , R its regular representation, N its natural representation as a subgroup of SU_2 , $\Delta(\Gamma)$ the associated Dynkin diagram (cf. Section 1.1.1), and define $M(\Gamma) = (\text{End}(R) \otimes N)^\Gamma$. Then $M(\Gamma)$ is the representation space of a *McKay quiver* Q , i.e. a quiver whose vertices are the vertices of the extended Dynkin diagram $\tilde{\Delta}(\Gamma)$, with two arrows (one in each direction) for any edge in $\tilde{\Delta}(\Gamma)$.

The group $G(\Gamma) = (\prod_{i=0}^r GL_{d_i}(\mathbb{C})) / \mathbb{C}^*$, with $(d_0, \dots, d_r)$ the dimension vector of $M(\Gamma)$, acts on $M(\Gamma)$ by simultaneous conjugation.

Using a formula due to G. Lusztig, one can define a non-degenerate $G(\Gamma)$ -invariant symplectic form $\langle \cdot, \cdot \rangle$ on $M(\Gamma)$ that induces a moment map

$$\mu_{CS} : M(\Gamma) \rightarrow (\text{Lie } G(\Gamma))^* \subset \bigoplus_{i=0}^r M_{d_i}(\mathbb{C}).$$

We identify $\text{Lie } G(\Gamma)$ and its dual $(\text{Lie } G(\Gamma))^*$.

Let Z be the dual of the center of $\text{Lie } G(\Gamma)$. Because the moment map is $G(\Gamma)$ -equivariant, $G(\Gamma)$ acts on any fiber $\mu_{CS}^{-1}(z)$ when $z \in Z$. Hence for any $z \in Z$, the quotient $\mu_{CS}^{-1}(z) // G(\Gamma)$ is well defined. It is then proved in [5] that the morphism

$$\mu_{CS}^{-1}(Z) // G(\Gamma) \longrightarrow Z$$

is the pullback of a semiuniversal deformation of the simple singularity $\mathbb{C}^2/\Gamma$ by the natural morphism $\mathfrak{h} \rightarrow \mathfrak{h}/W$, with $\mathfrak{h}$ a Cartan subalgebra of a Lie algebra of type $\Delta(\Gamma)$ and W the associated Weyl group.

1.3. Deformations of inhomogeneous simple singularities

This section is devoted to the extension conducted in [4] of the construction of Section 1.2 to the inhomogeneous simple singularities of type B_r ($r \geq 2$), C_r ($r \geq 3$), F_4 and G_2 .

Let $\Delta(\Gamma)$ be a Dynkin diagram of type A_{2r-1} ($r \geq 2$), D_r ($r \geq 4$) or E_6 , with Γ being the associated finite subgroup of SU_2 (cf. Section 1.1.1). Using the previous section, one obtains the following diagram:

$$\begin{array}{ccc} \mu_{CS}^{-1}(Z)/G(\Gamma) & = & X_\Gamma \times_{\mathfrak{h}/W} \mathfrak{h} \xrightarrow{\psi} X_\Gamma \\ \downarrow \tilde{\alpha} & \circlearrowleft & \downarrow \alpha \\ Z \cong \mathfrak{h} & \xrightarrow{\pi} & \mathfrak{h}/W \end{array}$$

with α a semiuniversal deformation of the simple singularity $\mathbb{C}^2/\Gamma$ of type $\Delta(\Gamma)$, $\mathfrak{h}$ a Cartan subalgebra of type $\Delta(\Gamma)$ and W the associated Weyl group.

Let Γ' be the finite subgroup of SU_2 such that there exists a simple singularity of inhomogeneous type $\Delta(\Gamma, \Gamma')$ (cf. Definition 1.1). Then $\Omega = \Gamma'/\Gamma$ acts on the singularity $X_{\Gamma,0} = \alpha^{-1}(0)$. Let us define $X_{\Gamma,\Omega} = \alpha^{-1}((\mathfrak{h}/W)^\Omega)$ and $\alpha^\Omega = \alpha|_{X_{\Gamma,\Omega}}$. Our aim is to define natural actions of Ω on X_Γ and $\mathfrak{h}/W$ such that α becomes Ω -equivariant, because if so, using results of P. Slodowy, one can show that the restriction $\alpha^\Omega : X_{\Gamma,\Omega} \rightarrow (\mathfrak{h}/W)^\Omega$ is a semiuniversal deformation of an inhomogeneous simple singularity of type $\Delta(\Gamma, \Gamma')$ (cf. [4] Section 4.3.1).

In [4] it is shown that if the action of Ω on $M(\Gamma)$ is symplectic, then $\tilde{\alpha}$ can be made into an Ω -equivariant map, and so is α . We then have the following result:

Theorem 1.4. *Let $M(\Gamma)$ be the representation space of a McKay quiver built on a Dynkin diagram of type A_{2r-1} , D_r or E_6 as explained in Section 1.2. Then there exists an action of $\Omega = \Gamma'/\Gamma$ on $M(\Gamma)$ that is both symplectic and induces the natural action on the singularity $\mathbb{C}^2/\Gamma$. This action then turns α into an Ω -equivariant morphism.*

The previous theorem enables us to explicitly compute semiuniversal deformations for the inhomogeneous types. For details on these computations, see [4] Sections 4.3 and 4.4.

2. Quotients of semiuniversal deformations of inhomogeneous simple singularities

In the previous section we have seen that the morphism $\alpha^\Omega : X_{\Gamma,\Omega} \rightarrow (\mathfrak{h}/W)^\Omega$ is Ω -invariant. Thus Ω acts on each fiber of α^Ω and the fibers can be quotiented. Furthermore it is known that $(\alpha^\Omega)^{-1}(\bar{0}) = X_{\Gamma,0} = \mathbb{C}^2/\Gamma$. Hence the quotient fiber is $(\alpha^\Omega)^{-1}(\bar{0})/\Omega = X_{\Gamma,0}/\Omega = (\mathbb{C}^2/\Gamma)/(\Gamma'/\Gamma) \cong \mathbb{C}^2/\Gamma'$. It is a simple singularity because Γ' is a finite subgroup of SU_2 . Therefore the family given by the quotient map $\overline{\alpha^\Omega} : X_{\Gamma,\Omega}/\Omega \rightarrow (\mathfrak{h}/W)^\Omega$ is a deformation of the simple singularity $\mathbb{C}^2/\Gamma'$ of type $\Delta(\Gamma')$. We will say that the deformation $\overline{\alpha^\Omega}$ is obtained through $\Delta(\Gamma) - \Delta(\Gamma, \Gamma') - \Delta(\Gamma')$ -procedure.

The results obtained in [4] Sections 4.3 and 4.4 lead to the following table:

Type of α^Ω	Type of $\overline{\alpha^\Omega}$	Rank of $\overline{\alpha^\Omega}$
B_r ($r \geq 2$)	D_{r+2}	r
C_r ($r \geq 3$)	D_{2r}	r
F_4	E_7	4
G_2	E_7	2

Table 3: Type and rank of $\overline{\alpha^\Omega}$

Here we slightly abuse notations by saying that a deformation is of type Δ if it is a deformation of a simple singularity of type Δ . Furthermore the *rank* of a deformation is the dimension of its base space.

Remark 2.1. In the case $\Delta(\Gamma, \Gamma') = G_2$, one can replace the symmetry group $\mathfrak{S}_3$ by $\mathbb{Z}/3\mathbb{Z}$. The rank of $\overline{\alpha^\Omega}$ remains 2, but the type of α^Ω becomes E_6 .

Examples 2.2. • For $\Delta(\Gamma, \Gamma') = B_2$, the map α^Ω is given by the projection

$$\begin{aligned} X_{\Gamma, \Omega} &= \{((x, y, z), (t_2, 0, t_4)) \in \mathbb{C}^3 \times \mathfrak{h}/W \mid z^4 + t_2 z^2 + t_4 + \frac{t_2^2}{8} = xy\} \\ &\downarrow \alpha^\Omega \\ (\mathfrak{h}/W)^\Omega &= \{(t_2, 0, t_4) \in \mathfrak{h}/W\} \end{aligned}$$

with the action of $\Omega = \mathbb{Z}/2\mathbb{Z} = \langle \sigma \rangle$ on a fiber $(\alpha^\Omega)^{-1}(t_2, 0, t_4)$ being $\sigma.(x, y, z) = (y, x, -z)$. Here t_i denotes the flat coordinate of degree i (cf. [13]). Therefore the quotient $\overline{\alpha^\Omega}$ is given by

$$\begin{aligned} X_{\Gamma, \Omega}/\Omega &= \{((X, Z, W), (t_2, 0, t_4)) \in \mathbb{C}^3 \times \mathfrak{h}/W \mid Z(X^2 - 4Z^2) + W^2 - 4t_2 Z^2 - 4(t_4 + \frac{t_2^2}{8})Z = 0\} \\ &\downarrow \overline{\alpha^\Omega} \\ (\mathfrak{h}/W)^\Omega &= \{(t_2, 0, t_4) \in \mathfrak{h}/W\} \end{aligned}$$

The map $\overline{\alpha^\Omega}$ is a deformation of a D_4 -singularity and is of rank 2.

• For $\Delta(\Gamma, \Gamma') = C_3$, the map α^Ω is given by the projection

$$\begin{aligned} X_{\Gamma, \Omega} &= \{((x, y, z), (t_2, t_4, t_6, 0)) \in \mathbb{C}^3 \times \mathfrak{h}/W \mid z^2 = xy(x + y) - \frac{t_2}{2}xy - \frac{t_4}{4}x + \frac{1}{4}(t_6 + \frac{t_2 t_4}{6} + \frac{t_2^3}{108})\} \\ &\downarrow \alpha^\Omega \\ (\mathfrak{h}/W)^\Omega &= \{(t_2, t_4, t_6, 0) \in \mathfrak{h}/W\} \end{aligned}$$

with the action of $\Omega = \mathbb{Z}/2\mathbb{Z} = \langle \sigma \rangle$ on a fiber $(\alpha^\Omega)^{-1}(t_2, t_4, t_6, 0)$ being $\sigma.(x, y, z) = (x, -x - y + \frac{t_2}{2}, -z)$. The coordinates on $\mathfrak{h}/W$ are once again the flat coordinates. Therefore the quotient $\overline{\alpha^\Omega}$ is given by

$$\begin{aligned} X_{\Gamma, \Omega}/\Omega &= \{((X, Y, W), (t_2, t_4, t_6, 0)) \in \mathbb{C}^3 \times \mathfrak{h}/W \mid -\frac{1}{64}X^5 + XY^2 - W^2 \\ &\quad + A_{X^4}X^4 + A_{X^3}X^3 + A_{X^2}X^2 + A_X X + A_Y Y + A_0 = 0\} \\ &\downarrow \overline{\alpha^\Omega} \\ (\mathfrak{h}/W)^\Omega &= \{(t_2, t_4, t_6, 0) \in \mathfrak{h}/W\} \end{aligned}$$

where the A_i are polynomials in t_2, t_4, t_6 without constant terms. So the map $\overline{\alpha^\Omega}$ is indeed a deformation of a D_6 -singularity and is of rank 3.

For details on the computations, see [4] Section 4.4.

E. Brieskorn showed ([3]) that a semiuniversal deformation of a simple singularity of type Δ_r ($\Delta = A, D$ or E) can be constructed such that the base space is the quotient $\mathfrak{h}/W$ with $\mathfrak{h}$ a Cartan subalgebra of a simple Lie algebra of type Δ_r and W the associated Weyl group. Hence the rank of a semiuniversal deformation of a simple singularity of type Δ_r is r . However one notices in Table 3 that it is not true for $\overline{\alpha^\Omega}$. Therefore $\overline{\alpha^\Omega}$ is not a semiuniversal deformation in any case.

We will now prove the following theorem regarding the regularity of the fibers of $\overline{\alpha^\Omega}$:

Theorem 2.3. *Let $\alpha^\Omega : X_{\Gamma, \Omega} \rightarrow (\mathfrak{h}/W)^\Omega$ be the semiuniversal deformation obtained in Section 1.3 of a simple singularity of inhomogeneous type B_r ($r \geq 2$), C_r ($r \geq 3$), F_4 or G_2 . Then every fiber of the quotient $\overline{\alpha^\Omega} : X_{\Gamma, \Omega}/\Omega \rightarrow (\mathfrak{h}/W)^\Omega$ is singular.*

Proof. In the following computations, the t_i 's will always be the flat coordinates on $\mathfrak{h}/W$ (cf. [13]), with i being the degree. For F_4 and G_2 , we will look at the explicit equation obtained in [4] Section 4.4 and see that each fiber of $\overline{\alpha^\Omega}$ is indeed singular. For the general types B_r and C_r , another approach is needed. We will show that each fiber of α^Ω has at least one isolated fixed point p . If a fiber has such a point p , then there are two possibilities:

- p is singular, so a neighbourhood of p is isomorphic to $\mathbb{C}^2/\Xi$ with Ξ a finite subgroup of SU_2 , because p is a singular point in a semiuniversal deformation of a simple singularity. Thus we have $\dim T_p(\mathbb{C}^2/\Xi) > \dim \mathbb{C}^2/\Xi$. Furthermore $T_{\pi(p)}((\mathbb{C}^2/\Xi)/\Omega) \cong T_p(\mathbb{C}^2/\Xi)/T_p(\Omega.p)$ with π the quotient map. As p is a fixed point, we obtain $\dim T_{\pi(p)}((\mathbb{C}^2/\Xi)/\Omega) = \dim T_p(\mathbb{C}^2/\Xi) > \dim \mathbb{C}^2/\Xi = \dim (\mathbb{C}^2/\Xi)/\Omega$ because Ω is a finite group. Hence $\dim T_{\pi(p)}((\mathbb{C}^2/\Xi)/\Omega) > \dim (\mathbb{C}^2/\Xi)/\Omega$, and $\pi(p)$ is a singular point of $(\mathbb{C}^2/\Xi)/\Omega$.

- p is smooth, then the fiber has a tangent space at this point and it is isomorphic to $\mathbb{C}^2$. So locally around p , the action of $\Omega = \mathbb{Z}/2\mathbb{Z}$ is the same as its action on $\mathbb{C}^2$ around the origin. Hence the quotient will be isomorphic to $\mathbb{C}^2/\Omega$, which is an A_1 -singularity. The quotient will then be singular.

- Type B_r ($r \geq 2$). The fiber of α^Ω above a point $(t_2, 0, t_4, 0, \dots, t_{2r}) \in (\mathfrak{h}/W)^\Omega$ is given by the points $(x, y, z) \in \mathbb{C}^3$ such that $z^{2r} + \sum_{i=1}^r f_{2i}(t_2, \dots, t_{2r}) z^{2(r-i)} = xy$ (cf. [4] Section 4.3.6) where, for all $1 \leq i \leq 2r$, f_i is a polynomial such that $f_i(t_2(\xi), \dots, t_{2r}(\xi)) = \epsilon_i(\xi)$, ϵ_i being the elementary symmetric polynomial of degree i and ξ being a coordinate vector on the Cartan subalgebra $\mathfrak{h}$ of type A_{2r-1} . The action of $\Omega = \mathbb{Z}/2\mathbb{Z} = \langle \sigma \rangle$ on the fiber is given by

$$\sigma.(x, y, z) = ((-1)^r y, (-1)^r x, -z).$$

Hence a fixed point is a point $p = (x, (-1)^r x, 0)$ such that $x^2 = (-1)^r f_2(t_2, \dots, t_{2r})$. As $f_2(t_2, \dots, t_{2r}) = t_2$, we define $x = (-1)^{\frac{r}{2}} \sqrt{t_2}$. Then $p = (x, (-1)^r x, 0)$ is a fixed point of the fiber. It is isolated because there are at most two possible values for x .

- Type C_r ($r \geq 3$). Based on results from A. Kas and M. Schlessinger ([8]), as well as computations from [4], the fiber of the restriction α^Ω of the semiuniversal deformation α of a singularity of type D_{r+1} above a fixed point $(t_2, t_4, \dots, t_{2r}, 0) \in (\mathfrak{h}/W)^\Omega$ has the following equation:

$$x^r + xy^2 + z^2 + \alpha_{r-1}x^{r-1} + \dots + \alpha_1x + \beta y + \alpha_0 = 0,$$

with β and the α_i 's being complex polynomials in variables $t_2, t_4, \dots, t_{2r}$. Here the system of flat coordinates on $\mathfrak{h}/W$ is given by $(t_2, t_4, \dots, t_{2r}, \psi)$, Ω fixes each t_i and sends ψ on $-\psi$. The preceding equation is a quasi-homogeneous polynomial with

$$\begin{cases} x \text{ of degree } 2, \\ y \text{ of degree } r-1, \\ z \text{ of degree } r, \\ t_i \text{ of degree } i, i = 2, 4, \dots, 2r. \end{cases}$$

From the study of the special fiber $(\alpha^\Omega)^{-1}(\bar{0})$ and the fact that the action of $\Omega = \mathbb{Z}/2\mathbb{Z} = \langle \sigma \rangle$ preserves the degree, one can see that

$$\begin{cases} \sigma.x = x + p(x, y, z, t_2, \dots, t_{2r}), \text{ of degree } 2, \\ \sigma.y = -y + q(x, y, z, t_2, \dots, t_{2r}), \text{ of degree } r-1, \\ \sigma.z = -z + s(x, y, z, t_2, \dots, t_{2r}), \text{ of degree } r, \\ \sigma.t_i = t_i, i = 2, 4, \dots, 2r, \end{cases}$$

where p, q and s are homogeneous polynomials with respect to the degrees of x, y, z and the t_i 's, and such that $p(x, y, z, 0, \dots, 0) = q(x, y, z, 0, \dots, 0) = s(x, y, z, 0, \dots, 0) = 0$ for any $(x, y, z) \in \mathbb{C}^3$.

Let us fix $(t_2, \dots, t_{2r}, 0) \in (\mathfrak{h}/W)^\Omega$ and study the fiber $(\alpha^\Omega)^{-1}(t_2, \dots, t_{2r}, 0)$ given by the equation given above.

Because x is of smallest possible degree and $\sigma^2 = \text{Id}$, the only possibility is $\sigma.x = x$. For the other variables, the action might depend on the parity of r .

■ Assume that r is even. y is of degree $r - 1$ which is odd, and all the others terms are of even degree. The only possibility is thus $\sigma.y = -y$. β is a polynomial of degree $2r - \deg y = r + 1$, which is odd. However β is a polynomial in the variables $t_2, \dots, t_{2r}$, all of even degree. Hence $\beta = 0$.

z is of degree r which is even, so $\sigma.z = -z + \sum_{i=0}^{\frac{r}{2}} \lambda_i x^i$ with the λ_i 's being polynomials in $t_2, \dots, t_{2r}$. One can check that $\sigma^2 z = z$.

We are then looking for the fixed points of the action

$$\begin{cases} \sigma.x = x, \\ \sigma.y = -y, \\ \sigma.z = -z + \sum_{i=0}^{\frac{r}{2}} \lambda_i x^i. \end{cases}$$

If $(x, y, z) \in (\alpha^\Omega)^{-1}(t_2, \dots, t_{2r}, 0)$, then $\sigma.(x, y, z) = (x, -y, -z + \sum_{i=0}^{\frac{r}{2}} \lambda_i x^i) \in (\alpha^\Omega)^{-1}(t_2, \dots, t_{2r}, 0)$. Therefore the following equation has to be verified:

$$\begin{aligned} & x^r + xy^2 + (-z + \sum_{i=0}^{\frac{r}{2}} \lambda_i x^i)^2 + \alpha_{r-1} x^{r-1} + \dots + \alpha_1 x + \alpha_0 = 0, \\ \Leftrightarrow & x^r + xy^2 + z^2 + \alpha_{r-1} x^{r-1} + \dots + \alpha_1 x + \alpha_0 + (\sum_{i=0}^{\frac{r}{2}} \lambda_i x^i)(-2z + \sum_{i=0}^{\frac{r}{2}} \lambda_i x^i) = 0, \\ \Leftrightarrow & (\sum_{i=0}^{\frac{r}{2}} \lambda_i x^i)(-2z + \sum_{i=0}^{\frac{r}{2}} \lambda_i x^i) = 0 \end{aligned}$$

because $(x, y, z) \in (\alpha^\Omega)^{-1}(t_2, \dots, t_{2r}, 0)$.

If $\lambda_i = 0$ for all $i \in \{0, \dots, \frac{r}{2}\}$, then the action is $\begin{cases} \sigma.x = x, \\ \sigma.y = -y, \\ \sigma.z = -z. \end{cases}$

If there exists $i \in \{0, \dots, \frac{r}{2}\}$ such that $\lambda_i \neq 0$, then the equation $\sum_{i=0}^{\frac{r}{2}} \lambda_i x^i = 0$ is non-trivial in x and has therefore at least one solution (and at most $\frac{r}{2}$). Given that in $(\alpha^\Omega)^{-1}(t_2, \dots, t_{2r}, 0)$ the variable x is not constant, there exists x such that $\sum_{i=0}^{\frac{r}{2}} \lambda_i x^i \neq 0$ and such that there exist $y, z \in \mathbb{C}$ with $(x, y, z) \in (\alpha^\Omega)^{-1}(t_2, \dots, t_{2r}, 0)$. Hence we get that $z = \frac{1}{2} \sum_{i=0}^{\frac{r}{2}} \lambda_i x^i \neq 0$. So for such an x , the value of z is uniquely determined and is not zero. However, based on the equation of $(\alpha^\Omega)^{-1}(t_2, \dots, t_{2r}, 0)$, we see that if (x, y, z) is a point on the surface, then so is $(x, y, -z)$. If a non-zero value of z exists, a second value exists also. This contradicts the unicity of the value of z .

We have shown that the action of Ω on $(\alpha^\Omega)^{-1}(t_2, \dots, t_{2r}, 0)$ is given by

$$\begin{cases} \sigma.x = x, \\ \sigma.y = -y, \\ \sigma.z = -z. \end{cases}$$

A fixed point is thus given by $(x, 0, 0)$ with x verifying the non-trivial equation $x^r + \alpha_{r-1}x^{r-1} + \dots + \alpha_1x + \alpha_0 = 0$. So a fixed point exists, there are at most r of them, and they are all isolated.

■ Assume that r is odd. z is of degree r which is odd, and all the others terms are of even degree. The only possibility is thus $\sigma.z = -z$.

β is a polynomial of degree $2r - \deg y = r + 1$, which is even. So it is possible that $\beta \neq 0$.

y is of degree $r - 1$ which is even, so $\sigma.y = -y + \sum_{i=0}^{\frac{r-1}{2}} \mu_i x^i$ with the μ_i 's being polynomials in $t_2, \dots, t_{2r}$. One can check that $\sigma^2 y = y$.

We are then looking for the fixed points of the action

$$\begin{cases} \sigma.x = x, \\ \sigma.y = -y + \sum_{i=0}^{\frac{r-1}{2}} \mu_i x^i, \\ \sigma.z = -z. \end{cases}$$

If $(x, y, z) \in (\alpha^\Omega)^{-1}(t_2, \dots, t_{2r}, 0)$, then $\sigma.(x, y, z) = (x, -y + \sum_{i=0}^{\frac{r-1}{2}} \mu_i x^i, -z) \in (\alpha^\Omega)^{-1}(t_2, \dots, t_{2r}, 0)$. Therefore the following equation has to be verified:

$$\begin{aligned} & x^r + x(-y + \sum_{i=0}^{\frac{r-1}{2}} \mu_i x^i)^2 + z^2 + \alpha_{r-1}x^{r-1} + \dots + \alpha_1x + \beta(-y + \sum_{i=0}^{\frac{r-1}{2}} \mu_i x^i) + \alpha_0 = 0, \\ \Leftrightarrow & x^r + xy^2 + z^2 + \alpha_{r-1}x^{r-1} + \dots + \alpha_1x + \beta y + \alpha_0 + (\sum_{i=0}^{\frac{r-1}{2}} \mu_i x^i)(\beta - 2xy + x \sum_{i=0}^{\frac{r-1}{2}} \mu_i x^i) = 2\beta y, \\ \Leftrightarrow & (\sum_{i=0}^{\frac{r-1}{2}} \mu_i x^i)(\beta - 2xy + x \sum_{i=0}^{\frac{r-1}{2}} \mu_i x^i) = 2\beta y \end{aligned}$$

because $(x, y, z) \in (\alpha^\Omega)^{-1}(t_2, \dots, t_{2r}, 0)$.

If $\mu_i = 0$ for all $i \in \{0, \dots, \frac{r-1}{2}\}$, then $2\beta y = 0$ for all y such that $(x, y, z) \in (\alpha^\Omega)^{-1}(t_2, \dots, t_{2r}, 0)$. But there exists at least one point in the surface for which $y \neq 0$. Hence $\beta = 0$, and the action is given by

$$\begin{cases} \sigma.x = x, \\ \sigma.y = -y, \\ \sigma.z = -z. \end{cases}$$

If there exists $i \in \{0, \dots, \frac{r-1}{2}\}$ such that $\mu_i \neq 0$, we have to consider two cases: $\beta = 0$ and $\beta \neq 0$.

▲ Assume that $\beta = 0$. Then $x(\sum_{i=0}^{\frac{r-1}{2}} \mu_i x^i)(-2y + \sum_{i=0}^{\frac{r-1}{2}} \mu_i x^i) = 0$. The equation $x(\sum_{i=0}^{\frac{r-1}{2}} \mu_i x^i) = 0$ is non-trivial in x and has therefore at least one solution (and at most $\frac{r+1}{2}$). Given that in $(\alpha^\Omega)^{-1}(t_2, \dots, t_{2r}, 0)$ the variable x is not constant, there exists x such that $x(\sum_{i=0}^{\frac{r-1}{2}} \mu_i x^i) \neq 0$ and such that there exist $y, z \in \mathbb{C}$ with $(x, y, z) \in (\alpha^\Omega)^{-1}(t_2, \dots, t_{2r}, 0)$. Hence we get that $y = \frac{1}{2} \sum_{i=0}^{\frac{r-1}{2}} \mu_i x^i \neq 0$. So for such an x , the value of y is uniquely determined and is not zero. However, based on the equation of $(\alpha^\Omega)^{-1}(t_2, \dots, t_{2r}, 0)$, we see that if (x, y, z) is a point on the surface, then so is $(x, -y, z)$ (because $\beta = 0$). If a non-zero value of y exists, a second value exists also. This contradicts the unicity of the value of y .

▲ Assume that $\beta \neq 0$. We have

$$\begin{aligned} & (\sum_{i=0}^{\frac{r-1}{2}} \mu_i x^i)(\beta - 2xy + x \sum_{i=0}^{\frac{r-1}{2}} \mu_i x^i) = 2\beta y, \\ \Leftrightarrow & (\beta + x \sum_{i=0}^{\frac{r-1}{2}} \mu_i x^i)(-2y + \sum_{i=0}^{\frac{r-1}{2}} \mu_i x^i) = 0. \end{aligned}$$

As $\beta \neq 0$, for any $x \in \mathbb{C}$, there exist an infinity of $(y, z) \in \mathbb{C}^2$ such that $(x, y, z) \in (\alpha^\Omega)^{-1}(t_2, \dots, t_{2r}, 0)$. Set $x \in \mathbb{C}$ such that $x(\beta + x \sum_{i=0}^{\frac{r-1}{2}} \mu_i x^i) \neq 0$ (we get rid of a finite number of possible values). Set $y \in \mathbb{C}$ such that $2xy + \beta \neq 0$ (we get rid of one possible value). Because of the equation of the surface, there exists $z \in \mathbb{C}$ such that $(x, y, z) \in (\alpha^\Omega)^{-1}(t_2, \dots, t_{2r}, 0)$. One can then check that the point $(x, y - \frac{\beta + 2xy}{x}, z)$ is well defined and belongs to the surface. But because of the choice of x , we know that $y = \frac{1}{2} \sum_{i=0}^{\frac{r-1}{2}} \mu_i x^i$. So for such an x , the value of y is uniquely determined. However, we have just seen that (x, y, z) and $(x, y - \frac{\beta + 2xy}{x}, z)$ are two distinct points of the surface. This contradicts the unicity of the value of y .

We have shown that the action of Ω on $(\alpha^\Omega)^{-1}(t_2, \dots, t_{2r}, 0)$ is given by

$$\begin{cases} \sigma.x = x, \\ \sigma.y = -y, \\ \sigma.z = -z. \end{cases}$$

A fixed point is thus given by $(x, 0, 0)$ with x verifying the non-trivial equation $x^r + \alpha_{r-1}x^{r-1} + \dots + \alpha_1x + \alpha_0 = 0$. Therefore a fixed point exists, there are at most r of them, and they are all isolated.

We have thus proved that, for any $r \geq 3$ and any $(t_2, \dots, t_{2r}, 0) \in (\mathfrak{h}/W)^\Omega$, the equation of $(\alpha^\Omega)^{-1}(t_2, \dots, t_{2r}, 0)$ is

$$x^r + xy^2 + z^2 + \alpha_{r-1}x^{r-1} + \dots + \alpha_1x + \alpha_0 = 0,$$

the action of Ω is given by

$$\begin{cases} \sigma.x = x, \\ \sigma.y = -y, \\ \sigma.z = -z, \end{cases}$$

and there is at least one isolated fixed point.

• Type F_4 . The equation of the fiber of α^Ω above a point $(t_2, 0, t_6, t_8, 0, t_{12}) \in (\mathfrak{h}/W)^\Omega$ is given by the points $(x, y, z) \in \mathbb{C}^3$ such that

$$-\frac{1}{4}x^4 + y^3 + z^2 - \frac{t_2}{4}x^2y + \frac{1}{48}(t_6 - \frac{t_2^3}{8})x^2 + \frac{1}{48}(-t_8 + \frac{t_6t_2}{4} - \frac{t_2^4}{192})y + \frac{1}{576}(t_{12} - \frac{t_8t_2^2}{8} - \frac{t_6^2}{8} + \frac{t_6t_2^3}{96}) = 0.$$

The action of $\Omega = \mathbb{Z}/2\mathbb{Z} = \langle \sigma \rangle$ on the fiber is then given by

$$\sigma.(x, y, z) = (-x, y, -z).$$

We can compute the equation of the quotient of the fiber and obtain

$$-\frac{1}{4}X^3 + XY^3 + Z^2 - \frac{t_2}{4}X^2Y + \frac{1}{48}(t_6 - \frac{t_2^3}{8})X^2 + \frac{1}{48}(-t_8 + \frac{t_6t_2}{4} - \frac{t_2^4}{192})XY + \frac{1}{576}(t_{12} - \frac{t_8t_2^2}{8} - \frac{t_6^2}{8} + \frac{t_6t_2^3}{96})X = 0.$$

One can check that for any $(t_2, 0, t_6, t_8, 0, t_{12}) \in (\mathfrak{h}/W)^\Omega$, the surface just obtained is indeed singular.

• Type $G_2 = (D_4, \mathbb{Z}/3\mathbb{Z})$. The equation of the fiber of α^Ω above a point $(t_2, 0, t_6, 0) \in (\mathfrak{h}/W)^\Omega$ is given by the points $(x, y, z) \in \mathbb{C}^3$ such that

$$z^2 = xy(x + y) - \frac{t_2}{2}xy + \frac{1}{4}(t_6 + \frac{t_2^3}{108}).$$

The action of $\Omega = \mathbb{Z}/3\mathbb{Z} = \langle \rho \rangle$ on the fiber is thus given by

$$\rho.(x, y, z) = (y, -x - y + \frac{t_2}{2}, z).$$

We can compute the equation of the quotient of the fiber and obtain

$$11664X^4 - Y^3 - Z^2 - 324t_2X^2Y - (189t_2^3 + 5832t_6)X^2 + (81t_2t_6 + \frac{15}{16}t_2^4)Y + \frac{11}{32}t_2^6 + \frac{189}{4}t_2^3t_6 + 729t_6^2 = 0.$$

One can easily check that for any $(t_2, 0, t_6, 0) \in (\mathfrak{h}/W)^\Omega$, the surface defined by the previous equation has at least one singular point.

• Type $G_2 = (D_4, \mathfrak{S}_3)$. The fixed point of $\Omega = \mathfrak{S}_3 = \langle \rho, \sigma \rangle$ in $\mathfrak{h}/W$ are the same as the ones by the action of $\mathbb{Z}/3\mathbb{Z}$: $\{(t_2, 0, t_6, 0) \in \mathfrak{h}/W\}$. The equation of the fiber of α^Ω above a fixed point in $\mathfrak{h}/W$ is then the same as in the case of $(D_4, \mathbb{Z}/3\mathbb{Z})$. The action of Ω on such a fiber is given by

$$\rho.(x, y, z) = (y, -x - y + \frac{t_2}{2}, z) \text{ and } \sigma.(x, y, z) = (x, -x - y + \frac{t_2}{2}, -z).$$

The quotient of the fiber has then the equation

$$X^3Y - 11664Y^3 + Z^2 + 324t_2XY^2 + (189t_2^3 + 5832t_6)Y^2 - (\frac{15}{16}t_2^4 + 81t_2t_6)XY - (\frac{11}{32}t_2^6 + \frac{189}{4}t_2^3t_6 + 729t_6^2)Y = 0.$$

Like before, for any $(t_2, 0, t_6, 0) \in (\mathfrak{h}/W)^\Omega$, the surface has at least one singular point. $\square$

3. Singular configurations and root systems

Let $\mathfrak{g}$ be a simple Lie algebra of type A_r , D_r or E_r , $\mathfrak{h}$ a Cartan subalgebra of $\mathfrak{g}$, W the associated Weyl group, and $e \in \mathfrak{g}$ a nilpotent element. As e is nilpotent, it can be included in an $\mathfrak{sl}_2$ -triple (e, f, h) by Jacobson-Morozov's theorem. Let us look at the restriction of the adjoint quotient $\chi : \mathfrak{g} \rightarrow \mathfrak{h}/W$ to the Slodowy slice $S_e = e + \mathfrak{z}_{\mathfrak{g}}(f)$. Consider the diagram

$$\begin{array}{ccc} & & S_e \\ & & \downarrow \chi|_{S_e} \\ \mathfrak{h} & \xrightarrow{\pi} & \mathfrak{h}/W \\ \bigcup_{\alpha \in \Phi^+} H_\alpha & \longmapsto & \mathbb{D} \end{array}$$

with π the natural projection, Φ^+ the set of positive roots of $\mathfrak{g}$, the H_α 's the reflection hyperplanes with respect to the roots $\alpha \in \Phi^+$, and $\mathbb{D}$ the discriminant of χ . E. Brieskorn proved in [3] that if e is a subregular (i.e. $\dim Z_G(e) = \text{rank } \mathfrak{g} + 2$, with G the simple Lie group associated to $\mathfrak{g}$) nilpotent element of $\mathfrak{g}$, then $\chi|_{S_e}$ is a semiuniversal deformation of a simple singularity of the same type as $\mathfrak{g}$, and is surjective ([15] Section 7.4, Corollary 1). Furthermore, it is shown in [15] (Section 6.5, Lemmas 1, 2, 3) that the type of the singular configuration which appears in S_e above a point $\pi(h) \in \mathbb{D}$ is given by the sub-root system $\{\alpha \in \Phi \mid h \in H_\alpha\}$.

Let $\mathfrak{g}'$ be a simple Lie algebra of type $\Delta(\Gamma')$ with root system Φ' , $\mathfrak{h}' \subset \mathfrak{g}'$ a Cartan subalgebra and W' the associated Weyl group. Set $e' \in \mathfrak{g}'$ a subregular nilpotent element and $S_{e'}$ the associated Slodowy slice. Hence the restriction $\chi'_{|S_{e'}} : S_{e'} \rightarrow \mathfrak{h}'/W'$ of the adjoint quotient χ' of $\mathfrak{g}'$ to $S_{e'}$ is a semiuniversal deformation of a simple singularity of type $\Delta(\Gamma')$. We have seen before that $\overline{\alpha^\Omega} : X_{\Gamma, \Omega} \rightarrow (\mathfrak{h}/W)^\Omega$ is a non-semiuniversal deformation of a simple singularity X_0 of type $\Delta(\Gamma')$. Therefore $\overline{\alpha^\Omega}$ is isomorphic to the pullback χ'_f of $\chi'_{|S_{e'}}$ by a base change $f : (\mathfrak{h}/W)^\Omega \rightarrow \mathfrak{h}'/W'$ whose differential at 0 is uniquely determined (cf. [14] Section 1.4). We thus have the following diagram:

$$\begin{array}{ccccccc}
& & g' & & (X_0, 0) & & \\
& & \cup & & \swarrow & & \searrow \\
(S_{e'}, e') & \xleftarrow{\varphi} & (S_{e'}, e') \times_{(\mathfrak{h}'/W', \bar{0})} ((\mathfrak{h}/W)^\Omega, \bar{0}) & \xleftarrow{\cong} & X_{\Gamma, \Omega}/\Omega & & \\
\downarrow \chi'_{|S_{e'}} & & \downarrow \chi'_f & & \downarrow \overline{\alpha^\Omega} & & \\
\mathfrak{h}' & \xrightarrow{\pi'} & (\mathfrak{h}'/W', \bar{0}) & \xleftarrow{f} & ((\mathfrak{h}/W)^\Omega, \bar{0}) & \xleftarrow{=} & ((\mathfrak{h}/W)^\Omega, \bar{0}) \\
\cup & & & & & & \\
\bigcup_{\alpha \in \Phi'^+} H_\alpha & \longrightarrow & \mathbb{D}' & & & &
\end{array}$$

Set $t \in (\mathfrak{h}/W)^\Omega$. Through the isomorphism, the fiber $(\overline{\alpha^\Omega})^{-1}(t)$ is the space

$$\{(\lambda, t) \in S_{e'} \times \{t\} \mid \chi'_{|S_{e'}}(\lambda) = f(t)\} \cong (\chi'_{|S_{e'}})^{-1}(f(t)).$$

Therefore the type of the singular configuration of $(\overline{\alpha^\Omega})^{-1}(t)$ is the same as the one of $(\chi'_{|S_{e'}})^{-1}(f(t))$, which is given by the sub-root system $\phi' = \{\alpha \in \Phi'^+ \mid \alpha(h) = 0, \text{ with } h \in \mathfrak{h}' \text{ a representative of } f(t)\}$.

The following conjecture describes the link between the singular configurations of the fibers of $\overline{\alpha^\Omega}$ and sub-root systems of the root system Φ' of type $\Delta(\Gamma')$:

Conjecture 3.1. *Let $\overline{\alpha^\Omega}$ be a deformation of a simple singularity of type $\Delta(\Gamma')$ obtained through $\Delta(\Gamma) - \Delta(\Gamma, \Gamma') - \Delta(\Gamma')$ -procedure. With the notations of the previous commutative diagram, there exists a subset Θ of the set simple roots of Φ' such that the base change f is an isomorphism on its image $f(\mathfrak{h}/W) = \pi'(\cap_{\alpha \in \Theta} H_\alpha)$, and such that if $h' \in \cap_{\alpha \in \phi'} H_\alpha \subset \mathfrak{h}'$ with $\Theta \subset \phi'$ and ϕ' maximal for h' (i.e. $\alpha(h') \neq 0$ if $\alpha \notin \phi'$), then the singular configuration of the fiber $(\overline{\alpha^\Omega})^{-1}(f^{-1}(\pi'(h')))$ is of the same type as the root system ϕ' . Furthermore, all singular configurations of the fibers of $\overline{\alpha^\Omega}$ are obtained in this way.*

Depending on the type of $\overline{\alpha^\Omega}$, the subset Θ is given in the following table (the numbering of the root systems is the one of Bourbaki (cf. [2])):

$\Delta(\Gamma) - \Delta(\Gamma, \Gamma') - \Delta(\Gamma')$	Θ	Θ in $\Delta(\Gamma')$
$A_{2r-1} - B_r - D_{r+2}$	$\{\alpha_{r+1}, \alpha_{r+2}\}$	
$D_{r+1} - C_r - D_{2r}$	$\{\alpha_{2i+1}\}_{0 \leq i \leq r-1}$	
$E_6 - F_4 - E_7$	$\{\alpha_2, \alpha_5, \alpha_7\}$	

$D_4 - G_2 - E_6$	$\{\alpha_1, \alpha_3, \alpha_5, \alpha_6\}$	
$D_4 - G_2 - E_7$	$\{\alpha_1, \alpha_2, \alpha_3, \alpha_5, \alpha_7\}$	

Table 4: Description of Θ for the different cases

The next statement validates the conjecture for small rank cases.

Theorem 3.2. *The Conjecture 3.1 is true for the types:*

- $A_3 - B_2 - D_4$,
- $A_5 - B_3 - D_5$,
- $D_4 - C_3 - D_6$,
- $D_4 - G_2 - E_6$,
- $D_4 - G_2 - E_7$,
- $E_6 - F_4 - E_7$.

4. Proof of Theorem 3.2

In what follows, the set of coordinates on $\mathfrak{h}/W$ and $\mathfrak{h}'/W'$ will always be the flat coordinates (cf. [11] and [12] for details).

4.1. Strategy of the proof

The strategy for the proof of each case is twofold:

- Part 1:
 - Compute the deformation α^Ω and its discriminant.
 - Decompose the discriminant depending on what singular configurations appear in the fibers of α^Ω .
 - Compute the deformation $\overline{\alpha^\Omega}$.
 - Determine the types of singular configurations appearing in the fibers of $\overline{\alpha^\Omega}$ above the previously obtained decomposition of $(\mathfrak{h}/W)^\Omega$.
- Part 2:
 - Compute all realisations of the sub-root systems of the root system of type $\Delta(\Gamma')$ containing Θ .
 - Compute the flat coordinates of the Cartan subalgebra $\mathfrak{h}'$ of type $\Delta(\Gamma')$, and the restriction of the map $\pi' : \mathfrak{h}' \rightarrow \mathfrak{h}'/W'$ to the subspace $\bigcap_{\alpha \in \Theta} H_\alpha$, with $H_\alpha = \{h \in \mathfrak{h}' \mid \alpha(h) = 0\}$.
 - Construct the base change $f : (\mathfrak{h}/W)^\Omega \longrightarrow \mathfrak{h}'/W'$.
 - Verify that $f : (\mathfrak{h}/W)^\Omega \xrightarrow{\cong} \pi'(\bigcap_{\alpha \in \Theta} H_\alpha)$.
 - Check that the morphism f realises a bijection between the singular configurations of the fibers of $\overline{\alpha^\Omega}$ and the sub-root systems of $\Delta(\Gamma')$ containing Θ .

4.2. Case $A_3 - B_2 - D_4$

In this section $\Gamma = \mathcal{C}_4$, $\Gamma' = \mathcal{D}_2$ and $\Omega = \mathbb{Z}/2\mathbb{Z} = \langle \sigma \rangle$. It is known from [4] that the map α^Ω is given by the projection

$$\begin{aligned} X_{\Gamma, \Omega} &= \{((x, y, z), (t_2, 0, t_4)) \in \mathbb{C}^3 \times \mathfrak{h}/W \mid z^4 + t_2 z^2 + t_4 + \frac{t_2^2}{8} = xy\} \\ &\downarrow \alpha^\Omega \\ (\mathfrak{h}/W)^\Omega &= \{(t_2, 0, t_4) \in \mathfrak{h}/W\} \end{aligned}$$

Furthermore, the action of Ω on a fiber is $\sigma.(x, y, z) = (y, x, -z)$, and so the quotient $\overline{\alpha^\Omega}$ is given by

$$\begin{aligned} X_{\Gamma, \Omega}/\Omega &= \{((X, Z, W), (t_2, 0, t_4)) \in \mathbb{C}^3 \times \mathfrak{h}/W \mid Z(X^2 - 4Z^2) + W^2 - 4t_2 Z^2 - 4(t_4 + \frac{t_2^2}{8})Z = 0\} \\ &\downarrow \overline{\alpha^\Omega} \\ (\mathfrak{h}/W)^\Omega &= \{(t_2, 0, t_4) \in \mathfrak{h}/W\} \end{aligned}$$

Before quotient, the discriminant of α^Ω is

$$\{(t_2, 0, t_4) \in \mathfrak{h}/W \mid (t_2^2 + 8t_4)(t_2^2 - 8t_4) = 0\}.$$

After quotient, it is known from Theorem 2.3 that all of $(\mathfrak{h}/W)^\Omega$ composes the discriminant. By studying explicitly the singularities in the fibers $(\alpha^\Omega)^{-1}(t_2, 0, t_4)$ and $(\overline{\alpha^\Omega})^{-1}(t_2, 0, t_4)$, we obtain the following types:

$(t_2, 0, t_4)$	Configuration of $(\alpha^\Omega)^{-1}(t_2, 0, t_4)$	Configuration of $(\overline{\alpha^\Omega})^{-1}(t_2, 0, t_4)$
Generic point	$p_1 + p_2$	$A_1 + A_1$
$t_4 = -\frac{t_2^2}{8} \neq 0$	A_1	A_3
$t_4 = \frac{t_2^2}{8} \neq 0$	$p_1 \quad + \quad p_2 \quad + \quad A_1 \quad + \quad A_1$	$A_1 + A_1 + A_1$
$t_2 = t_4 = 0$	A_3	D_4

Table 5: Configurations of the fibers of α^Ω and $\overline{\alpha^\Omega}$ for $\Delta(\Gamma) = A_3$

with p_i ($i = 1, 2$) denoting a smooth point, and if the action of Ω is not mentioned, the point (smooth or singular) is Ω -fixed.

Let $\Pi' = \{\alpha_1, \alpha_2, \alpha_3, \alpha_4\}$ be a set of simple roots of the root system of type D_4 (cf. Table 4 for the numbering). All the sub-root systems of D_4 containing $\Theta = \{\alpha_3, \alpha_4\}$ are either of type $A_1 + A_1$, $A_1 + A_1 + A_1$, A_3 or D_4 , and their realisations ϕ' are given in the following table:

Type of ϕ'	Realisations of ϕ'
$A_1 + A_1$	$\langle \alpha_3, \alpha_4 \rangle$
$A_1 + A_1 + A_1$	$\langle \alpha_1, \alpha_3, \alpha_4 \rangle, \langle \alpha_1 + 2\alpha_2 + \alpha_3 + \alpha_4, \alpha_3, \alpha_4 \rangle$
A_3	$\langle \alpha_2, \alpha_3, \alpha_4 \rangle, \langle \alpha_1 + \alpha_2, \alpha_3, \alpha_4 \rangle$
D_4	$\langle \alpha_1, \alpha_2, \alpha_3, \alpha_4 \rangle$

Table 6: Realisations of the sub-root systems containing Θ for $\Delta(\Gamma) = A_3$

where $\langle \dots \rangle$ means the root system generated by "...".

Set (e_1, e_2, e_3, e_4) an orthonormal basis of $\mathfrak{h}'$ of type D_4 , and define $(\epsilon_1, \epsilon_2, \epsilon_3, \epsilon_4)$ its dual basis. Following [2], the simple roots of D_4 are

$$\begin{cases} \alpha_1 &= \epsilon_1 - \epsilon_2, \\ \alpha_2 &= \epsilon_2 - \epsilon_3, \\ \alpha_3 &= \epsilon_3 - \epsilon_4, \\ \alpha_4 &= \epsilon_3 + \epsilon_4. \end{cases}$$

Let $h' = (\xi_1, \xi_2, \xi_3, \xi_4) \in \mathfrak{h}'$, with the coordinates being expressed in the basis (e_1, e_2, e_3, e_4) . Then $h' \in H_{\alpha_3} \cap H_{\alpha_4}$ if and only if $\xi_3 = \xi_4 = 0$.

Using [13], one can compute the flat coordinates of D_4 restricted to the subspace $H_{\alpha_3} \cap H_{\alpha_4} \subset \mathfrak{h}'$ and obtains

$$\begin{cases} \psi_2(\xi_1, \xi_2, 0, 0) &= \xi_1^2 + \xi_2^2, \\ \psi_4(\xi_1, \xi_2, 0, 0) &= -\frac{1}{4}(\xi_1^2 - \xi_2^2)^2, \\ \psi_6(\xi_1, \xi_2, 0, 0) &= -\frac{1}{6}(\xi_1^2 + \xi_2^2)\xi_1^2\xi_2^2 + \frac{7}{216}(\xi_1^2 + \xi_2^2)^3, \\ \psi(\xi_1, \xi_2, 0, 0) &= 0. \end{cases}$$

One notices that $\psi_6 = -\frac{1}{108}\psi_2^3 - \frac{1}{6}\psi_2\psi_4$.

Define

$$f : (\mathfrak{h}/W)^\Omega \rightarrow \mathfrak{h}'/W'$$

$$\begin{pmatrix} t_2 \\ 0 \\ t_4 \end{pmatrix} \mapsto \begin{pmatrix} t_2 \\ t_4 - \frac{t_2^2}{8} \\ \frac{5}{432}t_2^3 - \frac{1}{6}t_2t_4 \\ 0 \end{pmatrix}$$

The morphism f is clearly injective, and if $(\psi_2, \psi_4, \psi_6, 0) \in \pi'(H_{\alpha_3} \cap H_{\alpha_4})$, then by setting $t_2 = \psi_2$ and $t_4 = \psi_4 + \frac{1}{8}\psi_2^2$, it follows that $f(t_2, 0, t_4) = (\psi_2, \psi_4, \psi_6, 0)$. Hence

$$f : (\mathfrak{h}/W)^\Omega \xrightarrow{\cong} \pi'(H_{\alpha_3} \cap H_{\alpha_4}) \subset \mathfrak{h}'/W'.$$

Let ϕ' be a sub-root system in Table 6, and set $h' \in \cap_{\alpha \in \phi'} H_\alpha$. With the formulas given before, one can compute $(t_2, 0, t_4) = f^{-1}(\pi'(h'))$, and verify the following correspondence:

ϕ' of type $A_1 + A_1$	$\longleftrightarrow$	$(t_2, 0, t_4)$ generic,
ϕ' of type $A_1 + A_1 + A_1$	$\longleftrightarrow$	$(t_2, 0, t_4)$ verifies $t_4 = \frac{t_2^2}{8} \neq 0$,
ϕ' of type A_3	$\longleftrightarrow$	$(t_2, 0, t_4)$ verifies $t_4 = -\frac{t_2^2}{8} \neq 0$,
ϕ' of type D_4	$\longleftrightarrow$	$(t_2, 0, t_4) = (0, 0, 0)$.

With Table 5, we see that the singular configuration in the fiber of $\overline{\alpha^\Omega}$ above $f^{-1}(\pi'(h'))$ is of the same type as ϕ' . Therefore for the type $A_3 - B_2 - D_4$, the map f realises a bijection between the singular configurations of the fibers of $\overline{\alpha^\Omega}$ and the sub-root systems of D_4 containing $\Theta = \{\alpha_3, \alpha_4\}$.

4.3. Case $A_5 - B_3 - D_5$

In this part $\Gamma = \mathcal{C}_6$, $\Gamma' = \mathcal{D}_3$ and $\Omega = \mathbb{Z}/2\mathbb{Z} = \langle \sigma \rangle$. The map α^Ω is computed in [4] and is given by the projection

$$\begin{aligned} X_{\Gamma, \Omega} &= \{((x, y, z), (t_2, 0, t_4, 0, t_6)) \in \mathbb{C}^3 \times \mathfrak{h}/W \mid z^6 + t_2 z^4 + (t_4 + \frac{t_2^2}{4})z^2 + t_6 + \frac{t_2 t_4}{6} + \frac{t_2^3}{108} = xy\} \\ &\downarrow \alpha^\Omega \\ (\mathfrak{h}/W)^\Omega &= \{(t_2, 0, t_4, 0, t_6) \in \mathfrak{h}/W\} \end{aligned}$$

Furthermore, the action of Ω on a fiber is $\sigma.(x, y, z) = (-y, -x, -z)$. The quotient $\overline{\alpha^\Omega}$ is then given by

$$\begin{aligned} X_{\Gamma, \Omega}/\Omega &= \{((X, Z, W), (t_2, 0, t_4, 0, t_6)) \in \mathbb{C}^3 \times \mathfrak{h}/W \mid Z(X^2 + 4Z^3) + W^2 + 4t_2 Z^3 \\ &\quad + 4(t_4 + \frac{t_2^2}{4})Z^2 + 4(t_6 + \frac{t_2 t_4}{6} + \frac{t_2^3}{108})Z = 0\} \\ &\downarrow \overline{\alpha^\Omega} \\ (\mathfrak{h}/W)^\Omega &= \{(t_2, 0, t_4, 0, t_6) \in \mathfrak{h}/W\} \end{aligned}$$

Before quotient, the discriminant of α^Ω is given by $H_1 \cup H_2 \subset (\mathfrak{h}/W)^\Omega$ with

$$H_1 = \{t_6 + \frac{t_2 t_4}{6} + \frac{t_2^3}{108} = 0\}, H_2 = \{-\frac{t_2^6}{432} + \frac{t_2^4 t_4}{12} - \frac{t_2^2 t_4^2}{4} - 9t_2 t_4 t_6 + 4t_4^3 + 27t_6^2 = 0\}.$$

After quotient, Theorem 2.3 says that the discriminant is all of $(\mathfrak{h}/W)^\Omega$. By studying explicitly the singularities in the fibers $(\alpha^\Omega)^{-1}(t_2, 0, t_4, 0, t_6)$ and $(\overline{\alpha^\Omega})^{-1}(t_2, 0, t_4, 0, t_6)$, we obtain the following types:

$(t_2, 0, t_4, 0, t_6)$	Configuration of $(\alpha^\Omega)^{-1}(t_2, 0, t_4, 0, t_6)$	$(\overline{\alpha^\Omega})^{-1}(t_2, 0, t_4, 0, t_6)$
Generic point	$p_1 + p_2$	$A_1 + A_1$
$H_1 \cap \{t_4 \neq 0\}$	A_1	A_3
$H_1 \cap \{t_4 = 0, t_2 \neq 0\}$	$A_1 + A_1 \xrightarrow{\Omega} A_1$	$A_3 + A_1$
$H_1 \cap \{t_4 = -\frac{t_2^2}{4} \neq 0\}$	A_3	D_4
$H_2 \setminus \{0\}$	$p_1 + p_2 + A_1 + A_1 \xrightarrow{\Omega} A_1$	$A_1 + A_1 + A_1$
$H_2 \cap \{t_4 = \frac{t_2^2}{12} \neq 0\}$	$p_1 + p_2 + A_2 + A_2 \xrightarrow{\Omega} A_2$	$A_1 + A_1 + A_2$
$t_2 = t_4 = t_6 = 0$	A_5	D_5

Table 7: Configurations of the fibers of α^Ω and $\overline{\alpha^\Omega}$ for $\Delta(\Gamma) = A_5$

with p_i ($i = 1, 2$) denoting a smooth point, and if the action of Ω is not mentioned, the point (smooth or singular) is fixed.

Let $\Pi' = \{\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5\}$ be a set of simple roots of the root system of type D_5 (cf. Table 4 for the numbering). All the sub-root systems ϕ' of D_5 containing $\Theta = \{\alpha_4, \alpha_5\}$, as well as their realisations are given in the following table:

Type of ϕ'	Realisations of ϕ'
$A_1 + A_1$	$\langle \alpha_4, \alpha_5 \rangle$
$A_1 + A_1 + A_1$	$\langle \alpha_1, \alpha_4, \alpha_5 \rangle, \langle \alpha_2, \alpha_4, \alpha_5 \rangle, \langle \alpha_1 + \alpha_2 + \alpha_4, \alpha_5 \rangle, \langle \alpha_1 + 2\alpha_2 + 2\alpha_3 + \alpha_4 + \alpha_5, \alpha_4, \alpha_5 \rangle,$ $\langle \alpha_1 + \alpha_2 + 2\alpha_3 + \alpha_4 + \alpha_5, \alpha_4, \alpha_5 \rangle, \langle \alpha_2 + 2\alpha_3 + \alpha_4 + \alpha_5, \alpha_4, \alpha_5 \rangle$
A_3	$\langle \alpha_3, \alpha_4, \alpha_5 \rangle, \langle \alpha_2 + \alpha_3, \alpha_4, \alpha_5 \rangle, \langle \alpha_1 + \alpha_2 + \alpha_3, \alpha_4, \alpha_5 \rangle$
$A_1 + A_1 + A_2$	$\langle \alpha_1, \alpha_2, \alpha_4, \alpha_5 \rangle, \langle \alpha_1, \alpha_2 + 2\alpha_3 + \alpha_4 + \alpha_5, \alpha_4, \alpha_5 \rangle,$ $\langle \alpha_2, \alpha_1 + \alpha_2 + 2\alpha_3 + \alpha_4 + \alpha_5, \alpha_4, \alpha_5 \rangle, \langle \alpha_1 + \alpha_2, \alpha_2 + 2\alpha_3 + \alpha_4 + \alpha_5, \alpha_4, \alpha_5 \rangle$
$A_3 + A_1$	$\langle \alpha_1, \alpha_3, \alpha_4, \alpha_5 \rangle, \langle \alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_4, \alpha_5 \rangle, \langle \alpha_1 + 2\alpha_2 + 2\alpha_3 + \alpha_4 + \alpha_5, \alpha_3, \alpha_4, \alpha_5 \rangle,$ $\langle \alpha_1 + \alpha_2 + 2\alpha_3 + \alpha_4 + \alpha_5, \alpha_2 + \alpha_3, \alpha_4, \alpha_5 \rangle, \langle \alpha_1 + \alpha_2 + \alpha_3, \alpha_2, \alpha_4, \alpha_5 \rangle,$ $\langle \alpha_1 + \alpha_2 + \alpha_3, \alpha_2 + 2\alpha_3 + \alpha_4 + \alpha_5, \alpha_4, \alpha_5 \rangle,$
D_4	$\langle \alpha_2, \alpha_3, \alpha_4, \alpha_5 \rangle, \langle \alpha_1 + \alpha_2, \alpha_3, \alpha_4, \alpha_5 \rangle, \langle \alpha_1, \alpha_2 + \alpha_3, \alpha_4, \alpha_5 \rangle$
D_5	$\langle \alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5 \rangle$

Table 8: Realisations of the sub-root systems containing Θ for $\Delta(\Gamma) = A_5$

Set $(e_1, e_2, e_3, e_4, e_5)$ an orthonormal basis of $\mathfrak{h}'$ of type D_5 , and define $(\epsilon_1, \epsilon_2, \epsilon_3, \epsilon_4, \epsilon_5)$ its dual basis. Following [2], the simple roots of D_5 are

$$\begin{cases} \alpha_1 &= \epsilon_1 - \epsilon_2, \\ \alpha_2 &= \epsilon_2 - \epsilon_3, \\ \alpha_3 &= \epsilon_3 - \epsilon_4, \\ \alpha_4 &= \epsilon_4 - \epsilon_5. \\ \alpha_5 &= \epsilon_4 + \epsilon_5. \end{cases}$$

Let $h' = (\xi_1, \xi_2, \xi_3, \xi_4, \xi_5) \in \mathfrak{h}'$, with the coordinates being expressed in the basis $(e_1, e_2, e_3, e_4, e_5)$. Then $h' \in H_{\alpha_4} \cap H_{\alpha_5}$ if and only if $\xi_4 = \xi_5 = 0$.

Using [13], one can compute the flat coordinates of D_5 restricted to the subspace $H_{\alpha_4} \cap H_{\alpha_5} \subset \mathfrak{h}'$ and obtain

$$\begin{cases} \psi_2(\xi_1, \xi_2, \xi_3) &= \xi_1^2 + \xi_2^2 + \xi_3^2, \\ \psi_4(\xi_1, \xi_2, \xi_3) &= \xi_1^2 \xi_2^2 + \xi_1^2 \xi_3^2 + \xi_2^2 \xi_3^2 - \frac{5}{16}(\xi_1^2 + \xi_2^2 + \xi_3^2)^2, \\ \psi_6(\xi_1, \xi_2, \xi_3) &= \xi_1^2 \xi_2^2 \xi_3^2 - \frac{3}{8}(\xi_1^2 + \xi_2^2 + \xi_3^2)(\xi_1^2 \xi_2^2 + \xi_1^2 \xi_3^2 + \xi_2^2 \xi_3^2) + \frac{11}{128}(\xi_1^2 + \xi_2^2 + \xi_3^2)^3, \\ \psi_8(\xi_1, \xi_2, \xi_3) &= -\frac{1}{8}(\xi_1^2 + \xi_2^2 + \xi_3^2)\xi_1^2 \xi_2^2 \xi_3^2 - \frac{1}{16}(\xi_1^2 \xi_2^2 + \xi_1^2 \xi_3^2 + \xi_2^2 \xi_3^2)^2 \\ &\quad + \frac{9}{128}(\xi_1^2 \xi_2^2 + \xi_1^2 \xi_3^2 + \xi_2^2 \xi_3^2)(\xi_1^2 + \xi_2^2 + \xi_3^2)^2 - \frac{51}{4096}(\xi_1^2 + \xi_2^2 + \xi_3^2)^4, \\ \psi(\xi_1, \xi_2, \xi_3) &= 0. \end{cases}$$

One can see that $\psi_8 = -\frac{1}{2048}\psi_2^4 - \frac{1}{8}\psi_2\psi_6 - \frac{1}{64}\psi_2^2\psi_4 - \frac{1}{16}\psi_4^2$.

Define

$$f: (\mathfrak{h}/W)^\Omega \rightarrow \mathfrak{h}'/W'$$

$$\begin{pmatrix} t_2 \\ 0 \\ t_4 \\ 0 \\ t_6 \end{pmatrix} \mapsto \begin{pmatrix} t_2 \\ t_4 - \frac{1}{16}t_2^2 \\ t_6 - \frac{5}{24}t_2t_4 + \frac{5}{3456}t_2^3 \\ \frac{7}{110592}t_2^4 - \frac{1}{8}t_2t_6 + \frac{7}{384}t_2^2t_4 - \frac{1}{16}t_4^2 \\ 0 \end{pmatrix}$$

We see that f is an injective morphism, and if $(\psi_2, \psi_4, \psi_6, \psi_8, 0) \in \pi'(H_{\alpha_4} \cap H_{\alpha_5})$, then by setting

$$\begin{cases} t_2 = \psi_2, \\ t_4 = \psi_4 + \frac{1}{16}\psi_2^2, \\ t_6 = \psi_6 + \frac{5}{24}\psi_2\psi_4 + \frac{5}{432}\psi_2^3, \end{cases}$$

it follows that $f(t_2, 0, t_4, 0, t_6) = (\psi_2, \psi_4, \psi_6, \psi_8, 0)$. Hence

$$f: (\mathfrak{h}/W)^\Omega \xrightarrow{\cong} \pi'(H_{\alpha_4} \cap H_{\alpha_5}) \subset \mathfrak{h}'/W'.$$

Let ϕ' be a sub-root system in Table 8, and set $h' \in \bigcap_{\alpha \in \phi'} H_\alpha$. With the formulas just given, one can compute $(t_2, 0, t_4, 0, t_6) = f^{-1}(\pi'(h'))$, and check the following correspondence:

ϕ' of type $A_1 + A_1$	$\longleftrightarrow$	$(t_2, 0, t_4)$ generic,
ϕ' of type $A_1 + A_1 + A_1$	$\longleftrightarrow$	$(t_2, 0, t_4)$ verifies $(t_2, 0, t_4, 0, t_6) \in H_2 \setminus \{0\}$,
ϕ' of type A_3	$\longleftrightarrow$	$(t_2, 0, t_4)$ verifies $(t_2, 0, t_4, 0, t_6) \in H_1 \cap \{t_4 \neq 0\}$,
ϕ' of type $A_2 + A_1 + A_1$	$\longleftrightarrow$	$(t_2, 0, t_4, 0, t_6) \in H_2 \cap \{t_4 = \frac{t_2^2}{12} \neq 0\}$,
ϕ' of type $A_3 + A_1$	$\longleftrightarrow$	$(t_2, 0, t_4, 0, t_6) \in H_1 \cap \{t_4 = 0, t_2 \neq 0\}$,
ϕ' of type D_4	$\longleftrightarrow$	$(t_2, 0, t_4, 0, t_6) \in H_1 \cap \{t_4 = -\frac{t_2^2}{4} \neq 0\}$,
ϕ' of type D_5	$\longleftrightarrow$	$(t_2, 0, t_4, 0, t_6) = (0, 0, 0, 0, 0)$.

With Table 7, we see that the singular configuration in the fiber of $\overline{\alpha^\Omega}$ above $f^{-1}(\pi'(h'))$ is of the same type as ϕ' . Therefore for the type $A_5 - B_3 - D_5$, the map f realises a bijection between the singular configurations of the fibers of $\overline{\alpha^\Omega}$ and the sub-root systems of D_5 containing $\Theta = \{\alpha_4, \alpha_5\}$.

4.4. Case $D_4 - C_3 - D_6$

Here $\Gamma = \mathcal{D}_2$, $\Gamma' = \mathcal{D}_4$ and $\Omega = \mathbb{Z}/2\mathbb{Z} = \langle \sigma \rangle$. It is known from [4] that the map α^Ω is the projection

$$X_{\Gamma, \Omega} = \{((x, y, z), (t_2, t_4, t_6, 0)) \in \mathbb{C}^3 \times \mathfrak{h}/W \mid z^2 = xy(x+y) - \frac{t_2}{2}xy - \frac{t_4}{4}x + \frac{1}{4}(t_6 + \frac{t_2t_4}{6} + \frac{t_2^3}{108})\}$$

$$\downarrow \alpha^\Omega$$

$$(\mathfrak{h}/W)^\Omega = \{(t_2, t_4, t_6, 0) \in \mathfrak{h}/W\}$$

Furthermore, the action of Ω on a fiber is $\sigma.(x, y, z) = (x, -x - y + \frac{t_2}{2}, -z)$, and the quotient $\overline{\alpha^\Omega}$ is given by

$$\begin{aligned}
X_{\Gamma, \Omega} / \Omega &= \{((X, Y, W), (t_2, t_4, t_6, 0)) \in \mathbb{C}^3 \times \mathfrak{h}/W \mid -\frac{1}{64}X^5 + XY^2 - W^2 \\
&\quad + A_{X^4}X^4 + A_{X^3}X^3 + A_{X^2}X^2 + A_X X + A_Y Y + A_0 = 0\} \\
&\quad \downarrow \overline{\alpha^\Omega} \\
(\mathfrak{h}/W)^\Omega &= \{(t_2, t_4, t_6, 0) \in \mathfrak{h}/W\}
\end{aligned}$$

with

$$\begin{cases}
A_{X^4} = \frac{t_2}{32}, \\
A_{X^3} = -\frac{3}{128}t_2^2 - \frac{1}{32}t_4, \\
A_{X^2} = \frac{7}{192}t_2t_4 + \frac{1}{32}t_6 + \frac{7}{864}t_2^3, \\
A_X = -\frac{1}{32}t_6t_2 - \frac{5}{384}t_2^2t_4 - \frac{35}{27648}t_2^4 - \frac{1}{64}t_4^2, \\
A_Y = \frac{1}{4}t_6 + \frac{1}{24}t_2t_4 + \frac{1}{432}t_2^3, \\
A_0 = \frac{1}{128}t_6t_2^2 + \frac{1}{32}t_6t_4 + \frac{11}{6912}t_2^3t_4 + \frac{1}{192}t_2t_4^2 + \frac{1}{13824}t_2^5.
\end{cases}$$

Before quotient, the discriminant of α^Ω is given by $L \cup H \subset (\mathfrak{h}/W)^\Omega$ with

$$L = \{t_6 + \frac{t_2t_4}{6} + \frac{t_2^3}{108} = 0\}, H = \{\frac{t_2^6}{6912} - \frac{t_2^4t_4}{192} + \frac{t_2^2t_4^2}{64} + \frac{9}{16}t_2t_4t_6 - \frac{t_4^3}{4} - \frac{27}{16}t_6^2 = 0\}.$$

After quotient, it is known from Theorem 2.3 that all of $(\mathfrak{h}/W)^\Omega$ is the discriminant. By studying explicitly the singularities in the fibers $(\alpha^\Omega)^{-1}(t_2, t_4, t_6, 0)$ and $(\overline{\alpha^\Omega})^{-1}(t_2, t_4, t_6, 0)$, we obtain the following types:

$(t_2, t_4, t_6, 0)$	Configuration of $(\alpha^\Omega)^{-1}(t_2, t_4, t_6, 0)$	$(\overline{\alpha^\Omega})^{-1}(t_2, t_4, t_6, 0)$
$(\mathbb{C}^3 \times \{0\}) \setminus (L \cup H)$	$p_1 + p_2 + p_3$	$A_1 + A_1 + A_1$
$L \setminus \{t_4 = -\frac{t_2^2}{4}\}$	$\begin{array}{c} A_1 \quad + \quad A_1 \\ \quad \quad \quad \Omega \end{array} + p_1 + p_2 + p_3$	$A_1 + A_1 + A_1 + A_1$
$H \setminus ((L \cap \{t_4 = -\frac{t_2^2}{4}\}) \cup \{t_4 = \frac{t_2^2}{12}\})$	$A_1 + p$	$A_3 + A_1$
$L \cap \{t_4 = -\frac{t_2^2}{4} \neq 0\}$	$A_3 + p$	$D_4 + A_1$
$H \cap \{t_4 = \frac{t_2^2}{12} \neq 0\}$	A_2	A_5
$(L \cap H) \setminus \{t_4 = -\frac{t_2^2}{4}\}$	$\begin{array}{c} A_1 \quad + \quad A_1 \\ \quad \quad \quad \Omega \end{array} + A_1 + p$	$A_3 + A_1 + A_1$
$t_2 = t_4 = t_6 = 0$	D_4	D_6

Table 9: Configurations of the fibers of α^Ω and $\overline{\alpha^\Omega}$ for $\Delta(\Gamma) = D_4$ and $\Omega = \mathbb{Z}/2\mathbb{Z}$

with p_i ($i = 1, 2, 3, \emptyset$) denoting a smooth point, and if the action of Ω is not mentioned, the point (smooth or singular) is fixed.

Let $\Pi' = \{\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6\}$ be a set of simple roots of the root system of type D_6 (cf. Table 4 for the numbering). All the sub-root systems ϕ' of D_6 containing $\Theta = \{\alpha_1, \alpha_3, \alpha_5\}$, as well as their realisations are given in the following table:

Type of ϕ'	Realisations of ϕ'
$A_1 + A_1 + A_1$	$\langle \alpha_1, \alpha_3, \alpha_5 \rangle$
$A_1 + A_1 + A_1 + A_1$	$\langle \alpha_1, \alpha_3, \alpha_5, \alpha_6 \rangle, \langle \alpha_1, \alpha_3, \alpha_5, \alpha_1 + 2\alpha_2 + 2\alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6 \rangle,$ $\langle \alpha_1, \alpha_3, \alpha_5, \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6 \rangle$
$A_3 + A_1$	$\langle \alpha_1, \alpha_3, \alpha_5, \alpha_2 \rangle, \langle \alpha_1, \alpha_3, \alpha_5, \alpha_4 \rangle, \langle \alpha_1, \alpha_3, \alpha_5, \alpha_2 + \alpha_3 + \alpha_4 \rangle,$ $\langle \alpha_1, \alpha_3, \alpha_5, \alpha_2 + \alpha_3 + \alpha_4 + \alpha_6 \rangle, \langle \alpha_1, \alpha_3, \alpha_5, \alpha_4 + \alpha_6 \rangle,$ $\langle \alpha_1, \alpha_3, \alpha_5, \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6 \rangle$
$A_3 + A_1 + A_1$	$\langle \alpha_1, \alpha_3, \alpha_5, \alpha_2, \alpha_6 \rangle, \langle \alpha_1, \alpha_3, \alpha_5, \alpha_4, \alpha_1 + 2\alpha_2 + 2\alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6 \rangle,$ $\langle \alpha_1, \alpha_3, \alpha_5, \alpha_2 + \alpha_3 + \alpha_4, \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6 \rangle,$ $\langle \alpha_1, \alpha_3, \alpha_5, \alpha_2 + \alpha_3 + \alpha_4 + \alpha_6, \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6 \rangle,$ $\langle \alpha_1, \alpha_3, \alpha_5, \alpha_4 + \alpha_6, \alpha_1 + 2\alpha_2 + 2\alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6 \rangle,$ $\langle \alpha_1, \alpha_3, \alpha_5, \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6, \alpha_6 \rangle$
$D_4 + A_1$	$\langle \alpha_1, \alpha_3, \alpha_5, \alpha_2, \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6 \rangle, \langle \alpha_1, \alpha_3, \alpha_5, \alpha_4, \alpha_6 \rangle,$ $\langle \alpha_1, \alpha_3, \alpha_5, \alpha_2 + \alpha_3 + \alpha_4, \alpha_6 \rangle$
A_5	$\langle \alpha_1, \alpha_3, \alpha_5, \alpha_2, \alpha_4 \rangle, \langle \alpha_1, \alpha_3, \alpha_5, \alpha_2 + \alpha_3 + \alpha_4 + \alpha_6, \alpha_4 \rangle, \langle \alpha_1, \alpha_3, \alpha_5, \alpha_2, \alpha_4 + \alpha_6 \rangle,$ $\langle \alpha_1, \alpha_3, \alpha_5, \alpha_4 + \alpha_6, \alpha_2 + \alpha_3 + \alpha_4 \rangle$
D_6	$\langle \alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6 \rangle$

Table 10: Realisations of the sub-root systems containing Θ for $\Delta(\Gamma) = D_4$ and $\Omega = \mathbb{Z}/2\mathbb{Z}$

Set $(e_1, e_2, e_3, e_4, e_5, e_6)$ an orthonormal basis of $\mathfrak{h}'$ of type D_6 , and define $(\epsilon_1, \epsilon_2, \epsilon_3, \epsilon_4, \epsilon_5, \epsilon_6)$ its dual basis. Following [2], the simple roots of D_5 are

$$\begin{cases} \alpha_1 &= \epsilon_1 - \epsilon_2, \\ \alpha_2 &= \epsilon_2 - \epsilon_3, \\ \alpha_3 &= \epsilon_3 - \epsilon_4, \\ \alpha_4 &= \epsilon_4 - \epsilon_5, \\ \alpha_5 &= \epsilon_5 - \epsilon_6, \\ \alpha_6 &= \epsilon_5 + \epsilon_6. \end{cases}$$

Let $h' = (\xi_1, \xi_2, \xi_3, \xi_4, \xi_5, \xi_6) \in \mathfrak{h}'$, with the coordinates being expressed in the basis $(e_1, e_2, e_3, e_4, e_5, e_6)$. Then $h' \in H_{\alpha_1} \cap H_{\alpha_3} \cap H_{\alpha_5}$ if and only if $\xi_1 = \xi_2$, $\xi_3 = \xi_4$ and $\xi_5 = \xi_6$.

Using [13], one can compute the flat coordinates of D_6 restricted to the subspace $H_{\alpha_1} \cap H_{\alpha_3} \cap H_{\alpha_5} \subset \mathfrak{h}'$ and obtain

$$\left\{ \begin{array}{lcl} \psi_2(\xi_1, \xi_3, \xi_5) & = & 2(\xi_1^2 + \xi_3^2 + \xi_5^2), \\ \psi_4(\xi_1, \xi_3, \xi_5) & = & -\frac{2}{5}\xi_1^4 - \frac{2}{5}\xi_3^4 - \frac{2}{5}\xi_5^4 + \frac{6}{5}\xi_1^2\xi_3^2 + \frac{6}{5}\xi_1^2\xi_5^2 + \frac{6}{5}\xi_3^2\xi_5^2, \\ \psi_6(\xi_1, \xi_3, \xi_5) & = & 2\xi_1^2\xi_3^2\xi_5^2, \\ \psi_8(\xi_1, \xi_3, \xi_5) & = & \frac{4}{125}\xi_1^8 + \frac{4}{125}\xi_3^8 + \frac{4}{125}\xi_5^8 - \frac{14}{125}\xi_1^6\xi_3^2 - \frac{14}{125}\xi_1^6\xi_5^2 - \frac{14}{125}\xi_3^6\xi_5^2 - \frac{14}{125}\xi_1^4\xi_3^4 - \frac{14}{125}\xi_1^4\xi_5^4 - \frac{14}{125}\xi_3^4\xi_5^4 + \frac{98}{125}\xi_1^2\xi_3^2\xi_5^4 + \frac{98}{125}\xi_1^2\xi_3^4\xi_5^2, \\ \psi_{10}(\xi_1, \xi_3, \xi_5) & = & -\frac{108}{625}\xi_1^2\xi_3^2\xi_5^6 - \frac{108}{625}\xi_1^6\xi_3^2\xi_5^2 - \frac{108}{625}\xi_1^2\xi_3^6\xi_5^2 - \frac{22}{3125}\xi_1^{10} - \frac{22}{3125}\xi_3^{10} - \frac{22}{3125}\xi_5^{10} \\ & & - \frac{24}{625}\xi_1^6\xi_3^4 - \frac{24}{625}\xi_1^4\xi_3^6 - \frac{24}{625}\xi_1^4\xi_5^6 - \frac{24}{625}\xi_3^4\xi_5^6 - \frac{24}{625}\xi_5^4\xi_3^6 - \frac{24}{625}\xi_5^4\xi_1^6 \\ & & + \frac{18}{625}\xi_1^8\xi_3^2 + \frac{18}{625}\xi_1^8\xi_5^2 + \frac{18}{625}\xi_3^8\xi_5^2 + \frac{18}{625}\xi_1^2\xi_3^8 + \frac{18}{625}\xi_1^2\xi_5^8 + \frac{18}{625}\xi_3^2\xi_5^8 \\ & & + \frac{648}{625}\xi_1^4\xi_3^2\xi_5^4 + \frac{648}{625}\xi_1^2\xi_3^4\xi_5^4 + \frac{648}{625}\xi_1^4\xi_3^4\xi_5^2, \\ \psi(\xi_1, \xi_3, \xi_5) & = & \xi_1^2\xi_3^2\xi_5^2. \end{array} \right.$$

We can verify that $\left\{ \begin{array}{lcl} \psi_8 & = & \frac{1}{5}\psi_2\psi_6 - \frac{1}{100}\psi_2^2\psi_4 + \frac{1}{10}\psi_4^2, \\ \psi_{10} & = & -\frac{1}{50000}\psi_2^5 + \frac{1}{50}\psi_2^2\psi_6 - \frac{1}{50}\psi_2\psi_4^2 + \frac{2}{5}\psi_4\psi_6, \\ \psi & = & \frac{1}{2}\psi_6. \end{array} \right.$

Define

$$f : (\mathfrak{h}/W)^\Omega \rightarrow \mathfrak{h}'/W'$$

$$\begin{pmatrix} t_2 \\ t_4 \\ t_6 \\ 0 \end{pmatrix} \mapsto \begin{pmatrix} t_2 \\ \frac{1}{2}t_4 + \frac{1}{40}t_2^2 \\ \frac{1}{4}t_6 + \frac{1}{24}t_2t_4 + \frac{1}{432}t_2^3 \\ \frac{1}{20}t_2t_6 + \frac{7}{1200}t_2^2t_4 + \frac{119}{432000}t_2^4 + \frac{1}{40}t_2^2 \\ \frac{133}{3600000}t_2^5 + \frac{3}{400}t_2^2t_6 + \frac{131}{108000}t_2^3t_4 + \frac{1}{300}t_2t_4^2 + \frac{1}{20}t_4t_6 \\ \frac{1}{8}t_6 + \frac{1}{48}t_2t_4 + \frac{1}{864}t_2^3 \end{pmatrix}$$

It is clear that f is an injective morphism, and if $(\psi_2, \psi_4, \psi_6, \psi_8, \psi_{10}, \psi) \in \pi'(H_{\alpha_1} \cap H_{\alpha_3} \cap H_{\alpha_5})$, then by setting

$$\left\{ \begin{array}{l} t_2 = \psi_2, \\ t_4 = 2\psi_4 - \frac{1}{20}\psi_2^2, \\ t_6 = 4\psi_6 - \frac{1}{3}\psi_2\psi_4 - \frac{1}{1080}\psi_2^3, \end{array} \right.$$

it follows that $f(t_2, t_4, t_6, 0) = (\psi_2, \psi_4, \psi_6, \psi_8, \psi_{10}, \psi)$. Hence

$$f : (\mathfrak{h}/W)^\Omega \xrightarrow{\cong} \pi'(H_{\alpha_1} \cap H_{\alpha_3} \cap H_{\alpha_5}) \subset \mathfrak{h}'/W'.$$

Let ϕ' be a sub-root system in Table 10, and set $h' \in \bigcap_{\alpha \in \phi'} H_\alpha$. With the formulas just given, one can compute $(t_2, t_4, t_6, 0) = f^{-1}(\pi'(h'))$, and verify the following correspondence:

ϕ' of type $A_1 + A_1 + A_1$	$\longleftrightarrow$	$(t_2, t_4, t_6, 0)$ generic,
ϕ' of type $A_1 + A_1 + A_1 + A_1$	$\longleftrightarrow$	$(t_2, t_4, t_6, 0) \in L \setminus \{t_4 = -\frac{t_2^2}{4}\}$,
ϕ' of type $A_3 + A_1$	$\longleftrightarrow$	$(t_2, t_4, t_6, 0) \in H \setminus ((L \cap \{t_4 = -\frac{t_2^2}{4}\}) \cup \{t_4 = \frac{t_2^2}{12}\})$,
ϕ' of type $A_3 + A_1 + A_1$	$\longleftrightarrow$	$(t_2, t_4, t_6, 0) \in (L \cap H) \setminus \{t_4 = -\frac{t_2^2}{4}\}$,
ϕ' of type $D_4 + A_1$	$\longleftrightarrow$	$(t_2, t_4, t_6, 0) \in L \cap \{t_4 = -\frac{t_2^2}{4} \neq 0\}$,
ϕ' of type A_5	$\longleftrightarrow$	$(t_2, t_4, t_6, 0) \in H \cap \{t_4 = \frac{t_2^2}{12} \neq 0\}$,
ϕ' of type D_6	$\longleftrightarrow$	$(t_2, t_4, t_6, 0) = (0, 0, 0, 0)$.

With Table 9, we see that the singular configuration in the fiber of $\overline{\alpha^\Omega}$ above $f^{-1}(\pi'(h'))$ is of the same type as ϕ' . Therefore for the type $D_4 - C_3 - D_6$, the map f realises a bijection between the singular configurations of the fibers of $\overline{\alpha^\Omega}$ and the sub-root systems of D_5 containing $\Theta = \{\alpha_1, \alpha_3, \alpha_5\}$.

4.5. Case $D_4 - G_2 - E_6$

In this section $\Gamma = \mathcal{D}_2$, $\Gamma' = \mathcal{T}$ and $\Omega = \mathbb{Z}/3\mathbb{Z} = \langle \rho \rangle$. The restriction α^Ω of the semiuniversal deformation α of a simple singularity of type D_4 is

$$\begin{aligned} X_{\Gamma, \Omega} &= \{((x, y, z), (t_2, 0, t_6, 0)) \in \mathbb{C}^3 \times \mathfrak{h}/W \mid z^2 = xy(x+y) - \frac{t_2}{2}xy + \frac{1}{4}(t_6 + \frac{t_2^3}{108})\} \\ &\downarrow \alpha^\Omega \\ (\mathfrak{h}/W)^\Omega &= \{(t_2, 0, t_6, 0) \in \mathfrak{h}/W\} \end{aligned}$$

The action of Ω on a fiber is $\rho.(x, y, z) = (y, -x - y + \frac{t_2}{2}, z)$, and so the quotient $\overline{\alpha^\Omega}$ is given by

$$\begin{aligned} X_{\Gamma, \Omega}/\Omega &= \{((X, Y, Z), (t_2, 0, t_6, 0)) \in \mathbb{C}^3 \times \mathfrak{h}/W \mid 11664X^4 - Y^3 - Z^2 - 324t_2X^2Y \\ &\quad - (189t_2^3 + 5832t_6)X^2 + (81t_2t_6 + \frac{15}{16}t_2^4)Y + \frac{11}{32}t_2^6 + \frac{189}{4}t_2^3t_6 + 729t_6^2 = 0\} \\ &\downarrow \overline{\alpha^\Omega} \\ (\mathfrak{h}/W)^\Omega &= \{(t_2, 0, t_6, 0) \in \mathfrak{h}/W\} \end{aligned}$$

Before quotient, the discriminant of α^Ω is given by $\{(t_6 + \frac{t_2^3}{108})(t_6 - \frac{t_2^3}{108}) = 0\}$, and after quotient, it is all of $(\mathfrak{h}/W)^\Omega$ because of Theorem 2.3. By studying explicitly the singularities in the fibers $(\alpha^\Omega)^{-1}(t_2, 0, t_6, 0)$ and $(\overline{\alpha^\Omega})^{-1}(t_2, 0, t_6, 0)$, we obtain the following types:

$(t_2, 0, t_6, 0)$	Configuration of $(\alpha^\Omega)^{-1}(t_2, 0, t_6, 0)$	$(\overline{\alpha^\Omega})^{-1}(t_2, 0, t_6, 0)$
Generic point	$p_1 + p_2$	$A_2 + A_2$
$t_6 = -\frac{t_2^3}{108} \neq 0$		$A_2 + A_2 + A_1$
$t_6 = \frac{t_2^3}{108} \neq 0$	A_1	A_5
$t_2 = t_6 = 0$	D_4	E_6

Table 11: Configurations of the fibers of α^Ω and $\overline{\alpha^\Omega}$ for $\Delta(\Gamma) = D_4$ and $\Omega = \mathbb{Z}/3\mathbb{Z}$

with p_i ($i = 1, 2$) denoting a smooth point, and if the action of Ω is not mentioned, the point (smooth or singular) is fixed.

Let $\Pi' = \{\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6\}$ be a set of simple roots of the root system of type E_6 . All the sub-root systems ϕ' of E_6 containing $\Theta = \{\alpha_1, \alpha_3, \alpha_5, \alpha_6\}$, as well as their realisations are given in the following table:

Type of ϕ'	Realisations of ϕ'
$A_2 + A_2$	$\langle \alpha_1, \alpha_3, \alpha_5, \alpha_6 \rangle$
$A_2 + A_2 + A_1$	$\langle \alpha_1, \alpha_3, \alpha_5, \alpha_6, \alpha_2 \rangle, \langle \alpha_1, \alpha_3, \alpha_5, \alpha_6, \alpha_1 + \alpha_2 + 2\alpha_3 + 3\alpha_4 + 2\alpha_5 + \alpha_6 \rangle,$ $\langle \alpha_1, \alpha_3, \alpha_5, \alpha_6, \alpha_1 + 2\alpha_2 + 2\alpha_3 + 3\alpha_4 + 2\alpha_5 + \alpha_6 \rangle$
A_5	$\langle \alpha_1, \alpha_3, \alpha_5, \alpha_6, \alpha_4 \rangle, \langle \alpha_1, \alpha_3, \alpha_5, \alpha_6, \alpha_2 + \alpha_4 \rangle, \langle \alpha_1, \alpha_3, \alpha_5, \alpha_6, \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 \rangle$
E_6	$\langle \alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6 \rangle$

Table 12: Realisations of the sub-root systems containing Θ for $\Delta(\Gamma) = D_4$ and $\Omega = \mathbb{Z}/3\mathbb{Z}$

Set $h' = (\xi_1, \xi_2, \xi_3, \xi_4, \xi_5, \xi_6) \in \mathfrak{h}'$, with the coordinates being expressed in the basis of fundamental coweights $(\omega_1^\vee, \omega_2^\vee, \omega_3^\vee, \omega_4^\vee, \omega_5^\vee, \omega_6^\vee)$. Then $h' \in H_{\alpha_1} \cap H_{\alpha_3} \cap H_{\alpha_5} \cap H_{\alpha_6}$ if and only if $\xi_1 = \xi_3 = \xi_5 = \xi_6 = 0$.

Using [7] and [13], we compute the flat coordinates of E_6 restricted to the subspace $H_{\alpha_1} \cap H_{\alpha_3} \cap H_{\alpha_5} \cap H_{\alpha_6} \subset \mathfrak{h}'$ and obtain

$$\begin{cases} \psi_2(\xi_2, \xi_4) &= 2\xi_2^2 + 6\xi_2\xi_4 + 6\xi_4^2, \\ \psi_5(\xi_2, \xi_4) &= 0, \\ \psi_6(\xi_2, \xi_4) &= -\xi_2^6 - 9\xi_2^5\xi_4 - 30\xi_2^4\xi_4^2 - 45\xi_2^3\xi_4^3 - 30\xi_2^2\xi_4^4 - 9\xi_2\xi_4^5 - 3\xi_4^6, \\ \psi_8(\xi_2, \xi_4) &= \frac{1}{12}(\xi_2^2 + 3\xi_2\xi_4 + 3\xi_4^2)(5\xi_2^6 + 45\xi_2^5\xi_4 + 144\xi_2^4\xi_4^2 + 189\xi_2^3\xi_4^3 + 72\xi_2^2\xi_4^4 - 27\xi_2\xi_4^5 - 9\xi_4^6), \\ \psi_9(\xi_2, \xi_4) &= 0, \\ \psi_{12}(\xi_2, \xi_4) &= \frac{693}{4}\xi_2^2\xi_4^{10} + \frac{189}{2}\xi_2\xi_4^{11} - \frac{2277}{2}\xi_2^5\xi_4^7 - \frac{1947}{2}\xi_2^7\xi_4^5 - \frac{9}{2}\xi_2^{11}\xi_4 - \frac{143}{4}\xi_2^{10}\xi_4^2 - 165\xi_2^9\xi_4^3 \\ &\quad - \frac{1089}{2}\xi_2^4\xi_4^8 - \frac{5225}{4}\xi_2^6\xi_4^6 + \frac{63}{4}\xi_4^{12} - \frac{1}{4}\xi_2^{12} - \frac{979}{2}\xi_2^8\xi_4^4. \end{cases}$$

There are the following relations among the flat coordinates:

$$\begin{cases} \psi_8 &= -\frac{1}{192}\psi_2^4 - \frac{1}{4}\psi_2\psi_6, \\ \psi_{12} &= \frac{1}{1536}\psi_2^6 - \frac{1}{8}\psi_6^2 + \frac{1}{48}\psi_2^3\psi_6. \end{cases}$$

Define

$$f: (\mathfrak{h}/W)^\Omega \rightarrow \mathfrak{h}'/W'$$

$$\begin{pmatrix} t_2 \\ 0 \\ t_6 \\ 0 \end{pmatrix} \mapsto \begin{pmatrix} t_2 \\ 0 \\ -6t_6 - \frac{5}{72}t_2^3 \\ \frac{7}{576}t_2^4 + \frac{3}{2}t_2t_6 \\ 0 \\ -\frac{29}{20736}t_2^6 - \frac{9}{2}t_6^2 - \frac{11}{48}t_6t_2^3 \end{pmatrix}$$

The map f is an injective morphism, and if $(\psi_2, \psi_5, \psi_6, \psi_8, \psi_9, \psi_{12}) \in \pi'(H_{\alpha_1} \cap H_{\alpha_3} \cap H_{\alpha_5} \cap H_{\alpha_6})$, then by setting

$$\begin{cases} t_2 = \psi_2, \\ t_6 = -\frac{1}{6}\psi_6 - \frac{5}{432}\psi_2^3, \end{cases}$$

it follows that $f(t_2, 0, t_6, 0) = (\psi_2, \psi_5, \psi_6, \psi_8, \psi_9, \psi_{12})$. Hence

$$f : (\mathfrak{h}/W)^\Omega \xrightarrow{\cong} \pi'(H_{\alpha_1} \cap H_{\alpha_3} \cap H_{\alpha_5} \cap H_{\alpha_6}) \subset \mathfrak{h}'/W'.$$

Let ϕ' be a sub-root system in Table 12, and set $h' \in \bigcap_{\alpha \in \phi'} H_\alpha$. With the preceding expressions, we compute $(t_2, 0, t_6, 0) = f^{-1}(\pi'(h'))$, and verify the following correspondence:

$$\begin{array}{ll} \phi' \text{ of type } A_2 + A_2 & \longleftrightarrow (t_2, 0, t_6, 0) \text{ generic,} \\ \phi' \text{ of type } A_2 + A_2 + A_1 & \longleftrightarrow (t_2, 0, t_6, 0) \in \{t_6 = -\frac{t_2^3}{108} \neq 0\}, \\ \phi' \text{ of type } A_5 & \longleftrightarrow (t_2, 0, t_6, 0) \in \{t_6 = \frac{t_2^3}{108} \neq 0\}, \\ \phi' \text{ of type } E_6 & \longleftrightarrow (t_2, 0, t_6, 0) = (0, 0, 0, 0). \end{array}$$

With Table 11, we see that the singular configuration in the fiber of $\overline{\alpha^\Omega}$ above $f^{-1}(\pi'(h'))$ is of the same type as ϕ' . Hence for $D_4 - G_2 - E_6$, the map f realises a bijection between the singular configurations of the fibers of $\overline{\alpha^\Omega}$ and the sub-root systems of E_6 containing $\Theta = \{\alpha_1, \alpha_3, \alpha_5, \alpha_6\}$.

4.6. Case $D_4 - G_2 - E_7$

In this section $\Gamma = \mathcal{D}_2$, $\Gamma' = \mathcal{O}$ and $\Omega = \mathfrak{S}_3 = \langle \rho, \sigma \rangle$. The restriction α^Ω above the fixed points is the same as the one in the preceding part, ie

$$\begin{array}{c} X_{\Gamma, \Omega} = \{(x, y, z), (t_2, 0, t_6, 0) \in \mathbb{C}^3 \times \mathfrak{h}/W \mid z^2 = xy(x+y) - \frac{t_2}{2}xy + \frac{1}{4}(t_6 + \frac{t_2^3}{108})\} \\ \downarrow \alpha^\Omega \\ (\mathfrak{h}/W)^\Omega = \{(t_2, 0, t_6, 0) \in \mathfrak{h}/W\} \end{array}$$

and the action of Ω on a fiber is

$$\sigma.(x, y, z) = (x, -x - y + \frac{t_2}{2}, -z) \text{ and } \rho.(x, y, z) = (y, -x - y + \frac{t_2}{2}, z).$$

The quotient $\overline{\alpha^\Omega}$ is then given by

$$\begin{array}{c} X_{\Gamma, \Omega}/\Omega = \{(X, Y, Z), (t_2, 0, t_6, 0) \in \mathbb{C}^3 \times \mathfrak{h}/W \mid X^3Y - 11664Y^3 + Z^2 + 324t_2XY^2 \\ + (189t_2^3 + 5832t_6)Y^2 - (\frac{15}{16}t_2^4 + 81t_2t_6)XY - (\frac{11}{32}t_2^6 + \frac{189}{4}t_2^3t_6 + 729t_6^2)Y = 0\} \\ \downarrow \overline{\alpha^\Omega} \\ (\mathfrak{h}/W)^\Omega = \{(t_2, 0, t_6, 0) \in \mathfrak{h}/W\} \end{array}$$

The discriminant of the deformation before quotient is $\{(t_6 + \frac{t_2^3}{108})(t_6 - \frac{t_2^3}{108}) = 0\}$, and after quotient it is $(\mathfrak{h}/W)^\Omega$. The study of the fibers of α^Ω and $\overline{\alpha^\Omega}$ gives the following singular configurations:

$(t_2, 0, t_6, 0)$	Configuration of $(\alpha^\Omega)^{-1}(t_2, 0, t_6, 0)$	$(\overline{\alpha^\Omega})^{-1}(t_2, 0, t_6, 0)$
Generic point	smooth and no Ω -fixed point	$A_2 + A_1 + A_1 + A_1$
$t_6 = -\frac{t_2^3}{108} \neq 0$		$A_3 + A_2 + A_1$
$t_6 = \frac{t_2^3}{108} \neq 0$	A_1 (Ω -fixed)	$D_5 + A_1$
$t_2 = t_6 = 0$	D_4 (Ω -fixed)	E_7

Table 13: Configurations of the fibers of α^Ω and $\overline{\alpha^\Omega}$ for $\Delta(\Gamma) = D_4$ and $\Omega = \mathfrak{S}_3$

Let $\Pi' = \{\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6, \alpha_7\}$ be a set of simple roots of the root system of type E_7 . All the sub-root systems ϕ' of E_7 containing $\Theta = \{\alpha_1, \alpha_2, \alpha_3, \alpha_5, \alpha_7\}$, as well as their realisations are given in the following table:

Type of ϕ'	Realisations of ϕ'
$A_2 + A_1 + A_1 + A_1$	$\langle \alpha_1, \alpha_2, \alpha_3, \alpha_5, \alpha_7 \rangle$
$A_3 + A_2 + A_1$	$\langle \alpha_1, \alpha_2, \alpha_3, \alpha_5, \alpha_7, \alpha_6 \rangle, \langle \alpha_1, \alpha_2, \alpha_3, \alpha_5, \alpha_7, \alpha_1 + \alpha_2 + 2\alpha_3 + 3\alpha_4 + 2\alpha_5 + 2\alpha_6 + \alpha_7 \rangle$
$D_5 + A_1$	$\langle \alpha_1, \alpha_2, \alpha_3, \alpha_5, \alpha_7, \alpha_4 \rangle, \langle \alpha_1, \alpha_2, \alpha_3, \alpha_5, \alpha_7, \alpha_4 + \alpha_5 + \alpha_6 \rangle, \langle \alpha_1, \alpha_2, \alpha_3, \alpha_5, \alpha_7, \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6 \rangle$
E_7	$\langle \alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6, \alpha_7 \rangle$

Table 14: Realisations of the sub-root systems containing Θ for $\Delta(\Gamma) = D_4$ and $\Omega = \mathfrak{S}_3$

Set $\mathfrak{h}' = \{(\xi_1, \xi_2, \xi_3, \xi_4, \xi_5, \xi_6, \xi_7, \xi_8) \in \mathbb{C}^8 \mid \xi_7 + \xi_8 = 0\}$, $(e_1, \dots, e_8)$ the canonical basis of $\mathbb{C}^8$, and $(\epsilon_1, \dots, \epsilon_8)$ its dual basis. Following [2], the simple roots of E_7 are

$$\left\{ \begin{array}{lcl} \alpha_1 & = & \frac{1}{2}(\epsilon_1 + \epsilon_8) - \frac{1}{2}(\epsilon_2 + \epsilon_3 + \epsilon_4 + \epsilon_5 + \epsilon_6 + \epsilon_7), \\ \alpha_2 & = & \epsilon_1 + \epsilon_2, \\ \alpha_3 & = & \epsilon_2 - \epsilon_1, \\ \alpha_4 & = & \epsilon_3 - \epsilon_2, \\ \alpha_5 & = & \epsilon_4 - \epsilon_3, \\ \alpha_6 & = & \epsilon_5 - \epsilon_4, \\ \alpha_7 & = & \epsilon_6 - \epsilon_5. \end{array} \right.$$

Set $h' = (\xi_1, \xi_2, \xi_3, \xi_4, \xi_5, \xi_6, \xi_7) \in \mathfrak{h}'$ (we write here $(\xi_1, \xi_2, \xi_3, \xi_4, \xi_5, \xi_6, \xi_7)$ for $(\xi_1, \xi_2, \xi_3, \xi_4, \xi_5, \xi_6, \xi_7, -\xi_7)$). Then $h' \in H_{\alpha_1} \cap H_{\alpha_2} \cap H_{\alpha_3} \cap H_{\alpha_5} \cap H_{\alpha_7}$ if and only if $\xi_1 = \xi_2 = 0$, $\xi_3 = \xi_4$, $\xi_5 = \xi_6$ and $\xi_7 = -\xi_3 - \xi_5$.

Using [1] and [10], one can compute the flat coordinates of E_7 restricted to the subspace $H_{\alpha_1} \cap H_{\alpha_2} \cap H_{\alpha_3} \cap H_{\alpha_5} \cap H_{\alpha_7} \subset \mathfrak{h}'$. We will therefore look at the flat coordinates as functions of ξ_3 and ξ_5 . The expressions are

$$\left\{ \begin{array}{l}
\psi_2(\xi_3, \xi_5) = \frac{2}{5}(\xi_3^2 + \xi_3\xi_5 + \xi_5^2), \\
\psi_6(\xi_3, \xi_5) = \frac{32176}{225}\xi_3^6 + \frac{32176}{75}\xi_3^5\xi_5 + \frac{53552}{75}\xi_3^4\xi_5^2 + \frac{160432}{225}\xi_3^3\xi_5^3 + \frac{53552}{75}\xi_3^2\xi_5^4 + \frac{32176}{75}\xi_3\xi_5^5 + \frac{32176}{225}\xi_5^6, \\
\psi_8(\xi_3, \xi_5) = \frac{16}{30375}(\xi_3^2 + \xi_3\xi_5 + \xi_5^2)(550819\xi_3^6 + 1652457\xi_3^5\xi_5 + 1389264\xi_3^4\xi_5^2 + 24433\xi_3^3\xi_5^3 \\
+ 1389264\xi_3^2\xi_5^4 + 1652457\xi_3\xi_5^5 + 550819\xi_5^6), \\
\psi_{10}(\xi_3, \xi_5) = \frac{96}{109375}(20743\xi_3^6 + 62229\xi_3^5\xi_5 + 41208\xi_3^4\xi_5^2 - 21299\xi_3^3\xi_5^3 + 41208\xi_3^2\xi_5^4 + 62229\xi_3\xi_5^5 \\
+ 20743\xi_5^6)(\xi_3^2 + \xi_3\xi_5 + \xi_5^2)^2, \\
\psi_{12}(\xi_3, \xi_5) = -\frac{42062501701}{398671875}\xi_3^8\xi_5^4 - \frac{4423023418}{102515625}\xi_3^9\xi_5^3 - \frac{62899959716}{5980078125}\xi_3^{10}\xi_5^2 - \frac{347826674932}{1993359375}\xi_3^7\xi_5^5 \\
- \frac{3081278138}{17940234375}\xi_3^{12} - \frac{3081278138}{17940234375}\xi_3^{12} - \frac{175228928248}{854296875}\xi_3^6\xi_5^6 - \frac{6162556276}{5980078125}\xi_3^{11}\xi_5 \\
- \frac{347826674932}{1993359375}\xi_3^5\xi_5^7 - \frac{42062501701}{398671875}\xi_3^4\xi_5^8 - \frac{4423023418}{102515625}\xi_3^3\xi_5^9 - \frac{62899959716}{5980078125}\xi_3^2\xi_5^{10} \\
- \frac{6162556276}{5980078125}\xi_3\xi_5^{11}, \\
\psi_{14}(\xi_3, \xi_5) = -\frac{4}{30903847734375}(\xi_3^2 + \xi_3\xi_5 + \xi_5^2)(1511960253367\xi_3^{12} + 9071761520202\xi_3^{11}\xi_5 \\
+ 67786465629432\xi_3^{10}\xi_5^2 + 255774514211975\xi_3^9\xi_5^3 + 617323843488330\xi_3^8\xi_5^4 \\
+ 1034437665403692\xi_3^7\xi_5^5 + 1226835303782847\xi_3^6\xi_5^6 + 1034437665403692\xi_3^5\xi_5^7 \\
+ 617323843488330\xi_3^4\xi_5^8 + 255774514211975\xi_3^3\xi_5^9 + 67786465629432\xi_3^2\xi_5^{10} \\
+ 9071761520202\xi_3\xi_5^{11} + 1511960253367\xi_5^{12}), \\
\psi_{18}(\xi_3, \xi_5) = -\frac{49900582548245699977888}{128185297421220703125}\xi_3^{18} - \frac{49900582548245699977888}{128185297421220703125}\xi_3^{18} - \frac{1808994581776446325173376}{42728432473740234375}\xi_3^3\xi_5^{15} \\
- \frac{43351951625476282697248}{2848562164916015625}\xi_3^2\xi_5^{16} - \frac{49900582548245699977888}{14242810824580078125}\xi_3^{17} - \frac{49900582548245699977888}{14242810824580078125}\xi_3^{17}\xi_5 \\
- \frac{43351951625476282697248}{2848562164916015625}\xi_3^{16}\xi_5^2 - \frac{1808994581776446325173376}{42728432473740234375}\xi_3^{15}\xi_5^3 - \frac{1224969840491929611874048}{14242810824580078125}\xi_3^{14}\xi_5^4 \\
- \frac{283014291225008940645632}{2034687260654296875}\xi_3^{13}\xi_5^5 - \frac{8202907266598286263520384}{42728432473740234375}\xi_3^{12}\xi_5^6 - \frac{338389353113795266600128}{14242810824580078125}\xi_3^{11}\xi_5^7 \\
- \frac{349873020975151098628384}{1294800984052734375}\xi_3^{10}\xi_5^8 - \frac{3288297534494448854110112}{11653208856474609375}\xi_3^9\xi_5^9 - \frac{349873020975151098628384}{1294800984052734375}\xi_3^8\xi_5^{10} \\
- \frac{338389353113795266600128}{14242810824580078125}\xi_3^7\xi_5^{11} - \frac{8202907266598286263520384}{42728432473740234375}\xi_3^6\xi_5^{12} - \frac{283014291225008940645632}{2034687260654296875}\xi_3^5\xi_5^{13} \\
- \frac{1224969840491929611874048}{14242810824580078125}\xi_3^4\xi_5^{14}.
\end{array} \right.$$

We can verify that

$$\left\{ \begin{array}{l}
\psi_8 = -\frac{2252645}{81}\psi_2^4 + \frac{473}{27}\psi_2\psi_6, \\
\psi_{10} = -\frac{557383}{105}\psi_2^5 + \frac{111}{35}\psi_2^2\psi_6, \\
\psi_{12} = -\frac{43251895481}{24494400}\psi_2^6 + \frac{1079173}{1360800}\psi_2^3\psi_6 - \frac{1}{103680}\psi_6^2, \\
\psi_{14} = -\frac{573683065303}{145496736}\psi_2^7 + \frac{10112840293}{4688228160}\psi_2^4\psi_6 - \frac{17821}{89299584}\psi_2\psi_6^2, \\
\psi_{18} = -\frac{15896711538141155833}{4023348492240}\psi_2^9 + \frac{391876556269181513}{64820614597200}\psi_2^6\psi_6 - \frac{7868764351687}{3601145255400}\psi_2^3\psi_6^2 - \frac{5}{419904}\psi_6^3.
\end{array} \right.$$

Define

$$f: (\mathfrak{h}/W)^\Omega \rightarrow \mathfrak{h}'/W'$$

$$\begin{pmatrix} t_2 \\ 0 \\ t_6 \\ 0 \end{pmatrix} \mapsto \begin{pmatrix} t_2 \\ \frac{18610}{9}t_2^3 + 18000t_6 \\ \frac{2044595}{243}t_2^4 + \frac{946000}{3}t_2t_6 \\ \frac{6247}{5}t_2^5 + \frac{399600}{7}t_2^2t_6 \\ -\frac{877545367}{5248800}t_2^6 + \frac{30746815}{2268}t_2^3t_6 - 3125t_6^2 \\ -\frac{4251411945217}{12658216032}t_2^7 + \frac{42570028475}{1775844}t_2^4t_6 - \frac{556906250}{8613}t_2t_6^2 \\ -\frac{134750219913739937987}{150013422353520}t_2^9 - \frac{277874830706221330}{4910652621}t_2^6t_6 - \frac{488086012279345000}{666878751}t_2^3t_6^2 - \frac{625000000}{9}t_6^3 \end{pmatrix}$$

The morphism f is injective and if $(\psi_2, \psi_6, \psi_8, \psi_{10}, \psi_{12}, \psi_{14}, \psi_{18}) \in \pi'(H_{\alpha_1} \cap H_{\alpha_2} \cap H_{\alpha_3} \cap H_{\alpha_5} \cap H_{\alpha_7})$, then by setting

$$\begin{cases} t_2 = \psi_2, \\ t_6 = \frac{1}{18000}\psi_6 - \frac{1861}{16200}\psi_2^3, \end{cases}$$

it follows that $f(t_2, 0, t_6, 0) = (\psi_2, \psi_6, \psi_8, \psi_{10}, \psi_{12}, \psi_{14}, \psi_{18})$. Hence

$$f : (\mathfrak{h}/W)^\Omega \xrightarrow{\cong} \pi'(H_{\alpha_1} \cap H_{\alpha_2} \cap H_{\alpha_3} \cap H_{\alpha_5} \cap H_{\alpha_7}) \subset \mathfrak{h}'/W'.$$

Let ϕ' be a sub-root system in Table 14, and set $h' \in \cap_{\alpha \in \phi'} H_\alpha$. With the preceding expressions, we compute $(t_2, 0, t_6, 0) = f^{-1}(\pi'(h'))$, and verify the following correspondence:

$$\begin{array}{ll} \phi' \text{ of type } A_2 + A_1 + A_1 + A_1 & \longleftrightarrow (t_2, 0, t_6, 0) \text{ generic,} \\ \phi' \text{ of type } A_3 + A_2 + A_1 & \longleftrightarrow (t_2, 0, t_6, 0) \in \{t_6 = -\frac{t_2^3}{108} \neq 0\}, \\ \phi' \text{ of type } D_5 + A_1 & \longleftrightarrow (t_2, 0, t_6, 0) \in \{t_6 = \frac{t_2^3}{108} \neq 0\}, \\ \phi' \text{ of type } E_7 & \longleftrightarrow (t_2, 0, t_6, 0) = (0, 0, 0, 0). \end{array}$$

With Table 13, we see that the singular configuration in the fiber of $\overline{\alpha^\Omega}$ above $f^{-1}(\pi'(h'))$ is of the same type as ϕ' . Hence for $D_4 - G_2 - E_7$, the map f realises a bijection between the singular configurations of the fibers of α^Ω and the sub-root systems of E_7 containing $\Theta = \{\alpha_1, \alpha_2, \alpha_3, \alpha_5, \alpha_7\}$.

4.7. Case $E_6 - F_4 - E_7$

Set $\Gamma = \mathcal{T}$, $\Gamma' = \mathcal{O}$ and $\Omega = \mathbb{Z}/2\mathbb{Z} = \langle \sigma \rangle$. According to [4], a semiuniversal deformation α^Ω of a simple singularity of type F_4 is given by

$$\begin{aligned} X_{\Gamma, \Omega} &= \{((x, y, z), (t_2, 0, t_6, t_8, 0, t_{12})) \in \mathbb{C}^3 \times \mathfrak{h}/W \mid -\frac{1}{4}x^4 + y^3 + z^2 - \frac{t_2}{4}x^2y + \frac{1}{48}(t_6 - \frac{t_2^3}{8})x^2 \\ &\quad + \frac{1}{48}(-t_8 + \frac{t_6t_2}{4} - \frac{t_2^4}{192})y + \frac{1}{576}(t_{12} - \frac{t_8t_2^2}{8} - \frac{t_6^2}{8} + \frac{t_6t_2^3}{96}) = 0\} \\ &\quad \downarrow \alpha^\Omega \\ (\mathfrak{h}/W)^\Omega &= \{(t_2, 0, t_6, t_8, 0, t_{12}) \in \mathfrak{h}/W\} \end{aligned}$$

Furthermore, the action of Ω on a fiber of α^Ω is $\sigma.(x, y, z) = (-x, y, -z)$ and the quotient $\overline{\alpha^\Omega}$ is

$$\begin{aligned} X_{\Gamma, \Omega}/\Omega &= \{((X, Y, Z), (t_2, 0, t_6, t_8, 0, t_{12})) \in \mathbb{C}^3 \times \mathfrak{h}/W \mid -\frac{1}{4}X^3 + XY^3 + Z^2 - \frac{t_2}{4}X^2Y + \frac{1}{48}(t_6 - \frac{t_2^3}{8})X^2 \\ &\quad + \frac{1}{48}(-t_8 + \frac{t_6t_2}{4} - \frac{t_2^4}{192})XY + \frac{1}{576}(t_{12} - \frac{t_8t_2^2}{8} - \frac{t_6^2}{8} + \frac{t_6t_2^3}{96})X = 0\} \\ &\quad \downarrow \overline{\alpha^\Omega} \\ (\mathfrak{h}/W)^\Omega &= \{(t_2, 0, t_6, t_8, 0, t_{12}) \in \mathfrak{h}/W\} \end{aligned}$$

Before quotient, the discriminant of α^Ω is $\mathcal{H}_1 \cup \mathcal{H}_2$ with

- $\mathcal{H}_1 = \{t_2^{12} - 144t_2^9t_6 + 576t_2^8t_8 + 5184t_2^6t_6^2 - 13824t_2^5t_6t_8 - 138240t_2^4t_8^2 - 69120t_2^3t_6^3 - 331776t_{12}t_2^3t_6 + 829440t_2^2t_6^2t_8 + 3981312t_{12}t_2^2t_8 - 5308416t_2t_6t_8^2 - 248832t_6^4 + 3981312t_{12}t_6^2 + 7077888t_8^3 - 15925248t_{12}^2 = 0\},$
- $\mathcal{H}_2 = \{t_2^{12} + 144t_2^9t_6 + 576t_2^8t_8 + 5184t_2^6t_6^2 + 13824t_2^5t_6t_8 - 138240t_2^4t_8^2 + 69120t_2^3t_6^3 - 331776t_{12}t_2^3t_6 + 829440t_2^2t_6^2t_8 - 3981312t_{12}t_2^2t_8 + 5308416t_2t_6t_8^2 - 248832t_6^4 - 3981312t_{12}t_6^2 + 7077888t_8^3 - 15925248t_{12}^2 = 0 \text{ and } t_2 \neq 0\}.$

After quotient, it is known from Theorem 2.3 that all of $(\mathfrak{h}/W)^\Omega$ composes the discriminant. By studying explicitly the singularities in the fibers $(\alpha^\Omega)^{-1}(t_2, 0, t_6, t_8, 0, t_{12})$ and $(\overline{\alpha^\Omega})^{-1}(t_2, 0, t_6, t_8, 0, t_{12})$, we obtain the following types:

$(t_2, 0, t_6, t_8, 0, t_{12})$	Configuration of $(\alpha^\Omega)^{-1}(t_2, 0, t_6, t_8, 0, t_{12})$	$(\overline{\alpha^\Omega})^{-1}(t_2, 0, t_6, t_8, 0, t_{12})$
Generic point	$p_1 + p_2 + p_3$	$A_1 + A_1 + A_1$
$\mathcal{H}_1$	$A_1 + p$	$A_3 + A_1$
$\mathcal{H}_1 \cap \{t_8 = \frac{1}{96} \frac{t_2^6 + 96t_6^2}{t_2^2}\}$	$A_3 + p$	$D_4 + A_1$
$\mathcal{H}_1 \cap \{t_8 = \frac{1}{96} \frac{t_2^6 + 96t_6^2}{t_2^2}\} \cap \{t_2^3 + 8t_6 = 0\}$	$A_5 + p$	$D_5 + A_1$
$\mathcal{H}_1 \cap \{t_8 = \frac{1}{96} \frac{t_2^6 + 96t_6^2}{t_2^2}\} \cap \{t_2^3 - 8t_6 = 0\}$	D_4	D_6
$\mathcal{H}_1 \cap \{t_8 = -\frac{1}{192} t_2^4 + \frac{1}{4} t_2 t_6\}$	A_2	A_5
$\mathcal{H}_2 \setminus \mathcal{H}_1$	$\begin{array}{c} A_1 \quad + \quad A_1 + p_1 + p_2 + p_3 \\ \curvearrowright \\ \Omega \end{array}$	$A_1 + A_1 + A_1 + A_1$
$\mathcal{H}_2 \cap \{t_8 = -\frac{1}{192} t_2^4 - \frac{1}{4} t_2 t_6\}$	$\begin{array}{c} A_2 \quad + \quad A_2 + p_1 + p_2 + p_3 \\ \curvearrowright \\ \Omega \end{array}$	$A_2 + A_1 + A_1 + A_1$
$\mathcal{H}_1 \cap \mathcal{H}_2$	$\begin{array}{c} A_1 \quad + \quad A_1 + A_1 + p \\ \curvearrowright \\ \Omega \end{array}$	$A_3 + A_1 + A_1$
$\mathcal{H}_1 \cap \mathcal{H}_2 \cap \{t_8 = -\frac{1}{192} t_2^4 - \frac{1}{4} t_2 t_6\}$	$\begin{array}{c} A_2 \quad + \quad A_2 + A_1 + p \\ \curvearrowright \\ \Omega \end{array}$	$A_3 + A_2 + A_1$
$\mathcal{H}_1 \cap \mathcal{H}_2 \cap \{t_8 = -\frac{1}{192} t_2^4 + \frac{1}{4} t_2 t_6\}$	$\begin{array}{c} A_2 \quad + \quad A_1 \quad + \quad A_1 \\ \quad \quad \quad \curvearrowright \\ \quad \quad \quad \Omega \end{array}$	$A_5 + A_1$
$t_2 = t_6 = t_8 = t_{12} = 0$	E_6	E_7

Table 15: Configurations of the fibers of α^Ω and $\overline{\alpha^\Omega}$ for $\Delta(\Gamma) = E_6$

with p_i ($i = 1, 2, 3, \emptyset$) denoting a smooth point, and if the action of Ω is not mentioned, the point (smooth or singular) is fixed.

Let $\Pi' = \{\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6, \alpha_7\}$ be a set of simple roots of the root system of type E_7 . All the sub-root systems ϕ' of E_7 containing $\Theta = \{\alpha_2, \alpha_5, \alpha_7\}$, as well as their realisations are given in the following table (except in the first case $A_1 + A_1 + A_1$ and the last one E_7 , we will not repeat that ϕ' contains Θ , for the sake of readability):

Type of ϕ'	Realisations of ϕ'
$A_1 + A_1 + A_1$	$< \alpha_2, \alpha_5, \alpha_7 >$
$A_3 + A_1$	$< \alpha_4 >, < \alpha_3 + \alpha_4 >, < \alpha_1 + \alpha_3 + \alpha_4 >, < \alpha_1 + \alpha_2 + 2\alpha_3 + 3\alpha_4 + 2\alpha_5 + 2\alpha_6 + \alpha_7 >, < \alpha_4 + \alpha_5 + \alpha_6 >, \\ < \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6 >, < \alpha_1 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6 >, < \alpha_1 + \alpha_2 + 2\alpha_3 + 3\alpha_4 + 2\alpha_5 + \alpha_6 >, < \alpha_6 >, \\ < \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6 >, < \alpha_1 + \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6 >, < \alpha_1 + \alpha_2 + 2\alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6 >$

$A_5 + A_1$	$\langle \alpha_4, \alpha_6, \alpha_1 \rangle, \langle \alpha_4, \alpha_6, \alpha_1 + 2\alpha_2 + 3\alpha_3 + 4\alpha_4 + 3\alpha_5 + 2\alpha_6 + \alpha_7 \rangle,$ $\langle \alpha_4, \alpha_6, 2\alpha_1 + 2\alpha_2 + 3\alpha_3 + 4\alpha_4 + 3\alpha_5 + 2\alpha_6 + \alpha_7 \rangle, \langle \alpha_4, \alpha_1 + \alpha_2 + 2\alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6, \alpha_1 \rangle,$ $\langle \alpha_4, \alpha_1 + \alpha_2 + 2\alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6, \alpha_2 + \alpha_3 + 2\alpha_4 + 2\alpha_5 + 2\alpha_6 + \alpha_7 \rangle,$ $\langle \alpha_4, \alpha_1 + \alpha_2 + 2\alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6, \alpha_1 + \alpha_2 + \alpha_3 + 2\alpha_4 + 2\alpha_5 + 2\alpha_6 + \alpha_7 \rangle, \langle \alpha_3 + \alpha_4, \alpha_6, \alpha_1 + \alpha_3 \rangle,$ $\langle \alpha_3 + \alpha_4, \alpha_6, \alpha_1 + 2\alpha_2 + 2\alpha_3 + 4\alpha_4 + 3\alpha_5 + 2\alpha_6 + \alpha_7 \rangle,$ $\langle \alpha_3 + \alpha_4, \alpha_6, 2\alpha_1 + 2\alpha_2 + 3\alpha_3 + 4\alpha_4 + 3\alpha_5 + 2\alpha_6 + \alpha_7 \rangle, \langle \alpha_3 + \alpha_4, \alpha_1 + \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6, \alpha_1 + \alpha_3 \rangle,$ $\langle \alpha_3 + \alpha_4, \alpha_1 + \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6, \alpha_2 + \alpha_3 + 2\alpha_4 + 2\alpha_5 + 2\alpha_6 + \alpha_7 \rangle,$ $\langle \alpha_3 + \alpha_4, \alpha_1 + \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6, \alpha_1 + \alpha_2 + 2\alpha_3 + 2\alpha_4 + 2\alpha_5 + 2\alpha_6 + \alpha_7 \rangle,$ $\langle \alpha_1 + \alpha_3 + \alpha_4, \alpha_6, \alpha_3 \rangle, \langle \alpha_1 + \alpha_3 + \alpha_4, \alpha_6, \alpha_1 + 2\alpha_2 + 2\alpha_3 + 4\alpha_4 + 3\alpha_5 + 2\alpha_6 + \alpha_7 \rangle,$ $\langle \alpha_1 + \alpha_3 + \alpha_4, \alpha_6, \alpha_1 + 2\alpha_2 + 3\alpha_3 + 4\alpha_4 + 3\alpha_5 + 2\alpha_6 + \alpha_7 \rangle, \langle \alpha_1 + \alpha_3 + \alpha_4, \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6, \alpha_3 \rangle,$ $\langle \alpha_1 + \alpha_3 + \alpha_4, \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6, \alpha_1 + \alpha_2 + \alpha_3 + 2\alpha_4 + 2\alpha_5 + 2\alpha_6 + \alpha_7 \rangle,$ $\langle \alpha_1 + \alpha_3 + \alpha_4, \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6, \alpha_1 + \alpha_2 + 2\alpha_3 + 2\alpha_4 + 2\alpha_5 + 2\alpha_6 + \alpha_7 \rangle,$ $\langle \alpha_4 + \alpha_5 + \alpha_6, \alpha_1 + \alpha_2 + 2\alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6, \alpha_1 \rangle,$ $\langle \alpha_4 + \alpha_5 + \alpha_6, \alpha_1 + \alpha_2 + 2\alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6, \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 \rangle,$ $\langle \alpha_4 + \alpha_5 + \alpha_6, \alpha_1 + \alpha_2 + 2\alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6, \alpha_1 + \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 \rangle,$ $\langle \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6, \alpha_1 + \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6, \alpha_1 + \alpha_3 \rangle,$ $\langle \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6, \alpha_1 + \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6, \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 \rangle,$ $\langle \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6, \alpha_1 + \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6, \alpha_1 + \alpha_2 + 2\alpha_3 + 2\alpha_4 + \alpha_5 \rangle,$ $\langle \alpha_1 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6, \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6, \alpha_1 + \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 \rangle,$ $\langle \alpha_1 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6, \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6, \alpha_1 + \alpha_2 + 2\alpha_3 + 2\alpha_4 + \alpha_5 \rangle,$ $\langle \alpha_1 + \alpha_2 + 2\alpha_3 + 3\alpha_4 + 2\alpha_5 + \alpha_6, \alpha_6, \alpha_1 \rangle, \langle \alpha_1 + \alpha_2 + 2\alpha_3 + 3\alpha_4 + 2\alpha_5 + \alpha_6, \alpha_6, \alpha_3 \rangle,$ $\langle \alpha_1 + \alpha_2 + 2\alpha_3 + 3\alpha_4 + 2\alpha_5 + \alpha_6, \alpha_6, \alpha_1 + \alpha_3 \rangle,$ $\langle \alpha_4, \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6, \alpha_1 + \alpha_2 + 2\alpha_3 + 2\alpha_4 + \alpha_5 \rangle,$ $\langle \alpha_4, \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6, \alpha_1 + \alpha_2 + \alpha_3 + 2\alpha_4 + 2\alpha_5 + 2\alpha_6 + \alpha_7 \rangle,$ $\langle \alpha_4, \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6, 2\alpha_1 + 2\alpha_2 + 3\alpha_3 + 4\alpha_4 + 3\alpha_5 + 2\alpha_6 + \alpha_7 \rangle,$ $\langle \alpha_4, \alpha_1 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6, \alpha_1 + \alpha_2 + 2\alpha_3 + 2\alpha_4 + \alpha_5 \rangle,$ $\langle \alpha_4, \alpha_1 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6, \alpha_2 + \alpha_3 + 2\alpha_4 + 2\alpha_5 + 2\alpha_6 + \alpha_7 \rangle,$ $\langle \alpha_4, \alpha_1 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6, \alpha_1 + 2\alpha_2 + 3\alpha_3 + 4\alpha_4 + 3\alpha_5 + 2\alpha_6 + \alpha_7 \rangle,$ $\langle \alpha_3 + \alpha_4, \alpha_4 + \alpha_5 + \alpha_6, \alpha_1 + \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 \rangle,$ $\langle \alpha_3 + \alpha_4, \alpha_4 + \alpha_5 + \alpha_6, \alpha_1 + \alpha_2 + 2\alpha_3 + 2\alpha_4 + 2\alpha_5 + 2\alpha_6 + \alpha_7 \rangle,$ $\langle \alpha_3 + \alpha_4, \alpha_4 + \alpha_5 + \alpha_6, 2\alpha_1 + 2\alpha_2 + 3\alpha_3 + 4\alpha_4 + 3\alpha_5 + 2\alpha_6 + \alpha_7 \rangle,$ $\langle \alpha_3 + \alpha_4, \alpha_1 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6, \alpha_1 + \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 \rangle,$ $\langle \alpha_3 + \alpha_4, \alpha_1 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6, \alpha_2 + \alpha_3 + 2\alpha_4 + 2\alpha_5 + 2\alpha_6 + \alpha_7 \rangle,$ $\langle \alpha_3 + \alpha_4, \alpha_1 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6, \alpha_1 + 2\alpha_2 + 2\alpha_3 + 4\alpha_4 + 3\alpha_5 + 2\alpha_6 + \alpha_7 \rangle,$ $\langle \alpha_1 + \alpha_3 + \alpha_4, \alpha_4 + \alpha_5 + \alpha_6, \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 \rangle,$ $\langle \alpha_1 + \alpha_3 + \alpha_4, \alpha_4 + \alpha_5 + \alpha_6, \alpha_1 + \alpha_2 + 2\alpha_3 + 2\alpha_4 + 2\alpha_5 + 2\alpha_6 + \alpha_7 \rangle,$ $\langle \alpha_1 + \alpha_3 + \alpha_4, \alpha_4 + \alpha_5 + \alpha_6, \alpha_1 + 2\alpha_2 + 3\alpha_3 + 4\alpha_4 + 3\alpha_5 + 2\alpha_6 + \alpha_7 \rangle,$ $\langle \alpha_1 + \alpha_3 + \alpha_4, \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6, \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 \rangle,$ $\langle \alpha_1 + \alpha_3 + \alpha_4, \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6, \alpha_1 + \alpha_2 + \alpha_3 + 2\alpha_4 + 2\alpha_5 + 2\alpha_6 + \alpha_7 \rangle,$ $\langle \alpha_1 + \alpha_3 + \alpha_4, \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6, \alpha_1 + 2\alpha_2 + 2\alpha_3 + 4\alpha_4 + 3\alpha_5 + 2\alpha_6 + \alpha_7 \rangle$
E_7	$\langle \alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6, \alpha_7 \rangle$

Table 16: Realisations of the sub-root systems containing Θ for $\Delta(\Gamma) = E_6$

The description of E_7 will be the same as the one in Section 4.6. Therefore $h' \in H_{\alpha_2} \cap H_{\alpha_5} \cap H_{\alpha_7}$ if and only if $\xi_2 = -\xi_1$, $\xi_4 = \xi_3$ and $\xi_6 = \xi_5$.

Using [1] and [10], one can compute the flat coordinates of E_7 restricted to the subspace $H_{\alpha_2} \cap H_{\alpha_5} \cap H_{\alpha_7} \subset \mathfrak{h}'$, i.e. as functions of ξ_1, ξ_3, ξ_5 and ξ_7 . We will not give these expressions because of their extensive size. For example, ψ_{18} contains 220 terms.

It can be verified that

$$\left\{ \begin{array}{lcl} \psi_{10} & = & \frac{82928}{45}\psi_2^5 - \frac{4}{3}\psi_2^2\psi_6 + \frac{9}{35}\psi_2\psi_8, \\ \psi_{14} & = & -\frac{12190772504}{2219805}\psi_2^7 + \frac{7281979}{1522152}\psi_2^4\psi_6 - \frac{471547}{789264}\psi_2^3\psi_8 + \frac{3779}{6765120}\psi_2\psi_6^2 \\ & & + \frac{73490}{8613}\psi_{12}\psi_2 - \frac{1}{25920}\psi_6\psi_8, \\ \psi_{18} & = & -\frac{1271044247268145576}{94705443405}\psi_2^9 + \frac{256749355304}{25982289}\psi_2^6\psi_6 - \frac{12475637391961}{9093801150}\psi_2^5\psi_8 \\ & & + \frac{6360724111}{3117874680}\psi_2^3\psi_6^2 + \frac{470160383920}{31756131}\psi_{12}\psi_2^3 - \frac{5935967}{41810580}\psi_2^2\psi_6\psi_8 \\ & & - \frac{101699}{30970800}\psi_2\psi_8^2 - \frac{1}{52488}\psi_6^3 - \frac{20}{27}\psi_{12}\psi_6. \end{array} \right.$$

Define

$$f: (\mathfrak{h}/W)^\Omega \rightarrow \mathfrak{h}'/W' \left(\begin{array}{c} t_2 \\ 0 \\ t_6 \\ t_8 \\ 0 \\ t_{12} \end{array} \right) \mapsto \left(\begin{array}{c} t_2 \\ \frac{16735}{9}t_2^3 - 3000t_6 \\ \frac{4884005}{972}t_2^4 - \frac{360500}{9}t_2t_6 + 50000t_8 \\ \frac{13113}{20}t_2^5 - 6300t_2^2t_6 + \frac{90000}{7}t_2t_8 \\ -\frac{1533855367}{5248800}t_2^6 - \frac{5865805}{7776}t_2^3t_6 + \frac{2927375}{504}t_2^2t_8 - \frac{34375}{72}t_6^2 - 3125t_{12} \\ -\frac{4794135161101}{9205975296}t_2^7 + \frac{2226935425}{6088608}t_2^4t_6 + \frac{4765011875}{295974}t_2^3t_8 - \frac{103796875}{28188}t_2^2t_6^2 - \frac{229656250}{8613}t_2t_{12} + \frac{156250}{27}t_6t_8 \\ \frac{53845033157553005}{8418261636}t_2^6t_6 - \frac{2973773239515500}{545628069}t_2^5t_8 - \frac{2028480753289375}{155893734}t_2^3t_6^2 - \frac{3997692756437500}{95268393}t_2^3t_{12} - \frac{635618750000}{77427}t_2t_8^2 \\ -\frac{62500000}{9}t_{12}t_6 + \frac{32998989562500}{696843}t_2^2t_6t_8 - \frac{74647399081995101197}{218201341605120}t_2^9 - \frac{132812500}{243}t_6^3 \end{array} \right)$$

The morphism f is injective and if $(\psi_2, \psi_6, \psi_8, \psi_{10}, \psi_{12}, \psi_{14}, \psi_{18}) \in \pi'(H_{\alpha_2} \cap H_{\alpha_5} \cap H_{\alpha_7})$, then by setting

$$\left\{ \begin{array}{l} t_2 = \psi_2, \\ t_6 = \frac{3347}{5400}\psi_2^3 - \frac{1}{3000}\psi_6, \\ t_8 = \frac{1}{50000}\psi_8 + \frac{1283191}{3240000}\psi_2^4 - \frac{721}{2700000}\psi_2\psi_6, \\ t_{12} = \frac{18995770783}{43740000000}\psi_2^6 - \frac{342859}{972000000}\psi_2^3\psi_6 + \frac{23419}{630000000}\psi_2^2\psi_8 - \frac{11}{648000000}\psi_6^2 - \frac{1}{3125}\psi_{12}, \end{array} \right.$$

it follows that $f(t_2, 0, t_6, t_8, 0, t_{12}) = (\psi_2, \psi_6, \psi_8, \psi_{10}, \psi_{12}, \psi_{14}, \psi_{18})$. Hence

$$f: (\mathfrak{h}/W)^\Omega \xrightarrow{\cong} \pi'(H_{\alpha_2} \cap H_{\alpha_5} \cap H_{\alpha_7}) \subset \mathfrak{h}'/W'.$$

Let ϕ' be a sub-root system in Table 16, and set $h' \in \bigcap_{\alpha \in \phi'} H_\alpha$. With the preceding expressions, we compute $(t_2, 0, t_6, t_8, 0, t_{12}) = f^{-1}(\pi'(h'))$, and verify the following correspondence:

ϕ' of type $A_1 + A_1 + A_1$	$\longleftrightarrow$	(t_2, t_6, t_8, t_{12}) generic,
ϕ' of type $A_3 + A_1$	$\longleftrightarrow$	$(t_2, t_6, t_8, t_{12}) \in \mathcal{H}_1$,
ϕ' of type $D_4 + A_1$	$\longleftrightarrow$	$(t_2, t_6, t_8, t_{12}) \in \mathcal{H}_1 \cap \{t_8 = \frac{1}{96} \frac{t_2^5 + 96t_6^2}{t_2^2}\}$,
ϕ' of type $D_5 + A_1$	$\longleftrightarrow$	$(t_2, t_6, t_8, t_{12}) \in \mathcal{H}_1 \cap \{t_8 = \frac{1}{96} \frac{t_2^5 + 96t_6^2}{t_2^2}\} \cap \{t_2^3 + 8t_6 = 0\}$,
ϕ' of type D_6	$\longleftrightarrow$	$(t_2, t_6, t_8, t_{12}) \in \mathcal{H}_1 \cap \{t_8 = \frac{1}{96} \frac{t_2^5 + 96t_6^2}{t_2^2}\} \cap \{t_2^3 - 8t_6 = 0\}$,
ϕ' of type A_5	$\longleftrightarrow$	$(t_2, t_6, t_8, t_{12}) \in \mathcal{H}_1 \cap \{t_8 = -\frac{1}{192}t_2^4 + \frac{1}{4}t_2t_6\}$,
ϕ' of type $A_1 + A_1 + A_1 + A_1$	$\longleftrightarrow$	$(t_2, t_6, t_8, t_{12}) \in \mathcal{H}_2 \setminus \mathcal{H}_1$,

ϕ' of type $A_2 + A_1 + A_1 + A_1$	$\longleftrightarrow$	$(t_2, t_6, t_8, t_{12}) \in \mathcal{H}_2 \cap \{t_8 = -\frac{1}{192}t_2^4 - \frac{1}{4}t_2t_6\},$
ϕ' of type $A_3 + A_1 + A_1$	$\longleftrightarrow$	$(t_2, t_6, t_8, t_{12}) \in \mathcal{H}_1 \cap \mathcal{H}_2,$
ϕ' of type $A_3 + A_2 + A_1$	$\longleftrightarrow$	$(t_2, t_6, t_8, t_{12}) \in \mathcal{H}_1 \cap \mathcal{H}_2 \cap \{t_8 = -\frac{1}{192}t_2^4 - \frac{1}{4}t_2t_6\},$
ϕ' of type $A_5 + A_1$	$\longleftrightarrow$	$(t_2, t_6, t_8, t_{12}) \in \mathcal{H}_1 \cap \mathcal{H}_2 \cap \{t_8 = -\frac{1}{192}t_2^4 + \frac{1}{4}t_2t_6\},$
ϕ' of type E_7	$\longleftrightarrow$	$(t_2, t_6, t_8, t_{12}) = (0, 0, 0, 0).$

With Table 15, we see that the singular configuration in the fiber of $\overline{\alpha^\Omega}$ above $f^{-1}(\pi'(h'))$ is of the same type as ϕ' . Hence for $E_6 - F_4 - E_7$, the map f realises a bijection between the singular configurations of the fibers of $\overline{\alpha^\Omega}$ and the sub-root systems of E_7 containing $\Theta = \{\alpha_2, \alpha_5, \alpha_7\}$.

Conclusion

Conjecture 3.1 and the subsequent proof (Theorem 3.2) for small rank cases give a direct link between the singular configurations of the fibers of the deformation $\overline{\alpha^\Omega}$ of a simple singularity of type $\Delta(\Gamma')$ and sub-root systems of a root system of type $\Delta(\Gamma')$. As every fiber of $\overline{\alpha^\Omega}$ is singular (Theorem 2.3) and as the Dynkin diagram associated to the singular configuration of such a fiber has to contain the subset Θ of simple roots of a root system of type $\Delta(\Gamma')$, this subset Θ characterises how far the morphism $\overline{\alpha^\Omega}$ is from being isomorphic to the semiuniversal deformation $\chi'_{|S, e'}$ of the simple singularity of type $\Delta(\Gamma')$.

The full proof of Conjecture 3.1 requires another approach. Indeed the proof presented in Section 4 is based on direct computations of the singular configurations of the fibers of α^Ω and of $\overline{\alpha^\Omega}$. But these computations themselves stem from the equation of the deformation α^Ω , which is not known when α^Ω is the semiuniversal deformation of a simple singularity of type C_r , $r \geq 6$ (i.e. $\Gamma = \mathcal{D}_{r-1}$ and $\Gamma' = \mathcal{D}_{2(r-1)}$). This is due to the fact that in order to obtain α^Ω , we need to determine the independent cycles of $M(\Gamma)$ (cf. Section 1.2), and as the dimension vector of $M(\Gamma)$ is $(1, 1, 2, \dots, 2, 1, 1)$, the 2-dimensional vertices introduce a complexity in the relations between the cycles, complicating the computation of α^Ω when the rank r increases (cf. [4] Section 4.3.7 for details). The author is therefore working on a more general approach which would provide a proof of Conjecture 3.1 in all generality.

Acknowledgements

The author would like to thank Michaël Bulois for his helpful comments and remarks that helped improve the form of the present article, as well as Kenji Iohara for the constancy of his support and guidance.

References

- [1] Abriani, D., 2009. Frobenius manifolds associated to Coxeter groups of type E_7 and E_8 , Version 1, submitted on Arxiv, pp. 1–13. <https://arxiv.org/abs/0910.5453>.
- [2] Bourbaki, N., 1968. Éléments de mathématique. Fasc. XXXIV. Groupes et algèbres de Lie. Chapitre IV: Groupes de Coxeter et systèmes de Tits. Chapitre V: Groupes engendrés par des réflexions. Chapitre VI: systèmes de racines. Actualités Scientifiques et Industrielles, No. 1337, Hermann, Paris.
- [3] Brieskorn, E., 1971. Singular elements of semi-simple algebraic groups, in: Actes du Congrès International des Mathématiciens (Nice, 1970), Tome 2. Gauthier-Villars, Paris, pp. 279–284.
- [4] Caradot, A., 2017. Singularity and Lie Theory, Ph.D Thesis. Université de Lyon, pp. 1–279. <https://hal.archives-ouvertes.fr/tel-01556773v2>.
- [5] Cassens, H., Slodowy, P., 1998. On Kleinian singularities and quivers, in: Singularities (Oberwolfach, 1996). Birkhäuser, Basel. volume 162 of *Progr. Math.*, pp. 263–288. doi:10.1007/978-3-0348-8770-0_14.
- [6] Du Val, P., 1934. On isolated singularities of surfaces which do not affect the conditions of adjunction (part I, II, III). Mathematical Proceedings of the Cambridge Philosophical Society 04, 453–465. doi:10.1017/S030500410001269X.

- [7] Frame, J.S., 1951. The classes and representations of the groups of 27 lines and 28 bitangents. *Ann. Mat. Pura Appl.* (4) 32, 83–119. doi:10.1007/BF02417955.
- [8] Kas, A., Schlessinger, M., 1972. On the versal deformation of a complex space with an isolated singularity. *Math. Ann.* 196, 23–29. doi:10.1007/BF01419428.
- [9] Klein, F., 1884. *Vorlesungen über das Ikosaeder und die Auflösung der Gleichungen vom fünften Grade.* Teubner. URL: <http://eudml.org/doc/203220>.
- [10] Mehta, M.L., 1988. Basic sets of invariant polynomials for finite reflection groups. *Comm. Algebra* 16, 1083–1098. doi:10.1080/00927878808823619.
- [11] Saito, K., 1979. On the periods of primitive integrals, Harvard University.
- [12] Saito, K., 1983. Period mapping associated to a primitive form. *Publ. Res. Inst. Math. Sci.* 19, 1231–1264. doi:10.2977/prims/1195182028.
- [13] Saito, K., Yano, T., Sekiguchi, J., 1980. On a certain generator system of the ring of invariants of a finite reflection group. *Comm. Algebra* 8, 373–408. doi:10.1080/00927878008822464.
- [14] Slodowy, P., 1980a. Four lectures on simple groups and singularities. volume 11 of *Communications of the Mathematical Institute, Rijksuniversiteit Utrecht*. Rijksuniversiteit Utrecht, Mathematical Institute, Utrecht.
- [15] Slodowy, P., 1980b. Simple singularities and simple algebraic groups. volume 815 of *Lecture Notes in Mathematics*. Springer, Berlin.