Phytolith-rich biochar increases cotton biomass and silicon-mineralomass in a highly weathered soil

Zimin Li, Bruno Li, Johan Yans, Nicolas Dufour, David Houben,
Jean-Thomas Cornélis

► To cite this version:

Zimin Li, Bruno Li, Johan Yans, Nicolas Dufour, David Houben, et al.. Phytolith-rich biochar increases cotton biomass and silicon-mineralomass in a highly weathered soil. *Journal of Plant Nutrition and Soil Science*, 2018, 181 (4), pp.537-546. 10.1002/jpln.201800031 . hal-02176368

HAL Id: hal-02176368

<https://hal.science/hal-02176368>

Submitted on 8 Jul 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Phytolith-rich biochar increases cotton biomass and silicon-mineralomass in a highly weathered soil

Zimin Li^{1*}, Bruno Delvaux¹, Johan Yans², Nicolas Dufour¹, David Houben³, Jean-Thomas Cornelis⁴

¹Université catholique de Louvain, Earth and Life Institute, soil science, 1348 Louvain-la-Neuve, Belgium

²Department of Geology, NaGRIDD, University of Namur, 5000 Namur, Belgium

³HydrISE, Institut Polytechnique LaSalle Beauvais, rue Pierre Waguet 19, 60026 Beauvais Cedex, France

⁴BIOSE department, Gembloux Agro-Bio Tech, University of Liege, 5030 Gembloux, Belgium

***Corresponding author:** zimin.li@uclouvain.be (Zimin LI)

Address: Université catholique de Louvain, Earth and Life Institute, soil science, 1348 Louvain-la-Neuve, Belgium, Croix du Sud 2/L7.05.10,1348 Louvain-la-Neuve, Belgium

Resubmitted to the journal of *Journal of Plant Nutrition and Soil Science* (jpln.201700236)

Tables: **4**

Figures: **4**

Pages: **24**

Abstract

Non-essential silicon (Si) is beneficial to plants. It increases the biomass of Si-accumulator plants by improving photosynthetic activity and alleviating stresses. Desilication, however, takes place because of natural soil weathering and removal of harvested biomass. Pyrolysis transforms Si-rich biomass into biochar that can be used to supply bioavailable Si.

Here, we applied two biochar materials differing in Si content on soils differing in weathering stage: a young Cambisol and a highly weathered Nitisol. We studied the impact of biochar supply on the bioavailability of Si, cotton biomass and Si mineralomass. The biochar materials derived from, respectively: *Miscanthus x giganteus* –Mi– (34.6 g Si kg⁻¹ in biochar) and soft woody material –SW– (0.9 g Si kg⁻¹ in biochar). They were compared to conventional Si fertilizer wollastonite –Wo– (CaSiO₃). Amendments were incorporated in soils at the rate of 3% (w/w). The content of bioavailable Si in soil was determined through 0.01 M CaCl₂ extraction.

In the Cambisol, the proportion [CaCl₂ extractable Si: total Si content] was significantly smaller for Mi (0.9%) than for Wo (5.2%). In the Nitisol, this proportion was much larger for Mi (1.4%) than for Wo (0.7%). Mi-biochar significantly increased Si-mineralomass relatively to SW-biochar in both soils. This increase was, however, much larger in the Nitisol (5.9-fold) than in the Cambisol (2.2-fold).

Mi biochar is thus an alternative Si fertilizer to Wo to supply bioavailable Si, increase plant biomass, and promote the biological cycle of Si in the soil-plant system in the Nitisol. Besides, it increased soil fertility and soil organic carbon content.

Keywords biochar, bioavailable silicon, soil desilication, cotton

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Introduction

Silicon (Si) is ubiquitous in rocks and soils. Si is not essential, but beneficial to plants. It increases photosynthetic activity and plant resistance against various biotic and abiotic stresses (*Belanger, 1995; Fauteux et al., 2005; Liang et al., 2007*). Aqueous monosilicic acid (H_4SiO_4^0) is ubiquitous in soil solution where plant roots take it up. In plants, mass flow transports H_4SiO_4^0 upwards to transpiration sites. Therein, water loss increases H_4SiO_4^0 concentration, and induces polymerization through Si-O-Si bounds, and further precipitation of hydrated amorphous silica as phytoliths (amorphous $\text{SiO}_2 \cdot n\text{H}_2\text{O}$) (*Epstein, 1994; Ma et al., 2006*). Phytoliths (PhSi) are major biogenic silicate minerals (BSi). They return into the topsoil with dead plant materials, and further contribute to the pool of dissolved Si (DSi) in soil (*Keller et al., 2012*) because of their solubility (*Frayssé et al., 2009*). They thus supply the pool of bioavailable Si in soils under forests (*Cornelis et al., 2011; Gérard et al., 2008*), grasslands (*Alexandre et al., 2011; Blecker et al., 2006; Issaharou-Matchi et al., 2016; White et al., 2012*) and croplands either in temperate conditions (*Vandevenne et al., 2015*) or in the humid tropics (*Henriet et al., 2008*). The contribution of PhSi to DSi pool can be substantially impacted by land use (*Barão et al., 2014; Struyf et al., 2010*), harvest exportation (*Keller et al., 2012; Vandevenne et al., 2015; Vandevenne et al., 2012*), soil weathering stage (*Henriet et al., 2008*) and, overall, soil processes (*Cornelis and Delvaux, 2016*).

The regular removal of straws of high-Si accumulator plants combined with natural desilication in freely drained soils leads to a substantial decrease of PhSi and DSi pools in agroecosystems (*Keller et al., 2012; Struyf et al., 2010; Vandevenne et al., 2015*). Si supply may therefore be required to sustain continuous cropping in highly desilicated soils depleted in weatherable silicate minerals. The use of traditional amendments raises cost and environmental problems (*Berthelsen et al., 2001; Haynes, 2014; Haynes et al., 2013; Datnoff and Heckman, 2014*), prompting us to develop other ways to improve the use of available biomass.

Biochar is a black carbon used as a soil amendment. It is produced by heating organic matter under

oxygen-limited conditions. Pyrolysis converts organic substances into stable C compounds, hence preserves atmospheric C (Lehmann and Joseph, 2015; Lehmann et al., 2011). Applied to soil, biochar retains and supplies plant nutrients (Glaser et al., 2002; Laird et al., 2010; Lehmann et al., 2003), increases soil C stock (Lehmann and Joseph, 2015), soil pH and plant growth (Biederman and Harpole, 2013). Black carbon significantly increases soil cation exchange capacity (CEC) through its large surface charge density and surface area (Liang et al., 2006). Yet the effect of biochar on plant growth largely depends on soil properties and biochar types (Biederman and Harpole, 2013; Crane-Droesch et al., 2013; Jeffery et al., 2011).

Pyrolysis of plant residues concentrates PhSi minerals (Houben et al., 2014) which in turn contribute to supply DSi (Houben et al., 2014; Xiao et al., 2014), and increase Si uptake by plants (Liu et al., 2014). The impact of both the PhSi content of biochar and soil silicon availability on plant growth and Si uptake is, however, unknown. Here, we used two biochars differing in Si content, and apply them on soils differing in weathering stage, respectively, a young Cambisol and an old, highly weathered, Nitisol. We studied the release of DSi after biochar application. We further investigated the effect of biochar on soil fertility, plant biomass and Si mineralomass, and compared it to the one of CaSiO₃-wollastonite.

Materials and methods

Biochar, wollastonite and soils

Biochars and wollastonite. The biochars were produced by industrial pyrolysis (550°C, heating rate 65 °C/min, kiln residence time 12 min) at UK Biochar Research Centre (UKBRC), respectively from *Miscanthus* (*Miscanthus x giganteus*) –Mi– and soft wood (Mixed softwood pellets) –SW– (Table 1). The wollastonite –Wo– was provided by R.T Vanderbilt Company, Inc. (Norwalk, CT, USA) (Table 1). In the following text, the biochars and wollastonite are considered as soil amendments.

Soils and soil:amendment mixtures. The two soils key out as, respectively, a Dystric Cambisol (CA) and a

1
2
3 83 Rhodic Nitisol (NI) in the WRB system (IUSS, 2014). CA (Forest of Lauzelle, Louvain-la Neuve,
4
5 84 Belgium) developed from quaternary calcareous silty loess under humid temperate conditions (Titeux and
6
7 85 Delvaux, 2009). NI is a paleosol from the quarry of Transinne (Belgium), formed under humid tropical
8
9 86 paleo-conditions. The quarry exhibits a deep 65m saprolithic profile derived from Early Devonian
10
11 87 bedrock, containing kaolinite, resistant muscovite and quartz, and Fe oxyhydroxides (Thiry et al., 2006).
12
13 88 The soil samples were air-dried and sieved at 2-mm. The soil:amendment mixtures were made at 100:3
14
15 89 g:g (dry soil weight), and are named CA-Mi, CA-SW, CA-Wo, NI-Mi, NI-SW and NI-Wo.

16
17
18
19 90 *CaCl₂ extraction of Si*

20
21
22 91 The extraction was carried out on the Si amendments, soils and soil:amendment mixtures before the cotton
23
24 92 pot experiment. The CaCl₂-extractable Si content (CaCl₂-Si) was determined using a 5g:50ml (0.01 M
25
26 93 CaCl₂) soil:solution ratio, and a 5h continuous shaking time (Haysom and Chapman, 1975). After
27
28 94 centrifuged at 3,000×g for 20 min, the supernatant was separated for analysis. The residues were
29
30 95 submitted to five successive extractions during 5 days. The concentration of Si was measured in each
31
32 96 extract by Inductively Coupled Plasma-Atomic Emission Spectrometry (ICP-AES).

33
34
35
36 97 Analyses. pH was measured in, respectively, H₂O and KCl 1M, with 5g:25ml suspensions. The CEC and
37
38 98 content of exchangeable cations were determined using CH₃COONH₄ 1M buffered at pH=7 (Chapman,
39
40 99 1965). Elemental analysis was carried out after calcination and Li-metaborate + Li-tetraborate fusion
41
42 100 (Chao and Sanzolone, 1992). The elemental contents were determined by ICP-AES. Elemental contents of
43
44 101 alkaline and alkaline-earth cations were summed up as the total reserve in bases (TRB), which estimates
45
46 102 the content of weatherable minerals (Herbillon, 1986). Total carbon (C) and nitrogen (N) contents were
47
48 103 measured by dry combustion with a CNS analyzer (Flash EA1112 Series).

49
50
51
52 104 Scanning Electron Microscopy (SEM). The analyses of biochar particles were performed on specimens
53
54 105 using a JEOL FEG SEM 7600F, equipped with Energy Dispersive X-ray Spectrometry system (EDX, Jeol

JSM2300 with a resolution < 129 eV) operating at 15 keV with a working distance of 8 mm for the element detection. The acquisition time for the chemical spectra lasted 100 s with a probe current of 1 nA.

Plant material and analyses, pot experiment

Cotton plant. We used *Gossypium hirsutum*, var (STAM 59A-Bollgard II) from southern Burkina Faso.

Pot experiment in a phytotron (5 months). The soil:amendment mixtures (100:3 g:g) were irrigated with deionized water and allowed to equilibrate for a week at 70% of field water holding capacity before planting. After germination in deionized water, the plantlets were allowed to grow up to 5-cm. They were then transplanted into pots (7.5 L) containing the respective soil:amendment mixtures (3kg dry weight), in triplicate under 28°C and relative humidity at 80%. Each pot was watered every three days with 33 ml deionized water during the first two months, and 100 ml the last three months. No nutrients were supplied. We collected the aboveground plant part after 5 months for further drying at 55°C during 7 days prior to biomass weighing.

Analyses. Mineral analysis was carried out after calcination at 450°C for 1 day and fusion in Li-metaborate + Li-tetraborate at 1,000°C (Chao and Sanzalone, 1992), followed by ash dissolution with concentrated HNO₃. Element contents were measured by ICP-AES. Element mineralomass was computed from total element contents and cotton biomass.

Statistical analyses

We used the SPSS 17.0 statistical software to carry out the one-way analysis of variance (ANOVA) (at $p<0.05$ level) and LSD (at $p<0.05$ level).

Results

Properties of the biochars and wollastonite

1
2
3 127 As shown in Table 1, the OC content (%) was 84.9 in SW, 77.1 in Mi and 0.6 in Wo. The total N content
4
5 128 (%) was 1.2 in SW and 2.0 in Mi. The contents of CaCl_2 extractable Si (g kg^{-1}) were larger in Mi (11.4)
6
7 129 than in Wo (8.8) and SW (0.01). Si content of Wo amounted to 233.3 g kg^{-1} which was 7 times larger than
8
9 130 that of Mi (34.6 g kg^{-1}), and 259 times larger than in SW (0.9 g kg^{-1}). The contents of Ca, Mg in Wo were
10
11 131 also significantly larger than in Mi and WS. The contents of K ($\text{cmol}_c \text{ kg}^{-1}$) were larger in Mi (14.9) than
12
13 132 in SW (2.6) and Wo (1.1). Figure 1 illustrated the SEM micrographs and associated EDX spectra of
14
15 133 biochar materials SW and Mi. In particular, Fig. 1(f) showed the occurrence of dumbbell-shaped, fine silt-
16
17 134 sized phytoliths in Mi biochar. As inferred from Fig.1 (e), they were abundant and undamaged after
18
19 135 pyrolysis at 550°C .

20
21
22
23 136 *Soil and soil:amendment properties*
24
25

26 137 CA and NI are acid soils since their pH_{water} values were 4.6 and 5.2, respectively (Table 2). C and N
27
28 138 contents, CEC and contents of exchangeable cations were larger in CA than in NI. Of course, biochar
29
30 139 application resulted in a huge increase in C content while Wo did not. The latter application led to a
31
32 140 marked increase in pH and content of exchangeable Ca. SW application did not generate any change of
33
34 141 pH, CEC, contents of exchangeable cations and base saturation whatever the soil type. To the contrary, Mi
35
36 142 application induced a significant increase of these parameters. In particular, exchangeable K content
37
38 143 ($\text{cmol}_c \text{ kg}^{-1}$) increased from, respectively 0.2 to 0.97 in CA, and 0.05 to 0.81 in NI. From Total contents of
39
40 144 Al and Fe (g kg^{-1}) amounted to, respectively, 39.2 and 18.5 in CA, 96.5 and 30.6 in NI (Table 3). Total Si
41
42 145 content (g kg^{-1}) in CA and NI was 376.8 and 311.9, respectively. Computed values of $\text{Si}/(\text{Al}+\text{Fe})$ atomic
43
44 146 ratio were thus larger in CA (7.5) than in NI (2.7). TRB ($\text{cmol}_c \text{ kg}^{-1}$) amounted to 112 in CA and 99 in NI.
45
46 147 Residual weatherable minerals in CA were plagioclase and feldspar, and soil clay minerals. From Figure 2,
47
48 148 CA clay minerals consisted of kaolinite and an assemblage of 2:1:1 and 2:1 clay minerals involving
49
50 149 chlorite, illite, vermiculite and smectite. NI distinctly differed by the large dominance of well crystallized
51
52 150 muscovite and kaolinite. In the Nitisol, K was by far the largest dominant cation in TRB, in line with the
53
54 151 occurrence of muscovite, a dioctahedral mica resistant to weathering. The large amount of non-
55
56
57
58
59
60

exchangeable K (Table 3) thus hid the advanced weathering stage of NI because muscovite is stable and does not weather in soils. The reserve of weatherable minerals was then estimated by the difference between TRB and the content of non-exchangeable K, excluding K-bearing muscovite. [TRB-K] ($\text{cmol}_c \text{ kg}^{-1}$) amounted to 72.5 in CA and 22.1 in NI, indicating the strongly advanced weathering stage of NI. [TRB-K] did not increase following biochar application. To the contrary, wollastonite application generated a marked increase of total Ca content ($\text{cmol}_c \text{ kg}^{-1}$): from 15 to 63 in CA, and from 1 to 48 in NI. Cumulative $\text{CaCl}_2\text{-Si}$ (mg kg^{-1}) (Table 3, Figure 3) was significantly larger in CA (34) than in NI (24). $\text{CaCl}_2\text{-Si}$ significantly ($p < 0.01$) increased after Wo application in CA (398.7) and NI (71.7). Unlike SW, Mi significantly ($p < 0.01$) increased $\text{CaCl}_2\text{-Si}$ in CA (43.2) and NI (38.5).

Biomass and mineralomass of cotton plants

The cotton biomass (g plant^{-1}) was significantly larger in CA (8.6) than in NI (0.32) (Table 4). Mi significantly increased cotton biomass, by 84% in CA, and 368% in NI. Mi significantly increased plant mineralomass of Ca, K and Mg in both soils. Plant Si mineralomass (mg plant^{-1}) was, by far, significantly larger in CA (10.6) than in NI (0.8). Relatively to SW, Mi significantly increased the plant Si content ($\text{g Si kg}^{-1} \text{ DW}$): 0.65 for CA-Mi and 0.74 for NI-Mi, as well as plant Si mineralomass (mg Si plant^{-1}): 10.52 for CA-Mi and 1.12 for NI-Mi). Si mineralomass indeed increased by 120% in CA-Mi and 180% in NI-Mi relatively to the untreated soils CA and Ni.

Discussion

Weathering stage of soils

The values of TRB and $\text{Si}/(\text{Al}+\text{Fe})$ ratio as well as the nature of secondary clay minerals are used to assess soil weathering stage (Herbillon, 1986). Taking into account the stability of muscovite in soils, NI exhibited an advanced weathering stage characteristic of strongly weathered ferrallitic soils (Herbillon, 1986). In CA, the large dominance of quartz is characteristic of soils of the Belgian loess belt, in which the

1
2
3 175 reserve of weatherable minerals is concentrated in the clay fraction (Brahya et al., 2000). CA soil clay
4
5 176 minerals (kaolinite, chlorite, illite, vermiculite and smectite: Figure 2a) revealed a mineralogical
6
7 177 assemblage typical of moderately weathered soils under temperate climates. To the contrary, NI soil clay
8
9 178 minerals (Figure 2b) make an assemblage typical for desilicated, highly weathered soils formed under
10
11 179 humid tropical conditions. Consequently, CaCl₂-Si content was significantly larger in CA than in NI
12
13 180 (Figure 3).

14
15
16
17 181 *Mobility of Si in soil:amendment mixtures*
18
19

20 182 As it is linked to the DSi pool from which plant roots take up Si, CaCl₂-Si could quantify the pool of
21
22 183 bioavailable Si in soils (Sauer et al., 2006). Cumulative CaCl₂-Si (mg kg⁻¹) was significantly larger in
23
24 184 soil:Wo (398.7 in CA-Wo, 71.7 in NI-Wo) than in soil:biochar CA-Mi (43.2), NI-Mi (38.5), CA-SW
25
26 185 (30.9) and NI-SW (24.1) (Table 3; Figure 3). Wo is indeed an efficient Si fertilizer releasing bioavailable
27
28 186 Si. For each amendment, CaCl₂-Si was systematically smaller in NI:amendment mixtures. Part of H₄SiO₄^o
29
30 187 released by amendments was undoubtedly adsorbed by oxide surfaces in NI as it commonly occurs in
31
32 188 highly weathered soils (Herbillon et al., 1977).

33
34
35
36 189 To further assess Si mobility, we define the efficiency of each amendment (AM) using CaCl₂-Si and total
37
38 190 Si contents (Table 3) as follows:
39
40

41
42
$$\frac{[CaCl_2 - Si \text{ in soil: AM}] - [CaCl_2 - Si \text{ in soil}]}{\text{Total Si content of AM}} \times 100$$

43
44
45

46 191 In CA, the efficiency of MI was significantly smaller (0.9%) than that of Wo (5.2%). To the contrary, in
47
48 192 NI, Mi efficiency was larger (1.4%) than to the one of Wo (0.7%). Possibly phytolith solubility could be
49
50 193 larger in NI-Mi, because of a larger pH. Besides, the release of bioavailable Si was significantly larger for
51
52 194 Mi than for SW (Figure 3 c, d). The ample reserve of PhSi in Mi vs SW (34.6 vs 0.9 g Si kg⁻¹, Table 1) and
53
54 195 the accessibility of phytoliths in Mi (Figure 1 e, f) support that phytoliths readily contribute to DSi. They
55
56
57
58
59
60

resist to pyrolytic temperatures as high as 550°C, and further dissolve to release DSi given their solubility (Frayse et al., 2009). Since their dissolution rate increases with increasing pH in the range of pH 3-12 (Frayse et al., 2006), the ability of phytolith-rich Mi to supply DSi is enhanced by pH increase, especially in NI, where silica sorption increased pH of surrounding solution. Indeed, Fe oxides adsorb Si in soils (Jones and Handreck, 1963, 1967) and control DSi (McKeague and Cline, 1963).

The release of bioavailable Si in Mi-amended soils was not affected by soil type and weathering stage. The biochar derived from high-Si accumulator *Miscanthus* is thus an alternative Si fertilizer to wollastonite to supply bioavailable Si in the highly weathered Nitisol.

Effect of biochar amendment on soil properties

As already reported (Sohi et al., 2010; Van Zwieten et al., 2010), biochar application to soils affects their properties (Table 2). The impact depends, however, on the types of biochar and soil. Supplying ash biochar induces a change of pH and nutrient availability, (Clarholm, 1994; Mahmood et al., 2003). Here, applying biochar did not significantly modify CEC in CA. However, it slightly increased CEC in NI, but only after Mi application, which also increased the contents of exchangeable cations, namely K and Mg in CA-Mi, and Ca, K, Na and Mg in NI-Mi (Table 2). Such increases were not observed after SW addition. Since applying Mi increased the effective CEC (ECEC), we hypothesize that it is due to an increase of surface area and surface charge density of biochar particle, as previously reported for black C (Liang et al., 2006). The pH rise, in turn, undoubtedly led to increase ECEC (Glaser et al., 2002; Houben et al., 2014; Laird et al., 2010). Relatively to SW, Mi contained more ashes and initial exchangeable and non-exchangeable cations (Table 1 and Figure 1). Mi thus had a larger ability to increase pH and ECEC, as observed by Houben et al. (2014). Furthermore, the improvement of soil properties was more pronounced in NI-Mi than in CA-Mi. Base saturation (BS %) significantly increased in both CA-Mi and NI-Mi, reflecting the ability of Mi biochar to enhance nutrient availability (Table 3) as predicted by Liang et al. (2006) for black C. The larger BS after Mi amendment was due to the larger nutrient content (Ca, Mg and

1
2
3 220 K) in Mi relatively to SW (Table 1). The impact of Mi biochar on soil fertility was thus positive,
4
5 221 particularly in the highly weathered Nitisol, though at a very low level tested in non-fertilized soil.
6
7

8 222 *Effect of biochar amendment on cotton biomass and Si mineralomass*
9

10
11 223 The amounts of available nutrients (exchangeable Ca, K and Mg) and bioavailable Si were larger in CA
12
13 224 than in NI (Tables 2 and 3). Thus, the cotton biomass and mineralomass were significantly larger in CA
14
15 225 than in NI regardless the type of biochar amendment.
16
17

18
19 226 Plant response to biochar amendment varied with the type of biochar (*Spokas and Reicosky, 2009*) and
20
21 227 application rate (*Liu et al., 2013*). Mi-biochar increased both the plant biomass and element uptake, as a
22
23 228 result of elevated nutrient availability (Ca, Mg and K). Here, both the Ca, Mg and K mineralomass and
24
25 229 soil Ca, Mg and K contents were significantly larger, respectively, in CA-Mi and NI-Mi than in CA-SW
26
27 230 and NI-SW (Table 2 and Table 4). Nutrients were likely released in soil via the degradation of labile
28
29 231 organic compounds associated with biochar (*Rajkovich et al., 2012; Topoliantz et al., 2005; Yamato et al.,*
30
31 232 *2006*). Moreover, biochar amendment promotes plant productivity through the increase of pH, which in
32
33 233 turn increases nutrient availability and/or decrease mineral toxicities (*Lehmann et al., 2003; Rondon et al.,*
34
35 234 *2007; Yamato et al., 2006*). Here, the biochar-induced increase of ECEC likely promoted plant growth and
36
37 235 biomass (Table 2 and Table 4) (*Major et al., 2010*). The marked difference of soil fertility between CA and
38
39 236 NI, discussed above, induced a huge difference in cotton biomass (g plant^{-1}): 8.8 in CA vs 0.3 in NI (Table
40
41 237 3 and Table 4). The increase of cotton biomass was 4.7 fold from NI to NI-Mi and 1.8-fold from CA to
42
43 238 CA-Mi, whereas no significant effect was measured after SW amendment. Mi-biochar induced an increase
44
45 239 of CEC and ECEC, and thus improved soil fertility in the highly weathered Nitisol, thereby promoting
46
47 240 plant growth.
48
49
50

51 241 Si uptake by cotton plants markedly differed between CA and NI (Figure 4). After Mi application, Si
52
53 242 uptake increased by 5.9-fold in NI, and 2.2-fold in CA, keeping in mind that biomass (g.plant^{-1}) (0.2 – 1.5
54
55
56
57
58
59
60

g.plant⁻¹) was very low in NI (0.2–1.5) compared to CA (8.8–16.2) (Table 4). As discussed above, phytoliths were abundant in Mi; they dissolved and released DSi, thereby contributing to the pool of bioavailable Si, which was readily taken up by plant roots (Figure 1, 2, 3; Table 3). This effect was particularly important in the highly weathered Nitisol depleted in primary and secondary silicates.

Conclusion

Miscanthus biochar and wollastonite amendments significantly ($p<0.01$) enhanced the release of CaCl₂-extractable Si in a young Cambisol and an old, highly weathered Nitisol. Soft wood biochar amendment had no effect on the release of CaCl₂-extractable Si in both soils. *Miscanthus* biochar supply increased ECEC, particularly the content of exchangeable K, while soft wood biochar did not have any effect on these parameters whatever the soil type. Both biochar amendments induced a significant effect on C and N contents in both soils. *Miscanthus* biochar generated a significant increase of cotton biomass. That increase was, however, larger in the Nitisol than in the Cambisol, since it was 4.7-fold in the former and 1.8-fold in the latter. In terms of Si uptake after Mi application, the increase in the Nitisol (5.9-fold) was much larger than in the Cambisol (2.2-fold), supporting a larger efficiency of Si release from *Miscanthus* phytoliths in the Nitisol.

Our data thus highlight that *Miscanthus* biochar is an alternative Si amendment to wollastonite to supply bio-available Si, increase plant biomass and promote the biological cycle of Si in the soil-plant system. This positive impact of a Si-rich biochar is, however, larger in the old, desilicated, highly weathered Nitisol. Overall, it demonstrates that Si is progressively released from phytolith-rich biochar, and directly contributes to the pool of bioavailable Si for plant root uptake in soil. This is particularly important for agro-ecosystems in which high-Si accumulator plants grow in highly weathered, desilicated soils. Further field experiments are required to confirm the effect of phytolith-rich biochar as a Si fertilizer over the whole plant cycle. Furthermore, the efficiency of various biochar types to release Si and plant nutrients should be studied.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgements

We thank A. Iserentant and C. Givron for laboratory assistance (UCL). Z Li is supported by the “Fonds Spécial de Recherché” of the UCL in 2014-2015 and the Aspirant (ASP)-Fonds National de la Recherché Scientifique» (FNRS) of Belgium in 2015-2019.

References

Alexandre, A. , Bouvet, M. , Abbadie, L. (2011). The role of savannas in the terrestrial Si cycle: a case-study from Lamto, Ivory Coast. *Global and Planetary Change* **78**, 162-169.

Barão, L. , Clymans, W. , Vandevenne, F. , Meire, P. , Conley, D., Struyf, E. (2014). Pedogenic and biogenic alkaline-extracted silicon distributions along a temperate land-use gradient. *European Journal of Soil Science* **65**, 693-705.

Belanger, R. R. (1995). Soluble silicon: its role in crop and disease management of greenhouse crops. *Plant Dis* **79**, 329-336.

Berthelsen, S. , Noble, A. , Garside, A. (2001). Silicon research down under: past, present, and future. *Studies in Plant Science* **8**, 241-255.

Biederman, L. A. , Harpole, W. S. (2013). Biochar and its effects on plant productivity and nutrient cycling: a meta-analysis. *GCB bioenergy* **5**, 202-214.

Blecker, S. W. , McCulley, R. L. , Chadwick, O. A. , Kelly, E. F. (2006). Biologic cycling of silica across a grassland bioclimate sequence. *Global Biogeochemical Cycles* **20**.

Brahy, V. , Deckers, J. , Delvaux, B. (2000). Estimation of soil weathering stage and acid neutralizing capacity in a toposequence Luvisol–Cambisol on loess under deciduous forest in Belgium. *European Journal of Soil Science* **51**, 1-13.

Chao, T. , Sanzolone, R. (1992). Decomposition techniques. *Journal of Geochemical Exploration* **44**, 65-106.

Chapman, H. (1965). Cation-exchange capacity. Methods of soil analysis. Part 2. *Chemical and microbiological properties*, 891-901.

Clarholm, M. (1994). Granulated wood ash and a ‘N-free’ fertilizer to a forest soil—effects on P availability. *Forest Ecology and Management* **66**, 127-136.

Cornelis, J.-T. , Titeux, H. , Ranger, J. , Delvaux, B. (2011). Identification and distribution of the readily soluble silicon pool in a temperate forest soil below three distinct tree species. *Plant and soil* **342**, 369-378.

Cornelis, J. T. , Delvaux, B. (2016). Soil processes drive the biological silicon feedback loop. *Functional Ecology* **30**, 1298-1310.

Crane-Droesch, A. , Abiven, S. , Jeffery, S., Torn, M. S. (2013). Heterogeneous global crop yield response to biochar: a meta-regression analysis. *Environmental Research Letters* **8**, 044049.

Datnoff, L. E. , Heckman, J. R. (2014). Silicon fertilizers for plant disease protection. In "World Fertilizer Congress", Vol. **16**, pp. 36.

Epstein, E. (1994). The anomaly of silicon in plant biology. *Proceedings of the National Academy of Sciences* **91**, 11-17.

Fauteux, F. , Rémus-Borel, W. , Menzies, J. G. , Bélanger, R. R. (2005). Silicon and plant disease resistance against pathogenic fungi. *FEMS Microbiology letters* **249**, 1-6.

Fraysse, F. , Pokrovsky, O. S., Schott, J. , Meunier, J.-D. (2006). Surface properties, solubility and dissolution kinetics of bamboo phytoliths. *Geochimica et Cosmochimica Acta* **70**, 1939-1951.

Fraysse, F. , Pokrovsky, O. S., Schott, J. , Meunier, J.-D. (2009). Surface chemistry and reactivity of plant

- phytoliths in aqueous solutions. *Chemical Geology* **258**, 197-206.
- Gérard, F. , Mayer, K., Hodson, M., Ranger, J. (2008). Modelling the biogeochemical cycle of silicon in soils: application to a temperate forest ecosystem. *Geochimica et Cosmochimica Acta* **72**, 741-758.
- Glaser, B. , Lehmann, J. , Zech, W. (2002). Ameliorating physical and chemical properties of highly weathered soils in the tropics with charcoal—a review. *Biology and fertility of soils* **35**, 219-230.
- Haynes, R. J. (2014). A contemporary overview of silicon availability in agricultural soils. *Journal of Plant Nutrition and Soil Science* **177**, 831-844.
- Haynes, R. J. , Belyaeva, O. , Kingston, G. (2013). Evaluation of industrial wastes as sources of fertilizer silicon using chemical extractions and plant uptake. *Journal of Plant Nutrition and Soil Science* **176**, 238-248.
- Haysom, M. , Chapman, L. (1975). Some aspects of the calcium silicate trials at Mackay. In "Proceedings".
- Henriet, C., Bodarwé, L. , Dorel, M. , Draye, X., Delvaux, B. (2008). Leaf silicon content in banana (*Musa* spp.) reveals the weathering stage of volcanic ash soils in Guadeloupe. *Plant and Soil* **313**, 71-82.
- Herbillon, A. (1986). Chemical estimation of weatherable minerals present in the diagnostic horizons of low activity clay soils. In "Proceedings of the 8th International Clay Classification Workshop: Classification, Characterization and Utilization of Oxisols (part 1)[Beinroth, FH, Camargo, MN and Eswaran (ed.)][39–48](Rio de Janeiro, 1986)".
- Herbillon A., Gallez A., Juo S.R. (1977) Characteristics of Silica Sorption and Solubility as Parameters to Evaluate the Surface Properties of Tropical Soils: II. The Index of Silica Saturation. *Soil Science Society of America Journal* **41**, 1151-1154.
- Houben, D. , Sonnet, P. , Cornelis, J.-T. (2014). Biochar from *Miscanthus*: a potential silicon fertilizer. *Plant and soil* **374**, 871-882.
- Issaharou-Matchi, I. , Barboni, D. , Meunier, J.-D. , Saadou, M. , Dussouillez, P. , Contoux, C. , Zirihiguede, N. (2016). Intraspecific biogenic silica variations in the grass species *Pennisetum pedicellatum* along an evapotranspiration gradient in South Niger. *Flora-Morphology, Distribution, Functional Ecology of Plants* **220**, 84-93.
- IUSS (2014). World reference base for soil resources 2014 international soil classification system for naming soils and creating legends for soil maps. *FAO*, Rome.
- Jeffery, S. , Verheijen, F. G. , Van Der Velde, M. , Bastos, A. C. (2011). A quantitative review of the effects of biochar application to soils on crop productivity using meta-analysis. *Agriculture, ecosystems & environment* **144**, 175-187.
- Jones, L., Handreck, K. (1963). Effects of iron and aluminium oxides on silica in solution in soils.
- Jones, L. , Handreck, K. (1967). Silica in soils, plants, and animals. *Advances in agronomy* **19**, 107-149.
- Keller, C. , Guntzer, F. , Barboni, D. , Labreuche, J. , Meunier, J.-D. (2012). Impact of agriculture on the Si biogeochemical cycle: input from phytolith studies. *Comptes Rendus Geoscience* **344**, 739-746.
- Kloss, S. , Zehetner, F. , Wimmer, B. , Buecker, J. , Rempt, F., Soja, G. (2014). Biochar application to temperate soils: effects on soil fertility and crop growth under greenhouse conditions. *Journal of plant nutrition and soil science* **177**, 3-15.
- Laird, D. A., Fleming, P., Davis, D. D., Horton, R., Wang, B., Karlen, D. L. (2010). Impact of biochar amendments on the quality of a typical Midwestern agricultural soil. *Geoderma* **158**, 443-449.
- Lehmann, J. , da Silva Jr, J. P. , Steiner, C., Nehls, T. , Zech, W. , Glaser, B. (2003). Nutrient availability and leaching in an archaeological Anthrosol and a Ferralsol of the Central Amazon basin: fertilizer, manure and charcoal amendments. *Plant and soil* **249**, 343-357.
- Lehmann, J. , Joseph, S. (2015). "Biochar for environmental management: science, technology and implementation," Routledge.
- Lehmann, J. , Rillig, M. C. , Thies, J. , Masiello, C. A. , Hockaday, W. C. , Crowley, D. (2011). Biochar effects on soil biota—a review. *Soil Biology and Biochemistry* **43**, 1812-1836.
- Liang, B. , Lehmann, J. , Solomon, D. , Kinyangi, J. , Grossman, J. , O'Neill, B. , Skjemstad, J. , Thies, J. , Luizao, F. , Petersen, J. (2006). Black carbon increases cation exchange capacity in soils. *Soil*

1
2
3 361 *Science Society of America Journal* **70**, 1719-1730.
4 362 Liang, Y. , Sun, W. , Zhu, Y.-G. , Christie, P. (2007). Mechanisms of silicon-mediated alleviation of abiotic
5 363 stresses in higher plants: a review. *Environmental pollution* **147**, 422-428.
6 364 Liu, X. , Li, L., Bian, R. , Chen, D. , Qu, J. , Wanjiru Kibue, G. , Pan, G. , Zhang, X. , Zheng, J. , Zheng, J.
7 365 (2014). Effect of biochar amendment on soil-silicon availability and rice uptake. *Journal of plant*
8 366 *nutrition and soil science* **177**, 91-96.
9 367 Liu, X. , Zhang, A. , Ji, C., Joseph, S. , Bian, R. , Li, L. , Pan, G. , Paz-Ferreiro, J. (2013). Biochar's effect
10 368 on crop productivity and the dependence on experimental conditions—a meta-analysis of
11 369 literature data. *Plant and soil* **373**, 583-594.
12 370 Ma, J. F. , Tamai, K. , Yamaji, N. , Mitani, N. , Konishi, S. , Katsuhara, M. , Ishiguro, M. , Murata, Y. ,
13 371 Yano, M. (2006). A silicon transporter in rice. *Nature* **440**, 688-691.
14 372 Mahmood, S. , Finlay, R. D. , Fransson, A.-M. , Wallander, H. (2003). Effects of hardened wood ash on
15 373 microbial activity, plant growth and nutrient uptake by ectomycorrhizal spruce seedlings. *FEMS*
16 374 *Microbiology Ecology* **43**, 121-131.
17 375 Major, J. , Rondon, M. , Molina, D. , Riha, S. J. , Lehmann, J. (2010). Maize yield and nutrition during 4
18 376 years after biochar application to a Colombian savanna oxisol. *Plant and soil* **333**, 117-128.
19 377 McKeague, J. , Cline, M. (1963). Silica in soil solutions: II. The adsorption of monosilicic acid by soil and
20 378 by other substances. *Canadian Journal of Soil Science* **43**, 83-96.
21 379 Rajkovich, S. , Enders, A. , Hanley, K. , Hyland, C. , Zimmerman, A. R. , Lehmann, J. (2012). Corn growth
22 380 and nitrogen nutrition after additions of biochars with varying properties to a temperate soil.
23 381 *Biology and Fertility of Soils* **48**, 271-284.
24 382 Rondon, M. A. , Lehmann, J. , Ramirez, J. , Hurtado, M. (2007). Biological nitrogen fixation by common
25 383 beans (*Phaseolus vulgaris* L.) increases with bio-char additions. *Biology and fertility of soils* **43**,
26 384 699-708.
27 385 Sauer, D., Saccone, L. , Conley, D. J. , Herrmann, L. , Sommer, M. (2006). Review of methodologies for
28 386 extracting plant-available and amorphous Si from soils and aquatic sediments. *Biogeochemistry*
29 387 **80**, 89-108.
30 388 Sohi, S. , Krull, E. , Lopez-Capel, E., Bol, R. (2010). A review of biochar and its use and function in soil.
31 389 *Advances in agronomy* **105**, 47-82.
32 390 Spokas, K. , Reicosky, D. (2009). Impacts of Sixteen Different Biochars on Soil Greenhouse Gas
33 391 Production. *Annals of Environmental Science* **3**, 179-193.
34 392 Struyf, E. , Smis, A. , Van Damme, S. , Garnier, J. , Govers, G. , Van Wesemael, B. , Conley, D. J. ,
35 393 Batelaan, O. , Frot, E. , Clymans, W. (2010). Historical land use change has lowered terrestrial
36 394 silica mobilization. *Nature communications* **1**, 129.
37 395 Thiry, M. , Quesnel, F. , Yans, J. , Wyls, R. , Vergari, A. , Theveniaut, H. , Simon-Coinçon, R. , Ricordel,
38 396 C. , Moreau, M.-G. , Giot, D. (2006). Continental France and Belgium during the early
39 397 Cretaceous: paleoweatherings and paleolandforms. *Bulletin de la Société géologique de France*
40 398 **177**, 155-175.
41 399 Titeux, H. , Delvaux, B. (2009). Experimental study of DOC, nutrients and metals release from forest
42 400 floors developed under beech (*Fagus sylvatica* L.) on a Cambisol and a Podzol. *Geoderma* **148**,
43 401 291-298.
44 402 Topoliantz, S. , Ponge, J.-F. , Ballof, S. (2005). Manioc peel and charcoal: a potential organic amendment
45 403 for sustainable soil fertility in the tropics. *Biology and Fertility of Soils* **41**, 15-21.
46 404 Van Zwieten, L. , Kimber, S. , Morris, S. , Chan, K. , Downie, A. , Rust, J. , Joseph, S. , Cowie, A. (2010).
47 405 Effects of biochar from slow pyrolysis of papermill waste on agronomic performance and soil
48 406 fertility. *Plant and soil* **327**, 235-246.
49 407 Vandevenne, F. , Barão, L. , Ronchi, B. , Govers, G. , Meire, P. , Kelly, E. , Struyf, E. (2015). Silicon pools
50 408 in human impacted soils of temperate zones. *Global Biogeochemical Cycles* **29**, 1439-1450.
51 409 Vandevenne, F. , Struyf, E. , Clymans, W. , Meire, P. (2012). Agricultural silica harvest: have humans
52 410 created a new loop in the global silica cycle? *Frontiers in Ecology and the Environment* **10**, 243-
53 411 248.

- 1
2
3 412 White, A. F. , Vivit, D. V. , Schulz, M. S., Bullen, T. D. , Evett, R. R. , Agarwal, J. (2012). Biogenic and
4 413 pedogenic controls on Si distributions and cycling in grasslands of the Santa Cruz soil
5 414 chronosequence, California. *Geochimica et Cosmochimica Acta* **94**, 72-94.
6 415 Xiao, X. , Chen, B. , Zhu, L. (2014). Transformation, morphology, and dissolution of silicon and carbon in
7 416 rice straw-derived biochars under different pyrolytic temperatures. *Environmental science &*
8 417 *technology* **48**, 3411-3419.
9 418 Yamato, M. , Okimori, Y. , Wibowo, I. F. , Anshori, S. , Ogawa, M. (2006). Effects of the application of
10 419 charred bark of *Acacia mangium* on the yield of maize, cowpea and peanut, and soil chemical
11 420 properties in South Sumatra, Indonesia. *Soil science and plant nutrition* **52**, 489-495.
12 421
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table1. Selected properties of the soft wood (SW), *Miscanthus* (Mi) biochars and wollastonite (Wo): pH_{water}, total contents of Carbon (C), Nitrogen (N), CaCl₂ extractable Si (CaCl₂-Si), major alkaline and alkaline-earth cations (Ca, Mg, K, Na), and silicon (Si).

Amendments	pH	[§] C	[§] N	CaCl ₂ -Si	Ca	Mg	Na	K	Si
	H ₂ O	(%)		g kg ⁻¹			(g kg ⁻¹)		
SW	7.4	84.90	1.15	0.01(0.00)	2.79(0.79)	0.53(0.17)	0.15(0.05)	2.20(0.61)	0.77(0.22)
Mi	10.1	77.13	2.01	11.45(1.05)	6.92(0.10)	2.15(0.10)	0.58(0.03)	14.87(0.16)	34.65(0.23)
Wo	9.1	0.56	nd	8.87(1.04)	314.71(2.54)	10.80(0.06)	<0.01	1.10(0.06)	233.34(2.03)

Note: nd means not determined; [§]: Single measurements only.

Table 2. Selected physico-chemical properties of the untreated soils (Cambisol CA, Nitisol NI) and amended soils, using soft wood (CA-SW, NI-SW), *Miscanthus* (CA-Mi, NI-Mi) and wollastonite (CA-Wo, NI-Wo). The average values presented with distinct letters (a, b, c) in the Cambisol and (x, y, z) in the Nitisol are significantly different at the $p < 0.05$ level of confidence according to LSD's multiple comparison test.

Treatments	pH		C	N	CEC	Exchangeable cations				BS
	H ₂ O	KCl				(cmol _c kg ⁻¹)				
			g kg ⁻¹		Ca ²⁺	K ⁺	Na ⁺	Mg ²⁺	(%)	
CA	4.62(0.02)c	3.72(0.03)c	14.12(1.32)b	1.19(0.05)a	6.67(0.31)b	2.31(0.05)b	0.20(0.02)b	0.04(0.02)c	0.39(0.01)c	44.08
CA-SW	4.72(0.00)c	3.73(0.01)c	30.95(2.38)a	1.22(0.12)a	6.14(0.82)b	2.29(0.07)b	0.22(0.02)b	0.04(0.01)c	0.39(0.01)c	47.88
CA-Mi	5.03(0.03)b	4.08(0.02)b	35.85(2.56)a	1.27(0.35)a	5.98(0.17)b	2.36(0.06)b	0.97(0.20)a	0.07(0.01)b	0.47(0.04)a	64.72
CA-Wo	6.25(0.01)a	5.36(0.02)a	14.78(1.23)b	1.11(0.03)a	8.11(0.11)a	5.42(0.13)a	0.24(0.03)b	0.10(0.01)a	0.43(0.00)b	76.32
NI	5.18(0.04)z	4.32(0.01)z	0.58(0.35)y	0.27(0.02)y	0.68(0.08)z	0.15(0.01)z	0.05(0.02)y	0.01(0.01)z	0.30(0.00)y	75.00
NI-SW	5.45(0.00)z	4.44(0.06)z	26.41(4.56)x	0.53(0.12)x	0.87(0.13)z	0.16(0.02)z	0.07(0.04)y	0.02(0.00)z	0.30(0.00)y	63.22
NI-Mi	6.25(0.10)y	5.79(0.13)y	21.29(3.69)x	0.50(0.11)x	1.04(0.07)y	0.36(0.13)y	0.81(0.55)x	0.07(0.03)x	0.37(0.05)x	154.81
NI-Wo	7.84(0.12)x	6.53(0.10)x	0.85(0.32)y	0.27(0.01)y	1.82(0.03)x	2.89x(0.23)	0.03(0.03)z	0.06(0.01)y	0.28(0.02)z	179.12

Table 3. Total contents of selected major elements and Total Reserve in Bases (TRB) in CA and NI, CA-SW, CA-Mi, CA-Wo; NI-SW, NI-Mi, NI-Wo). The average values presented with distinct letters (a, b, c) in Cambisols and (x, y, z) in Nitisols are significantly different at the $p<0.05$ level of confidence according to LSD's multiple comparison test.

Treatments	CaCl ₂ -Si ¹ (mg kg ⁻¹)	Total Si (g kg ⁻¹)	Total elements (cmol _c kg ⁻¹)				TRB ²	[TRB-K] ³	Total elements (g kg ⁻¹)		Si/(Al+Fe)
			Ca	K	Na	Mg			Al	Fe	
CA	34.05(1.67)c	376.78	16.50	40.00	29.80	26.00	112.3	72.5	39.21	18.52	7.52
CA-SW	30.95(1.00)c	365.47	16.55	39.06	28.92	25.34	109.9	70.5	38.02	17.92	7.52
CA-Mi	43.22(1.90)b	366.50	17.08	40.00	28.99	25.34	111.4	72.4	38.01	17.93	7.54
CA-Wo	398.69(1.03)a	368.74	63.25	38.94	28.91	27.86	158.9	120.2	38.13	18.00	7.57
NI	24.03(0.68)z	311.92	1.00	77.60	7.70	13.40	99.7	22.1	96.52	30.61	2.69
NI-SW	24.11(1.04)z	302.57	1.46	75.70	7.47	13.13	97.7	22.0	93.62	29.71	2.69
NI-Mi	38.50(1.52)y	303.60	2.00	76.64	7.54	13.13	99.3	23.4	93.64	29.73	2.70
NI-Wo	71.70(2.64)x	305.84	48.16	75.57	7.46	15.64	146.8	71.3	93.73	29.72	2.72

¹: cumulative amount after 5 extractions (see also Figure 3);
²: TRB is the sum of total contents of Ca, Mg, Na and K in soil (*Herbillon*, 1986);
³: TRB-K is the sum of total contents of Ca, Mg and Na in soil, excluding non-exchangeable K [total K – exchangeable K], associated with resistant muscovite or oxidized biotite, both dioctahedral micaceous minerals.

Table 4. Experimental values of cotton biomass, plant contents of Ca, Mg, Na, K, and Si, and element mineralomass (i.e. uptake) in the cotton plants grown on the untreated soils (Cambisol CA, Nitisol NI), and biochar-amended soils (*Miscanthus* Mi; Soft Wood SW). The average values presented with distinct letters (a, b, c) in CA and (x, y, z) in NI are significantly different at the $p < 0.05$ level of confidence according to LSD's multiple comparison test.

Treatments		CA	CA-SW	CA-Mi	NI	NI-SW	NI-Mi
Biomass	g plant ⁻¹	8.82(1.10)b	8.63(2.51)b	16.24(2.11)a	0.32(0.07)y	0.24(0.11)y	1.51(0.40)x
Ca	g kg ⁻¹	14.64(4.59)a	7.79(0.62)b	13.22(2.05)a	3.16(1.09)y	3.82(1.39)y	11.42(2.46)x
Mg		2.76(0.48)a	2.13(0.30)b	2.48(0.27)a	1.54(0.00)y	1.30(0.10)y	2.25(0.05)x
Na		0.24(0.08)a	0.11(0.02)b	0.09(0.03)b	0.82(0.01)x	0.58(0.00)y	0.20(0.22)z
K		15.17(3.39)b	15.15(1.33)b	23.03(4.96)a	10.40(0.01)y	14.51(0.60)y	26.74(1.61)x
Si		0.54(0.16)b	0.46(0.11)b	0.65(0.03)a	0.58(0.00)y	0.54(0.00)y	0.74(0.11)x
Ca	mg plant ⁻¹	129.09(25.14)b	67.22(16.59)c	214.73(37.52)a	1.01(0.22)y	0.92(0.65)y	17.24(4.52)x
Mg		24.35(7.27)bc	18.42(3.05)c	40.25(2.64)a	0.49(0.11)y	0.31(0.16)y	3.41(0.89)x
Na		2.09(0.84)a	0.92(0.32)c	1.53(0.44)b	0.26(0.05)xy	0.14(0.07)y	0.30(0.07)x
K		133.76(45.57)b	130.78(28.06)b	374.05(79.87)a	3.33(0.72)y	3.48(1.73)y	40.38(10.59)x
Si		4.76(0.32)b	3.93(0.21)c	10.52(2.30)a	0.19(0.05)y	0.13(0.07)y	1.12(0.12)x

Figures

Figure 1. Scanning electron micrographs (SEM) images of biochar materials from soft wood (a, b), and *Miscanthus* (*Miscanthus* × *giganteus*) straws (c, d). e and f are the SEM images of *Miscanthus* phytoliths. The energy dispersive X-ray spectra (EDX) (g) collected from f, shows the dominance of C, O and Si, hence documenting the presence of phytoliths (SiO₂·nH₂O).

Figure 2. XRD patterns of the clay-size fraction ($<2\ \mu\text{m}$) of the Cambisol CA (a) and Nitisol NI (b) after K^+ -saturation at 20, 105, 300 and 550°C; Mg^{2+} -saturation at 20 °C, and saturation with ethylene-glycol.

Figure 3. Plot of the cumulative amount of CaCl_2 extractable Si content against the extraction number (the Si content at each number is added to the Si contents extracted following the preceding(s) extraction(s) in (a, b) soil:wollastonite (a, b: NI-Wo and CA-Wo); (c) NI, NI-Mi and NI-SW, and (d) CA, CA-Mi and CA-SW.

Figure 4. Plot of Si uptake by cotton plants against the total reserve of Ca, Mg and Na minerals (excluding K-bearing muscovite) in the Cambisol (CA) and Nitisol (NI).