Oxygen Consumption by Warm Ischemia-Injured Porcine Kidneys in Hypothermic Static and Machine Preservation

Jacques Kaminski, Pierre-Olivier Delpech, Sihem Kaaki-Hosni, Xavier Promeyrat, Thierry Hauet, Patrick Hannaert

► To cite this version:

Jacques Kaminski, Pierre-Olivier Delpech, Sihem Kaaki-Hosni, Xavier Promeyrat, Thierry Hauet, et al.. Oxygen Consumption by Warm Ischemia-Injured Porcine Kidneys in Hypothermic Static and Machine Preservation. Journal of Surgical Research, 2019, 242, pp.78-86. 10.1016/j.jss.2019.04.015 . hal-02177529

HAL Id: hal-02177529

<https://hal.science/hal-02177529>

Submitted on 22 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution - NonCommercial| 4.0 International License

Oxygen consumption by WI-injured porcine kidneys in hypothermic static and machine preservation

Jacques Kaminski¹, Pierre-Olivier Delpech^{1,2}, Sihem Kaaki-Hosni¹,

Xavier Promeprat³, Thierry Hauet^{1,4}, Patrick Hannaert¹

Affiliations :

¹, INSERM U1082-IRTOMIT, CHU de Poitiers, 2 rue de la Milétrie,
86000 Poitiers, France

², Service d'Urologie, CHU de Poitiers, 2 rue de la Milétrie,
86000 Poitiers, France

³, Service d'Urologie et de Chirurgie de la Transplantation,
Hôpital Édouard-Herriot, Université Claude-Bernard Lyon 1,
5, place d'Arsonval, 69003 Lyon, France

⁴, Service de Biochimie, CHU de Poitiers, 2 rue de la Milétrie,
86000 Poitiers, France

Corresponding author: Patrick HANNAERT

INSERM U1082 (IRTOMIT), CHU de la Milétrie
2, rue de la Milétrie - 86021 Poitiers Cedex, France

e-mail: patrick.hannaert@univ-poitiers.fr
tel: +33-5-49-44-49-54 - fax: +33-5-49-44-49-22

Authors contributions

J Kaminski performed experiments on isolated kidneys, analyzed data, and contributed to the writing of the paper. S Kaaki-Hosni contributed to oxygen determination experiments. PO Delpech and X Promeyrat performed all surgical procedures. T Hauet initiated the project and contributed to the writing of the paper. P Hannaert analyzed the data, performed bibliographical analysis and wrote the paper.

Abstract

Static cold storage (SCS) and hypothermic machine perfusion (HMP) are currently standard methods for renal grafts clinical preservation. Both methods are predominantly implemented without the active delivery of oxygen, even for DCD-like kidneys. However, even under severe hypothermia (4-6°C), kidneys can consume oxygen and produce ATP. What is not established, though, is to what extent and how SCS and HMP compare in terms of oxygen. Using a porcine preclinical model of renal warm ischemia (WI) to compare SCS and HMP methods, we continuously monitored and quantified oxygen level and consumption along preservation; we also determined pre and post-preservation cortical ATP level; values given as median and [min;max] range. One hour WI reduced ATP by ~90% (from 3.3 [1.7;4.5] mmol/L tissue in Controls). Oxygen consumption (QO_2 , $\mu\text{mol}/\text{min per } 100\text{g}$) was determined from initial solution PO_2 decrease (SCS and HMP), and from arterio-venous difference (HMP). In SCS and HMP, PO_2 decreased rapidly ($t_{1/2}$ ~1h), from atmospheric levels to 52.9 [38.0;65.9] and 8.2 [3.0,16.0] mmHg, respectively. In HMP, QO_2 was 2.7 [0.4;3.9] *versus* 0.5 [0.0;1.3] in SCS ($p < 0.05$); post-preservation ATP amounted to 5.8 [3.2;6.5] in HMP *versus* 0.1 [0.0;0.2] in SCS. Despite hypothermic conditions in SCS or HMP, DCD-like renal grafts require oxygen. Increased oxygen consumption, restored ATP level, and improved histological profile in HMP might explain the established HMP superiority over SCS. These results establish a rational basis for the use of oxygen in hypothermic preservation. Optimal levels required for preservation and graft type variants remain to be determined.

Running title

Oxygen use by suboptimal renal grafts in cold preservation

Key words

Ischemia-reperfusion, Kidney, Preservation, Hypothermic machine perfusion

Static cold storage, Oxygen, ATP

Abbreviations:

ATP, Adenosine Tri-Phosphate

DCD, Donation after Circulatory Death

HES, Hematoxylin-Eosin-Safran

HMP, Hypothermic Machine Perfusion

KPS, Kidney Perfusion Solution

NMP, Normothermic Machine Perfusion

PAS, Periodic Acid Schiff

PO₂, oxygen partial pressure

SCS, Static Cold Storage

~~**SEM**, Standard Error of the Mean~~

UW, University of Wisconsin

WI, warm ischemia

1. Introduction

Despite the progressive emergence of normothermic preservation and reconditioning methods, such as *ex vivo* NMP (normothermic machine perfusion) and *in situ* ECMO (extra-corporeal membrane oxygenation), hypothermic preservation remains the dominant strategy in renal transplantation. Hypothermic preservation is implemented either as *static cold storage* (SCS) or as *hypothermic machine perfusion* (HMP), the later being particularly efficient for higher-risk donors¹⁻³. Biochemical processes, including ATP-producing oxidative phosphorylation and glycolysis, are slowed by low temperature: for instance, at 7°C, cellular reactions occur at a residual ~12% of normothermic rates⁴. However, under hypothermic conditions, even at temperatures as low as 4-6°C, residual oxygen-dependent (and oxygen-independent) metabolic activities can be significant, oxygen has indeed been shown to be consumed, and ATP can be produced, at least under perfusion^{5,6}. Accordingly, the addition of active oxygenation (100%) to HMP preservation improves post-transplantation outcomes, in preclinical models and in clinical situations, as reviewed⁷. Using a porcine model of WI-injured renal grafts, our laboratory showed that, as compared to passive oxygenation, active oxygenation (100%) during HMP preservation significantly improved early functional and inflammatory scores, as well as chronic functional and fibrosis scores⁸. However, as in most other studies, oxygen levels were not measured, and one must surmise that, in the “passive” arm for instance, oxygen level of the preservation solution started at atmospheric levels (21%).

In SCS preservation, on the other hand, to our knowledge, no dedicated study evaluated the addition of active oxygen, nor specifically determined oxygen consumption^{2,7,9}. Of note though, one study from our laboratory showed that the addition of an oxygen carrier (marine extracellular hemoglobin, M101) during SCS preservation significantly improved functional and histological post-transplantation outcomes, with a notable reduction of fibrosis, in a Pig model of autotransplantation of “normal”, uninjured kidneys (WI time < 30min, SCS 24h, one month follow-up)¹⁰. This strongly suggests that the provision of additional oxygen can be beneficial to renal grafts, even under SCS conditions.

In current clinical settings, all SCS and most HMP methods share the common feature of being non-actively oxygenated, i.e. relying on whatever oxygen is present, usually atmospheric oxygen (~158 mmHg); this is the case of the LifePort Kidney Transporter (ORS®). Exceptions which provide for active oxygenation allow for supplementation with either 100% oxygen (760 mmHg, Kidney Assist-Transport from Organ Assist® and IGL® Waves) or atmospheric oxygen (IGL Waves).

Few mechanisms can be invoked to explain the benefits of active oxygenation (or oxygen carriers) in SCS or HMP. Obviously, the intrinsic ability of oxygen to sustain mitochondrial ATP production, represents a key player by maintaining cellular structures and repair processes. An increased ATP level during organ preservation is expected to lead to better outcomes and reduced risks of organ dysfunction after transplantation, in animals and human studies^{9,11}. However, both in SCS and HMP preservation protocols, and even in preclinical studies, actual levels of renal ATP remain largely undetermined -- much like oxygen. The determination of renal graft oxygen levels and consumption, together with tissue ATP levels, should translate into a better understanding of the graft status and its requirements for optimal preservation. Under this rationale, comparing SCS

and HMP, the two current, non-actively oxygenated, preservation standards, we determined i) oxygen levels simultaneously in the preservation solution and within the kidney, and ii) ATP levels before and after WI, and after SCS and HMP preservation. Because the current trend in transplantation is to further extend the donor acceptance criteria, leading to lower quality graft, we performed this experimental study with kidneys pre-injured by one-hour warm ischemia, mimicking DCD grafts (donation after circulatory death) ⁸.

2. Methods

Animals and ethics.

Large White male pigs (3 months) were bred at the IBiSA MOPICT platform (INRA, Surgères, Fr), where organ procurement and euthanasia were carried out. All surgical and experimental protocols were performed in accordance with the guidelines of the French Ministry of Agriculture (FMA) for the use and care of laboratory animals, after approval by the FMA, the INRA, and the Poitou-Charentes ethical committee for animal experimentation.

Surgery and warm ischemia.

Surgery has been described in detail ^{8,12–15}. Briefly, after general anesthesia, midline abdominal incision, and pedicle clamping, the right kidney was removed by dissection of the vascular pedicle and ureter and transferred into a sterile bag for 60 min warm ischemia WI, and maintained at 37°C in a water bath. For oxygen monitoring during warm ischemia, pedicle-clamped, $n = 6$ kidneys were used (see figure 1): they were kept in normothermia *via* a thermostated bath filled with saline, under continuous temperature monitoring (solution and renal cortex). After WI, the kidney was gravimetrically (1 m height) flushed with 500 mL ice-cold UW® for SCS or KPS® for HMP; all flushings occurred in <10 min and yielded whitened kidneys. Figure 1 summarizes the protocol and the use of kidneys.

Hypothermic preservation – Oxygen and temperature measurements.

For SCS (20h), the kidney was placed in 1 L ice-cold UW® (Bridge to Life®, London, UK), and placed itself into a styrofoam box, filled with ice. For HMP (20h) the kidney was connected to a LifePort® machine, filled with ice (Lifeport Kidney Transporter 1.0, ORS®, Brussels, Be), set at 35 mmHg pressure.

Cortical temperature was monitored, *via* 200 µm-glass thermocouple microsensor (TP200, Unisens®), inserted 3-5 mm below the capsule (signal was fed to a Unisens® MicroSensor multimeter). For oxygen monitoring in the HMP perfusate, fiber optic oxygen microsensors (IMP-Pst1, Presens®) were adapted into the perfusion circuit, at the arterial inlet and venous outlets. Cortical O₂ (SCS and HMP) was monitored with Needle-Type optode Oxygen Microsensor (NTH-PSt1, Presens®), introduced 3-5 mm final under the capsule. Fiber optic oxygen electrodes were connected to a PreSens® oxymeter OXY-4 Mini (4 min⁻¹ sampling). Oxygen signal were converted into pressure and concentration with an Excel® sheet (PreSens®). For O₂-sensors, two-points calibration was performed for each experiment (100% N₂ and 100% air-saturated saline).

Renal tissue ATP

The ATP content from renal cortical samples, immediately frozen in liquid nitrogen, was measured with the ATP Bioluminescent Assay kit (Sigma®), with 40-70 mg tissue samples. The final ATP concentration is expressed as mmol/L of tissue (assuming a tissue density of 1 g/mL). An additional group of n = 5 kidneys was used for control ATP (i.e. ATP before warm ischemia; see figure 1).

Histological analysis.

For histological observations (control, end-CS, end-HMP, end-WI) and for control, 0.5x0.5x1.0 cm³ tissue samples (cortex and medulla) were cut and immersed into 4% buffered formalin (VWR®) for fixation. Fixed samples were then colored with HES/PAS (Hematoxylin-Eosin-Safran/Periodic Acid Schiff) for expert examination and semi-quantitative analysis for i) brush-border loss and endoluminal detachment, with the following scoring scale: **0** = no lesion; **1**, <10% area; **2**, 11-25% area; **3**, 26-50 % area; **4**, 51-75% area; **5**, >76% area), and ii) vacuolization and tissue infiltration (number of features *per* optical field). For both stainings and for each parameter, five slides were read, with six different fields for each (n = 30).

Data and statistics.

Arterio-venous oxygen consumption in HMP was calculated as the product of the perfusion rate with the oxygen arterio-venous difference (expressed in $\mu\text{moles/min per}$ 100 g kidney weight). Oxygen consumption from solution concentration was calculated from oxygen decrease between consecutive time-points, under initial rate conditions (10-30 min). Data are presented as median with range [min, max] of n = 5 values, unless otherwise stated (e.g. for cortical temperature mean $\pm$ SD).

Statistical analysis of oxygen time-course was performed as repeated measures ANOVA (NCSS® v0.7.1.21). For ATP values and histological scoring, statistics were performed as non-parametric one-way ANOVA, with Tukey post-test (Graphpad Prism® v5.01). A *p* value < 0.05 was considered significant.

3. Results

Renal oxygen during warm ischemia

Overall, during warm ischemia (WI, 60 min), average cortical temperature was 36.0 ± 0.6 °C (mean $\pm$ SD, $n = 60$ time-points): starting from $32.6 [29.1;35.1]$ °C at zero time ($n = 6$), cortical temperature equilibrated at values between 36 and 37 °C after 10 min; the final WI time-point (60 min) was $36.3 [35.8;36.6]$ °C (equilibration reached after 5-7min). Oxygen levels of the solution, remained in the 140-160 mmHg range (atmospheric level) throughout WI.

Cortical oxygen level was monitored during the warm ischemic phase, simultaneously with oxygen tension in the solution. Figure 2 plots the bath and cortical oxygen levels ($n = 5$ kidneys, see Methods). It shows that oxygen level in the solution remained in the 140-160 mmHg range, corresponding to atmospheric levels (open symbols). On the other hand, cortical oxygen level was close to zero ($0.1 [0.0;2.5]$ mmHg, $n = 8$ time-points; hatched symbols). This confirms that kidneys were hypoxic, quasi-anoxic, during the *ex vivo* normothermic ischemic phase.

Renal oxygenation parameters in SCS and HMP

After WI, kidneys were preserved for 20 hours under SCS (UW), or HMP (KPS; LifePort®). During the SCS period, 20h-average cortical temperature was 2.9 ± 2.0 °C (mean $\pm$ SD, $n = 1,200$ values): starting and final values were $15.0 [9.3;22.2]$ °C ($n = 5$) and $2.7 [2.4;3.0]$ °C ($n = 5$), respectively, the stable value being reached after 5-6 hours.

During HMP, average cortical temperature was 8.2 ± 1.0 °C, dropping from 13.2 [$11.1;14.4$] °C to 7.8 [$7.7;7.9$] °C ($n = 5$), with an apparent half-time for stabilization of ~5 h to 6.9 ± 0.9 °C.

Renal oxygenation parameters during SCS. Figure 3 (main panel) shows the oxygen level (left axis: partial pressure, mmHg; right axis: concentration, $\mu\text{mol/L}$) in the preservation solution (open circles) and in the renal cortex (hatched circles), as monitored throughout the 20h SCS phase ($n = 5$ kidneys). Oxygen level in the solution, starting from atmospheric-equivalent levels (150-160 mmHg) decreased steadily and significantly ($F\text{-ratio} = 33.13$, $p < 0.05$) to stabilize at 51.7 [$51.4;52.9$] mmHg (last $n = 5$ time-points, between 960 and 1,200 min; half-time of decrease ~90 min).

In the cortex, the oxygen pressure started from 0.3-0.4 mmHg (between 1 and 30 min), close to the ischemic cortical PO_2 (see figure 2).

Since PO_2 markedly decreased in the solution, it was possible to calculate (point-to-point) the instantaneous rate of oxygen concentration decrease, i.e. oxygen consumption, QO_2 . The *initial* rate of oxygen consumption was extracted from the linear phase of PO_2 decrease (10 - 30 min time range; see figure 3, main panel), i.e. as the average of the 10-20 and 20-30 minutes intervals: initial QO_2 was 0.6 [$0.5;0.7$] $\mu\text{mol/min per } 100\text{g}$ ($n = 4$ kidneys, one outlier excluded according to Grubbs' test, GraphPad online). Solution-based QO_2 is plotted in the inset of figure 2, in the 0 to 400 min range.

Renal oxygenation parameters during HMP. Figure 4 plots oxygen levels in solution (open circles) and in renal cortex (hatched circles), during 20h LifePort machine preservation ($n = 5$ kidneys, main panel). Under hypothermic machine perfusion, oxygen level in the perfusion solution remained shortly (10 min) at the atmospheric level (~145

mmHg), then rapidly and significantly ($F = 59.44$, $p < 0.05$) decreased, to a final value of 6.8 [6.6;8.2] mmHg (last $n = 5$ time-points, between 960 and 1,200 min; half-time ~40 min). As for SCS, the *initial* rate of QO_2 in HMP was extracted from the linear phase of PO_2 decrease: solution-based QO_2 is plotted in the inset of figure 4, in the 0 to 400 min range: initial rate was 3.4 [2.2;4.6] $\mu\text{mol/min per } 100\text{g}$ ($n = 5$ kidneys).

In the cortex, oxygen levels started at 10.2 [0;18.9] mmHg (first time-point, 5min) and dropped ($F = 4.79$, $p < 0.05$) to effectively zero afterwards (0.0-0.3 mmHg) within 60 min.

In addition, machine perfusion allows for the determination of the “arterio-venous oxygen consumption”, based arterial and venous oxygen concentrations (as reported by respective O_2 micro-sensors, see Methods) and on perfusion rate (as reported by the HMP device). In the LifePort machine, the renal perfusion rate rapidly increased from 0 to 40 mL/min (in less than 10 min), to slowly stabilize at ~50 mL/min after 600 min; the 20h-average perfusion rate was 45.5 ± 4.5 mL/min (mean $\pm$ SEM-SD, $n = 20$), corresponding to a renal resistance of ~0.7 mmHg.min/mL. The main panel of figure 5 shows the time-dependent evolution of arterio-venous renal QO_2 during 20h HMP: from the [3.4;4.1] $\mu\text{mol/min per } 100\text{g}$ range, renal QO_2 significantly decreased ($F = 38.89$, $p < 0.05$) to reach 0.4 $\mu\text{mol/min per } 100\text{g}$ at 120 min (see inset, zoomed over 0-120 min), then further decreased, very slowly, down to 0.2 $\mu\text{mol/min per } 100\text{g}$. Initial value of arterio-venous (HMP) QO_2 was 3.4-4.1 $\mu\text{mol/min per } 100\text{g}$, NS *versus* solution-based initial consumption (3.4 [2.2;4.6] $\mu\text{mol/min per } 100\text{g}$, see above). Conversely, one-way ANOVA comparison of solution-based SCS (0.6 $\mu\text{mol/min per } 100\text{g}$) and HMP (3.4 [2.2;4.6] $\mu\text{mol/min per } 100\text{g}$) consumption indicated a significant difference ($p < 0.01$).

ATP level of renal cortical tissue. Figure 6 summarizes ATP results. Renal cortical ATP in the control (non-ischemic) group was 3.3 [1.7;4.6] mmol/L of tissue. After 60 minutes of warm ischemia, the ATP level was 0.3 [0.2;0.6] mmol/L ($p < 0.01$ vs control). After 20 hours of SCS, the ATP content of the kidneys was 0.06 [0.04;0.09] mmol/L (NS vs WI), whereas it was 5.8 [3.2;6.5] mmol/L in HMP ($p < 0.01$ vs WI).

Histological analysis. Figure 7a shows representative HES and PAS-colorations of cortical and medullary tissue sections of the four experimental groups. Figure 7b summarizes the corresponding histological scores. It can be seen that WI60 induced significant histological injuries *versus* control (brush border loss, endoluminal detachment, vacuolization and infiltration). All four parameters were significantly improved by HMP *versus* SCS, although not normalized to control (except for tissue infiltration). Finally, brush border loss was potentiated by SCS *versus* WI60 (by about two-fold), and HMP did not restore control level.

4. Discussion

Oxygen supplementation during hypothermic preservation is the focus of preclinical and clinical studies ⁷, but, paradoxically oxygenation parameters are often ignored and left unquantified (i.e. oxygen level and/or consumption). In non-actively oxygenated hypothermic methods (where either static solution is enclosed in a sealed recipient, or perfusate solution is recirculated without possible or significant re-equilibration with atmospheric oxygen), oxygen-related parameters are usually not measured, e.g. because graft requirements are assumed to be minimal.

On the other hand, we reported, with our porcine model of renal autotransplantation that i) the addition of an hemoglobin-based oxygen carrier to SCS preservation does improve

post-transplant outcomes ^{10,15}, and ii) active oxygenation (100% O₂) in HMP improves post-transplantation function recovery of WI-injured kidneys ⁸. In other words, as reviewed ⁷, renal grafts can draw benefit from oxygen under hypothermic preservation. On the other hand, due to the expanding use of suboptimal grafts, HMP preservation has now grown more frequent than the simpler and cheaper SCS preservation, at least in european and north-american countries. But, as for SCS, oxygen-related parameters are most generally ‘ignored’. Some rare preclinical studies addressed oxygen issues, though. For instance, one porcine study promoted the determination of oxygen consumption for viability assessment ⁶. At least in the cortex, oxygen can be considered as a proxy for ATP, the cell energetic currency. However, because of technical and logistical reasons, ATP is even more difficult to monitor. Accordingly, very few studies, mostly preclinical, did measure ATP ^{16,17}.

Thus, the monitoring of renal graft oxygen levels and consumption, together with tissue ATP levels, will help understand graft status and requirements for better, if not optimal, preservation. To address the current clinical context, we used renal grafts injured by 60 minutes warm ischemia (WI) ^{18,19}, UW for SCS, and KPS and the LifePort® system for HMP.

Oxygen is consumed and limiting in non-oxygenated SCS and HMP

In SCS and HMP, PO₂ of the preservation solution decreased rapidly, from atmospheric level, then stabilized. Decreases of perfusion PO₂ in HMP were reported in uninjured (WI < 10min) porcine kidneys ⁶ and (45min WI-injured) canine kidneys ²⁰. However, in the first case data are not presented (*‘...PO₂ of the perfusate approached zero within minutes..’*) ⁶, while in the second, only initial and final PO₂ value are reported (from 170 mmHg at

zero time to 25 mmHg after 24h HMP, LifePort) ²⁰. To our knowledge, oxygen level has not been reported in SCS.

SCS exploratory experiments confirmed that solution PO₂ was stable for 24h (150 mmHg), whereas it decreased and stabilized with a WI-injured kidney (not shown). We conclude that, in hypothermia, injured kidneys do consume oxygen, despite the low temperature and the static condition (no perfusion and no transport-related consumption). Of note, the SCS and the HMP systems used in the present study were open to atmosphere (electrodes, holding arm), which suggests that in clinics oxygen becomes limiting even faster than observed here (< 60-90min). Finally, our conclusion that oxygen is limiting in non-oxygenated SCS and HMP agrees with reports that a short, post-ischemic/pre-transplantation period of active graft oxygenation (1-2h, PO₂ > 500 mmHg) improves kidney function recovery ^{21,22}. This was also shown in HMP or retrograde persufflation of the liver ²³.

In SCS, solution-based oxygen consumption was 0.6 $\mu\text{mol}/\text{min}$ *per* 100g, whereas in HMP, it was 3.4 $\mu\text{mol}/\text{min}$ *per* 100g, the later being not significantly different from the arterio-venous calculation in HMP (av-QO₂ 3.7 $\mu\text{mol}/\text{min}$ *per* 100g). Of note, in HMP, the initial decrease of cortical oxygen (from 10-15 to 0 mmHg), occurred simultaneously with the initial decrease in the solution.

To our knowledge, QO₂ has not been reported in SCS, likely because this was considered irrelevant. Conversely, in HMP QO₂ of Pig kidneys has been reported to amount to ~1,6 $\mu\text{mol}/\text{min}$ *per* 100g (WI < 10min) ⁶, a value close to the present one (3.7 $\mu\text{mol}.\text{min}$ *per* 100g, WI = 60min). Starting from a total physiological oxygen consumption (transport-related + basal ~200 $\mu\text{mol}.\text{min}$ *per* kidney ²⁴), and applying a 20% basal fraction, one obtains ~40 $\mu\text{mol}/\text{min}$ *per* 100g for non-transport related costs. After

temperature-dependent reduction, hypothermic QO_2 would amount to $\sim 5 \mu\text{mol}/\text{min per } 100\text{g}$, very close to and, supporting our determinations. This further supports the notion that, in hypothermic preservation, normal or injured kidneys do consume oxygen. Finally, it also suggests that warm ischemia may not modify basal consumption, although this remains to be established.

In the preservation solution, the low oxygen steady-state indicates that the transfer capacity, *via* the solution and the capsule, to the cortical parenchyme is limited (and limiting) with respect to intrarenal consumption. This observation is supported by exploratory experiments from our group, showing that solution PO_2 not only decreases with time, but that this decrease is more pronounced if electrodes are positioned “close to” *versus* “far from” the kidney: for instance, between 10 and 15 h the ‘close’ electrodes (2-3 cm) reported 6-34 mmHg PO_2 lower than the distant electrodes (5-7 cm; $n = 2$ healthy kidneys and $n = 2$ WI-injured kidneys). This is also in line with the report of a kidney ‘surface PO_2 ’ of 5-10 mmHg in the SCS solution, *versus* 100-150 mmHg in HMP²⁵. Furthermore, we suggest that the significantly higher steady-state values of solution oxygen in SCS (~ 52 mmHg) *versus* HMP (~ 8 mmHg) indicates that HMP kidneys consume more oxygen than SCS kidneys. Indeed, at 35 mmHg perfusion pressure, filtration still occurs, proportional to renal flow (since perfusion pressure is below the auto-regulatory range), roughly amounting to $\sim 30\%$ of normal filtration. This is comforted by published reports of urine production during HMP preservation ($\sim 0.1\text{mL}/\text{min per } 100\text{g}$ ²⁶) and our unpublished observations. A crude estimation indicates that under our HMP conditions, oxygen consumption should range $6-7 \mu\text{mol}/\text{min per } 100\text{g}$, relatively close to our $3-4 \mu\text{mol}/\text{min per } 100\text{g}$ measurement (see above).

ATP level and histological evaluation in HMP and SCS

Beneficial effects of oxygen supplementation in hypothermic preservation are acknowledged, i) in HMP and for high-risk kidneys, and ii) as involving some restoration of ATP ²⁷. More surprisingly, even under SCS conditions, direct (100 % O₂ ²⁸) or indirect (M101 carrier ¹⁰) oxygen supplementation is able to provide improvements as well.

As opposed to the renal medulla, cortical ATP synthesis heavily relies on oxygen and mitochondrial phosphorylation. Normal porcine cortical tissue PO₂ is ~45 mmHg ²⁹. Here, shortly after the start of the WI period (< 5 min), cortical oxygen barely exceeded electrode sensitivity (0.4-0.8 mmHg), even though the solution was in contact with atmosphere. Accordingly, end-ischemic ATP was strongly reduced (nb: control ATP, was similar to other reports ²⁵). ATP levels (after SCS) of “marginal” kidneys (WI 40min) have been reported to amount to ~0.16 μmol/g tissue (or ~0.2 mM), in the Pig ²³; this represents a 80-90% reduction *versus* control ATP, as we observed here.

After SCS of WI-injured kidneys, ATP remained very low, indicating that these grafts were unable to resynthesize ATP, whereas ATP reached back control level after HMP. These results suggest that ATP was being produced during HMP preservation, in excess to consumption. Few studies addressed the ATP production-consumption balance in hypothermic conditions. A careful bibliographical analysis evidenced a patent lack of information about ATP levels in renal graft preservation. Most generally, neither absolute ATP levels nor healthy control value are given (see Bagul and coworkers for a porcine model ³⁰, and Ravaioli et al., for human discarded kidneys¹¹). Nevertheless, supporting our present observations, two porcine studies, performed on did show marked benefits of HMP *versus* SCS in terms of graft ATP levels. First, ATP resynthesis was observed in oxygenated HMP but not in SCS, in a WI-injured graft model ³¹. Second,

a 2.5-fold ATP post-HMP increase was reported *versus* post-SCS in a WI-injured kidneys (≤ 30 min) model ³². In HMP, the “steady-state” solution oxygen was lower than in SCS, whereas, end-preservation ATP was higher in HMP.

Finally, considering tissue integrity outcomes, by showing less injured tissues, histological analysis confirmed HMP superiority over SCS, in agreement with the increased oxygen consumption and higher levels of ATP. It is important to note though that the ATP ‘restoration’ may not be the sole factor involved in renal tissue “repair” during hypothermic preservation since SCS itself was able to partially improve the histological profile.

Conclusion

Grafts under SCS or HMP preservation are rarely biopsied, while three-months and one-year post-transplantation biopsies are common in clinical practice, allowing for histological analysis and grading. *Ex vivo* quality assessment of renal grafts is limited to visual inspection, before and after flushing (color, patchiness), and to renal resistance in HMP. Oxygen monitoring requires a dedicated setup, and no current perfusion system is equipped with oxygen sensors ⁷. Even when actively oxygenated, no available system allows for oxygen measurements. Thus, oxygen and ATP are seldom measured, much less comparatively in SCS and HMP. Oxygen measurement (and consumption calculation) on the one hand, and high-resolution NMR spectroscopy or electrode-based determinations of tissue ATP, as alternatives, on the other hand, would constitute significant improvements, widening the currently restricted panel of clinical evaluation criteria with physiologically relevant and quantitative indexes ³³.

5. Acknowledgments

This work was performed in the U1082 INSERM (Institut National de la Santé et de la Recherche Médicale) research unit (dir. Pr T. Hauet), by the Centre Hospitalo-Universitaire (CHU) de Poitiers, and by the Université de Poitiers. It was supported by the CNRS (Centre National de la Recherche Scientifique) for PH salarial cost. JK was supported by a grant from the ANR (2011, RPIB-013, HemoPerf).

We are indebted to Nicolas Chatauret for the initial setting up of the SCS and HMP oxygen measurements, and for helpful discussion. We acknowledge Abdesalam Kasil's help with animal surgery and kidney withdrawal and expert histological slides preparation by Virginie Ameteau. We are grateful to Dr Jérôme Roumy for interest and discussion. Finally, we are greatly indebted to the surgical platform MOPICT (INRA, Surgères) and members for excellent and expert technical support: William Hébrard and Irène Launay (INRA, Surgères), and Pierre Couturier (CHU, Service de Biochimie).

6. Conflict of interest statement

No conflict of interest, financial or otherwise are declared by all authors.

7. References:

1. Jochmans, I., Nicholson, M. L. & Hosgood, S. A. Kidney perfusion: Some like it hot others prefer to keep it cool. *Curr. Opin. Organ Transplant.* **22**, 260–266 (2017).
2. Golbo, N. & Monbaliu, D. Temperature and oxygenation during organ preservation: Friends or foes? *Curr. Opin. Organ Transplant.* **22**, 290–299 (2017).
3. Jochmans, I. *et al.* Past, Present, and Future of Dynamic Kidney and Liver Preservation and Resuscitation. *Am. J. Transplant.* **16**, 2545–2555 (2016).
4. Folkert O. Belzer, J. H. S. Principles of solid-organ preservation by cold storage. *Transplantation* **45**, 673–676 (1988).
5. Stubenitsky, B. M., Ametani, M., Danielewicz, R., Southard, J. H. & Belzer, F. O. Regeneration of ATP in kidney slices after warm ischemia and hypothermic preservation. *Transpl. Int.* **8**, 293–297 (1995).
6. Weegman, B. P. *et al.* Continuous real-time viability assessment of kidneys based on oxygen consumption. in *Transplantation Proceedings* **42**, 2020–2023 (2010).
7. O’Callaghan, J. M. *et al.* Supplemental oxygen during hypothermic kidney preservation: A systematic review. *Transplant. Rev.* **31**, 172–179 (2017).
8. Thuillier, R. *et al.* Benefits of active oxygenation during hypothermic machine perfusion of kidneys in a preclinical model of deceased after cardiac death donors. *J. Surg. Res.* **184**, 1174–1181 (2013).

9. Hosgood, S. A., Nicholson, H. F. L. & Nicholson, M. L. Oxygenated kidney preservation techniques. *Transplantation* **93**, 455–459 (2012).
10. Thuillier, R. *et al.* Supplementation with a new therapeutic oxygen carrier reduces chronic fibrosis and organ dysfunction in kidney static preservation. *Am. J. Transplant.* **11**, 1845–1860 (2011).
11. Ravaioli, M. *et al.* Strategies to restore adenosine triphosphate (ATP) level after more than 20 hours of cold ischemia time in human marginal kidney grafts. *Ann. Transplant.* **23**, 34–44 (2018).
12. Favreau, F. *et al.* Anti-thrombin therapy during warm ischemia and cold preservation prevents chronic kidney graft fibrosis in a DCD model. *Am. J. Transplant.* **10**, 30–39 (2010).
13. Allain, G. *et al.* Development of a preclinical model of donation after circulatory determination of death for translational application. *Transplant. Res.* **3**, 13 (2014).
14. Tillet, S. *et al.* Kidney graft outcome using an anti-Xa therapeutic strategy in an experimental model of severe ischaemia-reperfusion injury. *Br. J. Surg.* **102**, 132–142; discussion 142 (2015).
15. Mallet, V. *et al.* Dose-ranging study of the performance of the natural oxygen transporter HEMO2life in organ preservation. *Artif. Organs* **38**, 691–701 (2014).
16. Buchs, J. B. *et al.* DCD Pigs' Kidneys Analyzed by MRI to Assess Ex Vivo Their Viability. *Transplantation* **97**, 148–153 (2014).
17. Minor, T., Efferz, P. & Lüer, B. Hypothermic reconditioning by gaseous oxygen persufflation after cold storage of porcine kidneys. *Cryobiology* **65**, 41–44 (2012).

18. Hosgood, S. A., Shah, K., Patel, M. & Nicholson, M. L. The effect of prolonged of warm ischaemic injury on renal function in an experimental ex vivo normothermic perfusion system. *J. Transl. Med.* **13**, (2015).
19. Delpech, P. O. *et al.* Inhibition of complement improves graft outcome in a pig model of kidney autotransplantation. *J. Transl. Med.* **14**, (2016).
20. Lindell, S. L., Muir, H., Brassil, J. & Mangino, M. J. Hypothermic Machine Perfusion Preservation of the DCD Kidney: Machine Effects. *J. Transplant.* **2013**, 1–7 (2013).
21. Hoyer, D. P. *et al.* Influence of oxygen concentration during hypothermic machine perfusion on porcine kidneys from donation after circulatory death. *Transplantation* **98**, 944–950 (2014).
22. Kron, P. *et al.* Short, cool, and well oxygenated - HOPE for kidney transplantation in a rodent model. *Ann. Surg.* **264**, 815–822 (2016).
23. Minor, T., Von Horn, C. & Paul, A. Role of temperature in reconditioning and evaluation of cold preserved kidney and liver grafts. *Curr. Opin. Organ Transplant.* **22**, 267–273 (2017).
24. Juillard, L. *et al.* Validation of renal oxidative metabolism measurement by positron-emission tomography. in *Hypertension* **50**, 242–247 (2007).
25. Minor, T., Sitzia, M. & Dombrowski, F. Kidney transplantation from non-heart-beating donors after oxygenated low-flow machine perfusion preservation with histidine – tryptophan – ketoglutarate solution. *Transplant. Int.* **17**, 707–712 (2005).

26. Blum, M. F. *et al.* Comparison of normothermic and hypothermic perfusion in porcine kidneys donated after cardiac death. 35–45 (2018). doi:10.1016/j.jss.2017.04.008.Comparison
27. Taylor, M. J. & Baicu, C. Current State of Hypothermic Machine Perfusion Preservation of Organs : The Clinical Perspective. *Cryobiology* **60**, S20–S35 (2010).
28. Maathuis, M. H. J. *et al.* Improved kidney graft function after preservation using a novel hypothermic machine perfusion device. *Ann. Surg.* **246**, 982–989 (2007).
29. Warner, L. *et al.* Regional decreases in renal oxygenation during graded acute renal arterial stenosis: a case for renal ischemia. *Am J Physiol Regul Integr Comp Physiol* **296**, R67-71 (2009).
30. Bagul, A. *et al.* Experimental renal preservation by normothermic resuscitation perfusion with autologous blood. *Br. J. Surg.* **95**, 111–118 (2008).
31. Lazeyras, F. *et al.* Detection of ATP by ‘in line’³¹P magnetic resonance spectroscopy during oxygenated hypothermic pulsatile perfusion of pigs’ kidneys. *Magn. Reson. Mater. Phys. Biol. Med.* **25**, 391–399 (2012).
32. La Manna, G. *et al.* An in vivo autotransplant model of renal preservation: Cold storage versus machine perfusion in the prevention of ischemia/reperfusion injury. *Artif. Organs* **33**, 565–570 (2009).
33. Kerforne, T. & Saint-yves, T. Inhibition of coagulation proteases Xa and IIa decreases ischemia – reperfusion injuries in a preclinical renal transplantation model. 95–107 (2016). doi:10.1016/j.trsl.2016.07.014

34. Willford, D. C., Hill, E. P. & Moores, W. Y. THEORETICAL ANALYSIS OF OXYGEN TRANSPORT DURING HYPOTHERMIA David C. Willford, PhD,* Esther P. Hill, PhD,* and William Y. Moores, MD'p. *J Clin Monit* **2**, 30–43 (1986).

8. Figure and legends to figures

Figure 1: Protocol flow chart

Figure 2: Time-course of oxygen in the solution and cortex during warm ischemia

Figure 3: Renal oxygenation during 20 hours SCS - Oxygen consumption.

Figure 4: Renal oxygenation during 20 hours HMP preservation – Oxygen consumption

Figure 5: Arterio-venous consumption of oxygen during 20 hours HMP

Figure 6 : ATP in the renal cortex, in Control, End-WI, End-SCS and End-HMP

Figure 7a : HES and PAS colorations of Control, End-WI, End-SCS and End-HMP

Figure 7b : Semi-quantitative histological scoring of renal tissue sections

Legend to figure 1: Protocol flow chart

Legend to figure 2: Figure depicts the time-course of oxygen levels in the solution and in the cortex during the 60 minutes warm ischemia (median and [range], n = 5). Monitoring of renal temperature and oxygen level was performed with electrodes inserted into the cortex (3-5 mm depth). Open circles represent oxygen pressure in the solution; hatched symbols correspond to levels of oxygen pressure in the renal cortex.

Legend to figure 3: Figure shows the time-course of oxygen levels in the solution and in the cortex during 20 hours of SCS preservation (median and [range], n = 5): open symbols, solution; hatched symbols, cortex (inset). The left y-axis represents oxygen partial pressure, in mmHg, whereas the right one represents the corresponding oxygen concentration, in μM (calculated using a temperature-adjusted oxygen solubility, see Methods). The decrease in the solution features at least two components but this was not further analyzed. The apparent half-time of the decrease is ~ 90 min; the final, steady-state value is 51.7 [$51.4;52.9$] mmHg (n = 5 time-points, between 960 and 1,200 min). Average temperature (20h) was 2.9 ± 2.0 °C (mean $\pm$ SD, n = 1,200 values, see Results for details); it was $15\text{-}8$ °C during the first 60 min (corresponding to a temperature-dependent 5-8 fold reduction of reaction rates, assuming $Q_{10} = 2$, see ³⁴). The inset shows the oxygen consumption (based on initial decrease of oxygen in the solution), plotted between 0-420 min (0-7h); initial rate was in the $0.5\text{-}0.7$ $\mu\text{mol}/\text{min}$ *per* 100g.

Legend to figure 4: Figure plots oxygen levels in solution (open circles, $n = 5$) and in renal cortex (hatched circles, $n = 5$), during 20 hours HMP (LifePort®). The left y-axis represents PO_2 (mmHg), while the right one represents corresponding concentration (μM , see Methods). The apparent half-time of the decrease is ~ 40 min, and the steady-state value is 6.8 [6.6;8.2] mmHg. Time-averaged temperature (20h) was 8.2 ± 1.0 °C (mean $\pm$ SD); it was ~ 12 °C during the first 60 min (~ 6 -fold reduction of reaction rates). The inset plots the oxygen consumption (calculated from initial decrease of oxygen in the solution) between 0-400 min, which is in the range 2.2-4.6 $\mu mol/min$ *per* 100g.

Legend to figure 5: Figure plots the arterio-venous consumption in HMP preservation, calculated as the product of renal perfusion rate by the arterio-venous oxygen difference (see Methods). Please note that the vertical axis is not *oxygen* (as is in figure 2, 3 and 4) but *oxygen consumption*. Initial renal graft consumption in HMP was [3.4;4.1] $\mu\text{mol/min per } 100\text{g}$. The inset represents the same results, zoomed over the first 120 minutes

Legend to figure 6: ATP concentration in renal cortical tissue (median and [range], n = 5 *per* group). Control was measured before warm ischemia, while end-ischemic ATP was measured at the end of the 60 minutes period of warm ischemia (End-WI). End-SCS and End-HMP, correspond to renal cortical ATP of kidneys submitted to 60 min WI followed by 20 hours SCS and 20 hours HMP preservation, respectively. ATP is given as mmol/L of tissue (conservatively equivalent to μ moles ATP *per* g wet tissue).

Legends to figure 7a and 7b

Figure 7a: Representative HES and PAS colorations of renal tissue, split into cortical (upper lines of each quadrant) and medullary (lower lines) areas. Each quadrant is further split vertically, into left and right columns, respectively showing HES and PAS staining.

Arrows exemplify: brush border loss (red), endoluminal detachment (blue), vacuolization (black), and interstitially infiltrated cells (yellow). The black bar represents 100 μm .

Figure 7b: Semi-quantitative histological scoring (see Methods). Panel **A**, Brush border loss scores (upper left); panel **B**, Endoluminal detachment scores (upper right); panel **C**, Vacuolization scores (lower left); panel **D**, Infiltration (lower right). Statistics performed as ANOVA one-way, with Tukey's post-test for multiple comparisons: *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ns, not significant.