Mitochondrial complex III Qi -site inhibitor resistance mutations found in laboratory selected mutants and field isolates

Pierre Mounkoro, Thomas Michel, Rafik Benhachemi, Georgiana Surpateanu, Bogdan I. Iorga, Nicholas Fisher, Brigitte Meunier

► To cite this version:

Pierre Mounkoro, Thomas Michel, Rafik Benhachemi, Georgiana Surpateanu, Bogdan I. Iorga, et al.. Mitochondrial complex III Qi -site inhibitor resistance mutations found in laboratory selected mutants and field isolates. *Pest Management Science*, 2019, 75 (8), pp.2107-2114. 10.1002/ps.5264 . hal-02182323

HAL Id: hal-02182323

<https://hal.science/hal-02182323>

Submitted on 6 Jul 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Mitochondrial complex III Q_i-site inhibitor resistance mutations found in laboratory selected mutants and field isolates.

Pierre Mounkoro¹, Thomas Michel¹, Rafik Benhachemi^{1,a}, Georgiana Surpateanu², Bogdan I. Iorga², Nicholas Fisher³, Brigitte Meunier^{1*}

¹ Institute for Integrative Biology of the Cell (I2BC), CEA, CNRS, Université Paris Sud, Université Paris Saclay, Gif sur Yvette, France.

² Institut de Chimie des Substances Naturelles, CNRS, UPR 2301, Université Paris Saclay, Labex LERMIT, Gif-sur-Yvette, France

³ MSU-DOE Plant Research Laboratory, Michigan State University, East Lansing, MI 48824, USA

* Correspondence to: Brigitte Meunier, I2BC, avenue de la Terrasse, 91198, Gif sur Yvette, France. E-mail : brigitte.meunier@i2bc.paris-saclay.fr

^a current address: Laboratory of food microbiology, KULeuven, Belgium

Running title: Target site QiI resistance mutations

Keywords: QiI, bc₁ complex, yeast model, resistance, target site mutation, fungicides

Abstract

BACKGROUND: Complex III inhibitors targeting the Q_i-site have been known for decades; some are used or are being developed as anti-microbial compounds. Target site resistance mutations have been reported in laboratory-selected mutants and in field isolates. Here we presented a brief over-view of mutations found in laboratory selected resistant mutants; and a study of mutations observed in field isolates of *Plasmopara viticola*, in particular the ametoctradin resistance mutation, S34L that we analyzed in the yeast model.

RESULTS: The survey of laboratory mutants showed that resistance could be caused by a large number of substitutions in the Q_i-site. Four residues seemed key in term of resistance, N31, G37, L198 and K228. Their mutations caused resistance to several QiIs, in different organisms with, in yeast, no or moderate loss of complex III function. Two mutations have been reported in the field, in *P.viticola* resistant to either cyazofamid or ametoctradin. Using the yeast model, we analyzed the effect of the ametoctradin resistance mutation S34L. We showed that S34L caused a high level of resistance combined with a loss of complex III activity, resulting in a decreased growth competence.

CONCLUSION: The use of QiIs that target a single site encoded by a mitochondrial gene, presumably prone to mutations is expected to result in the selection of resistant mutants. However if the mutation is associated with a fitness penalty, as it seems the case with S34L, the development of the resistance might not be an insuperable obstacle but careful monitoring is required.

1 INTRODUCTION

The respiratory chain complex III (or *bc*₁ complex) with its two quinol binding sites, termed the Q_o- and Q_i-sites, is a proven target for anti-microbial compounds. The first generation of antimicrobial drugs targeted the Q_o-site: the antimalarial atovaquone and a large series of fungicides used against plant pathogen fungi, such as azoxystrobin.¹ Target site resistance mutations have often been reported, for instance Y279C/S in *Plasmodium falciparum* (see for example ^{2,3}) or G143A in many species of plant pathogen fungi that confers cross-resistance to all the >20 Q_o-site targeting fungicides (termed QoIs) on the market (see FRAC web site). The structural basis of the resistance has been well studied.^{4,5}

New generation of inhibitors, still in development, target the Q_i-site, such as the anti-malarial ELQs⁶ and the agrochemical fungicide fenpicoxamid.⁷ The only Q_i-site inhibitors

(or QiIs) already on the market are cyazofamid⁸ and amisulbrom, used in field against plant pathogen oomycetes (see www.frac.info). A third anti-oomycete in use, ametoctradin, was listed as QoSI (see www.frac.info) i.e. binding at the Q_o-site in a manner similar to the Q_o-site inhibitor stigmatellin but distinct to QoIs.⁹ On the basis of further spectroscopic studies, the compound would target both Q_o- and Q_i- sites.¹⁰ Other Q_i-site inhibitors have been known for a long time, such as funiculosin, diuron, HQNO and antimycin, or described more recently, such as ilicicolin H¹¹ or HDQ.¹² These compounds have proven useful in structural and mechanistic studies of complex III.

Q_i-site mutations conferring resistance (or suspected to cause resistance) to inhibitors have been observed after treatment of pathogens in laboratory assays or in the field, and also through the use of model organisms (mainly *Saccharomyces cerevisiae*).

Here, we briefly review these reported Q_i-site mutations and study (using the yeast model) a mutation (S34L) found in the plant pathogen oomycete *Plasmopara viticola* field isolates after ametoctradin treatment.¹³

2 MATERIALS AND METHODS

2.1 Materials and growth media

Ametoctradin was kindly provided by BASF. The following media were used for yeast growth: YPD (1% yeast extract, 2% peptone, 3% glucose), YPG (1% yeast extract, 2% peptone, 2% glycerol), YPEth (1% yeast extract, 2% peptone, 2% ethanol), YPGal (1% yeast extract, 2% peptone, 0.2% glucose, 3% galactose)

2.2 Yeast strains

Strains derived from AD1-9 that lacks several membrane transporters (α ura3 his1, yor1 Δ ::hisG, snq2 Δ ::hisG, pdr5 Δ ::hisG, pdr10 Δ ::hisG, pdr11 Δ ::hisG, ycf1 Δ ::hisG,

pdr3 Δ ::hisG, pdr15 Δ ::hisG, pdr1 Δ ::hisG) and kindly provided by M. Ghislain, UCL, Belgium. The mutations were introduced in yeast cytochrome *b* as described in ^{14,15}. In all experiments, control and mutants have identical nuclear and mitochondrial genomes with the exception of the cytochrome *b* mutations.

2.3 O₂ consumption activity of cells

Activities were measured using a Clarke-type oxygen electrode. Cells were grown in YPGal medium for 24h. The cells were then harvested. O₂ consumption rates in presence of 4 μ M CCCP were recorded for two minutes. Ametoctradin was then added and the rate measured after two minutes. The data, normalised per cell count, are presented as % of control rates (without ametoctradin). The measurements were repeated at least three times and averaged.

2.4 Complex III activity

Activities were determined spectrophotometrically by measuring the reduction of 20 μ M cytochrome *c* at 550 *minus* 540 nm over 1-minute time-course in 1 ml of 10 mM potassium phosphate pH 7.0, 0.01% (w/v) lauryl-maltoside and 1 mM KCN. Mitochondria prepared as in¹⁶ were added to obtain a final concentration of 2 - 10 nM complex III. Complex III concentration was determined from dithionite-reduced optical spectra, using ϵ =28.5 mM⁻¹.cm⁻¹ at 562nm *minus* 575nm. The reaction was initiated by the addition of 20 μ M decylubiquinol and initial rates were measured. Each measurement was repeated three to five times and the values obtained were averaged. Complex III activity was determined as the cytochrome *c* reduction rate per complex III per second. Mid-point inhibition concentrations were determined by inhibitor titration.

2.5 Molecular modelling

1
2
3 100 *In silico* model of *P.viticola* Q_i-site was built by comparative modelling using MODELLER
4
5 101 9v7¹⁷, with the yeast structure (PDB 3CX5) as template. Three-dimensional structure of
6
7 102 ametoctradin was generated using CORINA (Molecular Networks GmbH, Erlangen,
8
9 103 Germany, <http://www.molecular-networks.com>). The docking of ametoctradin on the
10
11 104 *P.viticola* model of the Q_i-site was performed with GOLD version 5.0.¹⁸ The binding site was
12
13 105 defined as a sphere with a 15 Å radius around the antimycin in the PDB structure 1PPJ.
14
15 106 GoldScore was used as a scoring function and all other parameters had default values. UCSF
16
17 107 CHIMERA¹⁹ was used for generating the molecular modelling images.
18
19
20
21

22 109 **3 RESULTS AND DISCUSSION**

23
24 110 **3.1 Laboratory selected mutants with Q_i-site mutations causing inhibitor resistance: An**
25
26 111 **over-view**

27
28 112 The Q_i-site is formed from the cytochrome *b* C-terminal region of surface helix a, the N-
29
30 113 terminal region of transmembrane helices A and E and the C-terminal regions of helix D
31
32 114 (Fig.1).
33
34 115 Mutations reported to cause (or suspected to cause) inhibitor resistance have been found at
35
36 116 positions I17, S20, Q22, I27, W30, N31, G33, S34, L35, G37, L198, H204, M221, F225,
37
38 117 K228, G232 (Table 1). Note that we used *S.cerevisiae* residues and positions all through
39
40 118 except if specified.
41
42
43

44 119 Of these, four positions seem to be particularly important in term of inhibitor
45
46 120 resistance, namely N31, G37, L198 and K228. Mutations of these residues have been reported
47
48 121 in different studies and organisms and/or after selection with different inhibitors. N31K have
49
50 122 been found in *S.cerevisiae* and *Kluyveromyces lactis* after selection on the structurally-related
51
52 123 antimycin and fenpicoxamid, but also on the unrelated compound diuron, (or DCMU). L198F
53
54 124 causes resistance to antimycin, fenpicoxamid, ilicicolin H and funiculosin in *S.cerevisiae*;
55
56
57
58
59
60

1
2
3 125 resistance to antimycin and GNF7686 in *Trypanosoma cruzi* the causative agent of Chagas
4
5 126 disease. K228M confers resistance to antimycin in *S.cerevisiae* and *K.lactis*. Replacement of
6
7 127 the threonine by a proline at the equivalent position in *Toxoplasma gondii* causes resistance to
8
9 128 antimycin and ELQ-316. The most striking are mutations of residue G37. Five substitutions
10
11 129 have been reported: G37A,C,D,S,V. Acquired resistance mutations at that position have been
12
13 130 found in fungi *S.cerevisiae*, *Schizosaccharomyces pombe* and *Zymoseptoria tritici*, in the
14
15 131 human malaria parasite *P.falciparum* and in mouse cells in culture. They confer resistance to
16
17 132 antimycin and fenpicoxamid, diuron or the macrocyclic compound ML238 and its derivative,
18
19 133 BRD6323.

20
21
22 134 The residues N31, G37, L198 and K228, might have a key role for the binding of
23
24 135 inhibitors and accommodate substitutions without a severe loss of complex III activity. In
25
26 136 yeast, the decrease in complex III activity resulting from the substitution L198F was only of
27
28 137 10%²⁰, and of around 50%, for G37S, K228M and N31S^{12,21}, with no or mild effect on
29
30 138 respiratory growth competence. G37 and K228 are very highly conserved residues, although
31
32 139 N31 is often replaced by glycine in plants and isoleucine in prokaryotic cytochrome *b*
33
34 140 sequence data. N31 does not show binding interactions with bound quinone in the atomic
35
36 141 structure of yeast complex III while L198 forms a stabilizing hydrophobic interaction with the
37
38 142 benzoquinone headgroup of bound quinone (PDB 1EZV), and presumably a similar class of
39
40 143 interaction is maintained in the L198F substituent. The sidechain amino group of K228 has
41
42 144 been suggested to be involved in proton transfer to bound quinone during redox chemistry at
43
44 145 the Q_i-site.²² As the mutation of this residue to methionine is well tolerated by yeast, it is
45
46 146 likely that nearby protonatable residues H202 and D229 can rescue.

47
48
49
50 147 Inhibitor resistant mutants harbouring mutations at position N31, G37, L198 and K228
51
52 148 have been selected (and maintained) in laboratory conditions. In the field, a moderate loss of
53
54 149 complex III activity, such as observed with the yeast G37S, K228M and N31S mutants, may
55
56
57
58
59
60

1
2
3 150 result in decreased organismal fitness, which could be protective against resistance evolution
4
5 151 or at least delay resistance evolution. Nevertheless the observed cross-resistance to unrelated
6
7 152 compounds caused by mutations of those residues calls for a close monitoring when anti-
8
9 153 microbial Q_i-site targeting compounds will be on the market. On the other hand, monitoring
10
11 154 of resistance might –and likely will- detect other possible resistance mutations. Studies with
12
13 155 model organisms and laboratory strains would come useful, as illustrated above especially for
14
15 156 the ametoctradin resistance mutation, S34L.
16
17
18 157

19 20 158 **3.2 Anti-oomycetes resistance mutations found in the fields and the study in the yeast** 21 22 159 **model**

23 24 160 **3.2.1 Cyazofamid resistance mutations**

25
26 161 The cyazofamid resistance mutation, observed in field isolates of *P. viticola* is intriguing. It is
27
28 162 a short sequence duplication of six nucleotides resulting in the insertion of two residues E203-
29
30 163 DE-V204.²³ The Q_i-site region between the conserved residues H202 and S206 differs
31
32 164 between species (Fig.2). However, structures of the enzyme with quinone bound at the Q_i-site
33
34 165 showed that the conserved residues H202 and S206 act as hydrogen bond donors to the
35
36 166 benzoquinone headgroup, orientating the quinone correctly. Mutations H202F,N and S206L
37
38 167 resulted in growth defect (unpublished data and ²⁴). Therefore it would not be unexpected that
39
40 168 a major modification of that short region between H202 and S206 could have a deleterious
41
42 169 effect the on respiratory competence.
43
44
45

46 170 In order to test such effect, we replaced, in yeast cytochrome *b*, the sequence I203-
47
48 171 H204 by the sequence E-D-E-V (Fig.2). We observed that the mutation had no effect on
49
50 172 respiratory growth, measured as the cell density reached after three day cultures in YPEth (not
51
52 173 shown), which indicated that the region could accommodate such modifications without a
53
54 174 severe loss of function. Unfortunately, as yeast complex III is insensitive to cyazofamid, the
55
56
57
58
59
60

effect of the mutation on the fungicide sensitivity could not be analysed in our model. A cyazofamid binding model has been reported.²⁵ It can be hypothesized that the mutation E203-DE-V204 (Fig.2) found in *P.viticola* would affect the compound binding or access into the Q_i-site. The insertion is in the linker region between helices D and E (Fig.1), which is proximal to the benzoquinone headgroup of Q_i-site bound quinone (PDB 1EZV), at approximately 7 Å separation at the closest approach. This is likely to perturb the local fold around this region, and/or interfere directly with cyazofamid binding.

In order to pursue the study of the effect of that mutation on complex III activity and reactivity to cyazofamid, a yeast mutant sensitive to the compound needs to be found. It might be possible to generate such mutant by genetically replacing the yeast residues of and around the Q_i-site by the oomycetes equivalents, providing the multiple changes would not cause a too severe loss of function.

3.2.2 The ametocetradin resistance mutation, S34L

Ametocetradin, first listed as a QoSI, was shown to be a dual Q_o- and Q_i-site binding compound based on spectroscopic analysis.¹⁰ The approach, however, could not determine whether the compound would bind at the Q_o- and Q_i-sites with similar affinity. If ametocetradin affinity for one of the sites is much higher than its affinity for the other, resistant isolates harbouring target site mutation could be selected. Indeed a mutation in the Q_i-site, S34L, has been reported in *P. viticola* after treatment¹³, suggesting that ametocetradin have *in vivo* a higher affinity for the Q_i-site. In order to confirm the effect of S34L on ametocetradin resistance, we studied the mutation in the yeast model.

Yeast complex III is naturally highly resistant to ametocetradin (and to the two other anti-oomycetes compounds amisulbrom and cyazofamid). As we have previously shown¹⁰, the respiratory growth of the WT strain can be inhibited by a high dose (300 µM) of

1
2
3 200 ametoctradin which binds at the Q_o-site, and not at the Q_i-site, as Q_o-site mutations I147V and
4
5 201 L275F increased the resistance. Thus, as expected, the introduction of S34L in yeast Q_i-site
6
7 202 had no effect on the ametoctradin sensitivity (not shown).
8

9 203 However a mutant highly sensitive to ametoctradin, namely PFQi3, was identified in a
10
11 204 collection of yeast mutants carrying point and multiple mutations in the Q_i-site. PFQi3
12
13 205 complex III was inhibited by ametoctradin in the low μ M range and behaved in a similar
14
15 206 manner to the oomycete enzyme in inhibitor binding assays.¹⁰ PFQi3 seemed thus a pertinent
16
17 207 model to study the effect of S34L. In addition of PFQi3, we tested here two other mutants,
18
19 208 PFQi1 and PFQi2 (PFQi2 was already described in ²⁶). These mutants, and three other mutant
20
21 209 strains described below, namely PFQi8, PFQi9 and PFQi10, combined several modifications
22
23 210 at the Q_i-site (Table 2).
24
25

26 211 First we tested the ametoctradin sensitivity and resistance of the respiratory growth of
27
28 212 the PFQi1,2 and 3. To that end, cell suspensions were spotted on respiratory medium
29
30 213 containing increasing concentrations of ametoctradin and the plates were incubated for three
31
32 214 to four days (Fig.3). PFQi3 was highly sensitive to ametoctradin as the respiratory growth was
33
34 215 fully inhibited at 1 μ M while the growth of PFQi2 and PFQi1 was blocked at 10 μ M and 100
35
36 216 μ M, respectively.
37
38

39 217 PFQi1 harbours six mutations in the Q_i-site; PFQi2 has two additional changes and
40
41 218 PFQi3 has two more changes and the substitution L198F instead of L198I present in PFQi1
42
43 219 and 2 (Table 2). L198F has been reported as resistance mutation (Table 1). We considered
44
45 220 whether that change could be responsible for the increased sensitivity of PFQi3. Therefore
46
47 221 mutant PFQi8 was produced that combined the six mutations of PFQi1 with L198F. As
48
49 222 shown in Fig.3, PFQi8 was also highly sensitive to ametoctradin, suggesting that L198F could
50
51 223 stabilize ametoctradin in the modified Q_i-site.
52
53
54
55
56
57
58
59
60

224 We then introduced the mutation S34L in PFQi1 and PFQi3. As shown in Fig.3, the
225 resulting mutants, PFQi9 and 10, were resistant to ametoctradin. By comparison, we
226 monitored the sensitivity of the PFQi strains to cyazofamid and amisulbrom. The strains
227 remained fully insensitive to cyazofamid (not shown) while a moderate increase in sensitivity
228 to amisulbrom could be observed with PFQi1 and 2. Their growth was inhibited at 100 μ M
229 amisulbrom (Fig.4). Interestingly, the substitution L198F that renders the cells more sensitive
230 to ametoctradin seems to have the opposite effect with amisulbrom as the respiratory growth
231 of PFQi3 and PFQi8 was not inhibited at 100 μ M amisulbrom. This suggests that the three
232 anti-oomycete compounds would not share the same binding mode in the Q_i-site. However the
233 introduction of the ametoctradin resistance mutation S34L in PFQi1 (Fig.4, mutant PFQi9)
234 conferred also resistance to amisulbrom, suggesting some overlap between ametoctradin and
235 amisulbrom binding sites. However to pursue the study of amisulbrom sensitivity and
236 resistance, other yeast mutants with modified Q_i-site highly reactive to the compound would
237 be required.

238 PFQi10, while clearly resistant to ametoctradin as compared to PFQi3, presented a
239 weaker respiratory growth as seen on YPG plate without inhibitor (Figs. 3 and 4), suggesting
240 a deleterious impact of S34L on the respiratory competence. In order to confirm the effect of
241 S34L, we first monitored the respiratory growth in YPEth liquid cultures with or without
242 ametoctradin at increasing concentrations. We compared PFQi1, PFQi9, PFQi3 and PFQi10.
243 The cell densities reached after three day culture in absence of ametoctradin are shown in
244 Fig.5A. PFQi9 presented a moderate defect as compared to PFQi1 (that reached the same cell
245 density than control (not shown)). The growth yield of PFQi3 was slightly decreased whereas
246 a severe defect was observed for PFQi10. Thus the introduction of S34L in PFQi1 and in
247 PFQi3 had an impact on the growth competence, which was more deleterious for PFQi3 /
248 PFQi10.

249 Growth assays in presence of ametoctradin confirmed that S34L conferred resistance.
250 A three-fold increase in resistance was observed in PFQi9 as compared to PFQi1. Mid-point
251 inhibitory concentrations were around 15 and 45 μM for PFQi1 and PFQi9, respectively
252 (Fig.5B). A much more dramatic difference was observed between PFQi3 and PFQi10. Mid-
253 point inhibitory concentration was around 0.3 μM for PFQi3 while PFQi10 growth was not
254 affected at 20 μM ametoctradin (Fig.5C). The high level of resistance of PFQi10 was further
255 confirmed by monitoring the rates of O_2 consumption of cells in presence of increasing
256 concentrations of ametoctradin. The mid-point inhibitory concentration obtained for PFQi3
257 was around 0.7 μM while PFQi10 respiration could not be inhibited at 50 μM ametoctradin
258 (not shown).

259 Finally we prepared mitochondria and monitored complex III activity (measured as
260 cytochrome *c* reduction rates) as a function of ametoctradin concentration. As expected,
261 PFQi3 complex III was sensitive and 50% inhibition was reached at around 15 μM
262 ametoctradin while PFQi10 complex III was highly resistant as only 10-15% inhibition was
263 observed at 100 μM , in agreement with the growth and O_2 consumption assays.

264 The deleterious effect of S34L was also clearly observed in the enzymatic assays.
265 Complex III activities of control, PFQi3 and PFQi10, in absence of ametoctradin, were
266 compared (Fig.6). PFQi3 enzyme had a two-fold decreased activity as compared to control. In
267 PFQi10, a further two-fold decrease as compared to PFQi3 was observed. Thus the
268 introduction of S34L in the Q_i -site of PFQi3 impaired the activity of complex III. That loss of
269 activity can explain the observed growth defect. The remaining 20-25% complex III activity
270 in PFQi10 (as compared to control) could only support a weak respiratory growth while the
271 50% activity in PFQi3 had only a mild effect on respiratory growth competence. This was in
272 agreement with our previous observations with cytochrome *b* mutants. For comparison, we

also tested the effect of S34L introduced in unmodified Q_i-site on complex III activity. The same assay was performed. We observed 25% loss of activity (not shown).

To summarize, the assays performed here showed that S34L conferred a high level of ametoctradin resistance when introduced in the most sensitive surrogate model, PFQi3. The resistance was associated with a severe loss of complex III activity and respiratory growth competence. Introduction of S34L in the unmodified (and naturally highly resistant) complex III resulted in a mild decrease in activity. S34L introduced in PFQi1 that presented a moderate ametoctradin sensitivity caused a moderate loss of respiratory competence.

Finally, *in silico* docking of ametoctradin into a homology model of the Q_i-site of *P.viticola* cytochrome *b* was performed (Fig.7). The analysis of the docking results suggests a binding mode similar to that of native quinone in the atomic structure of yeast complex III (PDB 1EZV). The aliphatic octyl- and ethyl substituents of ametoctradin are predicted to form stabilizing hydrophobic interactions with the sidechains of I17 (helix A, which is L17 in *P.viticola*), V194 and L198 (C-terminal region of helix D). Accordingly, we note that the L198F substitution in PFQi3 and PFQi8 may further strengthen the interaction with ametoctradin, explaining the increased sensitivity of these mutants to this inhibitor. In contrast, the interactions between the amino-substituted triazolo-pyrimidinyl headgroup of ametoctradin and the Q_i-site are predicted to be more hydrophilic in nature, with a putative H-bonding interaction (2.5 Å separation) between the amino group of the headgroup and the carboxylate sidechain of D229. This may take the form of a salt bridge, depending upon the local dielectric microenvironment. Significantly, we also predict a weaker H-bonding interaction between this amino moiety of ametoctradin and the serinyl sidechain of S34 (3.1 Å separation), which may explain the resistance conferred by the S34L mutation. In any case,

1
2
3 297 introduction of the bulky leucyl sidechain at this position may be expected to be sterically
4
5 298 destabilizing with respect to ametoctradin binding.
6
7 299

9 300 **4. CONCLUSION**

11 301 The brief overview of the reported Q_i-site resistance mutations found in laboratory models or
12
13 302 field isolates demonstrates the mutational plasticity of cytochrome *b* in the presence of QiIs,
14
15 303 single site competitive inhibitors that function as Q_i-site antagonists. Cytochrome *b* is
16
17 304 mitochondrially encoded, and it is apparent that mitochondrial genes seem to accumulate
18
19 305 mutations at a higher frequency than nuclear genes. Target site resistance mutations are hence
20
21 306 expected to arise under selective pressure. Dual Q_o- and Q_i-site inhibitors would be effective
22
23 307 as mutation of both quinone binding sites would be expected to be a very infrequent event
24
25 308 whilst still maintaining organismal fitness. However such dual acting inhibitors are rare²⁶. In
26
27 309 their absence, a combination of a QoI and a QiI could be used. Unfortunately, at least for
28
29 310 *P.viticola*, the Q_o-site mutation G143A is already present in field population and isolates
30
31 311 harbouring both G143A and S34L have been observed¹³, which hinders that strategy. Other
32
33 312 combined treatments might need to be devised, possibly targeting sites away from complex
34
35 313 III.

37
38
39 314 A fitness cost associated with the resistance mutations would also be protective, as the
40
41 315 mutants would be counter-selected in absence of treatment, as might be the case for the
42
43 316 ametoctradin resistance mutation S34L as reported by Fontaine et al.¹³. A fitness penalty
44
45 317 associated with the resistance was confirmed in our yeast model. PFQi3 with a modified Q_i-
46
47 318 site rendered sensitive to ametoctradin seemed thus a useful model. However its pertinence
48
49 319 should be further confirmed by studying the effect of S34L on the oomycete complex III
50
51 320 activity and inhibitor sensitivity.
52
53 321

ACKNOWLEDGEMENTS

This work was in part supported by funding from ANSES (French Agency for Food, Environmental and Occupational Health & Safety) in the framework of the program of ‘phytopharmacovigilance’. NF acknowledges support from Grant DE-FG02-11ER16220 from the Photosynthetic Systems program from Division of Chemical Sciences, Geosciences, and Biosciences, Office of Basic Energy Sciences of the U.S. Department of Energy.

References

- 1 Bartlett DW, Clough JM, Godwin JR, Hall AA, Hamer M, and Parr-Dobrzanski B, The strobilurin fungicides. *Pest Manag Sci.* **58**:649–662 (2002).
- 2 Cottrell G, Musset L, Hubert V, Le Bras J, and Clain J, Emergence of resistance to atovaquone-proguanil in malaria parasites: insights from computational modeling and clinical case reports. *Antimicrob Agents Chemother* **58**:4504–4514 (2014).
- 3 Musset L, Bouchaud O, Matheron S, Massias L, and Le Bras J, Clinical atovaquone-proguanil resistance of *Plasmodium falciparum* associated with cytochrome *b* codon 268 mutations. *Microbes Infect* **8**:2599–2604 (2006).
- 4 Birth D, Kao W-C, and Hunte C, Structural analysis of atovaquone-inhibited cytochrome *bc*₁ complex reveals the molecular basis of antimalarial drug action. *Nat Commun* **5**:4029 (2014).
- 5 Esser L, Quinn B, Li Y-F, Zhang M, Elberry M, Yu L, *et al.*, Crystallographic studies of quinol oxidation site inhibitors: a modified classification of inhibitors for the cytochrome *bc*₁ complex. *J Mol Biol* **341**:281–302 (2004).
- 6 Stickles AM, De Almeida MJ, Morrissey JM, Sheridan KA, Forquer IP, Nilsen A, *et al.*, Subtle changes in endochin-like quinolone structure alter the site of inhibition within

1
2
3 347 the cytochrome *bc*₁ complex of *Plasmodium falciparum*. Antimicrob Agents
4
5 348 Chemother **59**:1977–1982 (2015).
6
7 349 7 Young DH, Wang NX, Meyer ST, and Avila-Adame C, Characterization of the
8
9 350 mechanism of action of the fungicide fenpicoxamid and its metabolite UK-2A. Pest
10
11 351 Manag Sci **74**:489-498 (2018).
12
13 352 8 Mitani S, Araki S, Takii Y, Ohshima T, Matsuo N, and Miyoshi H, The biochemical
14
15 353 mode of action of the novel selective fungicide cyazofamid: Specific inhibition of
16
17 354 mitochondrial complex III in *Phythium spinosum*. Pestic Biochem Physiol **71**:107–115
18
19 355 (2001).
20
21
22 356 9 Fehr M, Wolf A, and Stammel G, Binding of the respiratory chain inhibitor
23
24 357 ametoctradin to the mitochondrial *bc*₁ complex. Pest Manag Sci **72**:591–602 (2016).
25
26 358 10 Dreinert A, Wolf A, Mentzel T, Meunier B, and Fehr M, The cytochrome *bc*₁ complex
27
28 359 inhibitor Ametoctradin has an unusual binding mode. Biochim Biophys Acta
29
30 360 **1859**:567-576 (2018).
31
32
33 361 11 Ding MG, Di Rago JP, and Trumpower BL, Investigating the Q_n site of the cytochrome
34
35 362 *bc*₁ complex in *Saccharomyces cerevisiae* with mutants resistant to ilicicolin H, a novel
36
37 363 Q_n site inhibitor. J Biol Chem **281**:36036–36043 (2006).
38
39 364 12 Vallières C, Fisher N, Antoine T, Al-Helal M, Stocks P, Berry NG, *et al.*, HDQ, a
40
41 365 potent inhibitor of *Plasmodium falciparum* proliferation, binds to the quinone reduction
42
43 366 site of the cytochrome *bc*₁ complex. Antimicrob Agents Chemother **56**:3739–3747
44
45 367 (2012).
46
47
48 368 13 Fontaine S, Remuson F, Caddoux L, and Barrès B, Using a combination of bioassay
49
50 369 and molecular tools to survey pesticide resistance to complex III inhibitor of
51
52 370 *Plasmopara viticola* populations in French vineyards. Pest Manag Sci submitted.
53
54 371 14 Hill P, Kessl J, Fisher N, Meshnick S, Trumpower BL, and Meunier B, Recapitulation

- 372 in *Saccharomyces cerevisiae* of cytochrome *b* mutations conferring resistance to
373 atovaquone in *Pneumocystis jiroveci*. Antimicrob Agents Chemother **47**:2725–2731
374 (2003).
- 375 15 Fisher N, Castleden CK, Bourges I, Brasseur G, Dujardin G, and Meunier B, Human
376 disease-related mutations in cytochrome *b* studied in yeast. J Biol Chem **279**:12951–
377 12958 (2004).
- 378 16 Lemaire C and Dujardin G, Preparation of respiratory chain complexes from
379 *Saccharomyces cerevisiae* wild-type and mutant mitochondria: activity measurement
380 and subunit composition analysis. Methods Mol Biol **432**:65–81 (2008).
- 381 17 Sali A and Blundell TL, Comparative protein modelling by statisfaction of spatial
382 restraints. J Mol Biol **234**:779–815 (1993).
- 383 18 Verdonk ML, Cole JC, Hartshorn MJ, Murray CW, and Taylor RD, Improved protein-
384 ligand docking using GOLD. Proteins Struct Funct Genet **52**:609–623 (2003).
- 385 19 Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, *et al.*,
386 UCSF Chimera - A visualization system for exploratory research and analysis. J
387 Comput Chem **25**:1605–1612 (2004).
- 388 20 Rotsaert FAJ, Ding MG, and Trumpower BL, Differential efficacy of inhibition of
389 mitochondrial and bacterial cytochrome *bc*₁ complexes by center N inhibitors
390 antimycin, ilicicolin H and funiculosin. Biochim Biophys Acta **1777**:211–219 (2008).
- 391 21 Song Z, Laleve A, Vallières C, McGeehan JE, Lloyd RE, and Meunier B, Human
392 mitochondrial cytochrome *b* variants studied in yeast: not all are silent polymorphisms.
393 Hum Mutat **37**:933-941 (2016).
- 394 22 Gao X, Wen X, Esser L, Quinn B, Yu L, Yu C-A, *et al.*, Structural basis for the
395 quinone reduction in the *bc*₁ complex: a comparative analysis of crystal structures of
396 mitochondrial cytochrome *bc*₁ with bound substrate and inhibitors at the Q_i site.

- 397 Biochemistry **42**:9067–9080 (2003).
- 398 23 CONIphy, Mildiou de la Vigne, Caracterisation du mécanisme de resistance aux QiI
399 Newsletter N°1 (2017).
- 400 24 Coppee JY, Brasseur G, Brivet-Chevillotte P, and Colson AM, Non-native intragenic
401 reversions selected from *Saccharomyces cerevisiae* cytochrome *b*-deficient mutants.
402 Structural and functional features of the catalytic center N domain. J Biol Chem
403 **269**:4221–4226 (1994).
- 404 25 Li H, Zhu XL, Yang WC, and Yang GF, Comparative kinetics of Qi site inhibitors of
405 cytochrome *bc*₁ complex: Picomolar antimycin and micromolar cyazofamid. Chem Biol
406 Drug Des **83**:71–80 (2014).
- 407 26 Song Z, Iorga BI, Mounkoro P, Fisher N, and Meunier B, The antimalarial compound
408 ELQ-400 is an unusual inhibitor of the *bc*₁ complex, targeting both Q_o and Qi sites.
409 FEBS Lett **592**:1346-1356 (2018).
- 410 27 di Rago JP, Perea X, and Colson AM, DNA sequence analysis of diuron-resistant
411 mutations in the mitochondrial cytochrome *b* gene of *Saccharomyces cerevisiae*. FEBS
412 Lett **208**:208–210 (1986).
- 413 28 Coria R, Garcfa M, and Brunner A, Mitochondrial cytochrome *b* genes with a six-
414 nucleotide deletion or single-nucleotide substitutions confer resistance to antimycin A
415 in the yeast *Kluyveromyces lactis*. Mol Microbiol **3**:1599–1604 (1989).
- 416 29 Brasseur G and Brivet-Chevillotte P, Characterization of mutations in the
417 mitochondrial cytochrome *b* gene of *Saccharomyces cerevisiae* affecting the quinone
418 reductase site (Q_N). Eur J Biochem **230**:1118–1124 (1995).
- 419 30 Schnauffer A, Sbicego S, and Blum B, Antimycin A resistance in a mutant *Leishmania*
420 *tarentolae* strain is correlated to a point mutation in the mitochondrial apocytochrome *b*
421 gene. Curr Genet **37**:234–241 (2000).

- 422 31 Di Rago JP and Colson AM, Molecular basis for resistance to antimycin and diuron, Q-
 423 cycle inhibitors acting at the Q_i site in the mitochondrial ubiquinol-cytochrome *c*
 424 reductase in *Saccharomyces cerevisiae*. J Biol Chem **263**:12564–12570 (1988).
- 425 32 Lukens AK, Heidebrecht RW, Mulrooney C, Beaudoin JA, Comer E, Duvall JR, *et al.*,
 426 Diversity-oriented synthesis probe targets *Plasmodium falciparum* cytochrome *b*
 427 ubiquinone reduction site and synergizes with oxidation site inhibitors. J Infect Dis
 428 **211**:1097–1103 (2015).
- 429 33 Howell N and Gilbert K, Mutational analysis of the mouse mitochondrial cytochrome *b*
 430 gene. J Mol Biol **203**:607–618 (1988).
- 431 34 Weber S and Wolf K, Two changes of the same nucleotide confer resistance to diuron
 432 and antimycin in the mitochondrial cytochrome *b* gene of *Schizosaccharomyces pombe*.
 433 FEBS Lett **237**:31–34 (1988).
- 434 35 di Rago JP, Perea J, and Colson AM, Isolation and RNA sequence analysis of
 435 cytochrome *b* mutants resistant to funiculosin, a center i inhibitor of the mitochondrial
 436 ubiquinol-cytochrome *c* reductase in *Saccharomyces cerevisiae*. FEBS Lett **263**:93–98
 437 (1990).
- 438 36 Khare S, Roach SL, Barnes SW, Hoepfner D, Walker JR, Chatterjee AK, *et al.*,
 439 Utilizing chemical genomics to identify cytochrome *b* as a novel drug target for Chagas
 440 disease. PLOS Pathog **11**:e1005058 (2015).
- 441 37 Coppée JY, Tokutake N, Marc D, di Rago JP, Miyoshi H, and Colson AM, Analysis of
 442 revertants from respiratory deficient mutants within the center N of cytochrome *b* in
 443 *Saccharomyces cerevisiae*. FEBS Lett **339**:1–6 (1994).
- 444 38 Doggett JS, Nilsen A, Forquer I, Wegmann KW, Jones-Brando L, Yolken RH, *et al.*,
 445 Endochin-like quinolones are highly efficacious against acute and latent experimental
 446 toxoplasmosis. Proc Natl Acad Sci U S A **109**:15936–15941 (2012).

1
2
3 447 39 Alday PH, Bruzual I, Nilsen A, Pou S, Winter R, Ben Mamoun C, *et al.*, Genetic
4
5 448 evidence for cytochrome *b* Q_i site inhibition by 4(1H)-quinolone-3-diarylethers and
6
7 449 antimycin in *Toxoplasma gondii*. Antimicrob Agents Chemother **61**:e01866-16 (2017).
8

9 450
10
11 451
12
13
14 452

15
16
17 453
18
19 454
20

21
22 455 **Table 1. Inhibitor resistance mutations in complex III Q_i-site**

Mutation ^a	Resistant to	Organism	References ^b
I17F	Diuron	<i>Saccharomyces cerevisiae</i>	27
S20L, T	Illicicolin H	<i>Saccharomyces cerevisiae</i>	11
Q22E, T	Illicicolin H	<i>Saccharomyces cerevisiae</i>	11
I26L	ELQ-300	<i>Plasmodium falciparum</i>	6
W30C ^c	Illicicolin H,Funiculosin	<i>Saccharomyces cerevisiae</i>	20
N31K	Diuron,	<i>Saccharomyces cerevisiae</i>	27
	Antimycin, Fenpicoxamid		7
N31K	Antimycin	<i>Kluyveromyces lactis</i>	28
G33A ^c	Funiculosin, HQNO	<i>Saccharomyces cerevisiae</i>	29
	HDQ		12
S34L	Ametoctradin	<i>Plasmopora viticola</i>	13
S(L)35I	Antimycin	<i>Leishmania torentola</i>	30
G37C	Fenpicoxamid, Antimycin	<i>Saccharomyces cerevisiae</i>	7
G37D, S	Illicicolin H	<i>Saccharomyces cerevisiae</i>	11
G37V	Antimycin	<i>Saccharomyces cerevisiae</i>	31

G37A	ML238, BRD6923	<i>Plasmodium falciparum</i>	32
G37V	Antimycin	<i>Zymoseptoria tritici</i>	9
G37V	ML238, BRD6323, Antimycin	<i>Plasmodium falciparum</i>	32
G37V	Antimycin	<i>Mus musculus</i>	33
A(G)37V	Antimycin	<i>Schizosaccharomyces pombe</i>	34
A(G)37G	Diuron	<i>Schizosaccharomyces pombe</i>	34
L198F	Funiculosin,	<i>Saccharomyces cerevisiae</i>	35
	Ilicolin H,		11
	Fenpicoxamid, Antimycin		7
L198F	GNF7686, Antimycin	<i>Trypanosoma cruzi</i>	36
E203-DE-V204	Cyazofamid	<i>Plasmopora viticola</i>	23
(I203--H204)			
H204Y ^c	HDQ	<i>Saccharomyces cerevisiae</i>	12
M221Q ^c	Antimycin,	<i>Saccharomyces cerevisiae</i>	37
	HQNO		29
	Ilicolin H		20
	ELQ-271, ELQ-319		26, 38
M221E ^c	Antimycin	<i>Saccharomyces cerevisiae</i>	37
	HQNO		29
	Ilicolin H,		20
F225S, L	Diuron	<i>Saccharomyces cerevisiae</i>	27, 31
K228M	Antimycin	<i>Saccharomyces cerevisiae</i>	31
K228M ^c	HDQ	<i>Saccharomyces cerevisiae</i>	12
K228I/M	Antimycin	<i>Kluyveromyces lactis</i>	28
T228P	ELQ-316, Antimycin	<i>Toxoplasma gondii</i>	39
G(T)232D	Antimycin, HQNO	<i>Mus musculus</i>	33

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

457 ^a Yeast numbering; (), equivalent residue in yeast cytochrome *b* if different

458 ^b Cited references are the first report or first study on the mutations

459 ^c These mutations were not identified in mutants selected after inhibitor treatment but
460 originally were found in “revertants” i.e. respiratory growth competent clones selected from
461 respiratory deficient mutants (the defect was due to a point mutation in cytochrome *b*)

462

For Peer Review

463 **Table 2. Mutations introduced in yeast Q_i-site**

position	WT	PFQi1	PFQi2	PFQi3	PFQi8	PFQi9	PFQi10
34	S					L	L
40	L		F				
41	V		F				
190	L			G			G
191	A	L	L	L	L	L	L
192	A			C			C
195	I	F	F	F	F	F	F
198	L	I	I	F	F	I	F
207	S			T			T
221	M	F	F	F	F	F	F
225	F	L	L	L	L	L	L
226	I	L	L	L	L	L	L

464

465 The PFQi mutants were constructed using the mitochondrial transformation technique. PFQi1,
466 2, 3 and 8: yeast amino-acids were replaced by their equivalents observed in the protozoan
467 *Plasmodium falciparum* (with the exception of L198F). PFQi9 and 10: the ametoctradin
468 resistance mutation S34L was introduced in the PFQi modified Q_i-site.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figures legends

Figure 1. A) Catalytic core of complex III showing the redox-groups, haems and the 2Fe2S cluster, formed by three of the complex subunits: cytochrome *c*, the Rieske Iron-Sulphur Protein and cytochrome *b*, which is mitochondrially-encoded and provides the Q_o- and Q_i- sites.

B) Schematic representation of cytochrome *b*. Positions of residues involved in inhibitor resistance are shown: Q_o-site G143 and Y279; Q_i-site I17, N31, S34, G37, L198, H204 and K228.

Figure 2. Sequence comparison of the Q_i-site region of the cyazofamid resistance mutation, from H197 to S207 (*S.cerevisiae* residues and numbering). The sequences of ten fungal cytochrome *b* are compared. In bold, the sequence of the cyazofamid resistant isolate of *P.viticola*. Four residues are indicated, the conserved residues L198, H202 and S206, and the variable residue H204.

Figure 3. Effect of mutations in complex III Q_i-site on the growth sensitivity to ametoctradin. Suspensions in water of cells pre-grown on YPD plates were spotted, at three different cell concentrations, on plates containing the respiratory medium YPG with increasing concentrations of ametoctradin.

Figure 4. Growth sensitivity to amisulbrom of PFQ_i mutants. Suspensions in water of cells pre-grown on YPD plates were spotted, at three different cell concentrations, on plates containing the respiratory medium YPG with or without 100 µM amisulbrom.

Figure 5. Growth competence and ametoctradin sensitivity of mutants PFQi1, 3, 9 and 10. Cells freshly grown on YPD plates are inoculated at an initial OD_{600nm} of 0.2 in YPEth without or with ametoctradin at increasing concentrations. The cultures were incubated three days at 28°C with vigorous agitation. The OD_{600nm} were then recorded. The growth experiments were repeated at least twice and the data averaged. The error bars represent standard deviation A) Cell densities reached in absence of ametoctradin. B and C) Effect of ametoctradin. The data are presented as percentage of the OD_{600nm} of the untreated cultures. B) Comparison between PFQi1 (solid line) and PFQi9 (dotted line); C) Comparison between PFQi3 (solid line) and PFQi10 (dotted line).

Figure 6. Complex III activity of control, PFQi3 and PFQi10. Cytochrome *c* reduction was monitored at 550-540 nm. The reaction was initiated by the addition of 20 µM decylubiquinol to the reaction mixture containing 20 µM cytochrome *c* and 2-10 nM complex III. Initial rates were measured. Each measurement was repeated three to five times and the values obtained were averaged. The activities are presented as the cytochrome *c* reduction rate per complex III per second

Figure 7. Docking conformation of ametoctradin in the Q_i-site of *P.viticola*. Ametoctradin is represented as sticks with partially transparent surface. Residues that are interacting with the ligand are shown as sticks. The haem is also visible in the background.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1. Catalytic core of complex III and schematic representation of cytochrome b
69x29mm (300 x 300 DPI)

<i>S.cerevisiae</i>	HLMALHI--HGSS
<i>Z.tritici</i>	HLIALHDT-AGSG
<i>B.cinerea</i>	HLIALHDS-AGSG
<i>P.viticola</i>	HLILLHE--VGSN
Cyazo-Res	HLILLHE DE VGSN
<i>C.albicans</i>	HLMALHV--HGSS
<i>A.fumigates</i>	HLIAMHDT-VGSG
<i>S.pombe</i>	HLIALHT--NGSS
<i>P.anserina</i>	HLIALHDT-AGSS
<i>U.maydis</i>	HLLALHE--HGSS
<i>P.graminis</i>	HLLTLHE--HGSN

Figure 2. Sequence comparison of the Q_i-site region of the cyazofamid resistance

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 3. Effect of mutations in complex III Q_i -site on the growth sensitivity to ametoctradin

Peer Review

Figure 4. Growth sensitivity to amisulbrom of PFQI mutants

Or Peer Review

Figure 5. Growth competence and ametoctradin sensitivity of mutants PFQi1, 3, 9 and 10

Figure 6. Complex III activity of control, PFQi3 and PFQi10

Figure 7. Docking conformation of ametoctradin in the Q_i-site of *P.viticola*

Or Peer Review