2 **Gradient of decomposition in sugarcane mulches of various thicknesses**

4 Guilherme Dietrich^a, Sylvie Recous^{b*}, Patrick Leal Pinheiro^a, Douglas Adams Weiler^a, Adriane Luiza
Schu^a, Mathias Roberto Leite Rambo^a, Sandro José Giacomini^a

6

^a Department of Soils, Federal University of Santa Maria, 97105-900 Santa Maria, RS, Brazil

8 ^b FARE Laboratory, INRA, Université de Reims Champagne-Ardenne, 2 Esplanade R. Garros, 51100
Reims, France

10

*Corresponding author:

12 Sylvie Recous, Tel : +33 (0)326 773 583, E-mail: sylvie.recous@inra.fr

14 **Highlights**

- The vertical decomposition gradient was characterized in mulches of various thicknesses
- Top mulch layers decomposed similarly, regardless of the thickness of the underlying mulch
- Low mulch layers decomposed faster as the thickness of the overlying mulch increased
- The decomposition gradient within the mulch was closely correlated with the moisture gradient

20 **Abstract**

Crop residues left on the soil surface as mulch influence many services provided in agrosystems, particularly soil protection, water dynamics and nutrient fluxes. For a given crop, the residue mass influences the mulch thickness, but the effect of thickness on decomposition is not well understood, and decomposition gradients within mulches in the field have not been described. This study aimed to quantify the decomposition of sugarcane straw mulches by varying their mass and thickness and determine the decomposition gradient within the mulches. The experiment was conducted on a first-ratoon sugarcane crop for one year, with 4, 8 and 12 Mg straw dry matter ha⁻¹. Mulches were displayed in 0.16-m² litter boxes and formed by straw layers of equivalent mass, i.e., 4 Mg ha⁻¹, stacked either as a single layer (Low/Top layer), 2 layers (Low and Top layers) or 3 layers (Low, Medium and Top layers) to reconstitute the 3 mulch quantities. We quantified the carbon (C) and nitrogen (N) contents, the chemical composition and the water content of the remaining mulch particles in each layer 10 times within the 360 days. The mulch degradation rates were proportional to the initial amount of straw, the mulch losses representing 75% of its initial C and 46% of its initial N after one year, regardless of the initial mass. Fertilizer-N addition did not change the k decomposition rate, with $k=0.0044$ d⁻¹. However the decomposition rate differed according to the layer position in the mulch, e.g., in the 12 Mg ha⁻¹ treatment $k=0.0064$ d⁻¹, 0.0046 d⁻¹ and 0.0033 d⁻¹ for the low, medium and top layers, respectively, indicating that the low layers in contact with the soil decomposed faster than the top layers in contact with the atmosphere, while the medium layers demonstrated intermediate behaviour. The top layers evolved similarly, regardless of mulch thickness, while the lower layers in

contact with the soil decomposed more quickly as the overlying straw thickness increased. Regardless of the thickness and position of a given layer within mulch, the one-year layer C loss was linearly correlated with the average water content of that layer over the year ($y=52.79 - 0.0694x$, $R^2=0.991$, $P<0.05$). Here, we demonstrate that the dynamics of mulch decomposition is primarily driven by the dynamics of water, which is itself driven by mulch characteristics, rain regime and evaporation.

Keywords: Straw litter layer, carbon decomposition, nitrogen, crop residue water properties, soil-residue contact

1. Introduction

No-till in cropping systems are increasingly used worldwide for protecting soil from physical degradation and enhancing soil fertility (Tiecher et al., 2017), and the development of perennial or semi-perennial crops (miscanthus, sugarcane, etc.) dedicated to bioenergy (Peyrard et al., 2017; Bordonal et al., 2018) leads to the generalization of agricultural situations with crop residue layers, i.e., mulch, decomposing on the soil surface. Predicting the decomposition rates of crop residue mulches according to system characteristics (crop and rotation types, climatic conditions) is important because of the crucial services these mulches provide, such as soil cover and protection against erosion and weed growth, water regulation, nutrient recycling, and habitats and resources for soil communities. Moreover, a better understanding of the decomposition of straw at the soil surface is important because this process determines the fluxes of carbon (C) and nitrogen (N) in the soil and to the atmosphere via the emission of greenhouse gases (Chen et al., 2013; Pinheiro et al., *accepted*; Vasconcelos et al., 2018).

Many studies have shown that the decomposition rate of straw at the soil surface is lower than that of residues incorporated into the soil (Douglas et al., 1980; Schomberg et al., 1994; Coppens et al., 2006; Curtin et al., 2008). This difference was attributed to the straw undergoing greater and more frequent fluctuations in moisture at the soil surface than within the soil, particularly frequent drying between rains, combined with a reduced diffusion of nutrients and water from the soil to decomposition sites in the mulch (Christensen, 1986). The soil-residue contact determined by the

characteristics of crop residues (morphology and particle size, mass) and the characteristics of the underlying soil layer (texture, water content, mineral N content) are considered important in driving decomposition (e.g., Angers and Recous, 1997; Henriksen and Bleland, 2002; Iqbal et al., 2013). The straw mass, together with the residue morphology, is the main factor that determines the proportion of the mulch that is in contact with the soil. For a given crop residue type, as the residue amount on the soil surface increases, the thickness of the produced mulch increases (Putuhena and Cordery, 1996), and the proportion of the residues that are in contact with the soil decreases. This was reflected in the fact that under a number of experimental conditions the rate of decomposition decreased with an increase in the mass of crop residues at the soil surface (e. g. Steiner et al., 1999). These results had led models that simulate the decomposition of straw at the soil surface to consider the degradation of the mulch to occur from a decomposing layer in contact with the soil fed by an unaltered layer of overlying residues (APSIM Thorburn et al., 2001; CANTIS Findeling et al., 2007). However, a number of recent studies have changed this perception by showing that mulch decomposition can be proportional to the initial straw mass, within a range of 3 to 12 Mg ha⁻¹ for cover crop mulches (Acosta et al., 2014; Halde and Entz, 2015), sugarcane (Dietrich et al., 2017), or 5 to 15 Mg ha⁻¹ for rye (Williams et al., 2018). These results are questioning the hypothesis that the undecomposed proportion of the top layer increases with increasing residue mass. Therefore, it is necessary to improve our understanding of the decomposition processes of residue mulch, particularly how these decomposition processes are affected by the mass and thickness of mulch.

To answer these questions, the objective of this work was to investigate the vertical heterogeneity of the decomposition process within sugarcane straw mulches with different initial masses, thicknesses, and N availabilities. We investigated the following aspects in field conditions over one year after sugarcane harvest: i) how decomposition develops at different locations within mulches of varied initial masses, ii) the consequences of mulch features on mulch C and N dynamics and on the conditions in the soil layer below the mulch, and iii) the factors driving mulch decomposition and the role of the contact between soil and mulch in this decomposition.

94 2. Materials and methods

2.1 Study site

96 The study was located at the Federal University of Santa Maria (29°42'54" S, 53°42'23" W,
approximately 90 m elevation) in the state of Rio Grande do Sul, Brazil. The climate, according to the
98 Köppen classification system, is a type Cfa subtropical humid climate. The mean annual temperature
is 19.3°C; June is the coldest month, with a mean minimum temperature of 9.3°C, and January is the
100 warmest month, with a mean maximum temperature of 30.4°C. The mean annual precipitation is 1660
mm, without a dry season. The soil is classified as Typic Paleudalf according to USDA taxonomy
102 (Soil Survey Staff 2010). Selected soil characteristics (0-20 cm layer) at the beginning of the
experiment were 110 g kg⁻¹ clay, 260 g kg⁻¹ silt, 630 g kg⁻¹ sand, soil bulk density = 1.63 g dm⁻³, pH
104 (water) = 5.4; Mehlich-I P = 33.4 mg dm⁻³, total C = 4.1 g kg⁻¹ and total N = 0.32 g kg⁻¹.

2.2 Main experimental design

106 The experiment was carried out at the first sugarcane ratoon from November 2015 to
November 2016 and began after the harvest of cane plants on November 20, 2015. The sugarcane
108 variety was RB 956911, and the harvest was manual, removing all aboveground biomass (stalk +
leaves + tops) and leaving the soil uncovered. An area was demarcated with 32 plots in 4 lines with
110 1.4 m of spacing between rows and length of 4 m. Each plot had an area of 22.4 m². We used a
randomized complete block design in a factorial scheme of 4x2 with four replicates. The first
112 treatment was the straw quantity treatment, with four amounts of straw on the soil surface: 0, 4, 8 and
12 Mg ha⁻¹ of dry matter (DM). The second treatment was the fertilizer N treatment, with 0 or 100 kg
114 N ha⁻¹ added as urea. The straw was homogeneously distributed by hand on the soil surface, resulting
in 8 treatments (0S, 0S+N, 4S, 4S+N, 8S, 8S+N, 12S, 12S+N). The urea N was applied as granules
116 broadcasted in a single dose on the straw on day 52 after the plant cane harvest.

2.3 Straw decomposition microplots

A litter box approach was used to monitor straw decomposition (Fig. 1). Air-dried straw was added into 0.16-m² wooden frames (40-cm length x 40-cm width x 8-cm height) between rows of sugarcane, as described by Dietrich et al. (2017). To evaluate the gradient of decomposition within the mulch, the mulch was formed by straw layers of 4 Mg ha⁻¹, and we stacked a single layer, 2 layers or 3 layers inside the litter box to reconstitute the 3 mulch quantities. The layers were separated by a malleable 10-mm synthetic mesh with a wide aperture so each layer could subsequently be sampled separately for analysis. The 4 Mg ha⁻¹ treatment (4S) had a single layer, which was, in essence, simultaneously in contact with the soil (low position: L) and in contact with the atmosphere (top position: T), the layer in this treatment is referred to as 4S L/T. The 8 Mg ha⁻¹ treatment was composed of two layers, a top layer (8S T) and a low layer (8S L); the 12 Mg ha⁻¹ treatment was composed of three layers, a top layer (12S T), a middle layer (12S M) and a low layer (12S L).

Destructive sampling of litter boxes was performed on days 0, 15, 36, 51, 72, 103, 168, 238, 296 and 360 after straw application from November 2015 to November 2016. At each sampling date, the straw from each layer in each destroyed litter box was collected separately, weighed wet, stored in paper bags and dried in an oven at 40°C. The straw was then separated from adhering soil by dry cleaning, and a subsample was dried in an oven at 65°C to determine the remaining straw DM. Straw ashes were measured by incinerating 0.4 g in a muffle furnace at 550°C for 3 h. The total C and N concentrations of the straw were determined using an elemental analyser (FlashEA 1112 model, ThermoFinnigan, Milan, Italy). Straw soluble (SOL), cellulose (CEL), hemicellulose (HEM) and lignin (LIG) fractions were determined in residues pre-dried at 40°C using the method described by Van Soest (1963). The water-soluble organic C (C_{sw}) and water-soluble total N (N_{sw}) were extracted using 2 g of chopped plant residue (1 cm) pre-dried at 40°C in 80 mL of water following the methodology described by Schmatz et al. (2017). The C_{sw} and N_{sw} contents were analysed using a total organic carbon analyser equipped with a kit for N analysis (Shimadzu TOC-LCPH, Shimadzu, Kyoto, Japan). The chemical characteristics of the straw at the beginning of the experiment were as follows: 228 ± 7.5 g kg⁻¹ SOL, 372 ± 2.7 g kg⁻¹ CEL, 335 ± 5.2 g kg⁻¹ HEM, 65 ± 1.6 g kg⁻¹ LIG, 425

$\pm 2.7 \text{ g kg}^{-1}$ total C, $8 \pm 0.2 \text{ g kg}^{-1}$ total N, $36.7 \pm 8.5 \text{ g kg}^{-1}$ C_{sw}, $1.2 \pm 0.2 \text{ g kg}^{-1}$ N_{sw} and a C:N ratio of 53 ± 1.4 .

Mulch thickness was evaluated in four litter boxes randomly selected by treatment (4S, 8S, 12S). In each box, nine non-destructive thickness measurements were performed using a graduate ruler. The measurements were performed at 0, 29, 54, 72, 103, 168, 238, 296 and 360 days after straw addition.

2.4 Soil water content and soil temperature

Sensors with two 30-cm-long stainless steel rods (FDR CS 616-L, Campbell Scientific, Logan, Utah, US) inserted at 2.5 and 7.5 cm soil depth were used to measure soil moisture. The type T copper-constantan thermocouple was installed at the same depths to measure soil temperature. Both sensors were installed between sugarcane rows and coupled to a datalogger (CR1000, Campbell Scientific, Logan, Utah, USA), and measurements were performed at a 10 min sampling interval. The air temperature at 30-cm depth inside of the sugarcane plantation was also evaluated using the same sensor model. The sensor was covered by a polyvinyl chloride pipe wrapped by asphalt adhesive tape with holes on the sides. Daily air temperature and rainfall data were obtained from an automatic weather station located 1.7 km away from the experiment site.

2.5 Calculations and statistical analysis

The remaining C and N of the straw at each sampling date were calculated as the remaining mass of each straw layer multiplied by its C and N concentrations (deduced from the ash content). In treatments 8S and 12S (with and without N), the amounts of C and N remaining were obtained by summing the amounts of C and N in each straw layer. The C remaining in each layer and in the total mulch over time was described with a simple one-compartment equation according to Plante and Parton (2007):

$$C_{\text{remaining}} = C_0 e^{-kt}$$

where C remaining is the mass of the remaining C (Mg C ha^{-1}), C_0 is the straw C pool at time 0 (Mg C ha^{-1}), k is the degradation rate constant (day^{-1}), and t is the time (days).

The mean residence time ($t_{1/2}$) of the straw, which corresponds to the time necessary to degrade half of the initial quantity of mulch, was calculated from k according to Plante and Parton (2007). The LCI (lignocellulose index) was calculated from the CEL, HEM and LIG fractions of sugarcane straw and corresponds to the relative enrichment of LIG in the plant cell walls (Whittinghill et al., 2012). This index was taken as an indicator of the degradation process of the lignocellulose substrates.

All data were subjected to analysis of variance (ANOVA). The effect of straw quantity or straw layer, N fertilization and time, and their interaction on mulch decomposition (C and N remaining, SOL, CEL, HEM and LIG remaining and LCI index) and physical (thickness and water content) variables, were tested. Similarly, we tested the effect of straw quantity or straw layer and N fertilization, and their interaction on decomposition model parameters (C_0 , k and $t_{1/2}$). In the straw layer data analysis, data from the 4S treatment were included in the statistical analysis. This treatment was considered to have a straw layer representing both L and T straw because the layer was in direct contact with the soil and the atmosphere simultaneously. The effect of straw quantity, soil depth and time, and their interaction on soil moisture and soil temperature also were tested. When ANOVA showed significant ($P < 0.05$), the differences between means of treatments were compared by the Tukey test ($P < 0.05$). Relationships between remaining C or N measured at the end of experiment and the average water content of straw layers in the mulch (T/L, L, M and T) for the three treatments (4S, 8S and 12S) over the 360 days of the experiment was fitted.

3. Results

3.1 Climatic and soil environmental conditions

The total annual rainfall was 1519 mm, distributed over 107 events, and the mean air temperature was 18.4 °C, similar to the historical average (Fig. 2a). The relative air humidity ranged from 39 to 99%, with most values in the 70-90% range (Fig. 2b). As the temperature and moisture at the 2.5 and 7.5 cm soil depths varied in the same way over time and between treatments, only the 2.5 cm soil depth data are shown here (Fig. 2c and 2d). The effect of straw amount on soil temperature

was observed only in the first three months, when the sugarcane canopy had still not completely covered the soil surface (Fig. 2c). The mean soil temperature during this period at 2.5 cm depth decreased with increasing amounts of straw, as 0S (28.2°C) > 4S (27.1°C) > 8S (26.4°C) = 12S (26.5°C), indicating an approximately 2°C difference in mean temperature between the highest (12S) and the lowest (0S) amount of straw. The soil water content responded to daily precipitation and was affected by straw quantity (Fig. 2d). Averaged over the experimental year, the soil volumetric water content was 0.17 for 0S, 0.19 for 4S, 0.21 for 8S and 0.23 g cm⁻³ for 12S. The soil water content was markedly lower in the treatments with lower straw quantities in the soil surface during periods with a small amount of rainfall or a short dry period, e.g., from day 46 to 67, day 190 to 217, and day 297 to 320.

3.2 Dynamic of thickness and water retention of mulch

The initial mulch thickness varied significantly with straw quantity, and, as expected, had a ranking of 12S (4.7 cm) > 8S (3.6 cm) > 4S (1.9 cm) at day 0 (Fig. 3a). The thickness was not strictly proportional to the mass, suggesting a layer settling with the larger mass and therefore a higher mulch density for 12S treatment. Then the evolution of mulch thickness was proportional to the initial thickness in the three straw treatments, with a 32% decrease in thickness at day 29, 50% decrease in thickness at day 103 and 68% decrease in thickness after 360 days. At that time, the mulch was 0.6, 1.2 and 1.5 cm thick for 4S, 8S and 12S, respectively.

The water content of the sampled mulches varied greatly over the course of the year, ranging between 10 and 80 g H₂O 100 g⁻¹ DM. The water content varied significantly as a function of straw quantity at some sampling dates (Fig. 3b) and differed according to straw layer position (Fig. 3c and Fig. S1). Comparing the layer positions for a given treatment, we observed that the water contents in the L layers of 8S and 12S were higher than those in the T layers of 8S and 12S, except for the samplings performed at days 51, 103 and 238. Comparing different mulch treatments for a given layer position, we observed that generally speaking, the thicker the mulch, the higher the moisture content of a layer in a given position was. On average over the year, the moisture content of the whole mulch

was finally low or negligible for the three quantities of mulch (Fig. 3b), suggesting compensation between the different layers according to thickness.

3.3 Dynamic of mulch decomposition

The straw C and N remaining as mulch, expressed as the percentage of the initial straw C or N added, decreased quickly over time but were not influenced by straw quantity, N fertilization or their interaction (Table S1); therefore, the mean data of the 0N and 100N treatments for each straw treatment are presented here (Fig. 4a and 4b). Notably, approximately half of the straw C and two-thirds of straw N lost during the 360-day period were lost in the first 36 days, regardless of the initial straw quantity. The straw C and N lost by day 360, averaged for the 12S, 8S and 4S treatments, amounted to 75 and 46% of the initial C and N, respectively. The C:N ratio of the remaining straw decreased from an initial C:N of 53 to a final C:N of 29, with no significant differences between straw treatments or between N treatments (Fig. S2).

Straw C_{sw} decreased very quickly, and by day 36, only 14, 12 and 19% of the initial C_{sw} remained in the 4S, 8S and 12S treatments, respectively, with significant transient differences between treatments (Fig. 4c). Straw N_{sw} decreased more slowly until day 72 and then remained stable until the end of the experiment (Fig. 4d). The chemical composition of the straw particles changed over time, with, at day 360, a relative increase in HEM and LIG and a strong decrease in CEL, while the SOL fraction remained stable (Fig. S2). This final composition was not influenced by straw quantity, N application or their interaction ($P > 0.05$) (Table S1). The C₀ straw pools calculated with the one-compartment equation was, as expected, close to the initial straw quantities, representing 1.47 Mg ha⁻¹ (4S), 2.99 Mg ha⁻¹ (8S) and 4.45 Mg ha⁻¹ (12S) (Table 1). The straw quantity and N addition did not change the k decomposition rates, with on average $k = 0.0044$ and $t_{1/2} = 158$ days.

3.4 Effect of straw layer position on decomposition

3.4.1. C and N evolution in the mulch layers

The dynamics of straw C and N in the various mulch layers were significantly influenced by the position of the straw layers within the mulch ($P < 0.001$) but were not significantly influenced by N

fertilization or the interaction between straw layer position and N fertilization ($P>0.05$) (Table S2); therefore, the means of the two N treatments are presented here (Fig. 5). In treatment 12S (Fig. 5a), the dynamics were characterized by a rapid decrease in straw C that was similar among all layer positions (0-36 days), followed by a slower decrease, during which the decomposition of 12 L>12M>12T. The same pattern of decomposition was also observed in the 8S treatment, with layer 8 L>layer 8T (Fig. 5c). Comparing the C loss of the top layers across the three straw treatments (4S L/T vs. 8S T vs. 12S T), we observed that the dynamics were very similar, except for at two dates, which demonstrated a transient difference that subsequently disappeared (Fig. 5e). The same comparison among low layers (4S L/T vs. 8S L vs. 12S L) showed that the 4S L/T layer decomposed less rapidly and had significantly less decomposition over the final two sampling dates than the 8S L and 12S L layers (Fig. 5g). As expected, the decomposition parameters of the adjusted model were significantly different only for mulch layer position (Table 1): the highest and lowest decomposition k rates were observed in the L and T layers, respectively.

The straw N kinetics followed the same pattern for each layer position (Fig. 5b,d,f,h). The dynamics were characterized by a rapid decrease in straw N (0-36 days) that was similar among all layer positions, followed by a slower decrease up to day 103, after which the remaining N values were relatively stable until day 360. Differences in the remaining N among the straw layers were observed mainly from day 238 in the 12S (T>M>L) and 8S (T>L) mulch. The N remaining in the mulch at the end of the experiment also differed between the low layers, with 4S L/T>12S L.

3.4.2. Changes in chemical composition

The chemical composition of the straw layers, i.e., the straw C:N ratio and chemical composition expressed by the LCI, taken as an indicator of the decomposition process, varied over time for all layer positions and depended on the position of layers in the mulch (Fig. 6 and Fig. S3). Considering the 12S treatment as an example, we observed that the pattern of evolution of the C:N ratio was similar among the three layers, but in general, the C:N decrease was more pronounced for low layers, with 12S L<12S M<12ST (Fig. 6a). The LCI increased from 0.08 to 0.17 over time for all

layers and was significantly different between layers at different positions until day 168 (e.g., 12L = 12 M>12T), but no difference persisted at day 360 (Fig. 6b).

3.4.3. Relationships with straw water content

The C and N remaining at day 360 in the L, M, T or L/T straw layers were significantly and linearly related to the average moisture content of these layers sampled over the year, regardless of the position of the layer within the mulch and the mulch quantity (Fig. 7). This relationship was also found when using the degradation constant k , which reflects all decomposition kinetics, with $k=0.0001x - 0.0005$ ($R^2=0.882$, $P<0.05$).

4. Discussion

4.1. Effect of mulch mass and thickness on mulch decomposition

A main hypothesis, based on recent published work, was that increasing the mass of mulch residues at the soil surface does not affect the decomposition rate of these mulches (Acosta et al., 2014; Halde and Entz, 2016; Dietrich et al., 2017). This situation was reflected in the present study with 3 amounts of sugarcane straw initially placed on the soil surface, which increased the initial mulch thickness (12S>8S>4S). We confirmed that the straw C and -N dynamics and calculated decomposition rates of the mulches were similar, regardless of the initial quantity of straw and whether they received 100 kg of fertilizer N 52 days after straw application. This result signifies that, although the proportion of the total mass in contact with the soil decreased as the mass increased, the decomposition was not modified. In fact, the initial thickness of the mulch varied from 1.9 cm to 4.7 cm, which is not a great difference. This may be due to our experimental system, which, although using hand-cut straw particles of a realistic decimetre length, minimized the thickness by "stacking" the straw in layers. In the field, mulches can be thicker but less dense. Meier et al. (2006) observed an initial sugarcane mulch thickness of about 11 cm for a 10 Mg ha⁻¹ mulch mass, decreasing rapidly in one year. Stott et al. (1990) measured that the initial wheat straw layer thickness ranged from 12 and 20 cm. Although mulch thickness is a parameter important for both the interpretation and the

modelling of mulch decomposition, data are still very rare. These results suggest that neither microbial colonization nor N availability were limiting factors of decomposition for the thickest mulches, which invalidates the hypothesis that soil-straw contact plays a major role as a key driver of mulch decomposition (Coppens et al., 2007; Thorburn et al., 2001). The initial thickness of the mulch was not proportional to the mass, due to a settling of the mulch with increasing weight of the residues. It should also be noted that the decrease was not linear over time, probably due to the effect of two combined factors, an initial possible and rather rapid settling under the influence of rainfall, combined with a more regular decrease under the effect of decomposition. It is noteworthy that the decrease in thickness was proportional to the initial thickness, reflecting the observed proportionality of the decomposition rate with the initial mass.

Strikingly, the N dynamics in mulch differed substantially from the C dynamics in mulch. After a synchronized, rapid C and N net loss at the beginning of decomposition, the N in the mulch remained stable while decomposition continued, and adding N onto the mulch in fertilized plots did not modify this dynamic. Sousa Jr et al. (2017) also observed little and transient effect of adding vinasse onto sugarcane mulch on decomposition rate. These results suggest the efficient use of straw N and recycling of microbial N by decomposers growing on the straw and that the availability of N to decomposers was not a limiting factor of decomposition, even with increased mulch masses and thicknesses and a lack of mineral N in the soil under the mulches in non-fertilized plot (Pinheiro et al., *accepted*). Some work has also highlighted the possibility of a significant contribution of asymbiotic N-fixation during decomposition of sugarcane litter. Lincoln and Vitousek (2016) calculated that the asymbiotic N fixed represented 17 to 40% of the N present in the litter. They pointed out that the potential N-fixation per unit litter depended on the thickness of the litter with an optimum for a 5cm-thick litter. This amount of fixed N could represent from 5 to 39 kg N ha⁻¹ year⁻¹ depending on the conditions tested. Patriquin (1982) earlier found that 10 to 30 kg N ha⁻¹ were potentially fixed during sugarcane litter decomposition. The possible contribution of non-symbiotic fixation to mulch N dynamics would merit further investigation, as it appears to be quantitatively important. The relatively high C:N ratio of sugarcane straw suggests also that fungi may be the main group of microorganisms

involved in decomposition, particularly for the upper layers of the mulch, due to their ability to translocate N from the soil through the hypha network to substrates that are recalcitrant or poor in N (Frey et al., 2000).

The amount of straw on the soil surface also affected soil temperature and moisture and had a substantial effect on the straw water content. Previous studies demonstrated that mulch moisture is driven by the water retention characteristics of residue particles and by rain and evaporation regimes (Lee et al., 2014; Iqbal et al., 2015). Few studies have evaluated the interaction between different amounts of straw and soil moisture (Cook et al., 2006). Our results clearly showed an increase in soil moisture with increasing straw masses on the soil surface, mainly in periods with low rainfall, when thicker mulch prevented soil water evaporation more effectively, confirming other studies (Sousa Jr et al. (2017). This is because the straw on the surface acts as a physical barrier to the transfer of heat and water to the atmosphere (Fuchs and Hadas, 2011). Despite rain water being retained by mulches, soil without mulch remained drier than soils covered by mulches. This is due to the fact that the increased evaporation of soil not covered by mulch (0S) was not offset by the higher amount of water reaching soil after the rains in such situation. It is understandable that this effect depends on several factors, not only on the intensity and frequency of rainfall, but also on the proportion of mulch-covered area and the characteristics of the mulch, and may vary according to agricultural and climatic conditions.

4.2. Water and decomposition gradients within the mulch

The amount of straw on the soil surface not only affected the overall soil and straw moisture but also created a moisture gradient within the mulch. Sampling straw layers in different positions in the mulch showed that the layers in contact with the soil in the thicker mulch (8S L and 12S L) were almost always wetter during the year than the more distant layers (8S T, 12S M and 12S T), which were exposed to solar radiation and wind. The higher moisture of the layers in contact with the soil was mainly due to the lower water loss by evaporation but may also be due to the absorption of water from soil by the straw in a phenomenon defined as the “sponge effect” (Kravchenko et al. 2017).

These moisture conditions favour high microbial activities and faster decomposition at the soil-mulch interface.

The kinetics of straw C and N for each layer position and each mulch mass treatment indicated that all layers decomposed, even at the top of the thicker mulch (12S T), which was the most distant layer from the soil and that the decomposition of the top layers started as quickly as the decomposition of the other layers. This suggests that part of the initial fast C and N losses could be due to the leaching of soluble compounds under rain. However, the change in the chemical composition of mulch over time, expressed here by the LCI index, occurred in all straw layers, regardless of position. The LCI corresponds to the relative enrichment of LIG in the plant cell walls of the residues (Whittinghill et al., 2012) because structural components such as CEL and HEM are more easily decomposed than LIG (Berg and McClaugherty, 2014). This confirms that the C or N loss of the top layers was not simply due to the possible physical transport of straw particles from an upper layer to a lower layer during experimental handling or by rain leaching; rather, remaining particles of each layer underwent decomposition. The high LCI value of the low 12S layer (12S L) also confirms that decomposition advanced more in this layer than in the other straw layers.

We observed that the top layers behaved similarly (12S T= 8S T= 4S L/T), while the decomposition speed of the low layers increased as the overlying mulch thickness increased (12S L>8S L>4S L/T). This pattern indicates that the top layers were under similar control of humectation-desiccation cycles, driven by climate, while the low layers differentiated from each other by the thickness of the overlying mulch, which in turn influenced the low-layer moisture. These findings are very well supported by the strong relationships between the amount of straw C or N remaining after 360 days (or the k degradation constant) and the mean straw moisture in each layer, irrespective of layer position and the mulch thickness. We conclude that a mulch system of various thicknesses is driven primarily by water dynamics, which is under the control of water retention-evaporation by the straw particles. The moisture gradient in the mulch depended on mulch thickness and determined decomposition gradient, which finally defined the decomposition rate of the whole mulch. This raises the question of understanding how the observed differences between layers perfectly balance in order

not to be able to find any difference in decomposition rate between mulches of different thicknesses.

Our assumption is that, in our experimental conditions (climate, range of DM), the higher decomposition rates of the low layers in thick mulches compensated for the large mass of these thick mulches not in direct contact with the soil. This “compensation” hypothesis is also supported by our results of water content of whole mulches (measured 9 times during the year), showing that although a gradient was established within the mulches, and transient differences appeared between mulches of increasing thickness, on average there were little or no differences in water content of the mulches regardless of their initial mass. These results confirm the hypothesis of Dietrich et al. (2017) that increasing mulch quantity has an antagonistic effect, increasing the proportion of straw less wet and not in direct contact with the soil but improving the environmental conditions of decomposition of the mulch at the soil-mulch interface. Williams et al. (2018), measuring the decomposition of varied quantities of surface rye straw, also obtained similar decomposition rates in the range of 5 to 15 Mg ha⁻¹. Conversely, Vasconcelos et al. (2018) and Sousa Jr et al. (2017), investigating sugarcane, showed that the decomposition rate increased with the amount of straw in the range 3-14 Mg DM ha⁻¹, suggesting that the positive effect of increased moisture prevailed in thicker mulches. However, the thickness and density of the mulches were not available in these studies; therefore, it is not possible to interpret the differences with our results. Thus, further research is needed to establish the mulch thickness range and rain patterns for which these results are valid and, in particular, better understand how the dynamics of mulch layers that are thicker than those explored here and in the literature could modify the above conclusions. It is expected that these results will strongly depend on climatic conditions and will particularly be affected by interactions between mulch properties and thickness and water dynamics in dry periods.

4.3. Consequences for much conceptual modelling

Many modules developed in several simulation models (i.e., APSIM, Thorburn et al., 2001; PASTIS, Findeling et al., 2007) represent residue mulch composed of two layers; one of these layers is in contact with the soil, undergoing decomposition with moisture, available N and microbial biomass

processes influenced by the underlying soil layer, and the other layer is undecomposed straw, which
"feeds" the lower layer as it empties. Moreover, in the model proposed by Findeling et al. (2007), the
water dynamics of mulch are simulated by considering the whole mass of straw that forms the mulch
to present the same moisture content. Our results demonstrated that upper mulch layers, which are not
in direct contact with the soil (exposed to air), also decompose and that there is a gradient of moisture
and decomposition (and probably temperature, although we did not measure this factor) throughout the
mulch volume. These findings suggest the need for more sophisticated models dedicated to the
simulation of mulch evolution, by considering the whole straw layer to be decomposing, albeit with a
gradient, and with an accurate estimation of the mulch water content over time, including knowledge
about water interception, retention and leaching for various crop residues (Iqbal et al. 2013; Sato et al.
2004).

5. Conclusions

By varying mulch thickness under similar climatic conditions, we demonstrated that not only
the mulch in contact with the soil but also the whole mulch decomposed, starting quickly after the
residues were deposited on the soil, with a vertical gradient of decomposition closely linked to the
moisture gradient. The moisture gradient of the mulch, which is under the control of humectation
(rain) and desiccation (evaporation) and depends on thickness, drives decomposition; therefore, it is
important to correctly model water properties of crop residues and water transport between soil, mulch
and atmosphere. Our results indicated the importance of soil-residue layer contact properties in driving
mulch decomposition. We also showed that, despite heterogeneity within the mulch, decomposition
was proportional to the initial amount of mulch mass for the studied crop. For the present range of
straw masses, as mulch thickness increased, the moisture and decomposition rate of the underlying
layers increased compensating for the increase in the fraction of mulch more distant from the soil and
drier. Considering the importance of these results for the dynamics of nutrient recycling and release in
non-tilled agrosystems, the ability of mulch decomposers to efficiently recycle N and the dynamics of
microbial colonization within the mulch should receive more attention. Finally, the accurate prediction

of water dynamics in soil-mulch systems is crucial because these systems act as hot spots for many processes, particularly gas emissions (NH_3 , N_2 and N_2O), which are under the control of moisture conditions.

Acknowledgements

This work was supported by the Brazil government through the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES). The bilateral Brazil-France collaboration was funded under Program CNPq-Ciência sem Fronteiras, Process Number 208415/2017-3 and by the Institut National de la Recherche Agronomique (INRA) - Environment & Agronomy Division.

Declarations of interest: none

References

- Acosta, J.A. de A., Amado, T.J.C., Silva, L.S. da, Santi, A., Weber, M.A., 2014. Fitomass decomposition and nitrogen release of cover crops in function of the level of residue input to soil under no-tillage system. *Cienc. Rural* 44, 801–809. doi: 10.1590/s0103-84782014005000002
- Angers, D.A., Recous, S., 1997. Decomposition of wheat straw and rye residues as affected by particle size. *Plant Soil* 189, 197-203. doi: 10.1023/A:1004207219678
- Berg, B. and McClaugherty, C., 2014. *Plant Litter: Decomposition, Humus Formation, Carbon Sequestration*, third ed. Springer-Verlag Berlin Heidelberg. doi: 10.1007/978-3-642-38821-7
- Bordonal, R.O., Carvalho, J.L.N., Lal, R., de Figueiredo, E.F., de Oliveira, B.G., Newton La Scala Jr, N., 2018. Sustainability of sugarcane production in Brazil. A review. *Agron. Sustain. Dev.* 38, 13. doi: 10.1007/s13593-018-0490-x
- Chen, H., Li, X., Hu, F., Shi, W., 2013. Soil nitrous oxide emissions following crop residue addition: A meta-analysis. *Glob. Change Biol.* 19, 2956–2964. doi:10.1111/gcb.12274

458 Christensen, B.T., 1986. Barley straw decomposition under field conditions - effect of placement and
initial nitrogen-content on weight-loss and nitrogen dynamics. *Soil Biol. Biochem.* 18, 523-529. doi:
460 doi.org/10.1016/0038-0717(86)90010-6

Cook, H.F., Valdes, G.S.B., Lee, H.C., 2006. Mulch effects on rainfall interception, soil physical
462 characteristics and temperature under *Zea mays* L. *Soil Tillage Res.* 91, 227-235. doi:
10.1016/j.still.2005.12.007

464 Coppens, F., Garnier, P., De Gryze, S., Merckx, R., Recous, S., 2006. Soil moisture, carbon and
nitrogen dynamics following incorporation and surface application of labeled crop residues in soil
466 columns. *Eur. J. Soil Sci.* 57, 894–905. doi: 10.1111/j.1365-2389.2006.00783.x

Coppens, F., Garnier, P., Findeling, A., Merckx, R., Recous, S., 2007. Decomposition of mulched
468 versus incorporated crop residues: Modelling with PASTIS clarifies interactions between residue
quality and location. *Soil Biol. Biochem.* 39, 2339–2350. doi:10.1016/j.soilbio.2007.04.005

470 Curtin, D., Francis, G.S., Mccallum, F.M., 2008. Decomposition rate of cereal straw as affected by
soil placement. *Aust. J. Soil Res.* 46, 152–160. doi: 10.1071/SR07085

472 Dietrich, G., Sauvadet, M., Recous, S., Redin, M., Pfeifer, I.C., Garlet, C.M., Bazzo, H., 2017.
Sugarcane mulch C and N dynamics during decomposition under different rates of trash removal.
474 *Agric. Ecosyst. Environ.* 243, 123-131. doi: 10.1016/j.agee.2017.04.013

Douglas Jr., C.L., Allmaras, R.R., Rasmussen, P.E., Ramig, R.E., Roager Jr., N.C., 1980. Wheat straw
476 composition and placement effects on decomposition in dryland agriculture of the Pacific Northwest.
Soil Sci. Soc. Am. J. 44, 833-837. doi: 10.2136/sssaj1980.03615995004400040035x

478 Findeling, A., Garnier, P., Coppens, F., Lafolie, F., Recous, S., 2007. Modelling water, carbon and
nitrogen dynamics in soil covered with decomposing mulch. *Eur. J. Soil Sci.* 58, 196–206.
480 doi:10.1111/j.1365-2389.2006.00826.x

Frey, S. D., Elliott, E.T., Paustian, K., Peterson, G.A., 2000. Fungal translocation as a mechanism for
 482 soil nitrogen inputs to surface residue decomposition in a no-tillage agroecosystem. *Soil Biol.*
Biochem. 32, 689–698, 2000. doi: 10.1016/S0038-0717(99)00205-9

484 Fuchs, M., Hadas, A., 2011. Mulch resistance to water vapor transport. *Agric. Water Manag.* 98, 990-
 998. doi: 10.1016/j.agwat.2011.01.008

486 Halde, C., Entz, M.H., 2016. Plant species and mulch application rate affected decomposition of cover
 crop mulches used in organic rotational no-till systems. *Can. J. Plant Sci.* 96, 59–71. doi:
 488 10.1139/cjps-2015-0095

Henriksen, T.M., Breland, T.A., 2002. Carbon mineralization, fungal and bacterial growth, and
 490 enzyme activities as affected by contact between crop residues and soil. *Biol. Fert. Soils* 35: 41-48.
 doi: 10.1007/s00374-001-0438-0

492 Iqbal A, Beaugrand J., Garnier P., Recous S., 2013. Tissue density determines the water storage
 characteristics of crop residues. *Plant Soil* 367, 285-299. doi: 10.1007/s11104-012-1460-8

494 Iqbal, A., Aslam, S., Alavoine, G., Benoit, P., Garnier, P., Recous, S., 2015. Rain regime and soil type
 affect the C and N dynamics in soil columns that are covered with mixed-species mulches. *Plant Soil*
 496 393, 319–334. doi:10.1007/s11104-015-2501-x

Kravchenko, A.N., Toosi, E.R., Guber, A.K., Ostrom, N.E., Yu, J., Azeem, K., Rivers, M.L.,
 498 Robertson, G.P., 2017. Hotspots of soil N₂O emission enhanced through water absorption
 by plant residue. *Nat. Geosci.* 10, 496–500. doi: 10.1038/ngeo2963

500 Lee, H., Fitzgerald, J., Hewins, D.B., McCulley, R.L., Archer, S.R., Rahn, T., Throop, H.L., 2014.
 Soil moisture and soil-litter mixing effects on surface litter decomposition: a controlled environment
 502 assessment. *Soil Biol. Biochem.* 72, 123–132. doi: 10.1016/j.soilbio.2014.01.027

Lincoln, N.K., Vitousek, P., 2016. Nitrogen fixation during decomposition of sugarcane (*Saccharum*
 504 *officinarum*) is an important contribution to nutrient supply in traditional agricultural systems of
 Hawai'i. *Int. J. Agr. Sustain.* 14, 2, 214-230. doi: 10.1080/14735903.2015.1071547

506 Meier E.A., Thorburn, P.J., Wegener, M.K., Basford, K.E., 2006. The availability of nitrogen from
sugarcane trash on contrasting soils in the wet tropics of North Queensland. *Nutr. Cycl. Agroecosyst.*
508 75, 101-114. doi: 10.1007/s10705-006-9015-0

Patriquin, D.G., 1982. Nitrogen fixation in sugar cane litter. *Biol. Agric. Hortic.*, 1, 39-64.
510 doi.org/10.1080/01448765.1982.9754376

Peyrard, C., Ferchaud, C. F., Mary, B., Gréhan, B.E., 2017. Management practices of *Miscanthus ×*
512 *giganteus* strongly influence soil properties and N₂O emissions over the long term. *Bioenergy Res.* 10,
208-224. doi: 10.1007/s12155-016-9796-1

514 Pinheiro, P.L., Recous, S., Dietrich, G., Weiler, D.A., Schu, A.L., Bazzo, H.L.S., Giacomini, S.J.,
N₂O emission increases with mulch mass in fertilized sugarcane cropping system. *Submitted to*
516 *Biol. Fert. Soils.* (accepted for publication)

Plantae, A.F., Parton, W.J., 2007. The dynamics of soil organic matter and nutrient cycling. In: Paul
518 E.A. (Ed.), *Soil microbiology, ecology, and biochemistry*. Academic Press, Amsterdam, pp 433-464

Putuhen, W., Cordery, I., 1996. Estimation of interception capacity of the forest floor. *J. Hydrol.* 180,
520 283-299. doi: 10.1016/0022-1694(95)02883-8

Sato, Y., Kumagai, T., Kume, A., Otsuki, K., Ogawa, S., 2004. Experimental analysis of moisture
522 dynamics of litter layers—the effects of rainfall conditions and leaf shapes. *Hydrol. Process.* 18, 3007-
3018. doi: 10.1002/hyp.5746

524 Schimel J. P., Gulledge, J.M., Clein-Curley, J.S., Lindstrom, J.E., Braddock, J.F., 1999. Moisture
effects on microbial activity and community structure in decomposing birch litter in the Alaskan taiga.
526 *Soil Biol. Biochem.* 31, 831-838. doi: 10.1016/S0038-0717(98)00182-5

Schmatz, R., Recous, S., Aita, C., Tahir, M.M., Schu, A.S., Chaves, B., Giacomini, S.J., 2017. Crop
528 residue quality and soil type influence the priming effect but not the fate of crop residue C. *Plant Soil*
414, 229–245. doi: 10.1007/s11104-016-3120-x

530 Schomberg, H.H., Steiner, J.L., Unger, P.W., 1994. Decomposition and nitrogen dynamics of crop
residues: residue quality and water effects. *Soil Sci. Soc. Am. J.* 58, 372–381. doi:
532 10.2136/sssaj1994.03615995005800020019x

Soil Survey Staff, 2010. *Keys to Soil Taxonomy*, 11th ed. USDA-Natural Resources Conservation
534 Service, Washington, DC.

Sousa Jr, J. G. de A., Cherubin, M.R., Cerri, C.E.P., Cerri, C.C., Feigl, B.J., 2017. Sugar cane straw
536 left in the field during harvest: decomposition dynamics and composition changes. *Soil Res.* 55, 758-
768. doi: 10.1071/SR16310

538 Steiner, J.L., Schomberg, H.H., Unger, P.W., Cresap, J., 1999. Crop residue decomposition in no-
tillage small-grain fields. *Soil Sci. Soc. Am. J.*, 63, 1817-1824. doi: 10.2136/sssaj1999.6361817x

540 Stott, D.E., Stroo, H.F., Elliott, L.F., Papendick, R.I., Unger, P.W., 1990. Wheat residue loss from
fields under no-till management. *Soil Sci. Soc. Am. J.*, 54, 92-98.
542 doi:10.2136/sssaj1990.03615995005400010014x

Thorburn, P.J., Probert, M.E., Robertson, F.A., 2001. Modelling decomposition of sugar cane surface
544 residues with APSIM–Residue. *Field Crops Res.* 70, 223–232. doi:10.1016/S0378-4290(01)00141-1

Tiecher, T., Calegari, A., Caner, L., Rheinheimer, D. dos S., 2017. Soil fertility and nutrient budget
546 after 23-years of different soil tillage systems and winter cover crops in a subtropical Oxisol.
Geoderma 308, 78-85. doi: 10.1016/j.geoderma.2017.08.028

548 Van Soest, P.J., 1963. Use of detergents in the analysis of fibrous feeds I: preparation of fiber residues
of low nitrogen content. *J. Assoc. Off. Anal. Chem.* 46, 825-835.

550 Vasconcelos, A.L.S., Cherubin, M.R., Feigl, B.J., Cerri, C.E.P., Gmach, M.R., Siqueira-Neto, M.,
2018. Greenhouse gas emission responses to sugarcane straw removal. *Biomass Bioenergy* 113, 15–
552 21. doi: 10.1016/j.biombioe.2018.03.002

- Whittinghill, K.A., Currie, W.S., Zak, D.R., Burton, A.J., Pregitzer, K.S., 2012. Anthropogenic N
554 Deposition Increases Soil C Storage by Decreasing the Extent of Litter Decay: Analysis of Field
Observations with an Ecosystem Model. *Ecosyst.* 15, 450–461. doi:10.1007/s10021-012-9521-7
- 556 Williams, A., Wells, M.S., Dickey, D.A., Hu, S., Maul, J., Raskin, D.T., Reberg-Horton, S.C.,
Mirsky, S.B., 2018. Establishing the relationship of soil nitrogen immobilization to cereal rye residues
558 in a mulched system. *Plant Soil* 426, 95-107. doi: 10.1007/s11104-018-3566-0

Figure Captions

Fig. 1. Schematic layout of the field plots demonstrating the locations of soil water content and soil temperature sensors (2.5 and 7.5 cm depth) and the litter boxes, with emphasis on the different straw layers (L: low, M: middle and T: top) forming the three quantities of straw (4S: 4 Mg ha⁻¹, 8S: 8 Mg ha⁻¹ and 12S: 12 Mg ha⁻¹).

Fig. 2. Precipitation and air temperature (a), relative humidity (b), soil temperature (c) and soil moisture (d) at 2.5 cm depth for four quantities of straw (0S, 4S, 8S and 12S, i.e., 0, 4, 8 and 12 Mg ha⁻¹, respectively) over the 360 days of the experiment.

Fig. 3. Evolution of the thickness (a) and water content of the mulch over the 360 days of the experiment (b) and water contents of the mulch layers at two sampling dates and on average for the year (c). 4S, 8S and 12S correspond to 4, 8 and 12 Mg ha⁻¹, respectively, and L, M and T correspond to the low, middle and top positions, respectively, in the mulch. The vertical bars indicate the minimum significant difference between treatments (Tukey's test at $P < 0.05$). Values are the mean of 36 replicates for thickness or 4 replicates for straw water content.

Fig. 4. Dynamics of total C (a), total N (b), water-soluble C (c) and water-soluble N (d) in the remaining mulch particles for three quantities of straw (4S: 4 Mg ha⁻¹, 8S: 8 Mg ha⁻¹ and 12S: 12 Mg ha⁻¹). Data are the average of the treatments with 0 (0N) and 100 (100N) kg N ha⁻¹. The vertical bars in the graph indicate the minimum significant difference between treatments (Tukey's test at $P < 0.05$). Values are the mean of 8 replicates per treatment and per date.

Fig. 5. C and N remaining in the mulch straw layers: (a,b) and (c,d) show different positions in the mulch (L: low, M: middle, T: top) for the 12S and 8S treatments, respectively; the straw layer is located on the top of the mulch in (e,f) and on the bottom of the mulch in (g,h) for the three treatments (4S: 4 Mg ha⁻¹, 8S: 8 Mg ha⁻¹ and 12S: 12 Mg ha⁻¹). The vertical bars in the graph indicate the minimum significant difference between treatments (Tukey's test at $P < 0.05$). The values are the mean of 8 replicates per treatment and per sampling date.

Fig. 6. Evolution of the C:N ratio (a) and of the lignocellulose index (LCI) (b) of the straw for the three layer positions in the mulch (L: low, M: middle, T: top) for the 12S treatment (12 Mg ha⁻¹). The vertical bars indicate the minimum significant difference between treatments (Tukey's test at $P < 0.05$). Data are the mean of 8 replicates per treatment and per sampling date.

Fig. 7. Relationships between straw C or straw N and the average water content of straw layers with different positions in the mulch (L: low, M: middle, T: top) for the three treatments (4S: Mg ha⁻¹, 8S: Mg ha⁻¹ and 12S: Mg ha⁻¹) over the 360 days of the experiment. Values are the mean of 4 replicates per layer on 9 sampling dates.

Fig. 1.

Fig. 2.

Fig. 3.

Fig. 4.

Fig. 5.

1

Fig. 6.

2

3

4

Fig. 7.

1 **Table 1**

2 Decomposition rate (k) and mean residence time ($t_{1/2}$) for the mulch with three straw quantity (4, 8 and 12 Mg ha⁻¹) and two N rate (0 and 100 kg ha⁻¹) and
3 mulch straw layers (L: low, M: middle, T: top).

Parameters	Initial straw quantity (Mg ha ⁻¹)			N rate (kg ha ⁻¹)		Straw layers					
	4	8	12	0	100	4S L/T	8S T	8S	12S T	12S M	12S L
C_0 (Mg ha ⁻¹) ^a	1.47c	2.99b	4.45b	2.97a	2.97a	1.47ab	1.51a	1.51a	1.43b	1.52a	1.52a
k (day ⁻¹) ^b	0.0042a	0.0040a	0.0046a	0.0045a	0.0044a	0.0042b	0.0037bc	0.0055a	0.0033c	0.0046b	0.0064a
$t_{1/2}$ (day ⁻¹) ^c	166a	155a	153a	158a	159a	166bc	188ab	126de	209 ^a	155cd	110e

4 Values followed by the same letters are not significantly different (Tukey at $P<0.05$).

5 ^a C_0 is straw C pool.

6 ^b k is the degradation rate constant.

7 ^c $t_{1/2}$ is the half-life of the mulch.

8