Assume-guarantee contracts for continuous-time systems

Adnane Saoud, Antoine Girard, Laurent Fribourg

► To cite this version:

Adnane Saoud, Antoine Girard, Laurent Fribourg. Assume-guarantee contracts for continuous-time systems. *Automatica*, 2021, 134, 10.1016/j.automatica.2021.109910 . hal-02196511v3

HAL Id: hal-02196511

<https://hal.science/hal-02196511v3>

Submitted on 24 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Assume-guarantee contracts for continuous-time systems [★]

Adnane Saoud ^{a,b} Antoine Girard ^a Laurent Fribourg ^b

^a *Université Paris-Saclay, CNRS, CentraleSupélec
Laboratoire des Signaux et Systèmes
91190, Gif-sur-Yvette, France*

^b *Université Paris-Saclay, CNRS, ENS Paris-Saclay
Laboratoire de Méthodes Formelles
91190, Gif-sur-Yvette, France*

Abstract

Many techniques for verifying properties of continuous-time systems are limited to systems of moderate size. In this paper, we propose an approach based on assume-guarantee contracts and compositional reasoning for verifying properties of a broad class of continuous-time systems consisting of interconnected components. The notion of assume-guarantee contracts makes it possible to divide responsibilities among the system components: a contract specifies the property that a component must fulfill under some assumptions on the behavior of its environment (i.e. of the other components). We define weak and strong semantics of assume-guarantee contracts.^c We then establish a certain number of results for compositional reasoning, which allow us to show that a global assume-guarantee contract of the whole system is satisfied when all components satisfy their own contracts. We show that the weak satisfaction of the contract is sufficient to deal with interconnections described by a directed acyclic graph, while strong satisfaction is needed to reason about general interconnections containing cycles. Specific results for systems described by differential inclusions and invariance assume-guarantee contracts are then developed. Finally, we show how the proposed assume-guarantee framework can recast different versions of the small-gain theorem as a particular case. Throughout the paper, the main results are illustrated using simple examples.

Key words: component-based design, assume-guarantee contracts, prefix-closed properties, small-gain theorem.

1 Introduction

Cyber-physical systems (CPS) result from integrations of computational devices with physical processes and are to become ubiquitous in modern societies (autonomous vehicles, smart buildings, robots, etc.) (see [25] and the references therein). Despite considerable progress in the field, current techniques apply to system of moderate complexity (the complexity is quantified by the number of interacting components). Thus, the design of complex

CPS requires to divide large design problems in smaller sub-problems that can be solved using existing tools.

Compositional approaches for the analysis and the design of dynamical systems have been long known in the field of control theory, where the celebrated small-gain theorem [12,19,21] makes it possible to prove stability of a system from the stability of its components. Other compositional approaches for the analysis and design of cyber-physical systems have been mainly initiated in the field of computer science [1,15,18].

The study of properties of dynamical systems using decentralized approaches has been an ongoing research area in recent years [3,8,9,13,29,30,37]. Other compositional approaches, using formal methods and symbolic techniques, are presented in [10,22,24,26–28,32,35,38,40]. All these works develop efficient computational techniques by making specific assumptions on the classes of dynamical systems and of properties to which they can be applied.

[★] A preliminary version of this paper was presented at the 16th European Control Conference. This research was supported by Labex DigiCosme (project ANR-11-LABEX-0045-DIGICOSME) operated by ANR as part of the program "Investissement d'Avenir" Idex Paris Saclay (ANR-11-IDEX-0003-02). This project was partially supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 725144). Corresponding author A. Saoud; e-mail: Adnane.Saoud@l2s.centralesupelec.fr, Antoine.Girard@l2s.centralesupelec.fr, fribourg@lsv.ens-cachan.fr

In the current work, we aim at proposing a general theoretical framework and thus we make weak assumptions on systems and properties. We initiate a high-level framework for verifying properties of complex systems, consisting of interconnected components, using a contract-based approach [6]. Each component is assigned an assume-guarantee contract, which specifies the property that the component must fulfill under assumption about its environment (i.e. the other components). We introduce contracts and define weak and strong semantics. We then establish results that allow us to reason compositionally using assume-guarantee contracts: i.e. if all components satisfy their own contract then a global contract of the whole system is satisfied. We show that the weak satisfaction of the contract is sufficient to deal with interconnections described by a directed acyclic graph, while strong satisfaction is needed to reason about general interconnections containing cycles. We then show that for systems described by differential inclusions and invariance assume-guarantee contracts, weak satisfaction of contracts is sufficient to reason on general interconnections. Finally, we show how the proposed assume-guarantee framework can recast different versions of the small-gain theorem as a particular case.

There are several advantages in using contract based design for CPS. Firstly, by dividing a complex design problem into several smaller sub-problems, one is able to address design challenges that would be out of reach of current state-of-the-art design methods. Secondly, contract-based design makes it possible to replace a component without jeopardizing the behavior of the overall system: one just has to make sure that the new component satisfies the assigned contract. Thirdly, components are reusable when similar contracts appear in the decomposition of a global contract.

Let us emphasize that the main objective of the paper is to provide compositionality rules on how to go from the satisfaction of local contracts for the components to the satisfaction of a global contract for the whole interconnected system. The question on how to construct the local contracts from the global one is not investigated in this paper, but have been explored in other works in the literature for continuous-time in [41] and [6, Section 8.6], and for discrete-time [7,14,16] systems.

The present paper focuses on the theoretical development of a general framework for contract-based reasoning. Applications of this framework to the design of symbolic controllers are reported in [32,33,41].

Related work: Contract based design for CPS have been widely studied in [6]. Our approach differs from the one proposed in [6] in several directions:

- In [6] the contracts need to be checked for the whole time domain on which trajectories are defined. In this

paper, the satisfaction of contracts is defined in a progressive and time-dependant way, which makes it possible to detect the instant at which the contract is violated.

- While in [6] assumptions and guarantees are defined over all the set of variables, namely inputs and outputs, in this paper, we focus on the case where assumptions are defined on the inputs and guarantees on the states and outputs. Indeed, this restriction allows us to provide simple compositionality results.
- The use of saturated contracts is crucial in applying the contract framework proposed in [6], which requires being able to compute unions and complements of different assertions. In this paper, saturation of the contracts is not required, and one can deal directly with the contracts as they are.

Assume-guarantee reasoning has been previously used in control theory. The authors in [20] presented a compositionality result for linear dynamical systems based on the notion of simulation introduced in [39]. In spirit, our work is closer to the framework presented in [23] for verifying general properties using parametric assume-guarantee contracts and compositional reasoning by means of small-gain theorems. Our approach differs from the one proposed in [23] in the following directions:

- The approach proposed in [23] follows the framework of [6], which makes a noticeable difference with our results;
- In [23], the results hold only for discrete-time systems, while our approach makes it possible to deal with both continuous and discrete-time systems.
- The main compositionality result in [23] requires to assume that at least one component satisfies a contract (for some parameter value), independently of the behavior of other components. This breaks the circularity of implications of the assume-guarantee contracts, which is arguably the main difficulty in contract-based design, and the reason why we introduce weak and strong semantics for assume-guarantee contracts, which has not been done in [23].
- In [23], the authors show how their framework makes it possible to reprove the classical bounded input bounded output small-gain theorem for discrete-time systems. In our paper we show how the proposed framework allows to recast the bounded input bounded output small gain theorem for continuous-time systems. Moreover, we provide a new small-gain result for the notion of growth bound, which has not been proposed before in the control literature.

The paper is organized as follows. In Section 2, we introduce the class of prefix-closed properties. In Section 3, we introduce the class of systems and interconnections considered throughout the paper. In Section 4, we introduce assume-guarantee contracts, their weak and strong semantics. We then establish compositionality results for reasoning about interconnected systems in Section 5.

In Section 6, we develop specific results for systems described by differential inclusions and invariance assume-guarantee contracts. Finally, in Section 7, we show that different versions of the classical small-gain theorem can be recast as particular applications of our framework. Throughout the paper, simple examples are used as illustrations of the main results.

Due to space constraints, the results for discrete-time systems and contracts are omitted and can be found in [36, Chapter 2]. A preliminary version of this work has been presented in the conference paper [34]. The current paper extends the approach in different directions. First, we generalize the approach from cascade and feedback compositions to any composition structure and from invariance to more general properties. Second, while in [34], we have shown that for systems described by Lipschitz differential inclusions, the notion of invariance relative to assume-guarantee contracts implies the weak satisfaction. In the current work, we show the converse result. Finally, in the current work, we show that different versions of the classical small-gain theorem can be recovered using our framework.

Notation: $\mathbb{R}, \mathbb{R}_0^+, \mathbb{R}^+$ and $\mathbb{N}$ denote the set of reals, non-negative reals, positive reals and positive integers respectively. The set of continuous-time domains is $\mathbb{E}(\mathbb{R}_0^+) = \{[0, a], a \in \mathbb{R}_0^+\} \cup \{[0, a), a \in \mathbb{R}^+\} \cup \{\mathbb{R}_0^+\}$. For $Z \subseteq \mathbb{R}^n$, we denote by $M(Z)$ the set of continuous-time maps $z : E \rightarrow Z$, where $E \in \mathbb{E}(\mathbb{R}_0^+)$. For $q \in \mathbb{N}_0^+$, $\mathcal{D}^q$ denotes the set of continuous-time maps q times differentiable. For $x \in \mathbb{R}^n$, $\|x\|$ denotes the infinity norm. For a continuous-time map $x : E \rightarrow \mathbb{R}^n$ and for $t \in E$, $\|x\|_{[0, t]}\infty$ denotes the essential supremum of $x : E \rightarrow \mathbb{R}^n$ on $[0, t]$. Given a space X equipped with a metric $d : X \times X \rightarrow \mathbb{R}_0^+$, the closure of the set $A \subseteq X$ is denoted $\text{cl}(A)$ and its complement is denoted $\overline{A}$. For $\varepsilon > 0$ and $x \in X$, the ball with center in x and a radius ε is $\mathcal{R}_\varepsilon(x) = \{y \in X \mid d(x, y) \leq \varepsilon\}$. A continuous function $\gamma : \mathbb{R}_0^+ \rightarrow \mathbb{R}_0^+$ is said to belong to class $\mathcal{K}$ if it is strictly increasing and $\gamma(0) = 0$; γ is said to belong to class $\mathcal{K}_\infty$ if γ is $\mathcal{K}$ and $\gamma(r) \rightarrow \infty$ as $r \rightarrow \infty$. A continuous function $\beta : \mathbb{R}_0^+ \times \mathbb{R}_0^+ \rightarrow \mathbb{R}_0^+$ is said to belong to class $\mathcal{KL}$ if, for each fixed s , the map $\beta(\cdot, s)$ belongs to class $\mathcal{K}$, and for each fixed nonzero r , the map $\beta(r, \cdot)$ is strictly decreasing and $\beta(r, s) \rightarrow 0$ as $s \rightarrow \infty$. Given a set X , 2^X is used to denote its power set.

2 Preliminaries on prefix-closed sets

Given a set $Z \subseteq \mathbb{R}^n$, prefix closed sets are subsets $P \subseteq M(Z)$ that can be defined as follows: if a trajectory $z : E \rightarrow Z$ belongs to the prefix-closed set P , then any prefix of z belongs to P . In this part, we first give a formal definition of a prefix-closed set, we then give a necessary and sufficient condition for a set to be prefix-closed, finally we give some examples of such sets.

Definition 1 Let $Z \subseteq \mathbb{R}^n$. Let $z : E \rightarrow Z$ and $z' : E' \rightarrow Z$ in $M(Z)$. z is said to be a prefix of z' and denoted $z \in \text{pref}(z')$ if $E \subseteq E'$ and for all $t \in E$, $z(t) = z'(t)$. In this case, z can be seen as a restriction of z' and is also denoted $z = z'|_E$.

This notion is generalized toward sets of continuous-time maps in the usual way: for $A \subseteq M(Z)$, $\text{pref}(A) = \bigcup_{z \in A} \text{pref}(z)$.

Definition 2 Let $Z \subseteq \mathbb{R}^n$ and $P \subseteq M(Z)$. P is said to be prefix-closed if the following logical implication is satisfied: $z \in P$ and $\hat{z} \in \text{pref}(z) \Rightarrow \hat{z} \in P$.

In the following we will give a characterization of prefix-closed sets.

Proposition 1 Let $Z \subseteq \mathbb{R}^n$ and $P \subseteq M(Z)$. P is prefix-closed if and only if $\text{pref}(P) = P$.

PROOF. Suppose that $\text{pref}(P) = P$ and let us prove that P is prefix-closed. Let $z \in P$ and $\hat{z} \in \text{pref}(z)$. Since $z \in P$, we have $\hat{z} \in \text{pref}(z) \subseteq \text{pref}(P) = P$. Then, $\hat{z} \in P$ and P is prefix-closed. Now suppose that P is prefix-closed and let us prove that $\text{pref}(P) = P$. The inclusion $P \subseteq \text{pref}(P)$ is verified by definition of the prefix. Let $\hat{z} \in \text{pref}(P)$, then there exists $z \in P$ such that $\hat{z} \in \text{pref}(z)$. Since P is prefix-closed we get $\hat{z} \in P$. Then, $\text{pref}(P) \subseteq P$ which ends the proof. $\square$

It can be seen that our notion of prefix-closed sets is relatively close to the notion of safety for linear time properties in [5]. Indeed, the only difference is that we are considering finite and infinite trajectories, while the linear time properties are defined in [5] over infinite words. In the following we give some examples of prefix-closed sets. This notion allows us to represent different type of properties such as invariance or systems described by differential inclusions.

Example 1 (Invariance) Let the set $S \subseteq \mathbb{R}^n$ such that $S \neq \emptyset$ and let us define:

$$A = \{z : E \rightarrow \mathbb{R}^n \in M(\mathbb{R}^n) \mid \forall t \in E, z(t) \in S\}.$$

The set A represents the set of continuous-time trajectories that belongs to the set S for all time.

Example 2 Let $S_1, S_2, \dots, S_q \subseteq \mathbb{R}^n$ such that for all $i \in \{1, 2, \dots, q\}$, $S_i \neq \emptyset$ and let us define:

$$A = \{z : E \rightarrow \mathbb{R}^n \in M(\mathbb{R}^n) \cap \mathcal{D}^q \mid \forall i \in \{1, 2, \dots, q\}, \forall t \in E, z^{(i)}(t) \in S_i\}$$

Where $z^{(i)}$ denotes the i th derivative of z . The set A in this example represents the set of continuous-time trajectories for which the i -th time derivative, $i \in \{1, 2, \dots, q\}$, belongs to the set S_i for all time.

Example 3 (Differential inclusion) Let $F : \mathbb{R}^n \rightarrow 2^{\mathbb{R}^n}$ be a set-valued map we define:

$$A = \{z : E \rightarrow \mathbb{R}^n \in M(\mathbb{R}^n) \cap \mathcal{D}^1 \mid \forall t \in E, \dot{z}(t) \in F(z(t))\}$$

An example of a non prefix-closed property is the reachability property described as follows:

Example 4 Let the set $K \subseteq \mathbb{R}^n$ such that $K \neq \emptyset$ and let us define:

$$A = \{z : E \rightarrow \mathbb{R}^n \in M(\mathbb{R}^n) \mid \exists t \in E, z(t) \in K\}.$$

3 Systems and interconnections

3.1 Systems

In this section, we introduce the classes of systems and interconnections considered throughout this paper, it is important to note that the classes of systems used in the paper are quite general, and includes deterministic and nondeterministic systems, described by difference or differential equations and inclusions and allows us to deal with phenomena such as sampling, time delays.

Definition 3 A continuous-time system is a tuple $\Sigma = (W_1, W_2, X, Y, \mathcal{T})$ where

- $W_1 \subseteq \mathbb{R}^{m_1}$, $W_2 \subseteq \mathbb{R}^{m_2}$, $X \subseteq \mathbb{R}^n$ and $Y \subseteq \mathbb{R}^p$, are the sets in which external and internal inputs, states, and outputs take their values;
- $\mathcal{T} \subseteq M(W_1 \times W_2 \times X \times Y)$ is a set of continuous-time trajectories.

3.2 Interconnections

We first introduce some notations for interconnected systems. A network of systems consists of a collection of $N \in \mathbb{N}_{>0}$ systems $\{\Sigma^1, \dots, \Sigma^N\}$, a set of vertices $I = \{1, \dots, N\}$ and a binary connectivity relation $\mathcal{I} \subseteq I \times I$ where each vertex $i \in I$ is labelled with the system Σ^i . For $i \in I$, we define $\mathcal{N}(i) = \{j \in I \mid (j, i) \in \mathcal{I}\}$ as the set of neighbouring components from which the incoming edges originate. We define $I_{\text{init}} = \{i \in I \mid \mathcal{N}(i) = \emptyset\}$ as the set of components for which there exist no incoming edge.

Given a directed graph $\mathcal{G} = (I, \mathcal{I})$ over the set of vertices $I = \{1, \dots, N\}$ and binary connectivity relation $\mathcal{I}$. A walk is a sequence $\sigma = a_1 a_2 \dots a_m$ such that for all

$i \in \{1, \dots, m-1\}$, $(a_i, a_{i+1}) \in \mathcal{I}$, in such case we say that a_i is an element of σ . For a walk σ , if the vertices $a_1, \dots, a_{m-1}$ are distinct and $a_1 = a_m$, then σ is a cycle. The set of all cycles is denoted $\zeta = \{\zeta_1, \dots, \zeta_p\}$, $p \in \mathbb{N}$.

Remark 1 We recall that for a directed graph, if we drop one edge of every cycle in the graph, a directed acyclic graph (DAG) denoted $\mathcal{G}^{\text{DAG}}$ is obtained. In such case, the set of initial vertices for the new directed acyclic graph is characterized by $I_{\text{init}}^{\text{DAG}} \subseteq I_{\text{init}} \cup A$, where $A \subseteq I$ is the set of vertices to which we dropped an edge. An illustration of this approach is given in Figure 1.

Fig. 1. Left: A graph $\mathcal{G}$ of four vertices, containing two cycles. The set of initial vertices is $I_{\text{init}} = \{4\}$. Right: A new DAG graph is constructed by removing dashed edges $\mathcal{G}^{\text{DAG}}$. The set of initial vertices is $I_{\text{init}}^{\text{DAG}} = \{4, 3\} \subseteq I_{\text{init}} \cup A = \{4\} \cup \{2, 3\}$.

In the following, we consider interconnections of continuous-time systems defined as follows:

Definition 4 Let $\{\Sigma^i\}_{i \in I}$ be a collection of systems and $\mathcal{I} \subseteq I \times I$ a binary connectivity relation. We say that $\{\Sigma^i\}_{i \in I}$ is compatible for composition with respect to $\mathcal{I}$, if for each $i \in I$, we have $\prod_{j \in \mathcal{N}(i)} Y^j \subseteq W_2^i$, i.e., the internal input space of Σ^i is a superset of the Cartesian product of the output spaces of all the neighbors in $\mathcal{N}(i)$. The composed system Γ denoted $\langle (\Sigma^i)_{i \in I}, \mathcal{I} \rangle$, is given by a tuple $\Gamma = (W_1, \{0\}, X, Y, \mathcal{T})$ where:

- $W_1 = \prod_{i \in I} W_1^i$;
- $X = \prod_{i \in I} X^i$;
- $Y = \prod_{i \in I} Y^i$;
- $(w_1, 0, x, y) : E \rightarrow W_1 \times \{0\} \times X \times Y \in \mathcal{T}$ is a trajectory of Γ if and only if for all $i \in I$, there exists a trajectory $(w_1^i, w_2^i, x^i, y^i) : E \rightarrow W_1^i \times W_2^i \times X^i \times Y^i \in \mathcal{T}^i$ of Σ^i with $w_1 = (w_1^1, \dots, w_1^N)$ and such that the internal inputs are constrained by the relation

$$w_2^i(t) = (y^{j_1}(t), \dots, y^{j_p}(t)) \text{ where } \mathcal{N}(i) = \{j_1, \dots, j_p\}$$

for all $i \in I$ and for all $t \in E$.

By abuse of notation, the constraint on the internal inputs will be written as $w_2^i = \prod_{j \in \mathcal{N}(i)} \{y^j\}$.

Remark 2 Let us remark that in the proposed interconnection structure, all the internal inputs of a system are outputs of other systems. Then, the composed system $\Gamma =$

Fig. 2. An illustration of the weak satisfaction of a contract. Top: trajectories of the external w_1 and internal w_2 inputs. Bottom: trajectories of the state x and output y .

$\langle (\Sigma^i)_{i \in I}, \mathcal{I} \rangle$ has trivial null internal inputs. Hence, with an abuse of notation, we will denote $\Gamma = (W_1, X, Y, \mathcal{T})$ and $(w_1, x, y) \in \mathcal{T}$, with $(w_1, x, y) : E \rightarrow W_1 \times X \times Y$ as a trajectory of Γ . Similarly, all initial elements Σ_i , where $i \in I_{init}$ have trivial null internal inputs and we use the same notation for their trajectories.

We should emphasize that trajectories of systems need not be defined for all time, $\mathbb{R}_0^+$. This makes it possible to avoid forward-completeness issues related to systems composition as shown in the following example.

Example 5 Let us consider the system $\Sigma^1 = (W_1, W_2, X, Y, \mathcal{T})$ where $W_1 = \{0\}$, $W_2 = X = Y = \mathbb{R}$. A trajectory of Σ^1 is a quadruple $(0, w_2, x, y) : E \rightarrow W_1 \times W_2 \times X \times Y$ in $\mathcal{T}$ where $E \in \mathbb{E}(\mathbb{R}_0^+)$, w_2 is continuous, x and y are differentiable and such that $x(0) = 1$ and for all $t \in E$,

$$\begin{cases} \dot{x}(t) = w_2(t) \\ y(t) = (x(t))^2. \end{cases}$$

Let $I = \{1\}$ and the interconnection relation $\mathcal{I} = \{(1, 1)\}$. It is clear that $\{\Sigma^i\}_{i \in I}$ is compatible for composition with respect to $\mathcal{I}$. It can be seen that Σ^1 , has trajectories defined on the whole time domain $\mathbb{R}_0^+$. However, if we only consider those trajectories, the set of trajectories $\mathcal{T}_\Gamma$ of the composed system $\Gamma = \langle (\Sigma^i)_{i \in I}, \mathcal{I} \rangle$ would be empty since the trajectories of $\mathcal{T}_\Gamma$ are of the form $(0, x, y) : E \rightarrow W_1 \times X \times Y$ where $E \subseteq [0, 1)$, and for all $t \in E$, $x(t) = \frac{1}{1-t}$ and $y(t) = \frac{1}{(1-t)^2}$.

4 Assume-guarantee contracts

An assume-guarantee contract is a compositional tool that specifies how a system behaves under assumptions about its inputs [6]. The use of assume-guarantee contracts makes it possible to reason on a global system

Fig. 3. An illustration of the strong satisfaction of a contract. Top: trajectories of the external w_1 and internal w_2 inputs. Bottom: trajectories of the state x and output y .

based on properties of its components. In this section, we introduce assume-guarantee contracts to reason on properties for continuous-time systems. These contracts are equipped with a weak and a strong semantics, which will allow us to establish compositionality results.

We define contracts for continuous-time systems as follows:

Definition 5 Let $\Sigma = (W_1, W_2, X, Y, \mathcal{T})$ be a continuous-time system, an assume-guarantee contract for Σ is a tuple $\mathcal{C} = (A_{W_1}, A_{W_2}, G_X, G_Y)$ where

- $A_{W_1} \subseteq M(W_1)$ and $A_{W_2} \subseteq M(W_2)$ are sets of assumptions on the external and internal inputs;
- $G_X \subseteq M(X)$ and $G_Y \subseteq M(Y)$ are sets of guarantees on the states and outputs.

We say that Σ (weakly) satisfies $\mathcal{C}$, denoted $\Sigma \models \mathcal{C}$, if for all trajectories $(w_1, w_2, x, y) : E \rightarrow W_1 \times W_2 \times X \times Y$ in $\mathcal{T}$:

- for all $t \in E$, if $w_1|_{[0,t]} \in A_{W_1}$ and $w_2|_{[0,t]} \in A_{W_2}$, then:
 - $x|_{[0,t]} \in G_X$;
 - $y|_{[0,t]} \in G_Y$.

We say that Σ strongly satisfies $\mathcal{C}$, denoted $\Sigma \models_s \mathcal{C}$, if for all trajectories $(w_1, w_2, x, y) : E \rightarrow W_1 \times W_2 \times X \times Y$ in $\mathcal{T}$:

- if $w_1|_{[0,0]} \in A_{W_1}$ then $y|_{[0,0]} \in G_Y$
- for all $t \in E$, if $w_1|_{[0,t]} \in A_{W_1}$ and $w_2|_{[0,t]} \in A_{W_2}$, then:
 - $x|_{[0,t]} \in G_X$;
 - there exists $\delta > 0$ such that for all $s \in [0, \delta]$ $y|_{[0,t+s] \cap E} \in G_Y$.

$\Sigma \models_s \mathcal{C}$ obviously implies $\Sigma \models \mathcal{C}$. An assume-guarantee contract for a continuous-time system states that if the

restriction of the external and internal inputs to the system up to a time $t \in \mathbb{R}_0^+$ belongs to A_{W_1} and A_{W_2} , respectively, then the restriction of the state of the system up to time t belongs to G_X , and the restriction of the output of the system up to time t (or up to a time $t + s$ with $s \in [0, \delta]$ and $\delta > 0$, in the case of strong satisfaction) belongs to G_Y . We must highlight that the value of δ may depend on the trajectory $(w_1, w_2, x, y) \in \mathcal{T}$ and on the value of the time instant $t \in E$. An illustration of the weak and strong satisfactions is given in Figures 2 and 3. One may remark that if the set of guarantees on the outputs G_Y is prefix closed, the notion of strong satisfaction of a contract can be defined by: Σ strongly satisfies $\mathcal{C}$, if for all trajectories $(w_1, w_2, x, y) : E \rightarrow W_1 \times W_2 \times X \times Y$ in $\mathcal{T}$:

- if $w_1|_{[0,0]} \in A_{W_1}$ then $y|_{[0,0]} \in G_Y$
- for all $t \in E$, if $w_1|_{[0,t]} \in A_{W_1}$ and $w_2|_{[0,t]} \in A_{W_2}$, then:
 - $x|_{[0,t]} \in G_X$;
 - there exists $\delta > 0$ such that $y|_{[0,t+\delta] \cap E} \in G_Y$.

Remark 3 Similarly to Remark 2, a contract for the composed system $\Gamma = \langle (\Sigma^i)_{i \in I}, \mathcal{I} \rangle$ has trivial null assumptions on internal inputs. Hence, with an abuse of notation, a contract for the composed system Γ will be denoted $\mathcal{C} = (A_{W_1}, G_X, G_Y)$.

5 Compositional reasoning

In this section we provide results allowing us to reason about interconnected systems based on contracts satisfied by the components.

5.1 Acyclic interconnections

We first provide the following result on the composition of assume-guarantee contracts, where the interconnection graph $\mathcal{G}$ between the components is a DAG.

Theorem 1 Consider a network of continuous-time components $\{\Sigma^i\}_{i \in I}$ compatible for composition with respect to $\mathcal{I}$. Let $\Gamma = \langle (\Sigma^i)_{i \in I}, \mathcal{I} \rangle$ be the composed system and assume that $\mathcal{G} = (I, \mathcal{I})$ is a DAG. To each component Σ^i we associate a contract $\mathcal{C}^i = (A_{W_1^i}, A_{W_2^i}, G_{X^i}, G_{Y^i})$, and let $\mathcal{C} = (\prod_{i \in I} A_{W_1^i}, \prod_{i \in I} G_{X^i}, \prod_{i \in I} G_{Y^i})$ be a contract for Γ . If for all $i \in I$, $\Sigma^i \models \mathcal{C}^i$ and $\prod_{j \in \mathcal{N}(i)} G_{Y^j} \subseteq A_{W_2^i}$ then $\Gamma \models \mathcal{C}$.

PROOF. Let $(w_1, x, y) : E \rightarrow W_1 \times X \times Y$ in $\mathcal{T}$ be a trajectory of the system Γ . Then, for all $i \in I$, there exists a trajectory $(w_1^i, w_2^i, x^i, y^i) : E \rightarrow W_1^i \times W_2^i \times X^i \times Y^i \in \mathcal{T}^i$ of Σ^i such that $w_2^i = \prod_{j \in \mathcal{N}(i)} \{y^j\}$. Let $t \in E$ such that $w_1|_{[0,t]} \in A_{W_1}$. Then we have for all $i \in I$, $w_1^i|_{[0,t]} \in A_{W_1^i}$. Since initial components $\{\Sigma^i\}_{i \in I_{\text{init}}}$ do

not have internal inputs, and from the satisfaction of contracts for all components, we have:

$$\forall i \in I_{\text{init}}, \quad x_{|[0,t]}^i \in G_{X^i} \text{ and } y_{|[0,t]}^i \in G_{Y^i}. \quad (1)$$

Let us assume the existence of $i \in I \setminus I_{\text{init}}$, such that $x_{|[0,t]}^i \notin G_{X^i}$ or $y_{|[0,t]}^i \notin G_{Y^i}$. Since $\Sigma^i \models \mathcal{C}^i$ and $w_1^i|_{[0,t]} \in A_{W_1^i}$, we have that $w_2^i|_{[0,t]} \notin A_{W_2^i}$, then using the fact that $w_2^i = \prod_{j \in \mathcal{N}(i)} \{y^j\}$ and $\prod_{j \in \mathcal{N}(i)} G_{Y^j} \subseteq A_{W_2^i}$, we have the existence of $j \in \mathcal{N}(i)$ such that $y_{|[0,t]}^j \notin G_{Y^j}$. Hence, using the structure of a DAG, we have by iterating this procedure, the existence of $k \in I_{\text{init}}$ such that $y_{|[0,t]}^k \notin G_{Y^k}$ which contradicts (1). Hence, we have for all $i \in I$, $x_{|[0,t]}^i \in G_{X^i}$ and $y_{|[0,t]}^i \in G_{Y^i}$, which implies that $x|_{[0,t]} \in G_X$ and $y|_{[0,t]} \in G_Y$. Then, $\Gamma \models \mathcal{C}$. $\square$

Remark 4 To analyse the tightness of the composition of contracts operator, we start from a collection of local contracts $\mathcal{C}^i = (A_{W_1^i}, A_{W_2^i}, G_{X^i}, G_{Y^i})$, $i \in I$, and we proceed as follows:

- To each contract $\mathcal{C}^i$ we can associate a component $\Sigma^i = (W_1^i, W_2^i, X^i, Y^i, \mathcal{T}^i)$, where the set $\mathcal{T}^i$ is the set of all possible trajectories that weakly satisfy the contract $\mathcal{C}^i$, i.e.,

$$\begin{aligned} \mathcal{T}^i &= \{(w_1^i, w_2^i, x^i, y^i) : E \rightarrow W_1^i \times W_2^i \times X^i \times Y^i \mid \\ &\quad \forall t \in E, w_1^i|_{[0,t]} \in A_{W_1^i} \text{ and } w_2^i|_{[0,t]} \in A_{W_2^i} \\ &\quad \implies x^i|_{[0,t]} \in G_{X^i} \text{ and } y^i|_{[0,t]} \in G_{Y^i}\} \end{aligned}$$

Consider the directed acyclic interconnection graph $\mathcal{G} = (I, \mathcal{I})$ and let $\Gamma = \langle (\Sigma^i)_{i \in I}, \mathcal{I} \rangle = (W_1, X, Y, \mathcal{T}_\Gamma)$ be the composed system;

- We construct the global contract $\mathcal{C} = (\prod_{i \in I} A_{W_1^i}, \prod_{i \in I} G_{X^i}, \prod_{i \in I} G_{Y^i})$ resulting from the interconnection of the local contracts $\mathcal{C}^i = (A_{W_1^i}, A_{W_2^i}, G_{X^i}, G_{Y^i})$, $i \in I$. To the contract $\mathcal{C}$, we associate the system $\Sigma = (W_1, X, Y, \mathcal{T}_\Sigma)$ where the set $\mathcal{T}_\Sigma$ is the set of all possible trajectories that weakly satisfy the contract $\mathcal{C}$, i.e.,

$$\begin{aligned} \mathcal{T}_\Sigma &= \{(w_1, x, y) : E \rightarrow W_1 \times X \times Y \mid \forall t \in E, \\ &\quad w_1|_{[0,t]} \in A_{W_1} \implies x|_{[0,t]} \in G_X \text{ and } y|_{[0,t]} \in G_Y\} \end{aligned}$$

First, one can see that $\mathcal{T}_\Gamma \subseteq \mathcal{T}_\Sigma$ and, in general, one may have that $\mathcal{T}_\Gamma \neq \mathcal{T}_\Sigma$. Moreover, one can also check that the proposed contracts composition operator is tight, in the sense that the system Σ is the smallest system defining an assume-guarantee contract and containing the trajectories of the system Γ .

5.2 Cyclic interconnections

In order to deal with cyclic interconnections, we need the following assumption on the set of guarantees on the

output G_Y . This assumption will be explained later on different examples.

Assumption 1 Let $\Sigma = (W_1, W_2, X, Y, \mathcal{T})$ be a continuous-time system, and $\mathcal{C} = (A_{W_1}, A_{W_2}, G_X, G_Y)$ an assume-guarantee contracts for Σ . For any trajectory $(w_1, w_2, x, y) : E \rightarrow W_1 \times W_2 \times X \times Y$ of the system Σ , the following logical implication is satisfied for all $t \in E$:

$$\forall s \in [0, t], (y_{|[0,s]} \in G_Y \Rightarrow y_{|[0,t]} \in G_Y).$$

Now we give some sufficient conditions on systems and contracts in order to satisfy Assumption 1 for different examples. Let $\Sigma = (W_1, W_2, X, Y, \mathcal{T})$ be a system, and $\mathcal{C} = (A_{W_1}, A_{W_2}, G_X, G_Y)$ an assume-guarantee contract for Σ .

- If the set G_Y is of the same form as the set A defined in Example 1 and if for any trajectory $(w_1, w_2, x, y) : E \rightarrow W_1 \times W_2 \times X \times Y$ for the system Σ , $y : E \rightarrow Y$ is left continuous and the set of guarantees G_Y is closed then Assumption 1 is satisfied.
- If the set G_Y is of the same form as the set A defined in Example 2 and similarly to the previous example, it can be shown that if for any trajectory $(w_1, w_2, x, y) : E \rightarrow W_1 \times W_2 \times X \times Y$ for the system Σ , $y : E \rightarrow Y$ is q times differentiable, the q th derivative $y^{(q)}$ is left continuous and the sets S_i , $i = 1, \dots, q$, are closed then Assumption 1 is satisfied.
- If the set G_Y is of the same form as the set A defined in Example 3 and if the set valued map $F : \mathbb{R}^n \rightarrow 2^{\mathbb{R}^n}$ is outer semicontinuous (see Definition 5.4 in [31]) and for any trajectory $(w_1, w_2, x, y) : E \rightarrow W_1 \times W_2 \times X \times Y$ of the system Σ , $y : I \rightarrow Y$ is differentiable and its derivative is left continuous, then Assumption 1 is satisfied.

The following result relates the satisfaction of Assumption 1 for a global system to its satisfaction for the components.

Claim 1 Given a collection of components $\{\Sigma^i\}_{i \in I}$, such that each component Σ^i satisfies Assumption 1 with respect to the contract $\mathcal{C}^i = (A_{W_1^i}, A_{W_2^i}, G_{X^i}, G_{Y^i})$. Then the composed system $\Gamma = \langle (\Sigma^i)_{i \in I}, \mathcal{I} \rangle$ satisfies Assumption 1 with respect to the contract $\mathcal{C} = (\prod_{i \in I} A_{W_1^i}, \prod_{i \in I} G_{X^i}, \prod_{i \in I} G_{Y^i})$.

PROOF. Let $(w_1, x, y) : E \rightarrow W_1 \times X \times Y$ in $\mathcal{T}$ be a trajectory of the interconnected system Γ . Let $t \in E$ and assume that for all $s \in [0, t]$, $y_{|[0,s]} \in G_Y$. Then we have for all $s \in [0, t]$ and for all $i \in I$, $y_{|[0,s]}^i \in G_{Y^i}$. Since each component Σ^i satisfies Assumption 1 with respect to the contract $\mathcal{C}^i = (A_{W_1^i}, A_{W_2^i}, G_{X^i}, G_{Y^i})$, we have for all $i \in I$, $y_{|[0,t]}^i \in G_{Y^i}$, which in turn implies that $y_{|[0,t]} \in G_Y$. $\square$

Theorem 2 Consider a network of continuous-time components $\{\Sigma^i\}_{i \in I}$ compatible for composition with respect to $\mathcal{I}$. Let the system $\Gamma = \langle (\Sigma^i)_{i \in I}, \mathcal{I} \rangle$ be the composed system. To each component Σ^i , we associate a contract $\mathcal{C}^i = (A_{W_1^i}, A_{W_2^i}, G_{X^i}, G_{Y^i})$ and let $\mathcal{C} = (\prod_{i \in I} A_{W_1^i}, \prod_{i \in I} G_{X^i}, \prod_{i \in I} G_{Y^i})$ be a contract for Γ . Let us assume the following:

- (i) for all $i \in I$, $\Sigma^i \models \mathcal{C}^i$;
- (ii) for all $i \in I$, $\prod_{j \in \mathcal{N}(i)} G_{Y^j} \subseteq A_{W_2^i}$;
- (iii) for all $i \in I$, Σ^i satisfies Assumption 1;
- (iv) for any cycle ζ_q in the interconnection graph $\mathcal{G} = (I, \mathcal{I})$, there exists an element $k \in \zeta_q$ such that $\Sigma^k \models_s \mathcal{C}^k$;
- (v) for all $i \in I$, $A_{W_1^i}$ is a prefix-closed set.

then $\Gamma \models \mathcal{C}$.

PROOF. Let $(w_1, x, y) : E \rightarrow W_1 \times X \times Y$ in $\mathcal{T}$ be a trajectory of the system Γ . Then, for all $i \in I$, there exists a trajectory $(w_1^i, w_2^i, x^i, y^i) : E \rightarrow W_1^i \times W_2^i \times X^i \times Y^i \in \mathcal{T}^i$ of Σ^i such that $w_2^i = \prod_{j \in \mathcal{N}(i)} \{y^j\}$. Let $t \in E$ such that $w_{1|[0,t]} \in A_{W_1}$. Then we have for all $i \in I$, $w_{1|[0,t]}^i \in A_{W_1^i}$. All initial components $\{\Sigma^i\}_{i \in I_{\text{init}}}$ do not have internal inputs, then from the satisfaction of contracts for all components and since $A_{W_1^i}$ is prefix-closed for all $i \in I$, we have:

$$\forall i \in I_{\text{init}}, \forall s \in [0, t], x_{|[0,s]}^i \in G_{X^i} \text{ and } y_{|[0,s]}^i \in G_{Y^i}. \quad (2)$$

First, let us prove that

$$\forall i \in I, y_{|[0,0]}^i \in G_{Y^i}. \quad (3)$$

We have the existence of an element k in any cycle ζ_q such that $\Sigma^k \models_s \mathcal{C}^k$, which implies from prefix closedness of $A_{W_1^k}$ that $y_{|[0,0]}^k \in G_{Y^k}$. To prove that this initial condition is satisfied by all the components Σ^i , $i \in I$, we proceed as follows: for any component Σ^i that strongly satisfies its contract, we drop the incoming edge into the vertex k in the cycle ζ_q . Then, in view of Remark 1, a new DAG, $\mathcal{G}^{\text{DAG}}$ is obtained. Then from (2) we have:

$$\forall i \in I_{\text{init}}^{\text{DAG}} \subseteq I_{\text{init}} \cup A, y_{|[0,0]}^i \in G_{Y^i} \quad (4)$$

where A is the set of vertices to which we dropped an edge (vertices corresponding to components that strongly satisfy their contracts). Now let an element $i \in I \setminus I_{\text{init}}^{\text{DAG}}$ and let us assume that $y_{|[0,0]}^i \notin G_{Y^i}$. From prefix-closedness of $A_{W_1^i}$ it follows that $w_{1|[0,0]}^i \in A_{W_1^i}$, moreover $\Sigma^i \models \mathcal{C}^i$, then we have that $w_{2|[0,0]}^i \notin A_{W_2^i}$, and using the fact that $w_2^i = \prod_{j \in \mathcal{N}(i)} \{y^j\}$ and $\prod_{j \in \mathcal{N}(i)} G_{Y^j} \subseteq A_{W_2^i}$, we have the existence of $j \in \mathcal{N}(i)$ such that $y_{|[0,0]}^j \notin G_{Y^j}$.

Hence, using the structure of a DAG, we have by iterating this procedure, the existence of $h \in I_{\text{init}}^{\text{DAG}}$ such that $y_{[0,0]}^h \notin G_Y^h$ which contradicts (4). Hence, we have for all $i \in I$, $y_{[0,0]}^i \in G_Y^i$, which implies that $y_{[0,0]} \in G_Y$. To prove that $\Gamma \models \mathcal{C}$, we proceed by contradiction. Let us define

$$\begin{aligned} T &= \sup\{s \in [0, t] \mid \forall s' \in [0, s], y_{[0,s']} \in G_Y\} \\ &= \sup\{s \in [0, t] \mid \forall i \in I, \forall s' \in [0, s], y_{[0,s']}^i \in G_Y^i\}. \end{aligned} \quad (5)$$

From (3) we have $y_{[0,0]} \in G_Y$, it then follows that $T \in [0, t]$. Let us remark that by (5), we have that $y_{[0,s]} \in G_Y$ for all $s \in [0, T)$.

We have $y_{[0,s]} \in G_Y$ for all $s \in [0, T)$. Then, from (iii) and using Claim 1, we have that $y_{[0,T]} \in G_Y$. We have the existence of an element k in any cycle ζ_q such that $\Sigma^k \models_s \mathcal{C}^k$. We have from prefix-closedness of the set $A_{W_1^k}$ that $w_{1[0,T]}^k \in A_{W_1^k}$. Then, since

$$w_{2[0,T]}^k = \prod_{j \in \mathcal{N}(k)} \{y_{[0,T]}^j\} \in \prod_{j \in \mathcal{N}(k)} G_{Y^j} \subseteq A_{W_2^k}$$

we have from (iv) the existence of $\delta_k > 0$ such that

$$\forall s_k \in [0, \delta_k], y_{[0, T+s_k] \cap E}^k \in G_{Y^k}$$

Let $\delta = \min_{k \in A} \delta_k$, where A is the set of vertices corresponding to components that strongly satisfy their contracts, we have that

$$\forall s \in [0, \delta], y_{[0, T+s] \cap E}^k \in G_{Y^k}$$

by using the same procedure as above (dropping the incoming edges into the vertex k in the cycle ζ_q), we have from (2):

$$\forall i \in I_{\text{init}}^{\text{DAG}} \subseteq I_{\text{init}} \cup A, \forall s \in [0, \delta], y_{[0, T+s] \cap [0, t]}^i \in G_{Y^i}. \quad (6)$$

Now let an element $i \in I \setminus I_{\text{init}}^{\text{DAG}}$ and let us assume the existence of $s' \in [0, \delta]$ such that $y_{[0, T+s'] \cap [0, t]}^i \notin G_{Y^i}$. From prefix-closedness of $A_{W_1^i}$ we have that $w_{1[0, T+s'] \cap [0, t]}^i \in A_{W_1^i}$. Then, since $\Sigma^i \models \mathcal{C}^i$ we have that $w_{2[0, T+s'] \cap [0, t]}^i \notin A_{W_2^i}$, then using the fact that $w_2^i = \prod_{j \in \mathcal{N}(i)} \{y^j\}$ and $\prod_{j \in \mathcal{N}(i)} G_{Y^j} \subseteq A_{W_2^i}$, we have the existence of $j \in \mathcal{N}(i)$ such that $y_{[0, T+s'] \cap [0, t]}^j \notin G_{Y^j}$. Hence, using the structure of a DAG, we have by iterating this procedure, the existence of $h \in I_{\text{init}}^{\text{DAG}}$ such that $y_{[0, T+s'] \cap [0, t]}^h \notin G_Y^h$ which contradicts (6). Hence, we have

$$\forall i \in I, \forall s \in [0, \delta], y_{[0, T+s] \cap [0, t]}^i \in G_{Y^i}$$

which by definition of T implies that $T = t$. Hence, we have for all $i \in I$ and for all $s \in [0, t]$, $y_{[0,s]}^i \in G_{Y^i}$, which implies that $y_{[0,s]} \in G_Y$ for all $s \in [0, t]$.

Let $i \in I$, we have $w_{1[0,t]}^i \in A_{W_1^i}$ and

$$w_{2[0,t]}^i = \prod_{j \in \mathcal{N}(i)} \{y_{[0,t]}^j\} \in \prod_{j \in \mathcal{N}(i)} G_{Y^j} \subseteq A_{W_2^i}$$

then we have from (i) that for all $i \in I$, $x_{[0,t]}^i \in G_{X^i}$, which implies that $x_{[0,t]} \in G_X$. Hence, $\Gamma \models \mathcal{C}$. $\square$

The proof of Theorem 2 can be interpreted as follows: from the satisfaction of contracts, the time interval on which the property holds for the output may be right-open $[0, T)$. Then from Assumption 1, we can ensure that the property holds on its closure $[0, T]$. Hence, one can use the strong satisfaction again to move further and to enlarge the time interval on which the property holds $[0, T + \delta]$. Thus, it has to be the whole time domain on which the trajectory is defined.

Remark 5 It was shown in Theorem 2 that prefix-closedness of the set of assumptions A_{W_1} is critical for the compositionality result for general interconnections containing cycles. Given a non prefix-closed set of assumptions A_{W_1} , the set $\text{pref}(A_{W_1})$ is prefix-closed. Hence, the result of Theorem 2 remain correct if we assign to each component Σ^i the contract $\mathcal{C}^i = (\text{pref}(A_{W_1^i}), A_{W_2^i}, G_{X^i}, G_{Y^i})$. This approach allows to overcome the prefix-closedness of the set of assumptions on the external inputs A_{W_1} , at the cost of an additional conservatism.

Remark 6 Proposition 1 in [29] can be recovered by the result of Theorem 2, where our prefix-closed sets and general sets corresponds to invariants and LTL specifications, respectively, in that work.

Let us point out that weak satisfaction is generally insufficient to reason about general compositions containing cycles, as shown by the following counter-example:

Example 6 Let us consider the system $\Sigma^1 = (W_1, W_2, X, Y, \mathcal{T})$ where, $W_1 = W_2 = X = Y = \mathbb{R}_0^+$. A trajectory of Σ^1 is a quadruple $(w_1, w_2, x, y) : E \rightarrow W_1 \times W_2 \times X \times Y$ in $\mathcal{T}$ where $E \in \mathbb{E}(\mathbb{R}_0^+)$, w_1 and w_2 are continuous, x and y are differentiable and such that $x(0) = 0$, and for all $t \in \mathbb{R}_0^+$,

$$\begin{cases} \dot{x}(t) = \sqrt{w_2(t)} + w_1(t) \\ y(t) = x(t). \end{cases}$$

Let $I = \{1\}$ and consider the interconnection relation $\mathcal{I} = \{(1, 1)\}$. It is clear that $\{\Sigma^i\}_{i \in I}$ is compatible for

composition with respect to $\mathcal{I}$. Let us consider the assume-guarantee contract $\mathcal{C} = (A_{W_1}, A_{W_2}, G_X, G_Y)$ for Σ^1 , given by:

$$\begin{aligned} A_{W_1} &= \{w_1 : E \rightarrow W_1 \in M(W_1) \mid \forall t \in E, w_1(t) = 0\} \\ A_{W_2} &= \{w_2 : E \rightarrow W_2 \in M(W_2) \mid \forall t \in E, w_2(t) = 0\} \\ G_X &= \{x : E \rightarrow X \in M(X) \mid \forall t \in E, x(t) = 0\} \\ G_Y &= \{y : E \rightarrow Y \in M(Y) \mid \forall t \in E, y(t) = 0\} \end{aligned}$$

Let $\mathcal{C}_\Gamma$ be the contract for the composed system $\Gamma = \langle (\Sigma^i)_{i \in I}, \mathcal{I} \rangle$ defined as in Theorem 2. We can easily check that $\Sigma^1 \models \mathcal{C}$ and that Assumption 1 holds. However, the conclusion of the previous theorem does not hold. Indeed, the map $(w_1, x, y) : \mathbb{R}_0^+ \rightarrow W_1 \times X \times Y$ defined by $w_1(t) = 0$ and $x(t) = y(t) = t^2/4$ for all $t \in \mathbb{R}_0^+$ is a trajectory of Γ and the contract $\mathcal{C}_\Gamma$ of the system Γ is not satisfied.

It is clear from the previous example that strong satisfaction is needed to reason about general interconnections containing cycles. We show two modifications of the previous example, based on sampling or time-delays, which lead to strong satisfaction of the contract.

Example 7 Consider the system $\Sigma^1 = (W_1, W_2, X, Y, \mathcal{T})$ where $W_1 = W_2 = X = Y = \mathbb{R}_0^+$. A trajectory of Σ^1 is a quadruple $(w_1, w_2, x, y) : E \rightarrow W_1 \times W_2 \times X \times Y$ in $\mathcal{T}$ where $E \in \mathbb{E}(\mathbb{R}_0^+)$, w_1 and w_2 are continuous, x and y are differentiable and such that $x(0) = 0$, and for all $t \in \mathbb{R}_0^+$,

$$\begin{cases} \dot{x}(t) = \sqrt{w_2(t)} + w_1(t) \\ y(t) = 0 \\ y(t) = x(t_k) \end{cases} \quad \begin{matrix} 0 \leq t \leq t_0 \\ t_k < t \leq t_{k+1}, k \in \mathbb{N}. \end{matrix}$$

where $(t_k)_{k \in \mathbb{N}}$ a strictly increasing sequence of sampling instants with $t_0 \geq 0$ and $t_k \rightarrow +\infty$ when $k \rightarrow +\infty$. We consider the same assume-guarantee contract as in the previous example. It can be seen that y is left-continuous and Assumption 1 is satisfied. We can easily check that $\Sigma^1 \models_s \mathcal{C}$, where the value of δ as in Definition 5 is given by $\delta = t_{k+1} - t$ if $t_k \leq t < t_{k+1}$. Let $I = \{1\}$ and consider the interconnection relation $\mathcal{I} = \{(1, 1)\}$. Let $\mathcal{C}_\Gamma$ be the contract for the composed system $\Gamma = \langle (\Sigma^i)_{i \in I}, \mathcal{I} \rangle$ defined as in Theorem 2. Now we can check that the conclusion of the previous theorem holds since the only trajectory $(w_1, x, y) : \mathbb{R}_0^+ \rightarrow W_1 \times X \times Y$ of the composed system Γ is given by $w_1(t) = x(t) = y(t) = 0$, for all $t \in \mathbb{R}_0^+$.

Example 8 Consider the system $\Sigma^1 = (W_1, W_2, X, Y, \mathcal{T})$ where $W_1 = W_2 = X = Y = \mathbb{R}_0^+$. A trajectory of Σ^1 is a quadruple $(w_1, w_2, x, y) : E \rightarrow W_1 \times W_2 \times X \times Y$ in $\mathcal{T}$ where $E \in \mathbb{E}(\mathbb{R}_0^+)$, w_1 and w_2 are continuous, x and y are differentiable and such that $x(0) = 0$, and for all

$t \in \mathbb{R}_0^+$,

$$\begin{cases} \dot{x}(t) = \sqrt{w_2(t)} + w_1(t) \\ y(t) = 0 \\ y(t) = x(t - T) \end{cases} \quad \begin{matrix} 0 \leq t \leq T \\ T < t. \end{matrix}$$

where $T > 0$ is a time delay. We consider the same assume-guarantee contract as in Example 6. It can be seen that y is left-continuous and Assumption 1 is satisfied. We can easily check that $\Sigma \models_s \mathcal{C}$, where the value of δ as in Definition 5 is given by $\delta = T$. Let $I = \{1\}$ and consider the interconnection relation $\mathcal{I} = \{(1, 1)\}$. Let the contract $\mathcal{C}_\Gamma$ for the composed system $\Gamma = \langle (\Sigma^i)_{i \in I}, \mathcal{I} \rangle$ defined as in Theorem 2. Then, we can check that the conclusion of the previous theorem holds since the only trajectory $(w_1, x, y) : \mathbb{R}_0^+ \rightarrow W_1 \times X \times Y$ of the composed system Γ is given by $w_1(t) = x(t) = y(t) = 0$, for all $t \in \mathbb{R}_0^+$.

It can be seen from the Examples 7 and 8 that our framework is suitable to reason about systems that include some sampled or delayed behaviors. Moreover, these examples suggest that by sampling or delaying the output of a component, strong satisfaction of a contract can be obtained. These examples also show how one can go from weak to strong satisfaction by slightly modifying the system. In the next section, we show that this is also possible by slightly modifying the contract.

It is noteworthy to mention that Theorem 2 does not hold in general without Assumption 1, as shown by the following counter-example.

Example 9 Consider the system $\Sigma^1 = (W_1, W_2, X, Y, \mathcal{T})$ where $W_1 = W_2 = X = Y = \mathbb{R}$. A trajectory of Σ^1 is a quadruple $(w_1, w_2, x, y) : E \rightarrow W_1 \times W_2 \times X \times Y$ in $\mathcal{T}$ where $E \in \mathbb{E}(\mathbb{R}_0^+)$, w_1, w_2, x, y are continuous, $w_2(0) = 0$ and for all $t \in \mathbb{R}_0^+$,

$$\begin{cases} x(t) = w_2(t) + w_1(t) \\ y(t) = x(t). \end{cases}$$

Let $I = \{1\}$ and consider the interconnection relation $\mathcal{I} = \{(1, 1)\}$. It is clear that $\{\Sigma^i\}_{i \in I}$ is compatible for composition with respect to $\mathcal{I}$. Let us consider the assume-guarantee contract $\mathcal{C} = (A_{W_1}, A_{W_2}, G_X, G_Y)$ for Σ^1 , given by:

$$\begin{aligned} A_{W_1} &= \{w_1 : E \rightarrow W_1 \in M(W_1) \mid \forall t \in E, w_1(t) = 0\} \\ A_{W_2} &= \{w_2 : E \rightarrow W_2 \in M(W_2) \mid \forall t \in E, w_2(t) \in (-1, 1)\} \\ G_X &= \{x : E \rightarrow X \in M(X) \mid \forall t \in E, x(t) \in (-1, 1)\} \\ G_Y &= \{y : E \rightarrow Y \in M(Y) \mid \forall t \in E, y(t) \in (-1, 1)\} \end{aligned}$$

Let $\mathcal{C}_\Gamma$ be the contract for the composed system $\Gamma = \langle (\Sigma^i)_{i \in I}, \mathcal{I} \rangle$ defined as in Theorem 2. We can easily check

from the continuity of y that $\Sigma^1 \models_s \mathcal{C}$ and that all the assumptions of Theorem 2 hold except for Assumption 1. Moreover, we can check that the conclusion of Theorem 2 does not hold. Indeed, for $w_1 : \mathbb{R}_0^+ \rightarrow W_1 \in A_{W_1}$ any map $(w_1, x, y) : \mathbb{R}_0^+ \rightarrow W_1 \times X \times Y$ satisfying $x(0) = y(0) = 0$ and $x(t) = y(t)$ for all $t \in \mathbb{R}_0^+$ is a trajectory of Γ and the contract $\mathcal{C}_\Gamma$ of the system Γ is not satisfied.

Remark 7 Theorems 1 and 2 apply to a very general class of systems. When considering more specific classes, one can sometimes reason on general interconnections without strong contract satisfaction. Such a case will be shown in Section 6, where we consider systems modeled by Lipschitz differential inclusions and invariance assume-guarantee contracts.

5.2.1 From weak to strong contract satisfaction

In this section, we show that under some additional assumptions, it is possible to reason about general compositions using the weak semantics of assume guarantee contracts.

In order to measure the distance between two continuous-time trajectories, which might not have the same time domain. We use the notion of ε -closeness of trajectories [17], which is related to the Hausdorff distance between the graphs of the trajectories.

Definition 6 (ε -closeness of trajectories) Let $Z \subseteq \mathbb{R}^n$. Given $\varepsilon > 0$ and two continuous-time trajectories $z_1 : E_1 \rightarrow Z$ and $z_2 : E_2 \rightarrow Z$ in $M(Z)$. z_2 is said to be ε -close to z_1 , if for all $t_1 \in E_1$, there exists $t_2 \in E_2$ such that $|t_1 - t_2| \leq \varepsilon$ and $\|z_1(t_1) - z_2(t_2)\| \leq \varepsilon$. We define the ε -expansion of z_1 by: $\mathcal{B}_\varepsilon(z_1) = \{z' : E' \rightarrow Z \mid z' \text{ is } \varepsilon\text{-close to } z_1\}$.

This notion is generalized toward sets of continuous-time maps in the usual way: For $A \subseteq M(Z)$, $\mathcal{B}_\varepsilon(A) = \bigcup_{z \in A} \mathcal{B}_\varepsilon(z)$.

Proposition 2 Let $\Sigma = (W_1, W_2, X, Y, \mathcal{T})$ be a continuous-time system and let $\mathcal{C} = (A_{W_1}, A_{W_2}, G_X, G_Y)$ be an assume-guarantee contract for Σ . Let us assume that for all trajectories $(w_1, w_2, x, y) : E \rightarrow W_1 \times W_2 \times X \times Y \in \mathcal{T}$, $y : E \rightarrow Y$ is continuous and $y|_{[0,0]} \in G_Y$ if $w_1|_{[0,0]} \in A_{W_1}$. If $\Sigma \models \mathcal{C}$, then for all $\varepsilon > 0$, $\Sigma \models_s \mathcal{C}_\varepsilon$ where $\mathcal{C}_\varepsilon = (A_{W_1}, A_{W_2}, G_X, \mathcal{B}_\varepsilon(G_Y) \cap M(Y))$.

PROOF. Let $(w_1, w_2, x, y) : E \rightarrow W_1 \times W_2 \times X \times Y \in \mathcal{T}$, then $y|_{[0,0]} \in G_Y \subseteq \mathcal{B}_\varepsilon(G_Y) \cap M(Y)$. Let $t \in E$, such that $w_1|_{[0,t]} \in A_{W_1}$ and $w_2|_{[0,t]} \in A_{W_2}$. Then, satisfaction of $\mathcal{C}$ gives that $x|_{[0,t]} \in G_X$ and $y|_{[0,t]} \in G_Y$. By continuity of y , there exists $\delta > 0$ such that for all $s \in [0, \delta]$, $y|_{[0,t+s]} \in \mathcal{B}_\varepsilon(G_Y)$. Also by definition,

Fig. 4. An illustration of the result of Proposition 2. Top: trajectories of the external w_1 and internal w_2 inputs. Bottom: trajectories of the state x and output y .

$y|_{[0,t+s]} \cap E \in M(Y)$ for all $s \in [0, \delta]$. Hence, $y|_{[0,t+s]} \cap E \in \mathcal{B}_\varepsilon(G_Y) \cap M(Y)$, for all $s \in [0, \delta]$. $\square$

A pictorial representation for the result of proposition 2 is shown in Figure 4. The following example shows an application of the previous corollary:

Example 10 Consider the system $\Sigma^1 = (W_1, W_2, X, Y, \mathcal{T})$ where $W_1 = W_2 = X = Y = \mathbb{R}_0^+$. A trajectory of Σ^1 is a triple $(w_1, w_2, x, y) : E \rightarrow W_1 \times W_2 \times X \times Y$ in $\mathcal{T}$ where $E = \mathbb{R}_0^+$, w_1 and w_2 are continuous, x and y are differentiable and such that $x(0) = 0$, and for all $t \in \mathbb{R}_0^+$,

$$\begin{cases} \dot{x}(t) = \sqrt{w_2(t)} - x(t) + w_1(t) \\ y(t) = x(t). \end{cases}$$

Let $I = \{1\}$ and consider the interconnection relation $\mathcal{I} = \{(1, 1)\}$. It is clear that $\{\Sigma^i\}_{i \in I}$ is compatible for composition with respect to $\mathcal{I}$. Let $a > 1$ and let us consider the assume-guarantee contract $\mathcal{C} = (A_{W_1}, A_{W_2}, G_X, G_Y)$ for Σ^1 , given by:

$$\begin{aligned} A_{W_1} &= \{w_1 : E \rightarrow W_1 \in M(W_1) \mid \forall t \in E, w_1(t) = 0\} \\ A_{W_2} &= \{w_2 : E \rightarrow W_2 \in M(W_2) \mid \forall t \in E, w_2(t) \in [0, a^2]\} \\ G_X &= \{x : E \rightarrow X \in M(X) \mid \forall t \in E, x(t) \in [0, a]\} \\ G_Y &= \{y : E \rightarrow Y \in M(Y) \mid \forall t \in E, y(t) \in [0, a]\} \end{aligned}$$

We can easily check that $\Sigma^1 \models \mathcal{C}$ and for all trajectories $(w_1, w_2, x, y) \in \mathcal{T}$, $y : E \rightarrow Y$ is continuous and $y|_{[0,0]} \in G_Y$. Then, from Proposition 2, we have that $\Sigma \models_s \mathcal{C}_\varepsilon$ for any $\varepsilon > 0$, where $\mathcal{C}_\varepsilon = (A_{W_1}, A_{W_2}, G_X, \mathcal{B}_\varepsilon(G_Y) \cap M(Y))$. Now let $\varepsilon > 0$, such that

$$\begin{aligned} \mathcal{B}_\varepsilon(G_Y) \cap M(Y) &= \{y : E \rightarrow Y \in M(Y) \mid \forall t \in E, y(t) \in [0, a + \varepsilon]\} \\ &\subseteq \{y : E \rightarrow Y \in M(Y) \mid \forall t \in E, y(t) \in [0, a^2]\} \\ &= A_{W_2} \end{aligned}$$

Then, since the system Σ^1 satisfies Assumption 1 (the output trajectory $y : E \rightarrow Y$ is continuous and the set $[0, a]$ is closed), we have from Theorem 2 that the composed system $\Gamma = \langle (\Sigma^i)_{i \in I}, \mathcal{I} \rangle$ satisfies the composed contract $\mathcal{C}_\Gamma = (A_{W_1}, G_X, \mathcal{B}_\varepsilon(G_Y) \cap M(Y))$. Let us remark that there exist trajectories of the composed system Γ given by: $(w_1, x, y) : \mathbb{R}_0^+ \rightarrow W_1 \times X \times Y$, where $w_1(t) = 0$ and $x(t) = y(t) = (1 - e^{-t/2})^2$, for all $t \in \mathbb{R}_0^+$.

We have shown how one can go from weak to strong satisfaction of a contract, by relaxing the guarantees on the output. In the next result, we show that it is also possible to do so by relaxing the assumptions.

Proposition 3 Let $\Sigma = (W_1, W_2, X, Y, \mathcal{T})$ be a continuous-time system and let $\mathcal{C} = (A_{W_1}, A_{W_2}, G_X, G_Y)$ be an assume-guarantee contract for Σ . Let us assume that for all trajectories $(w_1, w_2, x, y) : E \rightarrow W_1 \times W_2 \times X \times Y \in \mathcal{T}$, $w_1 : E \rightarrow W_1$ and $w_2 : E \rightarrow W_2$ are continuous and $y|_{[0,0]} \in G_Y$ if $w_1|_{[0,0]} \in A_{W_1}$. For an $\varepsilon > 0$, if $\Sigma \models \mathcal{C}^\varepsilon$, with $\mathcal{C}^\varepsilon = (\mathcal{B}_\varepsilon(A_{W_1}) \cap M(W_1), \mathcal{B}_\varepsilon(A_{W_2}) \cap M(W_2), G_X, G_Y)$. Then, $\Sigma \models_s \mathcal{C}$.

PROOF. Let $(w_1, w_2, x, y) : E \rightarrow W_1 \times W_2 \times X \times Y \in \mathcal{T}$, then $y|_{[0,0]} \in G_Y$. Let $t \in E$, such that $w_1|_{[0,t]} \in A_{W_1}$ and $w_2|_{[0,t]} \in A_{W_2}$. By continuity of w_1 and w_2 , there exists $\delta > 0$ such that for all $s \in [0, \delta]$, $w_1|_{[0,t+s] \cap E} \in \mathcal{B}_\varepsilon(A_{W_1})$ and $w_2|_{[0,t+s] \cap E} \in \mathcal{B}_\varepsilon(A_{W_2})$. Also by definition, $w_1|_{[0,t+s] \cap E} \in M(W_1)$ and $w_2|_{[0,t+s] \cap E} \in M(W_2)$ for all $s \in [0, \delta]$. Then, satisfaction of $\mathcal{C}^\varepsilon$ gives that $x|_{[0,t+s]} \in G_X$ and $y|_{[0,t+s]} \in G_Y$ for all $s \in [0, \delta]$. $\square$

We recall that the approach proposed in Proposition 3 to ensure strong satisfaction of contract has been used in [32] to construct symbolic controllers for sampled-data systems. Interestingly, this technique is useful in practice, since it allows to ensure strong satisfaction of contracts without reasoning in terms of δ which may depend on time and trajectory.

6 Compositional invariants for differential inclusions

In this section, we focus on continuous-time systems $\Sigma = (W_1, W_2, X, Y, \mathcal{T})$ defined by differential inclusions, and invariance assume-guarantee contracts, where assumptions and guarantees are defined as in Example 1. We use the classical characterization of invariant sets for differential inclusions developed using the concept of contingent cone (see [4] and the references therein) to derive necessary and sufficient conditions for weak satisfaction of assume-guarantee contracts. We also show that under some technical assumptions (Lipschitzness of the vector field and the output map), weak satisfaction

makes it possible to reason on general interconnections containing cycles.

A trajectory of Σ is a triple $(w_1, w_2, x, y) : E \rightarrow W_1 \times W_2 \times X \times Y$ in $\mathcal{T}$ where $E \in \mathbb{E}(\mathbb{R}_0^+)$, w_1 and w_2 are locally measurable, x and y are absolutely continuous and continuous, respectively, and satisfy for almost all $t \in E$:

$$\begin{cases} \dot{x}(t) \in F(x(t), w_1(t), w_2(t)), & x(0) \in X_0 \\ y(t) = h(x(t)) \end{cases} \quad (7)$$

where $F : \mathbb{R}^n \times \mathbb{R}^{m_1} \times \mathbb{R}^{m_2} \rightarrow 2^{\mathbb{R}^n}$ is a set-valued map, $h : \mathbb{R}^n \rightarrow \mathbb{R}^p$ is continuous and X_0 is the set of initial conditions. Let us introduce the following assumption on the system Σ :

Assumption 2 The set-valued map¹ $F : \mathbb{R}^n \times \mathbb{R}^{m_1} \times \mathbb{R}^{m_2} \rightarrow 2^{\mathbb{R}^n}$ is Lipschitz, has compact values and $X \times W_1 \times W_2 \subseteq \text{Int}(\text{dom}(F))$. The map² $h : \mathbb{R}^n \rightarrow \mathbb{R}^p$ satisfies $X \subseteq \text{Int}(\text{dom}(h))$ and $h(X) \subseteq Y$.

Definition 7 Consider the sets $S_{W_1} \subseteq \mathbb{R}^{m_1}$, $S_{W_2} \subseteq \mathbb{R}^{m_2}$, $S_X \subseteq \mathbb{R}^n$ and $S_Y \subseteq \mathbb{R}^p$. A contract $\mathcal{C} = (A_{W_1}, A_{W_2}, G_X, G_Y)$ is an invariance contract with respect to $(S_{W_1}, S_{W_2}, S_X, S_Y)$ if the set of assumptions and guarantees are described as follows:

- $A_{W_i} = \{w_i : E \rightarrow \mathbb{R}^{m_i} \in M(\mathbb{R}^{m_i}) \mid \forall t \in E, w_i(t) \in S_{W_i}\}$, $i \in \{1, 2\}$;
- $G_X = \{x : E \rightarrow \mathbb{R}^n \in M(\mathbb{R}^n) \mid \forall t \in E, x(t) \in S_X\}$;
- $G_Y = \{y : E \rightarrow \mathbb{R}^p \in M(\mathbb{R}^p) \mid \forall t \in E, y(t) \in S_Y\}$;

Consider a network of components $\{\Sigma^i\}_{i \in I}$, compatible for composition with respect to $\mathcal{I}$, where each component has the form of (7). Each component Σ^i has maps and initial sets F^i, h^i, X_0^i , $i \in I$, the composed system $\Gamma = \langle (\Sigma^i)_{i \in I}, \mathcal{I} \rangle$ can be written under the same form with maps F, h and initial set X_0 given by:

$$\begin{aligned} F(x, w_1) &= \prod_{i \in I} F^i(x^i, w_1^i, w_2^i), \quad w_2^i = \prod_{j \in \mathcal{N}(i)} \{h^j(x^j)\} \\ h(x) &= (h^1(x^1), \dots, h^N(x^N)), \\ X_0 &= \prod_{i \in I} X_0^i. \end{aligned} \quad (8)$$

¹ Given a set-valued map $F : \mathbb{R}^q \rightarrow 2^{\mathbb{R}^n}$, the domain of F is $\text{dom}(F) = \{z \in \mathbb{R}^q \mid F(z) \neq \emptyset\}$. F is said to be locally Lipschitz if for all $z \in \text{Int}(\text{dom}(F))$, there exists a neighborhood U of z and a constant $L \geq 0$ (the Lipschitz constant) such that for every $z_1, z_2 \in U \cap \text{dom}(F)$, $F(z_1) \subseteq F(z_2) + L\|z_1 - z_2\|\mathbb{B}$. F is said to be Lipschitz if the constant L is independent of $z \in \text{Int}(\text{dom}(F))$. It has compact values if for all $z \in \text{dom}(F)$, $F(z)$ is compact.

² Given a map $h : \mathbb{R}^n \rightarrow \mathbb{R}^p$, the domain of h is denoted $\text{dom}(h)$ and consists of elements $x \in \mathbb{R}^n$ such that $h(x)$ is defined.

Note that this representation is consistent with the one given in Definition 4. We have the following technical result whose proof is given in the appendix.

Claim 2 *If h^i is Lipschitz and Assumption 2 holds for all Σ^i , $i \in I$, then Assumption 2 holds for $\Gamma = \langle (\Sigma^i)_{i \in I}, \mathcal{I} \rangle$;*

6.1 Invariants relative to assume-guarantee contracts

We give necessary and sufficient conditions for weak satisfaction of assume-guarantee contracts based on the classical characterization of invariant sets for differential inclusions (see e.g. Theorem 5.3.4 in [4]).

Definition 8 *Let $K \subseteq \mathbb{R}^n$ and $x \in K$, the contingent cone to set K at point x , denoted $T_K(x)$, is given by:*

$$T_K(x) = \left\{ z \in \mathbb{R}^n \mid \liminf_{h \rightarrow 0^+} \frac{d_K(x + hz)}{h} = 0 \right\}$$

where $d_K(y)$ denotes the distance of y to K , defined by $d_K(y) = \inf_{y' \in K} \|y - y'\|$.

Definition 9 *Let $\Sigma = (W_1, W_2, X, Y, \mathcal{T})$ be a continuous-time system described by (7). Let $\mathcal{C} = (A_{W_1}, A_{W_2}, G_X, G_Y)$ be an invariance assume-guarantee contract for Σ with respect to $(S_{W_1}, S_{W_2}, S_X, S_Y)$, where the sets S_{W_1}, S_{W_2} are compact. A closed set $K \subseteq X$ is said to be an invariant of Σ relative to the contract $\mathcal{C}$ if the following conditions hold:*

- (i) $X_0 \subseteq K \subseteq S_X \cap h^{-1}(S_Y)$;
- (ii) for all $x \in K$, $F(x, S_{W_1}, S_{W_2}) \subseteq T_K(x)$.

where the set-valued map is given by: $F(\cdot, S_{W_1}, S_{W_2}) = \bigcup_{w_1 \in S_{W_1}} \bigcup_{w_2 \in S_{W_2}} F(\cdot, w_1, w_2)$.

We prove that the existence of an invariant of Σ relative to a contract $\mathcal{C}$ is equivalent to the weak satisfaction of this contract.

Proposition 4 *Let $\Sigma = (W_1, W_2, X, Y, \mathcal{T})$ be a continuous-time system described by (7) such that Assumption 2 holds. Let $\mathcal{C} = (A_{W_1}, A_{W_2}, G_X, G_Y)$ be an invariance assume-guarantee contract for Σ with respect to $(S_{W_1}, S_{W_2}, S_X, S_Y)$, where the sets S_{W_1}, S_{W_2} are compact. Then, $\Sigma \models \mathcal{C}$, if and only if there exists a closed set $K \subseteq X$ that is an invariant of Σ relative to the contract $\mathcal{C}$*

PROOF. First let us prove that the existence of an invariant of Σ relative to a contract $\mathcal{C}$ implies the weak satisfaction of this contract. Let $(w_1, w_2, x, y) : E \rightarrow W_1 \times W_2 \times X \times Y$ in $\mathcal{T}$. Let $t \in E$ and suppose that $w_i|_{[0, t]} \in A_{W_i}$, $i \in \{1, 2\}$. Then, we have for all $s \in [0, t]$, $w_i(s) \in S_{W_i}$, then for

almost all $s \in [0, t]$, $\dot{x}(s) \in F(x(s), S_{W_1}, S_{W_2})$. From Assumption 2, we have $X \subseteq \text{Int}(\text{dom}(F(\cdot, S_{W_1}, S_{W_2})))$ and then $K \subseteq \text{Int}(\text{dom}(F(\cdot, S_{W_1}, S_{W_2})))$. Moreover, from the compactness of S_{W_i} , $i \in \{1, 2\}$, it follows that the set-valued map $F(\cdot, S_{W_1}, S_{W_2})$ is Lipschitz and has compact values. Then, since for all $x \in K$, $F(x, S_{W_1}, S_{W_2}) \subseteq T_K(x)$, we have by Theorem 5.3.4 in [4] that for all $s \in [0, t]$, $x(s) \in K \subseteq S_X$ and then for all $s \in [0, t]$, $y(s) = h(x(s)) \in h(K) \subseteq S_Y$. Then, $x|_{[0, t]} \in G_X$ and $y|_{[0, t]} \in G_Y$. Hence, $\Sigma \models \mathcal{C}$.

We now deal with the second implication. Let us assume that $\Sigma \models \mathcal{C}$. Then for any trajectory $(w_1, w_2, x, y) : E \rightarrow W_1 \times W_2 \times X \times Y$ of the system Σ . We have for all $t \in E$, if for all $s \in [0, t]$, $w_1(s) \in S_{W_1}$ and $w_2(s) \in S_{W_1}$, then for all $s \in [0, t]$, $x(s) \in S_X$ and $y(s) \in S_Y$. Let us prove the existence of a non empty set $K \subseteq X$ satisfying the conditions of Definition 9. Let us define

$$\hat{K} = \{p \in X \mid \exists (w_1, w_2, x, y) : E \rightarrow S_{W_1} \times S_{W_2} \times X \times Y \in \mathcal{T} \text{ with } x(0) \in X_0 \text{ and } \exists t \in E \text{ with } x(t) = p\}. \quad (9)$$

The set $\hat{K}$ is the set of reachable states for the differential inclusion (7) initialized in X_0 , where the external and internal inputs belongs to S_{W_1} and S_{W_2} , respectively. From the satisfaction of the contract, we have that $X_0 \subseteq \hat{K} \subseteq S_X \cap h^{-1}(S_Y)$. Let $(w'_1, w'_2) \in S_{W_1} \times S_{W_2}$ and let us prove that $\hat{K}$ is an invariant for the differential inclusion

$$\dot{x}(t) \in F(x(t), w'_1, w'_2). \quad (10)$$

Let $z^0 \in \hat{K}$, and let $z : E' \rightarrow X$ be a solution of (10) with $z(0) = z^0$. Since $z^0 \in \hat{K}$, we have the existence of a trajectory $\sigma = (w_1, w_2, x, y) : [0, s] \rightarrow S_{W_1} \times S_{W_2} \times X \times Y$ of the system Σ described in (7) such that $x(0) \in X_0$ and $x(s) = z^0$ and for which the external and internal inputs belong to S_{W_1} and S_{W_2} , respectively. Let the time domain E_c defined as follows:

$$\begin{cases} E^c = [0, a + s] \text{ if } E' = [0, a] \\ \quad = [0, a + s] \text{ if } E' = [0, a) \\ \quad = \mathbb{R}_0^+ \text{ if } E' = \mathbb{R}_0^+ \end{cases}$$

and let the trajectory $\sigma^c = (w_1^c, w_2^c, x^c, y^c) : E^c \rightarrow S_{W_1} \times S_{W_2} \times X \times Y$ of the system Σ defined as follows: for all $t \in [0, s]$, $\sigma(t) = \sigma^c(t)$ and for all $t \in E^c \setminus [0, s]$ we have, $x^c(t) = z(t - s)$, $y^c(t) = h(x^c(t))$, $w_1^c(t) = w_1^c$ and $w_2^c(t) = w_2^c$. From construction of $\hat{K}$, we have that $x(t) \in \hat{K}$ for all $t \in E^c$. Hence, for all $t \in E'$, $z(t') = x(t' + s) \in \hat{K}$, where $t' + s \in E^c$. Hence, $\hat{K}$ is an invariant for the differential inclusion (10). Let us now prove that $K = \text{cl}(\hat{K})$ is also an invariant for (10). Let $v^0 \in K$, and let us assume the existence of $v : E \rightarrow X$ solution to (10) with $v(0) = v^0$ and $s \in E$ such that $v(s) \in \bar{K}$. Since, $\bar{K}$ is an open set, we have the existence of $\varepsilon > 0$ such that

$\mathcal{R}_\varepsilon(v(s)) \subseteq \bar{K}$. Then, using the continuity of solutions of (10) in initial conditions (see Corollary 5.3.3 in [4]), we have the existence of $\eta > 0$ and $x^0 \in \hat{K}$ such that $x^0 = x(0) \in \mathcal{R}_\eta(v^0)$ and $x(s) \in \mathcal{R}_\varepsilon(v(s)) \in \bar{K}$, which contradicts the invariance of $\hat{K}$. Hence, $K = \text{cl}(\hat{K})$ is an invariant for the differential inclusion (10).

For $(w'_1, w'_2) \in S_{W_1} \times S_{W_2}$, we have that K is closed. Moreover by Assumption 2, F and thus $F(\cdot, w'_1, w'_2)$ is Lipschitz and has compact values. Moreover, $X \times W_1 \times W_2 \subseteq \text{Int}(\text{dom}(F))$ and thus $X \subseteq \text{Int}(F(\cdot, w'_1, w'_2))$, which in turn implies that $K \subseteq \text{Int}(F(\cdot, w'_1, w'_2))$. Then, from Theorem 5.3.4 in [4], we have

$$\forall x \in K, F(x, w'_1, w'_2) \subseteq T_K(x). \quad (11)$$

Since equation (11) is verified for all $(w'_1, w'_2) \in S_{W_1} \times S_{W_2}$, we have

$$\forall x \in K, F(\cdot, S_{W_1}, S_{W_2}) = \bigcup_{w'_1 \in S_{W_1}} \bigcup_{w'_2 \in S_{W_2}} F(\cdot, w'_1, w'_2) \subseteq T_K(x)$$

Then, K is an invariant of the system Σ relative to the contract $\mathcal{C}$. $\square$

It is noteworthy to mention that in view of Proposition 4, the Lipschitzness property of the system Σ is needed only on a neighbourhood of the set of interest given by $S_X \times S_{W_1} \times S_{W_2}$.

6.2 Composition of invariants

We now provide results allowing us to reason about interconnected systems based on invariants of their components.

Theorem 3 *Consider a network of continuous-time components $\{\Sigma^i\}_{i \in I}$ compatible for composition with respect to $\mathcal{I}$, where the components Σ_i , $i \in I$, have the form of (7) and satisfies Assumption 2. Each component Σ^i have maps and initial sets $F^i, h^i, X_0^i, i \in I$, where h^i is Lipschitz. Let $\mathcal{C}^i = (A_{W_1^i}, A_{W_2^i}, G_{X^i}, G_{Y^i})$ be an invariance assume-guarantee contract for Σ^i with respect to $(S_{W_1^i}, S_{W_2^i}, S_{X^i}, S_{Y^i})$, where the sets $S_{W_1^i}, S_{W_2^i}$ are compact. Let $\mathcal{C} = (\prod_{i \in I} A_{W_1^i}, \prod_{i \in I} G_{X^i}, \prod_{i \in I} G_{Y^i})$ be a contract for the composed system $\Gamma = \langle (\Sigma^i)_{i \in I}, \mathcal{I} \rangle$ and assume the following:*

- (i) for all $i \in I$, $\Sigma^i \models \mathcal{C}^i$;
- (ii) for all $i \in I$, $\prod_{j \in \mathcal{N}(i)} S_{Y^j} \subseteq S_{W_2^i}$.

then $\Gamma \models \mathcal{C}$.

PROOF. First, since h^i is Lipschitz and Assumption 2 holds for all Σ^i , $i \in I$, then from Claim 2 we have that

Assumption 2 holds for $\Gamma = \langle (\Sigma^i)_{i \in I}, \mathcal{I} \rangle$. Hence, using the equivalence between the invariance relative to contracts and the weak satisfaction of contracts from Proposition 4, we have the existence of a closed set K^i that is invariant for Σ^i relative to the contract $\mathcal{C}^i$.

To complete the proof, let us show that the set $K = \prod_{i \in I} K^i$ is an invariant of Γ relative to the contract $\mathcal{C}$. Let $i \in I$ and consider a trajectory $x^i : \mathbb{R}_0^+ \rightarrow X^i$ with $x^i(0) \in K^i$ and satisfying $\dot{x}^i(t) \in F^i(x^i, S_{W_1^i}, S_{W_2^i})$, for all $t \in \mathbb{R}_0^+$. Using the fact that K^i invariant of Σ^i relative to the contract $\mathcal{C}^i$ and from Proposition 4 we have that $x^i(t) \in K^i$, for all $t \in \mathbb{R}_0^+$.

Consider the map $G : X \rightarrow 2^X$ defined for $x \in X$ by:

$$G(x) = \prod_{i \in I} F^i(x^i, S_{W_1^i}, S_{W_2^i}); \quad (12)$$

Let a trajectory $x : \mathbb{R}_0^+ \rightarrow X$ with $x(0) \in K$ and satisfying $\dot{x}(t) \in G(x(t))$ for all $t \in \mathbb{R}_0^+$. We have from above that $x(t) \in K$, for all $t \in \mathbb{R}_0^+$. Moreover, using the fact that G is Lipschitz, has compact values, $X \subseteq \text{Int}(\text{dom}(G))$ and K is a closed set, we have by Theorem 5.3.4 in [4] that $G(x) \subseteq T_K(x)$ for all $x \in K$.

Let us now consider the system described by the following differential inclusion:

$$\dot{x}(t) \in F(x(t), S_{W_1}). \quad (13)$$

Where $x(t) = (x_1(t), \dots, x_N(t))$ and $S_{W_1} = \prod_{i \in I} S_{W_1^i}$.

Let $x \in K$, then for all $i \in I$, we have that

$$\begin{aligned} F^i(x^i, S_{W_1^i}, w_2^i) &= F^i(x^i, S_{W_1^i}, \prod_{j \in \mathcal{N}(i)} \{h^j(x^j)\}) \\ &\subseteq F^i(x^i, S_{W_1^i}, \prod_{j \in \mathcal{N}(i)} \{h^j(K^j)\}) \\ &\subseteq F^i(x^i, S_{W_1^i}, \prod_{j \in \mathcal{N}(i)} \{S_{Y^j}\}) \\ &\subseteq F^i(x^i, S_{W_1^i}, S_{W_2^i}). \end{aligned}$$

Where the first equality comes from the definition of an interconnection relation, the second inclusion comes from (i) and the last inclusion comes from (ii). Then, we have that $F(x, S_{W_1}) \subseteq G(x) \subseteq T_K(x)$, for all $x \in K$. Finally, we have $X_0 = \prod_{i \in I} X_0^i \subseteq \prod_{i \in I} K^i = K$, $K = \prod_{i \in I} K^i \subseteq \prod_{i \in I} G_{X^i}$, and $K = \prod_{i \in I} K^i \subseteq \prod_{i \in I} (h^i)^{-1}(G_{Y^i}) = h^{-1}(G_Y)$. Hence, K is an invariant of Γ relative to the contract $\mathcal{C}$. Then from Proposition 4 and Claim 2, we get $\Gamma \models \mathcal{C}$. $\square$

We show an example to illustrate the application of the previous theorem.

Example 11 Consider systems $\Sigma^i = (W_1^i, W_2^i, X_i, Y_i, \mathcal{T}_i)$, $i = 1, 2$ where $W_1^i = W_2^i = X^i = Y^i = \mathbb{R}$. A trajectory of Σ^i is a triple $(w_1^i, w_2^i, x^i, y^i) : E \rightarrow W_1^i \times W_2^i \times X^i \times Y^i$ in $\mathcal{T}_i$ where $E = \mathbb{R}_0^+$, w_1^i and w_2^i are locally measurable, x^i and y^i are absolutely continuous and continuous, respectively, and satisfy for almost all $t \in E$:

$$\begin{cases} \dot{x}^i(t) = f^i(x^i(t), w_1^i(t), w_2^i(t)) \\ \quad = -a_i x^i(t) + a_i w_2^i(t) + w_1^i(t), \\ y^i(t) = h^i(x(t)) = x^i(t). \end{cases}$$

where $x_i(0) \in [0, b_i]$ with $a_i, b_i \in \mathbb{R}_0^+$, let $b = \max(b_1, b_2)$. It can be seen that h_i is Lipschitz and that Assumption 2 holds for Σ_i . Let the interconnection relation $\mathcal{I} = \{(1, 2), (2, 1)\}$. It is clear that $\{\Sigma^i\}_{i \in I}$ is compatible for composition with respect to $\mathcal{I}$. Let the invariance contract $\mathcal{C}^i = (A_{W_1^i}, A_{W_2^i}, G_{X^i}, G_{Y^i})$ for the system Σ^i with respect to $(S_{W_1^i}, S_{W_2^i}, S_{X^i}, S_{Y^i})$, with $S_{W_1^i} = \{0\}$, $S_{W_2^i} = S_{X^i} = S_{Y^i} = [0, b]$. We can easily check that for all $x^i \in [0, b]$, $f_i(x^i, [0, b], \{0\}) \subseteq T_{[0, b]}(x^i)$, since

$$T_{[0, b]}(x^i) = \begin{cases} \mathbb{R}^+ & \text{if } x^i = 0, \\ \mathbb{R}^- & \text{if } x^i = b, \\ \mathbb{R} & \text{if } x^i \in (0, b) \end{cases}$$

Then $[0, b]$ is an invariant of the system Σ_i , relative to the contract $\mathcal{C}^i$. By Theorem 3, $[0, b]^2$ is an invariant of the composed system $\Gamma = \langle (\Sigma^i)_{i \in I}, \mathcal{I} \rangle$ relative to the composed contract $\mathcal{C}$.

7 Small gain results

In this part, we show how the proposed framework can recover different versions of the classical small gain theorem as a particular case. Indeed, we show how the framework allows to recover the classical BIBO stability result [12]. Moreover, we construct a new small-gain result for the concept of growth bound [2]. To the best of our knowledge, this result is new and have not been investigated before in the literature. We suppose for the sake of simplicity that for each system $\Sigma = (W_1, W_2, X, Y, \mathcal{T})$, we have $W_1 = \{0\}$, $X = Y = W_2 = \mathbb{R}^n$ and for all $(w_1, w_2, x, y) : \mathbb{R}_0^+ \rightarrow W_1 \times W_2 \times W \times X \times Y$ in $\mathcal{T}$, $x(t) = y(t)$, for all $t \in \mathbb{R}_0^+$.

7.1 BIBO stability

Given a system Σ satisfying a BIBO stability condition [12], we show that if the gain of the system is lower than 1 then the feedback³ composed system is bounded

³ Given a system Σ^1 and a set of vertices $I = \{1\}$, the feedback composition of the system Σ^1 is the composition with an interconnection relation $\mathcal{I} = \{(1, 1)\}$.

for all the time domain.

Theorem 4 Consider a component $\Sigma^1 = (\{0\}, W_2, X, Y, \mathcal{T})$, $I = \{1\}$ and let the interconnection relation $\mathcal{I} = \{(1, 1)\}$ such that $\{\Sigma^i\}_{i \in I}$ is compatible for composition with respect to $\mathcal{I}$. Let $\gamma < 1$ and $\beta \in \mathbb{R}_0^+$ such that for any trajectory $(0, w_2, x, y) : \mathbb{R}_0^+ \rightarrow W_1 \times W_2 \times X \times Y$ in $\mathcal{T}$, $x : \mathbb{R}_0^+ \rightarrow X$ is continuous, $\|x(0)\| \leq \frac{\beta}{1-\gamma}$ and for all $t \in \mathbb{R}_0^+$ we have:

$$\|x_{|[0, t]}\|_\infty \leq \gamma \|w_{2|[0, t]}\|_\infty + \beta. \quad (14)$$

Then for any trajectory $(0, x, y) : \mathbb{R}_0^+ \rightarrow \{0\} \times X \times Y$ of the composed system $\Gamma = \langle (\Sigma^i)_{i \in I}, \mathcal{I} \rangle$, we have for all $t \in \mathbb{R}_0^+$: $\|x_{|[0, t]}\|_\infty \leq \frac{\beta}{1-\gamma}$.

PROOF. We first start by constructing a suitable contract for the system Σ^1 . Let the map $a : \mathbb{R}_0^+ \rightarrow \mathbb{R}_0^+$, a parameter $\varepsilon > 0$ and a parametrized contract $\mathcal{C}(\varepsilon) = (A_{W_1}^\varepsilon, A_{W_2}^\varepsilon, G_X^\varepsilon, G_Y^\varepsilon)$ for Σ , where:

- $A_{W_1}^\varepsilon = \{w_1 : \mathbb{R}_0^+ \rightarrow W_1 \in M(W_1) \mid \forall t \in \mathbb{R}_0^+, w_1(t) = 0\}$;
- $A_{W_2}^\varepsilon = \{w_2 : \mathbb{R}_0^+ \rightarrow W_2 \in M(W_2) \mid \forall t \in \mathbb{R}_0^+, \|w_{2|[0, t]}\|_\infty \leq a(\varepsilon)\}$;
- $G_X^\varepsilon = G_Y^\varepsilon = \{x : \mathbb{R}_0^+ \rightarrow X \in M(X) \mid \forall t \in \mathbb{R}_0^+, \|x_{|[0, t]}\|_\infty \leq \gamma a(\varepsilon) + \beta\}$.

Let us choose $a(\varepsilon) = \frac{\beta + \varepsilon}{1 - \gamma}$, where $\varepsilon > 0$. We have that

$$\|x(0)\| \leq \frac{\beta}{1 - \gamma} \leq \gamma a(\varepsilon) + \beta = \frac{\beta + \gamma \varepsilon}{1 - \gamma}$$

for any $\varepsilon > 0$. Hence, $x_{|[0, 0]} \in G_X^\varepsilon$. We also have from (14) that $\Sigma^1 \models \mathcal{C}(\varepsilon)$, and for all trajectories $(w_1, w_2, x, y) \in \mathcal{T}$, $x : \mathbb{R}_0^+ \rightarrow X$ is continuous. Then, from Proposition 2, we have that $\Sigma^1 \models_s \mathcal{C}'(\varepsilon)$ for any $\varepsilon > 0$, where $\mathcal{C}'(\varepsilon) = (A_{W_1}^\varepsilon, A_{W_2}^\varepsilon, G_X^\varepsilon, \mathcal{B}_\varepsilon(G_Y^\varepsilon) \cap M(Y))$. Now, using the fact that $\gamma a(\varepsilon) + \beta - a(\varepsilon) = -\varepsilon < 0$, we have that $\mathcal{B}_\varepsilon(G_Y^\varepsilon) \cap M(Y) \subseteq A_{W_2}^\varepsilon$. Moreover, from continuity of $x : \mathbb{R}_0^+ \rightarrow X$ Assumption 1 is satisfied. Then from Theorem 2, the composed system $\Gamma = \langle (\Sigma^i)_{i \in I}, \mathcal{I} \rangle$ satisfies the composed contract $\mathcal{C}_\Gamma^\varepsilon = (A_{W_1}^\varepsilon, G_X^\varepsilon, \mathcal{B}_\varepsilon(G_Y^\varepsilon) \cap M(Y))$. Then, we have for all $t \in \mathbb{R}_0^+$: $\|x_{|[0, t]}\|_\infty \leq \gamma a(\varepsilon) + \beta = \frac{\beta + \gamma \varepsilon}{1 - \gamma}$.

Since the last inequality is verified for all $\varepsilon > 0$ we have for all $t \in \mathbb{R}_0^+$, $\|x_{|[0, t]}\|_\infty \leq \frac{\beta}{1 - \gamma}$. $\square$

7.2 Growth bound

The notion of growth bound allows us to analyse the growth or contraction properties of a system;

particularly, it coincides with forward completeness (see Corollary 2.3 in [2]) for finite-dimensional systems described by nonlinear differential equations $\dot{x}(t) = F(x(t), w_1(t), w_2(t))$ and with a locally Lipschitz map F . Given a continuous-time system with a given growth bound, in the following we show how to characterize the growth bound of the feedback composed system.

Theorem 5 Consider a component $\Sigma^1 = (\{0\}, W_2, X, Y, \mathcal{T})$, $I = \{1\}$ and let $\mathcal{I} = \{(1, 1)\}$ be the interconnection relation such that $\{\Sigma^i\}_{i \in I}$ is compatible for composition with respect to $\mathcal{I}$. Let $\gamma_1, \gamma_2, \gamma_3$ be class $\mathcal{K}$ maps and constants $(c, \rho) \in \mathbb{R} \times [0, 1)$, where $\gamma_3(\cdot) < \rho \text{Id}(\cdot)$ ⁴ and such that for any trajectory $(0, w_2, x, y) : \mathbb{R}_0^+ \rightarrow W_1 \times W_2 \times X \times Y$ in $\mathcal{T}$, $x : \mathbb{R}_0^+ \rightarrow X$ is continuous, $\|x(0)\| \leq \gamma_2(\|x(0)\|) + c$ and for all $t \in \mathbb{R}_0^+$ we have:

$$\|x(t)\| \leq \gamma_1(t) + \gamma_2(\|x(0)\|) + \gamma_3(\|w_2|_{[0,t]}\|_\infty) + c. \quad (15)$$

Then there exist $\mathcal{K}$ functions α_1, α_2 and $c' \in \mathbb{R}$ such that for any trajectory $(0, x, y) : \mathbb{R}_0^+ \rightarrow \{0\} \times X \times Y$ of the composed system $\Gamma = \langle (\Sigma^i)_{i \in I}, \mathcal{I} \rangle$ we have for all $t \in \mathbb{R}_0^+$, $\|x(t)\| \leq \alpha_1(t) + \alpha_2(\|x(0)\|) + c'$, where $\alpha_1 = \frac{2}{1-\rho}\gamma_1$, $\alpha_2 = \frac{2}{1-\rho}\gamma_2$, and $c' = \frac{2c}{1-\rho}$.

PROOF. We first define the system $\Sigma_{x(0)}^1 = (\{0\}, W_2, X, Y, \mathcal{T}_{x(0)})$ where $(0, w_2, x, y) : \mathbb{R}_0^+ \rightarrow \{0\} \times W_2 \times X \times Y \in \mathcal{T}_{x(0)}$ is a trajectory of the system $\Sigma_{x(0)}^1$ is and only if it is a trajectory of the system Σ^1 initialized in $x(0) \in X$.

We start by constructing a suitable contract for the system $\Sigma_{x(0)}^1$. Let the map $a : \mathbb{R}_0^+ \times \mathbb{R}_0^+ \rightarrow \mathbb{R}_0^+$. A parameter $\varepsilon > 0$ and a parametrized contract $\mathcal{C}(\varepsilon) = (A_{W_1}^\varepsilon, A_{W_2}^\varepsilon, G_X^\varepsilon, G_Y^\varepsilon)$ for Σ^1 , where:

- $A_{W_1}^\varepsilon = \{w_1 : \mathbb{R}_0^+ \rightarrow W_1 \in M(W_1) \mid \forall t \in \mathbb{R}_0^+, w_1(t) = 0\}$;
- $A_{W_2}^\varepsilon = \{w_2 : \mathbb{R}_0^+ \rightarrow W_2 \in M(W_2) \mid \forall t \in \mathbb{R}_0^+, \|w_2(t)\| \leq a(t, \varepsilon)\}$;
- $G_X^\varepsilon = G_Y^\varepsilon = \{x : \mathbb{R}_0^+ \rightarrow X \in M(X) \mid \forall t \in \mathbb{R}_0^+, \|x(t)\| \leq \gamma_1(t) + \gamma_2(\|x(0)\|) + \gamma_3(\|a(t, \varepsilon)\|) + c\}$.

Let us choose the map $a : \mathbb{R}_0^+ \times \mathbb{R}_0^+ \rightarrow \mathbb{R}_0^+$ satisfying

$$a(t, \varepsilon) = (1 - \rho)^{-1}(\gamma_1(t) + \gamma_2(\|x(0)\|) + c + \varepsilon)$$

where $\varepsilon > 0$. First, since $\rho < 1$, we have that $(1 - \rho)\text{Id}$ is a class $\mathcal{K}$ map, which implies from Lemma 4.2 in [21]) that $(1 - \rho)^{-1}\text{Id}$ is class $\mathcal{K}$. Moreover, since γ_1 is class $\mathcal{K}$ we have for all $t \in \mathbb{R}_0^+$,

$$\|a|_{[0,t]}(\cdot, \varepsilon)\|_\infty = \|a(t, \varepsilon)\| \quad (16)$$

⁴ i.e., $\gamma_3(r) < \rho r$ for all $r > 0$

Let us now prove that $\Sigma^1 \models \mathcal{C}(\varepsilon)$. Let $t \in \mathbb{R}_0^+$ and assume that $\|w_2(s)\| \leq a(s, \varepsilon)$ for all $s \in [0, t]$. We have from (15) that for all $s \in [0, t]$

$$\begin{aligned} \|x(s)\| &\leq \gamma_1(s) + \gamma_2(\|x(0)\|) + \gamma_3(\|w_2|_{[0,s]}\|_\infty) + c \\ &\leq \gamma_1(s) + \gamma_2(\|x(0)\|) + \gamma_3(\|a|_{[0,s]}(\cdot, \varepsilon)\|_\infty) + c \\ &\leq \gamma_1(s) + \gamma_2(\|x(0)\|) + \gamma_3(\|a(s, \varepsilon)\|) + c \end{aligned}$$

where the last inequality comes from (16). Hence, $\Sigma^1 \models \mathcal{C}(\varepsilon)$. Moreover we have that for all trajectories $(w_1, w_2, x, y) \in \mathcal{T}$, $x : \mathbb{R}_0^+ \rightarrow X$ is continuous and using the fact that $\|x(0)\| \leq \gamma_2(\|x(0)\|) + c$, we have that $x|_{[0,0]} \in G_X^\varepsilon$, for all $\varepsilon > 0$. Then, from Proposition 2, we have that $\Sigma^1 \models_s \mathcal{C}'(\varepsilon)$ for any $\varepsilon > 0$, where $\mathcal{C}'(\varepsilon) = (A_{W_1}^\varepsilon, A_{W_2}^\varepsilon, G_X^\varepsilon, \mathcal{B}_\varepsilon(G_Y^\varepsilon) \cap M(Y))$. Now, using the fact that $\gamma_3 < \rho \text{Id}$ we have

$$\begin{aligned} \gamma_1(t) + \gamma_2(\|x(0)\|) + \gamma_3(a(t, \varepsilon)) + c - a(t, \varepsilon) &\leq \gamma_1(t) + \\ &\quad \gamma_2(\|x(0)\|) + \rho a(t, \varepsilon) + c - a(t, \varepsilon) \\ &= -\varepsilon < 0 \end{aligned}$$

which implies that $\mathcal{B}_\varepsilon(G_Y^\varepsilon) \cap M(Y) \subseteq A_{W_2}^\varepsilon$. Moreover, from continuity of $x : \mathbb{R}_0^+ \rightarrow X$, Assumption 1 is satisfied. Then from Theorem 2, the composed system $\Gamma = \langle (\Sigma^i)_{i \in I}, \mathcal{I} \rangle$ satisfies the composed contract $\mathcal{C}_\Gamma^\varepsilon = (A_{W_1}^\varepsilon, G_X^\varepsilon, \mathcal{B}_\varepsilon(G_Y^\varepsilon) \cap M(Y))$. Then, we have for all $t \in \mathbb{R}_0^+$:

$$\begin{aligned} \|x(t)\| &\leq \gamma_1(t) + \gamma_2(\|x(0)\|) + \gamma_3(a(t, \varepsilon)) + c \\ &\leq a(t, \varepsilon) = (1 - \rho)^{-1}(\gamma_1(t) + \gamma_2(\|x(0)\|) + c + \varepsilon). \end{aligned}$$

The last inequality is verified for all $\varepsilon > 0$, which implies from the continuity of $(1 - \rho)^{-1}\text{Id}$ that for all $t \in \mathbb{R}_0^+$:

$$\begin{aligned} \|x(t)\| &\leq (1 - \rho)^{-1}(\gamma_1(t) + \gamma_2(\|x(0)\|) + c) \\ &\leq \frac{2}{1 - \rho}\gamma_1(t) + \frac{2}{1 - \rho}\gamma_2(\|x(0)\|) \\ &\quad + \frac{2c}{1 - \rho} \end{aligned}$$

where the last inequality comes from the fact that $(1 - \rho)\text{Id}$ is a class $\mathcal{K}$ and the weak triangular inequality [19]. By choosing $\alpha_1 = \frac{2}{1-\rho}\gamma_1$, $\alpha_2 = \frac{2}{1-\rho}\gamma_2$, and $c' = \frac{2c}{1-\rho}$ where α_1 and α_2 are class $\mathcal{K}$ (see Lemma 4.2 in [21]), we have for all $t \in \mathbb{R}_0^+$, $\|x(t)\| \leq \alpha_1(t) + \alpha_2(\|x(0)\|) + c'$. $\square$

We can mention that for finite-dimensional systems, described by nonlinear differential equations and with a locally Lipschitz map F , the previous result states that if a system is forward complete with a gain γ_3 lower than identity, then the feedback composed system is forward complete.

Remark 8 Let us mention that using the same approach, and similar to the work of [11], one can generalize

Fig. 5. Summary of main results in the paper

different small-gain results to different interconnection structures.

8 Conclusion

In this paper, we proposed a contract based approach for verifying compositionally properties of continuous-time interconnected systems. The main notions considered in the paper and their relationships are sketched in Figure 5. The main contributions are summarized below. We introduced a notion of assume-guarantee contracts equipped with a weak and a strong semantics. We showed that weak semantics are sufficient to deal with acyclic interconnections (Theorem 1), strong semantics are required to reason on cyclic interconnections (Theorem 2 and Example 6) and that strong semantics of a contract can sometimes be obtained from weak ones (Propositions 2 and 3).

We then developed specific results for systems described by differential inclusions and invariance assume-guarantee contracts. We showed that sufficient and necessary conditions for weak satisfaction of contracts can be given using invariant sets (Proposition 4) and that invariants are compatible with cyclic interconnections (Theorem 3). Finally, we have shown how the proposed assume-guarantee framework can recast different versions of the small-gain theorem as a particular case (Theorems 4 and 5).

Appendix

Proof of Claim 2

Consider a network of components $\{\Sigma^i\}_{i \in I}$, compatible for composition w.r.t. $\mathcal{I}$, where each component has the form of (7). Each component Σ^i has maps and initial sets F^i , h^i , X_0^i , $i \in I$, and the composed system $\Gamma = \langle (\Sigma^i)_{i \in I}, \mathcal{I} \rangle$ can be written as in (8). Assume that h^i is Lipschitz and Assumption 2 holds for all Σ_i , $i \in I$, and let us show that $\Gamma = \langle (\Sigma^i)_{i \in I}, \mathcal{I} \rangle$ satisfies Assumption 2.

First, using the fact that for all $i \in I$, F^i has compact values and $X^i \times W_1^i \times W_2^i \subseteq \text{Int}(\text{dom}(F^i))$, we have

directly from (8) that F has compact values and $X \times W_1 \subseteq \text{Int}(\text{dom}(F))$.

Similarly, using the fact that for all $i \in I$, h^i satisfies $X^i \subseteq \text{Int}(\text{dom}(h^i))$ and $h^i(X^i) \subseteq Y^i$, we directly get from (8) that h satisfies $X \subseteq \text{Int}(\text{dom}(h))$ and $h(X) \subseteq Y$.

Now assume that for all $i \in I$, F^i and h^i are Lipschitz, and let us show that the interconnected system $\Gamma = \langle (\Sigma^i)_{i \in I}, \mathcal{I} \rangle$ is Lipschitz. From Lipschitzness of F^i , we have the existence of $L^i, L_1^i, L_2^i > 0$ such that for all $(x_a^i, x_b^i) \in X^i$, for all $(w_{1,a}^i, w_{1,b}^i) \in W_1^i$ and for all $(w_{2,a}^i, w_{2,b}^i) \in W_2^i$, we have :

$$F^i(x_a^i, w_{1,a}^i, w_{2,a}^i) \subseteq F^i(x_b^i, w_{1,b}^i, w_{2,b}^i) + L^i \|x_a^i - x_b^i\| \mathbb{B} \\ + L_1^i \|w_{1,a}^i - w_{1,b}^i\| \mathbb{B} + L_2^i \|w_{2,a}^i - w_{2,b}^i\| \mathbb{B}$$

where $\|\cdot\|$ is the infinity norm of appropriate dimension and $\mathbb{B}$ is the ball centred at the origin with radius 1 of appropriate dimension. Similarly, from Lipschitzness of h^i , we have the existence of $l^i > 0$ such that for all $(x_a^i, x_b^i) \in X^i$ we have:

$$\|h^i(x_a^i) - h^i(x_b^i)\| \leq l^i \|x_a^i - x_b^i\|$$

Consider $(x_a, x_b) \in X$ and $(w_{1,a}, w_{1,b}) \in W_1$ with $x_a = (x_a^1, \dots, x_a^N)$, $x_b = (x_b^1, \dots, x_b^N)$, $w_{1,a} = (w_{1,a}^1, \dots, w_{1,a}^N)$, and $w_{1,b} = (w_{1,b}^1, \dots, w_{1,b}^N)$. Choosing $L = \max_{i \in I} L^i$, $L_1 = \max_{i \in I} L_1^i$, $L_2 = \max_{i \in I} L_2^i$ and $l = \max_{i \in I} l^i$, we have the chain of inclusions in (17). Hence, the interconnected system $\Gamma = \langle (\Sigma^i)_{i \in I}, \mathcal{I} \rangle$ is Lipschitz.

References

- [1] R. Alur, R. Grosu, I. Lee, and O. Sokolsky. Compositional refinement for hierarchical hybrid systems. In *International Workshop on Hybrid Systems: Computation and Control*, pages 33–48. Springer, 2001.
- [2] David Angeli and Eduardo D Sontag. Forward completeness, unboundedness observability, and their lyapunov characterizations. *Systems & Control Letters*, 38(4-5):209–217, 1999.
- [3] M. Arcak, C. Meissen, and A Packard. *Networks of dissipative systems: compositional certification of stability, performance, and safety*. Springer, 2016.
- [4] J.-P. Aubin. *Viability theory*. Springer Science & Business Media, 2009.
- [5] Christel Baier and Joost-Pieter Katoen. *Principles of model checking*. MIT press, 2008.
- [6] Albert Benveniste, Benoît Caillaud, Dejan Nickovic, Roberto Passerone, Jean-Baptiste Raclet, Philipp Reinkemeier, Alberto Sangiovanni-Vincentelli, Werner Damm, Thomas A

$$\begin{aligned}
F(x_a, w_{1,a}) &= \prod_{i \in I} F^i(x_a^i, w_{1,a}^i, w_{2,a}^i) \\
&\subseteq \prod_{i \in I} \left(F^i(x_b^i, w_{1,b}^i, w_{2,b}^i) + L^i \|x_a^i - x_b^i\|_{\mathbb{B}} + L_1^i \|w_{1,a}^i - w_{1,b}^i\|_{\mathbb{B}} + L_2^i \|w_{2,a}^i - w_{2,b}^i\|_{\mathbb{B}} \right) \\
&\subseteq \prod_{i \in I} \left(F^i(x_b^i, w_{1,b}^i, w_{2,b}^i) + L \|x_a^i - x_b^i\|_{\mathbb{B}} + L_1 \|w_{1,a}^i - w_{1,b}^i\|_{\mathbb{B}} + L_2 \|w_{2,a}^i - w_{2,b}^i\|_{\mathbb{B}} \right) \\
&\subseteq \prod_{i \in I} \left(F^i(x_b^i, w_{1,b}^i, w_{2,b}^i) + L \|x_a^i - x_b^i\|_{\mathbb{B}} + L_1 \|w_{1,a}^i - w_{1,b}^i\|_{\mathbb{B}} + L_2 \left\| \prod_{j \in \mathcal{N}(i)} \{h^j(x_a^j)\} - \prod_{j \in \mathcal{N}(i)} \{h^j(x_b^j)\} \right\|_{\mathbb{B}} \right) \\
&\subseteq \prod_{i \in I} \left(F^i(x_b^i, w_{1,b}^i, w_{2,b}^i) + L \|x_a^i - x_b^i\|_{\mathbb{B}} + L_1 \|w_{1,a}^i - w_{1,b}^i\|_{\mathbb{B}} + L_2 \left\| \prod_{j \in I} \{h^j(x_a^j)\} - \prod_{j \in I} \{h^j(x_b^j)\} \right\|_{\mathbb{B}} \right) \\
&\subseteq \prod_{i \in I} \left(F^i(x_b^i, w_{1,b}^i, w_{2,b}^i) + L \|x_a^i - x_b^i\|_{\mathbb{B}} + L_1 \|w_{1,a}^i - w_{1,b}^i\|_{\mathbb{B}} + L_2 \|x_a - x_b\|_{\mathbb{B}} \right) \tag{17} \\
&\subseteq \prod_{i \in I} \left(F^i(x_b^i, w_{1,b}^i, w_{2,b}^i) + L \|x_a - x_b\|_{\mathbb{B}} + L_1 \|w_{1,a} - w_{1,b}\|_{\mathbb{B}} + L_2 \|x_a - x_b\|_{\mathbb{B}} \right) \\
&\subseteq \prod_{i \in I} \left(F^i(x_b^i, w_{1,b}^i, w_{2,b}^i) + (L + L_2 l) \|x_a - x_b\|_{\mathbb{B}} + L_1 \|w_{1,a} - w_{1,b}\|_{\mathbb{B}} \right) \\
&\subseteq \prod_{i \in I} \left(F^i(x_b^i, w_{1,b}^i, w_{2,b}^i) \right) + (L + L_2 l) \|x_a - x_b\|_{\mathbb{B}} + L_1 \|w_{1,a} - w_{1,b}\|_{\mathbb{B}} \\
&= F(x_b, w_{1,b}) + (L + L_2 l) \|x_a - x_b\|_{\mathbb{B}} + L_1 \|w_{1,a} - w_{1,b}\|_{\mathbb{B}}
\end{aligned}$$

-
- Henzinger, Kim G Larsen, et al. Contracts for system design. *Foundations and Trends® in Electronic Design Automation*, 12(2-3):124–400, 2018.
- [7] Yuxiao Chen, James Anderson, Karan Kalsi, Steven H Low, and Aaron D Ames. Compositional set invariance in network systems with assume-guarantee contracts. In *2019 American Control Conference (ACC)*, pages 1027–1034. IEEE, 2019.
- [8] C. Conte, N.R. Voellmy, M.N. Zeilinger, M. Morari, and C.N. Jones. Distributed synthesis and control of constrained linear systems. In *American Control Conference*, pages 6017–6022, 2012.
- [9] S. Coogan and M. Arcak. A dissipativity approach to safety verification for interconnected systems. *IEEE Transactions on Automatic Control*, 60(6):1722–1727, 2015.
- [10] E. Dallal and P. Tabuada. On compositional symbolic controller synthesis inspired by small-gain theorems. In *IEEE Conference on Decision and Control*, pages 6133–6138, 2015.
- [11] S. Dashkovskiy, B.S. Rüffer, and F.R. Wirth. An iss small gain theorem for general networks. *Mathematics of Control, Signals, and Systems*, 19(2):93–122, 2007.
- [12] C.A. Desoer and M. Vidyasagar. *Feedback systems: input-output properties*, volume 55. Siam, 1975.
- [13] A. El-Guindy, Y.C. Chen, and M. Althoff. Compositional transient stability analysis of power systems via the computation of reachable sets. In *American Control Conference*, pages 2536–2543. IEEE, 2017.
- [14] Alina Eqtami and Antoine Girard. A quantitative approach on assume-guarantee contracts for safety of interconnected systems. In *2019 18th European Control Conference (ECC)*, pages 536–541. IEEE, 2019.
- [15] Goran Fedja Frehse. *Compositional verification of hybrid systems using simulation relations*. [SI: sn], 2005.
- [16] Kasra Ghasemi, Sadra Sadraddini, and Calin Belta. Compositional synthesis via a convex parameterization of assume-guarantee contracts. In *Proceedings of the 23rd International Conference on Hybrid Systems: Computation and Control*, pages 1–10, 2020.
- [17] R. Goebel, R. G. Sanfelice, and A. R. Teel. *Hybrid Dynamical Systems: Modeling, Stability, and Robustness*. Princeton University Press, New Jersey, 2012.
- [18] T.A. Henzinger, M. Minea, and V. Prabhu. Assume-guarantee reasoning for hierarchical hybrid systems. In *International Workshop on Hybrid Systems: Computation and Control*, pages 275–290. Springer, 2001.
- [19] Z-P Jiang, Andrew R Teel, and L Praly. Small-gain theorem for iss systems and applications. *Mathematics of Control, Signals and Systems*, 7(2):95–120, 1994.
- [20] F. Kerber and A.J. Van Der Schaft. Assume-guarantee reasoning for linear dynamical systems. In *European Control Conference*, pages 5015–5020. IEEE, 2009.
- [21] Hassan K Khalil. *Nonlinear systems*. Prentice-Hall, New Jersey, 2(5):5–1, 1996.
- [22] E.S. Kim, M. Arcak, and S.A. Seshia. Compositional controller synthesis for vehicular traffic networks. In *IEEE Conference on Decision and Control*, pages 6165–6171, 2015.
- [23] E.S. Kim, M. Arcak, and S.A. Seshia. A small gain theorem for parametric assume-guarantee contracts. In *Proceedings of the 20th International Conference on Hybrid Systems: Computation and Control*, pages 207–216. ACM, 2017.
- [24] A. Le Coënt, L. Fribourg, N. Markey, F. De Vuyst, and L. Chamoin. Distributed synthesis of state-dependent

- switching control. In *International Workshop on Reachability Problems*, pages 119–133, 2016.
- [25] E.A. Lee and S.A. Seshia. *Introduction to embedded systems: A cyber-physical systems approach*. Mit Press, 2016.
 - [26] Kaushik Mallik, Anne-Kathrin Schmuck, Sadegh Soudjani, and Rupak Majumdar. Compositional synthesis of finite-state abstractions. *IEEE Transactions on Automatic Control*, 64(6):2629–2636, 2018.
 - [27] P.-J. Meyer and D.V. Dimarogonas. Compositional abstraction refinement for control synthesis. *Nonlinear Analysis: Hybrid Systems*, 27:437–451, 2018.
 - [28] Pierre-Jean Meyer, Antoine Girard, and Emmanuel Witrant. Compositional abstraction and safety synthesis using overlapping symbolic models. *IEEE Transactions on Automatic Control*, 63(6):1835–1841, 2017.
 - [29] P. Nilsson and N. Ozay. Synthesis of separable controlled invariant sets for modular local control design. In *American Control Conference*, pages 5656–5663, 2016.
 - [30] S.V. Raković, B. Kern, and R. Findeisen. Practical set invariance for decentralized discrete time systems. In *IEEE Conference on Decision and Control*, pages 3283–3288, 2010.
 - [31] R. Tyrrell Rockafellar and Roger J-B Wets. *Variational analysis*. Springer Science Business Media, 2009.
 - [32] A. Saoud, A. Girard, and L. Fribourg. Contract based design of symbolic controllers for interconnected multiperiodic sampled-data systems. In *Conference on Decision and Control*, pages 773–779, 2018.
 - [33] A. Saoud, A. Girard, and L. Fribourg. Contract based design of symbolic controllers for vehicle platooning. In *Proceedings of the 21st International Conference on Hybrid Systems: Computation and Control*, HSCC '18, pages 277–278, 2018.
 - [34] A. Saoud, A. Girard, and L. Fribourg. On the composition of discrete and continuous-time assume-guarantee contracts for invariance. In *European Control Conference*, pages 435–440, 2018.
 - [35] A. Saoud, P. Jagtap, M. Zamani, and A. Girard. Compositional abstraction-based synthesis for cascade discrete-time control systems. *IFAC-PapersOnLine*, 51(16):13 – 18, 2018. 6th IFAC Conference on Analysis and Design of Hybrid Systems.
 - [36] Adnane Saoud. *Compositional and Efficient Controller Synthesis for Cyber-Physical Systems*. PhD thesis, Université Paris Saclay, 2019.
 - [37] C. Sloth, G.J. Pappas, and R. Wisniewski. Compositional safety analysis using barrier certificates. In *International Conference on Hybrid Systems: Computation and Control*, pages 15–24, 2012.
 - [38] S. W. Smith, P. Nilsson, and N. Ozay. Interdependence quantification for compositional control synthesis with an application in vehicle safety systems. In *Conference on Decision and Control*, pages 5700–5707. IEEE, 2016.
 - [39] AJ Van der Schaft. Equivalence of dynamical systems by bisimulation. *IEEE transactions on automatic control*, 49(12):2160–2172, 2004.
 - [40] Majid Zamani and Murat Arcak. Compositional abstraction for networks of control systems: A dissipativity approach. *IEEE Transactions on Control of Network Systems*, 5(3):1003–1015, 2017.
 - [41] Daniele Zonetti, Adnane Saoud, Antoine Girard, and Laurent Fribourg. A symbolic approach to voltage stability and power sharing in time-varying dc microgrids. In *2019 18th European Control Conference (ECC)*, pages 903–909. IEEE, 2019.