



**HAL**  
open science

## Non-coding RNAs in cardiac autophagy following myocardial infarction

Annie Turkieh, Henri Charrier, Emilie Dubois-Deruy, Sina Porouchani,  
Marion Bouvet, Florence Pinet

### ► To cite this version:

Annie Turkieh, Henri Charrier, Emilie Dubois-Deruy, Sina Porouchani, Marion Bouvet, et al.. Non-coding RNAs in cardiac autophagy following myocardial infarction. *Oxidative Medicine and Cellular Longevity*, 2019, pp.8438650. 10.1155/2019/8438650 . hal-02262183

**HAL Id: hal-02262183**

**<https://hal.science/hal-02262183>**

Submitted on 2 Aug 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

## Review

### Non-coding RNAs in cardiac autophagy following myocardial infarction

Annie Turkieh<sup>1</sup>, Henri Charrier<sup>2</sup>, Emilie Dubois-Deruy<sup>3</sup>, Sina Porouchani<sup>4</sup>, Marion Bouvet<sup>5</sup>, and Florence Pinet<sup>6,\*</sup>

<sup>1</sup> Inserm, Université de Lille, Institut Pasteur de Lille, FHU-REMOD-VHF, U1167-RID-AGE, F-59000 Lille, France; [ani.turkieh@pasteur-lille.fr](mailto:ani.turkieh@pasteur-lille.fr)

<sup>2</sup> Inserm, Université de Lille, Institut Pasteur de Lille, FHU-REMOD-VHF, U1167-RID-AGE, F-59000 Lille, France; [henri.charrier@pasteur-lille.fr](mailto:henri.charrier@pasteur-lille.fr)

<sup>3</sup> Inserm, Université de Lille, Institut Pasteur de Lille, FHU-REMOD-VHF, U1167-RID-AGE, F-59000 Lille, France; [emilie.deruy@pasteur-lille.fr](mailto:emilie.deruy@pasteur-lille.fr)

<sup>4</sup> Inserm, Université de Lille, Institut Pasteur de Lille, FHU-REMOD-VHF, U1167-RID-AGE, F-59000 Lille, France; [sina.porouchani@gmail.com](mailto:sina.porouchani@gmail.com)

<sup>5</sup> Inserm, Université de Lille, Institut Pasteur de Lille, FHU-REMOD-VHF, U1167-RID-AGE, F-59000 Lille, France; [marion.bouvet@inserm.fr](mailto:marion.bouvet@inserm.fr)

<sup>6</sup> Inserm, Université de Lille, Institut Pasteur de Lille, FHU-REMOD-VHF, U1167-RID-AGE, F-59000 Lille, France; [florence.pinet@pasteur-lille.fr](mailto:florence.pinet@pasteur-lille.fr)

\*Correspondence: [florence.pinet@pasteur-lille.fr](mailto:florence.pinet@pasteur-lille.fr); Tel.: +33 (0) 3 20 87 72 15

## ABSTRACT

Macroautophagy is an evolutionarily conserved process of the lysosome-dependent degradation of damaged proteins and organelles and plays an important role in cellular homeostasis. Macroautophagy is upregulated after myocardial infarction (MI) and seems to be detrimental during reperfusion and protective during left ventricle remodeling. Identify new regulators of cardiac autophagy may help to maintain the activity of this process and protect the heart from MI effects. Recently, it was shown that non-coding RNAs (microRNAs and long non-coding RNAs) are involved on autophagy regulation in different cell types including cardiac cells. In this review, we summarized the role of macroautophagy in the heart following MI and we focused on the non-coding RNAs and their targeted genes reported to regulate autophagy in the heart under these pathological conditions.

## 1. INTRODUCTION

Myocardial infarction (MI) is a cardiovascular event caused by obstruction of one or more arteries supplying the heart. This area of the heart is therefore no longer supplied with oxygen and nutrients leading to the death of cardiomyocytes. Coronary reperfusion is the only recognized method to reduce the size of the infarct if it is performed within hours after MI. Despite its beneficial effect, several deleterious events such as increased oxidative stress and cell death are observed during the reperfusion process. If the infarcted zone is very extensive, there is a decrease in the contractile function of the heart. In order to compensate for this loss and maintain normal blood flow, the heart will undergo structural changes such as thinning of the infarcted zone, fibrosis, cardiomyocyte hypertrophy and left ventricle (LV) dilatation<sup>1</sup>. Left ventricle remodeling (LVR) is initially a protective mechanism but in the long term can lead to heart failure (HF)<sup>2-4</sup>. Despite current therapy, acute MI and HF, remain the leading causes of death and disability worldwide. New therapeutic strategies are

therefore required to protect the heart against the detrimental effects of acute ischemia/reperfusion (I/R) injury, in order to prevent cardiomyocyte death and reduce myocardial infarct size, preserve LV ventricle function, and prevent the onset of HF.

Macroautophagy is an important and non-selective proteolytic mechanism that regulates the homeostasis of long-lived proteins, macromolecules including lipids and cell organelles by surrounding them in a double-membrane vesicle, known as autophagosome in order to deliver them to lysosome for degradation<sup>5</sup>. It plays an essential role for maintaining heart structure and function under baseline conditions<sup>6-8</sup>. Several studies showed that macroautophagy is upregulated in heart following MI and suggested that this process may protect heart against MI effects<sup>9-11</sup>. Recently, it was shown that non-coding RNAs (microRNAs (miRNA) and long non-coding RNAs (lncRNA)) are involved on autophagy regulation in different cell types including cardiac cells<sup>12-14</sup>. In this review, we summarized the role of macroautophagy in the heart following MI and we focused on the non-coding RNAs and their targeted genes reported to regulate autophagy in the heart under pathological conditions.

## **2. MACROAUTOPHAGY MECHANISM**

Macroautophagy proceeds in several successive steps and involves different proteins as previously described<sup>5</sup>. In summary, autophagy induction is mainly regulated by the ULK (unc-51-like kinase) complex which is composed of ULK1/2, ATG13 (autophagy-related gene 13), ATG101, and FIP200 (focal adhesion kinase family interacting protein with a 200 kDa mass). Activation of the PI3K complex contributes to the vesicle nucleation, the first step of autophagosome formation. This complex is composed of Beclin-1, ATG14, VPS34 (Phosphatidylinositol 3-kinase vacuolar protein sorting 34) and VPS15. Finally, two ubiquitin-like protein conjugation systems are required to the vesicle elongation, the first to form ATG12-ATG5-ATG16L1 complex and the second to form LC3II (microtubule-associated protein 1 light chain II), the lipidated form of LC3. For this latter step, ATG4 cleaves pro-LC3 to LC3I before its conjugation to phosphatidylethanolamine by ATG7, ATG3 and ATG12-ATG5-ATG16L1 complex. Several pathways were shown to regulate autophagy by activation or inactivation of one of these ATG proteins. For example, mTOR (mammalian target of rapamycin) activation inhibited autophagy by decreasing ULK1 activity<sup>15</sup> and ATG14/VSP34-35 complex formation<sup>16</sup>. AMPK (adenosine monophosphate-activated protein kinase) positively regulated autophagy by increasing Beclin-1 phosphorylation leading to its interaction with VSP34<sup>17</sup>. However, Bcl-2 interacts with Beclin-1 for blocking its interaction with VSP34<sup>18</sup>.

### ***2. 1. Macroautophagy during ischemia/reperfusion***

The regulation of autophagy is different during ischemia and reperfusion<sup>10</sup>. During heart ischemia, nutrient and oxygen supplies to the cardiac cells decrease, inducing mitochondrial and cellular dysfunction that lead to cell death. To protect them, the cardiac cells induce autophagy via the AMPK/m-TOR pathway in order to degrade/eliminate damaged organelles and proteins and provide the substrates necessary for their survival. During reperfusion, there is an increase of reactive oxygen species (ROS) production inducing a strong expression of Beclin-1 which on one hand, promotes the formation of autophagosomes and on the other hand, inhibits the expression of genes involved in the fusion of autophagosomes with lysosomes<sup>19</sup>. In addition, ROS inhibit the expression of LAMP-2, a protein involved in the fusion of autophagosomes with lysosomes. Autophagy is then induced excessively during reperfusion but is inactive. Blocking the degradation of the contents of autophagosomes promotes oxidative stress, decreases mitochondrial permeability and causes cell death. Partial inhibition of Beclin-1 expression (heterozygous mice) has been shown

to protect against apoptosis induced during reperfusion while its total deletion is deleterious<sup>10</sup>. These data showed that autophagy is a protective mechanism during ischemia but its excessive induction during reperfusion is deleterious.

## **2.2. Macroautophagy during LVR in post-MI**

The activity of autophagy and its role in LVR post-MI have been studied in murine models with permanent ligation of the left coronary artery. Autophagy is induced in non-infarcted area of the heart during the sub-acute (1 week) and chronic (3 weeks) stages after MI<sup>11</sup>. Inhibition of autophagy by Bafilomycin (a pharmacological agent that blocks the fusion of the autophagosome with the lysosome) promoted LVR and worsens cardiac dysfunction. In contrast, administration of Trehalose (a non-naturally reduced disaccharide) in mice after ligation, activated autophagy, reduced LVR, and improved cardiac function at 4 weeks post-MI<sup>20</sup>. However, this protective effect of trehalose on the heart was not observed in mice invalidated for the Beclin-1 gene, but an increase in the activity of mTOR was observed in the non-infarcted area of the heart. It has been shown that the inhibition of mTOR activity induced autophagy leading to a decrease of LVR and an improvement in cardiac function in post-MI<sup>21</sup>. All these data showed a protective role of autophagy in later stages in post-MI but its activity remained insufficient to prevent LVR and cardiac dysfunction.

## **3. Macroautophagy regulation by non-coding RNAs during and following MI**

About 99% of the human genome do not encode proteins, but are transcriptionally highly active and give rise to a broad spectrum of non-coding RNAs (ncRNAs) with regulatory and structural functions. Based on the size criteria of 200 nucleotides (nt), ncRNAs are divided into long (>200 nt) and short ncRNAs (<200 nt).

The ncRNAs are modulated in some cardiovascular diseases including MI<sup>22,23</sup>. The significant changes in their expression pattern upon MI highlighted their contribution in regulation of pathogenesis of MI. Furthermore, it was shown that ncRNAs could regulate autophagy in some cardiac disorders including MI, hypertrophy and HF<sup>12-14</sup>. In this part, we summarized the non-coding RNAs which have been reported to regulate cardiac autophagy during and following MI and highlighted their specific autophagic targets and their importance as new therapeutic targets to protect heart against I/R injury and prevent cardiac remodeling and dysfunction (**Figure 1**).

### **3.1. Macroautophagy regulation by mRNAs**

MiRNAs are defined as single-stranded non-coding RNAs around 22 nucleotides and are highly conserved between species<sup>22</sup>. Once synthesized and matured through several steps, these miRNAs bind to the complementary 3'UTR of their target mRNA and either degrade or silence them. A near perfect match between the seed region of the miRNA (8 nucleotides at its 5' UTR end) and its target leads to complete degradation of mRNA, while a partial complementary results in the suppression of the gene expression. MiRNAs may have one or multiple mRNA targets and are involved in the regulation of numerous biological processes in the heart including autophagy.

#### **3.1.1. Antiautophagic-miRNAs with protective effects**

Several miRNAs were modulated during I/R and seems to have a protective effect by decreasing excessive autophagy induced-cell apoptosis by targeting one of the ATG genes. MiR-188-3p levels are reduced in cardiomyocytes treated with anoxia/reoxygenation and in MI-mice. Overexpression of miR-188-3p in MI-mice attenuated autophagy by targeting autophagy mediator Atg7 and decreased the infarcted area size<sup>24</sup>. It was shown that miR-638 suppressed the expression of Atg5 by targeting its 3'UTR region. It is down-regulated in human cardiomyocytes after hypoxia/reoxygenation (H/R)

and its overexpression improve the viability of these cells. However, enforced expression of Atg5 reversed the effect of miR-638 on autophagy and cell apoptosis suggesting that miR-638 attenuated the effects of H/R treatment by regulating ATG5-mediated autophagy in human cardiomyocytes<sup>25</sup>. Also, overexpression of miR-129-5p in H9c2 cells treated by hydrogen peroxide inhibited autophagy by targeting Atg14 gene and activating PI3K/AKT/mTOR pathway resulting in decreased cell apoptosis<sup>26</sup>.

Other miRNAs play their protective effect by regulating one of the pathways involved on autophagy regulation. The levels of miR-223 are significantly upregulated in the heart of post-MI HF-rats and in hypoxia-treated neonatal rat cardiomyocytes (NRCMs) and H9c2 cells. The increased miR-223 levels protect NRCMs and H9c2 cells from hypoxia-induced apoptosis whereas decreasing miR-223 expression had contrasting effects. This protective effect of miR-223 is explained by the decrease of its target gene expression PARP-1 (poly(ADP-ribose) polymerase 1) resulting in inhibition of excessive autophagy via the Akt/mTOR pathway<sup>27</sup>. However, miR-204 expression is decreased in the heart of rat upon I/R injury associated with increased autophagy as observed by the increased LC3II levels<sup>28</sup>. Also, it was shown that transfection of miR-204 in H9c2 cells attenuated cell apoptosis induced by H/R treatment. The protective effect of miR-204 is explained by targeting SIRT1-mediated autophagy<sup>29</sup>. The expression of miR-34a is also decreased during I/R and overexpression of this miR decreased TNF $\alpha$  expression resulting in reduced autophagy and apoptosis levels on NRCMs after H/R<sup>30</sup>. Lower miR-29b-3p levels were found in HF patients and in hypoxia-stimulated H9c2 cells. The overexpression of miR-29b-3p inhibited autophagy and apoptosis induced in hypoxic-induced H9c2 cells through targeting SPARC and inhibiting TGF $\beta$ -1/Smad3 pathway<sup>31</sup>.

### 3.1.2. Antiautophagic-miRNAs with deleterious effects

Some miRNAs contribute to ischemic/reperfusion injury by inhibiting the autophagy process. The miR-497 is dramatically down-regulated in infarcted heart and in hypoxic cardiomyocytes and its overexpression in murine MI model increased the infarcted size. It was shown that miR-497 inhibited autophagy by targeting LC3B gene and induced cell apoptosis by targeting Bcl-2 gene suggesting that decreasing miR-497 levels is a protective mechanism of the heart in response to MI<sup>32</sup>. The expression of miR-30e was also decreased after myocardial I/R. Its silencing in H9c2 cells increased autophagy and attenuated oxidative stress and cell apoptosis, that are reversed by treating the cells with 3-methyladenine, an inhibitor of macroautophagy. These results suggest that decreasing the miR-30e levels protected the heart against I/R injury by autophagy induction<sup>33</sup>.

### 3.1.3. Proautophagic-miRNAs with protective effects

Higashi *et al.*<sup>34</sup> showed that 30 min of coronary occlusion followed by 2 days of reperfusion caused a significant decrease in the rabbit cardiac tissue expression of miR-145 in the border and infarcted areas of the myocardium compared to the remote non-infarcted area. Injection of liposomes containing miR-145 after the beginning of reperfusion reduced the infarcted area size, improved the LV function and remodeling, these beneficial effects were abolished by chloroquine treatment. Further study showed that miR-145 promoted autophagy in cardiomyocyte by directly targeting FRS2 (fibroblast growth factor receptor substrate 2) mRNA resulting in the acceleration of the transition of LC3I to LC3II, an important step of autophagosome maturation<sup>34</sup>. The protective effect of miR-145 is also observed in H9c2 cells after H/R. In this study, the authors demonstrated that miR-145 inhibited H/R-induced apoptosis by promoting autophagy via Akt3/mTOR signaling pathway<sup>35</sup>. The miR-99a was shown to be down-regulated in infarcted heart and in neonatal mice ventricle myocytes exposed to hypoxia. Lentivirus-mediated overexpression of miR-99a in infarcted

heart inhibited cardiac remodeling and improved heart function at 1 and 4 weeks after its administration. It was shown that miR-99a decreased mTOR protein levels without any effect on its mRNA levels suggesting that miR-99a regulated mTOR expression at a post-transcriptional level. Consequently, the autophagy induced was associated with a decrease of cell apoptosis. This study demonstrated that overexpression of miR-99a improved post-MI cardiac function by up-regulating autophagy via targeting mTOR pathways, inhibiting apoptosis and attenuating pathological remodeling<sup>36</sup>. The miR-144 levels were reduced in the heart of MI mice with permanent left anterior descending artery (LAD) ligation. The miR-144 k/o mice showed a worse HF phenotype with ventricular dilatation and impaired contractility after LAD ligation. However, miR-144 administration decreased myocardial infarcted size and improved post-MI remodeling. Further study allowed authors to conclude that miR-144 increased autophagy and decreased fibrosis and apoptosis by targeting mTOR<sup>37</sup>.

### **3.2. Macroautophagy regulation by lncRNAs**

lncRNAs are non-coding RNAs longer than 200 nucleotides that regulate both gene expression and protein translation<sup>22</sup>. Nuclear localized lncRNAs can regulate gene expression at both the epigenetic and transcriptional levels. Cytosol-based lncRNAs can modify protein translation by blocking, stabilizing/destabilizing, or sponging miRNAs. The lncRNAs are involved in the regulation of numerous biological processes including autophagy in cardiac and non-cardiac cells.

#### **3.2.1. Antiautophagic- lncRNA with protective effects**

Liu *et al.*<sup>38</sup> showed that the expression of lncRNA CAIF (cardiac autophagy inhibitory factor) was significantly decreased in a mice model of I/R injury and in cardiomyocytes treated with H<sub>2</sub>O<sub>2</sub>. Conversely, overexpression of CAIF inhibited autophagy inducing cardiomyocyte cell death and cardiac dysfunction caused by I/R. In this study, the authors demonstrate that CAIF directly binds to p53 protein and blocks its interaction with the myocardin promoter. Myocardin, a smooth muscle and cardiac muscle-specific transcriptional activator, is upregulated after I/R and H<sub>2</sub>O<sub>2</sub> treatment, and is involved on autophagy regulation in cardiomyocytes by increasing Beclin 1 expression. These data suggest CAIF-P53-myocardin pathway as a novel regulator of autophagy in cardiomyocytes and as a potential therapeutic target in order to inhibit excessive autophagy and improve cardiac function after I/R<sup>38</sup>.

#### **3.2.2. Proautophagic- lncRNA with protective effects**

On the other hand, it was shown that the lncRNA H19 expression was decreased in a mice model of acute MI and that its overexpression decreased infarcted size and improved cardiac function associated with autophagy upregulation; however, the mechanisms by which autophagy is regulated by H19 is still unknown. These results suggest that H19 protects the heart from MI by increasing cardiac autophagy<sup>39</sup>.

#### **3.2.3. Proautophagic- lncRNAs with deleterious effects**

Some lncRNAs are upregulated after I/H and enhanced autophagy-target genes expression by inhibiting miRNAs expression. Yin *et al.*<sup>40</sup> showed that the lncRNA Galnt1 (GATA1 activated lncRNA) is upregulated in neonatal mice cardiomyocytes in response to anorexia/reoxygenation; however, miR-338 expression is downregulated. Overexpression of miR-338 directly decreased the formation of autophagic vesicles and induced cell death after anorexia/reoxygenation treatment without any effect on control cells. The antiautophagic effect of miR-338 is explained by its direct targeting of the

autophagic mediator Atg5. It was shown that Galnt1 directly bound to miR-338 and decrease its expression. Consequently, Atg 5 expression is increased resulting in excessive cardiac autophagy and cell death<sup>40</sup>. Also, the lncRNA APF (Autophagy promoting factor) enhances cardiac autophagy and cell death by inhibiting miR-188-3p expression resulting in the increase of its target gene expression, Atg7<sup>24</sup>. Furthermore, the lncRNA AK088388 is upregulated during reoxygenation in mouse cardiac myocytes associated with the decreased miR-30a expression. Overexpression of miR-30a decreased the expression of its target gene Beclin-1 resulting in inhibition of autophagy induction and decreased cell death. The co-overexpression of lncRNA AK088388 inhibited the protective effect of miR-30a. However, the mutation of the miR30-a binding site in AK088388 failed to block the effect of this miRNA on autophagy and cell survival. These results suggest that the lncRNA AK088388 regulates autophagy through miR-30a/Beclin-1 pathway to affect cardiomyocyte injury<sup>41</sup>. The lncRNA HRIM (hypoxia/reoxygenation injury-related factor in myocyte) was upregulated after H/R in H9c2 cells. HRIM silencing prevented death of cells by suppressing the autophagic activity in H/R-treated cells. However, the target genes of this lncRNA and the detailed mechanism of its autophagic effect need to be elucidated<sup>42</sup>. Other lncRNAs were highly expressed in diabetic murine heart and contributed to I/R injury by regulating autophagy. It was shown that Neat-1 (Nuclear-enriched abundant transcript 1) and AK139328 seemed to induce autophagy by upregulating Foxo1 expression and decreasing miR-204-3p levels, respectively<sup>43,44</sup>.

The lncRNA MALAT1 (metastasis-associated lung adenocarcinoma transcript 1) is expressed at high levels in patients with acute MI<sup>45</sup> and is closely associated with the pathogenesis of myocardial I/R injury<sup>46,47</sup>. It was shown, on one hand, that MALAT1 contained binding site for miR-204<sup>48</sup> and, in the other hand, that miR-204 protected the cardiomyocytes against I/R injury via inhibiting autophagic cell death<sup>28</sup>. Also, MALAT1 targeted miR-558 to enhance ULK1-mediated autophagy in isoproterenol treated-cardiomyocytes<sup>49</sup>. It will be important to know if lncRNA MALAT1 increased cardiomyocyte autophagy and myocardial injury during I/R by negatively regulating miR-204 or miR-558 expression.

#### **4. CONCLUSION**

Despite current therapies, acute MI and HF which often follows, remain the leading causes of death and disability worldwide. New therapeutic strategies are therefore required to protect the heart against the detrimental effects of acute ischemia/reperfusion injury. Inhibition of macroautophagy during reperfusion prevented cardiomyocyte death and reduced myocardial infarct size, however its induction during LVR preserved LV function and prevent the onset of HF. The most pharmacological agents used up to date for regulating macroautophagy are not specific and may interfere with other cellular processes, so it will be necessary to identify new therapeutic approaches to regulate autophagy. Several non-coding RNAs were shown to be modulated during I/R and involved on cardiac autophagy regulation. The tissue specific expression of some non-coding RNAs and their easy manipulation show their potential as novel targets for clinical developments to treat autophagy related-diseases. Identification of specific cardiac non-coding RNAs that regulate autophagy could be a good opportunity to protect heart from MI injury without affecting the autophagy activity in other organ.

#### **5. ACKNOWLEDGMENTS**

AT received a fellowship from ANR [ERA-NET Co-fund action ERA-CVD JTC2016 N° 680969] and EDD from "Fondation Lefoulon Delalande". FP is a MC substitute member of COST Action Transautophagy (CA15138) and is a MC member of COST action cardioRNA (CA1229).

## 6. CONFLICTS OF INTEREST


The authors declare no conflict of interest

## 7. REFERENCES

1. Pfeffer MA, Braunwald E. Ventricular remodeling after myocardial infarction. Experimental observations and clinical implications. *Circulation*. 1990;81:1161–72.
2. St John Sutton M, Pfeffer MA, Plappert T, Rouleau JL, Moyé LA, Dagenais GR, Lamas GA, Klein M, Sussex B, Goldman S. Quantitative two-dimensional echocardiographic measurements are major predictors of adverse cardiovascular events after acute myocardial infarction. The protective effects of captopril. *Circulation*. 1994;89:68–75.
3. de Groote P, Fertin M, Duva Pentiah A, Goéminne C, Lamblin N, Bauters C. Long-term functional and clinical follow-up of patients with heart failure with recovered left ventricular ejection fraction after  $\beta$ -blocker therapy. *Circ Heart Fail*. 2014;7:434–9.
4. Bauters C, Dubois E, Porouchani S, Saloux E, Fertin M, de Groote P, Lamblin N, Pinet F. Long-term prognostic impact of left ventricular remodeling after a first myocardial infarction in modern clinical practice. *PLoS One*. 2017;12:e0188884.
5. Ohsumi Y. Historical landmarks of autophagy research. *Cell Res*. 2014;24:9–23.
6. Nakai A, Yamaguchi O, Takeda T, Higuchi Y, Hikoso S, Taniike M, Omiya S, Mizote I, Matsumura Y, Asahi M, Nishida K, Hori M, Mizushima N, Otsu K. The role of autophagy in cardiomyocytes in the basal state and in response to hemodynamic stress. *Nat Med*. 2007;13:619–24.
7. Gan B, Peng X, Nagy T, Alcaraz A, Gu H, Guan J-L. Role of FIP200 in cardiac and liver development and its regulation of TNF $\alpha$  and TSC-mTOR signaling pathways. *J Cell Biol*. 2006;175:121–33.
8. Kaizuka T, Mizushima N. Atg13 Is Essential for Autophagy and Cardiac Development in Mice. *Mol Cell Biol*. 2016;36:585–95.
9. Yan L, Vatner DE, Kim S-J, Ge H, Masurekar M, Massover WH, Yang G, Matsui Y, Sadoshima J, Vatner SF. Autophagy in chronically ischemic myocardium. *Proc Natl Acad Sci*. 2005;102:13807–13812.
10. Matsui Y, Takagi H, Qu X, Abdellatif M, Sakoda H, Asano T, Levine B, Sadoshima J. Distinct roles of autophagy in the heart during ischemia and reperfusion: roles of AMP-activated protein kinase and Beclin 1 in mediating autophagy. *Circ Res*. 2007;100:914–22.
11. Kanamori H, Takemura G, Goto K, Maruyama R, Tsujimoto A, Ogino A, Takeyama T, Kawaguchi T, Watanabe T, Fujiwara T, Fujiwara H, Seishima M, Minatoguchi S. The role of autophagy emerging in postinfarction cardiac remodeling. *Cardiovasc Res*. 2011;91:330–339.
12. Gupta SK, Thum T. Non-coding RNAs as orchestrators of autophagic processes. *J Mol Cell Cardiol*. 2016;95:26–30.
13. Yang L, Wang H, Shen Q, Feng L, Jin H. Long non-coding RNAs involved in autophagy regulation. *Cell Death Dis*. 2017;8:e3073.
14. Sun T, Li M-Y, Li P-F, Cao J-M. MicroRNAs in Cardiac Autophagy: Small Molecules and Big Role. *Cells*. 2018;7:104.
15. Kim J, Kundu M, Viollet B, Guan K-L. AMPK and mTOR regulate autophagy through direct phosphorylation of Ulk1. *Nat Cell Biol*. 2011;13:132–41.
16. Yuan H-X, Russell RC, Guan K-L. Regulation of PIK3C3/VPS34 complexes by MTOR in nutrient stress-induced autophagy. *Autophagy*. 2013;9:1983–95.
17. Kim J, Kim YC, Fang C, Russell RC, Kim JH, Fan W, Liu R, Zhong Q, Guan K-L. Differential regulation of distinct Vps34 complexes by AMPK in nutrient stress and autophagy. *Cell*. 2013;152:290–303.
18. Kang R, Zeh HJ, Lotze MT, Tang D. The Beclin 1 network regulates autophagy and apoptosis. *Cell Death Differ*. 2011;18:571–580.

19. Ma X, Liu H, Foyil SR, Godar RJ, Weinheimer CJ, Hill JA, Diwan A. Impaired autophagosome clearance contributes to cardiomyocyte death in ischemia/reperfusion injury. *Circulation*. 2012;125:3170–81.
20. Sciarretta S, Yee D, Nagarajan N, Bianchi F, Saito T, Valenti V, Tong M, Del Re DP, Vecchione C, Schirone L, Forte M, Rubattu S, Shirakabe A, Boppana VS, Volpe M, Frati G, Zhai P, Sadoshima J. Trehalose-Induced Activation of Autophagy Improves Cardiac Remodeling After Myocardial Infarction. *J Am Coll Cardiol*. 2018;71:1999–2010.
21. Buss SJ, Muenz S, Riffel JH, Malekar P, Hagenmueller M, Weiss CS, Bea F, Bekeredjian R, Schinke-Braun M, Izumo S, Katus HA, Hardt SE. Beneficial Effects of Mammalian Target of Rapamycin Inhibition on Left Ventricular Remodeling After Myocardial Infarction. *J Am Coll Cardiol*. 2009;54:2435–2446.
22. Das A, Samidurai A, Salloum FN. Deciphering Non-coding RNAs in Cardiovascular Health and Disease. *Front Cardiovasc Med*. 2018;5:73.
23. Poller W, Dimmeler S, Heymans S, Zeller T, Haas J, Karakas M, Leistner D-M, Jakob P, Nakagawa S, Blankenberg S, Engelhardt S, Thum T, Weber C, Meder B, Hajjar R, Landmesser U. Non-coding RNAs in cardiovascular diseases: diagnostic and therapeutic perspectives. *Eur Heart J*. 2018;39:2704–2716.
24. Wang K, Liu C-Y, Zhou L-Y, Wang J-X, Wang M, Zhao B, Zhao W-K, Xu S-J, Fan L-H, Zhang X-J, Feng C, Wang C-Q, Zhao Y-F, Li P-F. APF lncRNA regulates autophagy and myocardial infarction by targeting miR-188-3p. *Nat Commun*. 2015;6:6779.
25. Zhao P, Zhang B-L, Liu K, Qin B, Li Z-H. Overexpression of miR-638 attenuated the effects of hypoxia/reoxygenation treatment on cell viability, cell apoptosis and autophagy by targeting ATG5 in the human cardiomyocytes. *Eur Rev Med Pharmacol Sci*. 2018;22:8462–8471.
26. Zhang H, Zhang X, Zhang J. MiR-129-5p inhibits autophagy and apoptosis of H9c2 cells induced by hydrogen peroxide via the PI3K/AKT/mTOR signaling pathway by targeting ATG14. *Biochem Biophys Res Commun*. 2018;506:272–277.
27. Liu X, Deng Y, Xu Y, Jin W, Li H. MicroRNA-223 protects neonatal rat cardiomyocytes and H9c2 cells from hypoxia-induced apoptosis and excessive autophagy via the Akt/mTOR pathway by targeting PARP-1. *J Mol Cell Cardiol*. 2018;118:133–146.
28. Xiao J, Zhu X, He B, Zhang Y, Kang B, Wang Z, Ni X. MiR-204 regulates cardiomyocyte autophagy induced by ischemia-reperfusion through LC3-II. *J Biomed Sci*. 2011;18:35.
29. Qiu R, Li W, Liu Y. MicroRNA-204 protects H9C2 cells against hypoxia/reoxygenation-induced injury through regulating SIRT1-mediated autophagy. *Biomed Pharmacother*. 2018;100:15–19.
30. Shao H, Yang L, Wang L, Tang B, Wang J, Li Q. MicroRNA-34a protects myocardial cells against ischemia-reperfusion injury through inhibiting autophagy via regulating TNF $\alpha$  expression. *Biochem Cell Biol*. 2018;96:349–354.
31. Zhou S, Lei D, Bu F, Han H, Zhao S, Wang Y. MicroRNA-29b-3p Targets SPARC Gene to Protect Cardiocytes against Autophagy and Apoptosis in Hypoxic-Induced H9c2 Cells. *J Cardiovasc Transl Res*. 2018. doi:10.1007/s12265-018-9858-1.
32. Li X, Zeng Z, Li Q, Xu Q, Xie J, Hao H, Luo G, Liao W, Bin J, Huang X, Liao Y. Inhibition of microRNA-497 ameliorates anoxia/reoxygenation injury in cardiomyocytes by suppressing cell apoptosis and enhancing autophagy. *Oncotarget*. 2015;6:18829–44.
33. Zheng J, Li J, Kou B, Yi Q, Shi T. MicroRNA-30e protects the heart against ischemia and reperfusion injury through autophagy and the Notch1/Hes1/Akt signaling pathway. *Int J Mol Med*. 2018;41:3221–3230.
34. Higashi K, Yamada Y, Minatoguchi S, Baba S, Iwasa M, Kanamori H, Kawasaki M, Nishigaki K, Takemura G, Kumazaki M, Akao Y, Minatoguchi S. MicroRNA-145 repairs infarcted myocardium by accelerating cardiomyocyte autophagy. *Am J Physiol Heart Circ Physiol*. 2015;309:H1813–26.
35. Yan L, Guo N, Cao Y, Zeng S, Wang J, Lv F, Wang Y, Cao X. miRNA-145 inhibits myocardial infarction-induced apoptosis through autophagy via Akt3/mTOR signaling pathway in vitro and in vivo. *Int J Mol Med*. 2018;42:1537–1547.

36. Li Q, Xie J, Li R, Shi J, Sun J, Gu R, Ding L, Wang L, Xu B. Overexpression of microRNA-99a attenuates heart remodelling and improves cardiac performance after myocardial infarction. *J Cell Mol Med*. 2014;18:919–28.
37. Li J, Cai SX, He Q, Zhang H, Friedberg D, Wang F, Redington AN. Intravenous miR-144 reduces left ventricular remodeling after myocardial infarction. *Basic Res Cardiol*. 2018;113:36.
38. Liu C-Y, Zhang Y-H, Li R-B, Zhou L-Y, An T, Zhang R-C, Zhai M, Huang Y, Yan K-W, Dong Y-H, Ponnusamy M, Shan C, Xu S, Wang Q, Zhang Y-H, Zhang J, Wang K. LncRNA CAIF inhibits autophagy and attenuates myocardial infarction by blocking p53-mediated myocardin transcription. *Nat Commun*. 2018;9:29.
39. Zhou M, Zou Y-G, Xue Y-Z, Wang X-H, Gao H, Dong H-W, Zhang Q. Long non-coding RNA H19 protects acute myocardial infarction through activating autophagy in mice. *Eur Rev Med Pharmacol Sci*. 2018;22:5647–5651.
40. Yin G, Yang X, Li Q, Guo Z. GATA1 activated lncRNA (Galont) promotes anoxia/reoxygenation-induced autophagy and cell death in cardiomyocytes by sponging miR-338. *J Cell Biochem*. 2018;119:4161–4169.
41. Wang J, Bie Z, Sun C. Long noncoding RNA AK088388 regulates autophagy through miR-30a to affect cardiomyocyte injury. *J Cell Biochem*. 2019. doi:10.1002/jcb.28300.
42. Huang Z, Ye B, Wang Z, Han J, Lin L, Shan P, Cai X, Huang W. Inhibition of LncRNA-HRIM Increases Cell Viability by Regulating Autophagy Levels During Hypoxia/Reoxygenation in Myocytes. *Cell Physiol Biochem*. 2018;46:1341–1351.
43. Ma M, Hui J, Zhang Q, Zhu Y, He Y, Liu X. Long non-coding RNA nuclear-enriched abundant transcript 1 inhibition blunts myocardial ischemia reperfusion injury via autophagic flux arrest and apoptosis in streptozotocin-induced diabetic rats. *Atherosclerosis*. 2018;277:113–122.
44. Yu S, Dong B, Fang Z, Hu X, Tang L, Zhou S. Knockdown of lncRNA AK139328 alleviates myocardial ischaemia/reperfusion injury in diabetic mice via modulating miR-204-3p and inhibiting autophagy. *J Cell Mol Med*. 2018;22:4886–4898.
45. Vausort M, Wagner DR, Devaux Y. Long noncoding RNAs in patients with acute myocardial infarction. *Circ Res*. 2014;115:668–77.
46. Zhao Z-H, Hao W, Meng Q-T, Du X-B, Lei S-Q, Xia Z-Y. Long non-coding RNA MALAT1 functions as a mediator in cardioprotective effects of fentanyl in myocardial ischemia-reperfusion injury. *Cell Biol Int*. 2017;41:62–70.
47. Hu H, Wu J, Li D, Zhou J, Yu H, Ma L. Knockdown of lncRNA MALAT1 attenuates acute myocardial infarction through miR-320-Pten axis. *Biomed Pharmacother*. 2018;106:738–746.
48. Li J, Wang J, Chen Y, Li S, Jin M, Wang H, Chen Z, Yu W. LncRNA MALAT1 exerts oncogenic functions in lung adenocarcinoma by targeting miR-204. *Am J Cancer Res*. 2016;6:1099–107.
49. Guo X, Wu X, Han Y, Tian E, Cheng J. LncRNA MALAT1 protects cardiomyocytes from isoproterenol-induced apoptosis through sponging miR-558 to enhance ULK1-mediated protective autophagy. *J Cell Physiol*. 2018;:jcp.27925.



**Figure 1:** Outline summarizing the non-coding RNAs regulating cardiac autophagy, their targets and function. Green and red arrows indicate activation and inhibition, respectively