



Annex to the proposition denitac.pdf

Denise Vella-Chemla

► **To cite this version:**

| Denise Vella-Chemla. Annex to the proposition denitac.pdf. 2019. hal-02263283

HAL Id: hal-02263283

<https://hal.archives-ouvertes.fr/hal-02263283>

Submitted on 3 Aug 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

One looks for decomposing an even number n as a sum of 2 prime numbers $p_1 + p_2$.

We can't make reference to $\zeta(-1)$ as we made it in [1]. We can, however, to obtain a minoration for the number of Goldbach decompositions of n , use the cardinal $|\mathcal{P}_{\frac{n}{2}}|$ of the set of prime numbers that are lesser than or equal to $\frac{n}{2}$ and multiply it by the product $\prod_{p \leq \sqrt{n}} \left(1 - \frac{1}{p}\right)$ that counts how many chances the prime number p_1 has not to share its division rest with n in each division by a module p lesser than \sqrt{n} (the fact that p_1 doesn't share its rest in a division by p with n involves that the complementary of p_1 to n (called p_2) is prime too).

The minoration¹ of $\pi(x)$ (the number of prime numbers lesser than x) by $\frac{x}{\log x}$ can be found in [2], page 69, for $x \geq 17$ (Corollary 1, (3.5), of Theorem 2, whose demonstration can be found at paragraph 7 of [2]).

As a consequence, one has $|\mathcal{P}_{\frac{n}{2}}| > \frac{\frac{n}{2}}{\log(\frac{n}{2})}$.

The minoration of $\prod_{p \leq \sqrt{n}} \left(1 - \frac{1}{p}\right)$ is also provided by [2], page 70 (it's the corollary (3.27) of Theorem 7 whose demonstration is provided at paragraph 8 of [2], with γ the Euler-Mascheroni constant).

$$(3.27) \quad \frac{e^{-\gamma}}{\log x} \left(1 - \frac{1}{\log^2 x}\right) < \prod_{p \leq x} \left(1 - \frac{1}{p}\right) \quad \text{for } 1 < x.$$

Multiplying those expressions together, one obtains that the number of Goldbach decomposants of n must be greater than :

$$\frac{n/2}{\log(n/2)} \frac{e^{-\gamma}}{\log \sqrt{n}} \left(1 - \frac{1}{\log^2 \sqrt{n}}\right)$$

which is strictly greater than 1 for $n \geq 24$.

Bibliography

[1] <http://denisevellachemla.eu/denitac.pdf>.

[2] J. B. Rosser et L. Schoenfeld, *Approximate formulas for some functions of prime numbers*, dedicated to Hans Rademacher for his seventieth birthday, Illinois J. Math., Volume 6, Issue 1 (1962), 64-94.

1. This minoration must be distinguished from the Prime Number Theorem, independently proved by Hadamard and by de la Vallée-Poussin, and that provides an asymptotic tendency for $\pi(x)$.