

Tsallis and Kaniadakis Entropic Measures in Stellar Polytropes

Amelia Carolina Sparavigna

► To cite this version:

Amelia Carolina Sparavigna. Tsallis and Kaniadakis Entropic Measures in Stellar Polytropes. Philica, 2015, 10.5281/zenodo.3366270 . hal-02265910

HAL Id: hal-02265910 https://hal.science/hal-02265910

Submitted on 12 Aug 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Tsallis and Kaniadakis Entropic Measures in Stellar Polytropes

Amelia Carolina Sparavigna

(Department of Applied Science and Technology, Politecnico di Torino)

Published in astro.philica.com

Abstract

Polytropes are self-gravitating fluid spheres used in astrophysics as crude approximation of more realistic stellar models. They possess equations that have scale parameters linked to mass, energy and entropy. Since Boltzmann distribution yields unphysical results, the use of generalized entropies, such as Tsallis and Kaniadakis entropies, had been proposed. Here we discuss how these entropies are related in polytrope solutions.

Keywords: Entropy, Generalized Entropies.

1. Introduction

In astrophysics, a polytrope refers to a solution of the Lane–Emden equation. This is an equation which gives the pressure as a function of density [1]. These solutions are modelling self-gravitating fluid spheres that are called "polytropes" too, which are objects used as crude approximation to more realistic stellar models [2]. The solution of the Lane-Emden equation, a dimensionless form of Poisson's equation for the gravitational potential, depends on a parameter which is the polytropic index *n*. It is written as $P=K\rho^{(n+1)/n}$, where P is pressure, ρ is density and K a constant. If stellar structure is approximated with a polytrope having a given index, then two scaling parameters are needed to express the structure in physical units [3]. The two parameters that we can use are a constant which is related to entropy and the stellar mass. Since Boltzmann distribution yields unphysical results, the Boltzmann entropy had been substituted by a generalized entropy, the Tsallis entropy [4]. Another generalized entropy, the Kaniadakis entropy, had been recently proposed too, in [5]. Here we discuss how these two entropies are related in polytrope solutions, and that the result given in [5] can be easily obtained from [4].

2. The entropies

Well-known is the entropy proposed by Claude Shannon in 1948 [6].

He defined the entropy H of a discrete random variable X, as the expected value of the information

content: $H(X) = -\Sigma_i p_i \log_b p_i$. The probability of i-event is p_i and b is the base of the used logarithm. However, several entropies exist which are generalizing Shannon entropy. Among them we have Tsallis and Kaniadakis entropies [7,8], which are defined, with a corresponding choice of measurement units equal to 1, as follow:

(1) Tsallis :
$$T = T_q = \frac{1}{q-1} \left(1 - \sum_i p_i^q \right)$$

(2) Kaniadakis (κ - entropy) : $K_{\kappa} = -\sum_i \frac{p_i^{1+\kappa} - p_i^{1-\kappa}}{2\kappa}$

In (1) and (2) we have the entropic indies q and κ . For its generalized additivity, the Kaniadakis entropy requires another function, defined as follow:

$$\Im = \sum_{i} (p_i^{1+\kappa} + p_i^{1-\kappa})/2$$

A detailed discussion of the generalized additivity of Tsallis and κ-entropy is given in [9]. Tsallis and Kaniadakis entropies are linked:

(3)
$$K_{\kappa} = \frac{T_{1+\kappa} + T_{1-\kappa}}{2},$$

where $T(q = 1+\kappa) = -\frac{1}{\kappa} \sum_{i} p_{i}^{1+\kappa} + \frac{1}{\kappa}; \quad T(q = 1-\kappa) = \frac{1}{\kappa} \sum_{i} p_{i}^{1-\kappa} - \frac{1}{\kappa}$

Eq.(3) is a simpler form of an expression given in [10,11]. However, besides this relation, because of the generalized additivity possessed by the Kaniadakis entropy, we need also another relation:

(4)
$$\Im_{\kappa} = \frac{\kappa}{2} \left(-T_{1+\kappa} + T_{1-\kappa} + \frac{2}{\kappa} \right)$$

In (3) and (4), we have Kaniadakis functions expressed by Tsallis entropy. As shown in [12], we can also write T expressed by means of Kaniadakis functions:

$$2K + \frac{2}{\kappa}\Im = T_{1+\kappa} + T_{1-\kappa} + \left(-T_{1+\kappa} + T_{1-\kappa} + \frac{2}{\kappa}\right)$$

And then: (5) $K_{\kappa} + \frac{1}{\kappa}\Im_{\kappa} = T_{1-\kappa} + \frac{1}{\kappa}$

Let us have: $\kappa=1-q$. From (5) we have immediately the relation between Tsallis and Kaniadakis functions:

(6)
$$T_q = K_{1-q} + \frac{\Im_{1-q} - 1}{(1-q)}$$

3. With polytropes

The relation (6) between Tsallis and Kaniadakis entropies can be useful in several problems. Here we consider its use in polytropes. In the previous equations, we have p_i denoting the probability distribution. In Ref.4, it is used letter f for probability. From now on, we will use this notation. In [4], the distribution from Tsallis entropy is:

(7)
$$C_q(f) = \frac{f(1-f^{q-1})}{q-1}$$

After Eq.6, we can write Eq.7 in the following manner:

$$(8) \quad C_{q}(f) = \frac{f - f^{q}}{q - 1} = \frac{1}{2(q - 1)}(f^{2-q} - f^{q}) - \frac{1}{2(q - 1)}(f^{2-q} + f^{q}) + \frac{f}{q - 1}$$

Of course, (7) and (8) are the same equation. As a consequence, Kaniadakis distributions are linked to Tsallis distribution by:

(9)
$$t_q(f) = C_q(f) = \frac{(f - f^q)}{q - 1} = k_{1-q} - g_{1-q} + \frac{f}{q - 1}$$

where $K_{1-q} = \sum_i k_{1-q}$, $\Im_{1-q} = \sum_i g_{1-q}$, $1 = \sum_i f$

From [4], a relation exists between polytrope index and entropic Tsallis index:

$$(10) \quad n = \frac{3}{2} + \frac{1}{q-1}$$

As a special case, for $q \rightarrow 1$, we find the isothermal situation. To have Eq.6, as shown in [12], we need $\kappa=1-q$ or $\kappa=q-1$. Then, from (10), considering that we have for the Kaniadakis index, $-1 < \kappa < 1$:

(11)
$$n = \frac{3}{2} + \frac{1}{\kappa}$$

And in fact, (11) is the relation that we find in [5]. Using then the relation between Tsallis and Kaniadakis entropies and distributions we can easily finds results concerning several applications. Polytropes are an example of such a possible approach.

References

[1] Horedt, G.P. (2004). Polytropes. Applications in Astrophysics and Related Fields, Kluwer. ISBN 1-4020-2350-2

[2] Chandrasekhar, S. (1958). An Introduction to the Study of Stellar Structure, Dover.

[3] Knapp, J. (2015). Polytropes, retrieved 9 November 2015 from http://www.astro.princeton.edu/~gk/A403/polytrop.pdf

[4] Plastino, A.R.; Plastino, A. (1993). Stellar Polytropes and Tsallis Entropy. Physics Letters A 174(5):384-386. DOI: 10.1016/0375-9601(93)90195-6

[5] Bento, E.P.; Silva, J.R.P.; Silva, R. (2013). Non-Gaussian Statistics, Maxwellian Derivation and Stellar Polytropes. Physica A: Statistical Mechanics and its Applications 392.4 (2013): 666-672. DOI: 10.1016/j.physa.2012.10.022

[6] Shannon, C.E. (1948). A Mathematical Theory of Communication. Bell System Technical Journal 2(3):379–423. DOI: 10.1002/j.1538-7305.1948.tb01338.x

[7] Tsallis, C. (1960). Possible Generalization of Boltzmann-Gibbs Statistics, Journal of Statistical Physics, 1988, 52: 479–487. DOI:10.1007/BF01016429

[8] Kaniadakis, G.(2002). Statistical Mechanics in the Context of Special Relativity, Phys. Rev. E, 2002, 66, 056125. DOI: 10.1103/physreve.66.056125

[9] Sparavigna, A.C. (2015). On the Generalized Additivity of Kaniadakis Entropy, Int. J. Sci. 4(2):44-48. DOI: 10.18483/ijSci.627

[10] Kaniadakis, G. (2001). Non-Linear Kinetics Underlying Generalized Statistics, Physica A 296(3-4):405-425. DOI: 10.1016/s0378-4371(01)00184-4

[11] Santos, P.; Silva R.; Alcaniz J.S.; Anselmo, D.H.A.L. (2011). Generalized Quantum Entropies, Physics Letters A 375:3119–3123. DOI: 10.1016/j.physleta.2011.07.001

[12] Sparavigna, A.C. (2015). Relations Between Tsallis and Kaniadakis Entropic Measures and Rigorous Discussion of Conditional Kaniadakis Entropy, Int. J. Sci. 4(10): in print.

Information about this Article

Published on Monday 9th November, 2015 at 19:30:37.

The full citation for this Article is:

Sparavigna, A. (2015). Tsallis and Kaniadakis Entropic Measures in Stellar Polytropes. PHILICA.COM Article number 541.

Author comment added 10th November, 2015 at 19:09:48

Reference 12 is published. Here the details:

Amelia Carolina Sparavigna (2015). Relations Between Tsallis and Kaniadakis Entropic Measures and Rigorous Discussion of Conditional Kaniadakis Entropy, International Journal of Sciences 4(10):47-50 DOI: 10.18483/ijSci.866