Giant Core–Shell Nanospherical Clusters Composed of 32 Co or 32 Ni Atoms Held by 6 p - tert -Butylthiacalix[4]arene Units

Alexandre Gehin, Sylvie Ferlay, Jack M Harrowfield, D. Fenske, Nathalie Kyritsakas, Mir Wais Hosseini

► To cite this version:

Alexandre Gehin, Sylvie Ferlay, Jack M Harrowfield, D. Fenske, Nathalie Kyritsakas, et al.. Giant Core–Shell Nanospherical Clusters Composed of 32 Co or 32 Ni Atoms Held by 6 p - tert -Butylthiacalix[4]arene Units. *Inorganic Chemistry*, 2012, 51 (9), pp.5481-5486. 10.1021/ic300550v . hal-02301250

HAL Id: hal-02301250

<https://hal.science/hal-02301250>

Submitted on 27 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Giant core-shell nanospherical clusters composed of 32 Co or 32 Ni atoms held by 6 *p*-*tert*-butylthiacalix[4]arene units

Alexandre Gehin,^a Sylvie Ferlay,^{a*} Jack M. Harrowfield,^b D. Fenske,^c Nathalie Kyritsakas^a and Mir Wais Hosseini^{a*}

^a Laboratoire de Chimie de Coordination Organique, UMR CNRS 7140, Université de Strasbourg, F-67000 Strasbourg, France

^b Laboratoire de Chimie Supramoléculaire, Institut de Science et d'Ingénierie Supramoléculaires, Université de Strasbourg, 8, allée Gaspard Monge, 67083 Strasbourg, France

^c Institut für Anorganische Chemie, Universität Karlsruhe, D 76128 Karlsruhe, Germany

RECEIVED DATE (will be automatically inserted after manuscript is accepted)

Using combinations of *p*-*tert*-butylthiacalix[4]arene (TCA) and $[M(\text{DMSO})_6(\text{BF}_4)_2]$ salts ($M = \text{Co(II)}$ or Ni(II)), two almost isostructural core-shell type thermally stable giant nanoclusters, composed of 32 metal centres, 6 deprotonated calix units binding the metal centres by both their O and S atoms, 24 *m*-oxo or *m*-hydroxo bridging groups and 6 MeOH molecules, have been prepared under mild and reproducible conditions. For both giant clusters, the oxidation state II ($M^{II}_{32}O_{16}(\text{OH})_8(\text{CH}_3\text{OH})_6\text{TCA}_6$ ($M = \text{Co, Ni}$)) for the metal centre was demonstrated by XPS and electronic absorption spectroscopies.

Introduction

Polynuclear transition-metal compounds, in particular *m*-oxo and/or hydroxo bridged polyoxometallate type compounds, are of interest for their chemical and physical properties.¹ Several elegant strategies have been reported for the preparation of giant polynuclear species such as of, for example, iron², molybdenum³, manganese⁴ and cadmium⁵.

The calix[4]arene backbone **1** (Fig. 1 left)⁶ is an interesting unit for the design of polynucleating ligands. In particular, calix[4]arene derivatives in the cone conformation (Fig. 1 right) offer four OH groups oriented in a convergent manner. This feature has been explored for the formation of koilands, which are face to face calix[4]arenes doubly fused by two Si(IV)⁷, two Ti(IV)⁸, one Si(IV) and one Ti(IV)⁹, two Al(III)¹⁰, two Zn(II)¹¹, two Eu(III)¹² or four Nb(V)¹³ ions.

The thiacalix[4]arene backbone **2**¹⁴ (TCA) for which the four CH₂ groups connecting the aromatic moieties are replaced by sulphur atoms, offers four additional coordinating sites, *i.e.* the four S atoms¹⁵ and is a

candidate of choice for the generation of high nuclearity complexes. From **2**, which also adopts the cone conformation,¹⁶ tetranuclear Cu(II)¹⁷ and tetra- and hexa-nuclear Hg(II)¹⁸ species have been generated. Using dimercapto-¹⁹ and tetramercapto-calix[4]arene derivatives²⁰ **3** and **4** respectively, mononuclear and binuclear Hg(II) complexes have been reported. The synthesis and structural analysis of the tetramercapto-tetrathiacalix[4]arene **5**²¹ and some Hg(II)¹⁸ and low valence Mo and W²² or Ir and Rh²³ organometallic complexes have been also published.

The coordination ability²⁴ of **2** has been used for the preparation of other polynuclear species such as Cu₄,¹⁷ Mn₄,²⁵ Ln₄,²⁶ Ni₆,²⁷ V₆,²⁸ Fe₈,²⁹ Mn₄Ln₄,³⁰ Cu₉,³¹ Cu₁₀,³² and Co₂₄.³³ Recently, a new heterometallic cationic cluster $[\text{Co}^{II}_{24}(\text{M}^{\text{VI}}\text{O}_4)_8\text{TCA}_6\text{Cl}_6]^{2+}$ combining 24 Co(II) and 8 M(VI) ($M = \text{Mo}$ or W) atoms was published.³⁴ The highest nuclearity homometallic clusters reported are those containing 32 cobalt³⁵ ($\text{Co}^{II}_{24}\text{Co}^{III}_8\text{O}_{24}(\text{OH})_{24}\text{TCA}_6$) or nickel atoms ($\text{Ni}^{II}_{32}(\text{OH})_{40}\text{TCA}_6$)²⁷.

Figure 1: Chemical formulae of calix[4]arene, thiacalix[4]arene and mercaptathiacalix[4]arene derivatives (left) and a schematic representation of their cone conformation (right).

Here, we report on the synthesis, structural studies, thermal and XPS characterization of two new core-shell nanospherical clusters composed of 32Co(II) or 32Ni(II) held by 6 *p*-*tert*-butylthiacalix[4]arenes.

Experimental section

Preparation of Co_{32} : In a crystallisation tube (1 cm diameter), exposed to the atmosphere and at room temperature, diffusion of a DMF solution (5 mL) of $\text{Co(DMSO)}_6(\text{BF}_4)_2$ (230 mg, 0.28 mmol) into a DMF solution (1 mL) containing the thiacalixarene **2** (20 mg, 2.78×10^{-2} mmol) and Et_3N (0.1 mL), provided a green solution after *ca* one hour. This was filtered before MeOH (1 mL) was carefully added. After a week, a fine pale green precipitate had formed at the bottom of the tube. Upon standing further for *ca* 3 weeks, large pink crystals, suitable for single crystal X-ray diffraction, were formed on the wall of the tube. 97 mg, (48 % yield). MS (ESI (DMSO), *m/z*): 2285.7 (calculated for $\text{Co}_{32}(\text{TCA})_6(\text{CH}_3\text{OH})_6\text{O}_{16}(\text{OH})_8(\text{KNa}_2)^{3+}$, *m/z* = 2285.6); IR (ν , cm^{-1}): 752.3; 1092.7; 1255.7; 1448.6; 1672.3; 2956.0.

Synthesis of Ni_{32} : In a crystallisation tube (1 cm diameter), exposed to the atmosphere and at room temperature, diffusion of a DMF solution (5 mL) of $\text{Ni(DMSO)}_6(\text{BF}_4)_2$ (230 mg, 0.28 mmol) into a DMF solution (1 mL) containing the thiacalixarene **2** (20 mg, 2.78×10^{-2} mmol) and Et_3N (0.1 mL), provided a green solution after *ca* one hour. This was filtered before MeOH (1 mL) was carefully added. After a week, a fine green precipitate had formed at the bottom of the tube. Upon standing further for *ca* 3 weeks, large green crystals, suitable for single crystal X-ray diffraction, were formed on the wall of the tube. 95 mg, (*ca* 45 % yield). MS (ESI (DMSO), *m/z*): 3385.7 (calculated for $\text{Ni}_{32}(\text{TCA})_6(\text{CH}_3\text{OH})_6\text{O}_{16}(\text{OH})_8(\text{LiH})^{2+}$, *m/z* = 3385.6) ; 6788.2 (calculated for $\text{Ni}_{32}(\text{TCA})_6(\text{CH}_3\text{OH})_6\text{O}_{16}(\text{OH})_8\text{Na}^+$, *m/z* = 6787.8). IR (ν , cm^{-1}): 750.3; 1058.9; 1257.6; 1438.9; 1666.5; 2958.9.

Physical measurements

Single-Crystal Studies.

Data were collected at 173(2) K on a Bruker APEX8 CCD Diffractometer equipped with an Oxford Cryosystem liquid N_2 device, using graphite-monochromated Mo-K α ($\lambda = 0.71073$ Å) radiation. For both structures, diffraction data were corrected for absorption. Structures were solved using SHELXS-97 and refined by full matrix least-squares on F^2 using SHELXL-97. The hydrogen atoms were introduced at calculated positions and not refined (riding model).³⁶ In the case of Ni_{32} , due the disorder of solvent molecules, the SQUEEZE command was used.³⁷ CCDC XXX and CCDC XXX contain supplementary crystallographic data for the Co_{32} and Ni_{32}

nanospheres, respectively. They can be obtained free of charge from the Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/datarequest/cif.

Powder diffraction studies (PXRD)

For both Co_{32} and Ni_{32} nanoclusters, PXRD diagrams for the crystalline and powdered materials were collected on a Bruker D8 diffractometer using monochromatic Cu-K α radiation with a scanning range between 3.8 and 30° and a scan step of 2°/mn.

The desolvated forms of Co_{32} and Ni_{32} nanospheres ($\text{Ni}_{32}(\text{des})$ and $\text{Co}_{32}(\text{des})$) were generated upon heating the sample at 170 °C and the crystallinity was studied by PXRD. The thermal stability of Co_{32} and Ni_{32} nanoclusters was studied by first heating the powder to 300 °C and then measuring PXRD patterns at room temperature.

Thermogravimetric (TGA) Studies.

TGA measurements have been performed on both powdered single crystal samples of Co_{32} and Ni_{32} nanoclusters using a Pyris 6 TGA Lab System (Perkin-Elmer), using a N_2 flow of 20 mL/min and a heat rate of 10 °C/min.

XPS measurements.

XPS (X-ray photoelectron Spectroscopy) measurements were performed on powdered single crystal samples of both Co_{32} and Ni_{32} nanoclusters using a ThermoVGScientific photoelectron spectrometer equipped with a twin anode providing both unmonochromated Al K α and Mg K α radiation (1486.6 and 1453.6 eV, respectively). The spectrometer, equipped with a multichannel detector, was used in the constant resolution mode with a pass energy of 20 eV. The total resolution of the system was estimated at 0.55 eV. Spectra were referenced to the aliphatic hydrocarbon C1s signal at 285 eV.

Solid state UV measurements

UV/Visible spectra were obtained on powdered samples of single crystals of Co_{32} and Ni_{32} nanoclusters using a Shimadzu UV3600 spectrometer (data were recorded in the reflection mode using a 150 mm integrating sphere, with a resolution of 4 nm and a sampling rate of 300 nm min⁻¹)

Infra-red Studies

IR spectra were recorded on powdered samples of single crystals of Co_{32} and of Ni_{32} nanoclusters using a Perkin Elmer Spectrometer RX1.

Results and discussion

Using the *p*-*tert*-butylthiacalix[4]arene **2** in its cone conformation (Fig. 1), a reproducible synthetic procedure using identical conditions was established for the formation of giant clusters composed of 32 Co or 32 Ni atoms. The formation of these polynuclear assemblies displaying almost identical structures was

achieved under mild conditions. Upon slow diffusion at room temperature of a DMF solution containing the tetrafluoroborate salt of Co(II) or Ni(II) into a DMF solution of the TCA **2** and Et₃N as base, a powder was obtained which upon standing in MeOH afforded pink and green crystalline materials in the case of Co and Ni, respectively (see experimental section). It is worth noting that these species could only be produced when using [M^{II}(DMSO)₆(BF₄)₂] (M = Co or Ni) salts. Interestingly, the same type of crystalline material was obtained using EtOH or *i*-PrOH, however with much lower crystalline quality. Both clusters have been characterized by X-ray diffraction on single crystals.[‡]

In both cases, the diffraction data obtained were treated supposing the presence of discrete molecular entities with the following formula ((32M)(16O²⁻)(8OH⁻)(62⁴⁻)(6CH₃OH)). The giant clusters, with the longest distance between opposite CH₃ groups of *ca* 24.7 and of 23.7 Å for **Co**₃₂ and **Ni**₃₂ species, respectively, display almost identical connectivity patterns (Fig. 2 left). Whereas for **Co**₃₂ ((32Co)(16O²⁻)(8OH⁻)(6 **TCA**⁴⁻)(6CH₃OH)·3DMSO·3DMF) the solvent molecules (3 DMF and 3 DMSO) could be localised, for **Ni**₃₂, due to disorder of solvents, the structure was refined using the SQUEEZE procedure.³⁸

Thus, although both species crystallise in the same space group (P2₁/n), since the solvent molecules could not be identified in the case of **Ni**₃₂, the two crystals may or may not be isomorphous. The overall assembly in both cases may be described as a cubic metallic core composed of 8 Co or Ni centres located within a shell composed of 24 metal cations (Co or Ni) held together by 6 deprotonated thiocalix **2**⁴⁻ occupying the apices of an octahedron and forming an organic envelope (Fig. 2 right).

Figure 2: Crystal structure of the **M**₃₂ (M = Co or Ni) nano cluster. For the sake of clarity, the 8 core (light blue) and 24 shell (violet) metal centres are differentiated by colour. The 24 O atoms of the calix units (red), 24 m-oxo and m-hydroxo (green) and of MeOH units (pink) are also differentiated by colour. The C atom of the 6 MeOH molecules is coloured in brown. Solvent molecules are omitted (left) and schematic representation of the giant **M**₃₂ nanoclusters (M = Co or Ni) composed of a cubic **M**₈ core (blue) inserted inside of a spherical **M**₂₄ shell (black) surrounded by 6 thiocalix[4]arenes (grey) (right).

The aesthetically appealing structure of both clusters is rather complex. They will be described starting from their organic envelope and zooming towards their core.

As for the parent compound **2**,¹⁴ all (here, deprotonated) calix units (**2**⁴⁻) adopt the cone conformation (Fig. 1 right) with typical C-O distances in the 1.333(7)-1.368(8) and 1.281(13)-1.382(11) Å range for Co and Ni respectively and CS distances in the 1.773(6)-1.804(8) and 1.705(17)-1.844(13) Å range for Co and Ni, respectively. Among the 6 calix units, the *t*-butyl groups were found to be disordered over two positions (two calix with 4 and the remaining four with 2 disordered groups located on adjacent aromatic moieties). In the case of the **Co**₃₂ cluster, the three DMF solvent molecules are located within the cavities of the six calix units but with an occupancy of 50 %, *i.e.* one DMF molecule is disordered over two calix units. The three DMSO solvent molecules are located in the proximity of the metallic core with a Co...O distance of *ca* 3.4 Å. One of them is disordered over two positions with an occupancy of 50 %.

Each calix moiety behaves as a cluster keeper and binds 4 metal centres forming a square through both their phenoxide groups (O-M distance in the 2.022(4)-2.061(4) Å and 1.968(8)-2.060(9) Å range for Co and Ni respectively) and their sulfur atoms (S-M distance in the 2.4432(17)-2.4543(18) Å and 2.409(3)-2.439(3) Å range for Co and Ni, respectively). This motif is reminiscent of that found in the reported tetranuclear Cu(II)¹⁷ and Hg(II)¹⁸ species (Fig. 3).

Figure 3: A portion of the crystal structure of the **M**₃₂ (M = Co or Ni) nano cluster showing the 24 metal centres (violet) forming the shell of the giant cluster held by the 6 thiocalix units. For sake of clarity, the 24 O atoms of the calix units (red), 24 m-oxo and m-hydroxo (green) are differentiated by colour. Solvent molecules are omitted.

The 6 tetranuclear **M**₄ units are further interconnected by 16 m-oxo and 8 m-hydroxo groups forming thus a metallic shell of the sodalite type with a diameter of *ca* 10.63 and 10.38 Å (longest M-M distance) for **Co**₃₂ and **Ni**₃₂ respectively. The protonation state of the above-mentioned bridging O atoms cannot be specified. The M-O distances are in the 2.031(5)-2.062(5) and 1.989(7)-2.061(6) Å range for Co and Ni respectively. The coordination sphere around the metal centres is composed of 5 oxygen and 1 sulphur atoms. All metal centres adopt a distorted octahedral coordination geometry (Fig. 4).

Figure 4: A portion of the crystal structure of the M_{32} ($M = \text{Co}$ or Ni) nano cluster showing the 24 metal centres (violet) forming the shell and their surroundings. The O atoms are differentiated by colour (red and green) for clarity. Solvent molecules are omitted.

The metallic shell defines a cavity occupied by a octanuclear M_8 core. The 8 metallic centres occupying the corners of a distorted cube (M-M distance in the *ca* 4.20-4.38 Å range) are connected to the shell through the 24 m-oxo or m-hydroxo bridging groups (three O atoms per metal centre) with a O-M distance in the 2.057(5)-2.323(14) Å and 1.989(7)-2.24(2) Å range for Co and Ni, respectively.

Figure 5: A simplified view of the M_{32} ($M = \text{Co}$ or Ni) nanocluster found in the crystal lattices, showing the 24 metal centres (violet) forming the sodalite type shell and the 8 metals (light blue) forming a cubic core located at the centre of the giant cluster. The O atoms are differentiated by colour (red and green) for clarity. Solvent molecules are omitted.

Finally, within the cubic core, the 8 metal centres are connected by 6 MeOH molecules, with M-O distances in the 2.169(14)- 2.323(14) and 2.072(7)-2.39(2) Å range for Co_{32} and Ni_{32} respectively.

In regard to the organic envelope and metallic shell, the two clusters are very similar but they diverge in their cubic cores (Fig. 6). For Co_{32} cluster, of the six disordered MeOH molecules occupying the faces of the cubic arrangement, the oxygen atoms of two of them are disordered over four positions, whereas for the remaining four molecules, the oxygen atoms are disordered over three positions (Fig. 6 left). In the case of Ni_{32} cluster, the reverse is observed (Fig. 6 right).

Figure 6: Portions of the crystal structures of the nanoclusters showing the Co_8 (left) and Ni_8 (right) cubic metallic core interconnected by 6 MeOH (disordered O atoms in pink and C in brown). H atoms are omitted. For differences see text.

In the Co species, the inner Co_8 unit atoms have distorted octahedral O_6 coordination geometry, while in the Ni analogue, six have this form and two have distorted square-pyramidal O_5 coordination.

The packing of the two clusters is identical with no specific interactions between the polynuclear species (Fig. S1).

It is interesting to note that for the reported analogous but mixed oxidation state Co cluster,³⁴ 48 m-oxo or m-hydroxo bridging anions are involved in the coordination sphere of the metal centres, whereas for the Ni analogue,²⁷ 40 m-hydroxo groups were found to be bound to the metal. For both Co_{32} and Ni_{32} species reported here, in addition to six MeOH molecules bound to the metallic core, only 24 m-oxo or m-hydroxo groups are connected to metal centres.

The crystallisation process used for the preparation of both Co- and Ni-based nanospheres led to single crystals attached to the walls and a powder deposited at the bottom of the tube. The crystals and powder were separated and dried in air. Both phases were analysed using PXRD (Fig. 8 and 9) and similar patterns were observed. For the Co_{32} nanosphere dried in air, the observed pattern (Fig. 7b) fits well with the simulated one (Fig. 7a). Interestingly, the powder sample heated at 170 °C to afford the desolvated material also displayed a similar pattern (Fig. 8c bottom) indicating the conservation of the crystallinity of the sample. Retention of the same pattern even after heating at 300 °C (Fig. 7d) showed the thermal stability of the sample. The same behaviour was observed for the Ni_{32} nanosphere (Fig.8).

Figure 7: Comparison of the simulated (a) and recorded PXRD patterns for **Co**₃₂ at RT (b), after heating at 170 °C (c) and after heating at 300 °C (d).

Figure 8: Comparison of the simulated (a) and recorded PXRD patterns for **Ni**₃₂ at RT (b), after heating at 170 °C (c) and after heating at 300 °C (d).

The thermal stability of both species was also investigated by thermogravimetric analysis. For both **Co**₃₂ and **Ni**₃₂ species, in the 30 °C - 460 °C temperature range, a weight loss of *ca* 25 % was observed prior to decomposition which appeared at 440 °C for **Ni**₃₂ and at *ca* 460 °C for **Co**₃₂ (see figure S2). Although the weight loss prior to decomposition was assumed to be due to volatilisation of solvents, steps corresponding to the loss of different species were not discerned.

In order to clarify the oxidation state of the Co and Ni cations, X-Ray Photoelectron Spectroscopy (XPS) was used. Measurements were carried out on single crystals, using as reference salts **Co**(OH)₂•6•(NO₃)₂ for **Co**₃₂ and **Ni**(OH)₂•6•SO₄ for **Ni**₃₂. For both **Co**₃₂ (Fig. 9 top) and **Ni**₃₂ (Fig. 10 bottom) giant clusters, the spectra displayed the M2p_{3/2}, M2p_{1/2} and M2s signatures. Significantly, the match, in terms of binding energy and shape, between the recorded spectra for the two clusters and the corresponding reference compounds indicates that, in both cases, the metal centres are in the oxidation state II with a deformed octahedral environment.

Figure 9: The X-Ray Photoelectron Spectra (XPS) of the polycrystalline samples **Co**₃₂ (top, a) and **Ni**₃₂ (bottom, a) exhibiting the M2p_{3/2}, M2p_{1/2} and M2s signatures (M = Co or Ni). The spectra are compared to those of **Co**(OH)₂•6•SO₄ (top, b) for **Co**₃₂ and **Ni**(OH)₂•6•(NO₃)₂ (bottom, b) for **Ni**₃₂.

The UV-Visible reflectance spectra of **M**₃₂ (M = Co or Ni) clusters were recorded in the solid state using polycrystalline samples (Fig. 10). In agreement with the observations made by XPS spectroscopy mentioned above, in both cases, the recorded spectra are typical of M(II) species. For the **Co**₃₂ giant cluster, considering the oxidation state II and an octahedral geometry, three transitions in the 250-800 nm region are expected. Indeed, three major bands at 320nm, assigned to the ⁴T_{1g} (F) → ⁴T_{2g} (F) transition and at 550 nm to the ⁴T_{1g} (F) → ⁴T_{1g} (P) and ⁴T_{1g} (F) → ⁴A_{2g} (F) transitions, were observed. For **Ni**₃₂, again considering a Ni(II) cation in octahedral geometry, two transitions in the 250-800 nm region are expected and were indeed observed at 330 nm for the high energy ³A_{2g} → ³T_{1g}(P) transition and at 670 nm for the low energy band assigned to the ³A_{2g} → ³T_{1g}(F) transition. The presence of shoulders at *ca* 400 nm probably results from the presence of M centres in a significantly deformed octahedral geometry as discussed in the section dealing with structural analysis.

Figure 10: UV-Vis reflectance spectra of **Ni**₃₂ (green) and of **Co**₃₂ (pink) recorded in the solid state at RT.

Finally, in support of the above mentioned experimental facts, using the observed M-O or M-S distances, a BVS calculation³⁸ on both clusters revealed that for none of the metal centres was an oxidation state higher than 2.3 appropriate (see supporting information).

Conclusions

A mild, reproducible procedure for obtaining thermally stable, giant core-shell nanoclusters containing 32 Co or Ni atoms enclosed by six tetrathiacalix[4]arene units has been established. The structure of both species was elucidated using X-ray diffraction techniques both on single crystals and on powdered crystalline samples. Both clusters are based on a cubic metallic core composed of 8 metal centres located in the centre of a sodalite type shell composed of 24 metals surrounded by 6 calix units as an organic envelope (Fig. 2 right). In marked contrast with a reported analogous cluster,³⁵ obtained under hydrothermal conditions and which contains both Co(II) and Co(III) centres, for the **Co**₃₂ and **Ni**₃₂ clusters reported here, the metal centres are clearly in the oxidation state II.

Using the same methodology, the formation of analogous nanoclusters using other calixarene derivatives and metal centres is currently under investigation.

ACKNOWLEDGMENT

This work was financially supported by the University of Strasbourg (UdS), the Centre National de la Recherche Scientifique (CNRS), Institut Universitaire de France (IUF) and the International Centre for Frontier Research in Chemistry (FRC), Strasbourg. Pierre Bernardt (LMSPC, Strasbourg) is warmly acknowledged for the XPS measurements

Supporting information

TGA traces for **Co**₃₂ and **Ni**₃₂, as well as BVS calculations.

† (a) Crystal data for **Co**₃₂: (C₄₀H₄₄O₄S₄)₆Co₃₂(CH₃OH)₆ O₁₆(OH)₈, (C₃H₇NO)₃ (C₂H₆OS)₃, M = 7225.70, monoclinic, space group P2₁/n, a = 24.7390(7) Å, b = 26.8037(7) Å, c = 32.6310(9) Å, β = 94.2370(10)°, V = 21578.4(10) Å³, T = 173(2) K, Z = 2, D_c = 1.112 g·cm⁻³, μ = 1.371 mm⁻¹, 149389 collected reflections, 48887 independent [R(int) = 0.0463], GOF = 1.052, R1 = 0.0909, wR2 = 0.2687 for I > 2σ(I) and R1 = 0.1398, wR2 = 0.3015 for all data ;

(b) Crystal data for **Ni**₃₂: (C₄₀H₄₄O₄S₄)₆Ni₃₂(CH₃OH)₆ O₁₆(OH)₈, M = 6732.73, monoclinic, space group P2₁/n, a = 24.5603(5) Å, b = 26.6097(6) Å, c = 32.2446(7) Å, β = 93.4090(10)°, V = 21035.9(8) Å³, T = 173(2) K, Z = 2, D_c = 1.040 g·cm⁻³, μ = 1.557 mm⁻¹, 163687 collected reflections, 48655 independent [R(int) = 0.0852], GOF = 1.366, R1 = 0.1074, wR2 = 0.2613 for I > 2σ(I) and R1 = 0.2169, wR2 = 0.2949 for all data.

References

- [1] (a) Fenske, D.; Ohmer, J.; Hachgenei, J. *Angew. Chem., Int. Ed. Engl.* **1985**, *24*, 993-995; (b) Liu, T.; Zhang, Y.-J.; Wang, Z.-M.; Gao, S. *J. Am. Chem. Soc.* **2008**, *130*, 10500-10501; (c) Alborés, P.; Rentschler, E. *Angew. Chem., Int. Ed.* **2009**, *48*, 9366-9370.
- [2] Taft, K. L.; Lippard, S. J. *J. Am. Chem. Soc.* **1990**, *112*, 9629-9630.
- [3] (a) Müller, A.; Kögerler, P. *Coord. Chem. Rev.* **1999**, *182*, 3-17; (b) Müller, A.; Roy, S. *Coord. Chem. Rev.*, **2003**, *245* 153-166; (c) Long, D. L.; Tsunashima, R.; Cronin, L. *Angew. Chem. Int. Ed.* **2010**, *49*, 1736-1758.
- [4] (a) Goldber, D. P.; Caneschi, A.; Delfs, C. D.; Sessoli, R.; and Lippard, S. J. *J. Am. Chem. Soc.* **1995**, *117*, 5789-5800; (b) Bagai, R.; Christou, G. *Chem. Soc. Rev.* **2009**, *38*, 1011-1026.
- [5] Argent, S. P.; Greenaway, A.; Gimenez-Lopez, M.; Lewis, W.; Nowell, H.; Khlobystov, A. N.; Blake, A. J.; Champness, N. R. and Schröder, M. *J. Am. Chem. Soc.*, **2012**, *134*, 55-58.
- [6] (a) Gutsche, C. D. *Calixarenes Revisited*: Monographs in Supramolecular Chemistry; Royal Society of Chemistry: Cambridge, U.K., **1998**; (b) *Calixarenes 2001*; Asfari, Z.; Böhmer, V.; Harrowfield, J.; Vicens, J., Eds.; Kluwer: Dordrecht, The Netherlands, **2001**.
- [7] Delaigue, X.; Hosseini, M. W.; De Cian, A.; Fischer, J.; Leize, E.; Kieffer, S.; Van Dorsselaer, A. *Tetrahedron Lett.* **1993**, *34*, 3285-3288.
- [8] Olmstead, M. M.; Sigel, G.; Hope, H.; Xu, X.; Power, P. P. *J. Am. Chem. Soc.* **1985**, *107*, 8087-8091.
- [9] Delaigue, X.; Hosseini, M. W. *Tetrahedron Lett.* **1993**, *34*, 7561-7564.
- [10] Atwood, J. L.; Bott, S. G.; Jones, C.; Raston, C. J. *J. Chem. Soc., Chem. Commun.* **1992**, 1349-1351.
- [11] Atwood, J. L.; Junk, P. C.; Lawrence, S. M.; Raston, C. L. *Supramol. Chem.* **1996**, *7*, 15-17.
- [12] Bilyk, A.; Harrowfield, J. M.; Skelton, B. W.; White, A. H. *J. Chem. Soc., Dalton Trans.* **1997**, 4251-4256.
- [13] Corazza, F.; Floriani, C.; Chiesi-Villa, A.; Guastini, C. *Chem. Commun.* **1990**, 1083-1084.
- [14] Kumagai, H.; Hasegawa, M.; Miyazaki, S.; Sugawa, Y.; Sato, Y.; Hori, T.; Ueda, T.; Kamiyama, H.; Miyano, S. *Tetrahedron Lett.* **1997**, *38*, 3971-3971.
- [15] Hosseini, M. W. in *Calixarenes 2001*; Asfari, Z.; Böhmer, V.; Harrowfield, J.; Vicens, J., Eds.; Kluwer: Dordrecht, The Netherlands, **2001**. 110-129.
- [16] Akdas, H.; Bringle, L.; Graf, E.; Hosseini, M. W.; Mislin, G.; Pansanel, J.; De Cian, A.; Fischer, J. *Tetrahedron Lett.* **1998**, *39*, 2311-2314.
- [17] Mislin, G.; Graf, E.; Hosseini, M. W.; Bilyk, A.; Hall, A. K.; Harrowfield, J. M.; Skelton, B. W.; White, A. H. *Chem. Commun.* **1999**, 373-374.
- [18] Akdas, H.; Graf, E.; Hosseini, M. W.; De Cian, A.; Bilyk, A.; Skelton, B. W.; Koutsantonis, G. A.; Murray, I.; Harrowfield, J. M.; White, A. H. *Chem. Commun.* **2002**, 1042-1043.
- [19] (a) Delaigue, X.; Hosseini, M. W.; Kyritsakas, N.; De Cian, A.; Fischer, J. *Chem. Commun.* **1995**, 609-610; (b) Rao, P.; Enger, O.; Graf, E.; Hosseini, M. W.; De Cian, A.; Fischer, J. *Eur. J. Inorg. Chem.* **2000**, 1503-1508.
- [20] Delaigue, X.; Harrowfield, J. M.; Hosseini, M. W.; De Cian, A.; Fischer, J.; Kyritsakas, N. *Chem. Commun.* **1994**, 1579-1580.
- [21] Rao, P.; Hosseini, M. W.; De Cian, A.; Fischer, J. *Chem. Commun.* **1999**, 2169-2170.
- [22] Buccella, D.; Parkin, G. *Chem. Commun.* **2009**, 289-291.
- [23] Hirata, K.; Suzuki, T.; Noya, A.; Takei I.; Hidai, M. *Chem. Commun.* **2005**, 371-3720.
- [24] Kajiwara, T.; Iki, N.; Yamashita, M. *Coord. Chem. Rev.* **2007**, *251*, 1734-1746.
- [25] (a) Desroches, C.; Pilet, G.; Borshch, S. A.; Parola, S.; Luneau, D. *Inorg. Chem.* **2005**, *44*, 9112-9120; (b) Karotsis, G.; Teat, S. J.; Wernsdorfer, W.; Piligkos, S.; Dalgarno, S. J.; Brechin, E. K. *Angew. Chem., Int. Ed.* **2009**, *48*, 8285-8288.
- [26] (a) Bi, Y. F.; Wang, X. T.; Liao, W. P.; Wang, X. W.; Deng, R. P.; Zhang, H. J.; Gao, S. *Inorg. Chem.* **2009**, *48*, 11743-11747 ; (b) Bilyk, A.; Dunlop, J. W.; Fuller, R. O.; Hall, A. K.; Harrowfield, J. M.; Wais Hosseini, M. W.; Koutsantonis, G. A.; Murray, I. W.; Skelton, B. W.; Sobolev, A. N.; Stamps, R. L.; White, A. H. *Eur. J. Inorg. Chem.* **2010**, 2127-XXX.

-
- [27] Bilyk, A.; Dunlop, J. W.; Fuller, R. O.; Hall, A. K.; Harrowfield, J. M.; Wais Hosseini, M. W.; Koutsantonis, G. A.; Murray, I. W.; Skelton, B. W.; Stamps, R. L.; White, A. H. *Eur. J. Inorg. Chem.* **2010**, 2106-2126.
- [28] Aronica, C.; Chastanet, G.; Zueva, E.; Borshch, S. A.; Clemente-Juan, J. M.; Luneau, D. *J. Am. Chem. Soc.* **2008**, *130*, 2365-2371.
- [29] Desroches, C.; Pilet, G.; Szilágyi, P.; Molnár, G.; Borshch, S. A.; Bousseksou, A.; Parola, S.; Luneau, D. *Eur. J. Inorg. Chem.* **2006**, 357-365.
- [30] Karotsis, G.; Kennedy, S.; Teat, S. J.; Beavers, C. M.; Fowler, D. A.; Morales, J. J.; Evangelisti, M.; Dalgarno, S. J.; Brechin, E. K. *J. Am. Chem. Soc.* **2010**, *132*, 12983-12990.
- [31] Karotsis, G.; Kennedy, S.; Dalgarno, S. J.; Brechin, E. K. *Chem. Commun.* **2010**, 3884-3886.
- [32] Kajiwar, T.; Kon, N.; Yokozawa, S.; Ito, T.; Iki, N.; Miyano, S. *J. Am. Chem. Soc.* **2002**, *124*, 11274-11275.
- [33] Bi, Y. F.; Xu, G. C.; Liao, W. P.; Du, S. C.; Wang, X. W.; Deng, R. P.; Zhang, H. J.; Gao, S. *Chem. Commun.* **2010**, 6362-6364.
- [34] Bi, Y. F.; Du, S. C.; Liao, W. P. *Chem. Commun.* **2011**, 4724-4726.
- [35] Bi, Y. F.; Wang, X. T.; Liao, W. P.; Wang, X. F.; Wang, X. W.; Zhang, H. J.; Gao, S. *J. Am. Chem. Soc.* **2009**, *131*, 11650-11651.
- [36] Sheldrick, G. M. : *Program for Crystal Structure Solution*; University of Göttingen: Göttingen, Germany, **1997**.
- [37] Spek A. *J. Appl. Cryst.* **2003**, *36*, 7-13
- [38] Willis, A. S. Brown, I. *Valist*, version 3.0.15; CEA: Saclay, France, **1999**.

Table of Contents :

Thermally stable, giant core-shell nanoclusters containing 32 Co or Ni atoms enclosed by six tetrathiacalix[4]arene units are generated under mild and reproducible conditions.