UV photofragmentation of cold Cytosine-M⁺ complexes (M⁺: Na⁺, K⁺, Ag⁺)

*Martín I. Taccone,^{a,b,c} Andrés Cruz,^{a,b,c} Jordan Dezalay,^d Satchin Soorkia,^d Michel Broquier,^{d,e}
Gilles Grégoire,^{d,*} Cristián G. Sánchez^{a,f,#} and Gustavo A. Pino^{a,b,c,*}*

^a INFIQC (CONICET), Ciudad Universitaria, Pabellón Argentina, 5000 Córdoba, Argentina.

^b Departamento de Fisicoquímica, Fac. de Ciencias Químicas, Universidad Nacional de Córdoba,
Ciudad Universitaria, Pabellón Argentina, X5000HUA Córdoba, Argentina

^c Centro Láser de Ciencias Moleculares, Universidad Nacional de Córdoba, Ciudad Universitaria,
Pabellón Argentina, X5000HUA Córdoba, Argentina

^d Institut des Sciences Moléculaires d'Orsay (ISMO), CNRS, Univ. Paris-Sud, Université Paris-Saclay, F-91405 Orsay, France

^e Centre Laser de l'Université Paris-Sud (CLUPS/LUMAT), Univ. Paris-Sud, CNRS, IOGS,
Université Paris-Saclay, F-91405 Orsay, France

^f Departamento de Química Teórica y Computacional, Facultad de Ciencias Químicas,
Universidad Nacional de Córdoba, X5000HUA Córdoba, Argentina

[#] Present address: CONICET & Facultad de Ciencias Exactas y Naturales, Universidad Nacional
de Cuyo, Mendoza CP5500, Argentina

ABSTRACT

The UV-photofragmentation spectra of cold Cytosine- M^+ complexes (M^+ : Na^+ , K^+ , Ag^+) were recorded and analyzed through comparison with geometry optimizations and frequency calculations of the ground and excited states at the SCS-CC2/Def2-SVPD level of theory. While in all complexes, the ground state minimum geometry is planar (C_s symmetry), the $\pi\pi^*$ state minimum geometry has the NH_2 group slightly twisted and an out-of-plane metal cation. This was confirmed by comparing the simulated $\pi\pi^*$ Franck-Condon spectra with the vibrationally resolved photofragmentation spectra of $CytNa^+$ and $CytK^+$. Vertical excitation transitions were also calculated to evaluate the energies of the CT states involving the transfer of an electron from the Cyt moiety to M^+ . For both $CytK^+$ and $CytNa^+$ complexes, the first CT state corresponds to an electron transfer from the cytosine aromatic π ring to the antibonding σ^* orbital centered on the alkali cation. This $\pi\sigma^*$ state is predicted to lie much higher in energy (> 6 V) than the band origin of the $\pi-\pi^*$ electronic transition (around 4.3 eV) unlike what is observed for $CytAg^+$ complex for which the first excited state has a $n\sigma^*$ electronic configuration. This is the reason for the absence of the $Cyt^+ + M$ charge transfer fragmentation channel for $CytK^+$ and $CytNa^+$ complexes.

INTRODUCTION

Purine and pyrimidine bases (Adenine, Guanine, Thymine and Cytosine) are the building blocks of DNA. The DNA damage upon UV photoexcitation is an important cause for mutagenic effects in biological systems¹⁻⁴ and further possible development of different types of cancer.⁵⁻⁷ A useful strategy to understand these processes is a bottom-up approach by means of studying the excited state photodynamics of the building blocks.⁸⁻¹⁶ It is also known that the interaction of the DNA bases with cations (H^+ or metal cations M^+) can induce the formation of non-canonical structures¹⁷⁻²⁰ as well as to facilitate proton transfer (PT) reactions within base pairs.^{21,22}

During the last years we have studied the UV spectroscopy and excited state dynamics of protonated DNA bases^{13,23,24} as well as the effect of substituting H^+ by Ag^+ on the structure and reactivity of Cytosine and its dimeric form.^{19,25,26} In this regard, it was recently shown that the first excited state of $CytAg^+$ complex is a charge transfer (CT) state in which an electron from a n_O orbital of Cyt is transferred to a σ^* orbital on Ag^+ . The excitation of the complex to the $n_O\sigma^*$ state at 3.93 eV leads to ionized cytosine Cyt^+ and neutral Ag fragments.²⁵ Since ionization of DNA bases could induce PT reactions by reducing the energy barrier,^{27,28} the presence of CT states producing Cyt^+ at low excitation energies could alter the fidelity of DNA inducing mutagenic effects.^{29,30}

Moreover, the ground state fragmentation of $CytAg^+$ by infrared multiphoton dissociation (IRMPD) produces HNCN loss as the only fragmentation channel³¹ at variance with the fragmentation by IRMPD of $CytM^+$ complexes ($M = Li, Na, K, Rb, Cs$) for which the only fragmentation channel was the neutral Cyt loss.³² Considering this difference and the remarkable CT character of the first excited state of $CytAg^+$, the goal of this work was to study the effect of some of the two most abundant metal cations in biological systems, e.g. Na^+ and K^+ on the UV

photofragmentation of the CytNa^+ and CytK^+ complexes as compared to CytAg^+ by means of photofragmentation spectroscopy of cold ions.

EXPERIMENTAL AND THEORETICAL METHODS

The experiments have been conducted at the Centre Laser of the Université Paris-Sud (CLUPS) in Orsay. The experimental setup has been previously presented³³ and is briefly summarized here. It comprises an electrospray ion source (ESI), a 3D-quadrupole ion trap QIT (Jordan TOF, Inc.) mounted on a cold head of a compressed helium cryostat (CH-204S, Sumitomo) that maintains the temperature of the trap around 10-15 K, and a linear time-of-flight mass spectrometer (TOF-MS). The complexes of interest are produced in the ESI source from a water/methanol solution (50/50 by volume) of Cytosine and alkali salt (NaCl and KI) or AgNO_3 at 100 μM . The complexes are first stored in an octupole trap for 100 ms and then extracted, mass-selected and transferred into the 3D-QIT where they are thermalized through collisions with helium buffer gas introduced by a pulsed valve (General valve, series 9) a few ms before the entrance of the ions. Two UV lasers were used for the photodissociation of the ions, either a ps OPA laser (EKSPLA PG 411) with 8 cm^{-1} bandwidth and 16 ps time resolution, or the output of a frequency doubled ns dye laser (Quantel TDL 90) whose resolution is 0.2 cm^{-1} . Both lasers operated at 10 Hz and are triggered at least 40 ms after the entrance of the ions in the QIT to ensure their complete cooling and pumping of the buffer gas. All of the ionic fragments and remaining parent ions are extracted and accelerated for mass analysis in the linear TOF-MS and finally detected by microchannel plates (Z-Gap, Jordan TOF, Inc.). The resonant absorption of the UV laser is observed by monitoring the appearance of the ionic fragments at lower masses. In case of CytNa^+ , because of the low m/z cut-off of the Paul Trap, the photofragment sodium cation cannot be trapped in the QIT along with the parent ions, so the ions are immediately

extracted from the trap (within 1 μ s) after interaction with the laser. For CytK^+ , the potassium cation is stored, and we did not observe any other photofragment even at long fragmentation delays up to 30 ms after laser excitation. For CytAg^+ , the main photofragment at all excitation wavelengths is the cytosine cation Cyt^+ along with Ag^+ and loss of 43 Da (HNCO loss) from the CytAg^+ complex as minor fragments. Action spectra are recorded by scanning the laser frequency and monitoring the cationic alkali signal normalized by the intensity of laser and the parent ion signal.

The equilibrium geometries and the vibrational frequencies of the complexes in their ground and excited states were calculated with TURBOMOLE package (v6.6)³⁴ at SCS-CC2³⁵ level of theory, making use of resolution of identity (RI)³⁶ and Karlsruhe “def2” split valence polarization basis set with diffuse functions (Def2-SVPD).³⁷ In particular, the split-component scaled approach (SCS) was applied since it improves the accuracy of CC2 calculations for the charge transfer states.³⁸ The calculated vibrationally-resolved electronic spectra were obtained with the PGOPHER spectra simulator package³⁹ using the calculated frequencies of the ground and excited states. For sake of comparison, the simulation was performed at 0 K and using a convoluted with a gaussian function of 8 cm^{-1} (full-width at half-maximum FWHM).

RESULTS AND DISCUSSION

The UV photofragmentation of the CytNa^+ and CytK^+ complexes result in the formation of neutral cytosine and the corresponding alkali cation Na^+ and K^+ at m/z 23 and 39, respectively. It is worthy to note that Na^+ and K^+ are the only fragments obtained upon laser irradiation and no other ionic fragment, in particular the cytosine cation, was observed. This is at opposite to the result found for the CytAg^+ complex, in which the complex is excited to a charge transfer (CT) state from where it dissociates to Cyt^+ as the main ionic fragment.²⁵ However, the present result

is consistent with those reported by Rodgers and co-workers on the ground state fragmentation of the same alkali complexes induced by multiple photon dissociation (IRMPD).³² This similarity strongly suggests that upon UV excitation, the complexes undergo internal conversion to the ground state followed by statistical fragmentation.

Figure 1. UV photodissociation spectra of CytNa⁺ (top) and CytK⁺ (bottom) recorded with a mid-resolution (8 cm⁻¹) ps OPO laser.

The vibrationally-resolved UV photofragmentation spectra of both complexes shown in Figure 1 were recorded in the spectral range of 34500 to 36500 cm⁻¹. Above this excitation energy, the vibronic spectra are unresolved, exhibiting a broad absorption band decaying in intensity until 40 000 cm⁻¹. The 0₀⁰ transition of CytNa⁺ is detected at 35028 cm⁻¹ (4.34 eV). The band origin

(BO) has the largest intensity among the other vibronic transitions, revealing the good overlap between the vibrational wave functions of the ground and excited states, thus small geometry changes between the equilibrium structures of both states. For CytK⁺ complex, the 0₀⁰ band is found at 34707 cm⁻¹ (4.30 eV) but at variance with CytNa⁺, the BO is not the most intense vibronic transition, suggesting a larger equilibrium geometry difference between the ground and excited states.

Scheme 1 Atom numbering of the Cyt-M⁺, cytosine in its keto-amino form.

The most stable form of the ground state geometries of all complexes is planar (Cs symmetry), containing the keto-amino tautomer of cytosine in which the metal cation simultaneously interacts with the carbonyl oxygen O₇ and the nitrogen N₃ atoms (see scheme 1) in agreement with a previous report by Rodgers and co-workers.^{32,40} For the alkali complexes, geometry optimization of the first excited state which corresponds to a $\pi\pi^*$ transition also led to planar structure for both complexes. Nevertheless, they exhibited two imaginary frequencies indicating that they do not correspond to true minima of the excited state PES. The modes associated to the imaginary frequencies correspond to twist motion of the NH₂ group and out-of-plane motion of

the metal cation. A re-optimization of this structure starting with a small geometry distortion along the pyrimidinic out-of-plane motion finally lead, in both complexes, to true minima of the PES whose structures exhibit a slightly out-of-plane metal cation and small twist of the NH_2 group. Figure 2 shows the optimized structures of the S_1 $\pi\pi^*$ excited states for CytNa^+ and CytK^+ , the most relevant structural parameters (interatomic distances and dihedral angles) are reported in Table 1. As it was presumed from the photofragmentation spectra (Fig. 1), a larger geometry distortion from the planar Cs structure is found for CytK^+ than for CytNa^+ .

Table 1: interatomic distance (Å) and dihedral angle (deg) of the optimized structures of Cyt-M^+ (M: Na, K, Ag) in the ground and excited electronic states.

	$\text{Na}^+\dots\text{Cyt}$		$\text{K}^+\dots\text{Cyt}$		$\text{Ag}^+\dots\text{Cyt}$		
	S_0 (Cs)	$\pi\pi^*$	S_0 (Cs)	$\pi\pi^*$	S_0 (Cs)	$n\sigma^*$ (Cs)	$\pi\pi^*$
$\text{N}_3\dots\text{M}$	2.466	2.351	2.874	2.738	2.287	2.259	2.195
$\text{O}_7\dots\text{M}$	2.234	2.374	2.558	2.74	2.394	2.170	2.514
$\text{O}_7\text{C}_2\text{N}_3\text{M}$	0	- 4	0	-7.4	0	0	1
$\text{H}_1\text{N}_8\text{C}_4\text{N}_3$	0	36	0	41	0	0	18
$\text{H}_2\text{N}_8\text{C}_4\text{N}_3$	0	15	0	15	0	0	12

Figure 2. S_1 optimized structures and molecular orbitals involved in excited states of CytNa⁺, CytK⁺ and CytAg⁺.

The calculated vertical (E_v) and adiabatic (E_{ad}) transition energies to the most relevant excited states for the photophysics of all complexes are shown in Table 2, together with the experimental energy for the 0_0^0 bands. In addition, the corresponding information for the CytAg⁺ complex is also reported for comparison. In all cases, adiabatic energies were corrected by difference in the zero-point energy between the ground and excited states (ΔZPE), allowing a direct comparison with the experimental BOs. The calculated adiabatic energies of the first excited states are in very good agreement with the experimental values of the three complexes, been the difference between the calculated and the experimental values as small as 0.7% for CytNa⁺, 1.2% for CytK⁺ and 1.3 % for CytAg⁺.

Table 2. Vertical (E_v) and adiabatic (E_{ad}) transition energies for the most relevant electronic excited states with the corresponding oscillator strengths (Osc. Str.) for CytNa⁺, CytK⁺ and CytAg⁺ complexes calculated at the SCS-CC2/Def2-SVPD level of theory. The experimental values (bold) are in parenthesis.

Complex	State	E_v	E_{ad}	Osc. Str.
CytNa ⁺	S ₁ ($\pi\pi^*$)	4.95	4.31 (4.34)	0.0940
	S ₄ ($\pi\sigma^*$)	6.45	-	0.0025
	S ₁₁ ($n_O\sigma^*$)	7.36	-	0.0529
	S ₁₂ ($n_N\sigma^*$)	7.81	-	0.1296
CytK ⁺	S ₁ ($\pi\pi^*$)	4.93	4.25 (4.30)	0.0859
	S ₄ ($\pi\sigma^*$)	6.46	-	0.0019
	S ₁₁ ($n_N\sigma^*$)	7.28	-	0.0578
	S ₁₂ ($n_O\sigma^*$)	7.70	-	0.0784
CytAg ⁺	S ₁ ($n_O\sigma^*$)	4.15	3.98 (3.93)	0.0077
	S ₂ ($n_N\sigma^*$)	4.84	-	0.0079
	S ₃ ($\pi\pi^*$)	4.89	4.39 (4.32)	0.0909
	S ₇ ($\pi\sigma^*$)	5.56	-	$3.7 \cdot 10^{-5}$

E_{ad} are corrected by the zero point energy difference (ΔZPE) between the two states. E_v and E_{ad} are given in eV.

Frequency calculations for the ground and excited states equilibrium geometries were used to simulate the vibronic spectra of both complexes. The comparison of the calculated and experimental photofragmentation spectra is shown in Figure 3. The active modes for both complexes are shown in Figure SI 1. For both complexes, the first two out-of-plane ν_1 and ν_2 modes which involve twist motion of the amino group have the strongest Franck-Condon

activities, while a second set of vibronic progression is built from the in-plane ν_{12} modes at 532 cm^{-1} and 528 cm^{-1} for CytNa^+ and CytK^+ , respectively. For the ν_2 mode, the alkali cation also shows an in-plane translation. The intensities of the two out-of-plane modes are stronger in CytK^+ than in CytNa^+ , in agreement with the larger structural deviation from the Cs structure in the former. In overall, the vibronic spectra are satisfyingly reproduced at this level of theory for both complexes, confirming that the locally excited $\pi\pi^*$ state has a slightly bent structure with the alkali atom out of the plane of cytosine.

Figure 3: Comparison of the experimental and simulated vibronic spectra of (a) CytNa^+ and (b) CytK^+ .

A close inspection to the spectra shows that each band is slightly broader than the bandwidth of the picosecond laser (full width at half maximum, $\text{FWHM} = 8 \text{ cm}^{-1}$). Thus, to get more information on the origin of this broadening the first 600-1000 cm^{-1} of each spectrum was also recorded with a nanosecond dye laser ($\text{FWHM} = 0.2 \text{ cm}^{-1}$) and a comparison of both is shown in Figure SI 2. The vibronic spectra recorded with the high-resolution UV laser exhibit similar transition widths for both complexes, revealing an homogeneous broadening in both complexes.

The bandwidth for each 0_0^0 band is larger than the broadening due to the rotational contour at 20 K (fitted with a Gaussian function of 2 cm^{-1} width) as expected and observed in systems with similar rotational constants.⁴¹ Besides, the vibronic transitions become even broader when the first out-of-plane modes (ν_1 and ν_2) are excited (See Figure SI 1 for the vibrational modes assignment). The lifetime broadening (LTB), reported in Table 3 is estimated from the width of the Gaussian function used to fit the vibronic transitions, which is at least twice larger than the expected value (2 cm^{-1}). These LTBs are used to obtain an estimation of the excited state lifetimes (τ) of each vibronic state according to the following relationship:

$$LTB (\text{cm}^{-1}) > (2 \pi c \tau)^{-1} = \frac{5.3 \times 10^{-12}}{\tau(\text{s})} \quad (1)$$

where c is the speed of the light in vacuum.

Table 3 shows that the excited state lifetimes at the band origin is 1.2 and 1.3 ± 0.2 ps for CytNa⁺ and CytK⁺, respectively and it is significantly reduced upon excitation of the ν_1 and ν_2 out-of-plane modes. It is noteworthy that no further broadening as compared to the band origin is observed following excitation of the ν_{12} in-plane mode at about 530 cm^{-1} in both complexes. The short excited state lifetimes estimated for these complexes together with the specificity on the out-of-plane modes strongly suggests that the main deactivation process is the internal conversion to the ground state and that the out-of-plane motion couples both states, as it has been already observed in other related systems.^{23,42–46}

Table 3: Lifetime broadening (LTB in cm^{-1}) of different transition of the Cyt-Na^+ and Cyt-K^+ complexes and minimum excited state lifetimes (τ_{min} in ps) for each transition.

Cyt-Na^+			Cyt-K^+		
mode	LBT (cm^{-1})	τ_{min} (ps)	mode	LTB (cm^{-1})	τ_{min} (ps)
0_0^0	4.5 ± 0.25	1.2	0_0^0	4 ± 0.25	1.3
ν_1 (49 cm^{-1})	11.9 ± 1	0.45	ν_1 (39 cm^{-1})	6.1 ± 0.25	0.8
ν_2 (122 cm^{-1})	6.8 ± 0.25	0.8	ν_2 (85 cm^{-1})	5.7 ± 0.25	0.8
$2 \nu_2$	6.7 ± 0.25	0.8	$\nu_1 + 2 \nu_2$	9 ± 0.5	0.6
ν_{12} (532 cm^{-1})	4.9 ± 0.25	1.1	ν_{12} (528 cm^{-1})	4.1 ± 0.25	1.3
$\nu_1 + \nu_{12}$	6.5 ± 2	0.8	$\nu_1 + \nu_{12}$	6.5 ± 0.25	0.8

The previous UV photofragmentation spectrum of the CytAg^+ complex was recorded in a MS2 scheme, the CytAg^+ complex being issued from the photodissociation of the Cyt_2Ag^+ complex at 273 nm .²⁵ In this work, the CytAg^+ is directly produced in the ESI source and is thus transferred in the cold QIT where cold ion spectroscopy is performed. (Figure SI 3). Interestingly, the two spectra do not significantly differ from each other. The onset of the photofragmentation for cold CytAg^+ complex is found at 31665 cm^{-1} (3.92 eV),⁴⁷ which is $\sim 0.4 \text{ eV}$ lower than the 0_0^0 bands of the first excited states of CytNa^+ and CytK^+ complexes. The adiabatic $\pi\pi^*$ transition of CytAg^+ lies in the range of $4.25\text{-}4.32 \text{ eV}$, i.e. in the same spectral region that the first excited $\pi\pi^*$ state of CytNa^+ and CytK^+ . This also suggests that the first excited states of the CytNa^+ and CytK^+ complexes studied in this work considerably differs from the first excited state of CytAg^+ .

A close inspection to Table 2 shows that the first excited state for the later complex is a CT ($n_O\sigma^*$) state, in which one electron from a lone pair orbital n_O of the O_7 atom of Cyt is promoted to the σ^* orbital localized on the Ag^+ moiety. On the contrary, the first excited state of $CytNa^+$ and $CytK^+$ is a $\pi\pi^*$ state, consequently the electron remains on the Cyt moiety. In these latter cases, the first CT involving the transfer of the electron from Cyt to Na^+/K^+ is the forth excited state ($\pi\sigma^*$) located 1.5 eV higher than the bright $\pi\pi^*$ state. The other CT states with $n_O\sigma^*$ or $n_N\sigma^*$ electronic configurations lie even higher in energy.

Also related to this fact, the other main difference between $CytNa^+$, $CytK^+$ and $CytAg^+$ complexes is the fragmentation pattern upon UV excitation. From the BO (4.3 eV) to the shortest excitation wavelength used in this work (210 nm, 5.9 eV, spectrum not shown), $CytNa^+$ and $CytK^+$ dissociate to Cyt and Na^+ or K^+ , respectively, while the $CytAg^+$ complex fragments to the CT dissociation channel Cyt^+ and Ag even at low excitation energy of 3.9 eV.

The dissociation energy threshold (E_{th}) for the CT dissociation channel can be estimated by Equation 2:

$$E_{th} \geq IP_{Cyt} - IP_M + E_b(S_0) \quad (2)$$

where IP_{Cyt} corresponds to the ionization potential of Cyt (8.73 eV),⁴⁸ IP_M is the ionization potentials of the metal atoms and $E_b(S_0)$ stands for the ground state binding energy of the complexes.^{25,32} All values are reported in Table 4 together with the calculated value of E_{th} .

As shown in Table 4, E_{th} for $CytNa^+$ and $CytK^+$ are approximately 2 eV larger than the corresponding value for $CytAg^+$ as a consequence of the lower IP of Na and K as compared to the IP of Ag. Then, the CT state is pushed up in the former complexes above the $\pi\pi^*$ state, to energies higher than the excitation energy region explored in this work. Therefore, not only the

first excited state of CytNa^+ and CytK^+ are not the same than in CytAg^+ , but also the excitation of the complexes in the 4.3 - 5.8 eV range is below the dissociation threshold for leading to Cyt^+ and M.²⁵

Table 4. Energy threshold for the CT fragmentation channel (E_{th}) for the CytM^+ (M = Ag, Na and K) complexes, calculated according to Eq. 2. The ionization potential of the M atoms (IP_M) and ground state binding energy $E_b(S_0)$ of the complexes are also shown.

Complex	IP_M (eV)	$E_b(S_0)$ (eV)	E_{th} (eV)
CytAg^+	7.58 ^a	2.86 ^b	4.01
CytNa^+	5.14 ^a	2.23 ^c	5.82
CytK^+	4.34 ^a	1.67 ^c	6.07

^a Ref.49 ^b Ref.25 ^c Ref.32

The energy threshold E_{th} leading the Cyt^+ ionic fragment in CytAg^+ (4.01 eV) is predicted in the same energy range yet slightly higher than the adiabatic excitation energy of the S_1 ($n\sigma^*$) calculated at 3.98 eV. The onset of the fragmentation spectrum of CytAg^+ is observed at 3.93 eV, with a long and bell-shape vibronic progression, which precludes the accurate determination of the 0_0^0 transition (see Fig SI4). The simulated Franck-Condon spectrum of the S_1 $n\sigma^*$ state of CytAg^+ is reported in Figure SI 4. Since the complex keeps the Cs planar structure in the optimized $n\sigma^*$ excited state, only the in-plane vibrational modes in combination with even quanta of out-of-plane bending modes are active. Such symmetry restriction leads to a rather simple vibronic spectrum in the first 500 cm^{-1} above the band origin, at odds with the experimental spectrum. A dense manifold of close lying vibronic transition as experimentally observed can be found at higher excitation energy, from 1000-1500 cm^{-1} above the band origin. This suggests that the band origin of the $n\sigma^*$ state might be located slightly below the energy E_{th} , impeding the dissociation of the complex near the band origin. Finally, the binding energy of

the complex is calculated at 2.86 eV,²⁵ but as already mentioned in ref 25, the ionic Ag⁺ fragment is barely detected even through excitation of the strong π - π^* transition. This implies that internal conversion from both $n\sigma^*$ and $\pi\pi^*$ excited states to the ground state has a very low quantum yield, otherwise silver cation would have been detected as photofragment.

CONCLUSIONS

The UV photofragmentation spectra of cold CytNa⁺ and CytK⁺ complexes were recorded and compared to the one of CytAg⁺. Both the spectroscopy and the photofragmentation channels differ for the alkali complexes, the only fragmentation channel being the alkali cation Na⁺/K⁺ without involvement of charge transfer from the cytosine to the metal cation. The first excited state for these alkali complexes is a $\pi\pi^*$ state. Ground and excited state optimizations at the SCS-CC2 level of theory were performed. The calculations show that in the first excited state of both complexes, the alkali cation lies out-of-plane of the cytosine which has the NH₂ group slightly twisted. Both the simulated Franck-Condon spectra and the adiabatic transition energies to the first excited state ($\pi\pi^*$) are in good agreement with the experimental values. Unlike CytAg⁺, the energy of the CT states involving the transfer of an electron from the Cyt moiety to M⁺ for both CytK⁺ and CytNa⁺ complexes were not experimentally observed, putting a lower limit of 5.9 eV for the first CT state. Therefore, unlike what happens with Ag⁺, the interaction of Cyt with these alkali metal cations (Na⁺ and K⁺) does not lead to the Cyt radical cation following photoexcitation at low energy UV photons. It is thus suggested that the alkali cations do not induce any photo damage process in DNA.

AUTHOR INFORMATION

Corresponding Author

*E-mail: gpino@fcq.unc.edu.ar; Tel: +54-351-5353866 ext. 53529

*E-mail: gilles.gregoire@u-psud.fr ; Tel: +33 169153103

Author Contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

ACKNOWLEDGMENTS

This work has been conducted within the International Associated Laboratory LEMIR (CNRS/CONICET) and was supported by CONICET, FONCyT, SeCyT-UNC and the ANR Research Grant (ANR2010BLANC040501-ESPEM). GAP thanks the Labex PALM (ANR-10-LABX-0039-PALM) for the invited Professor grant in 2019. We also acknowledge the use of the computing facilities Méso-LUM of the LUMAT federation (LUMAT FR 2764) and MAGI of University Paris 13.

REFERENCES

- (1) Schreier, W. J.; Gilch, P.; Zinth, W. Early Events of DNA Photodamage. *Annu. Rev. Phys. Chem.* **2015**, *66* (1), 497–519.
- (2) Pfeifer, G. P.; You, Y. H.; Besaratinia, A. Mutations Induced by Ultraviolet Light. *Mutat. Res. - Fundam. Mol. Mech. Mutagen.* **2005**, *571* (1-2 SPEC. ISS.), 19–31.
- (3) Bera, P. P.; Schaefer, H. F. (G-H)•-C and G-(C-H)• Radicals Derived from the Guanine-cytosine Base Pair Cause DNA Subunit Lesions. *Proc. Natl. Acad. Sci.* **2005**, *102* (19), 6698–6703.

- (4) Boudaïffa, Badia and Cloutier, Pierre and Hunting, Darel and Huels, Michael A. and Sanche, L. Resonant Formation of DNA Strand Breaks by Low-Energy (3 to 20 eV) Electrons. *Am. Assoc. Adv. Sci.* **2000**, 287 (5458), 1658--1660.
- (5) Taylor, J. S. Unraveling the Molecular Pathway from Sunlight to Skin Cancer. *Acc. Chem. Res.* **1994**, 27 (3), 76–82.
- (6) Setlow, P.; Li, L. Photochemistry and Photobiology of the Spore Photoproduct: A 50-Year Journey. *Photochem. Photobiol.* **2015**, 91 (6), 1263–1290.
- (7) Faraji, S.; Dreuw, A. Physicochemical Mechanism of Light-Driven DNA Repair by (6-4) Photolyases. *Annu. Rev. Phys. Chem.* **2013**, 65 (1), 275–292.
- (8) Ullrich, S.; Schultz, T.; Zgierski, M. Z.; Stolow, A. Electronic Relaxation Dynamics in DNA and RNA Bases Studied by Time-Resolved Photoelectron Spectroscopy. *Phys. Chem. Chem. Phys.* **2004**, 6 (10), 2796–2801.
- (9) Serrano-Andrés, L.; Merchán, M. Are the Five Natural DNA/RNA Base Monomers a Good Choice from Natural Selection?: A Photochemical Perspective. *J. Photochem. Photobiol. C Photochem. Rev.* **2009**, 10 (1), 21–32.
- (10) Barbatti, M.; Aquino, A. J. A.; Szymczak, J. J.; Nachtigallová, D.; Hobza, P.; Lischka, H. Relaxation Mechanisms of UV-Photoexcited DNA and RNA Nucleobases. *Proc. Natl. Acad. Sci.* **2010**, 107 (50), 21453–21458.
- (11) Middleton, C. T.; de La Harpe, K.; Su, C.; Law, Y. K.; Crespo-Hernández, C. E.; Kohler, B. DNA Excited-State Dynamics: From Single Bases to the Double Helix. *Annu. Rev. Phys. Chem.* **2009**, 60 (1), 217–239.

- (12) Improta, R.; Santoro, F.; Blancafort, L. Quantum Mechanical Studies on the Photophysics and the Photochemistry of Nucleic Acids and Nucleobases. *Chem. Rev.* **2016**, *116* (6), 3540–3593.
- (13) Berdakin, M.; Féraud, G.; Dedonder-Lardeux, C.; Jouvet, C.; Pino, G. A. Excited States of Protonated DNA/RNA Bases. *Phys. Chem. Chem. Phys.* **2014**, *16* (22), 10643–10650.
- (14) Kleineremanns, K.; Nachtigallová, D.; de Vries, M. S. Excited State Dynamics of DNA Bases. *Int. Rev. Phys. Chem.* **2013**, *32* (2), 308–342.
- (15) Boldissar, S.; de Vries, M. S. How Nature Covers Its Bases. *Phys. Chem. Chem. Phys.* **2018**, *20* (15), 9701–9716.
- (16) Gustavsson, T.; Improta, R.; Markovitsi, D. DNA/RNA: Building Blocks of Life Under UV Irradiation. *J. Phys. Chem. Lett.* **2010**, *1* (13), 2025–2030.
- (17) Lippert, B.; Gupta, D. Promotion of Rare Nucleobase Tautomers by Metal Binding. *Dalt. Trans.* **2009**, No. 24, 4619–4634.
- (18) Cristofari, C.; Rigo, R.; Greco, M. L.; Ghezzi, M.; Sissi, C. PH-Driven Conformational Switch between Non-Canonical DNA Structures in a C-Rich Domain of EGFR Promoter. *Sci. Rep.* **2019**, *9* (1), 1210.
- (19) Berdakin, M.; Steinmetz, V.; Maitre, P.; Pino, G. A. Gas Phase Structure of Metal Mediated (Cytosine)₂Ag⁺ Mimics the Hemiprotonated (Cytosine)₂H⁺ Dimer in i-Motif Folding. *J. Phys. Chem. A* **2014**, *118* (21), 3804–3809.

- (20) Berdakin, M.; Féraud, G.; Dedonder-Lardeux, C.; Jouvet, C.; Pino, G. A. Effect of Ag^+ on the Excited-State Properties of a Gas-Phase $(\text{Cytosine})_2\text{Ag}^+$ Complex: Electronic Transition and Estimated Lifetime. *J. Phys. Chem. Lett.* **2014**, 5 (13), 2295–2301.
- (21) Šponer, J.; Burda, J.; Leszczyński, J.; Šponer, J. Interaction of Metal Cations with Nucleic Acids and Their Building Units. In *Computational Studies of RNA and DNA*; 2006; pp 389–410.
- (22) Cruz-Ortiz, A. F.; Rossa, M.; Berthias, F.; Berdakin, M.; Maitre, P.; Pino, G. A. Fingerprints of Both Watson–Crick and Hoogsteen Isomers of the Isolated $(\text{Cytosine-Guanine})\text{H}^+$ Pair. *J. Phys. Chem. Lett.* **2017**, 8 (22), 5501–5506.
- (23) Broquier, M.; Soorkia, S.; Pino, G.; Dedonder-Lardeux, C.; Jouvet, C.; Grégoire, G. Excited State Dynamics of Cold Protonated Cytosine Tautomers: Characterization of Charge Transfer, Intersystem Crossing, and Internal Conversion Processes. *J. Phys. Chem. A* **2017**, 121 (34), 6429–6439.
- (24) Féraud, G.; Berdakin, M.; Dedonder-Lardeux, C.; Jouvet, C.; Pino, G. A. Excited States of Proton-Bound DNA/RNA Base Homodimers: Pyrimidines. *J. Phys. Chem. B* **2015**, 119 (6), 2219–2228.
- (25) Taccone, M. I.; Féraud, G.; Berdakin, M.; Dedonder-Lardeux, C.; Jouvet, C.; Pino, G. A. Communication: UV Photoionization of Cytosine Catalyzed by Ag^+ . *J. Chem. Phys.* **2015**, 143 (4), 41103.
- (26) Berdakin, M.; Féraud, G.; Dedonder-Lardeux, C.; Jouvet, C.; Pino, G. A. Effect of Ag^+ on the Excited-State Properties of a Gas-Phase $(\text{Cytosine})_2\text{Ag}^+$ Complex: Electronic Transition and Estimated Lifetime. *J. Phys. Chem. Lett.* **2014**, 5 (13), 2295–2301.

- (27) Bertran, J.; Blancafort, L.; Noguera, M.; Sodupe, M. Proton Transfer in DNA Base Pairs. In *Computational Studies of RNA and DNA*; Šponer, J., Lankaš, F., Eds.; Springer Netherlands: Dordrecht, 2006; pp 411–432.
- (28) Nir, E.; Kleinermanns, K.; de Vries, M. S. Pairing of Isolated Nucleic-Acid Bases in the Absence of the DNA Backbone . *Nature* **2000**, *408* (6815), 949–951.
- (29) Li, D.; Fedeles, B. I.; Singh, V.; Peng, C. S.; Silvestre, K. J.; Simi, A. K.; Simpson, J. H.; Tokmakoff, A.; Essigmann, J. M. Tautomerism Provides a Molecular Explanation for the Mutagenic Properties of the Anti-HIV Nucleoside 5-Aza-5,6-Dihydro-2'-Deoxycytidine. *Proc. Natl. Acad. Sci.* **2014**, *111* (32), E3252--E3259.
- (30) Wang, W.; Hellinga, H. W.; Beese, L. S. Structural Evidence for the Rare Tautomer Hypothesis of Spontaneous Mutagenesis. *Proc. Natl. Acad. Sci.* **2011**, *108* (43), 17644–17648.
- (31) Berdakin, M.; Steinmetz, V.; Maitre, P.; Pino, G. A. On the Ag⁺–Cytosine Interaction: The Effect of Microhydration Probed by IR Optical Spectroscopy and Density Functional Theory. *Phys. Chem. Chem. Phys.* **2015**, *17* (39), 25915–25924.
- (32) Yang, B.; Wu, R. R.; Polfer, N. C.; Berden, G.; Oomens, J.; Rodgers, M. T. IRMPD Action Spectroscopy of Alkali Metal Cation--Cytosine Complexes: Effects of Alkali Metal Cation Size on Gas Phase Conformation. *J. Am. Soc. Mass Spectrom.* **2013**, *24* (10), 1523–1533.
- (33) Broquier, M.; Soorkia, S.; Grégoire, G. A Comprehensive Study of Cold Protonated Tyramine: UV Photodissociation Experiments and Ab Initio Calculations. *Phys. Chem. Chem. Phys.* **2015**, *17* (39), 25854–25862.

(34) *TURBOMOLE V6.6*. a development of University of Karlsruhe and Forschungszentrum Karlsruhe GmbH, 1989-2007 TURBOMOLE GmbH, since 2007; available from <http://www.turbomole.com>.

(35) Grimme, S. Improved Second-Order Møller–Plesset Perturbation Theory by Separate Scaling of Parallel- and Antiparallel-Spin Pair Correlation Energies. *J. Chem. Phys.* **2003**, *118* (20), 9095–9102.

(36) Ahlrichs, R. Efficient Evaluation of Three-Center Two-Electron Integrals over Gaussian Functions. *Phys. Chem. Chem. Phys.* **2004**, *6* (22), 5119–5121.

(37) Weigend, F. Accurate Coulomb-Fitting Basis Sets for H to Rn. *Phys. Chem. Chem. Phys.* **2006**, *8* (9), 1057–1065.

(38) Hellweg, A.; Grün, S. A.; Hättig, C. Benchmarking the Performance of Spin-Component Scaled CC2 in Ground and Electronically Excited States. *Phys. Chem. Chem. Phys.* **2008**, *10* (28), 4119–4127.

(39) Western, C. M. PGOPHER: A Program for Simulating Rotational, Vibrational and Electronic Spectra. *J. Quant. Spectrosc. Radiat. Transf.* **2017**, *186*, 221–242.

(40) Yang, Z.; Rodgers, M. T. Tautomerization in the Formation and Collision-Induced Dissociation of Alkali Metal Cation-Cytosine Complexes. *Phys. Chem. Chem. Phys.* **2012**, *14* (13), 4517–4526.

(41) Nieuwjaer, N.; Desfrancois, C.; Lecomte, F.; Manil, B.; Soorkia, S.; Broquier, M.; Grégoire, G. Photodissociation Spectroscopy of Cold Protonated Synephrine: Surprising

Differences between IR–UV Hole-Burning and IR Photodissociation Spectroscopy of the O–H and N–H Modes. *J. Phys. Chem. A* **2018**, *122* (15), 3798–3804.

(42) Blancafort, L.; Cohen, B.; Hare, P. M.; Kohler, B.; Robb, M. A. Singlet Excited-State Dynamics of 5-Fluorocytosine and Cytosine: An Experimental and Computational Study. *J. Phys. Chem. A* **2005**, *109* (20), 4431–4436.

(43) Sobolewski, A. L.; Domcke, W. Photophysics of Malonaldehyde: An Ab Initio Study. *J. Phys. Chem. A* **1999**, *103* (23), 4494–4504.

(44) Sobolewski, A. L.; Domcke, W. On the Mechanism of Rapid Non-Radiative Decay in Intramolecularly Hydrogen-Bonded π Systems. *Chem. Phys. Lett.* **1999**, *300* (5), 533–539.

(45) Zgierski, M.; Patchkovskii, S.; Lim, E. Ab Initio Study of a Biradical Radiationless Decay Channel of the Lowest Excited Electronic State of Cytosine and Its Derivatives. *J. Chem. Phys.* **2005**, *123*, 81101.

(46) Merchán, M.; Serrano-Andrés, L. Ultrafast Internal Conversion of Excited Cytosine via the Lowest $\pi\pi^*$ Electronic Singlet State. *J. Am. Chem. Soc.* **2003**, *125* (27), 8108–8109.

(47) Taccone, M. I.; Féraud, G.; Berdakin, M.; Dedonder-Lardeux, C.; Jouvet, C.; Pino, G. A. Communication: UV Photoionization of Cytosine Catalyzed by Ag^+ . *J. Chem. Phys.* **2015**, *143* (4).

(48) Lobsiger, S.; Leutwyler, S. The Adiabatic Ionization Energy and Triplet T1 Energy of Jet-Cooled Keto-Amino Cytosine. *J. Phys. Chem. Lett.* **2012**, *3* (23), 3576–3580.

(49) *CRC Handbook of Chemistry and Physics, 2009–2010, 90th Ed.*; 2009; Vol. 131.

TOC Graphic: