

# Base-Mediated Generation of Ketenimines from Ynamides: Direct Access to Azetidinimines by an Imino Staudinger Synthesis

Eugénie Romero, Corinne Minard, Mohamed Benchekroun, Sandrine Ventre,

Pascal Retailleau, Robert H Dodd, Kevin Cariou

### ► To cite this version:

Eugénie Romero, Corinne Minard, Mohamed Benchekroun, Sandrine Ventre, Pascal Retailleau, et al.. Base-Mediated Generation of Ketenimines from Ynamides: Direct Access to Azetidinimines by an Imino Staudinger Synthesis. Chemistry - A European Journal, 2017, 23 (53), pp.12991-12994. 10.1002/chem.201702545. hal-02307210

# HAL Id: hal-02307210 https://hal.science/hal-02307210

Submitted on 7 Oct 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Base-Mediated Generation of Ketenimines from Ynamides: Direct Access to Azetidinimines by an Imino Staudinger Synthesis

Eugénie Romero,<sup>†</sup> Corinne Minard,<sup>†</sup> Mohamed Benchekroun,<sup>‡</sup> Sandrine Ventre,<sup>‡</sup> Pascal Retailleau, Robert H. Dodd,<sup>\*</sup> Kevin Cariou<sup>\*</sup>

**Abstract:** Ynamides were used as precursors for the *in situ* generation of highly reactive ketenimines which could be trapped with imines in a [2+2] cycloaddition. This imino Staudinger synthesis led to a variety of imino-analogs of  $\beta$ -lactams, namely azetidinimines (20 examples), that could be further functionalized through a broad range of transformations.

Ynamides (1) are versatile nitrogen-containing building blocks composed of an acetylene moiety directly attached to a nitrogen atom that generally bears at least one electron-withdrawing group that stabilizes the molecules.<sup>1</sup> The strong inherent polarization of the triple bond, that can be visualized by drawing the main mesomeric form 1', generally drives their reactivity towards nucleophilic addition on the  $\alpha$  position and reactions with electrophiles at the  $\beta$  position.<sup>1</sup> This trend can be further enhanced by acid activation, since protonation of the substrate yields keteniminium 2 (Scheme 1).<sup>2</sup>



Scheme 1. Ynamides: Inherent Polarization vs Acid and Base Activations

For our part, we have shown that using strongly basic conditions<sup>3</sup> an unprecedented nucleophilic addition of various heteroarenes at the  $\beta$  position could take place.<sup>4</sup> Moreover, depending on the substitution pattern of the nitrogen atom, the same basic conditions could lead to an  $\alpha$  nucleophilic addition, albeit with the loss of the electron-withdrawing group (EWG). In this latter case, base activation would trigger the cleavage of the EWG, leading to an amide that will eventually evolve towards ketenimine **3**.<sup>4,5</sup>

[a] Dr E. Romero, Dr C. Minard, Dr M. Benchekroun, Dr S.Ventre, Dr P. Retailleau, Dr R. H. Dodd,\* Dr K. Cariou\* Institut de Chimie des Substances Naturelles, CNRS UPR 2301, Université Paris-Sud, Université Paris-Saclay Avenue de la Terrasse, 91198 Gif-sur-Yvette, France E-mail: <u>kevin.cariou@cnrs.f</u> r

Supporting information for this article is given via a link at the end of the document.

In order to further expand the potential of this highly reactive intermediate (**3**), we sought to trap it in a [2+2] cycloaddition.<sup>6</sup> By analogy with the well-established Staudinger synthesis of  $\beta$ -lactams,<sup>7</sup> and based on the pioneering studies by Ghosez<sup>8</sup> and Regitz,<sup>9</sup> we envisioned that the *N*-phenylethenimine **3a** generated *in situ* from ynamide **1a** would be sufficiently electrophilic to react with imine **4** to ultimately give azetidinimine **5** (Scheme 2).<sup>10</sup> From our previous experience, an unsubstituted *N*,*N*-Boc-aryl-ynamide would be the best precursor for the generation of the ketenimine and we assumed an electron-rich nitrogen atom in the imine would favor the desired cycloaddition.



Scheme 2. In situ Generation of an Ethenimine and Trapping with an Imine

However, direct application of our previously disclosed reaction conditions<sup>4</sup> - i.e., 1.0 equiv. of ynamide, 1.0 equiv. of t-BuONa as base, DMF as solvent, with molecular sieves at room temperature - proved totally ineffective, even after prolonged reaction time (Table 1, entry 1). We reasoned that the cycloaddition process required thermal activation and thus, while traditional heating (entry 2) was unsuccessful, switching to microwave conditions<sup>11</sup> allowed the reaction to take place (entry 3). By replacing the sodium salt by potassium (entry 4) or lithium (entry 5) t-butanolate the desired azetidinimine 5aa could be isolated with 46% and 47% yield, respectively. Doubling the amount of both the ynamide and the base (X = 2) had little effect (entry 6) unless the temperature was raised to 100 °C (entry 7) which allowed 5aa to be isolated in 50% yield. This 50% threshold could be overcome by using silica gel as the additive instead of molecular sieves (entry 8, 54%). These optimized conditions were then applied to pchlorobenzaldehyde-derived imine 4b which led to the isolation of 64% of azetidinimine 5ab (entry 9). Improved formation of the latter was attempted by using a catalytic amount of a Lewis acid as the additive (entry 10, 50%) or as a co-additive (entry 11, 52%) but to no avail.

Having determined the optimal reaction conditions, we then explored the scope of this cycloaddition (Table 2). First, the influence of the substituents on the imine partners was assayed. Unsubstituted diphenyl aldimine **4c** allowed formation of the corresponding azetidinimine **5ac** in only 22% yield (entry 3).

|       | Boc<br>N-      |                    | Ar  | base, X equiv.<br>additive                   | Ar  |                             |                        |
|-------|----------------|--------------------|---|--|---|-----------------------------|------------------------|
|       | Ph<br>,<br>X e | <b>1a</b><br>quiv. | <b>4a</b> , Ar = Ph<br><b>4b</b> , Ar = <i>p</i> -ClC <sub>6</sub> H <sub>4</sub> | DMF Pi<br>time<br>heating                    | $h \sim N'$ PMP<br>5aa, Ar = Ph<br>5ab, Ar = $p$ -ClC <sub>6</sub> H <sub>4</sub> | PMP = <i>p</i> -methoxypher | nyl                    |
| Entry | Х              | 4                  | Base  | Additive                                     | Time  | Heating                     | Yield, 5               |
| 1     | 1              | 4a                 | <i>t</i> -BuONa   | M. S.  | 24 h  | RT                          | 0%                     |
| 2     | 1              | 4a                 | <i>t</i> -BuONa   | M. S.  | 24 h  | 60 °C                       | Traces                 |
| 3     | 1              | 4a                 | <i>t</i> -BuONa   | M. S.  | 1 h   | 80 °C μW                    | 20% conv.              |
| 4     | 1              | 4a                 | t-BuOK  | M. S.  | 1 h   | 80 °C μW                    | 46%, <b>5aa</b>        |
| 5     | 1              | 4a                 | t-BuOLi   | M. S.  | 1 h   | 80 °C μW                    | 47%, <b>5aa</b>        |
| 6     | 2              | 4a                 | <i>t</i> -BuOK  | M. S.  | 1 h   | 80 °C μW                    | 45%, <b>5aa</b>        |
| 7     | 2              | 4a                 | <i>t</i> -BuOLi   | M. S.  | 1 h   | 100 °C µW                   | 50%, <b>5aa</b>        |
| 8     | 2              | 4a                 | <i>t</i> -BuOLi   | SiO <sub>2</sub> (1 equiv.)                  | 1 h   | 100 °C μW                   | <mark>54%</mark> , 5aa |
| 9     | 2              | 4b                 | <i>t</i> -BuOLi   | SiO <sub>2</sub> (1 equiv.)                  | 1 h   | 100 °C μW                   | 64%, 5ab               |
| 10    | 2              | 4b                 | <i>t</i> -BuOLi <sup>[b]</sup>  | ZnOTf (10 mol%)                              | 1 h   | 100 °C μW                   | 50%, <b>5ab</b>        |
| 11    | 2              | 4b                 | <i>t</i> -BuOLi <sup>[b]</sup>  | SiO <sub>2</sub> (1 equiv.) + ZnOTf (10 mol% | 6) 1 h  | 100 °C μW                   | 52%, <b>5ab</b>        |

Table 1. Optimization of the [2+2] Cycloaddition between Ynamide 1a and Imines 4a,b.ª

[a] Isolated yields. [b] A 2.2 M solution of t-BuOLi in THF was used.

This result appears consistent with the positive effect of having an electron-donating group on the nitrogen (entries 1,2) and an electron-poor aryl group on the carbon of the imine partner (entry 2). Imines derived from *meta*- (4d) and *ortho*-anisidine (4e)

were then reacted with ynamide **1a**, providing cycloadducts **5ad** and **5ae** in 26% and 37% yields, respectively (entries 4,5).These yield variations can be accounted for by standard electronic (*meta* vs. *para/ortho*) and steric (*ortho* vs. *para*) effects.

 Table 2. Scope and limitations of the [2+2] Cycloaddition with Ynamides 1 and Imines 4. <sup>[a]</sup>

| a. Scop<br>Boc<br>N≡<br>Ar´<br>1, 2.0 e | e:<br>$= + \frac{Ar^{2}}{N} + \frac{r^{2}}{NAr^{3}} - \frac{t - BuOLi, 2.0}{SiO_{2}, 1.0}}{DMF, 1h, 100}$ quiv. 4 | b. Unreactive<br>equiv.<br>equiv.<br>$r \sim N$<br>$Ar^2$<br>$Ar^2$<br>$Br \sim N$<br>$Ar^3$<br>Ph<br>PG = Ts | $= Ynamides :$ $= Boc \\ N = Boc \\ R' Ph' $ $= R' \\ R' = Bn, allyl R' = Ph, n-Bu, TIPS$ |
|---|---|---|---|
| Entry                                   | Ar, <b>1</b>  | Ar <sup>2</sup> , Ar <sup>3</sup> , <b>4</b>  | Yield, <b>5</b> ª   |
| 1                                       | Ph, <b>1a</b>   | Ph, 4-MeO-C <sub>6</sub> H <sub>4</sub> , <b>4a</b>   | 54%, <b>5aa</b> <sup>[b] [c]</sup>  |
| 2                                       | Ph, <b>1a</b>   | 4-CI-C <sub>6</sub> H <sub>4</sub> , 4-MeO-C <sub>6</sub> H <sub>4</sub> , <b>4b</b>                          | 64%, <b>5ab</b> <sup>[b]</sup>  |
| 3                                       | Ph, <b>1a</b>   | Ph, Ph, <b>4c</b>   | 22%, <b>5ac</b> <sup>[d]</sup>  |
| 4                                       | Ph, <b>1a</b>   | 4-CI-C6H4, 3-MeO-C6H4, <b>4d</b>  | 26%, <b>5ad</b> <sup>[d]</sup>  |
| 5                                       | Ph, <b>1a</b>   | 4-CI-C <sub>6</sub> H <sub>4</sub> , 2-MeO-C <sub>6</sub> H <sub>4</sub> , <b>4e</b>                          | 37%, <b>5ae</b> <sup>[d]</sup>  |
| 6                                       | Ph, <b>1a</b>   | Ph, 3,4,5-triMeO-C <sub>6</sub> H <sub>4</sub> , <b>4f</b>  | 54%, <b>5af</b> <sup>[b]</sup>  |
| 7                                       | Ph, <b>1a</b>   | 4-MeO-C <sub>6</sub> H <sub>4</sub> , 4-MeO-C <sub>6</sub> H <sub>4</sub> , <b>4g</b>                         | 40%, <b>5ag</b> <sup>[b]</sup>  |
| 8                                       | Ph, <b>1a</b>   | 4-Br-C <sub>6</sub> H <sub>4</sub> , 4-MeO-C <sub>6</sub> H <sub>4</sub> , <b>4h</b>                          | 63%, <b>5ah</b>   |
| 9                                       | Ph, <b>1a</b>   | 4-I-C <sub>6</sub> H <sub>4</sub> , 4-MeO-C <sub>6</sub> H <sub>4</sub> , <b>4i</b>                           | 61%, <b>5ai</b> <sup>[c]</sup>  |
| 10                                      | Ph, <b>1a</b>   | 4-MeO <sub>2</sub> C-C <sub>6</sub> H <sub>4</sub> , Ph, <b>4j</b>  | 4%, <b>5aj</b> + 26%, <b>5aj'</b> <sup>[d]</sup>  |
| 11                                      | 4-CI-C <sub>6</sub> H <sub>4</sub> , <b>1b</b>  | Ph, 4-MeO-C <sub>6</sub> H₄, <b>4a</b>  | 10%, <b>5ba</b> <sup>[c]</sup>  |
| 12                                      | 4-CI-C <sub>6</sub> H <sub>4</sub> , <b>1b</b>  | 4-CI-C <sub>6</sub> H <sub>4</sub> , 4-MeO-C <sub>6</sub> H <sub>4</sub> , <b>4b</b>                          | 29%, <b>5bb</b>   |
| 13                                      | 4-Br-C <sub>6</sub> H <sub>4</sub> , <b>1c</b>  | Ph, 4-MeO-C <sub>6</sub> H₄, <b>4a</b>  | 11%, <b>5ca</b>   |
| 14                                      | 4-Br-C <sub>6</sub> H <sub>4</sub> , <b>1c</b>  | 4-CI-C <sub>6</sub> H <sub>4</sub> , 4-MeO-C <sub>6</sub> H <sub>4</sub> , <b>4b</b>                          | 40%, <b>5cb</b>   |
| 15                                      | 4-I-C <sub>6</sub> H <sub>4</sub> , 1d  | Ph, 4-MeO-C <sub>6</sub> H <sub>4</sub> , <b>4a</b>   | 10%, <b>5da</b>   |
| 16                                      | 4-I-C <sub>6</sub> H <sub>4</sub> , 1d  | 4-CI-C <sub>6</sub> H <sub>4</sub> , 4-MeO-C <sub>6</sub> H <sub>4</sub> , <b>4b</b>                          | 29%, <b>5db</b>   |
| 17                                      | 4-CF <sub>3</sub> -C <sub>6</sub> H <sub>4</sub> , <b>1e</b>  | Ph, 4-MeO-C <sub>6</sub> H <sub>4</sub> , <b>4a</b>   | 36%, <b>5ea</b>   |
| 18                                      | 4-CF <sub>3</sub> -C <sub>6</sub> H <sub>4</sub> , <b>1e</b>  | 4-CI-C <sub>6</sub> H <sub>4</sub> , 4-MeO-C <sub>6</sub> H <sub>4</sub> , <b>4b</b>                          | 36%, <b>5eb</b>   |
| 19                                      | 4-MeO-C <sub>6</sub> H <sub>4</sub> , 1f  | Ph, 4-MeO-C <sub>6</sub> H <sub>4</sub> , <b>4a</b>   | 22%, <b>5fa</b> <sup>[c]</sup>  |
| 20                                      | 4-MeO-C <sub>6</sub> H <sub>4</sub> , 1f  | 4-CI-C <sub>6</sub> H <sub>4</sub> , 4-MeO-C <sub>6</sub> H <sub>4</sub> , <b>4b</b>                          | 35%, <b>5fb</b>   |

[a] Isolated yields, using a 2.2 M solution of *t*-BuOLi in THF. [b] Solid *t*-BuOLi was used. [c] An X-ray structure was obtained, see Supporting Information for details.<sup>12</sup> [d] 10 mol% of ZnOTf<sub>2</sub> was used as the additive instead of silica gel.

Strongly electron-enriched 3,4,5-trimethoxylated imine 4f reacted smoothly with 1a to give 5af in 54% yield (entry 6) while dianisyl imine 4g led to 5ag in 40% yield (entry 7). In order to allow further functionalization, p-bromo- and p-iodobenzaldehydederived imines (4h and 4i) were subjected to the reaction conditions and the desired adducts were obtained in 63% and 61% yields, respectively (entries 8,9). Imine 4j bearing a methoxycarboxylate was then used, leading to an overall 30% yield of cycloaddition adduct 5a, most of which had been transesterified to the corresponding t-butoxycarboxylate 5aj' during the course of the reaction (entry 10). While various substitution patterns on the imine partner were well tolerated, this was not the case for the ketenimine precursor. Indeed, a series of ynamides was evaluated for which no cycloaddition took place (Table 2, topright b). These include: sulfonamides (GP = tosyl, mesyl and pnosyl) and N-alkyl (R = benzyl or allyl) as well as C-substituted (R' = phenyl, n-butyl and triisopropylsilyl) derivatives. The reaction did take place, however, when functionalized aryl groups such as para halo- (Cl, Br and I), trifluoromethyl- and methoxy-phenyl were incorporated on the ynamide partner (1b-f). All of these reacted with both imines 4a and 4b (entries11-20) providing yields of cycloadducts ranging from 10-36% with imine 4a (entries 11, 13, 15, 17 and 19) to 29-40% with imine 4b (entries 12, 14, 16, 18 and 20). Compared with ynamide 1a, yields were systematically lower and this can be attributed to a variety of factors difficult to unambiguously pinpoint since the substitution of this aryl presumably has an influence on both the generation of the active ketenimine and its subsequent stability and reactivity.

Having a wide panel of azetidinimines at our disposal we then turned our attention to their use for further functionalization. Satisfyingly, this unusual four-membered ring proved to be quite tolerant to a broad range of reaction conditions, obviating the necessity of extensive optimization. First, Suzuki coupling was used as a typical Pd-catalyzed cross-coupling reaction. Thus, an additional *p*-methoxyphenyl moiety could be added with excellent yields on both the *C*-aryl (6) or iminoaryl (7) part of the scaffold, using either bromo- (5ah & 5ca) or iodo-azetidinimines (5ai & 5da) (Scheme 3).



Scheme 3. Ynamides: Inherent Polarization vs Acid and Base Activations

A strong Lewis acid such as boron tribromide (Scheme 4, Eq. 1) could be employed, without any noticeable degradation of the amidine ring, to selectively cleave the methyl group of **5ab** to give phenolic azetidinimine **8**. The latter could in turn be alkylated with *t*-butyl bromoacetate to give **9** in 78% yield. Finally, treatment of carboxylate **5aj**' (Eq. 2) with a strong acid such as trifluoroacetic acid or a strong reducing agent such as lithium aluminum hydride gave respectively carboxylic acid **10** (96%) and primary alcohol **11** (82%), leaving the four-member ring untouched. Benzyl alcohol **11** could furthermore be transformed into the corresponding chloride **12** in 70% yield, using methanesulfonyl chloride.



Scheme 4. Further Functionalization around the Azetidinimine Ring

In conclusion, we have taken advantage a specific mode of activation of ynamides under basic conditions to generate *N*-arylethenimines from Boc-aryl-ynamides and react them with imines in a [2+2] cycloaddition. This reaction further ascertains the intermediacy of the ketenimine intermediate and paves the way for other types of cycloadditions. Moreover, the azetidinimines resulting from this imino Staudinger synthesis can undergo a broad array of chemical transformations that leave the 4-membered amidine ring intact. This work should allow for further exploration around this particular heterocyclic scaffold which can be viewed as an imino-analog of monocyclic  $\beta$ -lactams.<sup>13</sup> Studies in these directions are currently ongoing in our laboratory and will be reported in due course.

#### **Experimental Section**

Imine **4a** (63 mg, 0.3 mmol, 1.0 equiv.), ynamide **1a** (130 mg, 0.6 mmol, 2.0 equiv.) and SiO<sub>2</sub> (18 mg, 0.3 mmol, 1.0 equiv.) were successively added in a microwave sealable tube and placed under argon before the addition of *t*-BuOLi (48 mg, 0.6 mmol, 2.0 equiv.) followed by extra dry DMF (900  $\mu$ L). The sealed tube was placed in a microwave apparatus for 1 h at 100 °C. The crude material was purified by flash chromatography on silica gel (10% ethyl acetate in heptane as eluent) to provide azetidinimine **5aa** with 54 % yield (53 mg, yellow solid).

#### Acknowledgements

The authors thank CNRS, ICSN, Labex Lermit (ANR grant ANR-10-LABX-33 under the program Investissements d'Avenir ANR-11-IDEX-0003-01) and FCS Campus Saclay for financial support. E. R., C. M. and S. V thank Labex Lermit and M. B. thanks FCS Campus Saclay for fellowships.

#### **Author Contributions**

K. C. and R. H. D. designed and supervised the experiments and wrote the manuscript, C. M.<sup>†</sup> and S. V.<sup>‡</sup> optimized the reaction conditions and initiated the scope study, E. R.<sup>†</sup> and M. B.<sup>‡</sup> expanded the scope and performed the functionalization of the compounds. P. R. performed the X-ray crystallography experiments. † These authors contributed equally. ‡ These authors contributed equally.

**Keywords:** ynamides • ketenimines • [2+2] cycloaddition • azetidinimines • Staudinger synthesis

- For general reviews, see: a) G. Evano, A. Coste, K. Jouvin, Angew. Chem. 2010, 122, 2902.; Angew. Chem., Int. Ed. 2010, 49, 2840; b) K.
   A. DeKorver, H. Li, A. G. Lohse, R. Hayashi, Z. Lu, Y. Zhang, R. P. Hsung, Chem. Rev. 2010, 110, 5064; c) X.-N. Wang, H.-S. Yeom, L.-C. Fang, S. He, Z.-X. Ma, B. L. Kedrowski, R. P. Hsung, Acc. Chem. Res. 2014, 47, 560; d) G. Evano, C. Theunissen, M. Lecomte, Aldrichimica Acta 2015, 48, 59; e) A. M. Cook, C. Wolf, Tetrahedron Lett. 2015, 56, 2377; f) G. Evano, N. Blanchard, G. Compain, A. Coste, C. S. Demmer, W.Gati, C. Guissart, J. Heimburger, N. Henry, K. Jouvin, G. Karthikeyan, A. Loaouiti, M. Lecomte, A. Martin-Mingot, B. Metayer, A. Michelet, C. Theunissen, S. Thibaudeau, J. Wang, M. Zarca, C. Zhang, Chem. Lett. 2016, 45, 574.
- For recent examples, see: a) J. A. Mulder, R. P. Hsung, M. O. Frederick, [2] M. R. Tracev, C. Zificsak, Org. Lett. 2002. 4, 1383; b) J. A. Mulder, K. C. M. Kurtz, R. P. Hsung, H. Coverdale, M. O. Frederick, L. Shen, C. A. Zificsak, Org. Lett. 2003, 5, 1547; c) Y. Zhang, R. P. Hsung, X. Zhang, J. Huang, B. W. Slafer, A. Davis, Org. Lett. 2005, 7, 1047; d) Y. Zhang, Tetrahedron 2006, 62, 3917; e) A. H. Sato, K. Ohashi, T. Iwasawa, Tetrahedron Lett. 2013, 54, 1309; f) B. Peng, X. Huang, L.-G. Xie, N. Maulide, Angew. Chem. 2014, 126, 8893; Angew. Chem., Int. Ed. 2014, 53, 8718; g) C. Theunissen, B. Metayer, G. Compain, N. Henry, J. Marrot, A. Martin-Mingot, S. Thibaudeau, G. Evano, J. Am. Chem. Soc. 2014, 136, 12528; h) B. Metayer, G. Compain, K. Jouvin, A. Martin-Mingot, C. Bachmann, J. Marrot, G. Evano, S. Thibaudeau, J. Org. Chem. 2015, 80, 3397; i) L.-G. Xie, S. Niyomchon, A. J. Mota, L. Gonzalez, N. Maulide, Nat. Commun. 2016, 7, 10914; j) L.- G. Xie, S. Shaaban, X. Chen, N. Maulide, Angew. Chem. 2016, 128, 13056; Chem., Int. Ed. 2016, 55, 12864; k) V. Tona, S. A. Ruider, M. Berger, S. Shaaban, M. Padmanaban, L.-G. Xie, L. Gonzalez, N. Maulide, Chem. Sci. 2016, 7, 6032; I) M. Lecomte, G. Evano, Angew, Chem. 2016, 128, 4623; Angew, Chem., Int. Ed. 2016, 55, 4547; m) S. W. Kim, T.-W Um, S. Shin, Chem. Commun. 2017. 53. 2733.
- For a review specifically covering the anionic chemistry of ynamides, see:
   G. Evano, B. Michelet, C. Zhang, C. R. Chimie in press: http://dx.doi.org/10.1016/j.crci.2016.12.002.
- [4] A. Hentz, P. Retailleau, V. Gandon, K. Cariou, R. H. Dodd, Angew. Chem. 2014, 126, 8473; Angew. Chem., Int. Ed. 2014, 53, 8333.
- [5] For a general review on the chemistry of ketenimine see: a) P. Lu, Y. Wang, *Chem. Soc. Rev.* 2012, *41*, 5687. For other strategies to generate ketenimines from ynamides under basic conditions, see: b) A. Laouiti, F. Couty, J. Marrot, T. Boubaker, M. M. Rammah, M. B. Rammah, G. Evano, *Org. Lett.* 2014, *16*, 2252; c) Y. Kong, L. Yu, Y. Cui, J. Cao, *Synthesis*

2014, 183; d) L. Yu, J. Cao, *Org. Biomol. Chem.* 2014, *12*, 3986. Through a palladium-catalyzed allyl transfer process, see: e) Y. Zhang, K. A. DeKorver, A. G. Lohse, Y.-S. Zhang, J. Huang, R. P. Hsung, *Org. Lett.* 2009, *11*, 899. Through a thermic rearrangement, see: f) M. B. Hieu, M. D. E. Bolanos, F. Wudl, *Org. Lett.* 2005, *7*, 783.

- [6] For a survey of the use of ynamides in ring forming reactions, see: a) X.-N. Wang, H.-S. Yeom, L.-C. Fang, S. He, Z.-X. Ma, B. L. Kedrowski, R. P. Hsung, Acc. Chem. Res. 2014, 47, 560. For recent examples of the use of ynamides in [2+2] cycloaddition, see: b) S. F. Tlais, R. L. Danheiser, J. Am. Chem. Soc. 2014, 136, 15489; c) Y. Yuan, L. Bai, J. Nan, J. Liu, X. Luan, Org. Lett. 2014, 16, 4316; d) B. Alcaide, P. Almendros, C. Lázaro-Milla, Chem.-Eur. J. 2016, 22, 8998; e) X.-N. Wang, Z.-X. Ma, J. Deng, R. P. Hsung, Tetrahedron Lett. 2015, 56, 3463; f) K. Enomoto, H. Oyama, M. Nakada, Chem.-Eur. J. 2015, 21, 2798; g) L. Chen, J. Cao, Z. Xu, Z.-J. Zheng, Y.-M. Cui, L.-W. Xu, Chem. Commun. 2016, 52, 9574; h) C.-Z. Zhong, P.-T. Tung, T.-H. Chao, M.-C. P. Yeh, J. Org. Chem. 2017, 82, 481. For recent examples of the use of allenamides in [2+2] cycloaddition, see: R.-R. Liu, J.-P. Hu, J.-J. Hong, C.-J Lu, J.-R. Gao, Y.-X.Jia, Chem. Sci. 2017, 8, 2811 and references therein.
- a) F. P. Cossío, A. Arrieta, M. A. Sierra, *Acc. Chem. Res.* 2008, *41*, 925;
   b) G. S. Singh, S. Sudheesh, *ARKIVOC* 2014, 337; c) A. D. Allen, T. T. Tidwell, *Chem. Rev.* 2013, *113*, 7287; d) C. R. Pitts, T. Lectka, *Chem. Rev.* 2014, *114*, 7930.
- [8] a) M. De Poortere, J. Marchand-Brynaert,, L. Ghosez, Angew. Chem.
  1974, 86, 272; b) J. Marchand-Brynaert, M. Moya-Portuguez, D. Lesuisse, L. Ghosez, J. Chem.Soc., Chem. Commun. 1980, 173; c) A. Van Camp, D. Goossens, M. Moya-Portuguez, J. Marchand-Brynaert, L. Ghosez, Tetrahedron Lett. 1980, 21, 3081; d) J. Marchand-Brynaert, M. Moya-Portuguez, I. Huber, L. Ghosez, J. Chem.Soc., Chem. Commun. 1983, 818.
- a) B. Arnold, M. Regitz, Angew. Chem. 1979, 91, 337; Angew. Chem. Int. Ed. Eng. 1979, 18, 320; b) M. Regitz, B. Arnold, D. Danion, H. Schubert, G. Fusser, Bull. Soc. Chim. Belg. 1981, 90, 615.
- [10] For other methods to access azetidinimines, see: a) D. Mostowicz, W. Abramski, K. Piotowska, C. Belzecki, Pol. J. Chem. 1983, 57, 297; b) M. Alajarín, P. Molina, A. Vidal, Tetrahedron Lett. 1996, 37, 8945; c) M. Alaiarín, P. Molina, A. Vidal, F. Tovar, Tetrahedron 1997, 53, 13449; d) M. Alajarín, A. Vidal, F. Tovar, A. Arrieta, B. Lecea, F. P. Cossío, Chem. Eur. J. 1999, 5, 1106; e) F. P. Cossío, A. Arrieta, B. Lecea, M. Alajarín, A. Vidal, F. Tovar, J. Org. Chem. 2000, 65, 3633; f) M. Alajarín, A. Vidal, F. Tovar, M. C. R. de Arellano, F. P. Cossío, A. Arrieta, B. Lecea, J. Org. Chem. 2000, 65, 7512; g) B. Mendler, U.Kazmaier, V. Huch, M.Veith Org. Lett. 2005, 7, 2643; h) M. Whiting, V. V. Fokin, Angew. Chem. Int. Ed. 2006, 118, 3229; Angew. Chem. Int. Ed. 2006, 45, 3157; i) Tz. Gospodova, J. R. Rashkova, L. Z. Viteva, Bulg. Chem. Commun. 2008, 40, 568; i) S. Gouthaman, P. Shanmugam, A. B. Mandal, Tetrahedron Lett. 2013, 54, 3007; k) Y. Xing, H. Zhao, Q. Shang, J. Wang, P. Lu, Y. Wang, Org. Lett. 2013, 15, 2668.
- [11] See Supporting Information for details.
- [12] CCDC 1546588–1546592 (**5aa**, **5ba**, **5ai**, **5fa**, **6** respectively) contain the crystallographic data for this paper.
- [13] In the context of antibiotic resistance, monocyclic β-lactam antibiotics (monobactams) such as aztreonam hold a special position due to their unique properties, see: C. Ramsey, A. P. MacGowan, *J. Antimicrob. Chemother.* **2016**, *76*, 2704 and references therein.

### **Entry for the Table of Contents**

Layout 2:

### COMMUNICATION



**Base Jump:** Under strong basic conditions ynamides gave highly reactive ketenimines which could be trapped with imines in a [2+2] cycloaddition. This imino Staudinger synthesis led to a variety of imino-analogs of  $\beta$ -lactams, namely azetidinimines (20 examples), that could be further functionalized through a broad range of transformations.

Dr E. Romero, Dr C. Minard, Dr M. Benchekroun, Dr S. Ventre, Dr P. Retailleau, Dr R. H. Dodd, \* Dr K. Cariou\*

Page No. – Page No.

Base-Mediated Generation of Ketenimines from Ynamides: Direct Access to Azetidinimines by an Imino Staudinger Synthesis