Copper-Catalyzed Hydrocarboxylation of N -Allenyl Derivatives

Rémi Blieck, Marc Taillefer, Florian Monnier

► To cite this version:

Rémi Blieck, Marc Taillefer, Florian Monnier. Copper-Catalyzed Hydrocarboxylation of N -Allenyl Derivatives. *Journal of Organic Chemistry*, 2019, 84 (17), pp.11247-11252. 10.1021/acs.joc.9b01510 . hal-02307629

HAL Id: hal-02307629

<https://hal.science/hal-02307629>

Submitted on 7 Jan 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copper-Catalyzed Hydrocarboxylation of *N*-Allenyl Derivatives

Rémi Blicq,[†] Marc Taillefer,^{*,†} and Florian Monnier^{*,†,‡}

[†] Ecole Nationale Supérieure de Chimie de Montpellier, Institut Charles Gerhardt Montpellier UMR 5253 CNRS, AM2N, 8 rue de l'Ecole Normale, Montpellier 34296 Cedex 5, France.

[‡] Institut Universitaire de France, IUF, 1 rue Descartes, 75231 Paris cedex 5, France

Supporting Information Placeholder

ABSTRACT: The addition of carboxylic acids to allenes was performed with copper catalysis. This hydrocarboxylation reaction, occurring on the terminal carbon of the allenes, is totally regio- and stereoselective. It represents the first copper-catalyzed example of intermolecular C-O bond formation by allene hydrofunctionalization. This ligand-free system, is based on the use of catalytic amounts of copper combined with a base (10 mol % of K₂CO₃).

The direct and simple metal-catalyzed substitution of allylic compounds bearing leaving groups is a powerful method to obtain various functionalized allylic compounds.¹ Despite an impressive efficiency, the Tsuji-Trost reaction suffered from an atom-economical point of view, as one equivalent of leaving group is lost from the starting molecule. Therefore, alternatives appeared in the literature, and one of the most direct one is the hydrofunctionalization of allenes. For construction of C-N bonds, metal-catalyzed hydroamination of allenes were recently developed,² as for the formation of C-C bonds with various transition-metal-based catalysts.³ In the case of addition of O-nucleophiles as alcohols or phenols, few results were published these recent years.⁴ In the special case of the addition of carboxylic acids, few examples were reported with iridium-, rhodium-, silver- or palladium-catalysts.⁵⁻⁸ Depending on the nature of the metal, regioselectivity of the addition of the nucleophile varies as mentioned in Scheme 1. Systems based on Ir⁵ and on Rh⁶ respectively developed by Krische and Breit gave the branched allylic esters. On the other hand, Ag⁷ and Pd⁸-catalytic systems allow the selective formation of linear allylic molecules with the selective formation of *E* double bond. Nevertheless, these reported methodologies suffered from expensive and/or toxic catalytic systems.

Scheme 1. State of the Art: Metal-Catalyzed Addition of Carboxylic to Terminal Allenes

To pursue our studies on hydrofunctionalization of allenes catalyzed by simple and cheap catalyst based on copper,^{2g,3j,9} we engaged our efforts on the formation of C-O bonds, in the presence of copper salts playing the role of precatalyst, by the addition of carboxylic acids on terminal allenes.

Herein, we would like to report the first hydrocarboxylation of allenes catalyzed by a copper system. The latter leads selectively to the linear allylic esters in moderate to excellent yields (Scheme 2).

Scheme 2. Copper-Catalyzed Hydrocarboxylation of Terminal Allenes

First, a set of experiments were performed using allenamide **1** and benzoic acid **a** as model substrates. We then tested catalytic system developed by our group^{3j} for the addition of 1,3-dicarbonyl compounds on allenes (Table 1, entry 1). Only 13% of desired product **1a** was observed in the presence of 10% of Cu(CH₃CN)₄PF₆ and one equivalent of base Cs₂CO₃.

As mentioned in literature,⁵⁻⁷ base could be used in catalytic amount for the addition of carboxylic acid on allenes. Indeed, when we reduced the amount of Cs₂CO₃ to 10% (Table 1, entry 2) we observed an impressive increase of the formation of **1a** to 64%. Decreasing temperature from 50 °C to 25 °C affords to a loose of reactivity with only 15% of **1a** formed (Table 1, entry 3). Blank experiments revealed that combination of copper precatalyst and base is absolutely needed for the reaction. (Table 1, entries 5-6). In a second set of experiments, we showed that K₂CO₃ is the most suitable base to perform the hydrocarboxylation (Table 1, entries 4, 7-9). It is also important to note that others copper (I) precatalysts (CuI, CuBr, Cu₂O) did not allow better reactivity than 10% mol of Cu(CH₃CN)₄PF₆ (Table 1, entries 10-13). Thus the most suitable conditions to perform the hydrocarboxylation of **1** are those of the entry 9: one equivalent of allene **1** reacts with 1.2 equivalent of benzoic acid **a** in 1.5 mL of THF during 18 h at 50 °C in the presence of 10% of Cu(CH₃CN)₄PF₆ and 10% of K₂CO₃ acting as a base. These mild conditions allow the regio- and stereoselective formation of **1a** in 90% NMR yield.

Table 1 Hydrocarboxylation of 1 with Benzoic Acid a. Selected Data for Reaction Development^a

entry	[Cu] (x mol %)	Base (y eq.)	THF (mL)	Yield ^b (%)
1	Cu(CH ₃ CN) ₄ PF ₆ (10)	Cs ₂ CO ₃ (1)	2.5	13
2	Cu(CH ₃ CN) ₄ PF ₆ (10)	Cs ₂ CO ₃ (0.1)	2.5	64
3	Cu(CH ₃ CN) ₄ PF ₆ (10)	Cs ₂ CO ₃ (0.1)	2.5	15 ^c
4	Cu(CH ₃ CN) ₄ PF ₆ (10)	Cs ₂ CO ₃ (0.1)	1.5	80
5	Cu(CH ₃ CN) ₄ PF ₆ (10)	-	1.5	0
6	-	Cs ₂ CO ₃ (0.1)	1.5	0
7	Cu(CH ₃ CN) ₄ PF ₆ (10)	KOH (0.1)	1.5	85
8	Cu(CH ₃ CN) ₄ PF ₆ (10)	NaOH (0.1)	1.5	82
9	Cu(CH ₃ CN) ₄ PF ₆ (10)	K ₂ CO ₃ (0.1)	1.5	90
10	CuI (10)	K ₂ CO ₃ (0.1)	1.5	32
11	CuBr (10)	K ₂ CO ₃ (0.1)	1.5	43
12	Cu ₂ O (10)	K ₂ CO ₃ (0.1)	1.5	21
13	Cu(CH ₃ CN) ₄ PF ₆ (5)	K ₂ CO ₃ (0.1)	1.5	52

^a Reaction conditions: **1** (0.5 mmol), **a** (0.6 mmol), base (0.05 to 0.5 mmol) and catalyst (0.025 to 0.05 mmol) were placed in a screw tube under argon in THF for 18 h at 50 °C. ^b NMR yields using 1,3,5-trimethoxybenzene as internal standard. ^c At 25 °C.

Next, investigations on the scope and limitations of reaction were engaged. In a first set of experiments, we wanted to know if this selective reaction is tolerant toward various carboxylic acids (Scheme 3). Moderate to excellent yields were obtained when *N*-allenyl-2-pyrrolidinone **1** reacted with aromatic carboxylic acids such as benzoic acid **a**, 4-fluorophenylacetic acid **b**, *m*-methylbenzoic acid **c**, *o*-methylbenzoic acid **d** (Scheme 3). We also demonstrated that *N*-allenyl-2-pyrrolidinone **1** allowed the formation of the desired carboxylated product **1e** in 75% yield, when engaged with carboxylic

acid bearing heterocycle as 2-furoic acid **e**. To our delight, we also engaged successfully *N*-protected amino acids as source of carboxylic acids. To this purpose, boc-proline **f** and boc-β-alanine **g** reacted with allene **1** under standard conditions to give desired allylic esters compounds **1f** and **1g** respectively in 80% and 72% yields. Finally, cinnamic acid **h** with allene **1** gave the desired molecule **1h** in 72% yield. All products **1a-1h** were obtained in regio- and stereoselective manner without the generation of any other products or isomers.

Scheme 3. Copper-Catalyzed Hydrocarboxylation of Allene 1 with Various Carboxylic Acids a-j (Isolated Yields)

In a following set of experiments, we showed that diverse allenamides could be suitable substrates for this selective reaction (Scheme 4). *N*-allenylloxazolidinone **2** reacted with benzoic acid **a** to give **2a** in 65% yield. Other important class of allenes such as *N*-allenyl-sulfonamide was engaged successfully, and under standard reaction conditions we were able to obtain **3a**, **4a** and **5a** respectively in 43%, 51% and 47% yield. Finally, *N*-allenyl-azoles such as allenes bearing triazole **6** and benzotriazole **7** were also good candidates for this hydrofunctionalization reaction in the presence of benzoic acid **a**.

Scheme 4. Copper-Catalyzed Hydrocarboxylation of Various *N*-allenyl Derivatives 2-7 with Benzoic Acid a (Isolated Yields)

Considering our previous reports on hydroamination of allenes catalyzed by similar catalytic system,^{9a-b} we propose a mechanism involving cationic copper species, which in a first step could complex the allene to afford intermediate **A** (Scheme 5). The in-situ generated carboxylate anion $R^1CO_2^-$ could attack **A** in an antiperiplanar manner to afford a *Z* allenylcopper intermediate **B**. Another equivalent of carboxylic acid could perform a protodemetalation through intermediate **C**, which could release the desired *E* allylic ester and regenerate carboxylate anion $R^1CO_2^-$. The latter could be re-engaged in the catalytic cycle, thus explaining the use of only catalytic amount of base in this reaction. Work is in progress to assess the soundness of this mechanistic proposal.

Scheme 5. Proposed Mechanism for the Copper-Catalyzed Hydrocarboxylation of Terminal Allenes

In conclusion we have reported the first copper-catalyzed C-O bond formation performed by hydrofunctionalization of allenes, giving an efficient and selective access to allylic moieties with a total atom economy. Under mild conditions, using catalytic amount of base, we explored the powerfulness and the tolerance of this novel catalytic system for the regio- and stereoselective generation of allylic esters starting from simple substrates, and we demonstrated the high flexibility of copper-based catalytic systems in the functionalization of allenes.

EXPERIMENTAL SECTION

General considerations

All reactions were performed in schlenk tube closed with a Rodaviss® screw cap and PTFE liner (approx 25mL), using a freshly anhydrous solvents obtained from a Mbraun Pure Solv® solvent purification system. Copper salts were purchased from either Sigma-Aldrich®, Strem®, Alfa Aesar® or Acros Organic®, used as received, stored in a dessicator cabinet and weighed to air. $[Cu(NCMe)_4]PF_6$ was synthesized according to a previously reported method.¹⁰ Cinnamic acid was obtained by saponification of the corresponding ester.¹¹ Other reagents were purchased from either Sigma-Aldrich® or Alfa Aesar® or Acros Organic® and used as received. 1H and ^{13}C NMR spectra were recorded with a Bruker AC-400 MHz spectrometer in $CDCl_3$. For 1H NMR (400 MHz), $CHCl_3$ and TMS served as internal standards (δ = 7.27, 2.50 and 0 ppm) and data are reported as follows: chemical shift (in ppm), multiplicity (s = singlet, d = doublet, t = triplet, m = multiplet), coupling constant (in Hz), and integration. For ^{13}C NMR (100 MHz), $CHCl_3$ was used as internal standards (δ = 77.2) and spectra were obtained with complete proton decoupling. Gas chromatography–mass spectra (GC-MS) were recorded on a Shimadzu QP2012-SE with a Zebron ZB-5MS (20m $\times$ 0.18mm), capillary apolar column (Stationary phase: 0.18 μ m film). GC-MS method: Initial temperature: 50 °C; Initial time: 2 min; Ramp: 22 °C/min; Final Temperature: 280 °C; Final time: 15 min or 23min. Melting point were measured with a Stuart SMP50 Automatic Melting Point apparatus. All allenes were synthesized according to previously described methods and analysis were in accordance with the literature.⁹

General Procedure A for hydrocarboxylation of allenes

A Schlenk flask of appropriate size was charged with $Cu(NCMe)_4PF_6$ (0.10 eq), carboxylic acid (1.2 eq) and base (0.10 eq) and closed with a rubber septum. After evacuation and back-filling with argon three times, dry THF (3 mL per mmol of allene) and allene (1.0 equiv) were sequentially added. The vessel was closed and stirred at 50 °C, unless otherwise stated, until complete conversion. After allowing the reaction to cool to room temperature, 1,3,5-trimethoxybenzene (0.33 eq) dissolved in dichloromethane was added as internal standard. 10 volumes of dichloromethane were then added and the mixture was partitioned with 3 volumes of water. The aqueous phase was extracted with dichloromethane (3 $\times$ 3 volumes) and the combined organic phases washed with saturated NaCl aqueous solution, then dried over $MgSO_4$. Volatiles were evaporated under reduced pressure and the residue was purified by flash chromatography on Et_3N -treated silica gel. In order to avoid any decomposition, the silica gel was treated with three volumes of trimethylamine solution (10 vol% in the selected eluent) and then rinsed with five volumes of fresh eluent.

(*E*)-3-(2-oxopyrrolidin-1-yl)allyl benzoate (**1a**). The reaction of *N*-allenylpyrrolidin-2-one **1** with benzoic acid according to the general procedure A (flash chromatography eluent: Pentane/AcOEt 80:20) gave the title compound in the form of a colorless oil (80 %). 1H NMR (400 MHz, $CDCl_3$) δ 8.05 – 8.02 (m, 2H), 7.55 – 7.53 (m, 1H), 7.45 – 7.41 (m, 2H), 7.26 (d, J = 14.4 Hz, 2H), 5.20 (dt, J = 14.4, 7.3 Hz, 1H), 4.85 (dd, J = 7.3, 1.0 Hz, 2H), 3.56 – 3.53 (m, 2H), 2.50 (t, J = 8.1 Hz, 2H), 2.17 – 2.10 (m, 2H). ^{13}C { 1H } NMR (101 MHz, $CDCl_3$) δ 173.7, 166.6, 133.0, 130.3, 129.7, 129.1, 128.4, 104.9, 64.3, 45.1, 31.2, 17.5. Due to the limited stability of this compound in the analysis conditions, no meaningful HRMS data (either EI or ESI) could be obtained. However, comparison of spectral data with analogous compounds that could be fully characterized allowed establishing the structure beyond reasonable doubt.

(*E*)-3-(2-oxopyrrolidin-1-yl)allyl 2-(4-fluorophenyl)acetate (**1b**). The reaction of *N*-allenylpyrrolidin-2-one **1** with 4-fluoro-phenylacetic acid **b** according to the general procedure A (flash chromatography eluent: Cyclohexane/AcOEt 60:40) gave the title compound in the form of a colorless oil (81 %). ¹H NMR (400 MHz, CDCl₃) δ 7.26–7.21 (m, 2H), 7.16 (d, *J* = 14.4 Hz, 1H), 7.04–6.97 (m, 2H), 5.05 (dt, *J* = 14.4, 7.2 Hz, 1H), 4.62 (dd, *J* = 7.2, 1.0 Hz, 2H), 3.59 (s, 2H), 3.50 (t, *J* = 8.0 Hz, 2H), 2.49 (t, *J* = 8.1 Hz, 2H), 2.17–2.06 (m, 2H). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 173.6, 171.1, 162.6 (d, *J* = 228 Hz), 131.0, 130.9, 129.2, 115.7, 115.5, 104.6, 64.4, 45.2, 40.6, 31.2, 17.6. HRMS (ESI-TOF) *m/z*: [M + Na]⁺ Calcd for C₁₅H₁₆FNO₃Na 300.1012; Found 300.1012.

(*E*)-3-(2-oxopyrrolidin-1-yl)allyl 3-methylbenzoate (**1c**). The reaction of *N*-allenylpyrrolidin-2-one **1** with *m*-methylbenzoic acid **c** according to the general procedure A (flash chromatography eluent: Pentane/AcOEt 70:30) gave the title compound in the form of a white solid (61 %). ¹H NMR (400 MHz, CDCl₃) δ 7.85–7.82 (m, 2H), 7.37–7.29 (m, 2H), 7.26 (d, *J* = 14.4 Hz, 1H), 5.20 (dt, *J* = 14.4, 7.3 Hz, 1H), 4.84 (dd, *J* = 7.3, 1.0 Hz, 2H), 3.56–3.53 (m, 2H), 2.50 (dd, *J* = 8.6, 7.7 Hz, 2H), 2.39 (s, 3H), 2.14–2.08 (m, 2H). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 173.6, 166.7, 138.2, 133.7, 130.2, 130.2, 129.0, 128.3, 126.8, 104.9, 64.2, 45.1, 31.1, 21.3, 17.5. HRMS (ESI-TOF) *m/z*: [M + Na]⁺ Calcd for C₁₅H₁₇NO₃Na 282.1106; Found 282.1107. mp: 75–76 °C

(*E*)-3-(2-oxopyrrolidin-1-yl)allyl 2-methylbenzoate (**1d**). The reaction of *N*-allenylpyrrolidin-2-one **1** with *o*-methylbenzoic acid **d** according to the general procedure A (flash chromatography eluent: Pentane/AcOEt 70:30) gave the title compound in the form of a white solid (59 %). ¹H NMR (400 MHz, CDCl₃) δ 7.89 (dd, *J* = 8.1, 1.4 Hz, 1H), 7.37 (td, *J* = 7.4, 1.3 Hz, 1H), 7.29–7.17 (m, 3H), 5.19 (dt, *J* = 14.4, 7.3 Hz, 1H), 4.82 (dd, *J* = 7.3, 1.0 Hz, 2H), 3.57–3.49 (m, 2H), 2.58 (s, 3H), 2.49 (t, *J* = 8.2 Hz, 2H), 2.17–2.04 (m, 2H). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 173.6, 167.5, 140.3, 132.1, 131.8, 130.7, 129.7, 129.0, 125.8, 105.0, 64.1, 45.2, 31.2, 21.8, 17.5. HRMS (ESI-TOF) *m/z*: [M + Na]⁺ Calcd for C₁₅H₁₇NO₃Na 282.1106; Found 282.1109. mp: 72–73 °C

(*E*)-3-(2-oxopyrrolidin-1-yl)allyl furan-2-carboxylate (**1e**). The reaction of *N*-allenylpyrrolidin-2-one **1** with 2-furoic acid **e** according to the general procedure A (flash chromatography eluent: Pentane/AcOEt 60:40) gave the title compound in the form of an orange oil (75 %). ¹H NMR (400 MHz, CDCl₃) δ 7.56 (dd, *J* = 1.7, 0.9 Hz, 1H), 7.23 (d, *J* = 14.4 Hz, 1H), 7.17 (dd, *J* = 3.5, 0.9 Hz, 1H), 6.49 (dd, *J* = 3.5, 1.8 Hz, 1H), 5.15 (dt, *J* = 14.4, 7.4 Hz, 1H), 4.82 (dd, *J* = 7.4, 1.0 Hz, 2H), 3.57–3.48 (m, 2H), 2.48 (t, *J* = 8.1 Hz, 2H), 2.17–2.04 (m, 2H). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 173.7, 158.7, 146.4, 144.7, 129.6, 118.2, 112.0, 104.4, 64.4, 45.1, 31.2, 17.6. HRMS (ESI-TOF) *m/z*: [M + H]⁺ Calcd for C₁₂H₁₄NO₄ 236.0923, found 236.0925.

(*E*)-1-(*tert*-butyl) 2-(3-(2-oxopyrrolidin-1-yl)allyl) pyrrolidine-1,2-dicarboxylate (**1f**). The reaction of *N*-allenylpyrrolidin-2-one **1** with Boc-proline **f** according to the general procedure A (flash chromatography eluent: Pentane/AcOEt 50:50) gave the title compound in the form of a colorless oil (80 %). ¹H NMR (400 MHz, CDCl₃) δ 7.20–7.11 (m, 1H), 5.12–5.00 (m, 1H), 4.63 (t, *J* = 8.0 Hz, 2H), 4.33–4.16 (m, 1H), 3.58–3.31 (m, 4H), 2.51–2.46 (m, 2H), 2.28–2.05 (m, 3H), 2.00–1.79 (m, 3H), 1.42 (d, *J* = 20.7 Hz, 9H). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 173.6, 172.2, 129.3, 128.9, 104.9, 104.4, 80.0, 64.4, 64.3, 59.3, 59.0, 46.5, 45.1, 31.2, 31.0, 30.0, 28.6, 28.5, 24.5, 23.8, 17.6. HRMS (ESI-TOF) *m/z*: [M + Na]⁺ Calcd for C₁₇H₂₆N₂O₅Na 361.1739; Found 361.1739.

(*E*)-3-(2-oxopyrrolidin-1-yl)allyl 3-((*tert*-butoxycarbonyl)amino)propanoate (**1g**). The reaction of *N*-allenylpyrrolidin-2-one **1** with Boc-β-alanine **g** according to the

general procedure A (flash chromatography eluent: Pentane/AcOEt 50:50) gave the title compound in the form of a white solid (72 %). ¹H NMR (400 MHz, CDCl₃) δ 7.11 (d, *J* = 14.4 Hz, 1H), 5.08–4.96 (m, 2H), 5.01 (dt, *J* = 14.5, 7.3 Hz, 1H), 4.57 (dd, *J* = 7.3, 0.9 Hz, 2H), 3.52–3.43 (m, 2H), 3.35–3.31 (m, 2H), 2.51–2.40 (m, 4H), 2.14–2.01 (m, 2H), 1.37 (s, 9H). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 173.6, 172.3, 155.8, 129.1, 104.4, 64.0, 45.1, 36.1, 34.7, 31.1, 28.4, 17.5. HRMS (ESI-TOF) *m/z*: [M + Na]⁺ Calcd for C₁₅H₂₄N₂O₅Na 335.1583; Found 335.1584. mp: 67–68 °C.

(*E*)-3-(2-oxopyrrolidin-1-yl)allyl cinnamate (**1h**). The reaction of *N*-allenylpyrrolidin-2-one **1** with cinnamic acid **h** according to the general procedure A (flash chromatography eluent: Pentane/AcOEt 70:30) gave the title compound in the form of a white solid (72 %). ¹H NMR (400 MHz, CDCl₃) δ 7.68 (d, *J* = 16.0 Hz, 1H), 7.55–7.47 (m, 2H), 7.44–7.33 (m, 3H), 7.22 (d, *J* = 14.4 Hz, 1H), 6.43 (dd, *J* = 16.0, 0.6 Hz, 1H), 5.14 (dt, *J* = 14.4, 7.2 Hz, 1H), 4.73 (d, *J* = 7.3 Hz, 2H), 3.53 (t, *J* = 7.2 Hz, 2H), 2.49 (t, *J* = 8.2 Hz, 2H), 2.17–2.05 (m, 2H). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 173.6, 166.9, 145.1, 134.5, 130.4, 129.0, 128.2, 118.1, 104.9, 63.9, 45.2, 31.2, 17.6. HRMS (ESI-TOF) *m/z*: [M + Na]⁺ Calcd for C₁₆H₁₇NO₃Na 294.1106; Found 294.1106. mp: 92–93 °C.

(*E*)-3-(2-oxopyrrolidin-1-yl)allyl 4-iodobenzoate (**1i**). The reaction of *N*-allenylpyrrolidin-2-one **1** with *p*-iodobenzoic acid **i** according to the general procedure A (flash chromatography eluent: Cyclohexane/AcOEt 60:40) gave the title compound in the form of a white solid (71 %). ¹H NMR (400 MHz, CDCl₃) δ 7.85–7.75 (m, 2H), 7.75–7.69 (m, 2H), 7.25 (d, *J* = 14.3 Hz, 1H), 5.17 (dt, *J* = 14.3, 7.3 Hz, 1H), 4.83 (dd, *J* = 7.3, 0.7 Hz, 2H), 3.59–3.49 (m, 2H), 2.50 (t, *J* = 8.1 Hz, 2H), 2.20–2.05 (m, 2H). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 173.7, 166.1, 137.8, 131.2, 129.8, 129.4, 104.6, 100.9, 64.6, 45.2, 31.2, 17.6. mp: 127–128 °C. Due to the limited stability of this compound in the analysis conditions, no meaningful HRMS data (either EI or ESI) could be obtained. However, comparison of spectral data with analogous compounds that could be fully characterized allowed establishing the structure beyond reasonable doubt.

(*E*)-3-(2-oxopyrrolidin-1-yl)allyl 2-phenylacetate (**1j**). The reaction of *N*-allenylpyrrolidin-2-one **1** with phenylacetic acid **j** according to the general procedure A (flash chromatography eluent: Cyclohexane/AcOEt 60:40) gave the title compound in the form of a colorless oil (81 %). ¹H NMR (400 MHz, CDCl₃) δ 7.29–7.13 (m, 5H), 7.07 (d, *J* = 14.4 Hz, 1H), 4.98 (dt, *J* = 14.4, 7.2 Hz, 1H), 4.54 (dd, *J* = 7.2, 0.8 Hz, 2H), 3.54 (s, *J* = 3.2 Hz, 2H), 3.45–3.36 (m, 2H), 2.40 (t, *J* = 8.2 Hz, 2H), 2.08–1.95 (m, 2H). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 173.6, 171.5, 133.9, 129.3, 128.9, 128.6, 127.1, 104.7, 64.2, 45.1, 41.4, 31.1, 17.4. Due to the limited stability of this compound in the analysis conditions, no meaningful HRMS data (either EI or ESI) could be obtained. However, comparison of spectral data with analogous compounds that could be fully characterized allowed establishing the structure beyond reasonable doubt.

(*E*)-3-(2-oxooxazolidin-3-yl)allyl benzoate (**2a**). The reaction of *N*-allenylloxazolidinone **2** with benzoic acid **a** according to the general procedure A (flash chromatography eluent: Pentane/AcOEt 80:20 to 60:40) gave the title compound in the form of a white solid (65 %). ¹H NMR (400 MHz, CDCl₃) δ 8.05–8.03 (m, 2H), 7.56–7.54 (m, 1H), 7.44 (ddd, *J* = 8.0, 6.7, 1.1 Hz, 2H), 7.07 (d, *J* = 14.2 Hz, 1H), 5.11 (dt, *J* = 14.3, 7.3 Hz, 1H), 4.85 (dd, *J* = 7.3, 0.9 Hz, 2H), 4.49–4.45 (m, 2H), 3.77–3.73 (m, 2H). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 167.4, 156.2, 134.0, 131.1, 130.5, 130.5, 129.3, 105.1, 64.7, 63.2, 43.3. HRMS (ESI-TOF) *m/z*: [M + Na]⁺ Calcd for C₁₃H₁₃N₂O₄Na 270.0742; Found 270.0744. mp: 104–105 °C

(*E*)-3-((4-methyl-*N*-phenylphenyl)sulfonamido)allyl benzoate (**3a**). The reaction of 4-methyl-*N*-phenyl-*N*-(propa-1,2-

dien-1-yl)benzenesulfonamide **3** with benzoic acid according to the general procedure A (flash chromatography eluent: Cyclohexane/AcOEt 90:10) gave the title compound in the form of an orange solid (43 %). ^1H NMR (400 MHz, CDCl_3) δ 8.03 – 7.95 (m, 2H), 7.59 – 7.50 (m, 3H), 7.46 – 7.33 (m, 6H), 7.28 – 7.25 (m, 2H), 7.01 – 6.94 (m, 2H), 4.74 (dd, $J = 7.1$, 0.9 Hz, 2H), 4.58 (ddd, $J = 13.9$, 7.7, 6.9 Hz, 1H), 2.43 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) δ 166.5, 144.3, 136.1, 135.7, 134.8, 133.1, 130.4, 130.3, 129.8, 129.8, 129.7, 129.4, 128.5, 127.7, 104.4, 64.0, 21.8. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{23}\text{H}_{22}\text{NO}_4\text{S}$ 408.1270, found 408.1271. mp: 126–127 °C

(*E*)-3-(*N*-(4-methoxyphenyl)methylsulfonamido)allyl benzoate (**4a**). The reaction of *N*-(4-methoxyphenyl)-*N*-(propa-1,2-dien-1-yl)methanesulfonamide **4** with benzoic acid according to the general procedure A (flash chromatography eluent: Cyclohexane/AcOEt 80:20) gave the title compound in the form of a white solid (51 %). ^1H NMR (400 MHz, CDCl_3) δ 8.04 – 7.97 (m, 2H), 7.59 – 7.51 (m, 1H), 7.47 – 7.38 (m, 2H), 7.29 – 7.22 (m, 3H), 7.02 – 6.95 (m, 2H), 4.78 – 4.73 (m, 2H), 4.69 (ddd, $J = 13.3$, 8.4, 6.0 Hz, 1H), 3.84 (s, 3H), 3.03 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) δ 166.6, 160.4, 134.5, 133.1, 131.2, 130.4, 130.3, 129.7, 128.6, 128.5, 128.2, 115.5, 104.0, 64.0, 55.7, 39.3. HRMS (ESI-TOF) m/z : $[\text{M} + \text{Na}]^+$ Calcd for $\text{C}_{18}\text{H}_{19}\text{N}_1\text{O}_5\text{S}_1\text{Na}$ 384.0882; Found 384.0883. mp: 139–140 °C

(*E*)-3-((*N*-allyl-4-methylphenyl)sulfonamido)allyl benzoate (**5a**). The reaction of *N*-allyl-4-methyl-*N*-(propa-1,2-dien-1-yl)benzenesulfonamide **5** with benzoic acid according to the general procedure A (flash chromatography eluent: Cyclohexane/AcOEt 90:10) gave the title compound in the form of a white oil (47 %). ^1H NMR (400 MHz, CDCl_3) δ 8.06 – 7.98 (m, 2H), 7.72 – 7.65 (m, 2H), 7.60 – 7.51 (m, 1H), 7.48 – 7.36 (m, 2H), 7.31 – 7.27 (m, 2H), 7.11 (d, $J = 14.1$ Hz, 1H), 5.63 (ddt, $J = 17.2$, 10.5, 5.3 Hz, 1H), 5.22 – 5.12 (m, 2H), 5.05 (dt, $J = 14.2$, 7.3 Hz, 1H), 4.78 (dd, $J = 7.3$, 0.8 Hz, 2H), 4.03 (dt, $J = 5.3$, 1.6 Hz, 2H), 2.41 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) δ 166.6, 144.2, 136.2, 133.0, 132.2, 131.3, 130.5, 130.0, 129.7, 128.5, 127.2, 118.3, 103.9, 64.4, 48.3, 21.7. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{20}\text{H}_{22}\text{N}_1\text{O}_4\text{S}_1$ 372.1270, found 372.1271.

(*E*)-3-(1*H*-1,2,4-triazol-1-yl)allyl benzoate (**6a**). The reaction of 1-(propa-1,2-dien-1-yl)-1*H*-1,2,4-triazole **6** with benzoic acid according to the general procedure A (flash chromatography eluent: Pentane/AcOEt 50:50) gave the title compound in the form of a white solid (63 %). ^1H NMR (400 MHz, CDCl_3) δ 8.25 (s, 1H), 8.08 – 8.02 (m, 2H), 7.99 (s, 1H), 7.61 – 7.51 (m, 1H), 7.43 (ddd, $J = 8.0$, 6.8, 1.1 Hz, 2H), 7.20 (dd, $J = 14.0$, 1.2 Hz, 1H), 6.54 (dt, $J = 14.0$, 6.5 Hz, 1H), 4.97 (dd, $J = 6.5$, 1.4 Hz, 2H). $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) δ 166.2, 133.4, 130.1, 129.7, 129.7, 128.7, 128.6, 126.7, 115.3, 62.1. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{12}\text{H}_{12}\text{N}_3\text{O}_2$ 230.0930, found 230.0933. mp: 62–63 °C

(*E*)-3-(1*H*-benzo[d][1,2,3]triazol-1-yl)allyl benzoate (**7a**). The reaction of 1-(propa-1,2-dien-1-yl)-1*H*-benzo[d][1,2,3]triazole **7** with benzoic acid according to the general procedure A (flash chromatography eluent: Pentane/AcOEt 95:05) gave the title compound in the form of a white solid (52 %). ^1H NMR (400 MHz, CDCl_3) δ 8.14 – 8.06 (m, 3H), 7.75 – 7.66 (m, 2H), 7.63 – 7.52 (m, 2H), 7.51 – 7.38 (m, 3H), 6.74 (dt, $J = 14.4$, 6.7 Hz, 1H), 5.11 (dd, $J = 6.7$, 1.4 Hz, 2H). $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) δ 166.4, 146.5, 133.4, 131.5, 129.9, 129.8, 128.7, 128.6, 127.1, 124.9, 120.6, 114.6, 110.2, 62.9. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{16}\text{H}_{14}\text{N}_3\text{O}_2$ 280.1086, found 280.1091. mp: 132–133 °C

ASSOCIATED CONTENT

Supporting Information

^1H and ^{13}C NMR spectra for all compounds

AUTHOR INFORMATION

Corresponding Author

* Email: florian.monnier@enscm.fr, marc.taillefer@enscm.fr

Author Contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

Notes

Any additional relevant notes should be placed here.

ACKNOWLEDGMENT

Financial support provided by Région Languedoc-Roussillon (PhD fellowship for RB), IUF (Institut Universitaire de France to FM), ANR CD₂I (Agence Nationale de la Recherche) and CNRS are warmly acknowledged with thanks. We thank Racha Abed Ali Abdine for discussion and assistance.

REFERENCES

- (1) For reviews on transition-metal-catalyzed allylic substitutions, see: (a) Qu, J.; Helmchen G. Applications of Iridium-Catalyzed Asymmetric Allylic Substitution Reactions in Target-Oriented Synthesis. *Acc. Chem. Res.* **2017**, *50*, 2539–2555. (b) Butt, N. A.; Zhang, W. Transition metal-catalyzed allylic substitution reactions with unactivated allylic substrates. *Chem. Soc. Rev.* **2015**, *44*, 7929–7967. (c) Lu, Z.; Ma, S. Metal-Catalyzed Enantioselective Allylation in Asymmetric Synthesis. *Angew. Chem., Int. Ed.* **2008**, *47*, 258–297. (d) Trost, B. M.; Crawley, M. L. Asymmetric Transition-Metal-Catalyzed Allylic Alkylations: Applications in Total Synthesis. *Chem. Rev.* **2003**, *103*, 2921–2944. (e) Trost, B. M.; Van Vranken, D. L. Asymmetric Transition Metal-Catalyzed Allylic Alkylations. *Chem. Rev.* **1996**, *96*, 395–422.
- (2) For recent reports respectively with Pd, Pt, Au, Rh, Ni, Ag and Cu, see: (a) Tafazolian, H.; Samblanet, D. C.; Schmidt, J. A. R. Electronic Role of 3-Iminophosphine Ligands in Palladium-Catalyzed Intermolecular Hydroamination. *Organometallics*, **2015**, *34*, 1809–1817. (b) Toups, K. L.; Widenhoefer, R. A. Platinum(II)-catalyzed intermolecular hydroamination of monosubstituted allenes with secondary alkylamines. *Chem. Commun.* **2010**, *46*, 1712–1714. (c) Zeng, X.; Soleilhavoup, M.; Bertrand, G. Gold-Catalyzed Intermolecular Markovnikov Hydroamination of Allenes with Secondary Amines. *Org. Lett.* **2009**, *11*, 3166–3169. (d) Cooke, M. L.; Xu, K.; Breit, B. Enantioselective rhodium-catalyzed synthesis of branched allylic amines by intermolecular hydroamination of terminal allenes. *Angew. Chem., Int. Ed.* **2012**, *51*, 10876–10879. (e) Tafazolian, H.; Schmidt, J. A. R. Cationic [(Iminophosphine)Nickel(Allyl)]⁺ Complexes as the First Example of Nickel Catalysts for Direct Hydroamination of Allenes. *Chem. Eur. J.* **2017**, *23*, 1507–1511. (f) Wei, T.; Xie, M. S.; Qu, G.-R.; Niu H.-Y.; Guo, H.-M. A new strategy to construct acyclic nucleosides via Ag(I)-catalyzed addition of pronucleophiles to 9-allenyl-9H-purines. *Org. Lett.* **2014**, *16*, 900–903. (g) Blieck, R.; Bahri, J.; Taillefer, M.; Monnier, F. Copper-Catalyzed Hydroamination of Terminal Allenes. *Org. Lett.* **2016**, *18*, 1482–1485.
- (3) (a) Yamamoto, Y.; Al-Masum, M.; Asao, N. Palladium-Catalyzed Addition of Activated Methylene and Methyne Compounds to Allenes. *J. Am. Chem. Soc.* **1994**, *116*, 6019–6020. (b) Trost, B. M.; Gerusz, V. J. Palladium-Catalyzed Addition of Pronucleophiles to Allenes. *J. Am. Chem. Soc.* **1995**, *117*, 5156–5157. (c) Besson, L.; Goré, J.; Cazes, B. Palladium-catalyzed addition of malonate type compounds to allenes via a hydropalladation process. *Tetrahedron Lett.* **1995**, *36*, 3853–3856. (d) Yamamoto, Y.; Al-Masum, M. Palladium Catalyzed α -Addition of Certain Pronucleophiles to Alkoxyallenes. *Synlett*, **1995**, 969–970. (e) Yamamoto, Y.; Al-Masum, M.; Takeda, A. Palladium Catalyzed α -Addition of Certain

Pronucleophiles to Alkoxyallenes. *Chem. Commun.* **1996**, 831-832. (f) Trost, B. M.; Simas, A. B. C.; Plietker, B.; Jäkel, C.; Xie, J. Enantioselective palladium-catalyzed addition of 1,3-dicarbonyl compounds to an allene derivative. *Chem. Eur. J.* **2005**, *11*, 7075-7082. (g) Zhou, H.; Wang, Y.; Zhang, L.; Cai, M.; Luo, S. Enantioselective Terminal Addition to Allenes by Dual Chiral Primary Amine/Palladium Catalysis. *J. Am. Chem. Soc.* **2017**, *139*, 3631-3634. (h) Li, C.; Breit, B. Rhodium-Catalyzed Chemo- and Regioselective Decarboxylative Addition of β -Ketoacids to Allenes: Efficient Construction of Tertiary and Quaternary Carbon Centers. *J. Am. Chem. Soc.*, **2014**, *136*, 862-865. (i) Beck, T. M.; Breit, B. Regio- and Enantioselective Rhodium-Catalyzed Addition of 1,3-Diketones to Allenes: Construction of Asymmetric Tertiary and Quaternary All Carbon Centers. *Angew. Chem., Int. Ed.* **2017**, *56*, 1903-1907. (j) Blicke, R.; Abed Ali Abdine, R.; Taillefer, M.; Monnier, F. Regio- and Stereoselective Copper-Catalyzed Allylation of 1,3-Dicarbonyl Compounds with Terminal Allenes. *Org. Lett.* **2018**, *20*, 2232-2235.

(4) For selected reports on hydroalkoxylation of allenes, see: (a) Harris, R. J.; Carden, R. G.; Duncan, A. N.; Widenhofer, R. A. Kinetics and Mechanism of the Gold-Catalyzed Intermolecular Hydroalkoxylation of Allenes with Alcohols. *ACS Catal.* **2018**, *8*, 8941-8952. (b) Webster, S.; Sutherland, D. R.; Lee, A.-L. Chirality Transfer in Gold(I)-Catalysed Hydroalkoxylation of 1,3-Disubstituted Allenes. *Chem. Eur. J.* **2016**, *22*, 18593-18600. (c) Liu, Z.; Breit, B. Rhodium-Catalyzed Enantioselective Intermolecular Hydroalkoxylation of Allenes and Alkynes with Alcohols: Synthesis of Branched Allylic Ethers. *Angew. Chem., Int. Ed.*, **2016**, *55*, 8440-8443. (d) Jiang, L.; Jia, T.; Wang, M.; Liao, J.; Cao, P. Pd-Catalyzed Enantioselective Hydroalkoxylation of Alkoxyallenes with Phenol for Construction of Acyclic O,O- Acetals. *Org. Lett.* **2015**, *17*, 1070-1073. (e) Lim, W.; Kim, J.; Rhee, Y. H. Pd-Catalyzed Asymmetric Intermolecular Hydroalkoxylation of Allene: An Entry to Cyclic Acetals with Activating Group-Free and Flexible Anomeric Control. *J. Am. Chem. Soc.* **2014**, *136*, 13618-13621. (f) Zhang, Z.; Lee, S. D.; Fisher, A. S.; Widenhofer, R. A. Gold(I)-catalyzed hydration of allenes. *Tetrahedron*, **2009**, *65*, 1794-1798. (g) Paton, R. S.; Maseras, F. Gold(I)-Catalyzed Intermolecular Hydroalkoxylation of Allenes: A DFT study. *Org. Lett.* **2009**, *11*, 2237-2240. (h) Kawamoto, T.; Hirabayashi, S.; Guo, X.-X.; Nishimura, T.; Hayashi, T. Rhodium-catalyzed asymmetric hydroalkoxylation and hydrosulfenylation of diphenylphosphinylallenes. *Chem. Commun.* **2009**, 3528-3530. (i)

Nishina, N.; Yamamoto, Y. Gold-catalyzed intermolecular hydroalkoxylation of allenes; difference in mechanism between hydroalkoxylation and hydroamination. *Tetrahedron Lett.* **2008**, *49*, 4908-4911. For a general review, see: (j) Le Bras, J. Muzart, J. Palladium-catalysed inter- and intramolecular formation of C-O bonds from allenes. *Chem. Soc. Rev.* **2014**, *43*, 3003-3040.

(5) Kim, I. S.; Krische, M. J. Iridium-Catalyzed Hydrocarboxylation of 1,1-Dimethylallene: Byproduct-Free Reverse Prenylation of Carboxylic Acids. *Org. Lett.* **2008**, *10*, 513-515.

(6) (a) Koschker, P.; Lumbroso, A.; Breit, B. Enantioselective Synthesis of Branched Allylic Esters via Rhodium-Catalyzed Coupling of Allenes with Carboxylic Acids. *J. Am. Chem. Soc.* **2011**, *133*, 20746-20749. For macrolactonisation via allene hydrocarboxylation, see : (b) Ganss, S.; Breit, B. Enantioselective Rhodium-Catalyzed Atom-Economical Macrolactonization. *Angew. Chem. Int. Ed.* **2015**, *55*, 9738-9742.

(7) See ref 2f

(8) (a) Al-Masum, M.; Yamamoto, Y. Palladium-Catalyzed Hydrocarboxylation of Allenes. *J. Am. Chem. Soc.* **1998**, *120*, 3809-3810; (b) Patil, N. T.; Pahadi, N. K.; Yamamoto, Y. Pd(0)-PhCOOH catalyzed addition of oxygen pronucleophiles to allenes and internal alkyne. *Can. J. Chem.* **2005**, *83*, 569-574.

(9) (a) Perego, L.; Blicke, R.; Groué, A.; Monnier, F.; Taillefer, M.; Ciofini, I.; Grimaud, L. Copper-Catalyzed Hydroamination of Allenes: from Mechanistic Understanding to Methodology Development. *ACS Catal.* **2017**, *7*, 4253-4264. b) Perego, L.; Blicke, R.; Michel, J.; Ciofini, I.; Grimaud, L.; Taillefer, M.; Monnier, F. Copper-Catalyzed Hydroamination of N-Allenylazoles: Access to Amino-Substituted N-Vinylazoles. *Adv. Synth. Cat.* **2017**, *359*, 4388-4392. (c) Blicke, R.; Perego, L. A.; Ciofini, I.; Grimaud, L.; Taillefer, M.; Monnier, F. Copper-Catalysed Hydroamination of N-Allenylsulfonamides: The Key Role of Ancillary Coordinating Groups. *Synthesis*, **2019**, *51*, 1225-1234.

(10) Kubas, G. J.; Monzyk, B.; Crumblis, A. L. Tetrakis(Acetonitrile)Copper(1+) Hexafluorophosphate(1-). *Inorg. Synth.* 1990, **28**, 68.

(11) Khurana, J. M.; Chauhan, S.; Bansal, G. Facile Hydrolysis of Esters with KOH-Methanol at Ambient Temperature . *Monatshefte Für Chemie*, 2004, **135**, 83-87.