Adverse effects induced by chronic gamma irradiation in progeny of adult fish not affecting parental reproductive performance

Noémie Guirandy, Béatrice Gagnaire, Sandrine Frelon, Thomas Munch, Nicolas Dubourg, Virginie Camilleri, Isabelle Cavalie, Magali Floriani, Caroline Arcanjo, Sophia Murat El Houdigui, et al.

► To cite this version:

Noémie Guirandy, Béatrice Gagnaire, Sandrine Frelon, Thomas Munch, Nicolas Dubourg, et al.. Adverse effects induced by chronic gamma irradiation in progeny of adult fish not affecting parental reproductive performance. *Environmental Toxicology and Chemistry*, 2019, 38 (11), pp.2556-2567. 10.1002/etc.4562 . hal-02324157

HAL Id: hal-02324157

<https://hal.science/hal-02324157>

Submitted on 7 Jul 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Référence :

GUIRANDY Noemie, GAGNAIRE Beatrice, FRELON Sandrine, MUNCH Thomas, DUBOURG Nicolas, CAMILLERI Virginie, CAVALIE Isabelle, FLORIANI Magali, ARCANJO Caroline, MURAT EL HOUDIGUI Sophia, ARMANT Olivier, ADAM-GUILLERMIN Christelle, GONZALEZ Patrice, SIMON Olivier

Adverse effects induced by chronic gamma irradiation in progeny of adult fish not affecting parental reproductive performance

Environ. Toxicol. Chem., **38**, 11 (John Wiley & Sons Ltd 2019) 2556-2567 | 10.1002/etc.4562 (ACL)

Auteurs dans l'ordre de la publication :

1, GUIRANDY, Noemie

Institut de radioprotection et sûreté nucléaire, PSE-ENV/SRTE/LECO, Cadarache, France

2, GAGNAIRE, Beatrice

Institut de radioprotection et sûreté nucléaire, PSE-ENV/SRTE/LECO, Cadarache, France

3, FRELON, Sandrine

Institut de radioprotection et sûreté nucléaire, PSE-ENV/SRTE/LECO, Cadarache, France

4, MUNCH, Thomas

Institut de Radioprotection et de Sûreté Nucléaire, PSE-ENV/SRTE/LECO, France

5, DUBOURG, Nicolas

Institut de radioprotection et sûreté nucléaire, PSE-ENV/SRTE/LECO, Cadarache, France

6, CAMILLERI, Virginie

Institut de radioprotection et sûreté nucléaire, PSE-ENV/SRTE/LECO, Cadarache, France

7, CAVALIE, Isabelle

Institut de radioprotection et sûreté nucléaire, PSE-ENV/SRTE/LECO, Cadarache, France

8, FLORIANI, Magali

Institut de radioprotection et sûreté nucléaire, PSE-ENV/SRTE/LECO, Cadarache, France

9, ARCANJO, Caroline

Institut de Radioprotection et de Sûreté Nucléaire, PSE-ENV/SRTE/LECO, France

10, MURAT EL HOUDIGUI, Sophia

Institut de radioprotection et sûreté nucléaire, PSE-ENV/SRTE/LECO, Cadarache, France

11, ARMANT, Olivier

Institut de radioprotection et sûreté nucléaire, PSE-ENV/SRTE/LECO, Cadarache, France

12, ADAM-GUILLERMIN, Christelle

Institut de radioprotection et sûreté nucléaire, PSE-SANTE/SDOS/LMDN, Cadarache, France

13, GONZALEZ, Patrice

Université Bordeaux, UMR CNRS 5805, Avenue des Facultés - 33405 TALENCE

CEDEX, France

14, SIMON, Olivier, REFERENT , CORRESPONDING AUTHOR

Institut de radioprotection et sûreté nucléaire, PSE-ENV/SRTE/LECO, Cadarache, France

Running Head: Irradiation leads to drastic effects in progeny

Title: Adverse effects induced by chronic gamma irradiation in progeny of adult fish not affecting parental reproductive performance

Abstract: Multigenerational studies has become of great interest in ecotoxicology since the consequence of parental exposure to contaminants on offspring generations was established *in situ* or in laboratory conditions. This study mainly examined the chronic effects of external Cs-137 gamma irradiation exposure at 4 dose rates (control, 0.5, 5 and 50 mGy h⁻¹) on adult zebrafish (F0) exposed for 10 days and its progeny (F1) exposed or unexposed for 4/5 days. The main endpoints investigated included parental reproductive performance, embryo-larval survival, DNA alterations and ROS production in F0 and F1. No effects on reproductive success, fecundity or egg fertilization rate were observed. However, drastic effects were observed on F1 exposed to 50 mGy h⁻¹, resulting in a mortality rate of 100%. The drastic effects were also observed when the progeny was not irradiated. It was demonstrated that the sensitivity of the embryos was mainly due to parental irradiation. Moreover, these drastic effects induced by adult irradiation disappeared over time when 10 d- irradiated adults were placed in a non-irradiated condition. DNA alterations in larvae were observed for the three dose rates, and an increase of ROS production was also shown for the two lowest dose rates. This study improves our understanding of the consequences of parental exposure conditions to the progeny. Furthermore, it provides an incentive to take transmitted generational effects into account in ecological risk assessments.

Keywords: reproduction, irradiation exposure, zebrafish, risk assessment

INTRODUCTION

Gamma radiation represents a potential health risk to biota, due to its ability to ionize molecules in tissue. Ionizing radiation is known to induce oxidative stress, DNA damage and apoptosis, which therefore constitute usual molecular markers for evaluating toxicity mechanisms (Gagnaire et al. 2015; Jaafar et al. 2013; Knowles 2002; Praveen Kumar et al. 2017; Simon et al. 2011b; Sinha et al. 2018). Significant effects in animals (fish, nematode, *Daphnia*) in terms of survival, reproduction and development have been observed (Adam-Guillermin et al. 2012; Buisset-Goussen et al. 2014; Gagnaire et al. 2015; Knowles 2002; Parisot et al. 2015; Simon et al. 2011b). Moreover, damage effects in fish have mainly been observed for early life stages, which are considered as the most vulnerable to ionizing radiation. Embryogenesis, in addition to gametogenesis and organogenesis, can be affected by ionizing radiation due to the high rate of cell division, proliferation and differentiation (Hurem et al. 2017a; Hurem et al. 2018) and enable the consequences of irradiation to be assessed.

In terms of ecological risk assessment (ERA), endpoints on the impact on adult reproduction performance (reproductive success, fecundity, fertility) and on development of early stages (growth, survival) remain the main useful endpoints in characterising the ecological consequences of pollutants, including ionizing radiation. Radiological protection criteria are largely based on data from acute exposure experiments of adult organisms, thus information on the effects of ionizing radiation during sensitive life stages and after chronic exposure are lacking (Hurem et al. 2017a; Hurem et al. 2017b). Overall, few data are available concerning the effects of ionizing radiation in fish. The ecological screening benchmark for a generic

ecosystem of 10 $\mu\text{Gy h}^{-1}$ based on the HDR₅ (Hazardous Dose Rate for 5% of the species) for radioactive substances was built with five EDR₁₀ (Effect Dose Rate related to a change of 10% for a particular effect) from fish, which involved values ranging from 47 to 20,881 $\mu\text{Gy h}^{-1}$ (Garnier-Laplace et al. 2010). This generic benchmark dose rate was proposed by the European project, ERICA, for the screening of potential radiological effects (Brown et al. 2008). For freshwater and chronic γ external exposure, applying a safety factor of 50 to the lowest EDR₁₀ (reproductive endpoint: 516 $\mu\text{Gy h}^{-1}$) led to a predicted no-effect value of 10 $\mu\text{Gy h}^{-1}$ (Garnier-Laplace et al. 2006). Derived consideration reference levels (DCRLs) *i.e.*, “the dose rate band within which there is likely to be some chance of deleterious effects occurring to individuals of such type of a given type of Reference Animal and Plant (RAP)” indicate a possible reduction of reproductive success for dose rates ranging between 40 and 4,000 $\mu\text{Gy h}^{-1}$ for trout (ICRP 2012).

Firstly, the EDR₁₀ did not include the assessment of adult reproduction performance. Secondly, none of these fish species belong to the cyprinid family. Thirdly, the consequence of parental exposure for offspring generations have not yet been taken into account in the ERA. However, multigenerational studies have become of great interest (Buisset-Goussen et al. 2014; Hurem et al. 2017a; Lemos et al. 2017; Parisot et al. 2015). For fish, the progeny of adult zebrafish exposed to 53 mGy h^{-1} (^{60}Co gamma radiation) for 27 days showed a 100 % mortality rate occurring at the gastrula stage (Hurem et al. 2017a). Thus, parental exposure can lead to hereditary effects in offspring, probably due to epigenetic mechanisms (Herráez et al. 2017; Hurem et al. 2017a; Lemos et al. 2017).

As fish are known to be the most radiosensitive organisms among the poikilothermic aquatic animals (Garnier-Laplace et al. 2006), zebrafish, or *Danio rerio*, were chosen in the present research as a cyprinid model for assessing gamma radiation effects on reproduction performance and larval development. Zebrafish have been widely used for examining effects of ionizing radiation (Choi et al. 2010; Choi et al. 2015; Gagnaire et al. 2015; Hurem et al. 2017a; Hurem et al. 2017b; McAleer et al. 2005; Ng et al. 2017; Simon et al. 2011b; Yum et al. 2010) in addition to transgenerational effects (Hurem et al. 2017a; Kamstra et al. 2018). Indeed, breeding success, high fecundity and rapid development are the main advantages of this model in performing multigenerational studies (Lawrence 2007; Simon et al. 2014).

In this study, adult zebrafish were exposed to four dose rates (0, 0.5, 5 and 50 mGy h⁻¹, ¹³⁷Cs gamma radiation) over 10 days. The dose rates were higher than those in “hot spots” measured in freshwater ecosystems around Chernobyl (< 0.2 mGy h⁻¹) (Bonzom et al. 2016; Fuller et al. 2018; Lecomte-Pradines et al. 2014; Lerebours et al. 2018). Reproduction was initiated and the progeny were then exposed or not to the same dose rates over 96-120 hours. The objectives of this study were to evaluate (i) the adult reproductive performance, (ii) the effects in the progeny of irradiated parents and (iii) the responses at the molecular and organism level. Effects on adults were assessed by reproductive performance and genotoxicity (comet assay). Effects on the progeny were assessed using the survival rate (%) and the genotoxic and oxidative stress effects. The progeny were also placed in non-irradiated conditions (F1 recovery) over 120 hours to evaluate the impact of adult irradiation on the progeny (generational effects).

Complementary objectives concerned the increase of the exposure duration on the effects and the time necessary to obtain reproductive resilience: for the two lowest dose rates tested (lowDR: 0.5, 5 mGy h⁻¹), adult F0 exposure duration was increased to 24 days; for the highest

dose rate (highDR: 50 mGy h⁻¹), 10 day-irradiated adults (F0) were placed in non-irradiated conditions over 63 days (F0 recovery). Adult performance, progeny survival and molecular effects were also evaluated during these complementary experiments.

MATERIALS AND METHODS

Fish husbandry

The project (APAFIS#15821) was authorized by the IRSN ethics committee N°81 (EU 0520) in an application under the directive 2010/63/UE relating to animal care. The study was conducted on wild-type zebrafish that were kept, reproduced and irradiated in a zebrafish housing system (Zebtec Tecniplast stand Alone, Varese, Italy) with recirculating fresh water. Adult fish were acclimatised over 2-3 weeks to tap water + 20% demineralized water renewed daily (Aquadem, Veolia, France) (pH=7.4 ± 0.2, conductivity = 398 ± 2 µS cm⁻¹, T=28.4 ± 0.3 °C), with a 14 h light/10 h dark cycle photoperiod. The fish were fed *ab libitum* twice a day with commercial flakes (Tetramin®).

Adult and embryo exposure

Figure 1 shows the exposure duration and endpoints for the experiments. Nominal dose rates were 0.5 and 5 mGy h⁻¹ for the low dose experiment (lowDR) and 50 mGy h⁻¹ for the high dose experiment (highDR). As both experiments (lowDR and highDR) were not carried out at the same time, control fish conditions were implemented for each experiment. Gamma-rays were emitted from a ¹³⁷Cs source (444 GBq, 662 keV, IRSN-MICADO-Lab platform). Dose rates were simulated using MCNP5 software and measured using thermoluminescent dosimeters (Chiyoada Technologies, Japan) and the values represented between 91.1 and

100.8% and between 80 and 120% of the nominal low DR and high DR values, respectively.
Two control conditions were kept in a separate room (60-80 nGy h⁻¹).

The population density of adult fish was 0.7 L g⁻¹. For the low dose rate experiment (lowDR), 20 adult fish per condition (female: 0.66 ± 0.13 g, n=24, male: 0.50 ± 0.065 g, n=11) were irradiated over 10 and 24 days. For the high dose rate experiment (highDR), 24 adult fish per condition (female: 0.38 ± 0.06 g, n=6, male: 0.40 ± 0.07 g, n=6) were irradiated over 10 days. To keep the radiation exposure as long as possible, the daily control of abiotic parameters (pH, conductivity, Temperature), the animal welfare and the feeding process were carried out in less than one hour, five times per week. During this hour, irradiation was stopped in order to access to the tanks. Food was supplied to the fish twice a day by automatic suppliers. During the weekends, the control of these parameters was achieved through the use of cameras, allowing animal welfare to be monitored without interrupting the irradiation. The preparation of reproduction and egg collection required stopping the irradiation during four hours. After 24 d of exposure, the adults from lowDR were sacrificed and the gonads were collected. After 10 d of exposure, six adults from highDR were sacrificed and the gonads were collected. Six other adults were introduced into non-irradiation conditions (F0 Recovery).

For each reproduction event, embryos were obtained from 10 spawning genitors (*i.e.* replicates) for lowDR and 12 genitors (*i.e.* replicates) for highDR per condition. Each group, consisting of one male and one female, was placed in specific spawning aquariums to avoid egg predation in the zebrafish housing system. Adults were irradiated during reproduction. After spawning, the egg viability (3-4 hpf, hours post fertilisation) of each spawn was confirmed when the blastula stage was reached. The embryos (2.6 ml/egg) were then reintroduced into small beakers (Ø=9 cm) in the zebrafish housing system to be irradiated (at

the same dose rate as the adult groups (F1 irradiated group) or without irradiation (F1 Recovery group) and larvae were fed starting from 5 dpf (days post fertilisation) (artificial commercial food ST-1, Aqua Schwarz GmbH). The culture medium of the embryos was the same as that of the adults and 10% was replenished daily.

Ecologically-representative endpoints for adults

For the lowDR experiment, adult F0 reproductive performance (n=10 couples per condition) was assessed after 10 and 24 days of exposure for both dose rates tested. Endpoints concerned the reproductive success (number of couples that spawned), the fecundity (number of eggs per female) and the egg survival rate at 3 hpf (fertilization rate). For the highDR experiment, F0 reproductive performance was assessed after 10 days of irradiation (n=12 couples per condition) and after 6, 36 and 63 days in non-irradiated conditions (F0 Recovery, n=6 couples) (Figure 1).

Ecologically-representative endpoints and oxidative stress analysis for progeny

For both experiments, the progeny survival rate (%) was assessed after 24 and 48 h in the control, irradiated and non-irradiated (recovery) conditions. Complementary experiments consisted of irradiating 3 hpf-eggs at 50 mGy h⁻¹ over 96 h (30 eggs from three spawns per condition) to measure the survival rate.

For the lowDR experiment, two pseudo-replicates of 80 eggs originating from three spawns per condition including the control, irradiated and non-irradiated conditions, were followed to measure the progeny survival rate over 1, 2, 10 and 22 d. For the highDR experiment, 30 eggs from six spawns of the three conditions were tracked over 5 d (120h).

For the highDR experiment, photographs of 72 hpf-larvae were recorded. Morphology (whole body size, surface of yolk reserve), malformations and cardiac edema were then assessed using software (Danioscope image analysis system version 10.0, Noldus). Analyses were made on 10 72 hpf-larvae per condition after 10 d F0 exposure (F0Irr. 10d) and after F0 recovery at 6 days (T10d+R6d) and at 36 days (T10d+R36d).

Reactive oxygen species (ROS) production was measured in F1-progeny from the lowDR experiment using a protocol adapted from Hurem et al, (2017) and Gagnaire et al., 2015. Given the high larval mortality rate, no measurements were made during the highDR experiment. ROS production was determined using the fluorescent probe (2', 7' - dichlorodihydrofluorescein diacetate (H₂DCFDA, Sigma-Aldrich, St. Louis, USA)) in 10 larvae per condition (larvae exposed from 3-4 hpf to 4 or 10 days originating from 10 d-adult and 24 d-adult reproduction). Through the oxidation of H₂DCFDA in dimethylsulfoxide (DMSO, Sigma-Aldrich, St. Louis, USA) water by ROS (mainly H₂O₂) the molecule was converted into 2', 7' dichlorodihydrofluorescein (DCF), which is highly fluorescent. The reported wavelengths for the measurement of DCF fluorescence are 500 nm for excitation and 525 nm for emission (TECAN Infinite M1000, Switzerland).

Genotoxic effects in progeny and adult

The genotoxic effects induced by gamma irradiation in the developmental stages (1 d, 4 d, 10 d for lowDR and 1 d for highDR experiments) and in gonads (10 d, highDR experiment) of zebrafish were evaluated using the alkaline comet assay as described in Sing et al, (1998) (1988) with modifications (Simon et al. 2011a; Simon et al. 2018). After centrifugation (110 g, 10 min, 8 °C, Eppendorf, 5427R, Germany), pellets were suspended in 1 mL of L15-HEPES and used immediately. Cells were counted on a Malassez cell and their viability was

assessed using trypan blue. Two hundred nucleoids per slide were analysed at $\times 400$ magnification under a fluorescence microscope (Nikon Eclipse E600) equipped with a 515–560 nm excitation filter. Comet pictures were analysed using Comet IV software (Perceptive Instruments). This assay was performed on three pools of ten 24-hpf eggs, on 10 individual 96-hpf larvae and on three pools of whole gonads per sex and per condition. The Tail Moment was defined as the percentage of DNA in the tail multiplied by the length between the centre of the head and tail.

Statistical analysis

All data are presented as means $\pm$ SD with significance taken at $p < 0.05$. A comparison between the control and irradiation or recovery exposures was performed. Before the statistical analyses, the normality and homogeneity of the variance were tested using the Shapiro-Wilks and Fisher tests, respectively. The T-test and one or two ways-ANOVAs were used when data were normally distributed. The non-parametric test (Kruskal-Wallis) was used when data were not normally distributed.

RESULTS

There were no instances of adult fish mortality during the experimental periods (control, low DR and high DR experiments).

Cumulative doses after exposure conditions

For the LowDR experiment, cumulative doses in adults ranged between 120 and 2880 mGy after 10 and 24 days of exposure, respectively. For the highDR experiment, the cumulative dose in adults was 12 Gy (SD Table 1). For the progeny, maximal cumulative

doses ranged from 264 to 2640 mGy for the lowDR experiment and 6 Gy for the highDR experiment (SD Table 1).

Adult performance (reproductive success, fecundity, total egg number and 3hpf-egg mortality)

For the highDR experiment, all females (n=12 per condition) spawned after 10 days of exposure. For the lowDR experiment (n=10 per condition), reproductive success was 100% and 80% for both the irradiated conditions and control conditions, respectively.

Figure 2 shows the fecundity (number of eggs per female) after 10 days of exposure for both experiments. High inter-individual variability was observed in all conditions. A significant difference (Kruskal-Wallis, $p=0.03$) was observed between 0.5 mGy h⁻¹ (240 eggs $\pm$ 132) and control (364 eggs $\pm$ 123) conditions. In the control LowDR, 43% of all spawns ranged between 101 and 300 eggs and 57% of spawns had higher than 300 eggs. At 0.5 mGy h⁻¹, 20% of all spawns produced less than 100 eggs. At 5 mGy h⁻¹, no significant difference in fecundity (413 eggs $\pm$ 148) was observed even though 89% of spawns had more than 300 eggs. At the high dose rate (50 mGy h⁻¹), a change in the size of spawns was also observed, with an increase in the number of spawning events (75%, 9/12), providing between 101 and 300 eggs compared to the control (42%, 5/12).

A significant difference was observed between both control fish fecundity (lowDR: 364 eggs $\pm$ 123 per female (n=7) versus the highDR: 203 $\pm$ 117 eggs per female (n=12), $p=0.042$ Kruskal-Wallis test) and egg quantity distribution.

The quality of oocytes was assessed by the survival 3hpf-progeny, indicating the number of fertilized *versus* non-fertilized eggs. The survival rate at 3 hpf remained high (>89%) without any significant difference between exposure conditions (SD Figure 1). Over the 50 spawns

monitored, only two presented a survival rate of less than 80%. Almost 85% of the spawns presented a survival rate of 90-100%.

Reproductive success after 24 days of lowDR exposure was not impacted by irradiation; 80, 90 and 100% of couples reproduced for the controls, 0.5 and 5 mGy h⁻¹, respectively. The significant difference in fecundity previously observed after 10 days between the control and 0.5 mGy h⁻¹ was not observed after 24 d of exposure. Fecundity was 375 ± 102, 289 ± 138 and 357 ± 138 eggs per female for the controls, 0.5 and 5 mGy h⁻¹, respectively. In the controls, 37.5% of spawns contained between 101 and 300 eggs and 62.5% contained more than 301 eggs, as observed after 10 days. At 5 mGy h⁻¹, 75 % of spawns presented more than 301 eggs, as previously observed after 10 days of exposure. The survival rate at 3 hpf-egg was close to the one measured after 10 days of adult exposure.

Progeny survival and size

Figure 3 shows F1 progeny survival rate (%) measured at 24 and 48 h for all exposure conditions of both experiments. For the lowDR experiment, the survival rate was higher than 87%, with no significant difference between the conditions. For the highDR experiment, a significant difference was observed between the control and larvae exposed to 50 mGy h⁻¹ for 24 h (p=0.02) and 48 h (p=0.006). Significant differences were also observed between the control and F1 recovery conditions for both times (p=0.015 and p=0.006 for 24 and 48 h, respectively). Figure 4 shows the survival rate (%) in F1 progeny originating from the lowDR experiment. Survival rates were higher than 80% and 60% at 2 and 10 d, respectively, for all exposure conditions. No significant effect of treatments and control was observed in the progeny coming from 10 d-irradiated adults. Similar results were observed for the progeny

coming from 24 d-irradiated adults. A significant difference was observed between 5 and F1R5 (Kruskal-Wallis test: $p=0.049$, $n=6$) at 22 days of exposure. No significant differences were observed for the hatching rate (%) at 48 h (data not shown).

For the irradiated and non-irradiated (F1 Recovery) larvae (Figure 5), 100% died after 120 h of highDR exposure, although the average survival rate for the control condition was $84 \pm 11\%$. In the control condition, the mortality rate was low and remained constant from 24 hpf. F1 irradiated (51%) and F1 Recovery (58%) survival rate at 24 hpf was low compared to the control and ranged between 16 to 96%, indicating a high spawn variability (SD Figure 2). Drastic effects seemed to appear more rapidly for F1 irradiated larvae than for F1 Recovery larvae (SD Figure 2).

Figure 6 shows the survival rate (%) in F1 progeny coming from the highDR experiment where adults were placed in non-irradiated exposure conditions (F0 Recovery) over 63 days. Fish were induced to spawn after 6, 36 and 63 days. Drastic effects of adult irradiation on survival progeny rate lasted for at least 6 days (35 ± 22 , $n=11$ spawns) and until 36 days (48 ± 37 , $n=8$ spawns). All larvae alive at 48 h from F0R0d, F0 R6d and F0 R36d, died at 96 h. After 63 days in non-irradiated conditions, the progeny survival rate remained high ($>85\%$, $n=11$ spawns) and no significant difference was observed compared to the control until 96 h.

For the highDR experiment, the body sizes of F1 irradiated and F1 Recovery larvae decreased significantly compared to the control (Figure 7). Similar results were also observed for non-irradiated 72 h-larvae originating from F0 recovery conditions after 6 days. After 36 d in non-irradiated conditions (F0 R36d), the body sizes of non-irradiated progeny showed no significant difference compared to the controls although 100% of mortality at 120 h was

observed. The surface of vitellus and pericardia area also showed no statistical difference with the controls (data not shown).

Oxidative stress

Figure 8 shows ROS production in F1-progeny coming from the lowDR experiment. Irradiation at 5 mGy h⁻¹ led to an increase of ROS production in 4 d-exposed larvae coming from 24 d-adult exposure. However, F1 R progeny exposed over 4 d at both dose rates showed a significant increase in ROS production for larvae coming from 10 d- and 24 d-adult exposure. No effects were observed in larvae (irradiated and recovery) exposed over 10 d (Figure 10B).

Genotoxic effects

Figure 9 shows DNA damage measured in progeny coming from 24 d-adults in the lowDR experiment, expressed by the tail moment. The progeny were placed in irradiated and non-irradiated conditions (F1 Recovery) over 10 days. Significant DNA damage was noticed at 4 d for irradiated (0.5 mGy h⁻¹: 11.3 ± 4.3%; 5 mGy h⁻¹: 14.4 ± 2.1%) and non-irradiated larvae (F1R: 0.5 mGy h⁻¹: 8.4 ± 3%) compared to the controls (x10). At 10 d, the tail moment showed no difference between the conditions.

Figure 10A shows DNA damage expressed by the tail moment in the progeny coming from 10 d-adult fish from the highDR experiment. The progeny was collected in irradiated and non-irradiated conditions (F1 Recovery) for 24 h. The high mortality rate did not allow larval collection at 4 and 10 d. A significant increase of the tail moment was observed between the control, irradiated (x6.5) and non-irradiated (x5) conditions. 10B-C shows DNA damage in the gonads of males and females after 10 days of irradiation. DNA damage

increased for both sexes compared to the control. The effects were more pronounced for males (x3) than for females (x1.7).

DISCUSSION

In the present work, embryo-larval and adult fish stages were exposed to 0.5, 5 and 50 mGy h⁻¹ (¹³⁷Cs gamma radiation) to augment the dataset of gamma ray effects after parental and progeny exposure.

Zebrafish as a relevant model for multigenerational studies

Reproduction was achieved for almost all couples and with a high egg survival rate (93 ± 7 % at 3 hpf for both control conditions, n=19), confirming the relevance of this model for multigenerational studies (Lawrence et al. 2012; Simon et al. 2014). Differences in reproductive performances were observed between the controls used in both lowDR and highDR experiments. The body weights of the control adults in both experiments was significantly different and could explain the differences in fecundity. Indeed, zebrafish fecundity, and more generally reproduction performance, can be influenced by the quality of the diet, the female body size and environmental enrichment (Karga et al. 2017; Lawrence 2007; Lawrence et al. 2012; Wafer et al. 2016). It is important to note that although zebrafish reproduce easily in laboratory conditions, assessing inter-individual variability of reproductive performances requires a large number of couples.

No effect of irradiation was observed on adult reproduction performance

For the testes dose rates, irradiation did not alter the reproductive capacity. Many pollutants which contribute to oxidative stress are known to alter reproductive performance, indicating that this physiological function may be sensitive to changes in the environment

(Faßbender et al. 2013; Newman et al. 2015; Simon et al. 2014; Wang et al. 2011). Perturbation of reproduction was previously observed in other fish species. Woodhead (1977) demonstrated that the total fecundity of guppies (*Poecilia reticulata*) was markedly reduced at dose rates of 1.7, 4 and 12.7 mGy h⁻¹ without drastic impacts on the mortality rate and survival of offspring. Rackham et al. (1984) demonstrated that spermatogenesis of adult butterfly splitfins (*Ameca splendens*) was disrupted (no production of sperm) after 7.3 mGy h⁻¹ of exposure over 52 days and that developing oocytes were less sensitive to the effects of radiation than spermatogenesis. Knowles (2002) observed a significant decrease in adult zebrafish in the mean number of eggs per spawn after 30 days at 7.4 mGy h⁻¹. Compared to this study, the toxic modes of action seem to be strongly influenced by the biological model and the experimental conditions of the dose rates. Under our experimental conditions, adult zebrafish stage did not exhibit sensitivity.

Drastic effects from irradiation on progeny survival were observed and transferred from adult fish

Drastic effects on progeny were only observed after high dose rate exposure and certainly had a detrimental effect at the population level. After exposure of adults to 5 mGy h⁻¹ over 10 or 24 days (cumulative dose rates of 1.2 and 2.8 Gy, respectively), no effect on progeny survival rate was observed until after 22 days. The range of these induced effects (5-50 mGy h⁻¹) is narrow and encourages us to more precisely determine the EDR₁₀. The non-irradiated progeny of 27 d-F0 exposed to 53 mGy h⁻¹ (34.3 Gy) previously showed a 100% mortality rate occurring at the gastrulation stage (8 hpf) (Hurem et al. 2017a). In this study, drastic effects appeared at a lower cumulative dose (12 Gy) for the same dose rate. These results raise questions regarding the best criteria (cumulative dose *versus* dose rate) for measuring gamma radiation exposure in the ERA.

The experimental design for the lowDR and highDR experiments did not allow the identical cumulative dose to be obtained. It could potentially be beneficial to vary the duration of exposure (1d at 50 mGy h⁻¹; 100d to 0.5 mGy h⁻¹) to compare the effects obtained at the same cumulative dose.

Adverse effects seemed to be due to adult exposure since (i) irradiated and recovery progeny showed identical effects compared to the control, and (ii) direct irradiation of embryos at this high dose rate did not cause significant mortality as confirmed in this study and by Simon et al, 2011. Vertebrate embryos are particularly sensitive to ionizing radiation, due to a high rate of cell division and migration (Hu et al. 2016; Hurem et al. 2017a; Jarvis et al. 2003; Rhee et al. 2012). The sensitivity of early stages (development and antioxidant enzymes activities) has already been demonstrated; 6hpf-embryos were more sensitive than 12- and 24-hpf embryos to gamma irradiation (0.01-1 Gy) (Hu et al. 2016). However, we demonstrated here that the sensitivity of the progeny to ionizing radiation was particularly high when they originated from irradiated parents.

Parental effects observed in the highDR experiment were reversible, as they disappeared when adults were placed in non-irradiated conditions for between 36 and 63 days. These results suggest F0 recovery of reproductive capacity. The F0 adults zebrafish exposed to 53 mGy h⁻¹ during 27 days (31 Gy) and placed in non-irradiated conditions failed to produce viable offspring 1.5 year after irradiation (Hurem et al. 2017a). The differences in terms of experimental design and cumulative dose between the present study and that of Hurem et al. (2017) encouraged us to elucidate the mechanisms behind the transfer from adult to progeny and the F0 recovery induction over time.

The earliest stages of embryonic development rely on maternal products that are generated during oogenesis and supplied to the egg. Yolk lipoprotein nutrients, vitamins, hormones, mRNA transcripts, and DNA methylation statuses have been identified as markers of egg quality. The damage to these maternal products could also explain the drastic effects observed in the progeny. Note that the period of maternal control of embryonic development varies among animals and could explain difference of ionizing radiation effects between species (Abrams et al. 2009; Vastenhouw et al. 2019). Among transferred maternal products, cortisol is essential for early development (Faught et al. 2016; Nesan et al. 2013; Nesan et al. 2016). Then, *de novo* synthesis of cortisol starts after hatching (48 hpf) (Nesan et Vijayan 2016). Finally, elevated cortisol levels in embryos leads to the same effects (mortality, pericardial oedema and heart malformation) (Nesan et al. 2012) observed in this study. Moreover, high levels of cortisol in females may directly impact estradiol production, possibly affecting vitellogenin production and its incorporation into the oocyte, leading to drastic effects in the progeny. In this study, we hypothesize that the high dose rates of gamma irradiation could have led to an alteration of the cortisol deposition and its transfer to eggs, which may have resulted in altered offspring phenotypes.

ROS production and genotoxicity after irradiated and recovery conditions

From potentially toxic mechanisms, the genotoxicity and ROS induction were assessed. Gamma irradiation is known to induce ROS formation in zebrafish larvae (Gagnaire et al. 2015; Jarvis et Knowles 2003). In this study, ROS production compared to the control was increased after 4 d of exposure of larvae originating from adults exposed during 10 and 24 days. Recovery larvae showed ROS production, without being irradiated. ROS production was higher in F1 recovery larvae than in irradiated larvae (originating from F0-10d). Once again, adult exposure led to high ROS production in larvae. This would seem to be a

consequence of an effect inherited from the parents. Results were similar to those observed in other exposure conditions (dose rate, recovery duration) by Hurem et al, 2017.

As no effect was observed in 10 d-larvae, the oxidative stress induced by gamma irradiation seems to be transitory. The activities of antioxidant enzymes are known to change in a developmental stage-dependent manner. However, larval ROS production in 4 d larvae was increased by gamma irradiation and persisted in F1 recovery condition. We hypothesized that under our experimental conditions, irradiation of the progeny led to an induction of protective agents that could counteract ROS production and re-establish a healthy cellular redox balance. Antioxidant enzyme activities could be assessed in further experiments in 10 d-irradiated larvae to test this hypothesis.

Significant genotoxic effects were observed at 4 d-irradiated and non-irradiated progeny coming from adults exposed in the lowDR experiment (0.5 and 5 mGy h⁻¹) during 24 days. High inter-individual variability was however observed for the lowDR experiment, preventing the highlighting of significant differences.

Genotoxicity has previously been demonstrated in eggs exposed to gamma rays in chronic or acute experiments (Gagnaire et al. 2015; Praveen Kumar et al. 2017; Simon et al. 2011b) or to non-radioactive pollutants (Kosmehl et al. 2008; Vicquelin et al. 2011). In this study, tail moment values were comparable to those measured in eggs/larvae exposed over 4 d at 24 mGy h⁻¹ (Gagnaire et al. 2015). Thus, these results suggest an increase in the intensity of genotoxic effects due to parental irradiation and confirm the molecular effects inherited from the parents. The genotoxic effects decreased at 10 d, suggesting a better efficiency of DNA repair mechanisms at this larval stage, potentially related to a better efficiency of the

antioxidant system. It would be interesting to measure some of the DNA repair mechanisms (as gamma H2AX activity) in order to confirm this hypothesis.

Irradiation of adults at high dose rates led to significant genotoxic effects on irradiated and non-irradiated progeny as early as 1 d, unlike results obtained at lower dose rates. However, the levels of genotoxic effects at 1 d remained lower than the ones measured at 4 d in the lowDR experiment which did not lead to a high mortality rate. The comparison between the different stages of development remains difficult and we have to take into account possible differences in reparation mechanism induction between the different stages to explain the relationship between mortality and genotoxicity effects. We suggest that the early live stage could be more sensitive than the later life stage.

High irradiation exposure led to significant genotoxic effects on adult gonads. However, the effects were lower than those measured after low uranium waterborne exposure (Simon et al. 2018). Significant differences were also observed in the controls between males and females, as previously observed in other studies (Simon et al. 2018). Moreover, irradiation effects seemed more pronounced for males than for females. Further studies may be completed by the determination of genotoxicity levels in zebrafish sperm, allowing a better link with genotoxicity observed in the progeny, as proposed by Reinardy et al. (2013). Indeed, during spermatogenesis, chromatin is processed and packaged and contact between DNA and nuclear matrix is reorganized. Furthermore, the epigenetic pattern is also totally re-modelled and transcription is stopped. All these events contribute to the control of embryo development in the early-life stages (Herráez et al. 2017).

CONCLUSION

These results on fish reproductive performance after gamma irradiation provide additional information for ERAs. Drastic effects on progeny and consequently on population levels were observed at dose rates between 5 and 50 mGy h⁻¹, confirming the DCRL values. However, it is important to note a difference in the sensitivity between the trout reference organism and the *Danio rerio*, the model organism in our study. Moreover, this study highlights that parental exposure leads to significant effects in the progeny at the molecular level (ROS, genotoxicity), even at low dose rates (0.5 m Gy h⁻¹). The consequences of these molecular effects will need to be studied following chronic exposure. The results encourage us to take into account the consequences of transmitted generational (*i.e.* heritable) effects for the determination of DCRL values. This study improves our understanding of the consequences of multigenerational exposure conditions for a better radioprotection of aquatic ecosystems. It confirms the difference in sensitivity of model species and identifies a narrow range of occurrence of drastic effects. It also acts as an incentive to acquire new data from multigenerational chronic exposure at low dose levels.

REFERENCES

- Abrams EW, Mullins MC. 2009. Early zebrafish development: It's in the maternal genes. *Curr Opin Gen Dev* 19:396-403.
- Adam-Guillermin C, Pereira S, Della-Vedova C, Hinton T, Garnier-Laplace J. 2012. Genotoxic and reprotoxic effects of tritium and external gamma irradiation on aquatic animals. *Rev Environ Contam Toxicol* 220:67-103.

490 Bonzom JM, Hättenschwiler S, Lecomte-Pradines C, Chauvet E, Gaschak S, Beaugelin-
 491 Seiller K, Della-Vedova C, Dubourg N, Maksimenko A, Garnier-Laplace J, Adam-Guillermin
 492 C. 2016. Effects of radionuclide contamination on leaf litter decomposition in the Chernobyl
 493 exclusion zone. *Sci Total Environ* 562:596-603.

494 Brown JE, Alfonso B, Avila R, Beresford NA, Copplestone D, Pröhl Gulanovsky A. 2008.
 495 The ERICA Tool. *J Environ Radioact* 99: 1371-1383.

496 Buisset-Goussen A, Goussen B, Della-Vedova C, Galas S, Adam-Guillermin C, Lecomte-
 497 Pradines C. 2014. Effects of chronic gamma irradiation: A multigenerational study using
 498 *Caenorhabditis elegans*. *J Environ Radioact* 137:190-197.

499 Choi VWY, Cheng SH, Yu KN. 2010. Radioadaptive response induced by alpha-particle-
 500 induced stress communicated *in vivo* between zebrafish embryos. *Environ Sci Tech* 44: 8829-
 501 8834.

502 Choi VWY, Yu KN. 2015. Embryos of the zebrafish *Danio rerio* in studies of non-targeted
 503 effects of ionizing radiation. *Cancer Lett* 356:91-104.

504 Faßbender C, Braunbeck T. 2013. Reproductive and genotoxic effects in zebrafish after
 505 chronic exposure to methyl methanesulfonate in a multigeneration study. *Ecotoxicology*
 506 22:825-837.

507 Faught E, Best C, Vijayan MM. 2016. Maternal stress-associated cortisol stimulation may
 508 protect embryos from cortisol excess in zebrafish. *R Soc Open Sci* 3(2):160032.

509 Fuller N, Ford AT, Nagorskaya LL, Gudkov DI, Smith JT. 2018. Reproduction in the
 510 freshwater crustacean *Asellus aquaticus* along a gradient of radionuclide contamination at
 511 Chernobyl. *Sci Total Environ* 628-629:11-17.

512 Gagnaire B, Cavalié I, Pereira S, Floriani M, Dubourg N, Camilleri V, Adam-Guillermin C.
 513 2015. External gamma irradiation-induced effects in early-life stages of zebrafish, *Danio*
 514 *rerio*. *Aquat Toxicol* 169:69-78.

515 Garnier-Laplace J, Della-Vedova C, Andersson P, Copplestone D, Cailles C, Beresford NA,
 516 Howard BJ, Howe P, Whitehouse P. 2010. A multi-criteria weight of evidence approach for
 517 deriving ecological benchmarks for radioactive substances. *J Radiol Prot* 30:215-233.

518 Garnier-Laplace J, Della-Vedova C, Gilbin R, Copplestone D, Hingston J, Ciffroy P. 2006.
 519 First derivation of predicted-no-effect values for freshwater and terrestrial ecosystems
 520 exposed to radioactive substances. *Environ SciTechn* 40:6498-6505.

521 Herráez MP, Ausió J, Devaux A, González-Rojo S, Fernández-Díez C, Bony S, Saperas N,
 522 Robles V. 2017. Paternal contribution to development: Sperm genetic damage and repair in
 523 fish. *Aquaculture* 472:45-59.

524 Hu M, Hu N, Ding D, Zhao W, Feng Y, Zhang H, Li G, Wang Y. 2016. Developmental
 525 toxicity and oxidative stress induced by gamma irradiation in zebrafish embryos. *Radiat*
 526 *Environ Biophys* 55:441-450.

527 Hurem S, Gomes T, Brede DA, Lindbo Hansen E, Mutoloki S, Fernandez C, Mothersill C,
 528 Salbu B, Kassaye YA, Olsen AK, Oughton D, Aleström P, Lyche JL. 2017a. Parental gamma
 529 irradiation induces reprotoxic effects accompanied by genomic instability in zebrafish (*Danio*
 530 *rerio*) embryos. *Environ Res* 159:564-578.

531 Hurem S, Martín LM, Brede DA, Skjerve E, Nourizadeh-Lillabadi R, Lind OC, Christensen
532 T, Berg V, Teien HC, Salbu B, Oughton DH, Aleström P, Lyche JL. 2017b. Dose-dependent
533 effects of gamma radiation on the early zebrafish development and gene expression. *PLoS*
534 *ONE* 12(6):e0179259.

535 Hurem S, Martín LM, Lindeman L, Brede DA, Salbu B, Lyche JL, Aleström P, Kamstra JH.
536 2018. Parental exposure to gamma radiation causes progressively altered transcriptomes
537 linked to adverse effects in zebrafish offspring. *Environ Pollut* 234:855-863.

538 ICRP. 2012. Protection of the environment under different exposure situations, Annals of the
539 ICRP.

540 Jaafar L, Podolsky RH, Dynan WS. 2013. Long-Term Effects of Ionizing Radiation on Gene
541 Expression in a Zebrafish Model. *PLoS ONE* 8(7):e69445.

542 Jarvis RB, Knowles JF. 2003. DNA damage in zebrafish larvae induced by exposure to low-
543 dose rate γ -radiation: detection by the alkaline comet assay. *Mutat Res*: 41, 63-69.

544 Kamstra JH, Hurem S, Martin LM, Lindeman LC, Legler J, Oughton D, Salbu B, Brede DA,
545 Lyche JL, Aleström P. 2018. Ionizing radiation induces transgenerational effects of DNA
546 methylation in zebrafish. *Sci Rep* 8(1):15373.

547 Karga J, Mandal SC. 2017. Effect of different feeds on the growth, survival and reproductive
548 performance of zebrafish, *Danio rerio* (Hamilton, 1822). *Aquac Nutr* 23:406-413.

549 Knowles JF. 2002. An investigation into the effects of chronic radiation on fish. *Report*
550 *Environment Agency, Bristol, UK*.

551 Kosmehl T, Hallare AV, Braunbeck T, Hollert H. 2008. DNA damage induced by
552 genotoxicants in zebrafish (*Danio rerio*) embryos after contact exposure to freeze-dried
553 sediment and sediment extracts from Laguna Lake (The Philippines) as measured by the
554 comet assay. *Mut Res* 650: 1-14.

555 Lawrence C. 2007. The husbandry of zebrafish (*Danio rerio*): A review. *Aquaculture* 269, 1-
556 20.

557 Lawrence C, Best J, James A, Maloney K. 2012. The effects of feeding frequency on growth
558 and reproduction in zebrafish (*Danio rerio*). *Aquaculture* 368:103-108.

559 Lecomte-Pradines C, Bonzom JM, Della-Vedova C, Beaugelin-Seiller K, Villenave C,
560 Gaschak S, Coppin F, Dubourg N, Maksimenko A, Adam-Guillermin C, Garnier-Laplace J.
561 2014. Soil nematode assemblages as bioindicators of radiation impact in the Chernobyl
562 Exclusion Zone. *Sci Total Environ* 490: 161-170.

563 Lemos J, Neuparth T, Trigo M, Costa P, Vieira D, Cunha L, Ponte F, Costa PS, Metello LF,
564 Carvalho AP. 2017. Single Low-Dose Ionizing Radiation Induces Genotoxicity in Adult
565 Zebrafish and its Non-Irradiated Progeny. *Bull Environ Contam Toxicol* 98: 190-195.

566 Lerebours A, Gudkov D, Nagorskaya L, Kaglyan A, Rizewski V, Leshchenko A, Bailey EH,
567 Bakir A, Ovsyanikova S, Laptev G, Smith JT. 2018. Impact of Environmental Radiation on
568 the Health and Reproductive Status of Fish from Chernobyl. *Environ Sci Technol* 52:9442-
569 9450.

570 McAleer MF, Davidson C, Davidson WR, Yentzer B, Farber SA, Rodeck U, Dicker AP.
571 2005. Novel use of zebrafish as a vertebrate model to screen radiation protectors and
572 sensitizers. *Int J Radiat Oncol Biol Phys* 61:10-13.

573 Nesan D, Vijayan MM. 2012. Embryo exposure to elevated cortisol level leads to cardiac
574 performance dysfunction in zebrafish. *Mol Cell Endocrinol* 363: 85-91.

575 Nesan D, Vijayan MM. 2013. Role of glucocorticoid in developmental programming:
576 Evidence from zebrafish. *Gen Comp Endocrinol* 181:35-44.

577 Nesan D, Vijayan MM. 2016. Maternal Cortisol Mediates Hypothalamus-Pituitary-Interrenal
578 Axis Development in Zebrafish. *Scientific Reports* 6:22582.

579 Newman TAC, Carleton CR, Leeke B, Hampton MB, Horsfield JA. 2015. Embryonic
580 oxidative stress results in reproductive impairment for adult zebrafish. *Redox Biology* 6:648-
581 655.

582 Ng CYP, Cheng SH, Yu KN. 2017. Photon hormesis deactivates alpha-particle induced
583 bystander effects between zebrafish embryos. *Radiat Phys Chem* 133:72-80.

584 Parisot F, Bourdineaud JP, Plaire D, Adam-Guillermin C, Alonzo F. 2015. DNA alterations
585 and effects on growth and reproduction in *Daphnia magna* during chronic exposure to gamma
586 radiation over three successive generations. *Aquat Toxicol* 163:27-36.

587 Praveen Kumar MK, Shyama SK, Kashif S, Dubey SK, Avelyno D, Sonaye BH, Kadam
588 Samit B, Chaubey RC. 2017. Effects of gamma radiation on the early developmental stages of
589 Zebrafish (*Danio rerio*). *Ecotoxicol Environ Saf* 142:95-101.

590 Rackham BD, Woodhead DS. 1984. Effects of Chronic γ -irradiation on the gonads of adult
591 *Ameca Splendens* (Osteichthyes: Teleostei). *Int J Radiat Biol Relat Stud Phys Chem Med*
592 45:645-656.

593 Reinardy HC, Syrett JR, Jeffree RA, Henry TB, Jha AN. 2013. Cobalt-induced genotoxicity
 594 in male zebrafish (*Danio rerio*), with implications for reproduction and expression of DNA
 595 repair genes. *Aquat Toxicol* 126:224-230.

596 Rhee JS, Kim BM, Kang CM, Lee YM, Lee JS. 2012. Gamma irradiation-induced oxidative
 597 stress and developmental impairment in the hermaphroditic fish, *Kryptolebias marmoratus*
 598 embryo. *Environ Toxicol Chem* 31:1745-1753.

599 Simon O, Floc'h E, Geffroy B, Frelon S, 2014. Exploring ecotoxicological fish bioassay for
 600 the evaluation of uranium reprotoxicity. *Environ Toxicol Chem* 33:1817-1824.

601 Simon O, Floriani M, Cavalie I, Camilleri V, Adam C, Gilbin R, Garnier-Laplace J. 2011a.
 602 Internal distribution of uranium and associated genotoxic damages in the chronically exposed
 603 bivalve *Corbicula fluminea*. *J Environ Radioact* 102:766-773.

604 Simon O, Gagnaire B, Camilleri V, Cavalié I, Floriani M, Adam-Guillermin C. 2018.
 605 Toxicokinetic and toxicodynamic of depleted uranium in the zebrafish, *Danio rerio*. *Aquat*
 606 *Toxicol* 197:9-18.

607 Simon O, Massarin S, Coppin F, Hinton TG, Gilbin R. 2011b. Investigating the embryo/larval
 608 toxic and genotoxic effects of γ irradiation on zebrafish eggs. *J Environ Radioact* 102:1039-
 609 1044.

610 Singh NP, McCoy MT, Tice RR, Schneider EL. 1988. A simple technique for quantitation of
 611 low levels of DNA damage in individual cells. *Exp Cell Res* 175:184-191.

612 Sinha P, Arunachalam KD, Annamalai SK. 2018. Radio-protective dosimetry of *Pangasius*
613 *sutchi* as a biomarker, against gamma radiation dosages perceived by genotoxic assays.
614 *Ecotoxicol Environ Saf* 164:629-640.

615 Vastenhouw NL, Cao WX, Lipshitz HD. 2019. The maternal-to-zygotic transition revisited.
616 *Development* 146: 161471.

617 Vicquelin L, Leray-Forget J, Peluhet L, LeMenach K, Deflandre B, Anschutz P, Etcheber H,
618 Morin B, Budzinski H, Cachot J. 2011. A new spiked sediment assay using embryos of the
619 Japanese medaka specifically designed for a reliable toxicity assessment of hydrophobic
620 chemicals. *Aquat Toxicol* 105:235-245.

621 Wafer LN, Jensen VB, Whitney JC, Gomez TH, Flores R, Goodwin BS. 2016. Effects of
622 environmental enrichment on the fertility and fecundity of zebrafish (*Danio rerio*). *J Am*
623 *Assoc Lab Anim Sci* 55:291-294.

624 Wang J, Zhu X, Zhang X, Zhao Z, Liu H, George R, Wilson-Rawls J, Chang Y, Chen Y.
625 2011. Disruption of zebrafish (*Danio rerio*) reproduction upon chronic exposure to TiO₂
626 nanoparticles. *Chemosphere* 83:361-467.

627 Woodhead DS, 1977. The effects of chronic irradiation on the breeding performance of the
628 guppy, *Poecilia reticulata* (osteichthyes Teleostei). *Int J Radiat Biol* 32:1-22.

629 Yum EHW, Li VWT, Choi VWY, Cheng SH, Yu KN. 2010. Effects of alpha particles on
630 zebrafish embryos. *Appl Radiat Isot* 68:714-717.

631

632

633

Figure legends

Figure 1. Exposure duration, conditions and endpoints for lowDR and highDR experiments.

For each exposure condition, a control condition was included. For the lowDR experiment (0.5, 5 mGy h⁻¹), adults were exposed over 24 days. Reproduction was assessed after 10 and 24 days of irradiation. The progeny were then placed in irradiated and non-irradiated (F1 Recovery) exposure conditions over 10 and 22 days. For the highDR experiment (50 mGy h⁻¹), adults were exposed over 10 days. The progeny were then placed in irradiated and non-irradiated (F1 Recovery) exposure conditions over 120 hours. 10d-irradiated adults were placed in non-irradiated exposure (F0 Recovery) over 63 days. Reproduction was assessed after 6, 36 and 63 days in F0 recovery conditions. The progeny were placed in non-irradiated exposure conditions over 96 hours for each time point (6, 36 and 63 days).

Figure 2. Fecundity (number of egg per female) of F0 adults measured after 10 days of exposure for both lowDR and highDR experiments (Treatment versus Control, *, Kruskal-Wallis p=0.03). n=number of spawns.

Figure 3. Individual (points) and average (histograms, means $\pm$ sd) survival rates (%) of F1 progeny at 24 and 48 h. F1 progeny came from adults irradiated during 10 days for all experiments. F1 R: F1 recovery condition, *i.e* progeny spawned by irradiated adults and placed in non-irradiated conditions.

For lowDR experiment (control, 0.5, 5, R0.5, R5 mGy h⁻¹), survival rate was calculated from 3 spawns (with 2 pseudo-replicates, 1 male+1 female).

656 For highDR experiment (control, 50, R50 mGy h⁻¹), survival rate was calculated from 6
657 spawns (1 male+1 female). (Kruskal-Wallis test: Treatment versus Control, a: p=0.024, b:
658 p=0.015, c: p=0.006, d: p=0.006; a>b>c>d).

659

660 Figure 4. Survival rate (%) of F1 progeny over time for lowDR experiment. Data are means ±
661 SD of two replicates of 80 eggs for 3 spawns per condition. (Kruskal-Wallis test: 5 versus
662 F1R, a: p=0.049, n=6).

663 A. F1 progeny coming from adults irradiated during 10 days. Immediately after spawning, F1
664 progeny was placed in irradiated and in non-irradiated (F1R Recovery) exposure conditions
665 during 22 days.

666 B. F1 progeny coming from adults irradiated during 24 days. Immediately after spawning, F1
667 progeny was placed in irradiated exposure conditions during 10 d.

668

669 Figure 5. Survival rate (% , n=6) of F1 progeny (3-4 hpf) over time. Progeny came from 10d-
670 irradiated adults for highDR experimentation. F1 Progeny were placed in irradiated and non-
671 irradiated (F1R, Recovery) exposure conditions over 120 h. (Kruskal-Wallis test: Treatment
672 versus Control, a: p=0.006, b: p=0.006, c: p=0.003, d: p=0.003; e=0.003, f: p=0.003).

673

674 Figure 6. Individual (points) and average (histograms, means ± SD) survival rates (% , n=4-6)
675 of F1 progeny after 24 and 48 h. Progeny came from adults irradiated during 10 days and then
676 placed in non-irradiated conditions (F0R, Recovery) during 63 d. (Kruskal Wallis test,

677 Treatment versus Control, a: $p=0.015$ $n=12$, b: $p=0.0060$ $n=11$, c: $p=0.05$ $n=8$, e: $p=0.0058$
678 $n=11$, f: $p=0.05$ $n=11$).

679

680 Figure 7. Body length (mm) and typical aspect of F1 progeny at 72 h. F1 progeny came from
681 adults exposed during 10d (Irr. 10d) and 10d followed by 6 days (F0 Irr. 10d+F0R 6d) and 36
682 days in non-irradiated (F0 Irr. 10d+F0R 36d) conditions. (Kruskal Wallis test, Treatment
683 versus Control, *, $p<0.05$, F0Irr. 10d $p=0.002$ $n=20$; 0.0049 $n=20$, F0 Irr. 10d+F0R 6d,
684 $p=0.00007$ $n=33$, F0 irr. 10d+F0R 36d, $p=0.077$, $n=85$).

685

686 Figure 8. ROS production measured in larval exposed during 4 d coming from F0 adults
687 exposed during 10 (A1) and 24d (A2) and in larval exposed during 10 d coming from F0
688 adults exposed during 24 d (B) of lowDR experiments. Nd: not determinate $n=20$ per
689 condition, Anova test *, $p<0.05$.

690 Figure 9. DNA damage (Tail moment) in F1 progeny after 1, 4 and 10d of exposure. Progeny
691 came from adults irradiated during 24d for lowDR experiment. Progeny (3-4 hpf) were placed
692 in irradiated and non-irradiated (F1R, Recovery) conditions. Kruskal Wallis test, *, $p<0.05$.

693

694 Figure 10. DNA damage (tail moment) in progeny (A: 1d, irradiated and non-irradiated
695 exposure (F1R, Recovery)) and in gonads of F0 adults (B: female, C: male) coming from
696 highDR experiment. Kruskal Wallis test: *, $p<0.05$.

Figure 1. Exposure duration, conditions and endpoints for the lowDR and highDR experiments. For each exposure condition, a control condition was included. For the lowDR experiment (0.5, 5 mGy h⁻¹), adults were exposed over 24 days. Reproduction was assessed after 10 and 24 days of irradiation. The progeny were then placed in irradiated and non-irradiated (F1 Recovery) exposure conditions over 10 and 22 days. For the highDR experiment (50 mGy h⁻¹), adults were exposed over 10 days. The progeny were then placed in irradiated and non-irradiated (F1 Recovery) exposure conditions over 120 hours. 10 d-irradiated adults were placed in non-irradiated (F0 Recovery) exposure conditions over 63 days. Reproduction was assessed after 6, 36 and 63 days in F0 recovery conditions. The progeny were placed in non-irradiated exposure conditions over 96 hours for each time point (6, 36 and 63 days).

Figure 2. Fecundity (number of eggs per female) of F0 adults measured after 10 days of exposure for both the lowDR and highDR experiments (Kruskal-Wallis test: Treatment versus Control, * $p=0.03$). n=number of spawns.

Figure 3. Individual (points) and average (histograms, means $\pm$ sd) survival rates (%) of the F1 progeny at 24 and 48 h. F1 progeny came from adults irradiated during 10 days for all experiments. F1 R: F1 recovery condition, i.e. progeny spawned by irradiated adults and placed in non-irradiated conditions.

For the lowDR experiment (control, 0.5, 5, R0.5, R5 mGy h⁻¹), the survival rate was calculated from 3 spawns (with 2 pseudo-replicates, 1 male+1 female).

For the highDR experiment (control, 50, R50 mGy h⁻¹), the survival rate was calculated from 6 spawns (1 male+1 female). (Kruskal-Wallis test: Treatment versus Control, a: $p=0.024$, b: $p=0.015$, c: $p=0.006$, d: $p=0.006$; $a>b>c>d$).

Figure 4. Survival rate (%) of F1 progeny over time for the lowDR experiment. Data are means $\pm$ SD of two replicates of 80 eggs for 3 spawns per condition. Kruskal-Wallis test: 5 versus F1R 5, a, $n=6$, $p=0.049$

A. F1 progeny coming from adults irradiated over 10 days. Immediately after spawning, the F1 progeny was placed in irradiated and in non-irradiated (F1R Recovery) exposure conditions over 22 days.

B. F1 progeny coming from adults irradiated over 24 days. Immediately after spawning, the F1 progeny was placed in irradiated exposure conditions over 10 d.

Figure 5. Survival rate (%) of 6 F1 progeny (3-4 hpf) over 120 h. The progeny came from 10 d-irradiated adults from the highDR experiment. The F1 Progeny were placed in irradiated and non-irradiated (F1R, Recovery) exposure conditions over 120 h. (Kruskal-Wallis test: Treatment versus Control, a: $p=0.006$, b: $p=0.006$, c: $p=0.003$, d: $p=0.003$; e: $p=0.003$, f: $p=0.003$).

Figure 6. Individual (points) and average (histograms, means $\pm$ SD) survival rates (%; n=4-6) of F1 progeny after 24 and 48 h. The progeny came from adults irradiated over 10 days and then placed in non-irradiated conditions (FOR, Recovery) over 63 d. Kruskal Wallis test: Treatment versus Control a: $p=0.015$ n=12, b: $p=0.0060$ n=11, c: $p=0.05$ n=8, e: $p=0.0058$ n=11, f: $p=0.05$ n=11.

Figure 7. Body length (mm) and typical aspect of F1 progeny at 72 h. The F1 progeny came from adults exposed over 10 d (Irr. 10 d) and 10 d followed by 6 days (F0 Irr. 10d+F0R 6d) and 36 days in non-irradiated (F0 Irr. 10d+F0R 36d) conditions. Kruskal Wallis test, Treatment versus Control *, $p < 0.05$, F0Irr. 10d $p = 0.002$ $n = 20$; 0.0049 $n = 20$, F0 Irr. 10d+F0R 6d, $p = 0.00007$ $n = 33$, F0 irr. 10d+F0R 36d, $p = 0.077$, $n = 85$.

Figure 8. ROS production measured in larvae exposed over 4 d coming from F0 adults exposed over 10 (A1) and 24 d (A2) and in larvae exposed during 10 d coming from F0 adults exposed over 24 d (B) of lowDR experiment. Nd: not determinate n=20 per condition, Anova test *, $p < 0.05$.

Figure 9. DNA damage (Tail moment) in the F1 progeny after 1, 4 and 10 d of exposure. The progeny came from adults irradiated over 24 d for the lowDR experiment. The progeny (3-4 hpf) were placed in irradiated and non-irradiated (F1R, Recovery) conditions. Kruskal Wallis test, Treatment versus Control *, $p < 0.05$.

Figure 10. DNA damage (tail moment) in the progeny (A: 1 d, irradiated and non-irradiated exposure (F1R, Recovery)) and in gonads of F0 adults (B: female, C: male) coming from the highDR experiment. Kruskal Wallis test, Treatment versus Control *, $p < 0.05$.