Pregnancy exposure to atmospheric pollution and meteorological conditions and placental DNA methylation

Emilie Abraham, Sophie Rousseaux, Lydiane Agier, Lise Giorgis-Allemand, Jörg Tost, Julien Galineau, Agnès Hulin, Valérie Siroux, Daniel Vaiman, Marie-Aline Charles, et al.

► To cite this version:

Emilie Abraham, Sophie Rousseaux, Lydiane Agier, Lise Giorgis-Allemand, Jörg Tost, et al.. Pregnancy exposure to atmospheric pollution and meteorological conditions and placental DNA methylation. *Environment International*, 2018, 118, pp.334-347. 10.1016/j.envint.2018.05.007 . hal-02325488

HAL Id: hal-02325488

<https://hal.science/hal-02325488>

Submitted on 22 Oct 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Environ Int. 2018 Sep;118:334-347. doi: 10.1016/j.envint.2018.05.007. Epub 2018 Jun 21.

Pregnancy exposure to atmospheric pollution and meteorological conditions and placental DNA methylation

Emilie Abraham^a, Sophie Rousseaux^a, Lydiane Agier^a, Lise Giorgis-Allemand^a, Jörg Tost^b, Julien Galineau^c, Agnès Hulin^d, Valérie Siroux^a, Daniel Vaiman^e, Marie-Aline Charles^f, Barbara Heude^f, Anne Forhan^f, Joel Schwartz^g, Florent Chuffart^a, Ekaterina Flin^a, Saadi Khochbin^a, Rémy Slama^a, Johanna Lepeule^a on behalf of the EDEN mother-child cohort study group.

^a Univ. Grenoble Alpes, Inserm, CNRS, IAB, 38000 Grenoble, France; ^b Laboratory for Epigenetics and Environment, Centre National de Genotypage, CEA – Institut de Génomique, Evry, France; ^c Air Lorraine, Nancy, France; ^d ATMO Poitou-Charentes, La Rochelle, France; ^e Genomics, Epigenetics and Physiopathology of Reproduction, Institut Cochin, U1016 Inserm – UMR 8104 CNRS – Paris-Descartes University, Paris; ^f Inserm U1153, Early Origins of Child Health and Development team, Research Center for Epidemiology and Biostatistics Sorbonne Paris Cité (CRESS), Paris Descartes University, Villejuif, France; ^g Department of Environmental Health, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA.

Corresponding author:

Johanna Lepeule, IAB – Inserm U1209, Site Santé - Allée des Alpes, 38700 La Tronche.
Telephone: +33 4 76 54 94 66 – E-mail: johanna.lepeule@univ-grenoble-alpes.fr

Competing Financial Interests: The authors declare they have no present or potential competing financial interests.

Funding: This work was made possible thanks to a grant from the Fondation de France (n°2012-00031593 and 2012-00031617). This work was also supported by AVIESAN (Alliance nationale pour les sciences de la vie et de la santé, n°ISP09_2014), the Fonds de Recherche en Santé Respiratoire, the Région Rhône-Alpes, and the National Agency for Research (ANR-13-CESA-0011-02).

Additional supports to SK team projects were from Fondation ARC “Canc’air” project (RAC16042CLA), Plan Cancer (CH7-INS15B66) and Plan Cancer (ASC16012CSA), the “Université Grenoble Alpes” ANR-15-IDEX-02, LIFE and SYMER. The EDEN cohort has been funded by Foundation for Medical Research (FRM), National Agency for Research (ANR), National Institute for Research in Public Health (IRESP: TGIR cohorte santé 2008 program), French Ministry of Health (DGS), French Ministry of Research, Inserm Bone and Joint Diseases National Research (PRO-A) and Human Nutrition National Research Programs, Paris–Sud University, Nestlé, French National Institute for Population Health Surveillance (InVS), French National Institute for Health Education (INPES), the European Union FP7 programmes (FP7/2007-2013, HELIX, ESCAPE, ENRIECO, Medall projects), Diabetes National Research Program (through a collaboration with the French Association of Diabetic Patients (AFD)), French Agency for Environmental Health Safety (now ANSES), Mutuelle Générale de l’Education Nationale (MGEN), French National Agency for Food Security, and the French-speaking association for the study of diabetes and metabolism (ALFEDIAM). Funders had no influence of any kind on analyses or results interpretation.

ABSTRACT

Background: *In utero* air pollution exposure represents a major health threat to the developing foetus. DNA methylation is one of the most well-known molecular determinants of the epigenetic status of cells. Blood DNA methylation has been proven sensitive to air pollutants, but the molecular impact on new-borns has so far received little attention.

Objectives: We investigated whether nitrogen dioxide (NO₂), particulate matter (PM₁₀), temperature and humidity during pregnancy are associated with differences in genome-wide placental DNA methylation levels.

Methods: Whole-genome DNA-methylation was measured using the Illumina's Infinium HumanMethylation450 BeadChip in the placenta of in 668 individuals enrolled in the EDEN cohort. We designed an original strategy using *a priori* biological information to focus on candidate genes with a specific expression pattern in placenta (active or silent) combined with an agnostic epigenome-wide association study (EWAS). We used robust linear regression to identify CpGs and differentially methylated regions (DMR) associated with each exposure during short- and long-term time-windows.

Results: The candidate genes approach identified nine CpGs mapped to 8 genes associated with prenatal NO₂, PM₁₀ and humidity exposure [false discovery rate (FDR) $p < 0.05$]. Among these, the methylation level of 2 CpGs located on *ADORA2B* remained significantly associated with NO₂ exposure during the 2nd trimester exposure in the EWAS (FDR $p < 0.05$). EWAS further revealed associations between exposure and variations of DNA methylation of 4 other CpGs. We further identified 27 DMRs significantly (FDR $p < 0.05$) associated with air pollutants exposure and 13 DMRs with weather conditions exposure.

Conclusions: These findings demonstrate that air pollutants exposure at levels commonly experienced in the European population are associated with placental gene methylation and provide some mechanistic insight into some of the reported effect of air pollutants on pre-

eclampsia. Indeed, the methylation of *ADORA2B*, a gene whose expression was previously associated with hypoxia and pre-eclampsia, was consistently found here sensitive to atmospheric pollutants. In addition, a number of air pollutants associated DMRs pointed to genes previously implicated in preeclampsia and metabolic syndrome.

Key words: mother-child cohort, placenta, air pollution, epigenetics, temperature, humidity

1. Introduction

Despite significant improvements in air quality in past decades, 50% of the population in 2014 in Europe live in areas that do not meet the World Health Organisation guidelines (World Health Organization, 2006) for particulate matter less than 10 microns (PM_{10}), 85% for particulate matter less than 2.5 microns ($PM_{2.5}$), and 7% for nitrogen dioxide (NO_2) (Ortiz, 2017). Ambient air pollution includes gaseous pollutants, such as nitrogen oxides, sulphur dioxide, ozone, benzene, as well as particulate matter of various sizes, which are a mixture of solid and liquid droplets including black carbon, metals, and polycyclic aromatic hydrocarbons. Air pollutants exposure during pregnancy is a major health threat to children as it can cross the placenta and expose the developing foetus (Valentino et al., 2015; Wick et al., 2010). Exposure to air pollutants during pregnancy has been associated to a range of adverse health outcomes both in the short-term, including low birth weight and preterm birth and in the long term, including infant reduced lung function and neurodevelopmental disorders (Chiu et al., 2016; Clifford et al., 2016; Jedrychowski et al., 2010; Stieb et al., 2012). Air pollution pregnancy exposure is also a threat to the pregnant woman and is likely a risk factor of preeclampsia (M. Pedersen et al., 2014). More recently, high- or low-ambient temperatures have also been suggested to play a role in adverse pregnancy outcomes (Beltran et al., 2014; Giorgis-Allemand et al., 2017; Kloog et al., 2015).

The placenta plays a key role in foetal programming by supporting both the health of the mother and the development of the foetus. It conveys nutrients and oxygen to the foetus and regulates gas and waste exchanges as well as hormone interactions (Murphy et al., 2006). Alterations in placental physiology and function, potentially driven by epigenetic changes, may impact the health of the future individuals during their childhood as well as into their adulthood. Exposure to chemically and/or physically inappropriate environmental conditions during pregnancy can affect the placental function by modifying its epigenome (Nelissen et al., 2011). Indeed epigenetic mechanisms control the timing and levels of gene expression, by defining the extent of their activation or by maintaining them repressed. Hence in utero environmental exposures could result in epigenetic modifications of the placenta at birth, including changes in the DNA methylation profile. As a transient organ, the placenta may provide a unique record of exposures specifically occurring during pregnancy. While several studies have reported changes in global and gene-specific methylation patterns from adult blood associated with air pollution (Madrigano et al., 2011; Panni et al., 2016), temperature and relative humidity exposure (Bind et al., 2014), the role of maternal exposure to such environmental factors on new-borns DNA methylation patterns has received little attention so far. In cord blood, methylation in mitochondria-related genes was associated with NO₂ exposure during pregnancy in a recent meta-analysis (Gruzieva et al., 2017). In placenta, a few studies have linked global DNA methylation and gene candidate methylation to air pollution exposure during pregnancy (Cai et al., 2017; Janssen et al., 2013). However, no study has yet investigated the relationship between exposure to air pollutants and meteorological conditions during pregnancy and placental methylation patterns at a genome-wide scale.

We hypothesized that maternal exposure to air pollutants and meteorological conditions could alter the placental function through modifications of DNA methylation. Therefore we

investigated the relationship of air pollutants and meteorological conditions with global DNA methylation and gene-specific methylation in placentas at birth using the Illumina 450k array. In order to identify potentially relevant changes in genomic methylation sites related to environmental exposures, we used two complementary approaches combining a concept-driven analysis with an Epigenome Wide Association Study (EWAS) (Figure 1). The concept-driven analysis is an original strategy relying on the hypothesis that sensitivity to DNA methylation variations induced by environmental factors may depend on the activity of the genomic region considered (Rousseaux et al., 2013). Indeed, a change in the DNA methylation profile affecting a region involved in the epigenetic control of the expression level of a gene actively repressed or activated during pregnancy is more likely to be biologically relevant than a change occurring in another region of the genome. Therefore, based on available expression data in placenta, we identified genes whose expression is either predominant in full-term placenta or undergoes significant variations during placenta development. In addition, using our methylation data, we also established a list of genes with highly methylated CpG-rich promoters. Focussing on these genes with a specific status or pattern of expression in the placenta, enabled us to explore the “basal epigenome dynamics – epigenome response” relationship in the placenta. We then used an agnostic EWAS design, in an attempt to confirm the findings of the concept-driven analysis and to identify new regions associated with maternal air pollutants exposure.

Figure 1. Workflow of the study

2. Methods

2.1. Study population

The EDEN mother-child cohort included 2002 pregnant women, mainly Caucasian, enrolled before 24 weeks of gestation in Nancy and Poitiers university hospitals, France, between 2003 and 2006 (Heude et al., 2015). Exclusion criteria were multiple pregnancies, pre-pregnancy diabetes, French illiteracy and planned move outside the region in the following 3 years. Residential addresses, lifestyle, demographic and medical data were collected by questionnaires and interviews during pregnancy and after delivery. Among the 1301 women for whom placenta samples were collected, we focused on 668 women. Placenta samples were collected at delivery by the midwife or the technician of the study using a standardized procedure. Samples of around 5mm x 5mm were carried out in the centre of the placenta on the foetal side and were immediately frozen at -80°C. The EDEN cohort received approval from the ethics committee (CCPPRB) of Kremlin Bicêtre and from the French data privacy institution “*Commission Nationale de l’Informatique et des Libertés*” (CNIL). Written consent was obtained from the mother for herself and for the offspring.

2.2. Placental DNA methylation assessment

156 DNA from placental samples was extracted using the QIAasympyphony instrument (Qiagen,
157 Germany). The DNA methylation analysis was performed by The Centre National de
158 Gnotypage (CNG, Evry, France). The DNA samples were plated onto 96-well or 48-well
159 plates. In total, nine plates including 64 chips were used. These plates were analyzed in 4
160 batches. The ratios for sex (boy/girl) and recruitment centre (Poitiers/Nancy) were balanced
161 for each chip. The Illumina's Infinium HumanMethylation450 BeadChip, representing over
162 485,000 individual CpG sites, was used to assess levels of methylation in placenta samples
163 following the manufacturer's instructions (Illuminas, San Diego, CA, USA). Raw signals of
164 450K BeadChips were extracted using the GenomeStudio^{} software (v2011.1. Illumina). The
165 DNA methylation level of each CpG was calculated as the ratio of the intensity of fluorescent
166 signals of the methylated alleles over the sum of methylated and unmethylated alleles (β
167 value). All samples passed initial quality control and had on average more than 98 % of valid
168 data points (detection p-value <0.01). A refined version of the Subset Quantile Normalization
169 (SQN) pipeline (Touleimat and Tost, 2012) including a revised annotation file (Price et al.,
170 2013) was used for data processing, correction and normalization. Intensity values were
171 corrected for potential biases in fluorescent dye intensity and background corrected using the
172 *lumi* R package (Du et al., 2008) as implemented in the SQN pipeline. Probes potentially
173 influenced by SNPs underlying the entire sequence of the probe (+1 or + 2 bases depending
174 on the Infinium probe type) that are present in the EUR population of the 1000 Genome
175 project (<http://www.1000genomes.org>) at a frequency of more than 5% were removed from
176 the analysis. Probes previously reported to map to several genomic regions were removed
177 (Y.-A. Chen et al., 2013). The SQN pipeline uses the intensity signals of high-quality (i.e. low
178 detection p-value) Infinium I probes as "anchors" to estimate a reference distribution of
179 quantiles for probes in a biologically similar context based on the annotation file (Touleimat
180 and Tost, 2012). This reference was then used to estimate a target distribution of quantiles for

InfII probes as a means to provide an accurate normalization of InfI/InfII probes and correct for the shift. SQN is performed for each individual separately. A principal component analysis as well as a hierarchical clustering were applied and showed no overall difference in the methylation patterns across participants samples and control samples, so that a quantile normalization was performed for between sample normalization. After quality control and normalization steps, there were 426,049 CpG sites left. Methylation beta values ranged from 0 to 1. Data points with a detection p-value >0.01 were excluded from subsequent analyses. To reduce the influence of potential outliers, we excluded data points below the 25th percentile - 3*interquartile ranges or above the 75th percentile + 3*interquartile ranges for each probe, which removed 0.4% of all methylation beta values across participants. CpGs with more than 25% of missing data were removed, leaving 425,878 CpG sites for statistical analyses.

Global methylation was also evaluated by measuring methylation in four CpG sites of repetitive Alu elements (*Alu*) and long interspersed nucleotide elements 1 (*LINE-1*) using a previously published pyrosequencing methylation assay (A. S. Yang et al., 2004). We then used the median percent methylation of the four CpG sites.

2.3. Cellular composition of placenta samples

Cellular composition of biological samples is a potential confounder in epigenetic epidemiology studies. In the absence of reference methylomes for placental tissue, we used a reference-free method, the RefFreeEWAS package available in R (Houseman et al., 2016), to deconvolute cell-type proportions from DNA methylation array data. The method relies on the identification of latent variables as surrogates for cell-type mixture. From the 10,000 most variable CpGs, we identified the optimal number of cell-types to be 6. We then used the 425,878 CpGs to estimate the proportion of each cell-type per sample.

2.4. Identification of genes with specific patterns of expression

We developed an approach relying on the integration of biological knowledge on the “epigenome-expression” relationship in the placenta. We hypothesised that genes and chromatin regions whose activity is controlled by DNA methylation as well as dynamically changing chromatin regions should be more prone to be affected by alterations of DNA methylation following environmental insults. We therefore focused on two categories of genes defined below: genes that are particularly *active* in the placenta and genes that are *silent* in the placenta but poised to be activated in response to external stimuli.

We defined placenta-predominantly expressed (“active”) genes as genes highly expressed in the placenta at the end of pregnancy, and with little or no expression in other tissues. This list was defined through the exploitation of public transcriptomic data (<http://www.ncbi.nlm.nih.gov/geo/>: GSE3526, GSE18809, GSE7434, GSE9984 studies) in which 575 genes showed a predominant expression in placentas at birth after normal pregnancies. A placenta predominant expression was defined by a mean level of expression value in placenta samples being above a threshold defined as the mean expression values + 2 standard deviations of a series of 112 adult non-germline/non-placenta tissues of various origins. The pre-treatment of transcriptomic data and the strategy to identify genes with a predominant expression in specific tissue types have been described previously (Rousseaux et al., 2013; Wang et al., 2015). These 575 placenta-predominantly expressed genes corresponded to 9,003 CpGs measured by the Illumina beadchip (Table S1).

Placenta DNA methylation-dependent repressed (“silent”) genes included genes whose regulatory region is heavily methylated in placenta, based on the assumption that a high level of methylation in CpG-rich regions (or CpG islands) in the regulatory elements of these genes could be associated with a methylation dependent repression. We then hypothesized that these genes could potentially be reactivated upon demethylation. There were 9,969 CpGs located in

CpG islands (and which were not in the CpGs list corresponding to placenta-predominantly expressed genes) and whose mean methylation level in our dataset was above 80% (Table S1). Annotation of CpGs was based on information available in the R/Bioconductor package *IlluminaHumanMethylation450kanno.ilmn12.hg19*, version 0.6.0.

2.5. Environmental exposures assessment

NO₂ and PM₁₀ hourly concentrations were modelled at the women's home addresses during pregnancy using the quasi-Gaussian Atmospheric Dispersion Modelling System (ADMS) Urban 3.1 (Cambridge Environmental Research Consultants, Cambridge, United Kingdom) (Sellier et al., 2014). The model was implemented with the same methodology for both Poitiers and Nancy and provides hourly estimates on a 40x40 meters grid. Mean daily ambient temperature (T, in °C) and relative humidity (RH, in %) were obtained from the monitors of the French weather network *Météo France*. For each woman, the nearest monitor to her home address was used. Because the most biologically relevant time period for the effect of air pollution and weather conditions on methylation is unknown, we considered different exposure windows throughout pregnancy over which we averaged levels of NO₂, PM₁₀, T and RH: day 1, 2 and 3 before delivery, one week before delivery, one month before delivery, each trimester of pregnancy and the whole pregnancy.

2.6. Statistical analyses

2.6.1. Global methylation

We examined the impact of environmental exposures on the overall methylation level across the entire genome using two approaches, one relying on repetitive elements *Alu* and *LINE-1*

and the other relying on the global profile of methylation levels estimated for the 425,878 CpGs.

Association of *Alu* and *LINE-1* methylation levels with air pollutants and meteorological exposures were investigated using robust linear regression, which accounts for potential outliers and heteroscedasticity in the data. We adjusted for the following *a priori* selected covariates: child sex, parity (0, 1, ≥ 2 children), maternal age at end of education (≤ 18 , 19-20, 21-22, 23-24, ≥ 25 years), season of conception, study centre (Poitiers and Nancy), maternal body mass index before pregnancy (continuous), maternal age at delivery (linear and quadratic terms), maternal smoking during pregnancy (continuous) and gestational duration (linear and quadratic terms). We additionally adjusted for technical factors related to the methylation measurements (batch, plate and chip) and for the estimated cell-type proportions. Meteorological exposures - temperature and humidity (restricted cubic splines) - were investigated simultaneously. Each air pollutant was studied separately and further adjusted for meteorological exposures (temperature and humidity, with restricted cubic splines).

Complementary to the repetitive elements analysis, we conducted a global analysis of methylation profiles (GAMP). This method, described by Zhao et al. (2015), allows to test whether exposure changes the overall profile or distribution of CpG measurements of the participants instead of examining the effect on individual CpGs. Each individual's methylation profile was characterized by approximating the density function and the cumulative distribution function of the methylation distribution using B-splines basis functions. The B-spline coefficients were then used to represent each individual overall methylation distribution. A variance component score test from the kernel machine framework, which naturally accommodates the correlation between B-spline coefficients, was computed to test the association of B-spline coefficients with each exposure of interest (temperature, humidity, NO₂, PM₁₀) adjusted for confounders. Meteorological exposures were investigated

simultaneously. Each air pollutant was studied separately in models further adjusted for meteorological exposures (temperature and humidity).

2.6.2. Methylation at specific loci

We studied the association between environmental exposures (NO₂, PM₁₀, temperature, humidity) and CpG-specific methylation level using a robust linear regression model. We applied the same approach described for global methylation analysis by testing: the simultaneous effect of meteorological conditions (temperature and humidity) (1); the effect of each air pollutant adjusted for meteorological conditions (temperature and humidity) (2) using the following models:

$$(1) Y_{ij} = \beta_0 + f(\beta_1 \cdot \text{Temperature}_i) + f(\beta_2 \cdot \text{Humidity}_i) + \beta_Z^T Z_i + \varepsilon_{ij}.$$

$$(2) Y_{ij} = \beta_0 + \beta_1 \text{Pollutant}_i + f(\beta_2 \cdot \text{Temperature}_i) + f(\beta_3 \cdot \text{Humidity}_i) + \beta_Z^T Z_i + \varepsilon_{ij}.$$

Where Y_{ij} is the methylation measurement for CpG j in subject i , $f(\beta_2 \cdot \text{Temperature}_i)$ and $f(\beta_2 \cdot \text{Humidity}_i)$ are restricted cubic spline functions of temperature and humidity, Z_i is the set of adjustment factors similar to the one used for the global methylation analysis and ε_{ij} is the random error. Each exposure window was tested in a separate model. Multiple testing was accounted for by applying a Benjamini and Hochberg False Discovery Rate (FDR) correction to the p-values that were obtained (Benjamini and Hochberg, 1995). The FDR corrected p-values were calculated for the 18,972 CpGs included in the *active* and *silent* genes lists for the concept-driven analysis and for the 425,878 CpGs for the agnostic EWAS. An FDR-corrected p-value <0.05 was considered statistically significant. For all significant CpGs (FDR-corrected p-value <0.05), we further tested whether the association with the exposures differed by child sex by adding an interaction term in the

model. The genomic inflation factor (λ) has been widely used in GWAS to assess inflation/deflation of p-values (Devlin et al., 2001). A recent study demonstrated that the Bayesian Inflation Factor (BIF) was a more relevant approach for EWAS (van Iterson et al., 2017). We computed the λ and BIF for both the concept-driven analysis and the agnostic EWAS.

2.6.3. Region-based analysis

To identify Differentially Methylated Regions (DMRs) in our data, we used Comb-p, a method relying on the Stouffer-Liptak-Kechris correction that combines specific CpG p-values using sliding windows and accounting for correlation between CpGs (B. S. Pedersen et al., 2012). DMR p-values were adjusted for multiple testing by Šidák correction (Šidák, 1967). Significant DMRs (p-value<0.05) included at least 2 probes (p-value<0.001) at a maximum distance of 500 bp. All analyses were performed using the statistical software R (version 3.0.1) and Python (version 2.7.14).

3. Results

3.1. Population characteristics and environmental factors

On average ($\pm$ SD), the participating mothers were 29 ($\pm$ 5.1) years old, with a pre-pregnancy BMI of 22.9 ($\pm$ 4.2) kg/m² (Table 1). Mean gestational duration was 40 ($\pm$ 1.7) weeks and 32 babies (5%) were born preterm (<37 gestational weeks).

Table 1. Characteristics of the EDEN study population (n=668) and environmental exposures over the whole pregnancy period.

	mean $\pm$ SE	n (%)	Median pregnancy levels			
Characteristics		All	T	RH	NO ₂	PM ₁₀
Center						
Poitiers		287 (43)	12.2	77.3	12.2	15.6
Nancy		381 (57)	9.8	77.2	23.4	23.4
Sex of offspring						

Male	349 (52)	10.7	77.4	17.1	20.6
Female	319 (48)	11.1	76.8	17.4	21.5
Parity					
0	301 (45)	11.3	77.3	17.0	20.8
1	252 (38)	10.5	77.4	17.3	21.2
≥2	115 (17)	11.0	76.9	17.4	21.0
Maternal age at end of education (year)					
≤18	132 (20)	10.9	77.2	17.4	20.7
19-20	114 (17)	11.1	77.5	15.8	17.5
21-22	151 (23)	10.9	77.1	16.9	21.3
23-24	145 (22)	11.0	76.7	18.2	21.3
≥25	126 (19)	10.8	77.8	18.8	21.4
Season of conception					
January – March	143 (21)	13.5	73.8	14.6	17.2
April – June	153 (23)	12.3	77.3	18.6	21.8
July – September	189 (28)	8.7	80.1	19.3	22.0
October – December	183 (27)	10.1	76.7	16.9	20.5
BMI (kg/m²)	22.9±4.2	-	-	-	-
Maternal age (year)	29.0±5.1	-	-	-	-
Maternal smoking^a	1.7±3.5	-	-	-	-
Gestational duration (weeks)	39.8±1.7	-	-	-	-

T= Temperature (°C). RH= Relative humidity (%). NO₂= Nitrogen dioxide (µg/m³). PM₁₀= Particulate matter smaller than 10 µm (µg/m³). BMI= pre-pregnancy Body Mass Index.

^aAverage number of cigarette per day during pregnancy

The longer the exposure window was, the less variable was the exposure level (Figure 2).

Mean pregnancy levels were 19 (± 7.7) µg/m³ for NO₂ and 20 (± 4.5) µg/m³ for PM₁₀.

Figure 2. Environmental exposure across time window (d1, 2, 3 = day 1, 2, 3 before delivery; w = week before delivery; m = month before delivery; t1, 2, 3 = trimester 1, 2, 3 and P = whole pregnancy).

NO_2 and PM_{10} levels were highly correlated (range, 0.73 to 0.84 depending on averaging period, see Table S2). For each air pollutant, trimester-specific levels were strongly and

positively correlated (correlation ranged from 0.52 to 0.78 for NO₂ and from 0.79 to 0.86 for PM₁₀). Average temperature exposure during pregnancy varied from 5 °C minimum to 16 °C maximum while relative humidity varied from 66% minimum to 86% maximum. Temperature and humidity were strongly and negatively correlated (range between -0.71 and -0.80, depending on averaging period, see Table S2). As for correlation between air pollutants and meteorological conditions, NO₂ levels were moderately correlated with temperature and humidity ($|r|=0.15$ to 0.61) and PM₁₀ levels were weakly correlated with temperature and humidity ($|r|=0.04$ to 0.34).

3.2. Global DNA methylation

Average methylation level was 16.2 (± 1.0) for Alu and 26.1 (± 1.9) for LINE-1. The distribution of average methylation level by CpG site was bimodal, with a peak around 0.025 and another around 0.925 (Figure S1). Highest variability in methylation levels was observed for moderately methylated CpG sites (see Figure S2). PM₁₀ exposure the day before birth was positively associated with Alu methylation with a 0.08 regression coefficient (p-value=0.01) (Table S3). Temperature exposure during the 1st trimester of pregnancy was significantly associated with the density of methylation distribution (p-value=0.03). However, analyses of Alu and LINE-1 as well as the global analysis of methylation profiles mainly yielded not significant differences with p-values higher than 0.05 (Table S3).

3.3. Concept-driven approach: *silent* and *active* genes in placenta

Among the list of placenta-predominantly expressed genes (“active genes”), 3 CpGs mapping to 3 genes, whose average methylation levels varied between 0.03 and 0.58, were significantly associated with NO₂ exposure (FDR p-values <0.05, Table 2). Two of these 3

CpGs are located in the body of *ADORA2B*, encoding the Adenosine A2B Receptor, one is in a region shared by the promoter regions of two genes, *PXT1* (Peroxisomal testis specific 1) and *KCTD20* (potassium channel tetramerization domain containing 20). The methylation levels of the two CpGs of *ADORA2B* decreased on average with higher exposure to NO₂ during the 1st or 2nd trimester or during the whole pregnancy. Conversely, the methylation of the CpG located in the promoter of the *PXT1* and *KCTD20* genes was positively associated with NO₂ exposure during both the 2nd trimester and the whole pregnancy. Regarding exposure to PM₁₀, one CpG located in the promoter of *SLC44A5B* (solute carrier family 44 member 5) gene was positively associated with exposure the month before birth and another CpG located in the body of *KYNU* (kynureninase) was negatively associated with exposure 3 days before birth. No significant association with methylation was found for temperature and humidity levels (lowest FDR p-values, 1.00, 1.00, respectively).

Among the placenta DNA methylation-dependent repressed genes (“silent genes”), one CpG, located in the body of the *CAPN10* (Calcium-Activated Neutral Proteinase 10) gene, was positively associated with NO₂ exposure during the 3rd trimester. Exposure to PM₁₀ the month before birth was negatively associated with mean methylation of one CpG located in the body of *TUBGCP2* (tubulin gamma complex associated protein 2) and was positively associated with mean methylation of two CpGs, one located in the body of *TGM6* (transglutaminase 6) and another located in the body of *ADCK5* (aarF domain containing kinase 5). Regarding exposure to meteorological conditions, no significant associations with methylation of active genes were found (lowest FDR p-values, 1.00 for temperature and humidity).

The p-value distribution of the 18,972 CpGs included in the concept-driven analysis was close to the theoretical distribution as indicated by the BIF values (range from 0.95 to 1.1) which were substantially smaller compared to the lambda values (figures S3-S6).

383 Table 2. CpGs associated (FDR $p < 0.05$) with prenatal exposure to air pollutants or meteorological conditions

Exposure	CpG	Chr	Position (built 37)	Gene	Location in gene	Location of CpG	β^a	FDR p-value candidate	EWAS	Mean methylation level (SE)
<i>CpGs located in placenta-predominantly expressed genes (n=9,003 sites)</i>										
NO₂										
Pregnancy	cg10984505	chr6	36410951	PXT1;KCTD20	TSS1500;5'UTR	Island	0.002	0.04		0.031 (0.006)
	cg17580614	chr17	15849512	ADORA2B	Body	Island	-0.042	$<10^{-3}$	0.02	0.586 (0.138)
	cg07563400	chr17	15849556	ADORA2B	Body	S_Shore	-0.045	$<10^{-3}$		0.343 (0.125)
Trimester 1	cg17580614	chr17	15849512	ADORA2B	Body	Island	-0.037	$<10^{-3}$		0.586 (0.138)
	cg07563400	chr17	15849556	ADORA2B	Body	S_Shore	-0.042	$<10^{-3}$		0.343 (0.125)
Trimester 2	cg10984505	chr6	36410951	PXT1;KCTD20	TSS1500;5'UTR	Island	0.002	0.02		0.031 (0.006)
	cg17580614	chr17	15849512	ADORA2B	Body	Island	-0.044	$<10^{-4}$	$<10^{-3}$	0.586 (0.138)
	cg07563400	chr17	15849556	ADORA2B	Body	S_Shore	-0.047	$<10^{-4}$	$<10^{-3}$	0.343 (0.125)
PM₁₀										
Month before birth	cg12659128	chr1	76078176	SLC44A5	TSS1500	N_Shelf	0.037	0.03		0.587 (0.070)
Day 3 before birth	cg04112100	chr2	143701758	KYNU	Body	OpenSea	-0.012	0.04		0.257 (0.068)
<i>CpGs located in methylation-dependent repressed genes list (n=9,969 sites)</i>										
NO₂										
Trimester 3	cg01712700	chr2	241535695	CAPN10	Body	Island	-0.004	0.02		0.948 (0.013)
PM₁₀										
Month before birth	cg23075260	chr8	145617435	ADCK5	Body	Island	0.018	0.03		0.878 (0.034)
	cg05142592	chr10	135097800	TUBGCP2	Body	Island	-0.008	0.03		0.953 (0.018)
	cg06967014	chr20	2384240	TGM6	Body	Island	0.007	0.03		0.924 (0.015)
<i>CpGs not in active and silent placenta genes list (i.e. EWAS approach)</i>										
NO₂										
Pregnancy	cg20491726	chr2	242820622			N_Shelf	0.015		0.02	0.715 (0.051)
Trimester 2	cg20491726	chr2	242820622			N_Shelf	0.015		0.02	0.715 (0.051)
PM₁₀										
Trimester 1	cg03215416	chr8	18823341	PSD3	Body	OpenSea	-0.020		0.03	0.913 (0.024)
Humidity										
Trimester 2	cg16917193	chr12	54089295			Island			0.04 ^b	0.104 (0.114)
Month before birth	cg16075020	chr19	12606183	ZNF709	5'UTR	Island			$<10^{-3b}$	0.063 (0.077)

NO₂= Nitrogen dioxide ($\mu\text{g}/\text{m}^3$). PM₁₀= Particulate matter smaller than 10 μm ($\mu\text{g}/\text{m}^3$).

^a Results per 10 units increase in prenatal exposure.

^b p-value of the spline

387 3.4. Agnostic approach: EWAS examining each CpG

388 When we independently tested the adjusted effect of exposures on each of the 425,878 CpGs, six
389 showed significantly different methylation levels (Table 2). Among these, two CpGs had also been
390 identified in the concept-driven analysis: cg07563400 and cg17580614, both located in *ADORA2B*,
391 remained significantly associated with NO₂ exposure during the 2nd trimester of pregnancy after the
392 genome-wide FDR correction of p-values (Table 2; Figure 3). Other CpGs identified as
393 significantly associated with environmental exposures in the concept-driven approach did not
394 achieve genome-wide statistical significance in the EWAS, but all except one of them (cg05142592
395 on gene *TUBGCP2*) were still in the top 20 CpGs with the lowest p-values for association with
396 exposure (see Table S4 to S7).

397 The EWAS also revealed four new CpGs (not included in the concept-driven list of candidates)
398 whose average methylation was significantly (FDR p-value <0.05) associated with humidity, NO₂
399 and PM₁₀ exposure (Table 2). One CpG (cg20491726), mapped to a non genic region, was
400 significantly associated with a 10 µg/m³ increase in NO₂ exposure during both the second trimester
401 ($\beta=0.015$ FDR p=0.02) and the whole pregnancy ($\beta=0.015$ FDR p=0.02). Another CpG mapped to
402 *PSD3* (pleckstrin and Sec7 domain containing 3) was negatively associated with PM₁₀ during the 1st
403 trimester of pregnancy (Table 3). Two CpGs, one in the non genic region and one mapped to
404 *ZNF709* (zinc finger protein 709), were non-linearly associated with humidity in the 2nd trimester of
405 pregnancy and in the month before birth, respectively (Figure S7). The p-value distribution of the
406 425,878 CpGs was close to the theoretical distribution as indicated by the BIF values (range from
407 0.96 to 1.15) (Figure S8 to S11).

408 Although interaction terms with sex were significant for two CpGs (cg06967014, cg16917193), the
409 effect estimates (Table 8) and dose-response relationship (Figure S7) did not appear substantially
410 different between boys and girls.

411

412 Figure 3. Manhattan plots of p-values showing the association between environmental exposure
 413 (NO₂ (A, B), PM₁₀ (C), and humidity (D, E)) and 425,878 CpGs methylation using the agnostic
 414 EWAS. Each dot corresponds to the p-value of a CpG site and the horizontal lines indicates the
 415 level of statistical significance (FDR p < 0.05).

3.5. Regional analysis

418 Our regional analysis identified 27 DMRs significantly (Sidak-corrected P-value<0.05) associated
 419 with air pollutants exposure and 13 DMRs with weather conditions exposure (Table 3, Table S9).
 420 The *ADORA2B*-associated DMR was identified for the three trimesters and the whole pregnancy;
 421 the DMR included the two CpGs previously identified in the site-specific analyses. Among genes
 422 highly represented in the identified DMRs, several CpGs located in the *FAM38A* gene were

423 negatively associated with NO₂ exposure in each trimester, last month and pregnancy. The
424 *FAM124B*-associated DMR was hypomethylated across the 7 CpGs included in women with
425 increased exposure to NO₂ or PM₁₀, especially towards the end of pregnancy. NO₂ exposure,
426 especially towards the end of pregnancy (3rd trimester and last month) was negatively associated
427 with two DMRs located in *HSP90AA1* and *LOC254559* and positively associated with two DMRs
428 located in *LZTFL1*, and *P2RX4*. The top-ranked DMRs in terms of number of CpGs included were
429 located in *CD81* (13 CpGs) and in a non-genic region on chromosome 6 (15 CpGs).

430

431

432 Table 3. Regions identified using the Comb-P method as differentially methylated in the placenta in
 433 association with air pollutants exposure in pregnancy.

Gene ¹	Hg19	Exposure	Time window	No. of probes	Slk p-value	Sidak p-value	Direction of effect
ADORA2B	chr17:15849512-15849557	NO ₂	p	2	4.93E-12	4.67E-08	-
		NO ₂	t1	2	9.18E-12	8.69E-08	-
		NO ₂	t2	2	1.29E-14	1.22E-10	-
		NO ₂	t3	2	3.29E-09	3.11E-05	-
ANKRD45	chr1:173639044-173639136	NO ₂	m	4	1.61E-08	7.43E-05	-
		pm10	m	4	2.99E-08	1.39E-04	-
B3GALT4	chr6:33245717-33245805	NO ₂	p	6	1.04E-05	4.90E-02	-
C1orf86;LOC100128003	chr1:2125049-2125244	NO ₂	p	7	7.43E-08	1.62E-04	-
C22orf9	chr22:45608345-45608517	PM ₁₀	d2	7	4.92E-08	1.22E-04	+
CD81	chr11:2397486-2397832	PM ₁₀	d1	13	1.37E-16	1.37E-13	+
DAXX	chr6:33288180-33288373	PM ₁₀	d1	6	1.41E-10	3.10E-07	-
FAM124B	chr2:225266656-225266881	NO ₂	p	7	1.41E-08	2.67E-05	-
		NO ₂	t3	7	6.55E-11	1.24E-07	-
		NO ₂	m	7	7.62E-10	1.44E-06	-
		PM ₁₀	t3	7	4.80E-11	9.09E-08	-
FAM38A	chr16:88832476-88832573	NO ₂	t3	4	1.05E-08	4.62E-05	-
		NO ₂	m	4	1.51E-09	6.63E-06	-
FAM38A	chr16:88832476-88832701	NO ₂	p	6	1.16E-09	2.20E-06	-
		NO ₂	t1	6	6.15E-11	1.16E-07	-
FAM38A	chr16:88832532-88832573	NO ₂	t2	2	4.61E-07	4.78E-03	-
GNAS;GNASAS	chr20:57426240-57426275	NO ₂	p	3	1.64E-07	1.99E-03	+
GP1BB;SEPT5	chr22:19710880-19711052	PM ₁₀	d2	6	1.96E-09	4.86E-06	-
HSP90AA1	chr14:102554826-102554978	NO ₂	t3	4	8.60E-08	2.41E-04	-
		NO ₂	m	4	6.66E-09	1.87E-05	-
IL32	chr16:3114986-3115134	NO ₂	m	3	4.45E-09	1.28E-05	+
LIME1	chr20:62369445-62369606	PM ₁₀	d2	4	1.38E-08	3.64E-05	-
LMF1	chr16:967929-968025	NO ₂	t3	3	5.54E-08	2.46E-04	-
LOC254559	chr15:89921083-89921259	NO ₂	t3	6	3.97E-08	1.64E-04	-
		NO ₂	m	7	2.02E-09	4.90E-06	-
LZTFL1	chr3:45883529-45883735	NO ₂	p	7	1.16E-07	2.39E-04	+
		NO ₂	t3	8	2.75E-10	4.37E-07	+
		NO ₂	m	10	1.51E-11	1.94E-08	+
NOTCH3	chr19:15281200-15281518	PM ₁₀	d1	4	5.03E-12	6.73E-09	-
P2RX4	chr12:121647128-121647218	NO ₂	p	5	7.16E-08	3.39E-04	+
		NO ₂	t3	5	8.03E-08	3.80E-04	+
		NO ₂	m	5	5.51E-11	2.61E-07	+
PRRT1	chr6:32116538-32116781	PM ₁₀	d2	5	3.64E-09	6.38E-06	-
SOX2OT;SOX2	chr3:181428462-181428580	PM ₁₀	m	5	6.05E-08	2.18E-04	+
TRIO	chr5:14359544-14359615	PM ₁₀	m	3	1.44E-08	8.66E-05	+
ZNF563	chr19:12444484-12444742	PM ₁₀	m	6	3.21E-11	5.29E-08	+
	chr3:194650099-194650198	NO ₂	t3	2	6.78E-08	2.92E-04	+
	chr6:28583971-28584173	NO ₂	m	15	6.42E-09	1.35E-05	-

434 Hg19: Human Genome version 19; Slk: Stouffer-Liptak Kechris correction; NO₂= Nitrogen dioxide
 435 (µg/m³). PM₁₀= Particulate matter smaller than 10 µm (µg/m³). p: pregnancy. t1: trimester 1. t2:

trimester 2. t3: trimester 3. d1: day 1 before birth. d2: day 2 before birth. d3: day 3 before birth;¹
UCSC;² direction of effect was similar across all CpGs included in the DMR

3.6. Synthesis of the results

Our top finding was that one CpG (cg17580614) located in the body of *ADORA2B*, one of the placenta-predominantly expressed genes, showed a significantly lower methylation in placenta after NO₂ exposure during pregnancy. This result came out of the concept-driven analysis performed on 18,972 CpGs and was confirmed in the agnostic approach that used a more stringent FDR correction of the p-values on 425,878 CpGs and in the regional analysis. Furthermore, this CpG as well as another *ADORA2B* CpG (cg07563400) were in the top 20 of CpGs associated with NO₂ exposure during the 1st and 2nd trimesters and during the whole pregnancy (FDR p-values ≤0.1, Supplemental material, Table S6). Figure 4 shows the regression coefficient for each of the *ADORA2B* CpGs against its uncorrected p-value regarding the association with NO₂ exposure during the 2nd trimester and the whole pregnancy. Among the 19 CpGs of this gene, 16 were negatively associated with NO₂ exposure during the 2nd trimester and the whole pregnancy, and 11 (10, respectively) were in the top 1000 of the CpGs associated with placenta-predominantly expressed genes (Figure 4).

Figure 4: Visualisation of regional associations between NO₂ exposure during pregnancy and *ADORA2B* methylation indicating the direction of the effect (coefficient) and confidence interval for each CpG .

457 5. Discussion

458 To the best of our knowledge, the present study shows the results of the first genome-wide analysis
459 of differentially methylated probes and regions in placenta in relation to in utero exposure to PM₁₀,
460 NO₂, temperature and humidity. Associating a concept-driven analysis with an agnostic EWAS, we
461 showed lower methylation levels in two CpGs located in the body of *ADORA2B* associated with
462 increased exposure to NO₂ during pregnancy, especially in trimesters 1 and 2. The agnostic EWAS
463 further revealed new significant associations of prenatal exposure to PM₁₀, NO₂, humidity and
464 temperature during various time-windows and methylation of 4 CpGs (mapped to 2 genes (*PSD3*,
465 *ZNF709*) and 2 non genic regions) and 39 regions (Table 3, Table S9), which are candidates for
466 replication in future studies. Among these, several DMR were located in genes pointing towards
467 hypertensive and metabolic disorders.

468 *ADORA2B*, encoding the Adenosine A2B receptor, was shown to play a role in placental
469 development and possibly in the pathophysiology of hypoxia and preeclampsia (PE) in pregnant
470 women (Acurio et al., 2014; Darashchonak et al., 2014; Jia et al., 2012), a pathology which has also
471 been associated with air pollution exposure during pregnancy (M. Pedersen et al., 2014). In both
472 mice and humans, it has been demonstrated that activation of ADORA2B signalling contributes to
473 the pathogenesis of PE (Huang et al., 2017), to small foetuses, small placentas and foetal growth
474 restriction (Iriyama et al., 2015). The molecular basis of the functional link and relationship
475 between *ADORA2B* gene and placenta DNA methylation, and the involvement of the ADORA2B
476 protein in PE and birth outcomes are questions to be addressed in future investigations. Additional
477 work will also be required to assess the relevance of *ADORA2B* as a candidate marker and/or target
478 for the diagnosis, prevention and treatment of PE.

479 Of interest, four DMR were located in *CD81*, *DAXX*, *NOTCH3* and *P2RX4* genes, which have all
480 been implicated in PE phenotypes. *P2RX4*, a purinergic receptor highly expressed in normal
481 placentas, was shown to be overexpressed in both term and preterm preeclamptic placentas (Roberts

et al., 2007) and hypomethylated in preeclamptic placentas (Chu et al., 2014). In our study, NO₂ exposure in pregnancy, 3rd trimester and last month of pregnancy was positively associated with *P2RX4* methylation. In accordance with our result, a CpG of *P2RX4* was found hypermethylated in blood from adults exposed to PM_{2.5} in the previous month (Panni et al., 2016). In addition, our results indicated a positive association of PM₁₀ exposure the day before birth with methylation in a DMR located in *CD81* and a negative association with DMRs in *DAXX* and *NOTCH3*. *CD81*, a gene encoding a member of the transmembrane 4 superfamily, was shown to be mainly expressed in normal first trimester placentas and gradually downregulated with gestational age (Shen et al., 2017). In early-onset (≤ 34 weeks) severe preeclamptic placentas, *CD81* was up-regulated in syncytiotrophoblasts and extravillous cytotrophoblasts, which play a major role in maternal–fetal exchanges (Malassiné and Cronier, 2002) and placentation (Pijnenborg et al., 2006). The overexpression of *CD81* was also demonstrated in a PE-like phenotype in rats (Shen et al., 2017). A recent study demonstrated that *DAXX* methylation was involved in trophoblast differentiation, PE, and response to hypoxia, suggesting an important role of *DAXX* in placentation (Novakovic et al., 2017). Similarly, *NOTCH3* placental expression has been shown down regulated in PE (Fragkiadaki et al., 2015) and up regulated in early onset PE (Zhao et al., 2014). Notch signaling has been suggested to play a key role in cytotrophoblasts motility and differentiation (Haider et al., 2014). In contrast to our finding, *NOTCH3* methylation in blood from adults has been positively associated with PM_{2.5} exposure the previous month (Panni et al., 2016). Further studies are required in order to determine the functional relationship between these genes and their placental methylation. Hypertensive disorders, among which preeclampsia in pregnant women (M. Pedersen et al., 2014), are increasingly suspected as a potential mechanism to explain health effects of air pollutants (Shah et al., 2013).

Among air pollutants associated DMR in our study, four were located in *LMF1*, *GNAS*, *GNASAS*, and *IL-32* genes. These genes have been previously involved in metabolic syndrome and related phenotypes (Hong et al., 2017). Single nucleotide polymorphisms of *LMF1* is involved in severe

508 hypertriglyceridemia (Péterfy, 2012). *LMF1* blood methylation in adults has been positively
509 associated with PM_{2.5} exposure in the previous month (Panni et al., 2016), while our results
510 indicated a negative association of NO₂ exposure in the 3rd trimester with placental methylation of
511 this gene. *GNAS* is an imprinted gene with a complex imprinted pattern ; decreased placental
512 expression has been associated with intra uterine growth retardation (Dunk et al., 2012; McMinn et
513 al., 2006) and cord blood hypermethylation has been associated with gestational diabetes (D. Chen
514 et al., 2014). *GNASAS*, the *GNAS* antisense RNA 1, placental expression was related to newborns
515 neurobehavioral profile (Green et al., 2016), while its methylation in adult blood has been linked to
516 prenatal exposure to famine (Tobi et al., 2009). In rabbits, pregnancy exposure to diesel engine
517 exhaust has been associated with changes in triglyceride and cholesterol levels in the offspring
518 exposed *in utero* (Valentino et al., 2015) but further investigations are required to identify if this
519 association could be to some extent mediated by epigenetic changes. Taken together our findings
520 point to PE- and metabolic syndrome-associated genes, which is in line with the increasing
521 evidence on the relationship between air pollution and the metabolic syndrome (B.-Y. Yang et al.,
522 2018) and with the hypothesis that such effects might start early in life (Park, 2017).

523 Significant variations of the methylation levels of other CpGs and DMR, identified either from the
524 concept-driven analysis or from the agnostic EWAS approach, have occasionally been found
525 associated with NO₂, PM₁₀, temperature or humidity exposure at specific time windows of
526 pregnancy. However, the biological significance of the differential methylation observed for these
527 genes is unclear (Supplemental material, Table S10). These genes have not been associated with
528 NO₂ or proximity to traffic in the two recent EWAS conducted in newborns (Gruzieva et al., 2017;
529 Kingsley et al., 2016). Noteworthy, some of these results were in agreement with air pollution
530 association studies conducted on blood DNA methylation. This was the case for the positive
531 association of methylation in *LZTFL1*-DMR with average NO₂ exposure during pregnancy, 3rd
532 trimester and last month of pregnancy which was consistent with the relationship between *LZTFL1*
533 blood methylation and chronic PM_{2.5} exposure observed in an elderly population (Nwanaji-

534 Enwerem et al., 2016). The significant (FDR corrected) associations of PM_{2.5} exposure in the past
535 month and decreased blood methylation in *ANKRD45* and *LOC254559* and increased blood
536 methylation in *TRIO* found by Panni (Panni et al., 2016) was in agreement with our findings in the
537 placenta. However, the negative relationship they found between PM_{2.5} exposure and *ZNF563*
538 methylation disagree with our results. The small overlap between our results and those of previous
539 published EWAS can be explained by the fact that these studies relied on different tissues. Gruzieva
540 et al. investigated cord blood (Gruzieva et al., 2017) and Kingsley et al. examined placenta samples
541 from the maternal side (Kingsley et al., 2016). DNA methylation is tissue-specific (Ollikainen et
542 al., 2010) and it is likely that different tissue will respond differently to similar environmental
543 stimuli. The fact that DNA methylation is a dynamic mark also explains this absence of overlap
544 between the present results obtained in our population of newborns and other data obtained for
545 different populations and exposure levels.

546 Breton et al. (Breton et al., 2016) found that prenatal exposure to PM₁₀ in the first trimester was
547 associated with lower LINE-1 methylation in 459 newborn bloodspots. A similar result was found
548 in 471 placenta samples where LINE-1 was associated with distance to traffic (Kingsley et al.,
549 2016) and in 240 placenta samples where global methylation was linked to first trimester exposure
550 to PM_{2.5} (Janssen et al., 2013). These studies did not investigate shorter time-windows of exposure.
551 In our study, we also observed a negative relationship between PM₁₀ exposure during the first
552 trimester and LINE-1 and Alu CpG methylation, but our results did not reach statistical
553 significance. In the short-term, PM₁₀ exposure the day before birth was significantly and positively
554 associated with Alu repetitive elements methylation.

555 As for meteorological conditions, we observed a trend for temperature in the first trimester of
556 pregnancy being associated with the methylation distribution profiles (p-value for density function
557 = 0.03, p-value for cumulative distribution function = 0.11). We are aware of only one previous
558 study investigating the effects of meteorological conditions on DNA methylation in blood cells of
559 elderly men (Bind et al., 2014). The authors reported associations of ambient temperature with

560 DNA methylation in genes related to coagulation, inflammation, cortisol, and metabolic pathway
561 and associations of relative humidity with *LINE-1* hypomethylation and Alu hypermethylation. In
562 our study, we found two loci, one in *ZNF709* and one in a non genic region whose methylation was
563 associated with humidity exposure during the second trimester and the month preceding birth,
564 respectively. We also identified 13 DMRs mapping to 8 genes associated with meteorological
565 exposures. These genes do not seem involved in the inflammation or metabolic pathways
566 previously highlighted with blood methylation, and their association with prenatal meteorological
567 conditions had not been examined so far.

568 A novel feature of our study is the focus on genes with tissue specific patterns of expression, in
569 which we identified first a category of genes that are mostly active in the placenta and silent in
570 other tissues (“active genes”) and second a category of genes with a methylation-dependent
571 repressed state (“silent genes”) in the placenta. The definition of silent and active chromatin points
572 to regions with different dynamics and hence different capacity to respond to signals and to be
573 reprogrammed. In order to identify these regions, we used available transcriptomic data (for
574 selecting genes with specific patterns of expression in placenta) and the present methylome data
575 (list of silent genes with high methylation levels). This original approach not only allowed us to
576 identify potential target CpGs through the activity of their regulatory regions, but also to limit the
577 number of CpG sites to the most relevant ones to be tested in relation to environmental exposures,
578 without limiting the analysis to particular pathways previously identified in the literature. In the
579 context of a constantly expanding number of CpGs interrogated by microarrays and sequencing
580 techniques, which usually exceeds the sample size of the study, replication is often used to validate
581 the findings and limit the risk of false positives. Our strategy was rather to restrict our analysis to
582 the most biologically relevant genes and chromatin regions, which we defined as those prone to be
583 affected by alterations in methylation. Noteworthy, the results of the EWAS, which used a more
584 stringent correction of the p-value, and the DMR analyses strongly supports our finding from the

585 concept-driven analysis that *ADORA2B* placental methylation could be sensitive to prenatal air
586 pollution exposure.

587 We acknowledge several strengths and limitations to this study. First, our results demonstrate that
588 placental DNA methylation is associated with air pollutants exposure, but we cannot draw
589 conclusions on how gene expression relates to these methylation differences. Second, our study
590 focused on PM₁₀ and NO₂ air pollutants exposure, for which high temporally and spatially resolved
591 exposures were available. Air pollution is a complex mixture of compounds which are correlated
592 with the assessed pollutants. PM_{2.5} are of great concern because they can penetrate deep into the
593 lungs and enter the blood circulation. While our results identified mainly NO₂ exposure as
594 associated with placental DNA methylation, we rather interpret this association as an indication of
595 traffic-pollutants association with DNA methylation. Another limitation is the use of residential
596 address to estimate air pollution exposure, which is prone to misclassification due to the lack of
597 consideration of time activity patterns and of indoor exposures. However, we used a high spatially
598 and temporally resolved ambient dispersion model. Third, although we attempted to account for
599 many potential confounders including technical factors and cell heterogeneity, we cannot rule out
600 confounding from unaccounted factors such as physical activity, diet, or genetics. Moreover, since
601 our cohort consists of highly educated mainly Caucasian women, our findings may not be general-
602 izable to other populations. An important strength is that the sample size is one of the largest for a
603 study on placental DNA methylation and more generally for an environmental epigenetic study.
604 Samples were collected from the foetal side of the placenta, which is relatively easy to collect, non-
605 invasive and may allow a good reproducibility of results since the placenta lifetime is limited to the
606 pregnancy. In perinatal studies, the placenta represents a relevant tissue considered as an accurate
607 ‘record’ of children’s in-utero exposures (Maccani and Marsit, 2009) and as a health biomarker due
608 to its master regulator function of the fetal hormonal and endocrine milieu. EDEN is a well
609 characterized mother-child cohort, which allowed the consideration of a large range of potential
610 confounders, including data collected early in pregnancy.

611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635

6. Conclusions

Among the placenta methylated silent genes, we identified four CpGs, mapped to four genes, whose methylation levels in the foetal placenta were significantly associated with maternal exposures to NO₂ and PM₁₀. Among placenta active genes, we found five CpGs, mapped to five genes, whose methylation levels in the foetal placenta were significantly associated with maternal exposures to NO₂ and PM₁₀. One of these active genes, *ADORA2B*, had 2 CpGs whose methylation levels were also significantly associated with NO₂ exposure during the second trimester of pregnancy in the EWAS approach after FDR correction of the p-value. The DMRs analyses showed *ADORA2B* methylation as associated with NO₂ exposure in pregnancy and at each trimester of pregnancy. *ADORA2B* has been implicated in placental development, hypoxia and preeclampsia, a pathology which has been previously associated with air pollution exposure during pregnancy. This latter observation, together with our results, supports the hypothesis that placental DNA methylation is involved in the exposure-disease relationships during pregnancy. Whether the observed associations potentially bear long-term consequences for the health of the child and adult individual, or even possibly for future generations, remains to be investigated.

7. Acknowledgments

We thank the midwife research assistants (L. Douhaud. S. Bedel. B. Lortholary. S. Gabriel. M. Rogeon. and M. Malinbaum) for data collection and P. Lavoine for checking, coding, and entering data. The EDEN mother-child cohort study group includes: I Annesi-Maesano, JY Bernard, J Botton, M-A Charles, P Dargent-Molina, B de Lauzon-Guillain, P Ducimetière, M de Agostini, B Foliguet, A Forhan, X Fritel, A Germa, V Goua, R Hankard, B Heude, M Kaminski, B Larroque, N Lelong, J Lepeule, G Magnin, L Marchand, C Nabet, F Pierre, R Slama, MJ Saurel-Cubizolles, M

636 Schweitzer, O Thiebaugeorges. We thank Emie Seyve for her help in the preparation of the revised
637 manuscript and the reviewers for their insightful comments that improved the manuscript.

638 **8. References**

639 Acurio, J., Troncoso, F., Bertoglia, P., Salomon, C., Aguayo, C., Sobrevia, L., Escudero, C., 2014.
640 Potential role of A2B adenosine receptors on proliferation/migration of fetal endothelium
641 derived from preeclamptic pregnancies. *Biomed Res Int* 2014, 274507.
642 doi:10.1155/2014/274507

643 Beltran, A., Wu, J., Laurent, O., 2014. Associations of Meteorology with Adverse Pregnancy
644 Outcomes: A Systematic Review of Preeclampsia, Preterm Birth and Birth Weight. *IJERPH* 11,
645 91–172. doi:10.3390/ijerph110100091

646 Benjamini, Y., Hochberg, Y., 1995. Controlling the false discovery rate: a practical and powerful
647 approach to multiple testing. *Journal of the royal statistical society Series B* (... 57, 289–300.

648 Bind, M.-A., Zanobetti, A., Gasparrini, A., Peters, A., Coull, B., Baccarelli, A., Tarantini, L.,
649 Koutrakis, P., Vokonas, P., Schwartz, J., 2014. Effects of temperature and relative humidity on
650 DNA methylation. *Epidemiology* 25, 561–569. doi:10.1097/EDE.0000000000000120

651 Breton, C.V., Yao, J., Millstein, J., Gao, L., Siegmund, K.D., Mack, W., Whitfield-Maxwell, L.,
652 Lurmann, F., Hodis, H., Avol, E., Gilliland, F.D., 2016. Prenatal Air Pollution Exposures, DNA
653 Methyl Transferase Genotypes, and Associations with Newborn LINE1 and Alu Methylation
654 and Childhood Blood Pressure and Carotid Intima-Media Thickness in the Children's Health
655 Study. *Environ. Health Perspect.* 124, 1905–1912. doi:10.1289/EHP181

656 Cai, J., Zhao, Y., Liu, P., Xia, B., Zhu, Q., Wang, X., Song, Q., Kan, H., Zhang, Y., 2017. Exposure
657 to particulate air pollution during early pregnancy is associated with placental DNA
658 methylation. *Sci. Total Environ.* 607-608, 1103–1108. doi:10.1016/j.scitotenv.2017.07.029

659 Chen, D., Zhang, A., Fang, M., Fang, R., Ge, J., Jiang, Y., Zhang, H., Han, C., Ye, X., Huang, H.,
660 Liu, Y., Dong, M., 2014. Increased methylation at differentially methylated region of GNAS in
661 infants born to gestational diabetes. *BMC Med. Genet.* 15, 108. doi:10.1186/s12881-014-0108-
662 3

663 Chen, Y.-A., Lemire, M., Choufani, S., Butcher, D.T., Grafodatskaya, D., Zanke, B.W., Gallinger,
664 S., Hudson, T.J., Weksberg, R., 2013. Discovery of cross-reactive probes and polymorphic
665 CpGs in the Illumina Infinium HumanMethylation450 microarray. *Epigenetics* 8, 203–209.
666 doi:10.4161/epi.23470

667 Chiu, Y.-H.M., Hsu, H.-H.L., Coull, B.A., Bellinger, D.C., Kloog, I., Schwartz, J., Wright, R.O.,
668 Wright, R.J., 2016. Prenatal particulate air pollution and neurodevelopment in urban children:
669 Examining sensitive windows and sex-specific associations. *Environ Int* 87, 56–65.
670 doi:10.1016/j.envint.2015.11.010

671 Chu, T., Bunce, K., Shaw, P., Shridhar, V., Althouse, A., Hubel, C., Peters, D., 2014.
672 Comprehensive analysis of preeclampsia-associated DNA methylation in the placenta. *PLoS*
673 *ONE* 9, e107318. doi:10.1371/journal.pone.0107318

674 Clifford, A., Lang, L., Chen, R., Anstey, K.J., Seaton, A., 2016. Exposure to air pollution and
675 cognitive functioning across the life course - A systematic literature review. *Environ. Res.* 147,
676 383–398. doi:10.1016/j.envres.2016.01.018

677 Darashchonak, N., Sarisin, A., Kleppa, M.-J., Powers, R.W., Versen-Höynck, von, F., 2014.
678 Activation of adenosine A2B receptor impairs properties of trophoblast cells and involves
679 mitogen-activated protein (MAP) kinase signaling. *Placenta* 35, 763–771.
680 doi:10.1016/j.placenta.2014.06.369

681 Devlin, B., Roeder, K., Wasserman, L., 2001. Genomic control, a new approach to genetic-based
682 association studies. *Theor Popul Biol* 60, 155–166. doi:10.1006/tpbi.2001.1542

683 Du, P., Kibbe, W.A., Lin, S.M., 2008. lumi: a pipeline for processing Illumina microarray.
684 *Bioinformatics* 24, 1547–1548. doi:10.1093/bioinformatics/btn224

685 Dunk, C.E., Roggensack, A.M., Cox, B., Perkins, J.E., Åsenius, F., Keating, S., Weksberg, R.,
686 Kingdom, J.C.P., Adamson, S.L., 2012. A distinct microvascular endothelial gene expression
687 profile in severe IUGR placentas. *Placenta* 33, 285–293. doi:10.1016/j.placenta.2011.12.020
688 Fragkiadaki, P., Soultzis, N., Sifakis, S., Koutroulakis, D., Gourvas, V., Vrachnis, N., Spandidos,
689 D.A., 2015. Downregulation of notch signaling pathway in late preterm and term placentas
690 from pregnancies complicated by preeclampsia. *PLoS ONE* 10, e0126163.
691 doi:10.1371/journal.pone.0126163
692 Giorgis-Allemand, L., Pedersen, M., Bernard, C., Aguilera, I., Beelen, R.M.J., Chatzi, L., Cirach,
693 M., Danileviciute, A., Dedele, A., van Eijnden, M., Estarlich, M., Fernández-Somoano, A.,
694 Fernández, M.F., Forastiere, F., Gehring, U., Grazuleviciene, R., Gruzieva, O., Heude, B.,
695 Hoek, G., de Hoogh, K., van den Hooven, E.H., Håberg, S.E., Iñiguez, C., Jaddoe, V.W.V.,
696 Korek, M., Lertxundi, A., Lepeule, J., Nafstad, P., Nystad, W., Patelarou, E., Porta, D., Postma,
697 D., Raaschou-Nielsen, O., Rudnai, P., Siroux, V., Sunyer, J., Stephanou, E., Sørensen, M.,
698 Eriksen, K.T., Tuffnell, D., Varró, M.J., Vrijkotte, T.G.M., Wijga, A., Wright, J.,
699 Nieuwenhuijsen, M.J., Pershagen, G., Brunekreef, B., Kogevinas, M., Slama, R., 2017. The
700 Influence of Meteorological Factors and Atmospheric Pollutants on the Risk of Preterm Birth.
701 *Am J Epidemiol.* doi:10.1093/aje/kww141
702 Green, B.B., Karagas, M.R., Punshon, T., Jackson, B.P., Robbins, D.J., Houseman, E.A., Marsit,
703 C.J., 2016. Epigenome-Wide Assessment of DNA Methylation in the Placenta and Arsenic
704 Exposure in the New Hampshire Birth Cohort Study (USA). *Environ. Health Perspect.* 124,
705 1253–1260. doi:10.1289/ehp.1510437
706 Gruzieva, O., Xu, C.-J., Breton, C.V., Annesi-Maesano, I., Antó, J.M., Auffray, C., Ballereau, S.,
707 Bellander, T., Bousquet, J., Bustamante, M., Charles, M.-A., de Kluizenaar, Y., Dekker, den,
708 H.T., Duijts, L., Felix, J.F., Gehring, U., Guxens, M., Jaddoe, V.V.W., Jankipersadsing, S.A.,
709 Merid, S.K., Kere, J., Kumar, A., Lemonnier, N., Lepeule, J., Nystad, W., Page, C.M.,
710 Panasevich, S., Postma, D., Slama, R., Sunyer, J., Söderhäll, C., Yao, J., London, S.J.,
711 Pershagen, G., Koppelman, G.H., Melén, E., 2017. Epigenome-Wide Meta-Analysis of
712 Methylation in Children Related to Prenatal NO₂ Air Pollution Exposure. *Environ. Health*
713 *Perspect.* 125, 104–110. doi:10.1289/EHP36
714 Haider, S., Meinhardt, G., Velicky, P., Otti, G.R., Whitley, G., Fiala, C., Pollheimer, J., Knöfler,
715 M., 2014. Notch signaling plays a critical role in motility and differentiation of human first-
716 trimester cytotrophoblasts. *Endocrinology* 155, 263–274. doi:10.1210/en.2013-1455
717 Heude, B., Forhan, A., Slama, R., Douhaud, L., Bedel, S., Saurel-Cubizolles, M.-J., Hankard, R.,
718 Thiebaugeorges, O., De Agostini, M., Annesi-Maesano, I., Kaminski, M., Charles, M.-A.,
719 EDEN Mother–Child Cohort Study Group, 2015. Cohort Profile: The EDEN mother-child
720 cohort on the prenatal and early postnatal determinants of child health and development. *Int J*
721 *Epidemiol.* doi:10.1093/ije/dyv151
722 Hong, J.T., Son, D.J., Lee, C.K., Yoon, D.-Y., Lee, D.H., Park, M.H., 2017. Interleukin 32,
723 inflammation and cancer. *Pharmacol. Ther.* 174, 127–137.
724 doi:10.1016/j.pharmthera.2017.02.025
725 Houseman, E.A., Kile, M.L., Christiani, D.C., Ince, T.A., Kelsey, K.T., Marsit, C.J., 2016.
726 Reference-free deconvolution of DNA methylation data and mediation by cell composition
727 effects. *BMC Bioinformatics* 17, 259. doi:10.1186/s12859-016-1140-4
728 Huang, A., Wu, H., Iriyama, T., Zhang, Y., Sun, K., Song, A., Liu, H., Peng, Z., Tang, L., Lee, M.,
729 Huang, Y., Ni, X., Kellems, R.E., Xia, Y., 2017. Elevated Adenosine Induces Placental DNA
730 Hypomethylation Independent of A2B Receptor Signaling in Preeclampsia. *Hypertension* 70,
731 209–218. doi:10.1161/HYPERTENSIONAHA.117.09536
732 Iriyama, T., Sun, K., Parchim, N.F., Li, J., Zhao, C., Song, A., Hart, L.A., Blackwell, S.C., Sibai,
733 B.M., Chan, L.-N.L., Chan, T.-S., Hicks, M.J., Blackburn, M.R., Kellems, R.E., Xia, Y., 2015.
734 Elevated placental adenosine signaling contributes to the pathogenesis of preeclampsia.
735 *Circulation* 131, 730–741. doi:10.1161/CIRCULATIONAHA.114.013740
736 Janssen, B.G., Godderis, L., Pieters, N., Poels, K., Kiciński, M., Cuypers, A., Fierens, F., Penders,
737 J., Plusquin, M., Gyselaers, W., Nawrot, T.S., 2013. Placental DNA hypomethylation in

738 association with particulate air pollution in early life. *Part Fibre Toxicol* 10, 22.
739 doi:10.1186/1743-8977-10-22

740 Jedrychowski, W.A., Perera, F.P., Maugeri, U., Mroz, E., Klimaszewska-Rembiasz, M., Flak, E.,
741 Edwards, S., Spengler, J.D., 2010. Effect of prenatal exposure to fine particulate matter on
742 ventilatory lung function of preschool children of non-smoking mothers. *Paediatr Perinat*
743 *Epidemiol* 24, 492–501. doi:10.1111/j.1365-3016.2010.01136.x

744 Jia, R.-Z., Zhang, X., Hu, P., Liu, X.-M., Hua, X.-D., Wang, X., Ding, H.-J., 2012. Screening for
745 differential methylation status in human placenta in preeclampsia using a CpG island plus
746 promoter microarray. *Int. J. Mol. Med.* 30, 133–141. doi:10.3892/ijmm.2012.983

747 Kingsley, S.L., Eliot, M.N., Whitsel, E.A., Huang, Y.-T., Kelsey, K.T., Marsit, C.J., Wellenius,
748 G.A., 2016. Maternal residential proximity to major roadways, birth weight, and placental DNA
749 methylation. *Environ Int* 92-93, 43–49. doi:10.1016/j.envint.2016.03.020

750 Kloog, I., Melly, S.J., Coull, B.A., Nordio, F., Schwartz, J.D., 2015. Using Satellite-Based
751 Spatiotemporal Resolved Air Temperature Exposure to Study the Association between Ambient
752 Air Temperature and Birth Outcomes in Massachusetts. *Environ. Health Perspect.*
753 doi:10.1289/ehp.1308075

754 Maccani, M.A., Marsit, C.J., 2009. Epigenetics in the placenta. *Am. J. Reprod. Immunol.* 62, 78–
755 89. doi:10.1111/j.1600-0897.2009.00716.x

756 Madrigano, J., Baccarelli, A., Mittleman, M.A., Wright, R.O., Sparrow, D., Vokonas, P.S.,
757 Tarantini, L., Schwartz, J., 2011. Prolonged exposure to particulate pollution, genes associated
758 with glutathione pathways, and DNA methylation in a cohort of older men. *Environ. Health*
759 *Perspect.* 119, 977–982. doi:10.1289/ehp.1002773

760 Malassiné, A., Cronier, L., 2002. Hormones and human trophoblast differentiation: a review.
761 *Endocrine* 19, 3–11. doi:10.1385/ENDO:19:1:3

762 McMinn, J., Wei, M., Schupf, N., Cusmai, J., Johnson, E.B., Smith, A.C., Weksberg, R., Thaker,
763 H.M., Tycko, B., 2006. Unbalanced placental expression of imprinted genes in human
764 intrauterine growth restriction. *Placenta* 27, 540–549. doi:10.1016/j.placenta.2005.07.004

765 Murphy, V.E., Smith, R., Giles, W.B., Clifton, V.L., 2006. Endocrine regulation of human fetal
766 growth: the role of the mother, placenta, and fetus. *Endocr. Rev.* 27, 141–169.
767 doi:10.1210/er.2005-0011

768 Nelissen, E.C.M., van Montfoort, A.P.A., Dumoulin, J.C.M., Evers, J.L.H., 2011. Epigenetics and
769 the placenta. *Hum. Reprod. Update* 17, 397–417. doi:10.1093/humupd/dmq052

770 Novakovic, B., Evain-Brion, D., Murthi, P., Fournier, T., Saffery, R., 2017. VariableDAXXgene
771 methylation is a common feature of placental trophoblast differentiation, preeclampsia, and
772 response to hypoxia. *The FASEB Journal* 31, 2380–2392. doi:10.1096/fj.201601189RR

773 Nwanaji-Enwerem, J.C., Colicino, E., Trevisi, L., Kloog, I., Just, A.C., Shen, J., Brennan, K.,
774 Dereix, A., Hou, L., Vokonas, P., Schwartz, J., Baccarelli, A.A., 2016. Long-term ambient
775 particle exposures and blood DNA methylation age: findings from the VA normative aging
776 study. *Environ Epigenet* 2. doi:10.1093/eep/dvw006

777 Ollikainen, M., Smith, K.R., Joo, E.J.-H., Ng, H.K., Andronikos, R., Novakovic, B., Abdul Aziz,
778 N.K., Carlin, J.B., Morley, R., Saffery, R., Craig, J.M., 2010. DNA methylation analysis of
779 multiple tissues from newborn twins reveals both genetic and intrauterine components to
780 variation in the human neonatal epigenome. *Human Molecular Genetics* 19, 4176–4188.
781 doi:10.1093/hmg/ddq336

782 Ortiz, E.A.G., 2017. Air quality in Europe — 2016 report 1–88.

783 Panni, T., Mehta, A.J., Schwartz, J.D., Baccarelli, A.A., Just, A.C., Wolf, K., Wahl, S., Cyrus, J.,
784 Kunze, S., Strauch, K., Waldenberger, M., Peters, A., 2016. Genome-Wide Analysis of DNA
785 Methylation and Fine Particulate Matter Air Pollution in Three Study Populations: KORA F3,
786 KORA F4, and the Normative Aging Study. *Environ. Health Perspect.* 124, 983–990.
787 doi:10.1289/ehp.1509966

788 Park, S.K., 2017. Ambient Air Pollution and Type 2 Diabetes: Do the Metabolic Effects of Air
789 Pollution Start Early in Life? *Diabetes* 66, 1755–1757. doi:10.2337/dbi17-0012

790 Pedersen, B.S., Schwartz, D.A., Yang, I.V., Kechris, K.J., 2012. Comb-p: software for combining,

analyzing, grouping and correcting spatially correlated P-values. *Bioinformatics* 28, 2986–2988. doi:10.1093/bioinformatics/bts545

Pedersen, M., Stayner, L., Slama, R., Sørensen, M., Figueras, F., Nieuwenhuijsen, M.J., Raaschou-Nielsen, O., Dadvand, P., 2014. Ambient air pollution and pregnancy-induced hypertensive disorders: a systematic review and meta-analysis. *Hypertension* 64, 494–500. doi:10.1161/HYPERTENSIONAHA.114.03545

Péterfy, M., 2012. Lipase maturation factor 1: a lipase chaperone involved in lipid metabolism. *Biochim. Biophys. Acta* 1821, 790–794. doi:10.1016/j.bbalip.2011.10.006

Pijnenborg, R., Vercruysse, L., Hanssens, M., 2006. The uterine spiral arteries in human pregnancy: facts and controversies. *Placenta* 27, 939–958. doi:10.1016/j.placenta.2005.12.006

Price, M.E., Cotton, A.M., Lam, L.L., Farré, P., Emberly, E., Brown, C.J., Robinson, W.P., Kobor, M.S., 2013. Additional annotation enhances potential for biologically-relevant analysis of the Illumina Infinium HumanMethylation450 BeadChip array. *Epigenetics Chromatin* 6, 4. doi:10.1186/1756-8935-6-4

Roberts, V.H.J., Webster, R.P., Brockman, D.E., Pitzer, B.A., Myatt, L., 2007. Post-Translational Modifications of the P2X(4) purinergic receptor subtype in the human placenta are altered in preeclampsia. *Placenta* 28, 270–277. doi:10.1016/j.placenta.2006.04.008

Rousseaux, S., Debernardi, A., Jacquiau, B., Vitte, A.-L., Vesin, A., Nagy-Mignotte, H., Moro-Sibilot, D., Brichon, P.-Y., Lantuejoul, S., Hainaut, P., Laffaire, J., de Reyniès, A., Beer, D.G., Timsit, J.-F., Brambilla, C., Brambilla, E., Khochbin, S., 2013. Ectopic activation of germline and placental genes identifies aggressive metastasis-prone lung cancers. *Sci Transl Med* 5, 186ra66. doi:10.1126/scitranslmed.3005723

Sellier, Y., Galineau, J., Hulin, A., Caïni, F., Marquis, N., Navel, V., Bottagisi, S., Giorgis-Allemand, L., Jacquier, C., Slama, R., Lepeule, J., EDEN Mother–Child Cohort Study Group, 2014. Health effects of ambient air pollution: do different methods for estimating exposure lead to different results? *Environ Int* 66, 165–173. doi:10.1016/j.envint.2014.02.001

Shah, A.S.V., Langrish, J.P., Nair, H., McAllister, D.A., Hunter, A.L., Donaldson, K., Newby, D.E., Mills, N.L., 2013. Global association of air pollution and heart failure: a systematic review and meta-analysis. *Lancet* 382, 1039–1048. doi:10.1016/S0140-6736(13)60898-3

Shen, L., Diao, Z., Sun, H.-X., Yan, G.-J., Wang, Z., Li, R.-T., Dai, Y., Wang, J., Li, J., Ding, H., Zhao, G., Zheng, M., Xue, P., Liu, M., Zhou, Y., Hu, Y., 2017. Up-regulation of CD81 inhibits cytotrophoblast invasion and mediates maternal endothelial cell dysfunction in preeclampsia. *Proceedings of the National Academy of Sciences* 114, 1940–1945. doi:10.1073/pnas.1617601114

Stieb, D.M., Chen, L., Eshoul, M., Judek, S., 2012. Ambient air pollution, birth weight and preterm birth: A systematic review and meta-analysis. *Environ. Res.* 117, 100–111. doi:10.1016/j.envres.2012.05.007

Šidák, Z., 1967. Rectangular Confidence Regions for the Means of Multivariate Normal Distributions. *Journal of the American Statistical Association* 62, 626–633. doi:10.1214/aoms/1177728724

Tobi, E.W., Lumey, L.H., Talens, R.P., Kremer, D., Putter, H., Stein, A.D., Slagboom, P.E., Heijmans, B.T., 2009. DNA methylation differences after exposure to prenatal famine are common and timing- and sex-specific. *Human Molecular Genetics* 18, 4046–4053. doi:10.1093/hmg/ddp353

Touleimat, N., Tost, J., 2012. Complete pipeline for Infinium ®Human Methylation 450K BeadChip data processing using subset quantile normalization for accurate DNA methylation estimation. *Epigenomics* 4, 325–341. doi:10.2217/epi.12.21

Valentino, S.A., Tarrade, A., Aioun, J., Mourier, E., Richard, C., Dahirel, M., Rousseau-Ralliard, D., Fournier, N., Aubrière, M.-C., Lallemand, M.-S., Camous, S., Guinot, M., Charlier, M., Aujean, E., Adhami, A., H., Fokkens, P.H., Agier, L., Boere, J.A., Cassee, F.R., Slama, R., Chavatte-Palmer, P., 2015. Maternal exposure to diluted diesel engine exhaust alters placental function and induces intergenerational effects in rabbits. *Part Fibre Toxicol* 13, 39–39. doi:10.1186/s12989-016-0151-7

844 van Iterson, M., van Zwet, E.W., BIOS Consortium, Heijmans, B.T., 2017. Controlling bias and
845 inflation in epigenome- and transcriptome-wide association studies using the empirical null
846 distribution. *Genome Biol.* 18, 19. doi:10.1186/s13059-016-1131-9

847 Wang, J., Mi, J.-Q., Debernardi, A., Vitte, A.-L., Emadali, A., Meyer, J.A., Charmpi, K., Ycart, B.,
848 Callanan, M.B., Carroll, W.L., Khochbin, S., Rousseaux, S., 2015. A six gene expression
849 signature defines aggressive subtypes and predicts outcome in childhood and adult acute
850 lymphoblastic leukemia. *Oncotarget* 6, 16527–16542. doi:10.18632/oncotarget.4113

851 Wick, P., Malek, A., Manser, P., Meili, D., Maeder-Althaus, X., Diener, L., Diener, P.-A., Zisch,
852 A., Krug, H.F., Mandach, von, U., 2010. Barrier capacity of human placenta for nanosized
853 materials. *Environ. Health Perspect.* 118, 432–436. doi:10.1289/ehp.0901200

854 World Health Organization, 2006. Air quality guidelines: global update 2005.

855 Yang, A.S., Estécio, M.R.H., Doshi, K., Kondo, Y., Tajara, E.H., Issa, J.-P.J., 2004. A simple
856 method for estimating global DNA methylation using bisulfite PCR of repetitive DNA
857 elements. *Nucleic Acids Res.* 32, e38. doi:10.1093/nar/gnh032

858 Yang, B.-Y., Qian, Z.M., Li, S., Fan, S., Chen, G., Syberg, K.M., Xian, H., Wang, S.-Q., Ma, H.,
859 Chen, D.-H., Yang, M., Liu, K.-K., Zeng, X.-W., Hu, L.-W., Guo, Y., Dong, G.-H., 2018.
860 Long-term exposure to ambient air pollution (including PM1) and metabolic syndrome: The 33
861 Communities Chinese Health Study (33CCHS). *Environ. Res.* 164, 204–211.
862 doi:10.1016/j.envres.2018.02.029

863 Zhao, W.-X., Huang, T.-T., Jiang, M., Feng, R., Lin, J.-H., 2014. Expression of notch family
864 proteins in placentas from patients with early-onset severe preeclampsia. *Reprod Sci* 21, 716–
865 723. doi:10.1177/1933719113512530

866

867