Lipids in benthic diatoms: A new suitable screening procedure

Eva Cointet, Gaëtane Wielgosz-Collin, Vona Méléder, Olivier Gonçalves

► To cite this version:

Eva Cointet, Gaëtane Wielgosz-Collin, Vona Méléder, Olivier Gonçalves. Lipids in benthic diatoms: A new suitable screening procedure. *Algal Research - Biomass, Biofuels and Bioproducts*, 2019, 39, pp.101425. 10.1016/j.algal.2019.101425 . hal-02333689

HAL Id: hal-02333689

<https://hal.science/hal-02333689>

Submitted on 22 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution - NonCommercial| 4.0 International License

Lipids in benthic diatoms: a new suitable screening procedure

Short title: Screening and rapid selection of lipid rich benthic diatoms

Eva Cointet¹, Gaëtane Wielgosz-Collin¹, Vona Méléder¹, Olivier Gonçalves²✉

¹ Université de Nantes, Laboratoire Mer Molécules Santé, EA 21 60, BP 92208, 44322 Nantes, France

² Université de Nantes, GEPEA, UMR CNRS-6144, Bât.CRTT, 37 Boulevard de l'Université, BP406, F-44602 Saint-Nazaire Cedex, France

Abstract

The selection of suitable and indigenous microalgae species is a fundamental requirement in developing added-value bioactive compounds recoverable in the food, health, and cosmetics markets. In this work, an integrated screening approach was developed to characterize the lipid rate of 33 diatom species (including 15 species studied for the first time) belonging to 16 genera from the Nantes Culture Collection, with the main objective of discovering bioactive lipid producers. For that purpose, a simple reliable method for establishing growth kinetics of strains and semi-quantitative analysis of lipid rates was developed. Growth kinetics measurements were achieved by daily minimal measurement fluorescence (F0) whereas lipid rate analyses were performed by high-throughput Fourier Transform Infrared spectroscopy on entire cells and lipid extracts. Results indicated that the method could be used directly on entire cells in spite of the presence of silica for the FTIR approach (due to frustule). The total lipid rate was species-dependant and ranged from 3.7% to 30.5% DW. Six strains out of 33 were found to present a higher total lipid rate superior to 15% DW, and 11 showed medium lipid rates ranging from 10% to 15% DW. The results revealed that five diatom species i.e. *Amphora* sp. NCC169, *Nitzschia* sp. NCC109, *Nitzschia alexandrina* NCC33, *Opephora* sp NCC366 and *Staurosira* sp. NCC182 presented interesting growth capabilities and should be further investigated as potential sources for their original lipid rate.

Keywords: FTIR spectroscopy, growth kinetics, benthic Diatoms, lipid rate, bioactive fatty acids.

Highlights

- Minimal fluorescence measurement characterized growth capabilities
- FTIR spectroscopy on whole cells characterized lipid rates
- Lipid rates of 15 strains were highlighted for the first time
- Five strains of benthic diatoms were selected for further biotechnology applications

*Corresponding authors

Email: olivier.goncalves@univ-nantes.fr (Olivier Gonçalves), eva.cointet@univ-nantes.fr (Eva Cointet), vona.meleder@univ-nantes.fr (Vona Méléder), Gaetane.Wielgosz-Collin@univ-nantes.fr (Gaëtane Wielgosz-Collin).

Preprint submitted to Algal Research September 09, 2018

Conflict of interest

The authors declare that there is no conflict of interest. No conflicts, informed consent, human or animal rights are applicable to this work.

Author's contribution

Eva Cointet, Gaëtane Wielgosz-Collin, Vona Méléder and Olivier Gonçalves designed and supervised the research. Eva Cointet, Gaëtane Wielgosz-Collin, Vona Méléder and Olivier Gonçalves conducted experiments. Eva Cointet, Gaëtane Wielgosz-Collin, Vona Méléder and

51 Olivier Gonçalves analyzed and interpreted the data and drafted the manuscript. Eva Cointet,
52 Gaëtane Wielgosz-Collin, Vona Méléder and Olivier Gonçalves critically reviewed the
53 manuscript. All authors read and approved the final version of the manuscript.
54

55 Acknowledgements

56 This work was supported by the regional Atlantic Microalgae research program (AMI), funded
57 by the Pays de la Loire region. We also express our sincere thanks to GEPEA staff in particular
58 Remy Coat and Delphine Kucma for support and advice on the FTIR spectrometer.

59 Abbreviation

60 ATR: Attenuated Total Reflectance

61 AW: Ash weight

62 DW: Dry weight

63 DHA: Docosahexaenoic acid

64 EPA: Eicosapentaenoic acid

65 FA: Fatty acid

66 F0: Minimum chlorophyll fluorescence

67 HTSXT-FTIR: Fourier-transform infrared spectroscopy high-throughput screening extension

68 LED: Late exponential day

69 LED biomass: Biomass corresponding to culture harvest days

70 MUFA: Monounsaturated fatty acid

71 NCC: Nantes Culture Collection

72 NDVI: Normalized Difference Vegetation Index

73 N: Nitrogen

74 PFD: Photon flux density

75 PAM: Pulse amplitude modulation

76 PAR: Photosynthetically active radiation

77 PUFA: Polyunsaturated fatty acid

78 ρ : Reflectance

79 SD : Standard deviation

80 Si : Silica

81 SFA: Saturated fatty acid

82 TAG: Triacylglycerol

83 $\mu_{\max}$: Maximum growth rate

84

1 Introduction

In the last two decades, a large body of research has focused on finding new strains of microalgae capable of producing high lipid content for a wide range of applications including pharmaceutical, cosmetics and alternative biofuels [1–4]. In the kingdom of microalgae, diatoms are very accessible resources, since they are ubiquitously found in most aquatic environments (rivers, oceans, coastal areas). They constitute a unicellular eukaryotic group with a typical species-specific siliceous cell wall (also known as frustule). They present different life-forms that could be benthic (microphytobenthos) or planktonic (phytoplankton). Marine diatoms can grow quickly and store large amounts of lipids [5]. Their lipids are mainly composed of a neutral fraction with traces of sterols and polar lipids [6]. Neutral lipids constitute the reserve fraction, with triacylglycerol (TAG) accounting for more than 60% of the total lipids [7]. Their PUFAs are mainly composed of eicosapentaenoic acid (EPA, C20:5 n-3) [8] but some strains were also found to present docosahexaenoic acid (DHA, C22:6 n – 3) [9]. The biosynthesis of the lipids varies within the different diatom species, their growth stages, and environmental parameters [10,11]. Previous studies [5–9] have demonstrated their ability for lipid production, more specifically for the PUFA fraction (DHA and EPA), recognized for its broad spectrum bioactivities (anti-carcinogenic, immune modulation, anti-diabetic, anti-obesity and anti-thrombotic properties) [12]. Unfortunately, the Diatoms group is poorly studied and constitutes therefore an underexploited resource [13] even if the number of their genera and species is estimated to be between 250 and 100,000 [14]. Bioprospecting efforts should therefore be encouraged in order to assess this potential. Basically, bioprospecting would be achieved if the identified microalgae could be exploited at an industrial scale for their biomass or their high value lipid compounds [1]. Therefore, during the screening approach, specific focus should be performed on the efficient identification of the appropriate microalgae strains, i.e. those characterized by high productivity (biomass and lipids), high resistance to

110 contamination and high tolerance to a wide range of environmental parameters [15–18]. Native
111 species adapt to local environmental changes and should be thus resilient and competitive
112 enough regarding these criteria. However, systematic estimation of the growth rates requires a
113 time-consuming series of measurements in order to estimate biomass evolution. The cell count
114 approach is among the most widely used. Alternatively, other parameters could be measured as
115 proxy for the cell numbers, if it could be shown to be linearly correlated. Typical proxy
116 measurements are *in vivo* fluorescence [19], optical density and biomass direct estimation (such
117 as dry weight, pigment content) [20] and *in vivo* or in solvent spectroscopy [21]. The
118 concentrations of protein, carbohydrates and lipids in cultures could also be used as proxy
119 measures depending of the robustness of their linear relationship with either cell numbers or
120 biomass. Finally, the cell number is recognized as being a robust reference method if “counting
121 methods” are easily available and could be applied to a given biological system. Only *in vivo*
122 fluorescence and spectroscopy seem to be faster than cell counting and are easy to use. For
123 these reasons, one of the objectives of this study is to test both techniques as an alternative to
124 counting in the screening process. Conventional methods used for lipid determination,
125 systematically require solvent extraction and gravimetric determination. These methods are
126 time consuming, need extensive manipulations, use high amounts of biomass (10-15 mg [22])
127 and have a low throughput screening rates. Consequently, a faster measurement of the lipid
128 content is needed [23]. As for counting methods, alternative approaches exist, that can be
129 roughly classified into invasive and non-invasive techniques. Fluorescent based technologies
130 are the most commonly used and require fluorescent dyes like Bodipy. However, it is an indirect
131 measure that has several issues such as sample preparation before staining, careful choice of
132 dye especially in microalgae due to the presence of chlorophylls within chloroplasts, leading to
133 non-quantitative information [24]. Vibrational spectroscopy such as Raman [25] could be a
134 good alternative for investigating lipid content since it can image and chemically identify the

lipids without labelling. However, autofluorescence signals from the chloroplast will overwhelm the Raman signal. Increasing the acquisition time or incident power can solve this issue, although irreversible photo damage may result, leading to loss of the semi-quantitative information, limiting therefore its use in the context of high-throughput screening. Infrared spectroscopy has advantages over the above techniques, since it limits photodamage, is not influenced by autofluorescence, and presents robust systems such as high-throughput screening for chemical spectra acquisition for large quantities of samples. Coat *et al.* demonstrated [26] that it was robust and sensitive enough to quantify the lipids and that directly on entire microalgae cells. The second objective of this study was to test this technique as an alternative to gravimetric determination of lipid content.

In the present work, we propose to explore the potential for original lipid sourcing of the benthic diatoms hosted in the Nantes Culture Collection (NCC) bank. First – to identify the NCC strains that could potentially produce interesting fatty acids – a bibliographic inventory of current and past knowledge of the principal genus and species hosted in the NCC bank was conducted. It focused on the families of molecules with high added value for pharmacology, health, nutrition, and cosmetics such as PUFAs, taking also into account the influence of cultural conditions [27,28]. The selected species were then, investigated to gain basic knowledge on their biological characteristics, to highlight their chemo diversity (in terms of protein, carbohydrates, silica and lipid rates), and finally on their potential for high-value fatty acid molecule biosynthesis. To select the most promising NCC strains, an original workflow was developed, integrating different steps including the analysis of the NCC strain growth and their rapid biochemical profiling through HTSXT-FTIR. The results obtained in the present study are discussed hereafter.

158 2 Material and methods

159 2.1 Strains cultivation

160 Diatom strains from the NCC were selected and cultivated in 250 mL Erlenmeyer flasks filled
161 with 150 mL of F/2 culture medium, enriched in silica using 0.47 μm filtered natural sea water
162 [29, 30]. Vitamins and carbonates were added before autoclave sterilization (Autoclave Vertical
163 Lequeux AUV 100L), salinity was adjusted at 28 and pH was fixed at 7.8 to reduce nutrient
164 precipitation. Inoculation of each strain was performed from the stock cultures with an initial
165 concentration of 30 000 cells.mL⁻¹. The diatom strains were thus grown at 16°C with a photon
166 flux density (PFD) of 127 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ provided under continuous light to the bottom
167 of flasks by flat led pannel (LP EPURE-Chateaugiron, France).

168 2.2 Growth parameter estimations

169 To retrieve the growth parameters on the screened strains, the growth rate (μ_{max} in day⁻¹) and
170 the end of the exponential phase were identified. The growth rate identifies the faster growing
171 strains. The end of the exponential phase, known to be the period of lipid accumulation [31],
172 identifies the harvest day for lipid analyses. These parameters were retrieved using daily
173 biomass estimation. All erlen flasks and stir bars were autoclaved. Before sampling, flasks were
174 agitated by magnetic stirring for 2 minutes allowing cell and nutrient homogenization and
175 aggregate destruction.

176 Alternative techniques were tested to replace the cell counting by hemocytometer (here
177 Neubauer; $n \geq 300$) which is a time-consuming technique.,

178 - Minimum chlorophyll fluorescence measurement (F0) by fluorometry PAM (Water-PAM,
179 Waltz, Germany). This parameter, proportional to Chl *a* content, was used as a proxy of the
180 vegetal biomass [32]. Measurements were made directly on the microalgal suspension.

- Reflectance (ρ) of the cells by spectroradiometry (Jaz, Ocean Optics, USA) in the PAR domain (400-700 nm). Reflectance values were used to calculate the Normalized Difference Vegetation Index (NDVI) following the equation. (1), known to be proportional to Chl *a* content and used as a biomass proxy [21]. Measurements were made on filtered microalgae using CML microfiber filters with a 25 mm diameter and a 0.7 μ m pore diameter and a 25mm Whatman filter funnel.

$$NDVI = \frac{\lambda_{750} - \lambda_{675}}{\lambda_{750} + \lambda_{675}} \quad \text{eq. (1)}$$

With λ_{750} : maximal reflectance wavelength, and λ_{675} : Chl *a* absorption wavelength

Daily measurements were performed on each sample of culture to follow the growth kinetics (at the beginning of the growth phase and during three days, between 3 mL and 5 mL of cultures were sampled, and then 2 mL until the end of the growth). To extract and compare the growth parameters from the alternative techniques and the cell count approach, the Gompertz model [33] was used to fit the growth data (equation. 2) using MATLAB software. It consisted of a latency phase followed by an exponential phase and a stabilization of the curve at its maximum phase.

$$f(x) = A \times e^{-e(\mu_{max} \times \frac{e^1}{A} \times (\lambda - x) + 1)} \quad \text{eq. (2)}$$

With *A*: maximum cell concentration in the natural logarithm of the biomass (number of cells.mL⁻¹, F0 or NDVI); μ_{max} : Maximum growth rate (day⁻¹); λ : Latency (days).

The *A* parameter allows calculation of the biomass ($LED_{biomass}$) obtained at the late exponential day (LED) (equation. 3). This corresponded to the harvest day for the lipid analyses.

$$LED_{biomass} = \exp(A + \log(Bmin)) \quad \text{eq. (3)}$$

With *Bmin* = Initial biomass (number of cells.mL⁻¹, F0 or NDVI); $LED_{biomass}$ = biomass of the late exponential day (cell.mL⁻¹, F0 or NDVI)

The comparison of the three techniques, and the selection of the fastest one to estimate the growth parameters was done using a panel of 6 very different strains, chosen for their different growth kinetics and their different morphology (aggregate of cells, cells in chains or solitary cells) : *Amphora* sp. 1 NCC260, *Entomoneis* sp. 1 NCC350, *Entomoneis* sp. 6 NCC335, *Entomoneis paludosa* NCC18.2, *Extubocellulus cribriger* NCC229 and *Navicula* sp. 2 NCC226.

2.3 Diatom strain characterisation

At the end of the exponential phase (LED), when the cells were harvested for HTSXT-FTIR analyses (see §2.4), several analysis were performed to collect more information about the strains:

- the cell length and the width were estimated using the image of light microscopy (OLYMPUS CH40, $\times 400$; $n=150$).

- the dry weight (DW) of the biomass was estimated by filtering 150 mL of algal suspension through a microfiber filter, Whatman 47 mm diameter, 0.7 μm pore. The filters containing the cells were washed using 10 mL of ammonium formate (3%) to remove salt. The wet filters were frozen at -80°C and freeze-dried under a vacuum. DW (g.L^{-1}) and μ_{max} (day^{-1}) were then used to estimate the strain productivity (P_x) in $\text{g.L}^{-1}.\text{day}^{-1}$ (equation. 4).

$$P_x = \mu_{\text{max}} \times DW \quad \text{eq. (4)}$$

- the strains total lipid rate was assessed by gravimetric assay, to compare it to the infrared semi-quantitative measurements (HTSXT-FTIR, see § 2.4). Biomass filtered, washed and freeze-dried for dry weight estimation were used for lipid content estimation. The filters were macerated in flasks using 100 mL of solvent per gram of biomass (dichloromethane-methanol (1:1 V/V)) [34]. Maceration at ambient temperature was performed for 24H on a vibrating tray (Edmund Bühler GmbH, SM-30). After maceration, the mixture was filtered on pleated filters, 190 mm diameter, 10 μm pore, to remove the filter debris and the silica fragments. The filtrates were

transferred into a separatory funnel with 20 mL of distilled water and shaken for 5 min. The lipid fraction (organic phase) was then separated from the separatory funnel, dried using an anhydrous sodium sulfate salt, filtered, evaporated and weighed to obtain the crude lipid extract (CLE) value. Total lipid rate (TLR) was finally expressed in % of the DW (equation. 5).

$$TLR = \frac{CLE}{DW} \times 100 \quad \text{eq. (5)}$$

- the silica content of the cells was also determined to ensure the normalization of the FTIR semi-quantitative results (see § 2.4). Cultures were harvested by filtering new 150 mL of the algal suspension using the same recovery cell procedure used for dry weight estimation. Filters were then freeze-dried, weighed (DW) and heated at 400°C for four hours in a muffle oven and weighed (AW) and the silica proportion evaluated in % of DW by equation. 6 :

$$Silica\ content = \frac{AW}{DW} \times 100 \quad \text{eq. (6)}$$

2.4 Molecular profiles measured by infrared spectroscopy

The FTIR spectra acquisition on entire cells was performed according to Coat et al. 2014 recommendations [26]. It consisted in concentrating the cells up to 10^6 - 10^8 cells.mL⁻¹ by centrifugation (10 000g for 5 min) using Sigma 3K30 centrifuge. The supernatant was removed, and the pellet resuspended in an ammonium formiate isotonic solution (68 g.L⁻¹). This process was repeated twice to wash out the cells from the growth medium in order to avoid medium contribution on the FTIR spectra. The cells were thus resuspended in 1mL in an isotonic solution. Ammonium formiate solution prevents cell lyses during washing [35]. For the FTIR spectra acquisition, a Bruker tensor 27 FTIR spectrometer equipped with a HTSXT plate reader module coupled to OpusLab v 7.0.122 software (Bruker Optics, Germany) was used. Rinsed cell aliquots of 5 µL were deposited on a 384 well microplate, void-dried in a vacuum desiccator for at least 24 hours. FTIR spectra were then recorded in transmission mode directly on the microplate loaded with the dried samples. This method was chosen since it is fast and non-

invasive on intact diatom cells, their biochemical signatures expressed in term of total lipids, total proteins and total carbohydrates superimposed partially with the silica signal of the diatom frustul. To confirm the whole cells FTIR spectra results, estimation of the lipid rate was also performed on crude lipid extract, using attenuated total reflectance sampling system (ATR), that was more suitable for crude organic extracts. For the acquisition of the ATR spectra, 10 μ L of crude lipid extract (see §2.3. [34]) were deposited directly on the Bruker tensor 27 FTIR spectrometer lens. The absorbance spectra were all collected between 4 000 cm^{-1} and 700 cm^{-1} with 30 scans and averaged. The spectra were analyzed by relatively straightforward methods such as peak ratios or integral ratios [36]. The lipid signature was associated to the $\text{CH}_2\text{-CH}_3$ signal ($\sim 3000 - 2800 \text{ cm}^{-1}$) and the ester bond (Eb) signal ($\sim 1740 \text{ cm}^{-1}$). The carbohydrate signature was associated to the C-O-C signal of the polysaccharides ($\sim 1200\text{-}900 \text{ cm}^{-1}$) [37]. The protein signature was associated to the amide II band ($\sim 1540 \text{ cm}^{-1}$) of the N-H of the amides associated to the proteins. The silica signature was associated to the Si-O signal of the silicate frustule ($\sim 1068 \text{ cm}^{-1}$) [38]. To estimate the relative content of the lipids, carbohydrates and proteins, their respective peak heights (i.e. ester bond + ($\text{CH}_2 + \text{CH}_3$), $\sim 1159 \text{ cm}^{-1}$ and amide II) were standardized to the silica peak [39]. (equation 7). For the crude lipid extract spectra, the ratio used for that purpose was the ester bond and the $\text{CH}_2 + \text{CH}_3$ signals standardized with the total spectrum area (equation 8). FTIR and ATR ratio were expressed in arbitrary units abundance (a.u).

$$\text{FTIR} = \frac{\text{Peak height (S)}}{\text{Silica peak high}} \quad \text{eq. (7)}$$

S = lipids (eb+(CH_2+CH_3)) or carbohydrates ($\sim 1159 \text{ cm}^{-1}$) or amide II ($\sim 1540 \text{ cm}^{-1}$)

$$\text{ATR} = \frac{\text{Peak area (eb+CH}_2\text{+CH}_3\text{)}}{\text{Total spectra area}} \quad \text{eq. (8)}$$

2.5 Data processing

The Pearson product-moment correlation was carried out to test the positive correlation between the growth curves obtained with the cell count (Cells.mL⁻¹), fluorescence (F0) and reflectance (NDVI).

Comparison of the growth rate and LED estimated from the three techniques for the selected species were performed using ANOVA when the data presented normal distribution or the Kruskal – Wallis test when the data distribution was not normal. It was systematically followed by the Tuckey post hoc test. The ANOVA was also performed on the growth rate (μ_{max}), production (Px) and lipid rate (TLR) results to identify the strains with the highest performance.

A multivariate correspondence analysis was performed on the FTIR ratio normalized by silica for lipids, protein and carbohydrates to assess the dispersion of the biochemical information across the screened species and to identify if strains could be classified according to this information. This method was chosen since it analyzes binary, ordinal and nominal data without distributional assumptions (unlike traditional multivariate techniques) and also to preserve the categorical nature of the variables. The correspondence analysis provided a unique graphical display showing how the variable response categories were related [40].

The Pearson product-moment correlation was used to test the correlation between the calculation methods using the FTIR spectra semi-quantitative information with the total lipid quantification reference method (the gravimetric approach). The Pearson product-moment correlation and the comparison of the growth parameters were carried out using SigmaStat 3.1 software. The Past3 software was used for the correspondence analysis approach. All experiments were performed in triplicate.

Finally, all the information and tests were combined and used for the validation of the screening methodology to characterize the diatom strains of the NCC and the selection of candidates presenting the best potential in terms of growth capabilities and lipid rates.

3 Results

3.1 Determination of the growth parameters

The latency, exponential and stationnary phases were observable whatever the technique used to establish the growth curve for the 6 strains selected for their different kinetic behaviour (Fig. 1). Moreover, the cell count, NDVI and F0 data were all correlated (Tab.1) confirming the same growth pattern. Cell count and NDVI were positively correlated ($p<0.001$) as cell count and F0 ($p<0.001$). R values for the NDVI varied from 0.70 to 0.90 and for F0 measurements from 0.60 to 0.93.

Figure1. Example of growth curves measured for *Amphora* sp 1 NCC260 by (A) cell count, (B) fluorometry PAM (F0) and (C) radiometry (NDVI). Points corresponding to cell concentration according to time. Line curves corresponding to the Gompertz model fitted to cell concentration as a function of time. $n=3$, vertical bar = SD. The arrows indicate the late exponential day estimated by the Gompertz model.

Table 1. Pearson product correlation between radiometry (NDVI) and cell count and fluorometry PAM (F0) and cell count. Correlation was significant ($p < 0.001$)

Species	NDVI	F0
<i>Amphora</i> sp. 1 NCC260	0.90	0.93
<i>Entomoneis paludosa</i> NCC18.2	0.86	0.82
<i>Entomoneis</i> sp. 1 NCC335	0.86	0.81
<i>Entomoneis</i> sp. 6 NCC350	0.80	0.60
<i>Extubocellulus cf cribriger</i> NCC229	0.70	0.90
<i>Navicula</i> sp. 2 NCC226	0.80	0.78

There were no significant differences ($p > 0.05$) between the technique used to estimate the growth rate ($\mu_{\max}$) and the late exponential day (LED) for the six tested species, except for the $\mu_{\max}$ value estimated for *Entomoneis paludosa* (Tab. 2). For example, *Amphora* sp. 1 reached the late exponential phase at the day c.a. 13 (LED, Fig. 1, Tab. 2) with a mean $\mu_{\max}$ of c.a. 0.81 (Tab. 2). At this time, the maximum biomass was reached by the cell count indicator with $4.5 \pm 0.2 \times 10^6$ cells.mL⁻¹ by PAM fluorometry with F0 values 2785 ± 609 , and by radiometry with NDVI reaching 0.47 ± 0.05 . Because the fluorimetry is used extensively for the measurement of extracted Chl *a*, for the estimation of the phytoplankton Chl *a in vivo* [19,41] and do not need filtration of high amounts of culture (contrary to NDVI), it was selected as a fast and reliable alternative approach to the cell count to determine the growth rates of all selected diatom strains in this study. Regarding *Entomoneis paludosa*, a new experiment using only PAM measurements, in the same culture conditions but without counting and NDVI estimation were conducted. $\mu_{\max}$ values obtained were 0.52, 0.55, 0.55, leading to an average value of 0.54 ± 0.016 . A second statistical test performed on these new data concluded no significant difference for $\mu_{\max}$ and LED ($p = 0.47$).

Table 2. Values of the growth parameters (maximum growth rate, $\mu_{\max}$ (in d⁻¹) and late exponential day, LED (in days)) retrieved from the Gompertz model using six species employed for the comparison of the cell count approach with two alternative techniques: Fluorometry

PAM and radiometry. The calculated P value corresponded to the ANOVA test except for the (*) values that were obtained using the Kruskal-Wallis test. N=3, independent measurements, $\pm$ SD.

Species	Techniques	μ max (day ⁻¹)	LED (day)	P value (μ max)	P Value (LED)
<i>Amphora</i> sp. 1 NCC260	Cell count	0.76 $\pm$ 0.10	13 $\pm$ 1	P=0.8	P=0.16*
	Pam	0.78 $\pm$ 0.30	14 $\pm$ 1		
	Spectroradiometer	0.88 $\pm$ 0.25	12 $\pm$ 2		
<i>Entomoneis</i> <i>paludosa</i> NCC18.2	Cell count	0.50 $\pm$ 0.03	12 $\pm$ 1	P $\leq$ 0.05 ¹	P=0.07*
	Pam	0.95 $\pm$ 0.23 ¹	10 $\pm$ 0		
	Spectroradiometer	0.47 $\pm$ 0.01	10 $\pm$ 0		
<i>Entomoneis</i> sp. 6 NCC335	Cell count	0.53 $\pm$ 0.10	14 $\pm$ 1	P=0.27	P=0.14*
	Pam	0.54 $\pm$ 0.12	12 $\pm$ 1		
	Spectroradiometer	0.32 $\pm$ 0.14	12 $\pm$ 2		
<i>Entomoneis</i> sp. 1 NCC350	Cell count	0.24 $\pm$ 0.01	16 $\pm$ 2	P=0.29	P=0.09
	Pam	0.37 $\pm$ 0.12	14 $\pm$ 1		
	Spectroradiometer	0.35 $\pm$ 0.06	14 $\pm$ 1		
<i>Extubocellulus</i> <i>cf cribriger</i> NCC229	Cell count	0.68 $\pm$ 0.04	14 $\pm$ 1	P=0.10	P=0.14
	Pam	0.81 $\pm$ 0.01	9 $\pm$ 0		
	Spectroradiometer	0.83 $\pm$ 0.07	10 $\pm$ 3		
<i>Navicula</i> sp. 2 NCC226	Cell count	0.91 $\pm$ 0.04	8 $\pm$ 0	P=0.42*	P=0.78*
	Pam	1.02 $\pm$ 0.23	8 $\pm$ 2		
	Spectroradiometer	0.87 $\pm$ 0.02	9 $\pm$ 2		

¹ Entomoneis paludosa μ max value for the second run was 0.54 and associated P value was 0.47

3.1.1 Diatom strain characteristics

Among the 68 screened strains, 33 were cultivated successfully and 36 did not grow (supplementary data 1). This information is reported in Tab. 3, (with the NCC reference, the sampling location, the cell size, the LED, the growth rate, the silica content, the DW and lipid rate). Globally the cell length varied from 58 $\pm$ 14 μ m (*Craspedostauros* sp. 2 NCC218) to 4.8 $\pm$ 0.7 μ m (*Extubocellulus cf cribriger* NCC229). The cell width varied from 22.6 $\pm$ 4.1 μ m (*Lithodesmium* sp NCC353) to 2.9 $\pm$ 0.4 μ m (*Fallacia* sp. 1 NCC303). The end of the exponential phase, corresponding to the harvest day (LED), varied from day 6 for *Surirella* sp. 1 NCC270 to day 16 for *Craspedostauros britannicus* NCC228 and *Navicula* sp. 1 NCC113 (ANOVA, $p < 0.001$). Three groups were therefore identified depending on their growth rate: group A, included *Nitzschia alexandrina* NCC33 and *Entomoneis* sp. 5 NCC302 with respective growth rates of 1.89 $\pm$ 0.10 and 1.50 $\pm$ 0.18 day⁻¹, group B with 13 species (i.e 40% of the total number of strain) presented a growth rate ranging from 1.19 $\pm$ 0.08 to 0.81 $\pm$ 0.09

355 day⁻¹ and group C with 18 species (54%) had a growth rate below 0.8 day⁻¹ with a minimum at
356 0.22 ± 0.05 day⁻¹.

357 To ensure the standardization of the FTIR results for inter species comparison purposes, the
358 total silica rate was evaluated for the 33 species. It showed a significant difference (ANOVA,
359 $p < 0.01$), where five species (i.e. 15% of the total number of strains) had a silica content of c.a.
360 40% of the dry weight with a maximum of 46.11 ± 4.58% for *Entomoneis* sp 1 NCC350. Three
361 species (i.e. 9% of the total number of strains) presented a silica content of c.a. 23% of the dry
362 weight with a minimum of 22.9 ± 2.6% for *Conticriba weissflogii* CCMP1336. In the other
363 species, representing more than 75% of the total number of strains, the differences were not
364 significant (ANOVA, $p = 0.055$), supporting a stable silica content situated at around 35% DW.
365 The dry weight (DW) biomass values of the assessed strains ranged from 0.502 ± 0.039 g.L⁻¹
366 (*Craspedostauros britannicus* NCC195) to 0.065 ± 0.001 g.L⁻¹ (*Entomoneis* sp. 3 NCC351).
367 For three strains (9% of the total number of strains) the DW was greater than 0.30 g.L⁻¹. For 27
368 strains (81%) the biomass was greater than 0.10 g.L⁻¹ but lower than 0.30 g.L⁻¹. For the three
369 remaining strains (9%) the biomass was lower than 0.10 g.L⁻¹.

370 The lipid rate estimated with the gravimetric method ranged from 30.5 ± 0.7% DW (*Nitzschia*
371 sp. 5 NCC109) to 3.7 ± 1.1% DW (*Brockmaniella brockmanii* NCC161). For three strains (9%)
372 the lipid rate was greater than 20% DW. For 14 strains (42%) the lipid rate was greater than
373 10% DW. In the remaining 16 strains (48%) the lipid rate was lower than 10% DW.

374 Biomass productivity varied substantially among the tested strains (Fig. 2A) and ranged from
375 0.36 ± 0.02 g.L⁻¹.day⁻¹ (*Nitzschia alexandrina* NCC33) to 1.4 ± 0.3 × 10⁻² g.L⁻¹.day⁻¹
376 (*Entomoneis* sp. 3 NCC351). The results were found to be statistically different among those
377 strains (ANOVA, $p < 0.001$). Finally, the strains were clustered into three groups according to
378 the following parameters (Fig. 2): group 1 with productivity between 0.35 ± 0.02 and 0.19 ±
379 0.04 g.L⁻¹.day⁻¹; group 2 with productivity ranging from 0.17 ± 0.08 to 0.09 ± 0.02 g.L⁻¹.day⁻¹;

380 group 3 with lower productivity ranging from $7.5 \pm 0.4 \times 10^{-2}$ to $1.4 \pm 0.3 \times 10^{-2}$ (g.L⁻¹.day⁻¹).

381 The most productive strains were those exhibiting the highest $\mu_{\max}$ associated to the highest

382 DW. But they did not correspond to the richest in terms of total lipid rate (Fig. 2B).

383

384

385

386
387

Table 3. Characteristics and sampling locations of the investigated diatoms species. All data were obtained by experimental measurements. N=3, independent measurement, $\pm$ SD.

Species	NCC strain identification	Sampling location	Cell size (μ m)		LED (day)	μ max	Silica content (%)	Biomass (g.L ⁻¹)	TLR (% DW)	Ref
			Length	Width						
<i>Nitzschia alexandrina</i>	NCC33	France, NW Atlantic coast	11.0 $\pm$ 1.2	3.7 $\pm$ 0.4	11 $\pm$ 0	1.89 $\pm$ 0.10	A	27.2 $\pm$ 3.8	0.189 $\pm$ 0.004	[15,42–49]
<i>Entomoneis</i> sp. 5	NCC302	France, NW Atlantic coast	22.4 $\pm$ 3.5	15.0 $\pm$ 2.2	9 $\pm$ 0	1.50 $\pm$ 0.18		38.9 $\pm$ 0.7	0.132 $\pm$ 0.025	[50–54]
<i>Entomoneis</i> sp. 2	NCC20	France, NW Atlantic coast	30.5 $\pm$ 4.6	18.5 $\pm$ 3.2	8 $\pm$ 0	1.19 $\pm$ 0.08		34.3 $\pm$ 2.8	0.244 $\pm$ 0.042	[50–54]
<i>Staurosira</i> sp.	NCC182	France, NW Atlantic coast	6.9 $\pm$ 1.4	5.5 $\pm$ 1.2	8 $\pm$ 2	1.17 $\pm$ 0.09	B	28.9 $\pm$ 5.1	0.262 $\pm$ 0.084	[16]
<i>Fallacia</i> sp. 1	NCC303	Greenland, east coast	6.4 $\pm$ 0.7	2.9 $\pm$ 0.4	12 $\pm$ 2	1.12 $\pm$ 0.14		37.4 $\pm$ 4.5	0.154 $\pm$ 0.016	-
<i>Fallacia</i> sp. 2	NCC304	Greenland, east coast	6.4 $\pm$ 0.8	3.1 $\pm$ 0.4	12 $\pm$ 0	1.07 $\pm$ 0.15		31.8 $\pm$ 4.9	0.265 $\pm$ 0.028	-
<i>Entomoneis</i> sp. 7	NCC445	France, NW Atlantic coast	10.4 $\pm$ 1.5	4.1 $\pm$ 0.7	9 $\pm$ 2	1.02 $\pm$ 0.05		41.4 $\pm$ 2.9	0.122 $\pm$ 0.005	[50–54]
<i>Entomoneis paludosa</i>	NCC18.1.1	France, NW Atlantic coast	18.4 $\pm$ 3.3	10.7 $\pm$ 1.9	11 $\pm$ 1	0.96 $\pm$ 0.03		37.8 $\pm$ 4.0	0.130 $\pm$ 0.016	[50–54]
<i>Entomoneis</i> sp. 4	NCC301	France, NW Atlantic coast	23.9 $\pm$ 3.4	17.3 $\pm$ 2.2	7 $\pm$ 1	0.92 $\pm$ 0.14		26.7 $\pm$ 3.9	0.171 $\pm$ 0.012	[50–54]
<i>Pseudonitzschia americana</i>	PNA06 KER	France, NW Atlantic coast	6.6 $\pm$ 0.8	4.5 $\pm$ 0.5	9 $\pm$ 2	0.89 $\pm$ 0.09		36.4 $\pm$ 0.6	0.099 $\pm$ 0.019	-
<i>Surirella</i> sp. 1	NCC270	France, NW Atlantic coast	6.8 $\pm$ 1.3	5.0 $\pm$ 1.6	6 $\pm$ 1	0.85 $\pm$ 0.31		33.7 $\pm$ 3.5	0.173 $\pm$ 0.033	-
<i>Navicula</i> sp. 2	NCC226	France, NW Atlantic coast	9.1 $\pm$ 1.3	5.6 $\pm$ 0.7	8 $\pm$ 2	0.82 $\pm$ 0.05		41.4 $\pm$ 1.5	0.131 $\pm$ 0.015	[15,45,47,49,55,58]
<i>Opephora</i> sp. 1	NCC366	France, NW Atlantic coast	5.4 $\pm$ 0.8	3.7 $\pm$ 0.8	14 $\pm$ 1	0.82 $\pm$ 0.16		37.8 $\pm$ 8.4	0.233 $\pm$ 0.010	-
<i>Entomoneis paludosa</i>	NCC18.2.1	France, NW Atlantic coast	18.8 $\pm$ 1.8	11.2 $\pm$ 2.5	12 $\pm$ 1	0.81 $\pm$ 0.05	C	36.0 $\pm$ 3.9	0.150 $\pm$ 0.003	[50–54]
<i>Extubocellulus cf cribriger</i>	NCC229	France, NW Atlantic coast	4.8 $\pm$ 0.7	3.9 $\pm$ 0.5	9 $\pm$ 0	0.81 $\pm$ 0.09		33.7 $\pm$ 2.3	0.155 $\pm$ 0.003	[52,59,60]
<i>Amphora</i> sp. 1	NCC260	France, NW Atlantic coast	9.8 $\pm$ 1.3	3.7 $\pm$ 0.5	13 $\pm$ 0	0.78 $\pm$ 0.30		36.2 $\pm$ 2.6	0.176 $\pm$ 0.041	[49,55,58,61,62]
<i>Conticriba weissflogii</i>	CCMP1336	USA, NE Atlantic coast	15.6 $\pm$ 2.5	10.8 $\pm$ 1.4	12 $\pm$ 0	0.59 $\pm$ 0.15		22.9 $\pm$ 2.6	0.277 $\pm$ 0.035	-
<i>Brockmaniella brockmanii</i>	NCC161	France, NW Atlantic coast	9.2 $\pm$ 2.4	4.3 $\pm$ 0.6	10 $\pm$ 0	0.58 $\pm$ 0.01		37.5 $\pm$ 0.2	0.185 $\pm$ 0.025	-
<i>Craspedostauros britannicus</i>	NCC195	France, NW Atlantic coast	37.8 $\pm$ 8.8	11.0 $\pm$ 2.2	9 $\pm$ 0	0.53 $\pm$ 0.09		44.7 $\pm$ 0.1	0.502 $\pm$ 0.039	-
<i>Conticriba weissflogii</i>	NCC133	Morocco, Oum R'bia estuary	14.3 $\pm$ 1.7	12.1 $\pm$ 1.7	7 $\pm$ 0	0.53 $\pm$ 0.08		34.6 $\pm$ 1.2	0.181 $\pm$ 0.015	-
<i>Licmophora</i> sp. 1	NCC253	France, NW Atlantic coast	20.0 $\pm$ 3.9	12.4 $\pm$ 1.6	11 $\pm$ 0	0.51 $\pm$ 0.02		37.8 $\pm$ 5.7	0.161 $\pm$ 0.049	-
<i>Amphora</i> sp. 2	NCC169	France, NW Atlantic coast	9.0 $\pm$ 1.2	5.8 $\pm$ 1.2	13 $\pm$ 3	0.48 $\pm$ 0.14		33.8 $\pm$ 4.8	0.157 $\pm$ 0.024	[49,55,58,61,62]
<i>Craspedostauros</i> sp. 1	NCC228	France, NW Atlantic coast	55 $\pm$ 14	14.9 $\pm$ 4.2	16 $\pm$ 4	0.46 $\pm$ 0.22		25.2 $\pm$ 0.7	0.384 $\pm$ 0.042	-
<i>Craspedostauros</i> sp. 2	NCC218	France, NW Atlantic coast	58 $\pm$ 14	13.6 $\pm$ 3.4	12 $\pm$ 0	0.45 $\pm$ 0.18		26.0 $\pm$ 4.1	0.178 $\pm$ 0.047	-
<i>Cymatosira belgica</i>	NCC208	France, NW Atlantic coast	5.6 $\pm$ 0.6	4.7 $\pm$ 0.7	12 $\pm$ 0	0.44 $\pm$ 0.12		35.5 $\pm$ 3.4	0.118 $\pm$ 0.043	-
<i>Amphora acutiuscula</i>	NCC216	Viet Nam, South east coast	10.0 $\pm$ 1.0	5.6 $\pm$ 0.8	11 $\pm$ 0	0.43 $\pm$ 0.07		32.4 $\pm$ 0.7	0.144 $\pm$ 0.041	[49,55,58,61,62]
<i>Lithodesmium</i> sp	NCC353	Australia, Moreton bay	29.3 $\pm$ 5.4	22.6 $\pm$ 4.1	9 $\pm$ 1	0.43 $\pm$ 0.02		34.1 $\pm$ 6.1	0.129 $\pm$ 0.006	-
<i>Navicula</i> sp. 1	NCC113	Morocco, Oum R'bia estuary	17.1 $\pm$ 2.9	5.1 $\pm$ 1.2	16 $\pm$ 0	0.42 $\pm$ 0.02		25.0 $\pm$ 1.0	0.168 $\pm$ 0.035	[15,45,47,49,55,58]
<i>Entomoneis</i> sp. 6	NCC335	France, Mediterranean Sea	13.9 $\pm$ 1.9	7.7 $\pm$ 1.9	12 $\pm$ 1	0.38 $\pm$ 0.03		36.8 $\pm$ 4.7	0.226 $\pm$ 0.036	[50–54]
<i>Entomoneis</i> sp. 1	NCC350	France, Mediterranean Sea	29.0 $\pm$ 4.0	17.5 $\pm$ 2.7	14 $\pm$ 0	0.36 $\pm$ 0.13		46.1 $\pm$ 4.6	0.098 $\pm$ 0.048	[50–54]
<i>Craspedostauros britannicus</i>	NCC199	France, NW Atlantic coast	31.2 $\pm$ 3.2	8.8 $\pm$ 1.6	12 $\pm$ 2	0.35 $\pm$ 0.14		40.0 $\pm$ 2.9	0.421 $\pm$ 0.053	-
<i>Nitzschia</i> sp. 5	NCC109	France, NW Atlantic coast	33.1 $\pm$ 4.1	10.1 $\pm$ 1.5	12 $\pm$ 1	0.31 $\pm$ 0.05		23.0 $\pm$ 1.5	0.239 $\pm$ 0.026	[15,42–49]
<i>Entomoneis</i> sp. 3	NCC351	France, Mediterranean Sea	16.6 $\pm$ 3.7	12.3 $\pm$ 3.0	13 $\pm$ 0	0.22 $\pm$ 0.05		38.6 $\pm$ 5.8	0.065 $\pm$ 0.001	[50–54]

388

389

390

Figure 2. Values of the parameters measured for the screened strains includes strain productivity (A), lipid rates as measured with the gravimetric method (B), lipid ratio measured semi-quantitatively by the FTIR approaches, $[(eb+CH_3+CH_3)/si]$ (C) and $[area\ eb+CH_2+CH_3)/Total\ area]$ multiplied by 100 for scaling purposes (D). Notations a, b, and c correspond to the maximum and minimum values for groups 1, 2 and 3. N=3, independent measurements, $\pm$ SD.

3.2 FTIR analysis

3.2.1 FTIR spectrum interpretation

HTSXT-FTIR analysis was performed on the 33 assayed species (Tab. 3). The lipid rate was associated to three main signals on the recorded spectra (Fig. 3A), i.e. through the two vibrations of the fatty acid carbon chains ($\nu\ CH_2$ and $\nu\ CH_3$) ($\nu\ C-H \sim 2923$ and $2852\ cm^{-1}$) [26] and of the ester bond function (Eb) ($\nu\ C=O \sim 1750\ cm^{-1}$) [63]. The other major bands corresponded to the principal cellular components such as the proteins (the amide I band $\nu\ C=O \sim 1650\ cm^{-1}$; the amide II band $\delta\ N-H \sim 1540\ cm^{-1}$), the nucleic acids ($\nu\ P=O \sim 1230\ cm^{-1}$) and the carbohydrates band superimposed on the silica band ($\sim 900 - 1200\ cm^{-1}$). For details see Wagner et al., 2010 [64]. Whereas the infrared signature obtained on the whole cells showed superimposed bands of silica and carbohydrates at $1078\ cm^{-1}$ (Fig. 3A), the signature obtained on the crude lipid extract (Fig. 3B) did not exhibit this band, but a well defined ester bond (Eb) band at $1750\ cm^{-1}$ and well-defined bands for the CH_2-CH_3 signature at $3000-2800\ cm^{-1}$.

The lipid ratio estimated from the FTIR data measured on the entire cells (Fig.2C) ranged from 1.70 ± 0.59 (*Nitzschia* sp. 5 NCC109) to 0.49 ± 0.04 (*Entomoneis paludosa* NCC18.1.1). It was thus possible to cluster the assessed strains into three groups. Group 1 showed a maximum ratio at 1.69 ± 0.18 for *Craspedostauros britannicus* NCC195 and a minimum at 0.61 ± 0.01 for *Entomoneis* sp 5 NCC302, group 2 showed a maximum ratio at 1.34 ± 0.07 for *Fallacia* sp 1 NCC303 and a minimum at 0.49 ± 0.04 for *Entomoneis paludosa*

418 NCC18.1.1 and group 3 showed a maximum ratio at 1.70 ± 0.59 for *Nitzschia* sp NCC109
 419 and minimum at 0.56 ± 0.02 for *Lithodesmium* sp NCC35.3.

420
 421 **Figure 3.** Example of averaged FTIR spectra recorded on entire cells or on the corresponding
 422 lipid extract. (A) *Staurosira* sp NCC182 FTIR signature recorded on the entire cells and (B)
 423 *Staurosira* sp NCC182 FTIR signature recorded on a crude lipid extract. The grey area
 424 corresponds to the variation of the FTIR signal associated to the standard deviation for n=3
 425 independent measurements.

426
 427 The lipid ratio estimated from the FTIR data recorded on the lipid extracts (Fig.2D) ranged
 428 from 49.3 ± 8.5 (*Amphora* sp. 2 NCC169) to 16.5 ± 3.1 (*Entomoneis* sp. 6 NCC335). Three
 429 groups were also proposed regarding this criterion, group 1 showed a maximum ratio at 38.84
 430 ± 0.63 for *Staurosira* sp. NCC182 and a minimum ratio at 22.8 ± 1.8 for *Entomoneis* sp. 5
 431 NCC302, group 2 a maximum ratio at 39 ± 2 for *Extubocellulus* sp. NCC229 and a minimum
 432 at 16.5 ± 1.2 for *Craspedostauros britannicus* NCC228 and group 3 with a maximum ratio at
 433 49.3 ± 8.3 for *Amphora* sp. 2 NCC169 and a minimum at 23 ± 2 for *Entomoneis* sp. 3 NCC351.

The silica amount did not appear to significantly impact on the lipid FTIR signature except in three species for which the FTIR lipid ratio fell between the two FTIR signature sampling methods (i.e. on entire cells or on lipid extract). These were *Craspedostauros brittanicus* NCC195, *Navicula* sp. 4 NCC113 and *Conticriba weissflogii* CCMP1336. For the other species with different silica content, the lipid signature remained stable in both FTIR sampling methods.

3.2.2 Multivariate analysis of the FTIR spectra recorded on entire cells

In order to assess the main differences in terms of biochemical composition among the 33 screened strains, a correspondence analysis approach using the lipid, protein and carbohydrate bands normalized to the silica amounts was performed. The resulting map (Fig. 4) is a classification of the data on two main dimensions. Dimension 1 represents 81% of the initial information and could be associated to the variation of the lipid composition of the assessed strains, ranging from the lowest to the highest amount of total lipids from right to left on the map. Two strains, *Staurosira* sp. NCC182 and *Amphora* sp. 2 NCC169 presented the highest amount in total lipids. Along dimension 2, representing 19% inertia, the strains associated to that dimension were mainly opposed on the basis of their protein and carbohydrate content. *Craspedostauros* sp. 2 NCC218 was rich in proteins, whereas in *Brockmaniella brockmanii* NCC161 the main fraction was associated to the carbohydrates.

These results indicate that the 2 most notable differences or largest deviations in the sample were observed first between *Staurosira* sp. NCC182, *Amphora* sp. 2 NCC169 and the other species for their lipid rates, and secondly between the strains *Craspedostauros* sp. 2 NCC218, *Brockmaniella brockmanii* NCC161 and the other species by their respective protein and carbohydrate composition. This analysis summarizes the main biochemical characteristics of the strains hosted in the NCC bank in a single step. Although the distance between the

macromolecular content and the species were not mathematically defined, their closeness on the map could be used as a guideline to interpret their biochemical characteristics: the squares correspond to the strains particularly rich in lipids, the triangles, the strains rich in carbohydrates and the dots, the strains particularly rich in proteins.

Figure 4. Correspondence analysis map calculated on the basis of the macromolecular content as evaluated by FTIR on all the assayed strains of the NCC. N=3 independent measurements.

3.3 Comparison of the lipid amounts estimated by the gravimetric method and the FTIR approaches.

A significant positive correlation was found for all the techniques (gravimetric and FTIR) ($p < 1.10^{-7}$) with a Pearson correlation score R superior to 0.53. For the 33 analyzed species, 14 presented a high lipid ratio ($> 12\%$ DW) using the gravimetric measurements: *Nitzschia alexandrina* NCC33, *Staurosira* sp. NCC182, *Fallacia* sp. 1 NCC303, *Surirella* sp.

474 1 NCC270, *Extubocellulus cf cribriger* NCC229, *Amphora* sp. 1 NCC260, *Licmophora* sp.1
 475 NCC253, *Amphora* sp. 2 NCC169, *Cymatosira belgica* NCC208, *Amphora acutiuscula*
 476 NCC216, *Navicula* sp. 1 NCC113, *Craspedostauros britannicus* NCC199, *Nitzschia* sp. 5
 477 NCC109 and *Entomoneis* sp.3 NCC351 (Fig. 2B, Fig. 5). The FTIR method applied on entire
 478 cells identified 12 species rich in lipids (FTIR lipid ratio > 1.20) with nine species identified
 479 in common with the gravimetric method: NCC33, NCC182, NCC303, NCC270, NCC169,
 480 NCC216, NCC113, NCC199 and NCC109 (Fig. 2C, Fig. 5). FTIR for the lipid extract
 481 analyses identified 12 species rich in lipids (ATR lipid ratio > 30) with eight species identified
 482 in common with the gravimetric approach: NCC182, NCC303, NCC270, NCC229, NCC169,
 483 NCC216, NCC113 and NCC109 (Fig 2D, Fig. 5).
 484 The correspondence analyses (Fig.4) performed on the FTIR profiles obtained in the entire
 485 cells gave supplementary information and identified seven species particularly rich in lipids:
 486 NCC182, NCC270, NCC169, NCC113, NCC366, NCC109 and NCC33. These species were
 487 also identified by the gravimetric method with the exception of NCC366, only identified by
 488 the FTIR method on crude lipid extract.

489

Figure 5 Venn diagram showing the degree of overlap among the different approaches used to identify the lipid rich diatoms. In the gravimetric method circle, 14 strains were identified as rich in lipids: 4 only with this method (NCC260, NCC208, NCC253, NCC351), one strains were also identified as rich and lipid by the FTIR on lipid extract (NCC229), and two strains were also identified as rich in lipids by the FTIR on entire cells (NCC33, NCC199). One strain was identified by FTIR on lipid extract and FTIR on whole cells (NCC366) and seven strains were identified as rich in lipids by all three methods (NCC109, NCC182, NCC113, NCC270, NCC169, NCC303, NCC216).

3.4 Strain selection

The selection of the strains exhibiting both high biomass productivity and high lipid rates, was performed using whole sample distribution based on the lipid rates as estimated by the FTIR approaches, gravimetry and strain biomass productivity as estimated by the fluorometry. On the boxplots summarizing this data (Fig. 6), the colored dots represent the species with the highest potential for biotechnology applications based on lipid molecules: *Nitzschia alexandrina* NCC33, *Staurosira* sp NCC182 and *Opephora* sp NCC366 presented a lipid ratio whatever the considered technique and productivity above the median. *Amphora* sp 2 NCC169 and *Nitzschia* sp 5 NCC109 were also identified with an above median lipid ratio, but with lower productivity. Both these species were finally selected for their high lipid rate, even though their productivity needs to be improved.

Figure 6. Boxplots summarizing the sample distribution criteria as measured with FTIR methods, gravimetry for the lipid rate and fluorimetry for productivity. FTIR data was expressed in arbitrary units. Lipid rate in %DW and Px in $\text{g.L}^{-1}.\text{day}^{-1}$. FTIR results were

multiplied by 20 for scaling purposes. The Px was multiplied by 100 for scaling purposes. 33 strains were assayed in independent biological triplicates.

4 Discussion

4.1.1 Determination of growth by fluorimetry

In this study a rapid and precise method for the estimation of cell abundance was needed to efficiently establish the growth parameter of over 60 benthic diatom species. The use of microscope counting chambers is slow, tedious and imprecise. The use of particle counters is feasible only for those phytoplankton species within a certain size range that do not form chains and have long appendages [65]. *In vivo* fluorescence has been successfully used in the past to monitor the growth of phytoplankton culture [19, 20, 32, 41, 66]. *In vivo* fluorescence measurement is rapid, sensitive and can be used with all types of diatoms cell structures, i.e. on solitary cells such as *Entomoneis* and *Navicula* genera and on chain-forming diatoms, such as *Extubocellulus* genus or aggregates of cells such as *Amphora* genus. However, *in vivo* fluorescence is a measurement of the increase in chlorophyll and chlorophyll per cell varies greatly with light intensity and cellular nutritional status so it can only be used to measure growth when culture conditions are constant. Changes in light and other culture conditions can lead to a change in chlorophyll a content. In this case F0 will also change, introducing a bias in growth rate estimation. Reflectance can also be used to monitor the diatom growth rate, as demonstrated in this study; the NDVI was highly correlated with the cell count as *in vivo* fluorescence ($R > 0.70$). Méléder et al, 2003 [21] have already demonstrated this correlation on monospecific benthic diatom cultures with *Entomoneis paludosa* and *Navicula ramosissima*. Compared to fluorometry this technique required more culture sampling and a filtration step which took longer to set up. Fluorometry was more sensitive than spectroradiometry and required less material, which made it ideal for very low amounts of

biomass. *In vivo* fluorometry demonstrated several advantages: ease of use, real-time measurement, non-destructive sampling.

4.1.2 Determination of the lipid rate by FTIR

The total lipid evaluation by gravimetry has been used for more than 50 years [67], but this technique is clearly not compatible with screening efforts involving large numbers of samples because of processing time required particularly in the extraction phase. Furthermore, large volumes of samples are systematically needed for the measurements. It also requires sufficient amounts of dried biological sample, thus making it unsuitable for high frequency monitoring of small scale microalgal cultivation. Feng et al., (2013) [68] suggested that the presence of chlorophyll could also affect the accuracy of the method. In comparison to conventional chemical analysis, FTIR spectroscopy presented striking advantages due to its high reliability, sensitivity and speed of the measurement [64]. An IR spectrometer coupled to a microtiter plate reader open the possibility of high throughput analysis of a few nanograms of cell material [69]. Coat et al, (2014) [26] demonstrated that the repeatability of FTIR signal reached excellent values (10%), but only for a limited range of analyzed quantities of matter (10^6 to 10^8 cells. mL⁻¹).

In the present study we have demonstrated that FTIR was a suitable technique to evaluate the lipid ratio of diatoms. The use of the FTIR technique was even more rapid, due to the use of direct fresh biomass. The results obtained on entire cells and lipid crude extracts were similar, suggesting that measurement on entire cells did not improve the lipid quantification. Measurement on whole cells could be considered sufficient to get a first idea on their intracellular lipid rate. In addition, removing the silica did not seem necessary in view of the present results to improve the lipid semi-quantification.

However, there are some limitations with this non-invasive technique. Even though the FTIR results between entire cells and crude extracts were similar, we did not obtain a perfect linear

correlation with the traditional lipid extraction. Nevertheless, the FTIR method has the advantage of simultaneously detecting the relative amount of lipids, carbohydrates and proteins, even though those components overlapped, implying a certain degree of inaccuracy. The different steps required for the lipid extraction can also produce a bias in the evaluation of the lipid quantity, added to the inaccuracy of weighing of the lipid matter, making it less precise than the spectrometric approach. The heterogeneity observed in the present FTIR results could be also associated to the utilization of different strains (33 strains in the current study). These are not described in the previous studies evaluating lipid quantification with FTIR where only one species was used [26, 39, 70]. FTIR did not differentiate polar/apolar lipids, or the different types of fatty acids produced by the strains. Lipid analysis by chromatographic techniques coupled to mass spectrometry will be needed to further identify the presence of interesting molecules like EPA or DHA. Nevertheless, the present study opens the way to rapid and reliable semi-quantification of total amounts of intracellular lipids in diatoms using a fast, non-invasive approach.

4.1.3 Method assessments

This study highlighted the efficiency of PAM and FTIR measurements as fast techniques to characterize both the growth and lipid content of microalgae (Tab.4). The use of PAM fluorometry was three times faster than cell counting. Where 15 minutes were necessary to count 3 samples (8.25 hours for 99 samples) and 10 minutes to measure reflectance by spectroradiometry (5.5 hours for 99 samples), only 5 minutes were necessary to measure F0 by PAM fluorometry (2.75 hours for 99 samples).

Table 4 Evaluation of three methods for algal growth kinetic determination

Evaluation index	Counting	NDVI	PAM
Cultivation scale	Large	Large–small	Large–small
Monitoring frequency	High	Medium	High
Sample state	Liquid	Filtered	Liquid
Volume consumed	5 - 1 mL	5- 3 mL	>1 mL
Time consumed	8.25 h	5.5 h	2.75 h
Equipment	Counting chambers	Spectroradiometer	PAM fluorometer
Reliability	Low	High	High

For lipid extraction, dichloromethane methanol solvent is usually used [34, 71–73]. Even though extraction was very effective, this method is known to have environmental and health risks [71]. Moreover, it is expensive: for 99 extractions, 10 L of each solvent was necessary costing 234 € (9.68 €/L of methanol and 13.72 €/L of dichloromethane (Fischer Scientific). The time necessary for the lipid extraction using Bligh and Dyer method was time-consuming (Tab. 5) due to the time needed for maceration and lipid separation. 5.5 weeks were necessary to perform the extraction of 99 samples (ca. 7920 hours). Nile red, a lipid soluble fluorescent dye, is commonly used to evaluate the lipid content of animal cells, microorganisms, and especially microalgal strains. Required time to perform spectrophotometric measurements with Nile red and Bodipy was evaluated at 10 mins per sample (measurement before and after Nile red or Bodipy application) and 25 to 40 mins incubation time was necessary after

application of Nile red or Bodipy before reading spectrophotometric values [68,74]. In comparison FTIR is a fast and eco-friendly technique. Two plates of 384 well were necessary to evaluate the lipid content for the 33 strains (each strain needed 15 wells with 5 wells per replicate). It took 30 seconds to read each well to obtain the FTIR spectra leading to ca. and 4 hours to read an entire plate.

Table 5 Evaluation of three methods for algal lipid content determination

Evaluation index	Gravimetric	FT-IR	Nile Red / Bodipy 505/515
Cultivation scale	Large	Large–small	Large–small
Monitoring frequency	Low	Medium	High
Sample state	Dried	Liquid	Liquid
Biomass consumed	100 mg	5 μ L of 1.0 mg mL ⁻¹	3 mL of OD ₆₈₀ 0.4
Time consumed	>50 h	8 h	25 h
Equipment	Nitrogen evaporator	FT-IR	FS & FM ^a
Accuracy	Total lipid	Total lipid	Neutral lipid

^a Fluorescence spectrophotometer and Fluorescence microscopy

Nile red and Bodipy 505/515 staining are powerful quantification tools in terms of time and cost of biomass [75, 76], high throughput quantification method of lipids with Nile red or Bodipy 505/515 fluorescence can hardly been seen as a method for screening different species of microalgae, as the staining protocol is species specific. The significant disadvantages of Nile red were its limited photostability, interference with chlorophyll [77, 78], and difficulty of permeation for some species. Bodipy 505/515 produced a better marker than Nile red for visualizing neutral lipid content in fluorescence microscopy [75, 79] but some authors have reported disadvantages with these techniques such as background fluorescence of the dye in the medium and failure to quantify neutral lipids between rich and low oil strains. When microalgae were cultured on a large scale with a low-frequency monitoring requirement, any of the three methods could be adopted, although gravimetric determination might be preferable as it was an absolute method for quantification of both crude and neutral lipids without the need of specialized equipment. For the general laboratory culture of microalgae,

the FT-IR method for simultaneous characterization of total lipid, carbohydrate and protein content and the Nile Red method for both neutral lipid content and location can be used, both of which are relative quantification methods, but require special equipment.

Although these analyses demonstrate that FTIR and Nile Red were equally effective at measuring lipid accumulation, FTIR was likely to be a more efficient tool for this purpose because of its much faster analysis time and high reproducibility of results [80]. Furthermore, FTIR may also be more suitable than Nile Red for efficiently detecting large increases in lipid concentration. Nile Red does not appear to be efficient at accurately quantifying lipid concentration above 20 mg/ml [77] while FTIR can efficiently detect linear lipid concentration changes up to at least 250 mg [81]. Measurements with FTIR were more precise because of the technical quintuplicate performed for the acquisition of spectra.

4.2 Screening for lipid rich benthic diatom strains

The aim of this study was to investigate the growth characteristics and the lipid rates of the benthic marine diatom species hosted in the NCC bank in order to evaluate their potential for original lipid bioproducts of potential economic interests. First, a selection based on the genera identified in the literature was applied. Among the 134 strains hosted by the NCC, corresponding to 40 genera and 101 species, 23 genera (corresponding to 77 species, 105 strains in the NCC) were largely studied [15, 16, 42, 43, 46, 49, 52, 55, 58, 60, 82]. Among these 23 genera, only 13 genera (42 species, 47 strains) with high productivity and/or producing high lipid quantities were selected for the current study. Among these 47 strains, only 18 strains, (6 genera, 17 species) grew successfully. Among the NCC's 40 genera, 17 genera (corresponding to 24 species, 29 strains) were not previously reported in the literature in terms of productivity or ability to produce lipids or added value molecules. Thus, the 29 strains corresponding to these 17 genera were also selected and assayed for the current study. Among these strains, 15 strains (13 species, 10 genera) did successfully grow. Finally, at the

659 end of this first screening step based on the growth rates, 33 strains (18 previously described
660 and 15 that have never been described before in the literature) were selected for the second
661 step of the screening process. As reported in the supplementary data, 43 strains failed to grow
662 which may be associated to shear stress; some species like *Rhizosolenia setigera* cells were
663 broken during agitation.

664 The second step consisted in the determination of the lipid rate on the selected strains. In
665 microalgae, it can typically vary from 1 to 85% of the dry weight under adverse conditions
666 [15, 83, 84]. Factors such as temperature, irradiance and most markedly nutrient availability
667 have been shown to affect both lipid composition and lipid rate [17, 57, 85]. In general, high
668 irradiance stimulates TAG accumulation [57], while under low irradiance, the polar lipids
669 (phospholipids and glycolipids), structurally and functionally associated to cell membrane,
670 are preferentially synthesized [17]. The lipid rate found in the current study for the *Amphora*
671 genus was similar to that estimated by Renaud et al., (1999) [46]. The authors proposed 19%
672 DW vs. 14 to 19% DW in the current study. This similarity could be explained by the similar
673 culture conditions for light and nutrient although the temperature was different (25°C vs 16°C
674 in the current study). According to Chtourou et al., 2015 [61], the temperature could be an
675 important factor for *Amphora* genus lipid rate which can achieve lipid rates up to 24% of DW
676 at 20°C. Media enrichment could also be an important factor: *Amphora* cells grown in media
677 enriched with macronutrients and trace metal can also achieve lipid rates up to 32% under low
678 light conditions ($11.4 \mu\text{mol photon m}^{-2} \text{s}^{-1}$) [62] or under nitrogen deficiency [55]. These
679 factors could be used as guideline to improve lipid production of the selected *Amphora* strains
680 hosted in the NCC. The biomass measurements for the *Amphora* genus in Zhao et al study
681 [49] were in accordance with our results. The authors found 0.13 g.L^{-1} vs. 0.16 g.L^{-1} in the
682 current study. According to their measured lipid productivity, the *Amphora* genus appeared
683 to be a good candidate for lipid based potential applications.

684 The lipid rate found in our study for the *Nitzschia* genus was similar to the results found by
 685 Renaud et al., (1999) for one species [46] but very different for *Nitzschia sp 5* NCC109 in the
 686 current study. They found a lipid rate of between 13 and 16% DW for the *Nitzschia* genus. In
 687 the present study, *Nitzschia alexandrina* NCC33 showed a lipid rate of 13% DW but for
 688 *Nitzschia sp 5* NCC109 it was estimated up to 30% DW. However, this rate could only be
 689 found for *Nitzschia* genus under nitrogen or silica deficiency increasing up to 45-47% DW
 690 [44,45,47]. Since the amount of nitrogen was not monitored in the culture, it is possible that
 691 nitrogen depletion occurred, explaining the high lipid rate measured for *Nitzschia sp 5*
 692 NCC109. Further analysis for this strain is necessary to establish whether this was due to the
 693 absolute lipid richness of the strain, or to a bias in culture conditions.

694 The biomass found for the *Nitzschia* genus in Zhao et al (2016) [49] was lower than the
 695 measurements in the present study. The biomass estimated in the current study ranged from
 696 0.07 to 0.14 g.L⁻¹ vs. 0.19 to 0.23 g.L⁻¹. This difference could be due to the light/dark cycle
 697 culture conditions, suggesting that under continuous light the biomass production would be
 698 more significant for this specific genus [86].

699 It has been demonstrated that the species with a lipid rate of 30% DW and productivity under
 700 non-optimized conditions of around 0.30 g.L⁻¹ could be potential strains for lipid production
 701 [15, 87]. In the present study *Nitzschia sp 5* NCC109 had the highest lipid rate, 30.51% DW,
 702 *Nitzschia alexandrina* NCC33 with the highest productivity 0.35 g.L⁻¹. Both strains were thus
 703 selected as candidates for further analyses to assay their potential for lipid-based applications.

704 The *Navicula* genus also presented a lipid rate for the 2 tested species of between 9% and 23%
 705 DW. Zhao et al, (2016) [49] found a lipid range from 5 to 30% for 3 *Navicula* species
 706 (*Navicula ramosissima*, *Navicula molli* and *Navicula halophia*) and Scholz et al, (2013) [58]
 707 between 18-25% for 6 *Navicula* species (*Navicula digito-radiata*, *Navicula forcipata*,
 708 *Navicula gregaria*, *Navicula perminuta*, *Navicula phyllepta* and *Navicula salinicola*). The

data were estimated in spite of very different culture conditions (dark night cycle at $600 \mu\text{mol photon m}^{-2} \text{s}^{-1}$). This observation is important, since for that specific genus, the observed variability in the range of lipid rate for the different tested species is the same regardless of the culture conditions used. The biomass measurements found by Zhao et al., (2016) for the *Navicula* genus are in accordance with the present study, ranging from 0.10 to 0.17 g.L^{-1} vs. 0.13 to 0.17 g.L^{-1} suggesting that this genus could produce the identical biomass quantity under light/dark cycle or continuous light.

The lipid rate found for the *Extubocellulus* genus in Slocombe et al., (2015) [60] was 23% DW and the biomass equalled 0.06 g. L^{-1} . The lipid rate for this genus, 15% DW in the actual study was lower but the biomass obtained was higher 0.16 g.L^{-1} . The culture was grown under light/dark cycle in the Slocombe et al. study suggesting that this genus could grow better under continuous light but produce more lipids under light/dark cycle. Wahidin et al., (2013) [88] found the same trend for the microalgae genus *Nannochloropsis*.

Knuckey et al, (2002) [52] found 27% DW of lipid rate for the *Entomoneis* genera. In the present study 9 strains of *Entomoneis* presented lipid rates ranging from 3 to 13% DW. In the Knuckey et al, (2002) study the pH and the nutrients were monitored. The culture conditions were identical to the present study (continuous light, $120 \mu\text{mol photon m}^{-2} \text{s}^{-1}$). It is therefore possible, that for this genus a nutrient limitation occurred and instead of enhancing the lipid production, it has diminished it. It has been reported that this species produces EPA which is interesting for nutraceutical products [52].

In this study the species with the highest lipid rate ($\sim 20 \%$) and the greatest biomass productivity (up to $0.24 \text{ g.L}^{-1}.\text{day}^{-1}$) was *Staurosira* sp NCC182. This species was similar in productivity to the marine microalgae *Nannochloropsis* sp. with productivity higher than $0.21 \text{ g.L}^{-1}.\text{day}^{-1}$ and reached a total lipid rate of 30% DW, when cultivated in batch mode under continuous light and reached 68% DW of lipid production under nitrogen deprivation [89].

734 *Nannochloropsis* sp. was investigated for algal biofuel production due to its ease of growth
 735 and high oil rate. *Staurosira* sp was grown in raceway ponds by Huntley et al., (2015) [16]
 736 and the lipid quantification demonstrated that these strains could reach a lipid rate of 45.5%
 737 DW under low N content. In the present study, *Staurosira* sp. NCC182 presented all the
 738 characteristics (good productivity and oil content) to be produced on a large scale and was
 739 considered to be one of the most promising candidates for lipid-based applications.
 740 Even if the microalgae oil yield is strain-dependent it is generally superior to other vegetable
 741 crops [15, 87]. Oil content in Corn, Hemp, Soybean, Sunflower or Palm oil varied from 18
 742 (Soybean) to 44% DW (Corn). Christie et al., (2007) [15] demonstrated that microalgae with
 743 a lipid rate up to 30% of DW could produce 58 L/Ha of oil and microalgae up to 70% of DW
 744 could produce 136 L/Ha of oil. Corn can produce 172 L/Ha, soybean 446 L/Ha and oil palm
 745 5950 L/Ha but require a lot of land for production: 1540 M/Ha for Corn, 594 M/Ha for
 746 Soybean and 45 M/Ha for oil palm. Microalgae production only requires 2 to 5 M/Ha. In the
 747 NCC collection *Nitzschia* sp 5 NCC109 and *Staurosira* sp NCC182 had all the mandatory
 748 features to be grown on a large scale: good productivity and high lipid rate. In the present
 749 study, *Nitzschia alexandrina* NCC33 had the highest productivity in the NCC collection (0.36
 750 g.L⁻¹.day⁻¹) and presented a lipid rate superior to 10% DW. Despite its low productivity of
 751 0.08 g.L⁻¹.day⁻¹, *Amphora* sp 2 NCC169 was chosen for its lipid rate, superior to 15% DW.
 752 These 4 strains were selected for further analyses and to improve their productivity and lipid
 753 rate with the objective of supplying new resources for lipid based applications. *Entomoneis*
 754 *paludosa* was also selected despite its low lipid rate since it was already characterized for its
 755 ability to produce EPA.
 756 Among the strains not described in the literature, the measured lipid rate ranged from 3 to
 757 20% DW and productivity from 0.05 to 0.27 g.L⁻¹.day⁻¹. Only one genus was selected for
 758 further study: *Opephora* sp 1 NCC366. This species showed a mandatory balance between

relatively high biomass productivity (0.23 g.L^{-1}) and a high intracellular lipid rate (above 10% DW).

5 Conclusion

In this work, we focused on developing an easy to use screening method to explore the NCC bank for diatom strains with the highest relative lipid content. The experimental results showed that the combined use of water-PAM to estimate strain growth kinetics and FTIR on whole cells to estimate the semi-quantitative strain macromolecular content and more specifically lipids, could be rapid, reliable and accessible techniques. The developed methodology opens the way to a systematic, fast, and convenient screening of microorganisms (microalgae in this proof of concept). Moreover, the sensitivity and specificity of the method makes it suitable for a reasonable amount of biomass. This method could also be used in systematic studies for the optimization of culture conditions and to measure the influence of the environment on the metabolic plasticity of the assessed organism. Using this screening approach, 5 strains hosted in the NCC bank were selected for their high productivity and high lipid rate: *Nitzschia alexandrina* NCC33, *Staurosira sp* NCC182, *Opephora sp 1* NCC366, *Nitzschia sp 5* NCC109 and *Amphora sp 2* NCC169. The lipid rate achieved by these strains reached a maximum of 30% DW in the assayed cultivation conditions.

In order to improve the lipid quantity, the selected strains could be grown under different culture conditions. The impact of light, temperature and nutrients, especially nitrogen, could be assayed both in terms of lipid productivity and ecophysiology to ensure the highest growth rate possible. Once optimal conditions are found for those strains, the production in photobioreactors could be tested and productivity and lipid rates evaluated, in order to

estimate quantitatively if the selected strains can compete with the best ones found in the literature [90–92]. Finally, depending on their oil quality and original lipid activities, those strains may constitute new and original genetic resources that could have potential interesting applications (biodiesel, pharmaceutical, etc.).

6 References

- [1] T.M. Mata, A.A. Martins, N.S. Caetano, Microalgae for biodiesel production and other applications: A review, *Renew. Sustain. Energy Rev.* 14 (2010) 217–232. doi:10.1016/j.rser.2009.07.020.
- [2] P. Spolaore, C. Joannis-Cassan, E. Duran, A. Isambert, Commercial applications of microalgae, *J. Biosci. Bioeng.* 101 (2006) 87–96. doi:10.1263/jbb.101.87.

- [3] M.A. Borowitzka, High-value products from microalgae—their development and commercialisation, *J. Appl. Phycol.* 25 (2013) 743–756. doi:10.1007/s10811-013-9983-9.
- [4] M.I. Khan, J.H. Shin, J.D. Kim, The promising future of microalgae: current status, challenges, and optimization of a sustainable and renewable industry for biofuels, feed, and other products, *Microb. Cell Factories.* 17 (2018). doi:10.1186/s12934-018-0879-x.
- [5] Y.-F. Niu, M.-H. Zhang, D.-W. Li, W.-D. Yang, J.-S. Liu, W.-B. Bai, H.-Y. Li, Improvement of Neutral Lipid and Polyunsaturated Fatty Acid Biosynthesis by Overexpressing a Type 2 Diacylglycerol Acyltransferase in Marine Diatom *Phaeodactylum tricornutum*, *Mar. Drugs.* 11 (2013) 4558–4569. doi:10.3390/md11114558.
- [6] Z. Yi, M. Xu, X. Di, S. Brynjolfsson, W. Fu, Exploring Valuable Lipids in Diatoms, *Front. Mar. Sci.* 4 (2017). doi:10.3389/fmars.2017.00017.
- [7] E. Artamonova, J. Svenning, T. Vasskog, E. Hansen, H. Eilertsen, Analysis of phospholipids and neutral lipids in three common northern cold water diatoms: *Coscinodiscus concinnus*, *Porosira glacialis*, and *Chaetoceros socialis*, by ultra-high performance liquid chromatography-mass spectrometry, *J. Appl. Phycol.* 29 (2017) 1241–1249.
- [8] K.W. Chew, J.Y. Yap, P.L. Show, N.H. Suan, J.C. Juan, T.C. Ling, D.-J. Lee, J.-S. Chang, Microalgae biorefinery: High value products perspectives, *Bioresour. Technol.* 229 (2017) 53–62. doi:10.1016/j.biortech.2017.01.006.
- [9] G.A. Dunstan, J.K. Volkman, S.M. Barrett, J.-M. Leroi, S.W. Jeffrey, Essential polyunsaturated fatty acids from 14 species of diatom (Bacillariophyceae), *Phytochemistry.* 35 (1993) 155–161. doi:10.1016/S0031-9422(00)90525-9.
- [10] Y.-C. Chen, The biomass and total lipid content and composition of twelve species of marine diatoms cultured under various environments, *Food Chem.* 131 (2012) 211–219. doi:10.1016/j.foodchem.2011.08.062.
- [11] L. Chuecas, J.P. Riley, Component Fatty Acids of the Total Lipids of Some Marine Phytoplankton, *J. Mar. Biol. Assoc. U. K.* 49 (1969) 97. doi:10.1017/S0025315400046439.
- [12] K. Nagao, T. Yanagita, Conjugated fatty acids in food and their health benefits, *J. Biosci. Bioeng.* 100 (2005) 152–157. doi:10.1263/jbb.100.152.
- [13] M. Gross, The mysteries of the diatoms, *Curr. Biol.* 22 (2012) R581–R585. <http://www.sciencedirect.com/science/article/pii/S0960982212008664> (accessed September 12, 2016).
- [14] T. Lebeau, J.-M. Robert, Diatom cultivation and biotechnologically relevant products. Part II: Current and putative products, *Appl. Microbiol. Biotechnol.* 60 (2003) 624–632. doi:10.1007/s00253-002-1177-3.
- [15] Y. Chisti, Biodiesel from microalgae, *Biotechnol. Adv.* 25 (2007) 294–306. doi:10.1016/j.biotechadv.2007.02.001.
- [16] M.E. Huntley, Z.I. Johnson, S.L. Brown, D.L. Sills, L. Gerber, I. Archibald, S.C. Machesky, J. Granados, C. Beal, C.H. Greene, Demonstrated large-scale production of marine microalgae for fuels and feed, *Algal Res.* 10 (2015) 249–265. doi:10.1016/j.algal.2015.04.016.
- [17] Q. Hu, M. Sommerfeld, E. Jarvis, M. Ghirardi, M. Posewitz, M. Seibert, A. Darzins, Microalgal triacylglycerols as feedstocks for biofuel production: perspectives and advances, *Plant J.* 54 (2008) 621–639. doi:10.1111/j.1365-313X.2008.03492.x.
- [18] T.A. Dempster, M.R. Sommerfeld, Effects of environmental conditions on growth and lipid accumulation in *Nitzschia communis* (Bacillariophyceae), *J. Phycol.* 34 (1998) 712–721.

- [19] V. Vyhnaček, Z. Fišar, A. Fišarová, J. Komarková, In vivo fluorescence of chlorophyll a: Estimation of phytoplankton biomass and activity in Římov Reservoir (Czech Republic), *Water Sci. Technol.* 28 (1993) 29–33. <http://wst.iwaponline.com/content/28/6/29.abstract> (accessed October 10, 2017).
- [20] A.D. Steinman, G.A. Lamberti, P.R. Leavitt, D.G. Uzarski, Biomass and Pigments of Benthic Algae, in: *Methods Stream Ecol.* Vol. 1, Elsevier, 2017: pp. 223–241. doi:10.1016/B978-0-12-416558-8.00012-3.
- [21] V. Méléder, L. Barillé, P. Launeau, V. Carrère, Y. Rincé, Spectrometric constraint in analysis of benthic diatom biomass using monospecific cultures, *Remote Sens. Environ.* 88 (2003) 386–400. doi:10.1016/j.rse.2003.08.009.
- [22] L. Akoto, R. Pel, H. Irth, A.T. Udo, R.J. Vreuls, Automated GC–MS analysis of raw biological samples: Application to fatty acid profiling of aquatic micro-organisms, *J. Anal. Appl. Pyrolysis.* 73 (2005) 69–75.
- [23] K.E. Cooksey, J.B. Guckert, S.A. Williams, P.R. Callis, Fluorometric determination of the neutral lipid content of microalgal cells using Nile Red, *J. Microbiol. Methods.* 6 (1987) 333–345.
- [24] J. Koreivienė, Microalgae Lipid Staining with Fluorescent BODIPY Dye, (2017).
- [25] D. Jaeger, C. Pilger, H. Hachmeister, E. Oberländer, R. Wördenweber, J. Wichmann, J.H. Mussnug, T. Huser, O. Kruse, Label-free in vivo analysis of intracellular lipid droplets in the oleaginous microalga *Monoraphidium neglectum* by coherent Raman scattering microscopy, *Sci. Rep.* 6 (2016) 35340.
- [26] R. Coat, V. Montalescot, E.S. León, D. Kucma, C. Perrier, S. Jubeau, G. Thouand, J. Legrand, J. Pruvost, O. Gonçalves, Unravelling the matrix effect of fresh sampled cells for in vivo unbiased FTIR determination of the absolute concentration of total lipid content of microalgae, *Bioprocess Biosyst. Eng.* 37 (2014) 2175–2187.
- [27] J. Fan, Y. Cui, M. Wan, W. Wang, Y. Li, Lipid accumulation and biosynthesis genes response of the oleaginous *Chlorella pyrenoidosa* under three nutrition stressors, *Biotechnol. Biofuels.* 7 (2014) 17. doi:10.1186/1754-6834-7-17.
- [28] P. Metzger, C. Largeau, *Botryococcus braunii*: a rich source for hydrocarbons and related ether lipids, *Appl. Microbiol. Biotechnol.* 66 (2005) 486–496. doi:10.1007/s00253-004-1779-z.
- [29] R.R. Guillard, Culture of phytoplankton for feeding marine invertebrates, in: *Cult. Mar. Invertebr. Anim.*, Springer, 1975: pp. 29–60.
- [30] R.R. Guillard, J.H. Ryther, Studies of marine planktonic diatoms: I. *Cyclotella* Nana Hustedt, and *Detonula Confervacea* (CLEVE) Gran., *Can. J. Microbiol.* 8 (1962) 229–239.
- [31] Z.-K. Yang, Y.-F. Niu, Y.-H. Ma, J. Xue, M.-H. Zhang, W.-D. Yang, J.-S. Liu, S.-H. Lu, Y. Guan, H.-Y. Li, Molecular and cellular mechanisms of neutral lipid accumulation in diatom following nitrogen deprivation, *Biotechnol. Biofuels.* 6 (2013) 67. <https://biotechnologyforbiofuels.biomedcentral.com/articles/10.1186/1754-6834-6-67> (accessed October 6, 2017).
- [32] C. Honeywill, D. Paterson, S. Hagerthey, Determination of microphytobenthic biomass using pulse-amplitude modulated minimum fluorescence, *Eur. J. Phycol.* 37 (2002) 485–492. doi:10.1017/S0967026202003888.
- [33] B. Gompertz, XXIV. On the nature of the function expressive of the law of human mortality, and on a new mode of determining the value of life contingencies. In a letter to Francis Baily, Esq. FRS &c, *Philos. Trans. R. Soc. Lond.* 115 (1825) 513–583.
- [34] E.G. Bligh, W.J. Dyer, A RAPID METHOD OF TOTAL LIPID EXTRACTION AND PURIFICATION, *Can. J. Biochem. Physiol.* 37 (1959) 911–917. doi:10.1139/y59-099.

- [35] G. Breuer, W.A. Evers, J.H. de Vree, D.M. Kleinegris, D.E. Martens, R.H. Wijffels, P.P. Lamers, Analysis of fatty acid content and composition in microalgae, *J. Vis. Exp. JoVE*. (2013).
- [36] E.S. León, R. Coat, B. Moutel, J. Pruvost, J. Legrand, O. Gonçalves, Influence of physical and chemical properties of HTSXT-FTIR samples on the quality of prediction models developed to determine absolute concentrations of total proteins, carbohydrates and triglycerides: a preliminary study on the determination of their absolute concentrations in fresh microalgal biomass, *Bioprocess Biosyst. Eng.* 37 (2014) 2371–2380.
- [37] W. Zeroual, C. Choisy, S.M. Doglia, H. Bobichon, J.-F. Angiboust, M. Manfait, Monitoring of bacterial growth and structural analysis as probed by FT-IR spectroscopy, *Biochim. Biophys. Acta BBA - Mol. Cell Res.* 1222 (1994) 171–178. doi:10.1016/0167-4889(94)90166-X.
- [38] D.H. Williams, I. Fleming, *Spectroscopic methods in organic chemistry*, McGraw-Hill, 1980.
- [39] I. Schaub, H. Wagner, M. Graeve, U. Karsten, Effects of prolonged darkness and temperature on the lipid metabolism in the benthic diatom *Navicula perminuta* from the Arctic Adventfjorden, Svalbard, *Polar Biol.* (2017) 1–15. doi:10.1007/s00300-016-2067-y.
- [40] N. Sourial, C. Wolfson, B. Zhu, J. Quail, J. Fletcher, S. Karunanathan, K. Bandeen-Roche, F. Béland, H. Bergman, Correspondence analysis is a useful tool to uncover the relationships among categorical variables, *J. Clin. Epidemiol.* 63 (2010) 638–646. doi:10.1016/j.jclinepi.2009.08.008.
- [41] C.J. Lorenzen, A method for the continuous measurement of in vivo chlorophyll concentration, in: *Deep Sea Res. Oceanogr. Abstr.*, Elsevier, 1966: pp. 223–227. <http://www.sciencedirect.com/science/article/pii/0011747166911028> (accessed October 10, 2017).
- [42] G. Chen, Y. Jiang, F. Chen, Fatty acid and lipid class composition of the eicosapentaenoic acid-producing microalga, *Nitzschia laevis*, *Food Chem.* 104 (2007) 1580–1585. doi:10.1016/j.foodchem.2007.03.008.
- [43] M.C. Dalay, S.S. Gunes, others, Biodiesel from Microalgae: A Renewable Energy Source, *Middle East J. Sci. Res.* 22 (2014) 350–355. [http://idosi.org/mejsr/mejsr21\(12\)14/15.pdf](http://idosi.org/mejsr/mejsr21(12)14/15.pdf) (accessed February 19, 2016).
- [44] M.J. Griffiths, S.T.L. Harrison, Lipid productivity as a key characteristic for choosing algal species for biodiesel production, *J. Appl. Phycol.* 21 (2009) 493–507. doi:10.1007/s10811-008-9392-7.
- [45] J. Johansen, P. Lemke, N. Nagle, P. Chelf, P. Roessler, R. Galloway, S. Toon, Addendum to Microalgae Culture Collection 1986-1987, National Renewable Energy Lab.(NREL), Golden, CO (United States), 1987.
- [46] S.M. Renaud, L.-V. Thinh, D.L. Parry, The gross chemical composition and fatty acid composition of 18 species of tropical Australian microalgae for possible use in mariculture, *Aquaculture*. 170 (1999) 147–159.
- [47] J. Sheehan, T. Dunahay, J. Benemann, P. Roessler, Look Back at the U.S. Department of Energy's Aquatic Species Program: Biodiesel from Algae; Close-Out Report, National Renewable Energy Lab., Golden, CO. (US), 1998. doi:10.2172/15003040.
- [48] Z.-Y. Wen, F. Chen, Production potential of eicosapentaenoic acid by the diatom *Nitzschia laevis*, *Biotechnol. Lett.* 22 (2000) 727–733. <http://link.springer.com/article/10.1023/A:1005666219163> (accessed February 18, 2016).

- 956 [49] F.-Y. Zhao, J.-R. Liang, Y.-H. Gao, Q.-Q. Luo, Y. Yu, C.-P. Chen, L. Sun, Variations
957 in the Total Lipid Content and Biological Characteristics of Diatom Species for Potential
958 Biodiesel Production, *J. Fundam. Renew. Energy Appl.* 2016 (2016).
959 [http://www.omicsonline.com/open-access/variations-in-the-total-lipid-content-and-](http://www.omicsonline.com/open-access/variations-in-the-total-lipid-content-and-biological-characteristics-of-diatom-species-for-potential-biodiesel-production-2090-4541-1000201.php?aid=66659)
960 [biological-characteristics-of-diatom-species-for-potential-biodiesel-production-2090-](http://www.omicsonline.com/open-access/variations-in-the-total-lipid-content-and-biological-characteristics-of-diatom-species-for-potential-biodiesel-production-2090-4541-1000201.php?aid=66659)
961 [4541-1000201.php?aid=66659](http://www.omicsonline.com/open-access/variations-in-the-total-lipid-content-and-biological-characteristics-of-diatom-species-for-potential-biodiesel-production-2090-4541-1000201.php?aid=66659) (accessed February 9, 2016).
- 962 [50] T. Jauffrais, S. Drouet, V. Turpin, V. Méléder, B. Jesus, B. Cognie, P. Raimbault, R.P.
963 Cosson, P. Decottignies, V. Martin-Jézéquel, Growth and biochemical composition of a
964 microphytobenthic diatom (*Entomoneis paludosa*) exposed to shorebird (*Calidris alpina*)
965 droppings, *J. Exp. Mar. Biol. Ecol.* 469 (2015) 83–92.
- 966 [51] T. Jauffrais, B. Jesus, V. Méléder, V. Turpin, A.D.P.G. Russo, P. Raimbault, V.M.
967 Jézéquel, Physiological and photophysiological responses of the benthic diatom
968 *Entomoneis paludosa* (Bacillariophyceae) to dissolved inorganic and organic nitrogen in
969 culture, *Mar. Biol.* 163 (2016). doi:10.1007/s00227-016-2888-9.
- 970 [52] R.M. Knuckey, M.R. Brown, S.M. Barrett, G.M. Hallegraeff, Isolation of new
971 nanoplanktonic diatom strains and their evaluation as diets for juvenile Pacific oysters
972 (*Crassostrea gigas*), *Aquaculture*. 211 (2002) 253–274.
973 doi:[http://dx.doi.org/10.1016/S0044-8486\(02\)00010-8](http://dx.doi.org/10.1016/S0044-8486(02)00010-8).
- 974 [53] A.T. Soares, B.F. Silva, L.L. Fialho, M.A.G. Pequeno, A.A.H. Vieira, A.G. Souza, N.R.
975 Antoniosi Filho, Chromatographic characterization of triacylglycerides and fatty acid
976 methyl esters in microalgae oils for biodiesel production, *J. Renew. Sustain. Energy*. 5
977 (2013) 053111. doi:10.1063/1.4821289.
- 978 [54] T. Viriyayingsiri, P. Sittplangkoon, S. Powtongsook, K. Nootong, Continuous
979 production of diatom *Entomoneis* sp. in mechanically stirred tank and flat-panel airlift
980 photobioreactors, *Prep. Biochem. Biotechnol.* 46 (2016) 740–746.
981 doi:10.1080/10826068.2015.1135460.
- 982 [55] F.J. Fields, J.P. Kocielek, An evolutionary perspective on selecting high-lipid-content
983 diatoms (Bacillariophyta), *J. Appl. Phycol.* 27 (2015) 2209–2220. doi:10.1007/s10811-
984 014-0505-1.
- 985 [56] G.E. Fogg, D.M. Collyer, The accumulation of fats as a characteristic of certain classes
986 of algae, *Proc 8 Int Bot Congr.* 17 (1954) 28.
- 987 [57] P.G. Roessler, ENVIRONMENTAL CONTROL OF GLYCEROLIPID
988 METABOLISM IN MICROALGAE: COMMERCIAL IMPLICATIONS AND
989 FUTURE RESEARCH DIRECTIONS, *J. Phycol.* 26 (1990) 393–399.
990 doi:10.1111/j.0022-3646.1990.00393.x.
- 991 [58] B. Scholz, G. Liebezeit, Biochemical characterisation and fatty acid profiles of 25
992 benthic marine diatoms isolated from the Solthörn tidal flat (southern North Sea), *J.*
993 *Appl. Phycol.* 25 (2013) 453–465. doi:10.1007/s10811-012-9879-0.
- 994 [59] M. Islam, M. Magnusson, R. Brown, G. Ayoko, M. Nabi, K. Heimann, Microalgal
995 Species Selection for Biodiesel Production Based on Fuel Properties Derived from Fatty
996 Acid Profiles, *Energies*. 6 (2013) 5676–5702. doi:10.3390/en6115676.
- 997 [60] S.P. Slocombe, Q. Zhang, M. Ross, A. Anderson, N.J. Thomas, Á. Lapresa, C. Rad-
998 Menéndez, C.N. Campbell, K.D. Black, M.S. Stanley, Unlocking nature's treasure-
999 chest: screening for oleaginous algae, *Sci. Rep.* 5 (2015) 9844.
- 1000 [61] H. Chtourou, I. Dahmen, A. Jebali, F. Karray, I. Hassairi, S. Abdelkafi, H. Ayadi, S.
1001 Sayadi, A. Dhouib, Characterization of *Amphora* sp., a newly isolated diatom wild
1002 strain, potentially usable for biodiesel production, *Bioprocess Biosyst. Eng.* 38 (2015)
1003 1381–1392. doi:10.1007/s00449-015-1379-6.

- [62] M. De la Pena, Cell growth and nutritive value of the tropical benthic diatom, *Amphora* sp., at varying levels of nutrients and light intensity, and different culture locations, *J. Appl. Phycol.* 19 (2007) 647–655.
- [63] M. Giordano, M. Kansiz, P. Heraud, J. Beardall, B. Wood, D. McNaughton, Fourier transform infrared spectroscopy as a novel tool to investigate changes in intracellular macromolecular pools in the marine microalga *Chaetoceros muellerii* (Bacillariophyceae), *J. Phycol.* 37 (2001) 271–279.
- [64] H. Wagner, S. Dunker, Z. Liu, C. Wilhelm, Subcommunity FTIR-spectroscopy to determine physiological cell states, *Curr. Opin. Biotechnol.* 24 (2013) 88–94. doi:<https://doi.org/10.1016/j.copbio.2012.09.008>.
- [65] L.E. Brand, R.R. Guillard, L.S. Murphy, A method for the rapid and precise determination of acclimated phytoplankton reproduction rates, *J. Plankton Res.* 3 (1981) 193–201. <http://plankt.oxfordjournals.org/content/3/2/193.short> (accessed April 5, 2016).
- [66] M. Gilstad, E. Sakshaug, Growth rates of ten diatom species from the Barents Sea at different irradiances and day lengths., *Mar. Ecol. Prog. Ser. Oldendorf.* 64 (1990) 169–173.
- [67] C. Breil, M. Abert Vian, T. Zemb, W. Kunz, F. Chemat, “Bligh and Dyer” and Folch Methods for Solid–Liquid–Liquid Extraction of Lipids from Microorganisms. Comprehension of Solvation Mechanisms and towards Substitution with Alternative Solvents, *Int. J. Mol. Sci.* 18 (2017). doi:10.3390/ijms18040708.
- [68] G.-D. Feng, F. Zhang, L.-H. Cheng, X.-H. Xu, L. Zhang, H.-L. Chen, Evaluation of FT-IR and Nile Red methods for microalgal lipid characterization and biomass composition determination, *Bioresour. Technol.* 128 (2013) 107–112.
- [69] K. Stehfest, J. Toepel, C. Wilhelm, The application of micro-FTIR spectroscopy to analyze nutrient stress-related changes in biomass composition of phytoplankton algae, *Plant Physiol. Biochem.* 43 (2005) 717–726. doi:10.1016/j.plaphy.2005.07.001.
- [70] A.P. Dean, D.C. Sigee, B. Estrada, J.K. Pittman, Using FTIR spectroscopy for rapid determination of lipid accumulation in response to nitrogen limitation in freshwater microalgae, *Bioresour. Technol.* 101 (2010) 4499–4507. doi:10.1016/j.biortech.2010.01.065.
- [71] M. Mubarak, A. Shaija, T.V. Suchithra, A review on the extraction of lipid from microalgae for biodiesel production, *Algal Res.* 7 (2015) 117–123. doi:10.1016/j.algal.2014.10.008.
- [72] Y. Li, F.G. Naghdi, S. Garg, T.C. Adarme-Vega, K.J. Thurecht, W.A. Ghafor, S. Tannock, P.M. Schenk, A comparative study: the impact of different lipid extraction methods on current microalgal lipid research, *Microb. Cell Factories.* 13 (2014) 14.
- [73] S.J. Lee, B.-D. Yoon, H.-M. Oh, Rapid method for the determination of lipid from the green alga *Botryococcus braunii*, *Biotechnol. Tech.* 12 (1998) 553–556.
- [74] D. Xu, Z. Gao, F. Li, X. Fan, X. Zhang, N. Ye, S. Mou, C. Liang, D. Li, Detection and quantitation of lipid in the microalga *Tetraselmis subcordiformis* (Wille) Butcher with BODIPY 505/515 staining, *Bioresour. Technol.* 127 (2013) 386–390. doi:10.1016/j.biortech.2012.09.068.
- [75] H. De la Hoz Siegler, W. Ayidzoe, A. Ben-Zvi, R. Burrell, W. McCaffrey, Improving the reliability of fluorescence-based neutral lipid content measurements in microalgal cultures, *Algal Res.* 1 (2012) 176–184.
- [76] T. Mutanda, D. Ramesh, S. Karthikeyan, S. Kumari, A. Anandraj, F. Bux, Bioprospecting for hyper-lipid producing microalgal strains for sustainable biofuel production, *Bioresour. Technol.* 102 (2011) 57–70. doi:10.1016/j.biortech.2010.06.077.

- [77] W. Chen, C. Zhang, L. Song, M. Sommerfeld, Q. Hu, A high throughput Nile red method for quantitative measurement of neutral lipids in microalgae, *J. Microbiol. Methods*. 77 (2009) 41–47. doi:10.1016/j.mimet.2009.01.001.
- [78] L.M. Laurens, E.J. Wolfrum, Rapid compositional analysis of microalgae by NIR spectroscopy, *NIR News*. 23 (2012) 9–11.
- [79] M.S. Cooper, W.R. Hardin, T.W. Petersen, R.A. Cattolico, Visualizing “green oil” in live algal cells, *J. Biosci. Bioeng.* 109 (2010) 198–201. doi:10.1016/j.jbiosc.2009.08.004.
- [80] J.N. Murdock, D.L. Wetzell, FT-IR Microspectroscopy Enhances Biological and Ecological Analysis of Algae, *Appl. Spectrosc. Rev.* 44 (2009) 335–361. doi:10.1080/05704920902907440.
- [81] I. Dreissig, S. Machill, R. Salzer, C. Krafft, Quantification of brain lipids by FTIR spectroscopy and partial least squares regression, *Spectrochim. Acta. A. Mol. Biomol. Spectrosc.* 71 (2009) 2069–2075. doi:10.1016/j.saa.2008.08.008.
- [82] S.M. Renaud, H.C. Zhou, D.L. Parry, L.-V. Thinh, K.C. Woo, Effect of temperature on the growth, total lipid content and fatty acid composition of recently isolated tropical microalgae *Isochrysis* sp., *Nitzschia closterium*, *Nitzschia paleacea*, and commercial species *Isochrysis* sp.(clone T. ISO), *J. Appl. Phycol.* 7 (1995) 595–602. <http://link.springer.com/article/10.1007/BF00003948> (accessed August 25, 2016).
- [83] M.A. Borowitzka, L.J. Borowitzka, *Micro-algal biotechnology*, Cambridge University Press, 1988.
- [84] H.A. Spoehr, H.W. Milner, THE CHEMICAL COMPOSITION OF CHLORELLA; EFFECT OF ENVIRONMENTAL CONDITIONS, *Plant Physiol.* 24 (1949) 120–149. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC437916/>.
- [85] I.A. Guschina, J.L. Harwood, Lipids and lipid metabolism in eukaryotic algae, *Prog. Lipid Res.* 45 (2006) 160–186. doi:10.1016/j.plipres.2006.01.001.
- [86] L. Brand, R. Guillard, The effects of continuous light and light intensity on the reproduction rates of twenty-two species of marine phytoplankton, *J. Exp. Mar. Biol. Ecol.* 50 (1981) 119–132.
- [87] P.J. le B. Williams, L.M. Laurens, Microalgae as biodiesel & biomass feedstocks: review & analysis of the biochemistry, energetics & economics, *Energy Environ. Sci.* 3 (2010) 554–590.
- [88] S. Wahidin, A. Idris, S.R.M. Shaleh, The influence of light intensity and photoperiod on the growth and lipid content of microalgae *Nannochloropsis* sp., *Bioresour. Technol.* 129 (2013) 7–11. doi:10/f4mb36.
- [89] L. Rodolfi, G. Chini Zittelli, N. Bassi, G. Padovani, N. Biondi, G. Bonini, M.R. Tredici, Microalgae for oil: Strain selection, induction of lipid synthesis and outdoor mass cultivation in a low-cost photobioreactor, *Biotechnol. Bioeng.* 102 (2009) 100–112. doi:10.1002/bit.22033.
- [90] E.S. Shuba, D. Kifle, Microalgae to biofuels: ‘Promising’ alternative and renewable energy, review, *Renew. Sustain. Energy Rev.* 81 (2018) 743–755. doi:10.1016/j.rser.2017.08.042.
- [91] M.P. de Souza, M. Hoeltz, P.D. Gressler, L.B. Benitez, R.C.S. Schneider, Potential of Microalgal Bioproducts: General Perspectives and Main Challenges, *Waste Biomass Valorization*. (2018). doi:10.1007/s12649-018-0253-6.
- [92] M.-H. Liang, J. Zhu, J.-G. Jiang, High-value bioproducts from microalgae: strategies and progress, *Crit. Rev. Food Sci. Nutr.* 0 (2018) 01–53. doi:10.1080/10408398.2018.1455030.

7 Figure legends

Figure. 1 Example of growth curves measured for *Amphora* sp 1 NCC260 by (A) cell count, (B) fluorometry PAM (F0) and (C) radiometry (NDVI). Points corresponding to cell concentration according to time. Line curves corresponding to the Gompertz model fitted to cell concentration as a function of time. $n=3$, vertical bar = SD. The arrows indicate the late exponential day estimated by the Gompertz model.

Figure. 2 Values of the parameters measured for the screened strains. This includes strain productivity (A), lipid content as measured from the gravimetric method (B), the lipid ratio as measured semi-quantitatively by the FTIR approaches, $[(eb+CH_3+CH_3)/si]$ (C) or $[area\ eb+CH_2+CH_3]/Total\ area]$ multiplied by 100 for scaling purposes (D). The a, b, c notation corresponded to the maximum and minimum values for groups 1, 2 and 3. $N=3$, independent measurements, $\pm$ SD.

Figure. 3 Example of averaged FTIR spectra recorded on entire cells or on the corresponding lipid extract. (A) *Staurosira* sp NCC182 FTIR signature recorded on the entire cells and (B) *Staurosira* sp NCC182 FTIR signature recorded on a crude lipid extract. The grey area corresponds to the variation of the FTIR signal associated to the standard deviation for $n=3$ independent measurements.

Figure. 4 Correspondence analysis map calculated on the basis of the macromolecular content as evaluated by FTIR on all the assayed strains of the NCC. $N=3$ independent measurements.

Figure 5 Venn diagram showing the degree of overlap among the different approaches used to identify the lipid rich diatoms. In the gravimetric method circle, 14 strains were identified as rich in lipids: 4 only with this method (NCC260, NCC208, NCC253, NCC351), one strains were also identified as rich and lipid by the FTIR on lipid extract (NCC229), and two strains were also identified as rich in lipids by the FTIR on entire cells (NCC33, NCC199). One strain was identified by FTIR on lipid extract and FTIR on whole cells (NCC366) and seven strains were identified as rich in lipids by all three methods (NCC109, NCC182, NCC113, NCC270, NCC169, NCC303, NCC216).

Figure 6. Boxplots summarizing the sample distribution criteria as measured with FTIR methods, gravimetry for the lipid rate and with fluorimetry for productivity. FTIR data was expressed in arbitrary units. Lipid rates in %DW and Px in $\text{g.L}^{-1}.\text{day}^{-1}$. FTIR results was multiplied by 20 for scaling purposes. The Px was multiplied by 100 for scaling purposes. 33 strains were assayed in independent biological triplicates.

Figures

Figure 1.

Fig. 3

Fig. 4

1195 Fig. 5

1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208

Fig. 6

1234

1235 Table S1. Supplementary data. Strains who did not grown.

1236

1237

1238

1239

1240

1241

1242

1243

Species	NCC strain identification	Sampling location
<i>Amphora</i> sp. B2	NCC261	France, NW Atlantic coast
<i>Amphora</i> sp. AC16	NCC410	France, NW Atlantic coast
<i>Amphora</i> sp. AE8	NCC413	France, NW Atlantic coast
<i>Berkeleya rutilans</i>	NCC210.2	France, NW Atlantic coast
<i>Berkeleya rutilans</i>	NCC309	France, Mediterranean sea
<i>Brockmaniella brockmanii</i> 2	NCC403	France, NW Atlantic coast
<i>Caloneis</i> sp. 1	NCC180	France, NW Atlantic coast
<i>Catacombas</i> sp. 1	NCC337	France, Mediterranean sea
<i>Cocconeis scutellum</i> 1	NCC209.1	France, NW Atlantic coast
<i>Cocconeis scutellum</i> 2	NCC209.2	France, NW Atlantic coast
<i>Cocconeis scutellum</i> 2	NCC209.3	France, NW Atlantic coast
<i>Craspedostauros</i> sp. 3	NCC57	France, NW Atlantic coast
<i>Craspedostauros</i> sp. 4	NCC58	France, NW Atlantic coast
<i>Craspedostauros</i> sp. 5	NCC204	France, NW Atlantic coast
<i>Entomoneis alata</i> 1	NCC16	France, NW Atlantic coast
<i>Entomoneis alata</i> 2	NCC448	Portugal, NW Atlantic coast
<i>Entomoneis</i> sp. BAB2	NCC415	France, NW Atlantic coast
<i>Gyrosigma</i> sp. 1	NCC411	France, NW Atlantic coast
<i>Gyrosigma</i> sp. 2	NCC412	France, NW Atlantic coast
<i>Gyrosigma tenuissimum</i>	NCC258	France, NW Atlantic coast
<i>Halamphora coffeaformis</i>	UTCC58	Canada, NW Atlantic coast
<i>Helicotheca tamesis</i> 1	NCC59	, France, Mediterranean sea
<i>Helicotheca tamesis</i> 2	NCC60	France, NW Atlantic coast
<i>Lampriscus</i> sp.	NCC347	France, Mediterranean sea
<i>Leptocylindrus danicus</i> 1	NCC205	France, NW Atlantic coast
<i>Leptocylindrus danicus</i> 2	NCC206	France, NW Atlantic coast
<i>Melosira nummuloides</i> 1	NCC25	France, NW Atlantic coast
<i>Melosira nummuloides</i> 2	NCC25.1	France, NW Atlantic coast
<i>Navicula</i> sp. Z4	NCC224	France, NW Atlantic coast
<i>Navicula</i> sp. e1	NCC269	France, NW Atlantic coast
<i>Navicula cf ramosissima</i>	NCC449	France, NW Atlantic coast
<i>Nitzschia laevis</i>	NCC39	France, NW Atlantic coast
<i>Nitzschia salinicola</i>	NCC41	France, NW Atlantic coast
<i>Nitzschia</i> sp. B4	NCC114	France, NW Atlantic coast
<i>Opehora</i> sp. 2	NCC365	France, NW Atlantic coast
<i>Paralia sulcata</i>	NCC177	France, NW Atlantic coast
<i>Pleurosigma</i> sp. K	NCC339	Ukraine, Black sea
<i>Pleurosigma</i> sp. LM	NCC404	France, NW Atlantic coast
<i>Pleurosigma</i> sp. BC1	NCC423	France, NW Atlantic coast
<i>Pleurosigma</i> sp. BC7	NCC425	France, NW Atlantic coast
<i>Pleurosigma</i> sp. BC15	NCC428	France, NW Atlantic coast
<i>Rhizosolenia setigera</i>	NCC127	France, NW Atlantic coast
<i>Tabularia tabulata</i>	NCC338	France, NW Atlantic coast

Table S2: Lipid quantification evaluated by FTIR and ATR method for all the species.

Species	code	FTIR					ATR	
		Carb/si	Prot/si	Eb/si	CH2/si	CH3/si	eb	CH2 + CH3
<i>Amphora acutiuscula</i>	NCC216	53.6 ± 8.2	45.8 ± 5.7	22.1 ± 4.9	65.7 ± 8.0	53.5 ± 7.8	6.2 ± 0.9	24.3 ± 2.4
<i>Amphora</i> sp. 1	NCC260	63.6 ± 2.9	40.0 ± 2.0	12.2 ± 2.3	57.3 ± 3.3	39.2 ± 0.6	4.5 ± 0.7	18.1 ± 3.1
<i>Amphora</i> sp. 2	NCC169	58.1 ± 4.3	42.4 ± 1.1	17.2 ± 3.4	59.7 ± 3.4	49.7 ± 3.0	7.1 ± 0.7	42.1 ± 11.0
<i>Brockmaniella brockmanii</i>	NCC161	51.2 ± 5.5	25.2 ± 6.4	12.8 ± 4.0	57.7 ± 5.5	41.8 ± 5.5	4.1 ± 1.6	21.1 ± 5.6
<i>Conticriba weissflogii</i>	NCC133	47.4 ± 0.5	40.2 ± 0.8	5.03 ± 1.5	48.4 ± 2.6	41.7 ± 1.9	4.7 ± 0.3	20.0 ± 0.7
<i>Conticriba weissflogii</i>	CCMP1336	48.3 ± 2.8	43.5 ± 3.3	9.8 ± 1.8	48.7 ± 3.0	40.9 ± 2.7	5.0 ± 0.7	16.9 ± 2.2
<i>Craspedostauros britannicus</i>	NCC195	60.7 ± 2.4	69.8 ± 11	23.2 ± 3.3	80.7 ± 8.0	66.0 ± 6.9	4.8 ± 0.4	19.2 ± 1.8
<i>Craspedostauros britannicus</i>	NCC199	55.1 ± 3.4	47.5 ± 5.7	26.9 ± 3.4	62.2 ± 2.7	44.2 ± 3.3	5.4 ± 0.6	20.5 ± 1.9
<i>Craspedostauros</i> sp. 1	NCC228	44.3 ± 0.1	37.8 ± 4.3	6.2 ± 0.4	46.3 ± 6.4	37.8 ± 6.0	3.9 ± 0.8	15.2 ± 3.4
<i>Craspedostauros</i> sp. 2	NCC218	54.6 ± 2.6	49.4 ± 3.5	7.5 ± 0.3	46.0 ± 3.8	39.0 ± 4.0	4.7 ± 0.3	17.9 ± 2.5
<i>Cymatosira belgica</i>	NCC208	41.6 ± 4.8	25.4 ± 3.3	6.7 ± 1.1	55.4 ± 5.1	41.5 ± 3.7	5.9 ± 0.7	22.7 ± 3.2
<i>Entomoneis paludosa</i>	NCC18.1.1	39.6 ± 0.3	26.0 ± 2.7	2.9 ± 0.1	27.3 ± 2.4	19.4 ± 1.8	3.9 ± 0.7	23.1 ± 1.8
<i>Entomoneis paludosa</i>	NCC18.2.1	45.7 ± 0.6	40.9 ± 1.4	6.1 ± 0.1	34.5 ± 0.9	22.5 ± 0.6	3.1 ± 0.4	21.6 ± 1.1
<i>Entomoneis</i> sp. 1	NCC350	39.9 ± 0.1	30.4 ± 0.8	5.5 ± 1.5	28.6 ± 0.9	21.2 ± 1.7	2.7 ± 0.1	30.6 ± 4.3
<i>Entomoneis</i> sp. 2	NCC20	42.5 ± 0.5	32.5 ± 0.3	9.5 ± 0.3	40.0 ± 0.4	26.0 ± 0.6	4.7 ± 0.3	18.6 ± 0.4
<i>Entomoneis</i> sp. 3	NCC351	45.5 ± 2.9	41.4 ± 4.2	14.4 ± 3.4	37.2 ± 0.8	28.3 ± 0.6	3.9 ± 1.7	15.8 ± 5.6
<i>Entomoneis</i> sp. 4	NCC301	62.5 ± 4.8	56.7 ± 6.4	12.5 ± 2.5	64.0 ± 7.9	46.7 ± 5.6	2.9 ± 0.1	17.0 ± 0.8
<i>Entomoneis</i> sp. 5	NCC302	37.6 ± 0.7	36.8 ± 0.8	3.43 ± 0.1	33.9 ± 0.7	23.9 ± 0.5	3.3 ± 0.3	19.6 ± 1.5
<i>Entomoneis</i> sp. 6	NCC335	50.5 ± 0.6	35.3 ± 2.7	6.6 ± 0.7	29.9 ± 1.9	19.0 ± 1.1	3.1 ± 0.2	13.4 ± 3.0
<i>Entomoneis</i> sp. 7	NCC445	40.6 ± 0.9	27.0 ± 2.7	8.5 ± 0.2	37.7 ± 0.4	29.2 ± 0.3	4.1 ± 0.9	17.5 ± 4.3
<i>Extubocellulus cf cribriger</i>	NCC229	49.8 ± 3.7	38.1 ± 2.6	10.1 ± 1.2	48.0 ± 1.9	38.0 ± 1.1	4.9 ± 0.3	34.2 ± 1.7
<i>Fallacia</i> sp. 1	NCC303	64.7 ± 1.8	74.4 ± 7.0	14.2 ± 2.5	64.5 ± 4.1	55.5 ± 5.4	6.9 ± 0.3	26.4 ± 2.9
<i>Fallacia</i> sp. 2	NCC304	46.0 ± 6.2	45.7 ± 3.2	31.3 ± 4.0	72.8 ± 2.7	53.3 ± 2.1	5.1 ± 1.0	18.9 ± 3.4
<i>Licmophora</i> sp. 1	NCC253	53.1 ± 2.7	65 ± 10	4.4 ± 1.1	57.9 ± 5.4	50.3 ± 5.2	3.6 ± 0.3	22.6 ± 0.6
<i>Lithodesmium</i> sp	NCC353	34.2 ± 2.0	25.8 ± 0.8	2.4 ± 0.3	31.1 ± 1.1	23.1 ± 0.4	2.1 ± 0.3	28.0 ± 3.2
<i>Navicula</i> sp. 1	NCC113	63.3 ± 4.7	43.8 ± 8.9	33.4 ± 4.7	68.2 ± 9.6	52.4 ± 5.6	3.4 ± 0.4	28.8 ± 1.2
<i>Navicula</i> sp. 2	NCC226	54.2 ± 4.3	27.0 ± 3.0	5.2 ± 0.9	53.1 ± 3.8	39.5 ± 2.8	4.2 ± 0.7	22.4 ± 0.7
<i>Nitzschia</i> sp. 5	NCC109	62.0 ± 2.8	42.4 ± 6.7	22.0 ± 2.3	72.7 ± 4.6	61.3 ± 4.8	6.9 ± 0.4	45.1 ± 28.9
<i>Nitzschia alexandrina</i>	NCC33	58.2 ± 7.2	44.0 ± 4.6	20.7 ± 3.3	56.1 ± 4.0	45.8 ± 0.7	5.7 ± 0.3	21.3 ± 2.3
<i>Opephora</i> sp. 1	NCC366	58 ± 12	40.1 ± 8.8	31.6 ± 6.1	50.6 ± 4.5	42.0 ± 2.5	6.9 ± 1.1	26.3 ± 4.4
<i>Pseudonitzschia americana</i>	PNA06 KER	56.6 ± 1.4	60.6 ± 3.9	3.9 ± 1.0	44.0 ± 2.8	39.1 ± 3.2	4.0 ± 0.3	38 ± 28
<i>Staurosira</i> sp.	NCC182	47.7 ± 6.6	30.7 ± 4.1	30.9 ± 5.4	61.6 ± 4.0	49.0 ± 2.3	7.3 ± 0.1	31.6 ± 0.6
<i>Surirella</i> sp. 1	NCC270	58.5 ± 2.4	54.4 ± 8.4	22.0 ± 4.6	56.1 ± 5.3	51.3 ± 5.4	7.1 ± 0.4	27.1 ± 0.6