



HAL
open science

On the influence of gravity on density-dependent incompressible periodic fluids

Van-Sang Ngo, Stefano Scrobogna

► **To cite this version:**

Van-Sang Ngo, Stefano Scrobogna. On the influence of gravity on density-dependent incompressible periodic fluids. *Journal of Differential Equations*, 2019, 267 (2), pp.1510-1559. 10.1016/j.jde.2019.02.011 . hal-02334662

HAL Id: hal-02334662

<https://hal.science/hal-02334662>

Submitted on 22 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

On the influence of gravity on density-dependent incompressible periodic fluids

Van-Sang Ngo^{a,*}, Stefano Scrobogna^b

^a *Université de Rouen Normandie, Laboratoire de Mathématiques Raphaël Salem, UMR 6085 CNRS, 76801 Saint-Etienne du Rouvray, France*

^b *Basque Center for Applied Mathematics, Mazarredo, 14, E48009 Bilbao, Basque Country, Spain*

Abstract

The present work is devoted to the analysis of density-dependent, incompressible fluids in a 3D torus, when the Froude number ε goes to zero. We consider the very general case where the initial data do not have a zero horizontal average, where we only have smoothing effect on the velocity but not on the density and where we can have resonant phenomena on the domain. We explicitly determine the limit system when $\varepsilon \rightarrow 0$ and prove its global wellposedness. Finally, we prove that for large initial data, the density-dependent, incompressible fluid system is globally wellposed, provided that ε is small enough.

Keywords: Incompressible fluids, stratified fluids, parabolic systems, bootstrap

2010 MSC: 35Q30, 35Q35, 37N10, 76D03, 76D50, 76M45, 76M55

1. Introduction

In this paper, we want to study the motion of an incompressible, inhomogeneous fluid whose density profile is considered to be a perturbation around a stable state, which is described by the following system

$$(PBS_\varepsilon) \quad \begin{cases} \partial_t v^\varepsilon + v^\varepsilon \cdot \nabla v^\varepsilon - \nu \Delta v^\varepsilon - \frac{1}{\varepsilon} \rho^\varepsilon \vec{e}_3 = -\frac{1}{\varepsilon} \nabla \Phi^\varepsilon, \\ \partial_t \rho^\varepsilon + v^\varepsilon \cdot \nabla \rho^\varepsilon + \frac{1}{\varepsilon} v^{3,\varepsilon} = 0, \\ \operatorname{div} v^\varepsilon = 0, \\ (v^\varepsilon, \rho^\varepsilon)|_{t=0} = (v_0, \rho_0), \end{cases}$$

*Corresponding author

Email addresses: van-sang.ngo@univ-rouen.fr (Van-Sang Ngo), sscrobogna@bcamath.org (Stefano Scrobogna)

in the regime where the Froude number $\varepsilon \rightarrow 0$. Here, the vector field v^ε and the scalar function ρ^ε represent respectively the velocity and the density of the fluid and ν stands for the viscosity. For a more detailed discussion about the physical motivation and the derivation of the model, we refer to [40], [39], [41]. We also refer to the monographs [17] and [36] for a much wider survey on geophysical models.

Let us give some brief comments about the system (PBS_ε) . In a nutshell, there are two forces which constrain the motion of a fluid on a geophysical scale: the Coriolis force and the gravitational stratification. The predominant influence of one force, the other, or both gives rise to substantially different dynamics.

The *Coriolis force* (see [43] for a detailed analysis of such force) is due to the rotation of the Earth around its axis and acts perpendicularly to the motion of the fluid. If the magnitude of the force is sufficiently large (when the rotation is fast or the scale is large for example), the Coriolis force “penalizes” the vertical dynamics of the fluid and makes it move in rigid columns (the so-called *Taylor columns*). This tendency of a rotating fluid to displace in vertical homogeneous columns is generally known as *Taylor-Proudman theorem*, which was first derived by Sidney Samuel Hough (1870-1923), a mathematician at Cambridge in the work [29], but it was named after the works of G.I. Taylor [46] and Joseph Proudman [37]. On a mathematical point of view, the Taylor-Proudman effect for homogeneous, fast rotating fluids is a rather well understood after the works [3], [4], [15], [23] and [28]. In such setting, we mention as well the work [27] in which the authors consider an inhomogeneous rotation, the very recent work [20] in which fast rotation for nonhomogeneous fluids was considered and the works [21], [22] and [34] in which fast rotation was considered simultaneously with weak compressibility.

Beside the rotation, one can consider a fluid which is inhomogeneous and whose density profile is a linearization around a stable state, we refer to [7], [17] and references therein for a thorough derivation of the model. In such situation, we can imagine that the rotation effects and stratification effects are equally relevant: the system describing such effect is known as *primitive equations* (see [7] and [17]). The primitive equations and their asymptotic dynamic as stratification and rotation tend to infinity at a comparable rate are as well rather well understood on a mathematical viewpoint: we refer to the works [8], [9], [10], [11], [12], [14], [23], [42] and references therein.

Now, we will briefly discuss the *gravitational stratification* effect, which is the main physical phenomenon concerning the system (PBS_ε) for an inhomogeneous fluid, subjected to a gravitational force pointing downwards. Gravity force tends to lower the regions of the fluid with higher density and raise the regions with lower density, trying finally to dispose the fluid in horizontal stacks of vertically decreasing density. A fluid in such configuration (density profile which is a decreasing function of the variable x_3 only) is said to be in a *configuration of equilibrium*.

Let hence consider a fluid in a configuration of equilibrium and let us imagine to raise a small parcel of the fluid with high density in a region of low density. Since such parcel is much heavier (in average) than the fluid surrounding it, the gravity force will induce a downwards motion. Such motion does not stop until the parcel reaches a layer whose density is comparable to its own, and inertially it will continue to move downwards until sufficient buoyancy is provided to invert the motion, due to Archimedes principle. This kind of perturbation of an equilibrium state induces hence a pulsating motion which is described by the linear application

$$(1.1) \quad (u^{1,\varepsilon}, u^{2,\varepsilon}, u^{3,\varepsilon}, \rho^\varepsilon) \mapsto \frac{1}{\varepsilon} (0, 0, -\rho^\varepsilon, u^{3,\varepsilon}),$$

which appears in (PBS_ε) . The application (1.1) is called *stratification buoyancy* and we will base our analysis on the dispersive effects induced by such perturbation.

To the best of our knowledge, there are not many results concerning the effects of the stratification buoyancy. In [19], there was a first attempt to perform a multiscale analysis when Rossby and Froude number are in different regimes, while in [39] and [47], the authors studied the convergence and stability of solutions of (PBS_ε) when the Froude number $\varepsilon \rightarrow 0$ in the whole space \mathbb{R}^3 . In [40] the system (PBS_ε) is studied in nonresonant domains when the initial data has zero horizontal average. We mention as well the very recent works [31, 32, 33, 44, 45].

In this paper, the unknowns $(v^\varepsilon, \rho^\varepsilon)$ are considered to be functions in the variables $(x, t) \in \mathbb{T}^3 \times \mathbb{R}_+$ being the space domain \mathbb{T}^3 the three-dimensional periodic box

$$\mathbb{T}^3 = \prod_{i=1}^3 \mathbb{R} / a_i \mathbb{Z}, \quad a_i \in \mathbb{R}.$$

Compared to [40], we consider the much more general case with the following additional difficulties

1. Initial data are considered with generic horizontal average. This point seem marginal, but as showed in this paper, the dynamics induced by initial data with nonzero horizontal average create additional vertical gravitational perturbations, the control of which is highly non-trivial (see as well [26]).
2. Generic space domain may present resonant effects.
3. Density profiles are only transported and do not satisfy a transport-diffusion equation and so do not possess smoothing effects.

From now on we rewrite the system (PBS_ε) in the following more compact form

$$(\text{PBS}_\varepsilon) \quad \begin{cases} \partial_t V^\varepsilon + v^\varepsilon \cdot \nabla V^\varepsilon - \mathcal{A}_2(D) V^\varepsilon + \frac{1}{\varepsilon} \mathcal{A} V^\varepsilon = -\frac{1}{\varepsilon} \begin{pmatrix} \nabla \Phi^\varepsilon \\ 0 \end{pmatrix}, \\ V^\varepsilon = (v^\varepsilon, \rho^\varepsilon), \\ \operatorname{div} v^\varepsilon = 0, \\ V^\varepsilon|_{t=0} = V_0, \end{cases}$$

where

$$(1.2) \quad \mathcal{A} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix}, \quad \mathcal{A}_2(D) = \begin{pmatrix} \nu \Delta & 0 & 0 & 0 \\ 0 & \nu \Delta & 0 & 0 \\ 0 & 0 & \nu \Delta & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

The additional difficulties (1)–(3) listed above are the main difficulties in the present work and they modify significantly the dynamic of (PBS_ε) compared to the results proved in [40], as already mentioned. Let us hence start describing the effects induced by the hypothesis made in the point 1: we will see in the following that the dynamics of the solutions of (PBS_ε) in the limit regime $\varepsilon \rightarrow 0$ is essentially governed by the effects of the outer force $\varepsilon^{-1} \mathcal{A} V^\varepsilon$.

1.1. A survey on the adopted notation

All along this note we consider real valued vector fields, i.e. applications $V : \mathbb{R}_+ \times \mathbb{T}^3 \rightarrow \mathbb{R}^4$. We will often associate to a vector field V the vector field v which shall be simply the projection on the first three components of V . The vector fields considered are periodic in all their directions and they have zero global average $\int_{\mathbb{T}^3} V dx = 0$, which is equivalent to assume that the first Fourier coefficient $\widehat{V}(0) = 0$. We remark that the zero average propriety stated above is preserved in time t for both Navier-Stokes equations as well as for the system (PBS_ε) .

Let us define the Sobolev space $H^s(\mathbb{T}^3)$ as the set of all the tempered distributions f such that

$$(1.3) \quad \|f\|_{H^s(\mathbb{T}^3)} = \left(\sum_{n \in \mathbb{Z}^3} \left(1 + |\tilde{n}|^2\right)^s \left|\widehat{f}(n)\right|^2 \right)^{1/2} < \infty,$$

where $\widehat{f}(n)$ is the usual n -th Fourier mode of f and where we set

$$\tilde{n}_i = \frac{n_i}{a_i}, \quad |\tilde{n}|^2 = \sum_i \tilde{n}_i^2.$$

Since we always consider vector fields whose average is null, the Sobolev norm defined above is equivalent to the following semi-norm

$$\left\| (-\Delta)^{s/2} f \right\|_{L^2(\mathbb{T}^3)} \sim \|f\|_{H^s(\mathbb{T}^3)}, \quad s \in \mathbb{R},$$

which appears naturally in parabolic problems.

Throughout the paper, we also use \mathbb{P} to denote the three dimensional Leray operator which leaves untouched the fourth component, i.e.

$$(1.4) \quad \mathbb{P} = \left(\begin{array}{c|c} \mathbb{P}^{(3)} & 0 \\ \hline 0 & 1 \end{array} \right) = \left(\begin{array}{c|c} 1 - \Delta^{-1} \partial_i \partial_j & 0 \\ \hline 0 & 1 \end{array} \right)_{i,j=1,2,3},$$

where $\mathbb{P}^{(3)}$ is the usual Leray projection onto the subspace of divergence-free vector fields. The operator \mathbb{P} is then a pseudo-differential operator, in the Fourier space its symbol is

$$(1.5) \quad \widehat{\mathbb{P}}(n) = \left(\begin{array}{c|c} \delta_{i,j} - \frac{\tilde{n}_i \tilde{n}_j}{|\tilde{n}|^2} & 0 \\ \hline 0 & 1 \end{array} \right)_{i,j=1,2,3},$$

where $\delta_{i,j}$ is Kronecker's delta.

1.2. Local existence result

Being the operator \mathcal{A} skew-symmetric it is possible to apply energy methods to the system (PBS_ε) in the same fashion as it is done in [5, Chapter 4] for quasilinear symmetric hyperbolic systems. Being this the case we can deduce the following local existence result

Theorem 1.1. *Let $V_0 \in H^s(\mathbb{T}^3)$ where $s > 1 + 3/2$, there exist a $T^* > 0$ such that for every $T \in [0, T^*)$ the system (PBS_ε) admits a unique solution in the energy space*

$$\mathcal{C}([0, T]; H^s(\mathbb{T}^3)) \cap \mathcal{C}^1([0, T]; H^{s-1}(\mathbb{T}^3)).$$

Moreover there exist a positive constant c such that

$$T > \frac{c}{\|V_0\|_{H^s(\mathbb{T}^3)}},$$

and the maximal time of existence T^* is independent of ε and s and, if $T^* < \infty$, then

$$(1.6) \quad \int_0^{T^*} \|\nabla V^\varepsilon(t)\|_{L^\infty(\mathbb{T}^3)} dt = \infty.$$

1.3. Global existence result and limit dynamics

As in other singular perturbation problems (see [23], [24], [28], ... for instance), the difficulty to study the limit when $\varepsilon \rightarrow 0$ lies in the fact that $\partial_t V^\varepsilon$ is not uniformly bounded in ε , which is the result of the high oscillating free waves, described by the system

$$\begin{cases} \partial_\tau W + \mathbb{P}\mathcal{A}W = 0, \\ W|_{\tau=0} = W_0, \end{cases}$$

where the matrices \mathbb{P} and \mathcal{A} are defined respectively in (1.5) and (1.2). The standard procedure consists in filtering these oscillations. Let \mathcal{L} be the operator

$$(1.7) \quad \mathcal{L}(\tau) = e^{-\tau\mathbb{P}\mathcal{A}},$$

which send each initial datum W_0 to the solution of the above free wave system $W(t)$. Then, the filtered solution

$$U^\varepsilon = \mathcal{L}\left(-\frac{t}{\varepsilon}\right)V^\varepsilon,$$

is in fact solution of the filtered system

$$(S_\varepsilon) \quad \begin{cases} \partial_t U^\varepsilon + \mathcal{Q}^\varepsilon(U^\varepsilon, U^\varepsilon) - \mathcal{A}_2^\varepsilon(D)U^\varepsilon = 0, \\ U^\varepsilon|_{t=0} = V_0, \end{cases}$$

where

$$\begin{aligned} & \mathcal{Q}^\varepsilon(V_1, V_2) \\ &= \frac{1}{2}\mathcal{L}\left(\frac{t}{\varepsilon}\right)\mathbb{P}\left[\mathcal{L}\left(-\frac{t}{\varepsilon}\right)V_1 \cdot \nabla\mathcal{L}\left(-\frac{t}{\varepsilon}\right)V_2 + \mathcal{L}\left(-\frac{t}{\varepsilon}\right)V_2 \cdot \nabla\mathcal{L}\left(-\frac{t}{\varepsilon}\right)V_1\right] \end{aligned}$$

and

$$\mathcal{A}_2^\varepsilon(D)W = \mathcal{L}\left(\frac{t}{\varepsilon}\right)\mathcal{A}_2(D)\mathcal{L}\left(-\frac{t}{\varepsilon}\right)W.$$

The limit dynamics of (PBS_ε) will be described in Section 4, where we study in details the limit of \mathcal{Q}^ε and $\mathcal{A}_2^\varepsilon(D)$ as $\varepsilon \rightarrow 0$. For any vector field A , let $\underline{A} = \underline{A}(x_3)$ be the horizontal average of A , which is defined as

$$\underline{A}(x_3) = \frac{1}{4\pi^2 a_1 a_2} \int_{\mathbb{T}_h^2} A(y_h, x_3) dy_h.$$

Then, the horizontal oscillating part $\tilde{A} = \tilde{A}(x_h, x_3)$ of A (which is of zero horizontal average) will be

$$(1.8) \quad \tilde{A}(x_h, x_3) = A(x_h, x_3) - \underline{A}(x_3).$$

Now, as in [42], we will also decompose the horizontal oscillating part $\tilde{A} = \tilde{A}(x_h, x_3)$ into two parts: $\overline{A}(x)$ belongs to the kernel of $\mathbb{P}\mathcal{A}$ (we remark that \underline{A} also belongs to $\ker(\mathbb{P}\mathcal{A})$), and the other part is the 3D oscillating part $A_{\text{osc}}(x_h, x_3)$. To resume, we perform the following decomposition, for any vector field A ,

$$A = \underline{A}(x_3) + \overline{A}(x) + A_{\text{osc}}(x_h, x_3).$$

For any initial data V_0 of the system (PBS_ε) , we decompose the filtered counter part of V_0 as follows

$$U_0 = \mathcal{L}(0)V_0 = \underline{U}_0(x_3) + \overline{U}_0(x) + U_{0,\text{osc}}(x_h, x_3),$$

with

$$\begin{aligned}\underline{U}_0 &= \underline{V}_0 = \frac{1}{2\pi^2 a_1 a_2} \int_{\mathbb{T}_h^2} V_0(y_h, x_3) dy_h, \\ \bar{U}_0 &= (\bar{u}_0^h, 0, 0)^\top, \\ U_{0,\text{osc}} &= V_0 - \bar{U}_0 - \underline{U}_0,\end{aligned}$$

and with

$$\bar{u}_0^h = \begin{pmatrix} -\partial_2 \\ \partial_1 \end{pmatrix} (-\Delta_h)^{-1} (-\partial_2 V_0^1 + \partial_1 V_0^2).$$

Then, the main result of our paper is the following

Theorem 1.2. *Let V_0 be in $H^s(\mathbb{T}^3)$, $s > 9/2 (= \frac{d}{2} + 3)$, be the initial data of the system (PBS $_\varepsilon$). Then, there exists a $\varepsilon_0 > 0$ such that for any $\varepsilon \in (0, \varepsilon_0)$, the solution V^ε of (PBS $_\varepsilon$) globally exists in time and we have the following asymptotics*

$$V^\varepsilon = \underline{U} + \bar{U} + \mathcal{L}\left(\frac{t}{\varepsilon}\right) U_{\text{osc}} + o_\varepsilon(1)$$

in

$$\mathcal{C}(\mathbb{R}_+; H^{s-2}(\mathbb{T}^3)) \cap L_{\text{loc}}^\infty(\mathbb{R}_+; H^{s-2}(\mathbb{T}^3)).$$

Here, $\underline{U} = (\underline{u}^h, 0, \underline{U}^4) = (\underline{U}^1, \underline{U}^2, 0, \underline{U}^4)$, $\bar{U} = (\bar{u}^h, 0, 0) = (\bar{u}^1, \bar{u}^2, 0, 0)$ and U_{osc} respectively solve the following systems

$$(1.9) \quad \begin{cases} \partial_t \underline{u}^h(x_3, t) - \nu \partial_3^2 \underline{u}^h(x_3, t) = 0, \\ \underline{U}^4(x_3, t) = \underline{U}_0^4, \\ \underline{u}^h|_{t=0} = \underline{U}_0^h, \\ \underline{U}^4|_{t=0} = \underline{U}_0^4, \end{cases}$$

$$(1.10) \quad \begin{cases} \partial_t \bar{u}^h(t, x_h, x_3) + \bar{u}^h(t, x_h, x_3) \cdot \nabla_h \bar{u}^h(t, x_h, x_3) \\ \quad + \underline{u}^h(t, x_3) \cdot \nabla_h \bar{u}^h(t, x_h, x_3) - \nu \Delta_h \bar{u}^h(t, x_h, x_3) = -\nabla_h \bar{p}(t, x_h, x_3) \\ \text{div}_h \bar{u}^h(x_h, x_3) = 0, \\ \bar{u}^h(t, x_h, x_3)|_{t=0} = \bar{u}_0^h(x_h, x_3). \end{cases}$$

$$(1.11) \quad \begin{cases} \partial_t U_{\text{osc}} + \tilde{\mathcal{Q}}_1(U_{\text{osc}} + 2\bar{U}, U_{\text{osc}}) + \mathcal{B}(\underline{U}, U_{\text{osc}}) - \nu \Delta U_{\text{osc}} = 0, \\ \text{div} U_{\text{osc}} = 0, \\ U_{\text{osc}}|_{t=0} = U_{0,\text{osc}}, \end{cases}$$

where the explicit expressions of \mathcal{B} and $\tilde{\mathcal{Q}}_1$ are given by the equations (5.13) and (4.1) and are omitted here for the sake of simplicity.

Remark 1.3. It is interesting to remark that, despite V^ε is the solution of the parabolic-hyperbolic system (PBS_ε) , it is well defined in the space $\mathcal{C}([0, T]; H^s)$ for $s > 9/2$ and $T > 0$. This improvement of regularity is known as propagation of parabolicity: such name is motivated by the fact that the limit equations (1.10) and (1.11) are strictly parabolic and we can prove they are globally well posed in some suitable energy space of subcritical regularity. A similar phenomenon takes place as well in the study of the incompressible limit for weakly compressible fluids, we refer the reader to the works [18] and [25].

Remark 1.4. Equation (1.11) is a nonlinear three-dimensional parabolic equation. The nonlinearity $\tilde{\mathcal{Q}}_1$ is a modified, symmetric transport form which assumes the following explicit form

$$\tilde{\mathcal{Q}}_1(A, B) = \frac{1}{2} \chi(D) (A \cdot \nabla B + B \cdot \nabla A),$$

where χ is a Fourier multiplier of order zero which localizes bilinear interactions on a very specific frequency set. Following the theory of the three-dimensional Navier-Stokes equations we do not expect hence (1.11) to be globally well-posed. Despite this, we shall see that the zero-order Fourier multiplier $\chi(D)$ has in fact a nontrivial smoothing effect, which makes the bilinear interaction $\tilde{\mathcal{Q}}_1(U_{\text{osc}}, U_{\text{osc}})$ smoother than $U_{\text{osc}} \cdot \nabla U_{\text{osc}}$, and then global well-posedness possible for (1.11).

1.4. Organization of the paper

The paper will be organized as follows. In the next section, we analyse the spectral properties of the perturbation operator $\mathbb{P}\mathcal{A}$. Section 3 consists in the study of the filtered system (S_ε) . Section 4 is devoted to the determination of the limit system and Section 5 to the detailed study of limits of the bilinear term \mathcal{Q}^ε and of the linear operator $\mathcal{A}_2^\varepsilon(D)$, as $\varepsilon \rightarrow 0$. In Section 6, we prove the global propagation of smoothness for the limit system and finally, in the last section, we prove the main Theorem 1.2. At the end of our paper, we give brief recall of some elements of Littlewood-Paley theory.

2. Analysis of the linear perturbation operator $\mathbb{P}\mathcal{A}$

We recall that throughout this paper, we always use upper-case letters to represent vector fields on \mathbb{T}^3 , with four components, the first three components of which form a divergence-free vector field (denoted by the same lower-case letter). More precisely, for a generic vector field A , we have

$$\begin{aligned} A(x_1, x_2, x_3) &= (A^1(x_1, x_2, x_3), A^2(x_1, x_2, x_3), A^3(x_1, x_2, x_3), A^4(x_1, x_2, x_3)) \\ &= (a(x_1, x_2, x_3), A^4(x_1, x_2, x_3)), \end{aligned}$$

where

$$a = (a^1, a^2, a^3) \equiv (A^1, A^2, A^3), \quad \text{and} \quad \text{div } a = 0.$$

As explained in the introduction, the time derivative $\partial_t V^\varepsilon$ is not uniformly bounded. In order to take the limit $\varepsilon \rightarrow 0$, we need to filter the high oscillating terms out of the system (PBS $_\varepsilon$). To this end, we consider the following linear, homogeneous Cauchy problem, which describes the internal waves associated to (PBS $_\varepsilon$)

$$(2.1) \quad \begin{cases} \partial_\tau W + \mathbb{P}\mathcal{A} W = 0, \\ W|_{\tau=0} = W_0 \in L_\sigma^2(\mathbb{T}^3), \end{cases}$$

where

$$L_\sigma^2 \stackrel{def}{=} \left\{ U = (u, U^4) \in L^2, \operatorname{div} u = 0 \text{ and } \int_{\mathbb{T}^3} U(x) dx = 0 \right\}.$$

In [40], a detailed analysis of (2.1) was given in a particular case where the vector fields are supposed to have zero horizontal average. In this section, we will provide a complete spectral analysis of the operator $\mathbb{P}\mathcal{A}$ in the general case where there is no such assumption, which allows to get a detailed description of the solution of (2.1) in $L_\sigma^2(\mathbb{T}^3)$. Using the decomposition (1.8), we can write

$$L_\sigma^2 = \widetilde{L}_\sigma^2 \oplus \underline{L}_\sigma^2,$$

where

$$\begin{aligned} \widetilde{L}_\sigma^2 &= \left\{ U = (u, U^4) \in L_\sigma^2 \mid \int_{\mathbb{T}_h^2} U(x_h, x_3) dx_h = 0 \right\}, \\ \underline{L}_\sigma^2 &= \left\{ U = (u, U^4) \in L_\sigma^2 \mid U = U(x_3) \text{ and } u^3 \equiv 0 \right\}. \end{aligned}$$

Let us remark that, for $U \in \underline{L}_\sigma^2$, since $\operatorname{div} u = 0$ and $u = u(x_3)$, we deduce that $\partial_3 u^3 = 0$, which in turn implies that $u^3 = 0$, taking into account the zero average of u^3 in \mathbb{T}^3 .

Writing the first equation of (2.1) in the Fourier variables, we have

$$(2.2) \quad \begin{cases} \partial_\tau \widehat{W}(\tau, n) + \widehat{\mathbb{P}\mathcal{A}}(n) \widehat{W}(\tau, n) = 0, \\ \widehat{W}(0, n) = \widehat{W}_0(n), \end{cases}$$

where

$$(2.3) \quad \widehat{\mathbb{P}\mathcal{A}}(n) = \begin{pmatrix} 0 & 0 & 0 & -\frac{\check{n}_1 \check{n}_3}{|\check{n}|^2} \\ 0 & 0 & 0 & -\frac{\check{n}_2 \check{n}_3}{|\check{n}|^2} \\ 0 & 0 & 0 & 1 - \frac{\check{n}_3^2}{|\check{n}|^2} \\ 0 & 0 & -1 & 0 \end{pmatrix}.$$

Standard calculations show that the matrix $\widehat{\mathbb{P}\mathcal{A}}(n)$ possesses very different spectral properties in the case where $\check{n}_h = 0$ and in the case where $\check{n}_h \neq 0$.

1. In the case where $\check{n}_h \neq 0$: The matrix $\widehat{\mathbb{P}\mathcal{A}}(n)$ admits an eigenvalue $\omega^0(n) \equiv 0$ of multiplicity 2 and two other conjugate complex eigenvalues

$$(2.4) \quad i\omega^\pm(n) = \pm i\omega(n),$$

where $\omega(n) = \frac{|\check{n}_h|}{|\check{n}|}$. Associated to each eigenvalue, there is a unique unit eigenvector, orthogonal to the frequency vector ${}^t(\check{n}_1, \check{n}_2, \check{n}_3, 0)$, which is explicitly given as follows

$$(2.5) \quad e_0(n) = \frac{1}{|\check{n}_h|} \begin{pmatrix} -\check{n}_2 \\ \check{n}_1 \\ 0 \\ 0 \end{pmatrix}, \quad e_\pm(n) = \frac{1}{\sqrt{2}} \begin{pmatrix} \pm i \frac{\check{n}_1 \check{n}_3}{|\check{n}_h| |\check{n}|} \\ \pm i \frac{\check{n}_2 \check{n}_3}{|\check{n}_h| |\check{n}|} \\ \mp i \frac{|\check{n}_h|}{|\check{n}|} \\ 1 \end{pmatrix}.$$

Since $\{e_\alpha\}_{\alpha=0,\pm}$ form an orthonormal basis of the subspace of \mathbb{C}^4 which is orthogonal to ${}^t(\check{n}_1, \check{n}_2, \check{n}_3, 0)$, the classical theory of ordinary differential equations implies that the solution $\widehat{W}(\tau, n)$ of (2.2), with $\check{n}_h \neq 0$, writes

$$(2.6) \quad \widehat{W}(\tau, n) = \sum_{\alpha \in \{0,\pm\}} e^{i\tau\omega^\alpha(n)} \langle \widehat{W}_0(n), e_\alpha(n) \rangle_{\mathbb{C}^4} e_\alpha(n).$$

2. In the case where $\check{n}_h = 0$: The matrix $\widehat{\mathbb{P}\mathcal{A}}(n)$ becomes

$$\widehat{\mathbb{P}\mathcal{A}}(0, 0, n_3) = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 \end{pmatrix},$$

and admits only one eigenvalue $\omega(0, 0, n_3) = 0$ of multiplicity 4, and three associate unit eigenvectors, orthogonal to the frequency vector ${}^t(0, 0, \check{n}_3, 0)$

$$(2.7) \quad f_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \quad f_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \quad f_3 = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}.$$

The solution $\widehat{W}(\tau, n_3) \equiv \widehat{W}(\tau, 0, 0, n_3)$ of (2.2) writes

$$(2.8) \quad \widehat{W}(\tau, n_3) = \sum_{j \in \{1,2,3\}} \langle \widehat{W}_0(0, 0, n_3), f_j \rangle_{\mathbb{C}^4} f_j.$$

The expressions of the Fourier modes $\widehat{W}(n)$ given in (2.6) and (2.8) imply the following result

Lemma 2.1. *Let $W_0 \in L^2_\sigma(\mathbb{T}^3)$. The unique solution W of the system (2.1) accepts the following decomposition*

$$(2.9) \quad W(\tau, x) = \underline{W}(x_3) + \overline{W}(x) + W_{\text{osc}}(\tau, x),$$

where

$$\begin{aligned} \underline{W}(x_3) &= \sum_{n_3 \in \mathbb{Z}} \sum_{j \in \{1, 2, 3\}} \left\langle \widehat{W}_0(0, 0, n_3), f_j \right\rangle_{\mathbb{C}^4} e^{in_3 x_3} f_j \\ \overline{W}(x) &= \sum_{\substack{n \in \mathbb{Z}^3 \\ n_h \neq 0}} \left\langle \widehat{W}_0(n), e_0(n) \right\rangle_{\mathbb{C}^4} e^{i\tilde{n} \cdot x} e_0(n) \\ W_{\text{osc}}(\tau, x) &= \sum_{\substack{n \in \mathbb{Z}^3 \\ n_h \neq 0}} \sum_{\alpha \in \{\pm\}} \left\langle \widehat{W}_0(n), e_\alpha(n) \right\rangle_{\mathbb{C}^4} e^{i\tau \omega^\alpha(n)} e^{i\tilde{n} \cdot x} e_\alpha(n). \end{aligned}$$

Now, we set

$$\begin{aligned} E_\alpha(n, x) &= e^{i\tilde{n} \cdot x} e_\alpha(n), & \forall n \in \mathbb{Z}^3, n_h \neq 0, \forall \alpha \in \{0, \pm\} \\ F_j(n_3, x_3) &= e^{in_3 x_3} f_j, & \forall n_3 \in \mathbb{Z}, \forall j \in \{1, 2, 3\}, \end{aligned}$$

then, $\{E_\alpha(n, \cdot), F_j(n_3, \cdot)\}$ forms an orthonormal basis of $L^2_\sigma(\mathbb{T}^3)$, and we have the following decomposition

Definition 2.2. *For any vector field $V \in L^2_\sigma(\mathbb{T}^3)$, we have*

$$(2.10) \quad V(x) = \underline{V}(x_3) + \overline{V}(x) + V_{\text{osc}}(x),$$

where

$$\begin{aligned} \underline{V}(x_3) &= \sum_{n_3 \in \mathbb{Z}} \sum_{j \in \{1, 2, 3\}} \left\langle \widehat{V}(0, 0, n_3), f_j \right\rangle_{\mathbb{C}^4} F_j(n_3, x_3) \\ \overline{V}(x) &= \sum_{\substack{n \in \mathbb{Z}^3 \\ n_h \neq 0}} \left\langle \widehat{V}(n), e_0(n) \right\rangle_{\mathbb{C}^4} E_0(n, x) \\ V_{\text{osc}}(\tau, x) &= \sum_{\substack{n \in \mathbb{Z}^3 \\ n_h \neq 0}} \sum_{\alpha \in \{\pm\}} \left\langle \widehat{V}(n), e_\alpha(n) \right\rangle_{\mathbb{C}^4} E_\alpha(n, x). \end{aligned}$$

We also have the following result

Proposition 2.3. *Let Π_X be the projection onto the subspace X of $L^2_\sigma(\mathbb{T}^3)$. For any vector field $V \in L^2_\sigma(\mathbb{T}^3)$, we have*

1. $\Pi_{L^2_\sigma} V = \underline{V}(x_3)$.
2. $\Pi_{\widetilde{L^2_\sigma}} V = \widetilde{V}(x) = V(x) - \underline{V}(x_3) = \overline{V}(x) + V_{\text{osc}}(x)$.

$$3. \underline{V}(x_3) + \overline{V}(x) = \Pi_{\ker(\mathbb{P}\mathcal{A})}V = \Pi_{\ker(\mathcal{L}-\text{Id})}V.$$

Thus, the operator $\mathcal{L}(\tau)$ only acts on the oscillating part V_{osc} of V .

Proof. The points (1) and (2) are immediate consequences of the identity (1.8) and of Definition (2.2). The only non evident point is (3), the proof of which simply follows the lines of the proof of [28, Proposition 4.1]. \square

3. Analysis of the filtered equation

In this section, we will use the method of [38, 28, 23] or [35] to filter out the high oscillation term in the system (PBS_ε) . In order to do so, we first decompose the initial data in the same way as in (2.10), *i.e.*, we write

$$V_0 = \underline{V}_0 + \tilde{V}_0 = \underline{V}_0 + \overline{V}_0 + V_{\text{osc},0}.$$

We recall that in the introduction, we defined \mathcal{L} as the operator which maps $W_0 \in L^2_\sigma(\mathbb{T}^3)$ to the solution W of the linear system (2.1). Using this operator, we now define the following auxiliary vector field

$$U^\varepsilon = \mathcal{L}\left(-\frac{t}{\varepsilon}\right)V^\varepsilon.$$

Replacing $V^\varepsilon = \mathcal{L}\left(\frac{t}{\varepsilon}\right)U^\varepsilon$ into the initial system (PBS_ε) , straightforward computations show that U^ε satisfies the following “filtered” system

$$(S_\varepsilon) \quad \begin{cases} \partial_t U^\varepsilon + \mathcal{Q}^\varepsilon(U^\varepsilon, U^\varepsilon) - \mathcal{A}_2^\varepsilon(D)U^\varepsilon = 0, \\ U^\varepsilon|_{t=0} = V_0, \end{cases}$$

where

$$(3.1) \quad \begin{aligned} & \mathcal{Q}^\varepsilon(V_1, V_2) \\ &= \frac{1}{2}\mathcal{L}\left(\frac{t}{\varepsilon}\right)\mathbb{P}\left[\mathcal{L}\left(-\frac{t}{\varepsilon}\right)V_1 \cdot \nabla \mathcal{L}\left(-\frac{t}{\varepsilon}\right)V_2 + \mathcal{L}\left(-\frac{t}{\varepsilon}\right)V_2 \cdot \nabla \mathcal{L}\left(-\frac{t}{\varepsilon}\right)V_1\right], \end{aligned}$$

and

$$(3.2) \quad \mathcal{A}_2^\varepsilon(D)W = \mathcal{L}\left(\frac{t}{\varepsilon}\right)\mathcal{A}_2(D)\mathcal{L}\left(-\frac{t}{\varepsilon}\right)W.$$

In this section, we will consider the evolution of (S_ε) as the superposition of its projections onto the subspace of horizontal independent (or average) vector fields \underline{L}^2_σ and the subspace of horizontal oscillating vector fields \widetilde{L}^2_σ . Always

denoting \underline{V} and \widetilde{V} the projection of $V \in L_\sigma^2$ onto \underline{L}_σ^2 and \widetilde{L}_σ^2 , we formally decompose (\mathbf{S}_ε) as sum of two following systems

$$\begin{cases} \partial_t \underline{U}^\varepsilon + \underline{\mathcal{Q}}^\varepsilon(U^\varepsilon, U^\varepsilon) - \underline{\mathcal{A}}_2^\varepsilon(D)U^\varepsilon = 0, \\ \underline{U}^\varepsilon|_{t=0} = \underline{V}_0, \end{cases}$$

and

$$\begin{cases} \partial_t \widetilde{U}^\varepsilon + \widetilde{\mathcal{Q}}^\varepsilon(U^\varepsilon, U^\varepsilon) - \widetilde{\mathcal{A}}_2^\varepsilon(D)U^\varepsilon = 0, \\ \widetilde{U}^\varepsilon|_{t=0} = \widetilde{V}_0, \end{cases}$$

where, for the sake of the simplicity, we identify

$$\begin{aligned} \underline{\mathcal{Q}}^\varepsilon(U^\varepsilon, U^\varepsilon) &\equiv \underline{\mathcal{Q}}^\varepsilon(U^\varepsilon, U^\varepsilon) & \underline{\mathcal{A}}_2^\varepsilon(D)U^\varepsilon &\equiv \underline{\mathcal{A}}_2^\varepsilon(D)U^\varepsilon, \\ \widetilde{\mathcal{Q}}^\varepsilon(U^\varepsilon, U^\varepsilon) &\equiv \widetilde{\mathcal{Q}}^\varepsilon(\widetilde{U}^\varepsilon, \widetilde{U}^\varepsilon) & \widetilde{\mathcal{A}}_2^\varepsilon(D)U^\varepsilon &\equiv \widetilde{\mathcal{A}}_2^\varepsilon(D)\widetilde{U}^\varepsilon. \end{aligned}$$

We want to provide explicit formulas of $\underline{\mathcal{Q}}^\varepsilon(U^\varepsilon, U^\varepsilon)$, $\underline{\mathcal{A}}_2^\varepsilon(D)U^\varepsilon$, $\widetilde{\mathcal{Q}}^\varepsilon(U^\varepsilon, U^\varepsilon)$ and $\widetilde{\mathcal{A}}_2^\varepsilon(D)U^\varepsilon$, and to decompose the vectors $\underline{\mathcal{Q}}^\varepsilon(U^\varepsilon, U^\varepsilon)$ and $\underline{\mathcal{A}}_2^\varepsilon(D)U^\varepsilon$ in the L_σ^2 basis

$$\{E_\alpha(n, x), F_j(n_3, x_3)\}_{\substack{n \in \mathbb{Z}^3 \\ n_3 \in \mathbb{Z} \\ \alpha = 0, \pm \\ j = 1, 2, 3}},$$

given in the previous section. To this end, we will introduce some additional notations. For any vector field $V \in L_\sigma^2$, we set

$$V^a(n) = \left(\widehat{V}(n) \Big| e_a(n) \right)_{\mathbb{C}^4} e_a(n), \quad \forall a = 0, \pm, \forall n = (n_h, n_3) \in \mathbb{Z}^3, n_h \neq 0,$$

and

$$V^j(0, n_3) = \left(\widehat{V}(0, n_3) \Big| f_j \right)_{\mathbb{C}^4} f_j, \quad \forall j = 1, 2, 3, \forall n_3 \in \mathbb{Z}.$$

We also define the following quantities in order to shorten as much as possible the forthcoming expressions

$$(3.3) \quad \begin{cases} \omega_{k,m,n}^{a,b,c} = \omega^a(k) + \omega^b(m) - \omega^c(n), & a, b, c = 0, \pm, \\ \omega_n^{a,b} = \omega^a(n) + \omega^b(n), & a, b = 0, \pm, \\ \widetilde{\omega}_{m,n}^{b,c} = \omega^b(m) - \omega^c(n), & b, c = 0, \pm, \\ \omega_{k,m}^{a,b} = \omega^a(k) + \omega^b(m), & a, b = 0, \pm, \end{cases}$$

where the eigenvalues $\omega^0(\cdot)$ and $\omega^\pm(\cdot)$ are defined in the previous section.

Following the lines of [23] or [35], we deduce that, for $c = 0, \pm$, the projection of $\underline{\mathcal{Q}}^\varepsilon(V_1, V_2)$ onto the subspace generated by $E_c(n, \cdot)$, for any $n \in \mathbb{Z}^3$ such that

$n_h \neq 0$, is

$$\begin{aligned}
& \left\langle \mathcal{Q}^\varepsilon (V_1, V_2) \mid E_c(n, x) \right\rangle_{\widetilde{L}_\sigma^2} E_c(n, x) \\
&= \left\langle \mathcal{F} (\mathcal{Q}^\varepsilon (V_1, V_2)) (n) \mid e_c(n) \right\rangle_{\mathbb{C}^4} e^{i\tilde{n} \cdot x} e_c(n), \\
&= \sum_{k=1}^2 \left\langle \mathcal{F} \left(\widetilde{\mathcal{Q}}_k^\varepsilon (V_1, V_2) \right) (n) \mid e_c(n) \right\rangle_{\mathbb{C}^4} e^{i\tilde{n} \cdot x} e_c(n),
\end{aligned}$$

where using the divergence-free property, we can write the Fourier coefficients of the bilinear forms $\widetilde{\mathcal{Q}}_k^\varepsilon$, $k = 1, 2$, as follows

$$\begin{aligned}
& \mathcal{F} \left(\widetilde{\mathcal{Q}}_1^\varepsilon (V_1, V_2) \right) (n) \\
&= \sum_{\substack{k+m=n \\ k_h, m_h \neq 0 \\ a, b, c=0, \pm}} e^{i \frac{t}{\varepsilon} \omega_{k, m, n}^{a, b, c}} \left\langle \widehat{\mathbb{P}}(n) \begin{pmatrix} \tilde{n} \\ 0 \end{pmatrix} \cdot \mathbb{S} (V_1^a(k) \otimes V_2^b(m)) \mid e_c(n) \right\rangle_{\mathbb{C}^4} e_c(n)
\end{aligned}$$

and

$$\begin{aligned}
& \mathcal{F} \left(\widetilde{\mathcal{Q}}_2^\varepsilon (V_1, V_2) \right) (n) \\
&= \sum_{\substack{(0, k_3)+m=n \\ m_h \neq 0 \\ b, c=0, \pm \\ j=1, 2, 3}} e^{i \frac{t}{\varepsilon} \omega_{m, n}^{b, c}} \left\langle \widehat{\mathbb{P}}(n) \begin{pmatrix} \tilde{n} \\ 0 \end{pmatrix} \cdot \mathbb{S} \left(V_1^j(0, k_3) \otimes V_2^b(m) \right) \mid e_c(n) \right\rangle_{\mathbb{C}^4} e_c(n) \\
&+ \sum_{\substack{(0, k_3)+m=n \\ m_h \neq 0 \\ b, c=0, \pm \\ j=1, 2, 3}} e^{i \frac{t}{\varepsilon} \omega_{m, n}^{b, c}} \left\langle \widehat{\mathbb{P}}(n) \begin{pmatrix} \tilde{n} \\ 0 \end{pmatrix} \cdot \mathbb{S} \left(V_1^b(m) \otimes V_2^j(0, k_3) \right) \mid e_c(n) \right\rangle_{\mathbb{C}^4} e_c(n).
\end{aligned}$$

Here, $\begin{pmatrix} \tilde{n} \\ 0 \end{pmatrix}$ stands for the four-component vector ${}^t(\tilde{n}_1, \tilde{n}_2, \tilde{n}_3, 0)$ and \mathbb{P} is the Leray projection, defined in (1.4). In order to shorten the notations, we use \mathbb{S} for the following symmetry operator

$$\mathbb{S} (V_1 \otimes V_2) = V_1 \otimes V_2 + V_2 \otimes V_1,$$

for any $V_1 = (v_1, V_1^4)$ and $V_2 = (v_2, V_2^4)$. We remark that in the above summation formula there is no bilinear interaction which involves elements of the form $V^{j_1}(0, k_3) \otimes V^{j_2}(0, m_3)$ since, a priori, these represent vector fields the horizontal average of which is not zero and hence they do not belong to \widetilde{L}_σ^2 .

The projection of $\mathcal{Q}^\varepsilon (V_1, V_2)$ onto the subspace generated by $F_j(n_3, \cdot)$, for

any $j = 1, 2, 3$, for any $n_3 \in \mathbb{Z}$, can be computed in a similar way and we get

$$\begin{aligned}
& \langle \mathcal{Q}^\varepsilon(V_1, V_2) \mid F_j(n_3, x_3) \rangle_{L^2_\sigma} F_j(n_3, x_3) \\
&= \langle \mathcal{F}(\mathcal{Q}^\varepsilon(V_1, V_2))(0, n_3) \mid f_j \rangle_{\mathbb{C}^4} e^{i\tilde{n}_3 x_3} f_j, \\
&= \sum_{\substack{k+m=(0, n_3) \\ k_h, m_h \neq 0 \\ a, b \in \{0, \pm\}}} e^{i\frac{t}{\varepsilon} \omega_{k, m}^{a, b}} \left\langle \widehat{\mathbb{P}}(0, n_3) \begin{pmatrix} 0 \\ 0 \\ \tilde{n}_3 \\ 0 \end{pmatrix} \cdot \mathbb{S}(V_1^a(k) \otimes V_2^b(m)) \mid f_j \right\rangle_{\mathbb{C}^4} e^{i\tilde{n}_3 x_3} f_j, \\
&= \langle \mathcal{F}(\underline{\mathcal{Q}}^\varepsilon(V_1, V_2))(0, n_3) \mid f_j \rangle e^{i\tilde{n}_3 x_3} f_j.
\end{aligned}$$

Hence,

$$\begin{aligned}
\mathcal{Q}^\varepsilon(V_1, V_2)(x) &= \sum_{\substack{n \in \mathbb{Z}^3 \\ n_h \neq 0}} \mathcal{F}(\tilde{\mathcal{Q}}_1^\varepsilon(V_1, V_2) + \tilde{\mathcal{Q}}_2^\varepsilon(V_1, V_2))(n) e^{i\tilde{n} \cdot x} \\
&\quad + \sum_{n_3 \in \mathbb{Z}} \mathcal{F}(\underline{\mathcal{Q}}^\varepsilon(V_1, V_2))(0, n_3) e^{i\tilde{n}_3 x_3}.
\end{aligned}$$

The decomposition of $\mathcal{A}_2^\varepsilon(D)W$ can also be calculated as in [23]. We have

$$\begin{aligned}
& \langle \mathcal{A}_2^\varepsilon(D)W \mid E_b(n, x) \rangle_{\widetilde{L^2_\sigma}} E_b(n, x) \\
&= \sum_{a=0, \pm} e^{i\frac{t}{\varepsilon} \omega_n^{a, b}} \langle \mathcal{F}\mathcal{A}_2(n)W^a(n) \mid e_b(n) \rangle_{\mathbb{C}^4} e^{i\tilde{n} \cdot x} e_b(n),
\end{aligned}$$

and

$$\langle \mathcal{A}_2^\varepsilon(D)W \mid F_j(n_3, x_3) \rangle_{L^2_\sigma} F_j(n_3, x_3) = \langle \mathcal{F}\mathcal{A}_2(0, n_3)W^j(0, n_3) \mid f_j \rangle_{\mathbb{C}^4} e^{i\tilde{n}_3 \cdot x_3} f_j.$$

Thus,

$$\begin{aligned}
\widetilde{\mathcal{A}}_2^\varepsilon(D)W &= \widetilde{\mathcal{A}}_2^\varepsilon(D)\widetilde{W} \\
&= \sum_{\substack{n \in \mathbb{Z}^3 \\ n_h \neq 0}} \sum_{a, b=0, \pm} e^{i\frac{t}{\varepsilon} \omega_n^{a, b}} \left\langle \mathcal{F}\mathcal{A}_2(n)W^a(n) \mid e_b(n) \right\rangle_{\mathbb{C}^4} e^{i\tilde{n} \cdot x} e_b(n),
\end{aligned}$$

and

$$\begin{aligned}
\underline{\mathcal{A}}_2^\varepsilon(D)W &= \underline{\mathcal{A}}_2^0(D_3)\underline{W} = \sum_{n_3 \in \mathbb{Z}} \sum_{j=1, 2, 3} \langle \mathcal{F}\mathcal{A}_2(0, n_3)W^j(0, n_3) \mid f_j \rangle_{\mathbb{C}^4} e^{i\tilde{n}_3 x_3} f_j \\
&= (\nu \partial_3^2 \underline{W}^1, \nu \partial_3^2 \underline{W}^2, 0, 0),
\end{aligned}$$

where $\underline{\mathcal{A}}_2^0(D_3)$ is defined in (4.5). We remark that the operator $\underline{\mathcal{A}}_2^\varepsilon$ does not present oscillations, *i.e.* it is independent of ε .

Combining all the above calculations, we get the following decomposition of the system (\mathbf{S}_ε)

Lemma 3.1. *Let $V_0 \in H^s(\mathbb{T}^3)$, $s > 5/2$. We set*

$$\underline{V}_0(x_3) = \frac{1}{4\pi^2 a_1 a_2} \int_{\mathbb{T}_h^2} V_0(y_h, x_3) dy_h,$$

and

$$\tilde{V}_0 = V_0 - \underline{V}_0.$$

Then, the local solution U^ε of (\mathbf{S}_ε) can be written as the superposition

$$U^\varepsilon = \tilde{U}^\varepsilon + \underline{U}^\varepsilon,$$

where \tilde{U}^ε and $\underline{U}^\varepsilon$ are local solutions of the equations

$$(3.4) \quad \begin{cases} \partial_t \tilde{U}^\varepsilon + \tilde{\mathcal{Q}}_1(\tilde{U}^\varepsilon, \tilde{U}^\varepsilon) + \tilde{\mathcal{Q}}_2(U^\varepsilon, U^\varepsilon) - \tilde{\mathcal{A}}_2^\varepsilon(D) \tilde{U}^\varepsilon = 0, \\ \operatorname{div} \tilde{U}^\varepsilon = 0, \\ \tilde{U}^\varepsilon|_{t=0} = \tilde{V}_0, \end{cases}$$

and

$$(3.5) \quad \begin{cases} \partial_t \underline{U}^\varepsilon + \underline{\mathcal{Q}}(\tilde{U}^\varepsilon, \tilde{U}^\varepsilon) - \underline{\mathcal{A}}_2^0(D_3) \underline{U}^\varepsilon = 0, \\ \underline{U}^\varepsilon|_{t=0} = \underline{V}_0. \end{cases}$$

4. The limit system

In this short section, the convergence of suitable subsequences of local strong solutions $(U^\varepsilon)_{\varepsilon>0}$ of the system (\mathbf{S}_ε) will be put in evidence. Moreover, using the similar methods as in [23] or [40], we can determine the systems which describe the evolution of such limits. The explicit formulation of these limiting systems will be given in the next section. We first introduce the limit forms $\tilde{\mathcal{Q}}_1$, $\tilde{\mathcal{Q}}_2$, $\underline{\mathcal{Q}}$, $\tilde{\mathcal{A}}_2^0$ and $\underline{\mathcal{A}}_2^0$ such that

$$(4.1) \quad \begin{aligned} & \tilde{\mathcal{Q}}_1(V_1, V_2) \\ &= \sum_{\substack{n \in \mathbb{Z}^3 \\ n_h \neq 0}} \sum_{\substack{\omega_{k,m,n}^{a,b,c} = 0 \\ k+m=n \\ k_h, m_h, n_h \neq 0 \\ a,b,c \in \{0, \pm\}}} \left\langle \hat{\mathbb{P}}(n) \binom{n}{0} \cdot \mathbb{S}(V_1^a(k) \otimes V_2^b(m)) \Big| e_c(n) \right\rangle_{\mathbb{C}^4} e^{i\tilde{n} \cdot x} e_c(n), \end{aligned}$$

$$\begin{aligned}
(4.2) \quad & \tilde{\mathcal{Q}}_2(V_1, V_2) \\
&= \sum_{\substack{n \in \mathbb{Z}^3 \\ n_h \neq 0}} \sum_{\substack{(0, k_3) + m = n \\ m_h, n_h \neq 0 \\ \tilde{\omega}_{m, n}^{b, c} = 0 \\ b, c \in \{0, \pm\} \\ j = 1, 2, 3}} \left\langle \hat{\mathbb{P}}(n) \binom{n}{0} \cdot \mathbb{S} \left(V_1^j(0, k_3) \otimes V_2^b(m) \right) \Big| e_c(n) \right\rangle_{\mathbb{C}^4} e^{i\tilde{n} \cdot x} e_c(n) \\
&+ \sum_{\substack{n \in \mathbb{Z}^3 \\ n_h \neq 0}} \sum_{\substack{(0, k_3) + m = n \\ m_h, n_h \neq 0 \\ \tilde{\omega}_{m, n}^{b, c} = 0 \\ b, c \in \{0, \pm\} \\ j = 1, 2, 3}} \left\langle \hat{\mathbb{P}}(n) \binom{n}{0} \cdot \mathbb{S} \left(V_1^b(m) \otimes V_2^j(0, k_3) \right) \Big| e_c(n) \right\rangle_{\mathbb{C}^4} e^{i\tilde{n} \cdot x} e_c(n),
\end{aligned}$$

$$\begin{aligned}
(4.3) \quad & \underline{\mathcal{Q}}(V_1, V_2) \\
&= \sum_{n_3 \in \mathbb{Z}} \sum_{\substack{k+m=(0, n_3) \\ k_h, m_h \neq 0 \\ \omega^a(k) + \omega^b(m) = 0 \\ a, b \in \{0, \pm\} \\ j = 1, 2, 3}} \left\langle \hat{\mathbb{P}}(0, n_3) \binom{0}{\tilde{n}_3} \cdot \mathbb{S} \left(V_1^a(k) \otimes V_2^b(m) \right) \Big| f_j \right\rangle_{\mathbb{C}^4} e^{i\tilde{n}_3 x_3} f_j,
\end{aligned}$$

$$(4.4) \quad \widetilde{\mathcal{A}}_2^0(D)W = \sum_{\substack{n \in \mathbb{Z}^3 \\ n_h \neq 0}} \sum_{\substack{\omega_n^{a, b} = 0 \\ a, b \in \{0, \pm\}}} \left\langle \mathcal{F}\mathcal{A}_2(n) \widetilde{W}^a(n) \Big| e_b(n) \right\rangle_{\mathbb{C}^4} e^{i\tilde{n} \cdot x} e_b(n),$$

and

$$(4.5) \quad \underline{\mathcal{A}}_2^0(D)W = \sum_{n_3 \in \mathbb{Z}} \sum_{j=1, 2, 3} \left\langle \mathcal{F}\mathcal{A}(0, n_3) W^j(0, n_3) \Big| f_j \right\rangle_{\mathbb{C}^4} e^{i\tilde{n}_3 x_3} f_j.$$

Using the method of the non-stationary phases (see for instance [1]), we can prove the following convergence result (for more details, see [23, 28, 35] or [42]).

Lemma 4.1. *Let $T > 0$ and V_1, V_2 and W be zero-average smooth vector fields in $C^\infty([0, T] \times \mathbb{T}^3)$. Then, we have the following convergence in the sense of distributions*

$$\begin{aligned}
& \tilde{\mathcal{Q}}_k^\varepsilon(V_1, V_2) \xrightarrow{\varepsilon \rightarrow 0} \tilde{\mathcal{Q}}_k(V_1, V_2), \quad k = 1, 2 \\
& \underline{\mathcal{Q}}^\varepsilon(V_1, V_2) \xrightarrow{\varepsilon \rightarrow 0} \underline{\mathcal{Q}}(V_1, V_2), \\
& \widetilde{\mathcal{A}}_2^\varepsilon(D)W \xrightarrow{\varepsilon \rightarrow 0} \widetilde{\mathcal{A}}_2^0(D)W, \\
& \underline{\mathcal{A}}_2^\varepsilon(D_3)W \xrightarrow{\varepsilon \rightarrow 0} \underline{\mathcal{A}}_2^0(D_3)W.
\end{aligned}$$

Let us now define the (limit) bilinear forms

$$\begin{aligned}
(4.6) \quad & \mathcal{Q}(V_1, V_2) = \tilde{\mathcal{Q}}_1(V_1, V_2) + \tilde{\mathcal{Q}}_2(V_1, V_2) + \underline{\mathcal{Q}}(V_1, V_2), \\
& \mathcal{A}_2^0(D)W = \widetilde{\mathcal{A}}_2^0(D)W + \underline{\mathcal{A}}_2^0(D)W.
\end{aligned}$$

Then, the limit dynamics of (PBS_ε) as $\varepsilon \rightarrow 0$ can be described as follows

Proposition 4.2. *Let $V_0 \in H^s(\mathbb{T}^3)$, $s > 5/2$ and $T \in [0, T^*[$, where $T^* > 0$ is defined in Theorem 1.1. The sequence $(U^\varepsilon)_{\varepsilon>0}$ of local strong solutions of (S_ε) , which is uniformly bounded in $\mathcal{C}([0, T]; H^s(\mathbb{T}^3))$, is compact in the space $\mathcal{C}([0, T]; H^\sigma(\mathbb{T}^3))$ where $\sigma \in (s-2, s)$. Moreover each limit point U of $(U^\varepsilon)_{\varepsilon>0}$ solves the following limit equation*

$$(S_0) \quad \begin{cases} \partial_t U + \mathcal{Q}(U, U) - \mathcal{A}_2^0(D)U = 0, \\ U|_{t=0} = V_0, \end{cases}$$

in $\mathbb{T}^3 \times [0, T]$, where \mathcal{Q} and $\mathcal{A}_2^0(D)$ are given in (4.6).

Proof. The proof of the compactness of the sequence $(U^\varepsilon)_{\varepsilon>0}$ is a standard argument. Thanks to Theorem 1.1 we know that $(U^\varepsilon)_{\varepsilon>0}$ is uniformly bounded in $\mathcal{C}([0, T]; H^s(\mathbb{T}^3))$. Since

$$\|\mathcal{Q}^\varepsilon(U^\varepsilon, U^\varepsilon)\|_{H^{s-1}} \leq C \|U^\varepsilon\|_{H^s}^2,$$

and

$$\|\mathcal{A}_2^\varepsilon(D)U^\varepsilon\|_{H^{s-2}} \leq C \|U^\varepsilon\|_{H^s},$$

we deduce that $(\partial_t U^\varepsilon)_{\varepsilon>0}$ is uniformly bounded in $\mathcal{C}([0, T]; H^{s-2})$. Remarking that

$$\mathcal{C}([0, T]; H^{s-2}) \hookrightarrow L^1([0, T]; H^{s-2}),$$

we can hence apply Aubin-Lions lemma [2] to deduce the compactness of the sequence.

Finally, we need to prove that a limit point U (weakly) solves the limit system (S_0) . We remark that Lemma 4.1 cannot be directly applied since the sequence $(U^\varepsilon)_{\varepsilon>0}$ is not smooth enough. However, by mollifying the data as in [28, 38] or [42], we can deduce that U solves (S_0) in $\mathcal{D}'(\mathbb{T}^3 \times [0, T])$. We choose now $\bar{\sigma} \in (\frac{5}{2}, s)$. Since $U \in \mathcal{C}([0, T]; H^{\bar{\sigma}})$, Sobolev embeddings imply that $U \in \mathcal{C}([0, T]; \mathcal{C}^{1,1})$, and so it solves the system (S_0) in $\mathbb{T}^3 \times [0, T]$ \square

Remark 4.3. A similar result to Proposition 4.2 was proved in [40, Lemma 3.3]. We underline though that the proof of [40, Lemma 3.3] cannot be adapted to the present setting since it strongly used the regularity induced by the uniform parabolic smoothing effects.

5. Explicit formulations of the limit system

In this section, we determine the explicit formulation of the limit system (S_0) . More precisely, for each limit point U of the sequence of solutions $(U^\varepsilon)_{\varepsilon>0}$ of (S_ε) , we decompose the bilinear term $\mathcal{Q}(U, U)$ and the linear term $\mathcal{A}_2^0(D)U$ by mean of the projections onto $\ker \mathbb{P}\mathcal{A}$ defined in Proposition 2.3 and onto its

orthogonal $(\ker \mathbb{P}\mathcal{A})^\perp$. Such approach is generally used for hyperbolic symmetric systems with skew-symmetric perturbation in periodic domains, and we refer the reader to [23, 24, 25, 35, 40] or [42], for instance, for some other related systems. Since the spectral properties of the operator $\mathbb{P}\mathcal{A}$ are rather different in the Fourier frequency subspaces $\{n_h = 0\}$ and $\{n_h \neq 0\}$, as in Section 3, we write the limit system (S_0) as the superposition of the following systems

$$(5.1) \quad \begin{cases} \partial_t \underline{U} + \underline{\mathcal{Q}}(\tilde{U}, \tilde{U}) - \underline{\mathcal{A}}_2^0(D_3) \underline{U} = 0, \\ \underline{U}^3 \equiv 0, \\ \underline{U}|_{t=0} = \underline{V}_0, \end{cases}$$

and

$$(5.2) \quad \begin{cases} \partial_t \tilde{U} + \tilde{\mathcal{Q}}_1(\tilde{U}, \tilde{U}) + \tilde{\mathcal{Q}}_2(U, U) - \tilde{\mathcal{A}}_2^0(D) \tilde{U} = 0, \\ \operatorname{div} \tilde{U} = 0, \\ \tilde{U}|_{t=0} = \tilde{V}_0, \end{cases}$$

where the limit terms $\underline{\mathcal{Q}}(\tilde{U}, \tilde{U})$, $\underline{\mathcal{A}}_2^0(D_3) \underline{U}$, $\tilde{\mathcal{Q}}_1(\tilde{U}, \tilde{U})$, $\tilde{\mathcal{Q}}_2(\tilde{U}, \tilde{U})$ and $\tilde{\mathcal{A}}_2^0(D) \tilde{U}$ are defined as in Lemma 4.1. In what follows, we will separately study the systems (5.1) and (5.2).

5.1. The dynamics of \underline{U}

We will prove the following result, which allows to simplify the system (5.1).

Proposition 5.1. *Let $V_0 \in H^s(\mathbb{T}^3)$, $s > 5/2$, and $V_0 = \underline{V}_0 + \tilde{V}_0$ as in Lemma 3.1. Then, the horizontal average $\underline{U} = (\underline{u}^h, 0, \underline{U}^4)$ of the solution U of the limit system (S_0) solves the homogeneous diffusion system (1.9) in $[0, T] \times \mathbb{T}_\nu^1$ for each $T \in [0, T^*[$.*

In order to prove Proposition 5.1, we only need to prove the following lemma

Lemma 5.2. *The following identity holds true*

$$\underline{\mathcal{Q}}(\tilde{U}, \tilde{U}) = 0.$$

We recall that in Section 3, we already introduce, for any vector field $V \in L_\sigma^2$,

$$V^a(n) = \left(\widehat{V}(n) \Big| e_a(n) \right)_{\mathbb{C}^4} e_a(n), \quad \forall a = 0, \pm, \forall n = (n_h, n_3) \in \mathbb{Z}^3, n_h \neq 0.$$

Here, we also denote

$$\widehat{V}^a(n) = \left(\widehat{V}(n) \Big| e_a(n) \right)_{\mathbb{C}^4},$$

and $V^{a,l}$, $l = 1, 2, 3, 4$ the l -th component and respectively $V^{a,h}$ the first two components of the vector V^a . Using the definition of \underline{Q} and the particular form of the vector ${}^t(0, 0, \tilde{n}_3, 0)$ given in (4.3), we have

$$\begin{aligned} \mathcal{F}(\underline{Q}(U, U))(0, n_3) &= \sum_{\substack{k+m=(0, n_3) \\ k_h, m_h \neq 0 \\ \omega^a(k) + \omega^b(m) = 0 \\ a, b = 0, \pm \\ j=1, 2, 3}} \left\langle \widehat{\mathbb{P}}(0, n_3) n_3 U^{a,3}(k) U^b(m) \middle| f_j \right\rangle_{\mathbb{C}^4} f_j, \\ &= \sum_{a, b = 0, \pm} \sum_{j=1}^3 \sum_{\mathcal{I}_{a,b}(n_3)} \left\langle \widehat{\mathbb{P}}(0, n_3) n_3 U^{a,3}(k) U^b(m) \middle| f_j \right\rangle_{\mathbb{C}^4} f_j, \end{aligned}$$

where the set $\mathcal{I}_{a,b}(n_3)$ contains the following resonance frequencies

$$(5.3) \quad \mathcal{I}_{a,b}(n_3) = \left\{ (k, m) \in \mathbb{Z}^6, k_h, m_h \neq 0 \mid k + m = (0, n_3), \omega^a(k) + \omega^b(m) = 0 \right\}.$$

In order to prove Lemma 5.2, we will show that, for each couple $(a, b) \in \{0, \pm\}^2$, the contribution

$$(5.4) \quad \mathcal{J}_{a,b}(n_3) = \sum_{\mathcal{I}_{a,b}(n_3)} \widehat{\mathbb{P}}(0, n_3) n_3 U^{a,3}(k) U^b(m),$$

is null, which implies that

$$(5.5) \quad \mathcal{F}(\underline{Q}(U, U))(0, n_3) = \sum_{a, b = 0, \pm} \sum_{j=1}^3 \left\langle \mathcal{J}_{a,b}(n_3) \middle| f_j \right\rangle_{\mathbb{C}^4} f_j = 0.$$

Case 1: $(a, b) = (0, 0)$. We have

$$\mathcal{J}_{0,0}(n_3) = \sum_{k+m=(0, n_3)} \widehat{\mathbb{P}}(0, n_3) n_3 U^{0,3}(k) U^0(m) = 0,$$

since $U^{0,3} \equiv 0$ (see (2.5) and (2.7)).

Case 2: $(a, b) = (\pm, 0)$ or $(a, b) = (0, \pm)$. If $(a, b) = (\pm, 0)$, then,

$$\mathcal{J}_{\pm,0}(n_3) = \sum_{\substack{k+m=(0, n_3) \\ \omega^\pm(k)=0}} \widehat{\mathbb{P}}(0, n_3) n_3 U^{\pm,3}(k) U^0(m).$$

The condition $\omega^\pm(k) = 0$ implies that $k_h \equiv 0$, while the condition $k+m = (0, n_3)$ implies that $m_h \equiv 0$. Then from (2.7), we have $U^{a,3}(0, k_3) \equiv 0$, which shows that $\mathcal{J}_{\pm,0}(n_3)$ gives a null contribution in (5.5). The same approach can be applied to the case $(a, b) = (0, \pm)$.

Case 3: $(a, b) = (+, +)$ or $(a, b) = (-, -)$. In this case, $\mathcal{J}_{a,b}(n_3)$ writes

$$\mathcal{J}_{\pm,\pm}(n_3) = \sum_{\substack{k+m=(0,n_3) \\ \omega^\pm(k)+\omega^\pm(m)=0}} \widehat{\mathbb{P}}(0, n_3) n_3 U^{\pm,3}(k) U^\pm(m).$$

Since $k + m = (0, n_3)$, we can set $|\check{k}_h| = |\check{m}_h| = \lambda$. We deduce from the constraint $\omega^\pm(k) + \omega^\pm(m) = 0$ and the explicit formulation of the eigenvalues (2.4) that

$$\frac{\lambda}{\sqrt{\lambda^2 + \check{k}_3^2}} + \frac{\lambda}{\sqrt{\lambda^2 + \check{m}_3^2}} = 0,$$

which implies that $\lambda \equiv 0$. Then the similar argument as in Cases 1 and 2 shows that $\mathcal{J}_{\pm,\pm}(n_3) = 0$.

Case 4: $(a, b) = (+, -)$ or $(a, b) = (-, +)$. This is the most delicate case to treat. We write

$$(5.6) \quad \mathcal{J}_{\pm,\mp}(n_3) = \sum_{\substack{k+m=(0,n_3) \\ \omega^\pm(k)=\omega^\pm(m)}} \widehat{\mathbb{P}}(0, n_3) n_3 U^{\pm,3}(k) U^\mp(m).$$

The conditions $k + m = (0, n_3)$ and $\omega^\pm(k) = \omega^\pm(m)$ imply now that $k_h = -m_h$, $|\check{k}_h| = |\check{m}_h| = \lambda$ and

$$\frac{\lambda}{\sqrt{\lambda^2 + \check{k}_3^2}} = \frac{\lambda}{\sqrt{\lambda^2 + \check{m}_3^2}}.$$

Then, it is obvious that $m_3 = \pm k_3$.

If $m_3 = -k_3$ then the convolution constraint $k_3 + m_3 = n_3$ in (5.6) implies that $n_3 \equiv 0$, and hence there is no contributions of $\mathcal{J}_{\pm,\mp}(n_3)$ in (5.5). So, we concentrate on the case where $k_h = -m_h$ and $k_3 = m_3 = \frac{n_3}{2}$ and we will deal with the interaction will be of the form, for any $n_3 \in 2\mathbb{Z}$,

$$\mathcal{J}_{\pm,\mp}(n_3) = \sum_{m_h \in \mathbb{Z}^2} \widehat{\mathbb{P}}(0, n_3) n_3 U^{\pm,3} \left(-m_h, \frac{n_3}{2} \right) U^\mp \left(m_h, \frac{n_3}{2} \right).$$

For any $n_3 \in 2\mathbb{Z}$, we set

$$B_{n_3}^{\pm,\mp} = \sum_{m_h \in \mathbb{Z}^2} n_3 U^{\pm,3} \left(-m_h, \frac{n_3}{2} \right) U^{\mp,h} \left(m_h, \frac{n_3}{2} \right),$$

$$C_{n_3}^{\pm,\mp} = \sum_{m_h \in \mathbb{Z}^2} n_3 U^{\pm,3} \left(-m_h, \frac{n_3}{2} \right) U^{\mp,4} \left(m_h, \frac{n_3}{2} \right).$$

Taking into account the form of the vectors f_j in (2.7), we deduce that

$$\sum_{j=1}^3 \langle \mathcal{J}_{+,-}(n_3) + \mathcal{J}_{-,+}(n_3) | f_j \rangle_{\mathbb{C}^4} f_j = \widehat{\mathbb{P}}(0, n_3) \begin{pmatrix} B_{n_3}^{+,-} + B_{n_3}^{-,+} \\ 0 \\ C_{n_3}^{+,-} + C_{n_3}^{-,+} \end{pmatrix}.$$

Then, we can prove that the sum $\mathcal{J}_{+,-}(n_3) + \mathcal{J}_{-,+}(n_3)$ have no contribution in (5.5) and conclude Case 4 if we prove the following lemma

Lemma 5.3. *For any $n_3 \in 2\mathbb{Z}$, we have the following identities*

$$(5.7) \quad B_{n_3}^{+,-} = -B_{n_3}^{-,+},$$

and

$$(5.8) \quad C_{n_3}^{+,-} = -C_{n_3}^{-,+}.$$

Proof. Identity (5.8) is quite easy to prove. Indeed, we have

$$C_{n_3}^{\pm,\mp} = \sum_{m_h \in \mathbb{Z}^2} n_3 e_{\pm}^3 \left(-m_h, \frac{n_3}{2} \right) \widehat{U}^{\pm} \left(-m_h, \frac{n_3}{2} \right) \widehat{U}^{\mp} \left(-m_h, \frac{n_3}{2} \right).$$

Then,

$$(5.9) \quad \begin{aligned} C_{n_3}^{+,-} + C_{n_3}^{-,+} &= \sum_{m_h \in \mathbb{Z}^2} \frac{n_3}{2} \left[e_{\pm}^3 \left(-m_h, \frac{n_3}{2} \right) \widehat{U}^{\pm} \left(-m_h, \frac{n_3}{2} \right) \widehat{U}^{\mp} \left(-m_h, \frac{n_3}{2} \right) \right. \\ &\quad \left. + e_{\mp}^3 \left(m_h, \frac{n_3}{2} \right) \widehat{U}^{\mp} \left(-m_h, \frac{n_3}{2} \right) \widehat{U}^{\pm} \left(-m_h, \frac{n_3}{2} \right) \right]. \end{aligned}$$

The explicit expression of $e_{\mp}(n)$ in (2.5) yields

$$(5.10) \quad e_{\pm}^3 \left(-m_h, \frac{n_3}{2} \right) = -e_{\mp}^3 \left(m_h, \frac{n_3}{2} \right),$$

which, combined with (5.9), implies (5.8).

To prove Identity (5.7), we consider the quantities

$$(5.11) \quad \begin{aligned} \beta(m_h, n_3) &= \frac{n_3}{2} \left[U^{+,3} \left(-m_h, \frac{n_3}{2} \right) U^{-,h} \left(m_h, \frac{n_3}{2} \right) \right. \\ &\quad + U^{-,3} \left(-m_h, \frac{n_3}{2} \right) U^{+,h} \left(m_h, \frac{n_3}{2} \right) \\ &\quad + U^{+,3} \left(m_h, \frac{n_3}{2} \right) U^{-,h} \left(-m_h, \frac{n_3}{2} \right) \\ &\quad \left. + U^{-,3} \left(m_h, \frac{n_3}{2} \right) U^{+,h} \left(-m_h, \frac{n_3}{2} \right) \right], \end{aligned}$$

which allows to write

$$(5.12) \quad B_{n_3}^{+,-} + B_{n_3}^{-,+} = \sum_{m_h \in \mathbb{Z}^2} \beta(m_h, n_3).$$

Now, we decompose

$$\beta(m_h, n_3) = \beta^+(m_h, n_3) + \beta^-(m_h, n_3),$$

where

$$\beta^+(m_h, n_3) = \frac{n_3}{2} \left[U^{+,3} \left(-m_h, \frac{n_3}{2} \right) U^{-,h} \left(m_h, \frac{n_3}{2} \right) + U^{-,3} \left(m_h, \frac{n_3}{2} \right) U^{+,h} \left(-m_h, \frac{n_3}{2} \right) \right],$$

and

$$\beta^-(m_h, n_3) = \frac{n_3}{2} \left[U^{-,3} \left(-m_h, \frac{n_3}{2} \right) U^{+,h} \left(m_h, \frac{n_3}{2} \right) + U^{+,3} \left(m_h, \frac{n_3}{2} \right) U^{-,h} \left(-m_h, \frac{n_3}{2} \right) \right].$$

By definition, we have

$$\begin{aligned} & U^{+,3} \left(-m_h, \frac{n_3}{2} \right) U^{-,h} \left(m_h, \frac{n_3}{2} \right) \\ &= \left\langle \widehat{U} \left(-m_h, \frac{n_3}{2} \right) \middle| e_+ \left(-m_h, \frac{n_3}{2} \right) \right\rangle_{\mathbb{C}^4} e_+^3 \left(-m_h, \frac{n_3}{2} \right) \\ & \quad \times \left\langle \widehat{U} \left(m_h, \frac{n_3}{2} \right) \middle| e_- \left(m_h, \frac{n_3}{2} \right) \right\rangle_{\mathbb{C}^4} e_-^h \left(m_h, \frac{n_3}{2} \right), \end{aligned}$$

and

$$\begin{aligned} & U^{-,3} \left(m_h, \frac{n_3}{2} \right) U^{+,h} \left(-m_h, \frac{n_3}{2} \right) \\ &= \left\langle \widehat{U} \left(m_h, \frac{n_3}{2} \right) \middle| e_- \left(m_h, \frac{n_3}{2} \right) \right\rangle_{\mathbb{C}^4} e_-^3 \left(m_h, \frac{n_3}{2} \right) \\ & \quad \times \left\langle \widehat{U} \left(-m_h, \frac{n_3}{2} \right) \middle| e_+ \left(-m_h, \frac{n_3}{2} \right) \right\rangle_{\mathbb{C}^4} e_+^h \left(-m_h, \frac{n_3}{2} \right). \end{aligned}$$

The explicit formula (2.5) implies

$$\begin{aligned} e_-^h \left(m_h, \frac{n_3}{2} \right) &= e_+^h \left(-m_h, \frac{n_3}{2} \right) = A_{m_h, n_3}^h, \\ e_+^3 \left(-m_h, \frac{n_3}{2} \right) &= -e_-^3 \left(m_h, \frac{n_3}{2} \right) = A_{m_h, n_3}^3. \end{aligned}$$

Setting

$$\begin{aligned} C_{m_h, n_3} &= \left\langle \widehat{U} \left(-m_h, \frac{n_3}{2} \right) \middle| e_+ \left(-m_h, \frac{n_3}{2} \right) \right\rangle_{\mathbb{C}^4} \\ & \quad \times \left\langle \widehat{U} \left(m_h, \frac{n_3}{2} \right) \middle| e_- \left(m_h, \frac{n_3}{2} \right) \right\rangle_{\mathbb{C}^4}, \end{aligned}$$

we obtain

$$\begin{aligned} U^{+,3} \left(-m_h, \frac{n_3}{2} \right) \widehat{U}^{-,h} \left(m_h, \frac{n_3}{2} \right) &= -C_{m_h, n_3} A_{m_h, n_3}^h A_{m_h, n_3}^3, \\ U^{-,3} \left(m_h, \frac{n_3}{2} \right) \widehat{U}^{+,h} \left(-m_h, \frac{n_3}{2} \right) &= C_{m_h, n_3} A_{m_h, n_3}^h A_{m_h, n_3}^3, \end{aligned}$$

which imply

$$\beta^+(m_h, n_3) \equiv 0.$$

By the similar argument, we also get

$$\beta^-(m_h, n_3) \equiv 0,$$

which yields

$$\beta(m_h, n_3) \equiv 0.$$

Thus, Identity (5.12) implies (5.7)

□

5.2. The dynamics of \tilde{U}

In the previous paragraph, the dynamics of \underline{U} is well understood and turns out to follow quite a simple heat equation. To complete the study of the limit system (S_0) , we now give an explicit expression of the system (5.2). As in Lemma 2.1 and Proposition 2.3 (2), we will study the evolution of \tilde{U} as the superposition of

$$\tilde{U} = \bar{U} + U_{\text{osc}}.$$

The main technical difficulty of the study consists in giving a close formulation of the projection of the bilinear interactions, which was considered in [40]. In what follows, we only mention the main steps of the study, without going into technical calculations.

5.2.1. Derivation of the evolution of \bar{U}

We recall that \bar{U} is the projection of \tilde{U} onto the nonoscillating subspace generated by $\{E_0(n, \cdot)\}_n$. The derivation of the evolution of \bar{U} can be done in three steps.

Step 1: We explicitly compute the projections of $\tilde{Q}_1(\tilde{U}, \tilde{U})$ and $\tilde{Q}_2(U, U)$ onto $\text{Span}\{E_0(n, \cdot)\}$, i.e., for any $n \in \mathbb{Z}^3$, $n_h \neq 0$, we compute the following quantities

$$\begin{aligned} \overline{\tilde{Q}_1(\tilde{U}, \tilde{U})} &= \sum_{\substack{n \in \mathbb{Z}^3 \\ n_h \neq 0}} \left\langle \mathcal{F}\tilde{Q}_1(\tilde{U}, \tilde{U})(n) \mid e_0(n) \right\rangle_{\mathbb{C}^4} e^{i\tilde{n} \cdot x} e_0(n) \\ \overline{\tilde{Q}_2(U, U)} &= \sum_{\substack{n \in \mathbb{Z}^3 \\ n_h \neq 0}} \left\langle \mathcal{F}\tilde{Q}_2(U, U)(n) \mid e_0(n) \right\rangle_{\mathbb{C}^4} e^{i\tilde{n} \cdot x} e_0(n). \end{aligned}$$

The projection of $\tilde{Q}_1(\tilde{U}, \tilde{U})$ is a mere horizontal transport interaction of elements in the kernel of $\mathbb{P}\mathcal{A}$ as it is showed in the following lemma, the proof of which can be found in [40, Lemma 4.2].

Lemma 5.4. *The following identity holds true*

$$\overline{\tilde{\mathcal{Q}}_1(\tilde{U}, \tilde{U})} = \begin{pmatrix} \bar{u}^h \cdot \nabla_h \bar{u}^h \\ 0 \\ 0 \end{pmatrix} + \begin{pmatrix} \nabla_h \bar{p}_1 \\ 0 \\ 0 \end{pmatrix},$$

where

$$\begin{aligned} \bar{u}^h &= \nabla_h^\perp (-\Delta_h)^{-1} (-\partial_2 U^1 + \partial_1 U^2), \\ \bar{p}_1 &= (-\Delta_h)^{-1} \operatorname{div}_h \operatorname{div}_h (\bar{u}^h \otimes \bar{u}^h). \end{aligned}$$

For the projection of $\tilde{\mathcal{Q}}_2(U, U)$, we remark that the matrix $\widehat{\mathbb{P}}(n)$ is real and symmetric, so we can write

$$\begin{aligned} \mathcal{F}\tilde{\mathcal{Q}}_2(U, U)(n) &= \mathcal{F}\tilde{\mathcal{Q}}_2(\underline{U}, \tilde{U})(n) \\ &= 2 \sum_{\substack{(0, k_3) + m = n \\ m_h, n_h \neq 0 \\ \tilde{\omega}_{m, n}^{b, c} = 0 \\ b, c = 0, \pm \\ j = 1, 2, 3}} \left\langle \widehat{\mathbb{P}}(n) \begin{pmatrix} \tilde{n} \\ 0 \end{pmatrix} \cdot \mathbb{S}(U^j(0, k_3) \otimes U^b(m)) \Big| e_c(n) \right\rangle_{\mathbb{C}^4} e_c(n), \end{aligned}$$

whence $\tilde{\mathcal{Q}}_2$ in (5.2) acts as a non-local transport between the vectors \underline{U} and \tilde{U} . We have

Lemma 5.5. *Let U be as in Proposition 4.2, and let $\tilde{\mathcal{Q}}_2$ be defined as in (4.2). Then,*

$$\sum_{\substack{n \in \mathbb{Z}^3 \\ n_h \neq 0}} \left\langle \mathcal{F}\tilde{\mathcal{Q}}_2(\underline{U}, \tilde{U})(n) \Big| e_0(n) \right\rangle_{\mathbb{C}^4} e_0(n) = \begin{pmatrix} \underline{u}^h \cdot \nabla_h \bar{u}^h \\ 0 \\ 0 \end{pmatrix} + \begin{pmatrix} \nabla_h \bar{p}_2 \\ 0 \\ 0 \end{pmatrix},$$

where

$$\bar{p}_2 = (-\Delta_h)^{-1} \operatorname{div}_h (\underline{u}^h \cdot \nabla_h \bar{u}^h).$$

Proof. For $c = \pm$, from (2.5), we have $e^c \perp e_0$, which implies that

$$\begin{aligned} &\left\langle \mathcal{F}\tilde{\mathcal{Q}}_2(\underline{U}, \tilde{U})(n) \Big| e_0(n) \right\rangle_{\mathbb{C}^4} \\ &= 2 \sum_{\substack{(0, k_3) + m = n \\ m_h, n_h \neq 0 \\ \tilde{\omega}_{m, n}^{b, 0} = 0 \\ b = 0, \pm \\ j = 1, 2, 3}} \left\langle \widehat{\mathbb{P}}(n) \begin{pmatrix} \tilde{n} \\ 0 \end{pmatrix} \cdot \mathbb{S}(U^j(0, k_3) \otimes U^b(m)) \Big| e_0(n) \right\rangle_{\mathbb{C}^4}. \end{aligned}$$

The condition $\tilde{\omega}_{m,n}^{b,0} = 0$ implies that

$$\tilde{\omega}_{m,n}^{b,0} = \omega^b(m) - \omega^0(n) = \pm i \frac{|m_h|}{|m|} = 0,$$

hence $m_h = 0$. This consideration combined with the convolution constraint $(0, k_3) + m = n$ implies $n_h = 0$, which contradicts the definition of the form $\tilde{\mathcal{Q}}_2$. Then, in the expression of $\mathcal{F}\tilde{\mathcal{Q}}_2(\underline{U}, \tilde{U})$, we should only take $c = 0$ and Lemma 5.5 follows standard explicit computations. \square

Now, setting

$$\bar{p} = \bar{p}_1 + \bar{p}_2,$$

Lemma 5.4 and 5.5 imply the following

Corollary 5.6. *Let U be as in Proposition 4.2, and let $\tilde{\mathcal{Q}}_1$ and $\tilde{\mathcal{Q}}_2$ be defined as in (4.1) and (4.2) respectively. Then*

$$\begin{aligned} \sum_{\substack{n \in \mathbb{Z}^3 \\ n_h \neq 0}} \left\langle \mathcal{F}\tilde{\mathcal{Q}}_1(\tilde{U}, \tilde{U}) + \mathcal{F}\tilde{\mathcal{Q}}_2(U, U) \mid e_0(n) \right\rangle_{\mathbb{C}^4} e_0(n) \\ = \begin{pmatrix} \bar{u}^h \cdot \nabla_h \bar{u}^h + \underline{u}^h \cdot \nabla_h \bar{u}^h \\ 0 \\ 0 \end{pmatrix} + \begin{pmatrix} \nabla_h \bar{p} \\ 0 \\ 0 \end{pmatrix}. \end{aligned}$$

Step 2: The computation of the projection of $\widetilde{\mathcal{A}}_2^0(D)\tilde{U}$ onto $\text{Span}\{E_0(n, \cdot)\}$ is given in the following lemma, the proof of which can be found in [40, Lemma 4.4].

Lemma 5.7. *Let U be as in Proposition 4.2 and $\widetilde{\mathcal{A}}_2^0(D)$ be defined as in (4.4). Then*

$$\overline{\widetilde{\mathcal{A}}_2^0(D)\tilde{U}} = \sum_{\substack{n \in \mathbb{Z}^3 \\ n_h \neq 0}} \left\langle \mathcal{F}\left(\widetilde{\mathcal{A}}_2^0(D)\tilde{U}\right) \mid e_0(n) \right\rangle_{\mathbb{C}^4} e_0(n) = \begin{pmatrix} \nu \Delta \bar{u}^h \\ 0 \\ 0 \end{pmatrix}.$$

Step 3: Projecting the system (5.2) onto $\text{Span}\{E_0(n, \cdot)\}$ yields the following equation which describes the evolution of \bar{U}

$$\partial_t \bar{U} + \overline{\tilde{\mathcal{Q}}_1(\tilde{U}, \tilde{U})} + \overline{\tilde{\mathcal{Q}}_2(U, U)} - \overline{\widetilde{\mathcal{A}}_2^0(D)\tilde{U}} = 0.$$

Then, Corollary 5.6 and Lemma 5.7 imply

Proposition 5.8. *Let U be as in Proposition 4.2 and let*

$$\bar{V}_0 = \sum_{\substack{n \in \mathbb{Z}^3 \\ n_h \neq 0}} \langle \mathcal{F}V_0 \mid e_0(n) \rangle_{\mathbb{C}^4} e_0(n) \in H^s(\mathbb{T}^3),$$

for $s > 5/2$. Then, the projection \bar{U} of U onto $\text{Span}\{E_0(n, \cdot)\}$ belongs to the energy space

$$\mathcal{C}([0, T]; H^\sigma(\mathbb{T}^3)), \quad \sigma \in (s-2, s),$$

for each $T \in [0, T^*[$, and \bar{U} solves the Cauchy problem (1.10) almost everywhere in $\mathbb{T}^3 \times [0, T]$.

5.2.2. Derivation of the evolution of U_{osc}

As for \bar{U} , the study of U_{osc} also consists in three steps.

Step 1: Computation of

$$\begin{aligned} \left(\tilde{\mathcal{Q}}_1(\tilde{U}, \tilde{U}) \right)_{\text{osc}} &= \sum_{\substack{n \in \mathbb{Z}^3 \\ n_h \neq 0}} \sum_{c=\pm} \left\langle \mathcal{F}\tilde{\mathcal{Q}}_1(\tilde{U}, \tilde{U})(n) \mid e_c(n) \right\rangle_{\mathbb{C}^4} e^{i\tilde{n} \cdot x} e_c(n) \\ \left(\tilde{\mathcal{Q}}_2(\underline{U}, \tilde{U}) \right)_{\text{osc}} &= \sum_{\substack{n \in \mathbb{Z}^3 \\ n_h \neq 0}} \sum_{c=\pm} \left\langle \mathcal{F}\tilde{\mathcal{Q}}_2(\underline{U}, \tilde{U})(n) \mid e_c(n) \right\rangle_{\mathbb{C}^4} e^{i\tilde{n} \cdot x} e_c(n). \end{aligned}$$

Since $\tilde{U} = \bar{U} + U_{\text{osc}}$, we can decompose

$$\left(\tilde{\mathcal{Q}}_1(\tilde{U}, \tilde{U}) \right)_{\text{osc}} = \left(\tilde{\mathcal{Q}}_1(\bar{U}, \bar{U}) \right)_{\text{osc}} + 2 \left(\tilde{\mathcal{Q}}_1(\bar{U}, U_{\text{osc}}) \right)_{\text{osc}} + \left(\tilde{\mathcal{Q}}_1(U_{\text{osc}}, U_{\text{osc}}) \right)_{\text{osc}}.$$

The first term was already calculated in [40, Lemma 4.6], and we have

Lemma 5.9. *The following identity holds true*

$$\left(\tilde{\mathcal{Q}}_1(\bar{U}, \bar{U}) \right)_{\text{osc}} = 0.$$

To obtain the bilinear term of the equation (1.11), it remains to find the explicit expression of $\left(\tilde{\mathcal{Q}}_2(\underline{U}, \tilde{U}) \right)_{\text{osc}}$, which is in fact the bilinear term $\mathcal{B}(\underline{U}, U_{\text{osc}})$ of the equation (1.11).

Lemma 5.10. *We have the following explicit expression*

$$\begin{aligned} (5.13) \quad \mathcal{FB}(\underline{U}, U_{\text{osc}}) &= \mathcal{F} \left(\tilde{\mathcal{Q}}_2(\underline{U}, \tilde{U}) \right)_{\text{osc}} \\ &= \sum_{\substack{b, c=\pm \\ j=1,2,3}} \left\langle \hat{\mathbb{P}}(n) \begin{pmatrix} \tilde{n} \\ 0 \end{pmatrix} \cdot \mathbb{S}(U^j(0, 2n_3) \otimes U^b(n_h, -n_3)) \mid e_c(n) \right\rangle_{\mathbb{C}^4} e_c(n). \end{aligned}$$

Proof. Since $\tilde{U} = \bar{U} + U_{\text{osc}}$, we have

$$\left(\tilde{\mathcal{Q}}_2(\underline{U}, \tilde{U})\right)_{\text{osc}} = \left(\tilde{\mathcal{Q}}_2(\underline{U}, \bar{U})\right)_{\text{osc}} + \left(\tilde{\mathcal{Q}}_2(\underline{U}, U_{\text{osc}})\right)_{\text{osc}}.$$

According to the definition (4.2), we can write

$$\begin{aligned} & \left(\tilde{\mathcal{Q}}_2(\underline{U}, \bar{U})\right)_{\text{osc}} \\ &= \sum_{\substack{(0, k_3) + m = n \\ m_h, n_h \neq 0 \\ \tilde{\omega}_{m, n}^{0, c} = 0 \\ c = \pm \\ j = 1, 2, 3}} \left\langle \hat{\mathbb{P}}(n) \begin{pmatrix} \tilde{n} \\ 0 \end{pmatrix} \cdot \mathbb{S}(U^j(0, k_3) \otimes U^0(m)) \Big| e_c(n) \right\rangle_{\mathbb{C}^4} e_c(n). \end{aligned}$$

Let us remark that the above formulation differs to the one given in (4.2) since the projection onto the oscillating subspace forces the parameter c to be equal to \pm only, and the fact that \bar{U} is the second argument of the bilinear form forces the parameter b in (4.2) to be zero. Hence the bilinear interaction constraint $\tilde{\omega}_{m, n}^{0, c} = c \frac{|n_h|}{|n|} = 0$ combined with the convolution constraint $(0, k_3) + m = n$ implies that $n_h = 0$, that contradicts the definition of $\tilde{\mathcal{Q}}_2$. We deduce that

$$\left(\tilde{\mathcal{Q}}_2(\underline{U}, \bar{U})\right)_{\text{osc}} = 0.$$

It now remains to prove (5.13). According to the above argument we can argue that

$$\begin{aligned} & \left(\tilde{\mathcal{Q}}_2(\underline{U}, \tilde{U})\right)_{\text{osc}} = \left(\tilde{\mathcal{Q}}_2(\underline{U}, U_{\text{osc}})\right)_{\text{osc}} \\ &= \sum_{\substack{(0, k_3) + m = n \\ \tilde{\omega}_{m, n}^{b, c} = 0 \\ b, c = \pm \\ j = 1, 2, 3}} \left(\hat{\mathbb{P}}(n) \begin{pmatrix} \tilde{n} \\ 0 \end{pmatrix} \cdot \mathbb{S}(U^j(0, k_3) \otimes U^b(m)) \Big| e_c(n) \right)_{\mathbb{C}^4} e_c(n). \end{aligned}$$

In this case, $n_h = m_h$ and using (2.4), the equality

$$\tilde{\omega}_{m, n}^{b, c} = \omega^\pm(n_h, n_3) - \omega^\pm(n_h, m_3) = 0$$

becomes

$$\frac{|\tilde{n}_h|}{\sqrt{\tilde{n}_1^2 + \tilde{n}_2^2 + \tilde{m}_3^2}} = \frac{|\tilde{n}_h|}{\sqrt{\tilde{n}_1^2 + \tilde{n}_2^2 + \tilde{n}_3^2}}.$$

The above equality is satisfied if $m_3 = \pm n_3$. Let us suppose $m_3 = n_3$, if this is the case the convolution condition $k_3 + m_3 = n_3$ implies that $k_3 = 0$, in this case the term $\hat{U}(0, 0)$ denotes the average of the element u^h , which is identically zero by hypothesis since (PBS_ε) propagates the global average which is supposed to be zero since the beginning. Thus, we get $m_3 = -n_3$ and $k_3 = 2n_3$ and we recover the expression in (5.13). □

Step 2: Computation of $\left(\widetilde{\mathcal{A}}_2^0(D)\widetilde{U}\right)_{\text{osc}}$.

Lemma 5.11. *We have*

$$\left(\widetilde{\mathcal{A}}_2^0(D)\widetilde{U}\right)_{\text{osc}} = \nu\Delta U_{\text{osc}}.$$

Proof. We will calculate

$$\mathcal{F}\left(\widetilde{\mathcal{A}}_2^0(D)U_{\text{osc}}\right)_{\text{osc}}(n) = \sum_{\substack{a,b=\pm \\ \omega_n^{a,b}=0}} \widehat{U}^a(n) \langle \mathcal{A}_2(n)e_a(n) \mid e_b(n) \rangle_{\mathbb{C}^4} e_b(n),$$

where the matrix \mathcal{A}_2 is defined in (1.2).

If $a = -b$, the condition $\omega_n^{a,b} = 0$ becomes

$$\omega_n^{a,-a} = 2\omega^a(n) = 2a i \frac{|n_h|}{|n|} = 0,$$

which implies $n_h = 0$, contradicting the definition of $\widetilde{\mathcal{A}}_2^0$. Thus, we deduce that $a = b$. The expression of the eigenvalues given in (2.4) implies

$$\langle \mathcal{A}_2(D)e_a \mid e_a \rangle_{\mathbb{C}^4} = -\nu|n|^2.$$

Lemma 5.11 is then proved. □

Remark 5.12. We want to emphasize that Lemma 5.7 and 5.11 imply the strict (total) parabolicity of the operator $\widetilde{\mathcal{A}}_2^0(D)$ for vector fields with zero horizontal average. This is remarkable since the operator $\mathcal{A}_2(D)$ appearing in (PBS_ε) is *not* strictly parabolic.

Step 3: Projecting the system (5.2) onto $\text{Span}\{E_{\pm}(n, \cdot)\}$ and using Lemma 5.9 and 5.11, we deduce that the evolution of U_{osc} is given by

$$\partial_t U_{\text{osc}} + \widetilde{\mathcal{Q}}_1(U_{\text{osc}}, U_{\text{osc}} + 2\bar{U}) + \mathcal{B}(\underline{U}, U_{\text{osc}}) - \nu\Delta U_{\text{osc}} = 0,$$

and we hence prove the following result

Proposition 5.13. *Let $U_{\text{osc},0} \in H^s(\mathbb{T}^3)$, $s > 5/2$, the projection of U onto the oscillating subspace $\text{Span}\{E_{\pm}(n, \cdot)\}$ belongs to the energy space*

$$\mathcal{C}([0, T]; H^\sigma(\mathbb{T}^3)), \quad \sigma \in (s - 2, s),$$

for each $T \in [0, T^*[$, and U_{osc} solves the Cauchy problem (1.11) in $\mathbb{T}^3 \times [0, T]$.

6. Global propagation of smooth data for the limit system

We proved in Propositions 5.1, 5.8 and 5.13, if the initial data $U_0 \in H^s$, $s > 5/2$, then the decomposition $U = \underline{U} + \overline{U} + U_{\text{osc}}$ holds in $\mathcal{C}([0, T], H^\sigma)$, $s - 2 < \sigma < s$ and $0 \leq T < T^*$, and where \underline{U} , \overline{U} and U_{osc} are respectively solutions of the systems (1.9), (1.10) and (1.11). The aim of this section is to prove the *global* propagation of the $H^s(\mathbb{T}^3)$ -regularity, $s > 5/2$, by the limit system (S_0) , more precisely by the systems (1.9)–(1.11). This propagation can be resumed in the following propositions. We remark that for \underline{U} and \overline{U} , we need much less regularity, and the $H^s(\mathbb{T}^3)$ -regularity, $s > 5/2$, is especially needed for U_{osc} .

Proposition 6.1. *Let $U_0 \in H^s(\mathbb{T}^3)$, $s \geq 0$, then the solution $\underline{U} = (\underline{u}^h, 0, \underline{U}^4)$ of the equation (1.9) globally and uniquely exists in time variable*

$$\underline{u}^h \in \mathcal{C}(\mathbb{R}_+; H^s(\mathbb{T}_v^1)) \cap L^2(\mathbb{R}_+; H^{s+1}(\mathbb{T}_v^1)),$$

and

$$\underline{U}^4 \in \mathcal{C}(\mathbb{R}_+; H^s(\mathbb{T}_v^1)).$$

Proposition 6.2. *Let $s > 1/2$, $\sigma > 0$ and $U_0 \in H^s(\mathbb{T}^3) \cap L^\infty(\mathbb{T}_v; H^\sigma(\mathbb{T}_h^2))$ such that $\nabla_h U_0 \in L^\infty(\mathbb{T}_v; H^\sigma(\mathbb{T}_h^2))$. Then the system (1.10) possesses a unique solution in*

$$\overline{u}^h \in \mathcal{C}(\mathbb{R}_+; H^s(\mathbb{T}^3)) \cap L^2(\mathbb{R}_+; H^{s+1}(\mathbb{T}^3)).$$

Moreover for each $t > 0$ the following estimate holds true

$$(6.1) \quad \|\overline{u}^h(t)\|_{H^s(\mathbb{T}^3)}^2 + \nu \int_0^t \|\overline{u}^h(\tau)\|_{H^{s+1}(\mathbb{T}^3)}^2 d\tau \leq \mathcal{E}_1(U_0),$$

where

$$(6.2) \quad \mathcal{E}_1(U_0) = C \|\overline{u}_0^h\|_{H^s(\mathbb{T}^3)}^2 \exp \left\{ \frac{CK \Phi(U_0)}{c\nu} \|\nabla_h \overline{u}_0^h\|_{L_v^p(H_h^\sigma)} + \frac{C}{\nu} \|\underline{u}_0^h\|_{H^s(\mathbb{T}_v^1)}^2 \right\}$$

and

$$\Phi(U_0) = \exp \left\{ \frac{CK^2 \|\nabla_h \overline{u}_0^h\|_{L_v^\infty(L_h^2)}^2}{c\nu} \right. \\ \left. \times \exp \left\{ \frac{K}{c\nu} \left(1 + \|\overline{u}_0^h\|_{L_v^\infty(L_h^2)}^2 \right) \|\nabla_h \overline{u}_0^h\|_{L_v^\infty(L_h^2)}^2 \right\} \right\}.$$

Proposition 6.3. *Let $s > 5/2$ and $U_0 \in H^s(\mathbb{T}^3)$. For each $T > 0$, we have*

$$U_{\text{osc}} \in \mathcal{C}([0, T]; H^s(\mathbb{T}^3)) \cap L^2([0, T]; H^{s+1}(\mathbb{T}^3)),$$

and the following bound holds true for each $0 \leq t \leq T$

$$\|U_{\text{osc}}(t)\|_{H^s(\mathbb{T}^3)}^2 + \nu \int_0^t \|\nabla U_{\text{osc}}(\tau)\|_{H^s(\mathbb{T}^3)}^2 d\tau \leq \mathcal{E}_{3,\nu,T}(U_0),$$

where

$$(6.3) \quad \mathcal{E}_{3,\nu,T}(U_0) = \|U_{\text{osc},0}\|_{H^s}^2 \exp \left\{ \frac{\mathcal{E}_1(U_0)}{\nu} + T \|\underline{U}_0\|_{L^2(\mathbb{T}_v^1)}^2 + \frac{(\mathcal{E}_{2,U_0}(T))^2}{\nu} \right\},$$

and

$$(6.4) \quad \mathcal{E}_{2,U_0}(T) = C \|U_{\text{osc},0}\|_{L^2(\mathbb{T}^3)}^2 \exp \left\{ \frac{\mathcal{E}_1(U_0)}{\nu} + T \|\underline{U}_0\|_{H^s(\mathbb{T}_v^1)}^2 \right\}$$

and where $\mathcal{E}_1(U_0)$ is defined as in Proposition 6.2.

6.1. Proof of Proposition 6.1

The system (1.10) is a classical heat equation, the solution of which is well known in the literature. Here, we only remark that classical energy estimates imply

$$\|\underline{u}^h(t)\|_{H^s(\mathbb{T}_v^1)}^2 + 2\nu \int_0^t \|\partial_3 \underline{u}^h(\tau)\|_{H^s(\mathbb{T}_v^1)}^2 = \|\underline{u}_0^h\|_{H^s(\mathbb{T}_v^1)}^2.$$

Since \underline{u}^h has zero vertical average, $\underline{u}^h \in L^2(\mathbb{R}_+; H^{s+1}(\mathbb{T}_v^1))$ as well. \square

Remark 6.4. We would like to mention that

$$(6.5) \quad \|\underline{u}^h\|_{H^s(\mathbb{T}_v^1)} = \|\underline{u}^h\|_{H^s(\mathbb{T}^3)},$$

hence even if \underline{u}^h depends on the vertical variable only it still inherits the same isotropic regularity.

6.2. Proof of Proposition 6.2

We start by recalling a result proved in [40, Proposition 5]

Proposition 6.5. *Let \bar{u}^h be a solution of (1.10) with initial data \bar{u}_0^h and $\nabla_h \bar{u}_0^h$ belonging to $L_v^\sigma(H_h^\sigma)$, for some $\sigma \geq 1$. Then, we have*

$$\bar{u}^h \in L^2(\mathbb{R}_+; L^\infty(\mathbb{T}^3)),$$

and in particular

$$\|\bar{u}^h\|_{L^2(\mathbb{R}_+; L^\infty(\mathbb{T}^3))} \leq \frac{CK}{c\nu} \Phi(U_0) \|\nabla_h \bar{u}_0^h\|_{L_v^p(H_h^\sigma)},$$

where $\Phi(U_0)$ is defined as in Proposition 6.2 and c, C, K are positive constants.

Remark 6.6. The reader may notice that [40, Proposition 5] is applied on a limit system which is slightly different than (1.10), i.e. on the system

$$(6.6) \quad \begin{cases} \partial_t \bar{u}^h + \bar{u}^h \cdot \nabla_h \bar{u}^h - \nu \Delta \bar{u}^h = -\nabla \bar{p}, \\ \operatorname{div} \bar{u}^h = 0. \end{cases}$$

The only difference between (6.6) and (1.10) is the presence in (1.10) of the term $\underline{u}^h \cdot \nabla_h \bar{u}^h$. Such term though does not pose an obstruction to the application of [40, Proposition 5] to the limit system (1.10); the proof of such result is in fact based on the fact that the following nonlinear cancellation

$$\int_{\mathbb{R}_h^2} (\bar{u}^h \cdot \nabla_h \bar{u}^h) \cdot \bar{u}^h dy_h = 0,$$

holds true for (6.6) (and hence as well for (1.10)) since $\operatorname{div} \bar{u}^h = 0$. Indeed, the term $\underline{u}^h \cdot \nabla_h \bar{u}^h$ enjoys as well a nonlinear cancellation, since

$$\int_{\mathbb{R}_h^2} (\underline{u}^h \cdot \nabla_h \bar{u}^h) \cdot \bar{u}^h dy_h = \frac{1}{2} \underline{u}^h \int_{\mathbb{R}_h^2} \nabla_h |\bar{u}^h|^2 dy_h = 0,$$

being the vector field periodic. Whence [40, Proposition 5] can be applied to the limit system (1.10).

Next, we need the following estimate

Lemma 6.7. *Let \bar{u}^h be the solution of (1.10) and \underline{u}^h the solution of (1.9), then, for $s > 1/2$, we have*

$$(6.7) \quad \left| \langle \underline{u}^h \cdot \nabla_h \bar{u}^h \mid \bar{u}^h \rangle_{H^s(\mathbb{T}^3)} \right| \leq C \left(\|\underline{u}^h\|_{H^s(\mathbb{T}_v^1)} + \|\underline{u}^h\|_{H^{s+1}(\mathbb{T}_v^1)} \right) \|\bar{u}^h\|_{H^s(\mathbb{T}^3)} \|\nabla_h \bar{u}^h\|_{H^s(\mathbb{T}^3)}.$$

Proof. Applying the dyadic cut-off operator Δ_q to $\underline{u}^h \cdot \nabla_h \bar{u}^h$, taking the L^2 -scalar product of the obtain quantity with $\Delta_q \bar{u}^h$ and applying the Bony decomposition, we get

$$\left| \langle \Delta_q (\underline{u}^h \cdot \nabla_h \bar{u}^h) \mid \Delta_q \bar{u}^h \rangle_{L^2(\mathbb{T}^3)} \right| \leq B_q^1 + B_q^2,$$

where

$$\begin{aligned} B_q^1 &= \sum_{|q-q'| \leq 4} \left| \langle \Delta_q (S_{q'-1} \underline{u}^h \Delta_{q'} \nabla_h \bar{u}^h) \mid \Delta_q \bar{u}^h \rangle_{L^2} \right| \\ B_q^2 &= \sum_{q' > q-4} \left| \langle \Delta_q (\Delta_{q'} \underline{u}^h S_{q'+2} \nabla_h \bar{u}^h) \mid \Delta_q \bar{u}^h \rangle_{L^2} \right|. \end{aligned}$$

Applying Hölder inequality and using (A.3) on the term B_q^1 , we deduce

$$B_q^1 \leq C b_q 2^{-2qs} \|S_{q'-1} \underline{u}^h\|_{L^\infty} \|\nabla_h \bar{u}^h\|_{H^s(\mathbb{T}^3)} \|\bar{u}^h\|_{H^s(\mathbb{T}^3)}.$$

Since \underline{u}^h only depends on the vertical variable, thanks to the embedding

$$H^s(\mathbb{T}_v^1) \hookrightarrow L^\infty(\mathbb{T}_v^1), \quad \forall s > 1/2,$$

we deduce

$$\|S_{q'-1}\underline{u}^h\|_{L^\infty} \leq \|\underline{u}^h\|_{L^\infty(\mathbb{T}_v^1)} \leq \|\underline{u}^h\|_{H^s(\mathbb{T}_v^1)},$$

and whence,

$$(6.8) \quad B_q^1 \leq C b_q 2^{-2qs} \|\underline{u}^h\|_{H^s(\mathbb{T}_v^1)} \|\nabla_h \bar{u}^h\|_{H^s(\mathbb{T}^3)} \|\bar{u}^h\|_{H^s(\mathbb{T}^3)}.$$

Next, we apply Hölder inequality to the term B_q^2 and get

$$B_q^2 \leq \sum_{q' > q-4} \|\Delta_q \bar{u}^h\|_{L^2(\mathbb{T}^3)} \|\Delta_{q'} \underline{u}^h\|_{L_v^2(L_h^\infty)} \|\nabla_h \bar{u}^h\|_{L_v^\infty(L_h^2)}.$$

Bernstein inequality (given in Lemma [Appendix A.1](#)) and Estimates [\(A.3\)](#) and [\(6.5\)](#) yield

$$\|\Delta_{q'} \underline{u}^h\|_{L_v^2(L_h^\infty)} \leq C c_q 2^{q'-(q'+1)s} \|\underline{u}^h\|_{H^{s+1}(\mathbb{T}^3)} = C c_q 2^{-q's} \|\underline{u}^h\|_{H^{s+1}(\mathbb{T}_v^1)}.$$

Since $H^s(\mathbb{T}^3) \hookrightarrow H^{0,s} \hookrightarrow L_v^\infty(L_h^2)$, $s > 1/2$, we have

$$\|\nabla_h \bar{u}^h\|_{L_v^\infty(L_h^2)} \leq C \|\nabla_h \bar{u}^h\|_{H^s(\mathbb{T}^3)}.$$

Applying once again Estimate [\(A.3\)](#), we deduce

$$(6.9) \quad B_q^2 \leq C b_q 2^{-2qs} \|\underline{u}^h\|_{H^{s+1}(\mathbb{T}_v^1)} \|\nabla_h \bar{u}^h\|_{H^s(\mathbb{T}^3)} \|\bar{u}^h\|_{H^s(\mathbb{T}^3)}.$$

Now, combining [\(6.8\)](#) and [\(6.9\)](#) finally implies

$$\begin{aligned} & \left| \langle \Delta_q (\underline{u}^h \cdot \nabla_h \bar{u}^h) \mid \Delta_q \bar{u}^h \rangle_{L^2(\mathbb{T}^3)} \right| \\ & \leq C b_q 2^{-2qs} \left(\|\underline{u}^h\|_{H^s(\mathbb{T}_v^1)} + \|\underline{u}^h\|_{H^{s+1}(\mathbb{T}_v^1)} \right) \|\bar{u}^h\|_{H^s(\mathbb{T}^3)} \|\nabla_h \bar{u}^h\|_{H^s(\mathbb{T}^3)}. \end{aligned}$$

□

Proof of Proposition 6.2. We multiply [\(1.10\)](#) by $(-\Delta)^s \bar{u}^h$, integrate the obtained quantity over \mathbb{T}^3 . Using Inequality [\(6.7\)](#) and the following inequality

$$\left| \langle \bar{u}^h \cdot \nabla_h \bar{u}^h \mid \bar{u}^h \rangle_{H^s(\mathbb{T}^3)} \right| \leq C \|\bar{u}^h\|_{L^\infty} \|\bar{u}^h\|_{H^s(\mathbb{T}^3)} \|\nabla \bar{u}^h\|_{H^s(\mathbb{T}^3)},$$

we deduce that

$$(6.10) \quad \begin{aligned} & \frac{1}{2} \frac{d}{dt} \|\bar{u}^h\|_{H^s(\mathbb{T}^3)}^2 + \nu \|\bar{u}^h\|_{H^{s+1}(\mathbb{R}^3)}^2 \\ & \leq C \left(\|\bar{u}^h\|_{L^\infty(\mathbb{T}^3)} + \|\underline{u}^h\|_{H^s(\mathbb{T}_v^1)} + \|\underline{u}^h\|_{H^{s+1}(\mathbb{T}_v^1)} \right) \|\bar{u}^h\|_{H^s(\mathbb{T}^3)} \|\bar{u}^h\|_{H^{s+1}(\mathbb{R}^3)}. \end{aligned}$$

Then, Young inequality implies

$$\begin{aligned} \|\bar{u}^h\|_{L^\infty(\mathbb{T}^3)} \|\bar{u}^h\|_{H^s(\mathbb{T}^3)} \|\bar{u}^h\|_{H^{s+1}(\mathbb{R}^3)} &\leq \frac{\nu}{2} \|\bar{u}^h\|_{H^{s+1}(\mathbb{R}^3)}^2 \\ &+ C \left(\|\bar{u}^h\|_{L^\infty(\mathbb{T}^3)}^2 + \|\underline{u}^h\|_{H^s(\mathbb{T}_\nu^1)}^2 + \|\underline{u}^h\|_{H^{s+1}(\mathbb{T}_\nu^1)}^2 \right) \|\bar{u}^h\|_{H^s(\mathbb{T}^3)}^2, \end{aligned}$$

which, together with (6.10) and Gronwall lemma, leads to

$$\begin{aligned} \|\bar{u}^h(t)\|_{H^s(\mathbb{T}^3)}^2 + \nu \int_0^t \|\bar{u}^h(\tau)\|_{H^{s+1}(\mathbb{T}^3)}^2 d\tau \\ \leq C \|\bar{u}_0^h\|_{H^s(\mathbb{T}^3)}^2 \exp \left\{ \int_0^t \|\bar{u}^h(\tau)\|_{L^\infty(\mathbb{T}^3)}^2 \right. \\ \left. + \|\underline{u}^h(\tau)\|_{H^s(\mathbb{T}_\nu^1)}^2 + \|\underline{u}^h(\tau)\|_{H^{s+1}(\mathbb{T}_\nu^1)}^2 d\tau \right\}. \end{aligned}$$

Using Proposition 6.5 and Proposition 6.1, we finally obtain

$$\begin{aligned} \|\bar{u}^h(t)\|_{H^s(\mathbb{T}^3)}^2 + \nu \int_0^t \|\bar{u}^h(\tau)\|_{H^{s+1}(\mathbb{T}^3)}^2 d\tau \\ \leq C \|\bar{u}_0^h\|_{H^s(\mathbb{T}^3)}^2 \exp \left\{ \frac{CK}{c\nu} \Phi(U_0) \|\nabla_h \bar{u}_0^h\|_{L_v^p(H_h^s)} + \frac{C}{\nu} \|\underline{u}_0^h\|_{H^s(\mathbb{T}_\nu^1)} \right\}, \end{aligned}$$

where Φ is defined as in Proposition 6.2. \square

6.3. Proof of Proposition 6.3

We first remark that, if \bar{U} and \underline{U} are smooth enough, the system (1.11) admits global weak solutions *à la Leray* in the same fashion as for the incompressible Navier-Stokes equations (see [16] for instance).

Lemma 6.8. *Let $s > 1/2$, $\bar{U} \in L^2(\mathbb{R}_+; H^{s+1}(\mathbb{T}^3))$ and $\underline{U} \in L^\infty(\mathbb{R}_+; H^s(\mathbb{T}_\nu^1))$. Then, for any initial data $U_{\text{osc},0} \in L^2(\mathbb{T}^3)$, there exists a global weak solution of the system (1.11) such that*

$$U_{\text{osc}} \in \mathcal{C}(\mathbb{R}_+; L^2(\mathbb{T}^3)) \cap L_{\text{loc}}^\infty(\mathbb{R}_+; L^2(\mathbb{T}^3)) \cap L_{\text{loc}}^2(\mathbb{R}_+; H^1(\mathbb{T}^3)).$$

Moreover for any $t^* \in \mathbb{R}_+$ and for any $0 \leq t \leq t^* < \infty$, the following estimate holds true

$$(6.11) \quad \|U_{\text{osc}}(t)\|_{L^2(\mathbb{T}^3)}^2 + \nu \int_0^t \|\nabla U_{\text{osc}}(\tau)\|_{L^2(\mathbb{T}^3)}^2 d\tau \leq \mathcal{E}_{2,U_0}(t^*).$$

where

$$\mathcal{E}_{2,U_0}(t^*) = C \|U_{\text{osc},0}\|_{L^2(\mathbb{T}^3)}^2 \exp \left\{ \frac{\mathcal{E}_1(U_0)}{\nu} + t^* \|\underline{U}_0\|_{H^s(\mathbb{T}_\nu^1)}^2 \right\}.$$

Proof. We define the frequency cut-off operator

$$J_n W = \sum_{|k| \leq n} \widehat{W}(n) e^{i\tilde{n} \cdot x},$$

and consider the approximate system
(6.12)

$$\begin{cases} \partial_t U_{\text{osc},n} + J_n \tilde{\mathcal{Q}}_1 (U_{\text{osc},n} + 2\bar{U}, U_{\text{osc},n}) + J_n \mathcal{B}(\underline{U}, U_{\text{osc},n}) - \nu \Delta U_{\text{osc},n} = 0, \\ \operatorname{div} U_{\text{osc},n} = 0, \\ U_{\text{osc},n}|_{t=0} = J_n U_{\text{osc},0}. \end{cases}$$

The Cauchy-Lipschitz theorem implies the existence of a local solution for (6.12) in the space

$$U_{\text{osc},n} \in \mathcal{C}([0, T_n]; L_n^2),$$

where

$$L_n^2 = \left\{ f \in L^2(\mathbb{T}^3), \operatorname{supp} \hat{f} \subset \mathcal{B}(0, n) \right\}.$$

Since $U_{\text{osc},n}$ is of divergence-free we deduce that

$$\left\langle J_n \tilde{\mathcal{Q}}_1 (U_{\text{osc},n}, U_{\text{osc},n}) \mid U_{\text{osc},n} \right\rangle_{L^2(\mathbb{T}^3)} = 0.$$

Moreover, using the embedding $H^s \hookrightarrow L^\infty$, for $s > \frac{d}{2}$, we obtain the following inequalities

$$\begin{aligned} \left\langle \tilde{\mathcal{Q}}_1 (\bar{U}, U_{\text{osc},n}) \mid U_{\text{osc},n} \right\rangle &\leq C \|\nabla \bar{U}\|_{H^s(\mathbb{T}^3)} \|U_{\text{osc},n}\|_{L^2(\mathbb{T}^3)} \|\nabla U_{\text{osc},n}\|_{L^2(\mathbb{T}^3)}, \\ \left\langle \mathcal{B}(\underline{U}, U_{\text{osc},n}) \mid U_{\text{osc},n} \right\rangle &\leq C \|\underline{U}\|_{H^s(\mathbb{T}_\downarrow^1)} \|U_{\text{osc},n}\|_{L^2(\mathbb{T}^3)} \|\nabla U_{\text{osc},n}\|_{L^2(\mathbb{T}^3)}, \end{aligned}$$

which yield, for any $t^* \in \mathbb{R}_+$ and $t \in [0, t^*]$,

$$\begin{aligned} \|U_{\text{osc},n}(t)\|_{L^2(\mathbb{T}^3)}^2 + \nu \int_0^t \|\nabla U_{\text{osc},n}(\tau)\|_{L^2(\mathbb{T}^3)}^2 d\tau \\ \leq C \|U_{\text{osc},0}\|_{L^2(\mathbb{T}^3)}^2 \exp \left\{ \int_0^t \|\nabla \bar{U}(\tau)\|_{H^s(\mathbb{T}^3)}^2 + \|\underline{U}(\tau)\|_{H^s(\mathbb{T}_\downarrow^1)}^2 d\tau \right\}, \\ \leq C \|U_{\text{osc},0}\|_{L^2(\mathbb{T}^3)}^2 \exp \left\{ \frac{\mathcal{E}_1(U_0)}{\nu} + t^* \|U_0\|_{H^s(\mathbb{T}_\downarrow^1)}^2 \right\}, \end{aligned}$$

where \mathcal{E}_1 is defined in (6.2). Hence, by a continuation argument, we deduce that $T_n = \infty$ and for each $T > 0$, the sequence $(U_{\text{osc},n})_n$ is uniformly bounded in the space

$$\mathcal{C}([0, T]; L^2(\mathbb{T}^3)) \cap L^2([0, T]; H^1(\mathbb{T}^3)).$$

Standard product rules in Sobolev spaces show that the sequence $(\partial_t U_{\text{osc},n})_n$ is uniformly bounded in the space $L^2([0, T]; H^{-N})$ for $N \in \mathbb{N}$ large enough. Finally, applying Aubin-Lions lemma (see [2]), we deduce that the sequence $(U_{\text{osc},n})_n$ is compact in $L^2([0, T]; L^2)$, and each limit point of $(U_{\text{osc},n})_n$ weakly solves (1.11). □

Remark 6.9. We point out that the above construction of global weak solutions is possible thanks to the presence of the uniformly parabolic smoothing effect on the limit system (1.11), hence the importance of the propagation of parabolicity mentioned in Remark 1.3.

Next, we study the “purely bilinear” interactions of highly oscillating perturbations in (5.2) given by the term

$$\left(\tilde{\mathcal{Q}}_1 (U_{\text{osc}}, U_{\text{osc}}) \right)_{\text{osc}}.$$

Bilinear interactions of the above form, in general, prevent us from obtaining global-in-time energy subcritical and critical estimates. However, as pointed out in Remark 1.4, we can actually prove that the bilinear interaction $\tilde{\mathcal{Q}}_1 (U_{\text{osc}}, U_{\text{osc}})$ is in fact smoother than the vector $U_{\text{osc}} \cdot \nabla U_{\text{osc}}$. To do so, we introduce the following *resonant set*.

Definition 6.10. 1. *The resonant set \mathcal{K}^* is the set of frequencies such that*

$$\begin{aligned} \mathcal{K}^* &= \{ (k, m, n) \in \mathbb{Z}^9, k_h, m_h, n_h \neq 0 \mid \omega^a(k) + \omega^b(m) = \omega^c(n), \\ &\quad k + m = n, (a, b, c) \in \{-, +\} \}, \\ &= \{ (k, n) \in \mathbb{Z}^6, k_h, n_h \neq 0 \mid \omega^a(k) + \omega^b(n - k) = \omega^c(n), \\ &\quad (a, b, c) \in \{-, +\} \}, \end{aligned}$$

where ω^j , $j = \pm$ are the eigenvalues given in (2.4).

2. *The resonant set of the frequency $n : n_h \neq 0$, is defined as*

$$\mathcal{K}_n^* = \{ (k, m) \in \mathbb{Z}^6 \mid \omega^a(k) + \omega^b(m) = \omega^c(n), k + m = n, (a, b, c) \in \{-, +\} \}.$$

The resonant set is introduced in order to express the term $\left(\tilde{\mathcal{Q}}_1 (U_{\text{osc}}, U_{\text{osc}}) \right)_{\text{osc}}$ in a more concise way. Indeed, considering the explicit definition of the bilinear form $\tilde{\mathcal{Q}}_1$ given in (4.1) we can immediately deduce that

$$\left(\tilde{\mathcal{Q}}_1 (U_{\text{osc}}, U_{\text{osc}}) \right)_{\text{osc}} = \mathcal{F}^{-1} (1_{\mathcal{K}^*} \mathcal{F} (U_{\text{osc}} \cdot \nabla U_{\text{osc}})).$$

In other words, the resonant set \mathcal{K}^* is the set of frequencies on which the bilinear interaction $\left(\tilde{\mathcal{Q}}_1 (U_{\text{osc}}, U_{\text{osc}}) \right)_{\text{osc}}$ is localized.

We now define the following Fourier multiplier of order zero

$$\chi_{\mathcal{K}^*} (D) (a \ b) = \mathcal{F}^{-1} (1_{\mathcal{K}^*} \mathcal{F} (a \ b)).$$

We can hence rewrite

$$\left(\tilde{\mathcal{Q}}_1 (U_{\text{osc}}, U_{\text{osc}}) \right)_{\text{osc}} = \text{div} [\chi_{\mathcal{K}^*} (D) (U_{\text{osc}} \otimes U_{\text{osc}})].$$

We state the following technical lemma which is a simple variation of [16, Lemma 6.6, p.150], [35, Lemma 6.4, p.222] or [42, Lemma 8.4]. The proof is based on the fact that, for fixed (k_h, n) , the fiber

$$\mathcal{J}(k_h, n) = \{k_3 \in \mathbb{Z}, (k, n) \in \mathcal{K}^*\}$$

is a finite set.

Lemma 6.11. *Let $\phi_1, \phi_2 \in H^{1/2}(\mathbb{T}^3)$ and $\phi_3 \in L^2(\mathbb{T}^3)$ be vector fields of zero horizontal average on \mathbb{T}_h^2 . Then there exists a constant C which only depends on a_1/a_2 such that*

$$(6.13) \quad \left| \sum_{(k,n) \in \mathcal{K}^*} \widehat{\phi}_1(k) \widehat{\phi}_2(n-k) \widehat{\phi}_3(n) \right| \leq \frac{C}{a_3} \|\phi_1\|_{H^{1/2}(\mathbb{T}^3)} \|\phi_2\|_{H^{1/2}(\mathbb{T}^3)} \|\phi_3\|_{L^2(\mathbb{T}^3)}.$$

Proof. We first prove Lemma 6.11 when $\mathbb{T}^3 = [0, 2\pi]^3$. We write

$$(6.14) \quad \begin{aligned} I_{\mathcal{K}^*} &= \left| \sum_{(k,n) \in \mathcal{K}^*} \widehat{\phi}_1(k) \widehat{\phi}_2(n-k) \widehat{\phi}_3(n) \right| \\ &\leq \sum_{(k_h, n) \in \mathbb{Z}^2 \times \mathbb{Z}^3} \sum_{\{k_3: (k,n) \in \mathcal{K}^*\}} \left| \widehat{\phi}_1(k) \widehat{\phi}_2(n-k) \widehat{\phi}_3(n) \right|, \\ &\leq \sum_{(k_h, n) \in \mathbb{Z}^2 \times \mathbb{Z}^3} \left| \widehat{\phi}_3(n) \right| \sum_{\{k_3: (k,n) \in \mathcal{K}^*\}} \left| \widehat{\phi}_1(k) \right| \left| \widehat{\phi}_2(n-k) \right|. \end{aligned}$$

By Cauchy-Schwarz inequality, we have

$$\begin{aligned} &\sum_{\{k_3: (k,n) \in \mathcal{K}^*\}} \left| \widehat{\phi}_1(k) \right| \left| \widehat{\phi}_2(n-k) \right| \\ &\leq \left(\sum_{\{k_3: (k,n) \in \mathcal{K}^*\}} \left| \widehat{\phi}_1(k) \right|^2 \left| \widehat{\phi}_2(n-k) \right|^2 \right)^{1/2} \left(\sum_{\{k_3: (k,n) \in \mathcal{K}^*\}} 1 \right)^{1/2}. \end{aligned}$$

Now, fixing $(k_h, n) \in \mathbb{Z}^2 \times \mathbb{Z}^3$ there exists only a finite number of resonant modes k_3 , more precisely,

$$(6.15) \quad \#(\{k_3 : (k, n) \in \mathcal{K}^*\}) \leq 8.$$

Indeed, recalling the notations (3.3), we can write explicitly the resonant condition

$$\omega_{k, n-k, n}^{+,+,+} = 0$$

(the same procedure holds for the generic case $\omega_{k, n-k, n}^{a,b,c} = 0, a, b, c \neq 0$) as follows

$$\left(\frac{|k_h|^2}{|k_3|^2 + |k_h|^2} \right)^{1/2} + \left(\frac{|n_h - k_h|^2}{|n_3 - k_3|^2 + |n_h - k_h|^2} \right)^{1/2} = \left(\frac{|n_h|^2}{|n_3|^2 + |n_h|^2} \right)^{1/2}.$$

After some algebraic calculations, the above equation of k_3 (k_h and n being fixed) becomes a polynomial equation of the form

$$R(k_3) = 0,$$

where R is a real polynomial of degree eight, hence (6.15) follows from the fundamental theorem of algebra. Thus,

$$\sum_{\{k_3:(k,n) \in \mathcal{K}^*\}} |\widehat{\phi}_1(k)| |\widehat{\phi}_2(n-k)| \leq \sqrt{8} \left(\sum_{\{k_3:(k,n) \in \mathcal{K}^*\}} |\widehat{\phi}_1(k)|^2 |\widehat{\phi}_2(n-k)|^2 \right)^{1/2},$$

which, combined with Inequality (6.14), gives

$$I_{\mathcal{K}^*} \leq \sqrt{8} \sum_{k_h, n_h} \sum_{n_3} |\widehat{\phi}_3(n)| \left(\sum_{k_3} |\widehat{\phi}_1(k)|^2 |\widehat{\phi}_2(n-k)|^2 \right)^{1/2}.$$

Moreover

$$\begin{aligned} \sum_{n_3} |\widehat{\phi}_3(n)| \left(\sum_{k_3} |\widehat{\phi}_1(k)|^2 |\widehat{\phi}_2(n-k)|^2 \right)^{1/2} \\ \leq \left(\sum_{n_3} |\widehat{\phi}_3(n)|^2 \right)^{1/2} \left(\sum_{n_3, k_3} |\widehat{\phi}_1(k)|^2 |\widehat{\phi}_2(n-k)|^2 \right)^{1/2}, \end{aligned}$$

and hence

$$(6.16) \quad I_{\mathcal{K}^*} \leq \sqrt{8} \sum_{(k_h, n) \in \mathbb{Z}^2 \times \mathbb{Z}^3} \left(\sum_{n_3} |\widehat{\phi}_3(n)|^2 \right)^{1/2} \\ \times \left(\sum_{p_3} |\widehat{\phi}_2(n_h - k_h, p_3)|^2 \right)^{1/2} \left(\sum_{k_3} |\widehat{\phi}_1(k)|^2 \right)^{1/2}.$$

Let us denote at this point the quantities

$$\begin{aligned} \widetilde{\phi}_1(n_h) &= \left(\sum_{n_3} |\widehat{\phi}_1(n)|^2 \right)^{1/2}, \\ \widetilde{\phi}_2(n_h) &= \left(\sum_{n_3} |\widehat{\phi}_2(n)|^2 \right)^{1/2}, \\ \widetilde{\phi}_3(n_h) &= \left(\sum_{n_3} |\widehat{\phi}_3(n)|^2 \right)^{1/2}, \end{aligned}$$

and the following distributions

$$\begin{aligned}\widetilde{\phi}_1(x_h) &= \mathcal{F}_h^{-1} \left(\widehat{\phi}_1 \right) (x_h), \\ \widetilde{\phi}_2(x_h) &= \mathcal{F}_h^{-1} \left(\widehat{\phi}_2 \right) (x_h), \\ \widetilde{\phi}_3(x_h) &= \mathcal{F}_h^{-1} \left(\widehat{\phi}_3 \right) (x_h).\end{aligned}$$

Then, applying Plancherel theorem and the product rules for Sobolev spaces, Inequality (6.16) can be read as

$$\begin{aligned}I_{\mathcal{K}^*} &\leq \left\langle \widetilde{\phi}_1 \widetilde{\phi}_2 \mid \widetilde{\phi}_3 \right\rangle_{L^2(\mathbb{T}_h^2)} \leq \left\| \widetilde{\phi}_1 \widetilde{\phi}_2 \right\|_{L^2(\mathbb{T}_h^2)} \left\| \widetilde{\phi}_3 \right\|_{L^2(\mathbb{T}_h^2)} \\ &\leq \left\| \widetilde{\phi}_1 \right\|_{H^{1/2}(\mathbb{T}_h^2)} \left\| \widetilde{\phi}_2 \right\|_{H^{1/2}(\mathbb{T}_h^2)} \left\| \widetilde{\phi}_3 \right\|_{L^2(\mathbb{T}_h^2)} \\ &= \left\| \phi_1 \right\|_{H^{1/2,0}(\mathbb{T}^3)} \left\| \phi_2 \right\|_{H^{1/2,0}(\mathbb{T}^3)} \left\| \phi_3 \right\|_{L^2(\mathbb{T}^3)}, \\ &\leq \left\| \phi_1 \right\|_{H^{1/2}(\mathbb{T}^3)} \left\| \phi_2 \right\|_{H^{1/2}(\mathbb{T}^3)} \left\| \phi_3 \right\|_{L^2(\mathbb{T}^3)}.\end{aligned}$$

Finally, to lift this argument to a generic torus $\prod_{i=1}^3 [0, 2\pi a_i]$, it suffices to set

$$\Phi(x_1, x_2, x_3) = \phi(a_1 x_1, a_2 x_2, a_3 x_3),$$

and to use the identity

$$\|\Phi\|_{L^2([0,2\pi]^3)} = (a_1 a_2 a_3)^{-1/2} \|\phi\|_{L^2(\prod_{i=1}^3 [0,2\pi a_i])}.$$

□

Remark 6.12. Lemma 6.11 can be applied on U_{osc} , by taking

$$\phi_1 = \phi_2 = \phi_3 = U_{\text{osc}},$$

since the projection on the oscillating subspace defined in (2.9) has zero horizontal average.

Now, we can prove the energy bound required on the problematic trilinear term.

Lemma 6.13. *Let $s > 0$, then*

$$\begin{aligned}(6.17) \quad &\left\langle \left(\widetilde{\mathcal{Q}}_1(U_{\text{osc}}, U_{\text{osc}}) \right)_{\text{osc}} \mid U_{\text{osc}} \right\rangle_{H^s(\mathbb{T}^3)} \\ &\leq C \|U_{\text{osc}}\|_{L^2(\mathbb{T}^3)}^{1/2} \|\nabla U_{\text{osc}}\|_{L^2(\mathbb{T}^3)}^{1/2} \|U_{\text{osc}}\|_{H^s(\mathbb{T}^3)}^{1/2} \|\nabla U_{\text{osc}}\|_{H^s(\mathbb{T}^3)}^{3/2}.\end{aligned}$$

Proof. We recall that, for vector fields whose average is null, the usual Sobolev norm is equivalent to the semi-norm

$$\|f\|_{H^s(\mathbb{T}^3)} \sim \left\| (-\Delta)^{s/2} f \right\|_{L^2(\mathbb{T}^3)}, \quad s \in \mathbb{R},$$

we will always use $\|(-\Delta)^{s/2}(\cdot)\|_{L^2(\mathbb{T}^3)}$ to denote the $H^s(\mathbb{T}^3)$ -norm. First, we have

$$\left\langle \left(\tilde{\mathcal{Q}}_1(U_{\text{osc}}, U_{\text{osc}}) \right)_{\text{osc}} \mid U_{\text{osc}} \right\rangle_{H^s(\mathbb{T}^3)} = \left\langle \tilde{\mathcal{Q}}_1(U_{\text{osc}}, U_{\text{osc}}) \mid U_{\text{osc}} \right\rangle_{H^s(\mathbb{T}^3)}.$$

The definition of $\tilde{\mathcal{Q}}_1$ yields

$$\begin{aligned} \left\langle \tilde{\mathcal{Q}}_1(U_{\text{osc}}, U_{\text{osc}}) \mid U_{\text{osc}} \right\rangle_{H^s} &= - \langle \chi_{\mathcal{K}^*}(D)(U_{\text{osc}} \otimes U_{\text{osc}}) \mid \nabla U_{\text{osc}} \rangle_{H^s} \\ &= - \left\langle (-\Delta)^{s/2}(U_{\text{osc}} \otimes U_{\text{osc}}) \mid (-\Delta)^{s/2} \nabla U_{\text{osc}} \right\rangle_{\chi_{\mathcal{K}^*}}, \end{aligned}$$

where for any ϕ_1, ϕ_2, ϕ_3 as in Lemma 6.11, we set

$$\langle \phi_1 \phi_2, \phi_3 \rangle_{\chi_{\mathcal{K}^*}} = \sum_{(k,n) \in \mathcal{K}^*} \widehat{\phi}_1(k) \widehat{\phi}_2(n-k) \widehat{\phi}_3(n).$$

By a dyadic decomposition, we also have

$$\begin{aligned} &\left| \left\langle (-\Delta)^{s/2}(U_{\text{osc}} \otimes U_{\text{osc}}) \mid (-\Delta)^{s/2} \nabla U_{\text{osc}} \right\rangle_{\chi_{\mathcal{K}^*}} \right| \\ &\quad \sim \sum_q 2^{2qs} \left| \langle \Delta_q(U_{\text{osc}} \otimes U_{\text{osc}}) \mid \Delta_q \nabla U_{\text{osc}} \rangle_{\chi_{\mathcal{K}^*}} \right|. \end{aligned}$$

For each dyadic bloc in the above estimate, using Bony's decomposition, we can write

$$I_q = \left| \langle \Delta_q(U_{\text{osc}} \otimes U_{\text{osc}}) \mid \Delta_q \nabla U_{\text{osc}} \rangle_{\chi_{\mathcal{K}^*}} \right| \leq I_q^1 + I_q^2,$$

where

$$\begin{aligned} I_q^1 &= \sum_{|q-q'| \leq 4} \left| \langle \Delta_q(S_{q'} U_{\text{osc}} \otimes \Delta_{q'} U_{\text{osc}}) \mid \Delta_q \nabla U_{\text{osc}} \rangle_{\chi_{\mathcal{K}^*}} \right| \\ I_q^2 &= \sum_{q' > q-4} \left| \langle \Delta_q(\Delta_{q'} U_{\text{osc}} \otimes S_{q'+2} U_{\text{osc}}) \mid \Delta_q \nabla U_{\text{osc}} \rangle_{\chi_{\mathcal{K}^*}} \right|. \end{aligned}$$

Combining (6.13) with some classical computations with the dyadic blocs finally leads to, for any $k = 1, 2$,

$$I_q^k \leq C 2^{-2qs} b_q \|U_{\text{osc}}\|_{L^2(\mathbb{T}^3)}^{1/2} \|\nabla U_{\text{osc}}\|_{L^2(\mathbb{T}^3)}^{1/2} \|U_{\text{osc}}\|_{H^s(\mathbb{T}^3)}^{1/2} \|\nabla U_{\text{osc}}\|_{H^s(\mathbb{T}^3)}^{3/2},$$

where the sequence $(b_q)_q \in \ell^2$ depends on U_{osc} , concluding the proof. \square

Lemma 6.14. *Let $s > 1/2$, then*

$$(6.18) \quad \left\langle \tilde{\mathcal{Q}}_1(\bar{U}, U_{\text{osc}}) \mid U_{\text{osc}} \right\rangle_{H^s(\mathbb{T}^3)} \leq C \|\nabla \bar{U}\|_{H^s(\mathbb{T}^3)} \|\nabla U_{\text{osc}}\|_{H^s(\mathbb{T}^3)} \|U_{\text{osc}}\|_{H^s(\mathbb{T}^3)},$$

$$(6.19) \quad \langle \mathcal{B}(\underline{U}, U_{\text{osc}}) \mid U_{\text{osc}} \rangle_{H^s(\mathbb{T}^3)} \leq C \|\underline{U}\|_{L^2(\mathbb{T}^3_\downarrow)} \|\nabla U_{\text{osc}}\|_{H^s(\mathbb{T}^3)} \|U_{\text{osc}}\|_{H^s(\mathbb{T}^3)}.$$

Proof. The proof of Lemma 6.14 relies on direct estimates performed on both bilinear terms. For the first one, we have

$$\begin{aligned} \left| \left\langle \tilde{\mathcal{Q}}_1(\bar{U}, U_{\text{osc}}) \mid U_{\text{osc}} \right\rangle_{H^s(\mathbb{T}^3)} \right| &\leq \left| \left\langle \operatorname{div}(\bar{U} \otimes U_{\text{osc}}) \mid U_{\text{osc}} \right\rangle_{H^s(\mathbb{T}^3)} \right|, \\ &\leq \|\bar{U} \otimes U_{\text{osc}}\|_{H^{s+1}(\mathbb{T}^3)} \|U_{\text{osc}}\|_{H^s(\mathbb{T}^3)}, \\ &\leq C \|\nabla \bar{U}\|_{H^s(\mathbb{T}^3)} \|\nabla U_{\text{osc}}\|_{H^s(\mathbb{T}^3)} \|U_{\text{osc}}\|_{H^s(\mathbb{T}^3)}, \end{aligned}$$

where in the last inequality, we used the fact that $H^{s+1}(\mathbb{T}^3)$, $s > 1/2$ is a Banach algebra.

For the second one we use the explicit definition of the limit bilinear form \mathcal{B} given in (5.13) in order to deduce the identity

$$\begin{aligned} \langle \mathcal{B}(\underline{U}, U_{\text{osc}}) \mid U_{\text{osc}} \rangle_{H^s(\mathbb{T}^3)} &= \left\langle (-\Delta)^{s/2} \mathcal{B}(\underline{U}, U_{\text{osc}}) \mid (-\Delta)^{s/2} U_{\text{osc}} \right\rangle_{L^2(\mathbb{T}^3)}, \\ &= \left\langle \mathcal{B}(\underline{U}, (-\Delta)^{s/2} U_{\text{osc}}) \mid (-\Delta)^{s/2} U_{\text{osc}} \right\rangle_{L^2(\mathbb{T}^3)}, \end{aligned}$$

which implies inequality (6.19). □

Proof of Proposition 6.3. We have now all the ingredients to prove Proposition 6.3. Performing rather standard $H^s(\mathbb{T}^3)$ -energy estimates on the equation (1.11) with the energy bounds (6.17), (6.18) and (6.19), we obtain

$$\begin{aligned} &\frac{1}{2} \frac{d}{dt} \|U_{\text{osc}}(t)\|_{H^s(\mathbb{T}^3)}^2 + \nu \int_0^t \|\nabla U_{\text{osc}}(\tau)\|_{H^s(\mathbb{T}^3)}^2 d\tau \\ &\leq C \left(\|\nabla \bar{U}\|_{H^s(\mathbb{T}^3)} + \|\underline{U}\|_{L^2(\mathbb{T}^3_\downarrow)} \right) \|\nabla U_{\text{osc}}\|_{H^s(\mathbb{T}^3)} \|U_{\text{osc}}\|_{H^s(\mathbb{T}^3)} \\ &\quad + \|U_{\text{osc}}\|_{L^2(\mathbb{T}^3)}^{1/2} \|\nabla U_{\text{osc}}\|_{L^2(\mathbb{T}^3)}^{1/2} \|U_{\text{osc}}\|_{H^s(\mathbb{T}^3)}^{1/2} \|\nabla U_{\text{osc}}\|_{H^s(\mathbb{T}^3)}^{3/2} \end{aligned}$$

Then, Young inequality and Gronwall lemma imply

$$\begin{aligned} &\|U_{\text{osc}}(\tau)\|_{H^s(\mathbb{T}^3)}^2 + \nu \int_0^t \|\nabla U_{\text{osc}}(\tau)\|_{H^s(\mathbb{T}^3)}^2 d\tau \\ &\leq \|U_{\text{osc},0}\|_{H^s(\mathbb{T}^3)}^2 \exp \left\{ \int_0^t \|\nabla \bar{u}^h(\tau)\|_{H^s(\mathbb{T}^3)}^2 d\tau + \int_0^t \|\underline{U}(\tau)\|_{L^2(\mathbb{T}^3_\downarrow)}^2 d\tau \right. \\ &\quad \left. + \int_0^t \|U_{\text{osc}}(\tau)\|_{L^2(\mathbb{T}^3)}^2 \|\nabla U_{\text{osc}}(\tau)\|_{L^2(\mathbb{T}^3)}^2 d\tau \right\}. \end{aligned}$$

Thus, using Estimates (6.1), (6.11) and the result in Proposition 6.1, we deduce that, for each $T > 0$, the following bound holds true

$$\begin{aligned} & \|U_{\text{osc}}(\tau)\|_{H^s(\mathbb{T}^3)}^2 + \nu \int_0^t \|\nabla U_{\text{osc}}(\tau)\|_{H^s(\mathbb{T}^3)}^2 d\tau \\ & \leq \|U_{\text{osc},0}\|_{H^s(\mathbb{T}^3)}^2 \exp \left\{ \frac{1}{\nu} \mathcal{E}_1(U_0) + T \|\underline{U}_0\|_{L^2(\mathbb{T}^3)}^2 + \frac{1}{\nu} (\mathcal{E}_{2,U_0}(T))^2 \right\}, \\ & \leq C_\nu \exp \{C_\nu \exp \{C_\nu T\}\}, \end{aligned}$$

where \mathcal{E}_1 and \mathcal{E}_2 are respectively defined in (6.2) and (6.4). \square

7. Convergence as $\varepsilon \rightarrow 0$ and proof of the main result

As in the work [25], the lack of a complete parabolic smoothing effect on the system (PBS_ε) will prevent us to obtain a uniform global-in-time control for U^ε . Nonetheless we will be able to prove that, for each $T > 0$ arbitrary and $\varepsilon > 0$, the solutions of (S_ε) belong to the space $\mathcal{C}(\mathbb{R}_+; H^{s-2}(\mathbb{T}^3)) \cap L_{\text{loc}}^\infty(\mathbb{R}_+; H^{s-2}(\mathbb{T}^3))$ for $s > 9/2$ and converge in the same topology to the global solution of (S_0) . The idea to prove this convergence result is to use the method of Schochet (see [38]), which consists in a smart change of variable, which cancels some perturbations that we cannot control. We will use results and terminology introduced by I. Gallagher in [24] in the context of quasilinear hyperbolic symmetric systems with skew-symmetric singular perturbation.

Let us recall the following definition [24, Definition 1.2]

Definition 7.1. Let $T, \varepsilon_0 > 0$, $p \geq 1$ and $\sigma > d/2$. Let $\vec{k}_q = (k_1, \dots, k_q)$ where $k_i \in \mathbb{Z}^d$ and let

$$|\vec{k}_q| = \max_{1 \leq i \leq q} |k_i|.$$

Then a function $R_{\text{osc}}^\varepsilon(t)$ is said to be (p, σ) -oscillating function if it can be written as

$$R_{\text{osc}}^\varepsilon = \sum_{q=1}^p R_{q,\text{osc}}^\varepsilon(t),$$

where

$$R_{q,\text{osc}}^\varepsilon(t) = \mathcal{F}^{-1} \left(\sum_{\vec{k}_q \in K_q^n} e^{i \frac{t}{\varepsilon} \beta_q(n, \vec{k}_q)} r_0(n, \vec{k}_q) f_1^\varepsilon(t, k_1) \dots f_q^\varepsilon(t, k_q) \right),$$

with

$$K_q^n = \left\{ \vec{k}_q \in \mathbb{Z}^{dq} \mid \sum_{i=1}^q k_i = n \text{ and } \beta_q(n, \vec{k}_q) \neq 0 \right\},$$

and where r_0 and f_i^ε satisfy

- there exist $(\alpha_i)_{i \in \{1, \dots, q\}}$, $\alpha_i \geq 0$ such that

$$r_0 \left(n, \vec{k}_q \right) \leq C \prod_{i=1}^q (1 + |k_i|)^{\alpha_i},$$

- $(\mathcal{F}^{-1} f_i^\varepsilon)_{0 < \varepsilon < \varepsilon_0}$ is uniformly bounded in $\mathcal{C}([0, T]; H^{\sigma + \alpha_i}(\mathbb{T}^d))$, for any $i \in \{1, \dots, q\}$,
- there exists a $\sigma_i > -\sigma$ for which, $(\mathcal{F}^{-1} \partial_t f_i^\varepsilon)_{0 < \varepsilon < \varepsilon_0}$ is uniformly bounded in $\mathcal{C}([0, T]; H^{\sigma_i}(\mathbb{T}^d))$.

The abstract concept in Definition 7.1 is required in order to introduce the following result, see [24, Lemma 2.1] or [25, Lemma 2.1] for more details.

Lemma 7.2. *Let $T > 0$ and $\sigma > \frac{d}{2} + 2$, let $(b^\varepsilon)_\varepsilon$ be a family of functions, bounded in $\mathcal{C}([0, T]; H^\sigma(\mathbb{T}^d))$ and let $a_0^\varepsilon \rightarrow 0$ as $\varepsilon \rightarrow 0$ in $H^{\sigma-1}(\mathbb{T}^d)$. Let \mathcal{Q}^ε , $\mathcal{A}_2^\varepsilon$ be as in (3.1), (3.2), let $R_{\text{osc}}^\varepsilon$ be a $(p, \sigma - 1)$ -oscillating function and finally let $F^\varepsilon \rightarrow 0$ as $\varepsilon \rightarrow 0$ in $\mathcal{C}([0, T]; H^{\sigma-1}(\mathbb{T}^d))$. Then the function a^ε , solution of*

$$\begin{cases} \partial_t a^\varepsilon + \mathcal{Q}^\varepsilon(a^\varepsilon, b^\varepsilon) - \mathcal{A}_2^\varepsilon(D) a^\varepsilon = R_{\text{osc}}^\varepsilon + F^\varepsilon, \\ a^\varepsilon|_{t=0} = a_0^\varepsilon, \end{cases}$$

is an $o_\varepsilon(1)$ in the $\mathcal{C}([0, T]; H^{\sigma-1}(\mathbb{T}^d))$ topology.

Now, to prove our main result, we subtract (S_0) from (S_ε) , and we denote the difference unknown by $W^\varepsilon = U^\varepsilon - U$. Some basic algebra calculations lead to the following difference system

$$(7.1) \quad \begin{cases} \partial_t W^\varepsilon + \mathcal{Q}^\varepsilon(W^\varepsilon, W^\varepsilon + 2U) - \mathcal{A}_2^\varepsilon(D) W^\varepsilon = -(\mathcal{R}_{\text{osc}}^\varepsilon + \mathcal{S}_{\text{osc}}^\varepsilon), \\ \text{div } W^\varepsilon = 0, \\ W^\varepsilon|_{t=0} = 0, \end{cases}$$

where

$$\begin{aligned} \mathcal{R}_{\text{osc}}^\varepsilon &= \mathcal{Q}^\varepsilon(U, U) - \mathcal{Q}(U, U), \\ \mathcal{S}_{\text{osc}}^\varepsilon &= -(\mathcal{A}_2^\varepsilon(D) - \mathcal{A}_2^0(D)) U. \end{aligned}$$

We remark that $\mathcal{R}_{\text{osc}}^\varepsilon$ and $\mathcal{S}_{\text{osc}}^\varepsilon$ are highly oscillating functions which converge to zero in $\mathcal{D}'(\mathbb{T}^3 \times \mathbb{R}_+)$ only. Thanks to the results proved in Section 4, namely Lemma 4.1 and equation (4.6), we can compute the explicit value of $\mathcal{R}_{\text{osc}}^\varepsilon$ and $\mathcal{S}_{\text{osc}}^\varepsilon$ which is given by

$$\mathcal{R}_{\text{osc}}^\varepsilon = \mathcal{R}_{\text{osc,I}}^\varepsilon + \mathcal{R}_{\text{osc,II}}^\varepsilon + \mathcal{R}_{\text{osc,III}}^\varepsilon,$$

and

$$\begin{aligned}
\mathcal{FR}_{\text{osc,I}}^\varepsilon &= \sum_{\substack{\omega_{k,m,n}^{a,b,c} \neq 0 \\ k+m=n \\ k_h, m_h, n_h \neq 0 \\ a,b,c \in \{0, \pm\}}} e^{i \frac{t}{\varepsilon} \omega_{k,m,n}^{a,b,c}} \left(\mathbb{P}_n(n, 0) \cdot \mathbb{S}(U^a(k) \otimes U^b(m)) \Big| e_c(n) \right)_{\mathbb{C}^4} e_c(n), \\
\mathcal{FR}_{\text{osc,II}}^\varepsilon &= 2 \sum_{\substack{(0, k_3) + m = n \\ m_h, n_h \neq 0 \\ \tilde{\omega}_{m,n}^{b,c} \neq 0 \\ b,c=0, \pm \\ j=1,2,3}} e^{i \frac{t}{\varepsilon} \tilde{\omega}_{m,n}^{b,c}} \left(\mathbb{P}_n(n, 0) \cdot \mathbb{S}(U^j(0, k_3) \otimes U^b(m)) \Big| e_c(n) \right)_{\mathbb{C}^4} e_c(n), \\
\mathcal{FR}_{\text{osc,III}}^\varepsilon &= \sum_{\substack{k+m=(0, n_3) \\ k_h, m_h \neq 0 \\ \omega_{k,m}^{a,b} \neq 0 \\ a,b \in \{0, \pm\} \\ j=1,2,3}} e^{i \frac{t}{\varepsilon} \omega_{k,m}^{a,b}} \left(\mathbb{P}_{(0, n_3)}(0, 0, n_3, 0) \cdot \mathbb{S}(\tilde{V}_1^a(k) \otimes \tilde{V}_2^b(m)) \Big| f_j \right)_{\mathbb{C}^4} f_j, \\
\mathcal{FS}_{\text{osc}}^\varepsilon &= 1_{n_h \neq 0} \sum_{\substack{\omega_n^{a,b} \neq 0 \\ a,b=0, \pm}} e^{i \frac{t}{\varepsilon} \omega_n^{a,b}} \left(\mathcal{FA}_2(n) U^a(n) \Big| e_b(n) \right)_{\mathbb{C}^4} e_b(n).
\end{aligned}$$

The following result is immediate.

Proposition 7.3. *Under the assumption of Theorem 1.2 the function $\mathcal{R}_{\text{osc}}^\varepsilon$ is a $(2, s-1)$ -oscillating function, $\mathcal{S}_{\text{osc}}^\varepsilon$ is a $(1, s-2)$ -oscillating function and hence $\mathcal{R}_{\text{osc}}^\varepsilon + \mathcal{S}_{\text{osc}}^\varepsilon$ is a $(2, s-2)$ -oscillating function.*

We can now conclude by applying Lemma 7.2, with $\sigma = s-2$ and with the substitutions

$$\begin{aligned}
a^\varepsilon &= W^\varepsilon, & b^\varepsilon &= W^\varepsilon + 2U, \\
R_{\text{osc}}^\varepsilon &= -(\mathcal{R}_{\text{osc}}^\varepsilon + \mathcal{S}_{\text{osc}}^\varepsilon), & F^\varepsilon &= 0.
\end{aligned}$$

We deduce that for each $T \in [0, T^*]$, the function W^ε is an $o_\varepsilon(1)$ function in $\mathcal{C}([0, T]; H^{s-2})$. Setting hence

$$\widetilde{T}^* = \sup \left\{ t \in [0, T^*] \mid \|U^\varepsilon(t')\|_{H^{s-2}} < K \left(\mathcal{E}_1(V_0) + \mathcal{E}_{3,\nu,T}(V_0) \right), \forall t' \in [0, t] \right\},$$

where \mathcal{E}_1 and $\mathcal{E}_{3,\nu,T}$ are defined in Proposition 6.2 and 6.3, and K is a positive (possibly large) fixed, finite constant. Since $W^\varepsilon = U^\varepsilon - U$ we deduce that for any $t \in [0, \widetilde{T}^*]$

$$\|U^\varepsilon(t)\|_{H^{s-2}} \leq \|U(t)\|_{H^{s-2}} + \|W^\varepsilon(t)\|_{H^{s-2}} \leq \frac{K}{2} \left(\mathcal{E}_1(V_0) + \mathcal{E}_{3,\nu,T}(V_0) \right) + \frac{1}{2},$$

since

$$\begin{aligned} \|U(t)\|_{H^{s-2}} &\leq \|\overline{U}(t)\|_{H^{s-2}} + \|U_{\text{osc}}(t)\|_{H^{s-2}} + \|\underline{U}(t)\|_{H^{s-2}} \\ &\leq \mathcal{E}_1(V_0) + \mathcal{E}_{3,\nu,T}(V_0) + \|\underline{U}_0\|_{H^{s-2}} \\ &\leq \frac{K}{2} \left(\mathcal{E}_1(V_0) + \mathcal{E}_{3,\nu,T}(V_0) \right), \end{aligned}$$

for $K > 4$ and $\|W^\varepsilon(t)\|_{H^{s-2}} \leq 1/2$ thanks to the result of Lemma 7.2. Thus, $\widetilde{T}^* = T^*$, and supposing $T^* < \infty$, we deduce

$$\begin{aligned} \lim_{t \nearrow T^*} \int_0^t \|\nabla U^\varepsilon(t')\|_{L^\infty} dt' &\leq \lim_{t \nearrow T^*} \int_0^t \|U^\varepsilon(t')\|_{H^{s-2}} dt' \\ &\leq K \left(\mathcal{E}_1(V_0) + \mathcal{E}_{3,\nu,T}(V_0) \right) T^* < \infty, \end{aligned}$$

which indeed contradicts (1.6). We conclude that $T^* = \infty$.

Appendix A. Elements of Littlewood-Paley theory.

Appendix A.1. Dyadic decomposition

A tool that has been widely used all along the paper is the theory of Littlewood-Paley, which consists in doing a dyadic cut-off of the frequencies. Let us define the (non-homogeneous) truncation operators as follows:

$$\begin{aligned} \Delta_q u &= \sum_{n \in \mathbb{Z}^3} \hat{u}_n \varphi\left(\frac{|\check{n}|}{2^q}\right) e^{i\check{n} \cdot x}, & \text{for } q \geq 0, \\ \Delta_{-1} u &= \sum_{n \in \mathbb{Z}^3} \hat{u}_n \chi(|\check{n}|) e^{i\check{n} \cdot x}, \\ \Delta_q u &= 0, & \text{for } q \leq -2, \end{aligned}$$

where $u \in \mathcal{S}'(\mathbb{T}^3)$ and \hat{u}_n are the Fourier coefficients of u . The functions φ and χ represent a partition of the unity in \mathbb{R} , which means that are smooth functions with compact support such that

$$\text{supp } \chi \subset B\left(0, \frac{4}{3}\right), \quad \text{supp } \varphi \subset \mathcal{C}\left(\frac{3}{4}, \frac{8}{3}\right),$$

and such that for all $t \in \mathbb{R}$,

$$\chi(t) + \sum_{q \geq 0} \varphi(2^{-q}t) = 1.$$

Let us define further the low frequencies cut-off operator

$$S_q u = \sum_{q' \leq q-1} \Delta_{q'} u.$$

Appendix A.2. Paradiifferential calculus.

The dyadic decomposition is very useful when it comes to study the product of two tempered distributions. We can in fact, at least formally, write for two tempered distributions u and v

$$(A.1) \quad u = \sum_{q \in \mathbb{Z}} \Delta_q u; \quad v = \sum_{q' \in \mathbb{Z}} \Delta_{q'} v; \quad u \cdot v = \sum_{\substack{q \in \mathbb{Z} \\ q' \in \mathbb{Z}}} \Delta_q u \cdot \Delta_{q'} v.$$

Paradiifferential calculus consists in splitting the above sum in three parts, using the following Bony's decomposition (see [5, 6, 13] for the isotropic case and [15, 30] for the anisotropic one)

$$u \cdot v = T_u v + T_v u + R(u, v),$$

where

$$\begin{aligned} T_u v &= \sum_q S_{q-1} u \Delta_q v, \\ T_v u &= \sum_{q'} S_{q'-1} v \Delta_{q'} u, \\ R(u, v) &= \sum_k \sum_{|\nu| \leq 1} \Delta_k u \Delta_{k+\nu} v. \end{aligned}$$

Here,

- The first part concerns the indices (q, q') for which the size of $\text{supp } \mathcal{F}(\Delta_q u)$ is small compared to $\text{supp } \mathcal{F}(\Delta_{q'} v)$.
- The second part contains the indices corresponding to those frequencies of u which are large compared to those of v .
- In the last part $\text{supp } \mathcal{F}(\Delta_{q'} v)$ and $\text{supp } \mathcal{F}(\Delta_q u)$ have comparable sizes.

The following almost orthogonality properties hold

$$\begin{aligned} \Delta_q (S_q a \Delta_{q'} b) &= 0, & \text{if } |q - q'| \geq 5, \\ \Delta_q (\Delta_{q'} a \Delta_{q'+\nu} b) &= 0, & \text{if } q' > q - 4, |\nu| \leq 1, \end{aligned}$$

which implies the following identity

$$(A.2) \quad \Delta_q (u \cdot v) = \sum_{|q-q'| \leq 4} \Delta_q (S_{q'-1} v \Delta_{q'} u) + \sum_{q' > q-4} \Delta_q (S_{q'+2} u \Delta_{q'} v).$$

Using the frequency localization properties of the dyadic blocks, one can characterize the Sobolev norm in the following way

$$(A.3) \quad \|\Delta_q f\|_{L^2(\mathbb{T}^3)} \leq C c_q(f) 2^{-qs} \|f\|_{H^s(\mathbb{T}^3)},$$

with $\{c_q(f)\}$ is a square-summable sequence of positive numbers with

$$\sum_{q \geq -1} c_q(f)^2 = 1.$$

In this paper, we have also been using b_q to denote a sequence of positive numbers such that $\sum_q |b_q| \leq 1$.

The interest in the use of the dyadic decomposition is that the derivative of a function, the Fourier transform of which is localized in frequencies in a ball of size 2^q , acts like the multiplication with the factor 2^q (up to a constant independent of q). This property is given in the following lemma.

Lemma Appendix A.1 (Bernstein-type inequality). *Let u be a function such that $\mathcal{F}u$ is supported in $2^q\mathcal{C}$, where \mathcal{F} denotes the Fourier transform. For all integers k the following relation holds*

$$2^{qk} C^{-k} \|u\|_{L^p(\mathbb{T}^3)} \leq \|(-\Delta)^{k/2} u\|_{L^p(\mathbb{T}^3)} \leq 2^{qk} C^k \|u\|_{L^p(\mathbb{T}^3)}.$$

Let now $r \geq r' \geq 1$ be real numbers. Let $\text{supp}\mathcal{F}u \subset 2^q B$, then

$$\|u\|_{L^r} \leq C \cdot 2^{3q(\frac{1}{r'} - \frac{1}{r})} \|u\|_{L^{r'}}.$$

Let us consider now a function u such that $\mathcal{F}u$ is supported in $2^q\mathcal{C}_h \times 2^{q'}\mathcal{C}_v$. Let us define $D_h = (-\Delta_h)^{1/2}$, $D_3 = |\partial_3|$, then

$$C^{-q-q'} 2^{qs+q's'} \|u\|_{L^p(\mathbb{T}^3)} \leq \|D_h^s D_3^{s'} u\|_{L^p(\mathbb{T}^3)} \leq C^{q+q'} 2^{qs+q's'} \|u\|_{L^p(\mathbb{T}^3)},$$

and given $1 \leq p' \leq p \leq \infty$, $1 \leq r' \leq r \leq \infty$, then

$$\begin{aligned} \|u\|_{L_h^p L_v^r} &\leq C^{q+q'} 2^{2q(\frac{1}{p'} - \frac{1}{p}) + q'(\frac{1}{r'} - \frac{1}{r})} \|u\|_{L_h^{p'} L_v^{r'}}, \\ \|u\|_{L_v^r L_h^p} &\leq C^{q+q'} 2^{2q(\frac{1}{p'} - \frac{1}{p}) + q'(\frac{1}{r'} - \frac{1}{r})} \|u\|_{L_v^{r'} L_h^{p'}}. \end{aligned}$$

For a proof of the above lemma in the anisotropic (hence as well isotropic) setting we refer to the work [30]. For the sake of self-completeness we state the result in both isotropic and anisotropic setting.

Acknowledgments

The research of S.S. is supported by the Basque Government through the BERC 2018-2021 program and by Spanish Ministry of Economy and Competitiveness MINECO through BCAM Severo Ochoa excellence accreditation SEV-2013-0323 and through project MTM2017-82184-R funded by (AEI/FEDER, UE) and acronym "DESFLU".

References

- [1] Alinhac, S., Gérard, P., 1991. Opérateurs pseudo-différentiels et théorème de Nash-Moser. *Savoirs Actuels*, InterEditions, Paris; Éditions du Centre National de la Recherche Scientifique (CNRS), Meudon.
- [2] Aubin, J.P., 1963. Un théorème de compacité. *C. R. Acad. Sci. Paris* 256, 5042–5044.
- [3] Babin, A., Mahalov, A., Nicolaenko, B., 1996. Global splitting, integrability and regularity of 3d euler and navier-stokes equations for uniformly rotating fluids. *European J. Mech. B Fluids* 15, 291–300.
- [4] Babin, A., Mahalov, A., Nicolaenko, B., 2000. Global regularity of 3D rotating Navier-Stokes equations for resonant domains. *Appl. Math. Lett.* 13, 51–57.
- [5] Bahouri, H., Chemin, J.Y., Danchin, R., 2011. Fourier analysis and non-linear partial differential equations. volume 343 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer, Heidelberg.
- [6] Bony, J.M., 1981. Calcul symbolique et propagation des singularités pour les équations aux dérivées partielles non linéaires. *Ann. Sci. École Norm. Sup.* (4) 14, 209–246.
- [7] Charve, F., 2004a. Étude de phénomènes dispersifs en mécanique des fluides géophysiques. Ph.D. thesis. École Polytechnique.
- [8] Charve, F., 2004b. Global well-posedness and asymptotics for a geophysical fluid system. *Comm. Partial Differential Equations* 29, 1919–1940.
- [9] Charve, F., 2005. Convergence of weak solutions for the primitive system of the quasigeostrophic equations. *Asymptot. Anal.* 42, 173–209.
- [10] Charve, F., 2006. Asymptotics and vortex patches for the quasigeostrophic approximation. *J. Math. Pures Appl.* (9) 85, 493–539.
- [11] Charve, F., 2008. Global well-posedness for the primitive equations with less regular initial data. *Ann. Fac. Sci. Toulouse Math.* (6) 17, 221–238.
- [12] Charve, F., Ngo, V.S., 2011. Global existence for the primitive equations with small anisotropic viscosity. *Rev. Mat. Iberoam.* 27, 1–38.
- [13] Chemin, J.Y., 1995. Fluides parfaits incompressibles. volume 230 of *Astérisque*.
- [14] Chemin, J.Y., 1997. À propos d’un problème de pénalisation de type anti-symétrique. *J. Math. Pures Appl.* (9) 76, 739–755.

- [15] Chemin, J.Y., Desjardins, B., Gallagher, I., Grenier, E., 2000. Fluids with anisotropic viscosity. *M2AN Math. Model. Numer. Anal.* 34, 315–335.
- [16] Chemin, J.Y., Desjardins, B., Gallagher, I., Grenier, E., 2006. Mathematical geophysics. An introduction to rotating fluids and the Navier-Stokes equations. volume 32 of *Oxford Lecture Series in Mathematics and its Applications*. The Clarendon Press, Oxford University Press, Oxford.
- [17] Cushman-Roisin, B., Beckers, J.M., 2011. Introduction to geophysical fluid dynamics. Physical and numerical aspects. volume 101 of *International Geophysics Series*. Elsevier/Academic Press.
- [18] Danchin, R., 2000. Global existence in critical spaces for compressible Navier-Stokes equations. *Invent. Math.* 141, 579–614.
- [19] Embid, P.F., Majda, A.J., 1998. Low Froude number limiting dynamics for stably stratified flow with small or finite Rossby numbers. *Geophys. Astrophys. Fluid Dynam.* 87, 1–50.
- [20] Fanelli, F., Gallagher, I., . Asymptotics of fast rotating density-dependent incompressible fluids in two space dimensions. To appear in *Rev. Mat. Iberoam.*
- [21] Feireisl, E., Gallagher, I., Gerard-Varet, D., Novotný, A., 2012a. Multi-scale analysis of compressible viscous and rotating fluids. *Comm. Math. Phys.* 314, 641–670.
- [22] Feireisl, E., Gallagher, I., Novotný, A., 2012b. A singular limit for compressible rotating fluids. *SIAM J. Math. Anal.* 44, 192–205.
- [23] Gallagher, I., 1998a. Applications of Schochet’s methods to parabolic equations. *J. Math. Pures Appl.* (9) 77, 989–1054.
- [24] Gallagher, I., 1998b. Asymptotic of the solutions of hyperbolic equations with a skew-symmetric perturbation. *J. Differential Equations* 150, 363–384.
- [25] Gallagher, I., 2000. A remark on smooth solutions of the weakly compressible periodic Navier-Stokes equations. *J. Math. Kyoto Univ.* 40, 525–540.
- [26] Gallagher, I., Saint-Raymond, L., 2006a. Mathematical study of the beta-plane model: equatorial waves and convergence results. *Mém. Soc. Math. Fr. (N.S.)* , v+116 pp. (2007).
- [27] Gallagher, I., Saint-Raymond, L., 2006b. Weak convergence results for inhomogeneous rotating fluid equations. *J. Anal. Math.* 99, 1–34.
- [28] Grenier, E., 1997. Oscillatory perturbations of the Navier-Stokes equations. *J. Math. Pures Appl.* (9) 76, 477–498.

- [29] Hough, S.S., 1897. On the Application of Harmonic Analysis to the Dynamical Theory of the Tides. Part I. On Laplace's "Oscillations of the First Species," and on the Dynamics of Ocean Currents. *Proceedings of the Royal Society of London* 61, 236–238.
- [30] Iftimie, D., 1999. The 3D Navier-Stokes equations seen as a perturbation of the 2D Navier-Stokes equations. *Bull. Soc. Math. France* 127, 473–517.
- [31] Iwabuchi, T., Mahalov, A., Takada, R., 2016. Stability of time periodic solutions for the rotating Navier-Stokes equations, in: *Recent developments of mathematical fluid mechanics*. Birkhäuser/Springer, Basel. *Adv. Math. Fluid Mech.*, pp. 321–335.
- [32] Iwabuchi, T., Mahalov, A., Takada, R., 2017. Global solutions for the incompressible rotating stably stratified fluids. *Math. Nachr.* 290, 613–631.
- [33] Lee, S., Takada, R., 2017. Dispersive estimates for the stably stratified Boussinesq equations. *Indiana Univ. Math. J.* 66, 2037–2070.
- [34] Ngo, V.S., Scrobogna, S., 2018. Dispersive effects of weakly compressible and fast rotating inviscid fluids. *Discrete Contin. Dyn. Syst.* 38, 749–789.
- [35] Paicu, M., 2004. Étude asymptotique pour les fluides anisotropes en rotation rapide dans le cas périodique. *J. Math. Pures Appl.* (9) 83, 163–242.
- [36] Pedlosky, J., 1987. *Geophysical Fluid Dynamics*. Springer-Verlag.
- [37] Proudman, J., 1916. On the Motion of Solids in a Liquid Possessing Vorticity. *Proceedings of the Royal Society of London Series A* 92, 408–424.
- [38] Schochet, S., 1994. Fast singular limits of hyperbolic PDEs. *J. Differential Equations* 114, 476–512.
- [39] Scrobogna, S., . Global existence and convergence for non-dimensionalized incompressible Navier-Stokes equations in low Froude number regime. Submitted, <https://arxiv.org/abs/1702.07564>.
- [40] Scrobogna, S., 2017a. Derivation of limit equations for a singular perturbation of a 3D periodic Boussinesq system. *Discrete Contin. Dyn. Syst.* 37, 5979–6034.
- [41] Scrobogna, S., 2017b. On some models in geophysical fluids. Ph.D. thesis. Université de Bordeaux.
- [42] Scrobogna, S., 2018. Highly rotating fluids with vertical stratification for periodic data and anisotropic diffusion. *Rev. Mat. Iberoam.* 34, 1–58.
- [43] Stommel, Henry M. and Moore, Dennis W. , 1989. *An Introduction to the Coriolis Force*. Cambridge University Press.

- [44] Takada, R., 2016. Remarks on lower bounds for the maximal existence time to the 3D rotating Euler equations, in: Harmonic analysis and nonlinear partial differential equations. Res. Inst. Math. Sci. (RIMS), Kyoto. RIMS Kôkyûroku Bessatsu, B56, pp. 93–106.
- [45] Takada, R., 2018. Strongly Stratified Limit for the 3D Inviscid Boussinesq Equations. Arch. Ration. Mech. Anal. doi:[10.1007/s00205-018-01347-4](https://doi.org/10.1007/s00205-018-01347-4).
- [46] Taylor, G.I., 1917. Motion of Solids in Fluids When the Flow is Not Irrotational. Proceedings of the Royal Society of London Series A 93, 99–113.
- [47] Widmayer, K., . Convergence to Stratified Flow for an Inviscid 3D Boussinesq System. To appear in Comm. Math. Sci.