In the insurance business risky investments are dangerous: the case of negative risk sums

Yuri Kabanov, Serguei Pergamenshchikov

► To cite this version:

Yuri Kabanov, Serguei Pergamenshchikov. In the insurance business risky investments are dangerous: the case of negative risk sums. *Finance and Stochastics*, 2016, 20, pp.355 - 379. 10.1007/s00780-016-0292-4 . hal-02334871

HAL Id: hal-02334871

<https://hal.science/hal-02334871>

Submitted on 29 Oct 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

In the Life Insurance Business Risky Investments are Dangerous

Yuri Kabanov · Serguei Pergamenshchikov

Received: date / Accepted: date

Dedicated to the memory of Marc Yor.

Abstract We investigate models of the life annuity insurance when the company invests its reserve into a risky asset with price following a geometric Brownian motion. Our main result is an exact asymptotic of the ruin probabilities for the case of exponentially distributed benefits. As in the case of non-life insurance with exponential claims, the ruin probabilities are either decreasing with a rate given by a power function (the case of small volatility) or equal to unit identically (the case of large volatility). The result allows us to quantify the share of reserve to invest into such a risky asset to avoid a catastrophic outcome: the ruin with probability one. We address also the question of smoothness of the ruin probabilities as a function of the initial reserve for generally distributed jumps.

Keywords Life annuity insurance · Financial markets · Smoothness of the exit probability · Exponential functional of a Brownian motion · Autoregression with random coefficients

Mathematics Subject Classification (2000) 60G44

JEL Classification G22 · G23

Laboratoire de Mathématiques, Université de Franche-Comté,
16 Route de Gray, 25030 Besançon, cedex, France, and
International Laboratory of Quantitative Finance, National Research University Higher School of Economics,
Moscow, Russia
E-mail: Youri.Kabanov@univ-fcomte.fr

Laboratoire de Mathématiques Raphaël Salem,
Université de Rouen, Avenue de l'Université, BP.12 Technopôle du Madrillet F76801 Saint-Étienne-du-Rouvray, France, and
International Laboratory of Quantitative Finance, National Research University Higher School of Economics,
Moscow, Russia
E-mail: Serge.Pergamenchtchikov@univ-rouen.fr

1 Introduction

In the modern world insurance companies operate in a financial environment. In particular, they invest their reserves into various assets and this may add more risk to the business. To our knowledge, the first model of an insurance company investing its capital into a risky asset has appeared in the short note [12] where the author provided arguments showing that the asymptotic behavior of ruin probability is radically different from that in the classical Lundberg–Cramér model. A rigorous analysis in [13] confirmed the conjecture. Using Kalashnikov’s estimates for linear finite difference equations with random coefficients, [19], it was shown that, independently of the safety loading, the ruin is imminent with unit probability when the volatility σ of the stock price is large with respect to the instantaneous rate of return a (namely, when $2a/\sigma^2 < 1$), and the ruin probability is decreasing as a power function, when the volatility is small (namely, when $2a/\sigma^2 > 1$). For the model with exponentially distributed claims the exact asymptotic was found. Threshold case $2a/\sigma^2 = 1$ was studied in the papers [27], [28] where it was shown, using techniques based on a renewal theorem, that the ruin is imminent with unit probability. The setting of [13] and [27] is of the so-called non-life insurance: the company receives a flow of contributions and pay primes. The ruin occurs when a new claim arrives and its value is too large to be covered by the reserve: the risk process exits from the positive half-axis by a jump. The model can be studied in the discrete-time framework but the method of differential equations happens to be more efficient in the case of exponential claims where it allows to get the exact asymptotic.

In the present note we investigate the setting of life annuity, or pension, insurance, when the company pays to the policyholder a rent and earns a premium when the insured person dies, see, e.g., [16], p. 8 or [30]; in the classical literature such models are called *models with negative risk sums*. The highly stylized model of the collective risk theory may describe the situation where annuitant, entering the contract, pays a lump sum, e.g., the savings during his working life, getting in exchange regular payments until the death. One can interpret the premium as the reserve release. If the company has a large “stationary” portfolio of such contracts, with new arriving customers, then it is reasonable to think that the dates of reserve releases form a Poisson process and the sizes of premiums are independent random variables of the same distribution. A special type of life annuity, called *le viager* exists in France. This practice is, basically, an exchange of property, a house or an apartment, for life-long regularly payments. The annuitant enjoys a regularly income without having to give a house. For the company having a portfolio of viagers the amount of premiums depends on the real estate market.

For this classical model, where the reserve process has jumps upwards and may leave the positive half-axis only in a continuous way, the ruin problem can be easily reduced to the ruin problem for the non-life insurance using the so-called duality method, [2]. Its idea is to define the “dual” process by replacing the line segments of the original process between consecutive jumps by downward jumps of the same depth and the upward jumps by line segments of the same height with positive fixed slope. Note that in the literature the models with upward jumps are often referred to as *dual models*, [3], [1].

The duality method does not work in our setting where the capital of the company or its fraction is invested into a risky asset. The change of two signs to the opposite ones in the equation defining the dynamics of the reserve leads to technical complications. In particular, now the ruin may happen before the instant of the first jump and the latter is no more the instant of the regeneration, after which the process starts afresh provided being strictly positive.

Nevertheless, a suitable modification of arguments of [13] and [27] combined with some ideas allows us to obtain the asymptotic of the ruin probability as the initial reserve tends to infinity. Expectedly, it is the same as in the non-life insurance case. In many countries there are rules allowing insurance companies to invest only a small share of their reserves into risky assets. Our simple model confirms that it is reasonable and even provides a quantitative answer. To avoid the situation when the ruin happens with probability one, the proportion of investment into the risky asset should be strictly less than $2a/\sigma^2$.

It should be emphasized that the life insurance case is rather different and its study is not a straightforward exercise. The main difficulty in deriving the integro-differential equation is to prove the smoothness of the ruin probability and the integrability of derivatives. This issue is already delicate in the non-life insurance case. Unfortunately, it was not discussed in [13] where the reader was directed towards the literature. Now we are not sure that we disposed at that time a clean reference. The smoothness of the exit probability is discussed in many papers, see, e.g., an interesting article [31] where the explicit formula for an exponential functional of Brownian motion due to Marc Yor is used. Unfortunately, the needed smoothness property was established only under constraints on coefficients. One of the authors of the present note had a fruitful discussion with Marc on a possibility to deduce smoothness of the ruin probability function and the integrability of its derivatives without using complicated explicit formulae. His suggestions are realized here for the life insurance case. Note that in the literature on non-life insurance one can find other methods to establish the smoothness, for example an approach similar to verification theorems in the stochastic control theory, see [5].

The structure of the paper is as follows. Section 2 contains the formulation of the main results. In Section 3 we establish an upper asymptotic bound for the exit probability (from $]0, \infty[$) for a solution of a non-homogeneous linear stochastic equation and a lower asymptotic bound for a small volatility case. As a tool we use the Dufresne theorem rather than Goldie's renewal theorem from [15], which was the key ingredient of arguments in [27]. The proof of Theorem 2 asserting that in the case of large volatility the ruin is imminent is given in Section 4. The regularity of the non-ruin probability Φ is studied in Section 5 using a method based on integral representations. At the end of this section we derive the integro-differential equation for Φ . Section 6 contains the proof of the main theorem. Finally, in the appendix we provide a formulation of an ergodic theorem for an autoregression with random coefficients proven in [27].

Kalashnikov's approach (developed further in his joint work with Ragnar Norberg [20]) plays an important role in our study. Our techniques is elementary. More profound and general results can be found in [25], [24], [26], [22], [23] *et al.*

2 The model

We are given a stochastic basis $(\Omega, \mathcal{F}, \mathbf{F} = (\mathcal{F}_t)_{t \geq 0}, \mathbf{P})$ with a Wiener process W independent of the integer-valued random measure $p(dt, dx)$ with the compensator $\tilde{p}(dt, dx)$.

Let us consider a process $X = X^u$ of the form

$$X_t = u + a \int_0^t X_s ds + \sigma \int_0^t X_s dW_s - ct + \int_0^t \int xp(ds, dx), \quad (1)$$

where a and σ are arbitrary constants and $c \geq 0$.

We shall assume that $\tilde{p}(dt, dx) = \alpha dt F(dx)$ where $F(dx)$ is a probability distribution on $]0, \infty[$. In this case the integral with respect to the jump measure is simply a compound

Poisson process. It can be written as $\sum_{i=1}^{N_t} \xi_i$ where N is a Poisson process with intensity α and ξ_i are random variables with common distribution F ; the random variables W , N , ξ_i , $i \in \mathbb{N}$, are independent. We denote by T_n the successive jumps of N ; the inter-arrival time intervals $T_i - T_{i-1}$ are independent and exponentially distributed with the parameter α .

In our main result (Theorem 1) we assume that F is the exponential distribution with parameter μ .

Let $\tau^u := \inf\{t : X_t^u \leq 0\}$ (the instant of ruin), $\Psi(u) := P(\tau^u < \infty)$ (the ruin probability), and $\Phi(u) := 1 - \Psi(u)$ (the survival probability).

The parameter values $a = 0$, $\sigma = 0$, correspond to the life insurance (i.e. the dual) version of the Lundberg–Cramér model for which the risk process is usually written as

$$r_t := u - ct + \sum_{i=1}^{N_t} \xi_i. \quad (2)$$

In the considered case the capital evolves due to continuously outgoing cash flow with rate c and incoming random payoffs ξ_i at times forming an independent Poisson process N with intensity α . For the classical model with positive safety loading and F having a “non-heavy” tail, the Lundberg inequality provides an encouraging information: the ruin probability decreases exponentially as the initial capital u tends to infinity. Moreover, for the exponentially distributed claims the ruin probability admits an explicit expression, see [2] or [16].

The more realistic case $a > 0$, $\sigma = 0$, corresponding to non-risky investments, does not pose any problem.

We study here the case $\sigma > 0$. Now the equation (1) describes the evolution of the reserve of an insurance company which pays a rent and continuously reinvests its capital into an asset with the price following a geometric Brownian motion. The same model can be used for the description of the capital of a venture company funding R&D and selling innovations, [4].

Notations. Throughout the paper we shall use the following abbreviations:

$$\kappa := a - \frac{1}{2}\sigma^2, \quad \beta := \frac{2\kappa}{\sigma^2} = \frac{2a}{\sigma^2} - 1, \quad \eta_t := \kappa t + \sigma W_t.$$

The solution of the linear stochastic equation (1) can be written using the Cauchy formula:

$$X_t = e^{\eta_t} \left(u + \int_{[0,t]} e^{-\eta_s} dr_s \right). \quad (3)$$

Theorem 1 Let $F(x) = 1 - e^{-x/\mu}$, $x > 0$. Assume that $\sigma > 0$.

(i) If $\beta > 0$, then for some $K > 0$

$$\Psi(u) = Ku^{-\beta}(1 + o(1)), \quad u \rightarrow \infty. \quad (4)$$

(ii) If $\beta \leq 0$, then $\Psi(u) = 1$ for all $u > 0$.

The formulation of this theorem is exactly the same as in [13] for the non-life insurance model (the case $\beta = 0$ was analyzed in [27], [28]).

One must admit that in the life insurance models, especially, in the case of a company keeping a portfolio of viager contracts, the hypothesis that the benefits (i.e. prices of houses) follow the exponential distribution, are highly unrealistic. Nevertheless, we can claim, without any assumption on the distribution, that the ruin probabilities lay between two power functions, see Propositions 1 and 2.

The next result (implying the statement (ii) above) says that for $\delta > 0$ the ruin is imminent. It requires only that the distribution F has moments of positive order.

Theorem 2 *Assume that there is $\delta > 0$ such that $\mathbb{E}\xi_1^\delta < \infty$. If $\beta \leq 0$, then $\Psi(u) = 1$ for any $u > 0$.*

The same model serves well in the situation where only a fixed part $\gamma \in]0, 1]$ of the capital is invested in the risky asset: one should only replace the parameters a and σ in (1) by $a\gamma$ and $\sigma\gamma$. Theorem 1 implies that the ruin with probability one will be avoided only if $2a\gamma/(\sigma\gamma)^2 > 1$, i.e. when the share of investment into the risky asset is strictly less than $2a/\sigma^2$.

It is worth mentioning that our conclusion are robust and holds for more general models. The reader may contest that intensity of outgoing payments c is constant. Indeed, after the death of the annuitant the payments stop and the intensity must decrease while with a new customer it increases. Easy comparison arguments show that the above statements hold if $c = (c_t)$ is a random process, such that $0 < C_1 \leq c \leq C_2$ where C_1 and C_2 are constants.

The crucial part of the asymptotic analysis in Theorem 1 is based on the fact that for the Markov process given by (1) the non-exit probability $\Phi(u)$ is smooth and satisfies the following equation:

$$\frac{1}{2}\sigma^2 u^2 \Phi''(u) + (au - c)\Phi'(u) - \alpha\Phi(u) + \alpha \int_0^\infty \Phi(u + y)dF(y) = 0. \quad (5)$$

With $\sigma > 0$ this equation is of the second order and, hence, requires two boundary conditions — in contrast to the classical case ($a = 0, \sigma = 0$) where it degenerates to an equation of the first order requiring a single boundary condition, see [16]. The estimate given in Proposition 1 shows that $\Phi(\infty) = 1$.

There is an extensive literature concerning of the regularity of the survival probability for process with jumps in the context of non-life insurance models and models based on the Lévy processes, see, e.g., [31], [14], [17], [6], [7]. Our Theorem 3 requiring only smoothness of F' and the integrability of F'' seems to be the first result on the regularity of the survival probability in the considered setting.

3 Asymptotic bounds for the small volatility case

3.1 Upper asymptotic bound

Let us consider the exit probability problem for a more general process $X = X^u$ of the form

$$X_t = u + a \int_0^t X_s ds + \sigma \int_0^t X_s dW_s - ct + Z_t, \quad (6)$$

where $a, \sigma, c > 0$ are arbitrary constants and $Z = (Z_t)_{t \geq 0}$ is an increasing adapted càdlàg process starting from zero.

Proposition 1 *Assume that in the general model (1) the parameter $\beta > 0$. Then*

$$\limsup_{u \rightarrow \infty} u^\beta \Psi(u) \leq \frac{2^\beta c^\beta}{\sigma^{2\beta} \beta \Gamma(\beta)}.$$

Proof. Let Y be a solution of the linear stochastic equation

$$Y_t = u + a \int_0^t Y_s ds + \sigma \int_0^t Y_s dW_s - ct.$$

Introducing the notation

$$R_t := c \int_0^t e^{-\eta v} dv \quad (7)$$

we express the solution as

$$Y_t := e^{\eta t} (u - R_t).$$

The difference $X - Y$ satisfies the linear equation with zero initial condition. Since Z is increasing, we have the inequality $X \geq Y$ showing that the exit of X from $]0, \infty[$ implies the exit of Y . Thus,

$$\Psi(u) \leq P(R_\infty > u). \quad (8)$$

and the asymptotic behavior of the ruin probability for this general model can be estimated by the tail behavior of the distribution function of R_∞ . Using the change of variable $v = (4/\sigma^2)t$ and observing that $B_t := -(1/2)\sigma W_{(4/\sigma^2)t}$ is a Wiener process we obtain the representation

$$R_\infty = c \int_0^\infty e^{-(a-\sigma^2/2)v - \sigma W_v} dv = \frac{4c}{\sigma^2} \int_0^\infty e^{-2\beta t + 2B_t} dt =: \frac{4c}{\sigma^2} A_\infty^{(-\beta)}.$$

The Dufresne theorem (see, [9] or [21], Theorem 6.2) claims that $A_\infty^{(-\beta)}$ is distributed as the random variable $1/(2\gamma)$ where γ has the gamma distribution with parameter β . Thus,

$$\begin{aligned} \mathbf{P}(R_\infty > u) &= \mathbf{P}(2c/(\sigma^2\gamma) > u) = \mathbf{P}(\gamma < 2c/(\sigma^2 u)) = \frac{1}{\Gamma(\beta)} \int_0^{2c/(\sigma^2 u)} x^{\beta-1} e^{-x} dx \\ &\sim \frac{2^\beta c^\beta}{\sigma^{2\beta} \beta \Gamma(\beta)} u^{-\beta} \end{aligned}$$

and the result follows from (8). $\square$

3.2 Lower asymptotic bound

The next result shows that the ruin probability decreases as the initial capital tends to infinity not faster than a certain power function.

Proposition 2 *Assume that $\beta > 0$. Then there exists $\beta_* > 0$ such that*

$$\liminf_{u \rightarrow \infty} u^{\beta_*} \Psi(u) > 0.$$

Proof. Let $Y = (Y_k)_{k \geq 1}$ be the imbedded discrete-time process, that is the sequence of random variables defined recursively as follows:

$$Y_k = M_k Y_{k-1} + Q_k, \quad Y_0 = u, \quad (9)$$

where

$$M_k = e^{\eta T_k - \eta T_{k-1}} \quad \text{and} \quad Q_k = \xi_k - c \int_{T_{k-1}}^{T_k} e^{\eta T_k - \eta v} dv.$$

Let $\theta^u := \inf\{k : Y_k \leq 0\}$. It is clear that $X_{T_k} = Y_k$ for any $k \geq 1$. So, for any $u > 0$

$$\Psi(u) = \mathbf{P}(\tau^u < \infty) \geq \mathbf{P}(\theta^u < \infty). \quad (10)$$

Take $\varrho \in]0, 1[$ and chose B sufficiently large to ensure that

$$B_1 = B - \frac{1}{\varrho^2(1-\varrho)} > 0.$$

Define the sets

$$\Gamma_k = \{M_k \leq \varrho\} \cap \{Q_k \leq \varrho^{-1}\}, \quad D_k = \{M_k \leq \varrho^{-1}\} \cap \{Q_k \leq -B\}. \quad (11)$$

On the set $\cap_{k=1}^n \Gamma_k$ we have

$$Y_n = u \prod_{j=1}^n M_j + \sum_{k=1}^n Q_k \prod_{j=k+1}^n M_j \leq u\varrho^n + \frac{1}{\varrho(1-\varrho)}.$$

Therefore, on the set $\cap_{k=1}^n \Gamma_k \cap D_{n+1}$

$$Y_{n+1} = M_{n+1}Y_n + Q_{n+1} \leq u\varrho^{n-1} + \frac{1}{\varrho^2(1-\varrho)} - B = u\varrho^{n-1} - B_1.$$

It is easy to check that $u\varrho^{n-1} \leq B_1$, when $u > B_1$ and

$$n = 3 + \left\lfloor \frac{\ln(u/B_1)}{|\ln \varrho|} \right\rfloor,$$

where $\lfloor \dots \rfloor$ means the integer part. Therefore,

$$\mathbf{P}(\theta^u < \infty) \geq \mathbf{P}(\cap_{k=1}^n \Gamma_k \cap D_{n+1}) = (\mathbf{P}(\Gamma_1))^n \mathbf{P}(D_1).$$

Taking into account that $\mathbf{P}(\Gamma_1) > 0$ and $\mathbf{P}(D_1) > 0$, we obtain that

$$\lim_{u \rightarrow \infty} u^{\beta_*} \mathbf{P}(\theta^u < \infty) = \infty$$

for any

$$\beta_* > \frac{\ln \mathbf{P}(\Gamma_1)}{\ln \varrho}.$$

This implies the claim. $\square$

4 Large volatility: proof of Theorem 2

We consider separately two cases: $\beta < 0$ and $\beta = 0$ and show that in both cases the ruin probability is equal to one.

Proposition 3 Assume that $\beta < 0$ and $\mathbf{E} \xi_1^\delta < \infty$ for some $\delta > 0$. Then $\Psi(u) = 1$ for all $u > 0$.

Proof. As in the proof of Proposition 2 we consider the imbedded discrete-time process $Y = (Y_k)_{k \geq 1}$ defined by (9). By virtue of (10) it is sufficient to show that $\mathbf{P}(\theta^u < \infty) = 1$. Note that for $\delta \in]0, -\beta[$

$$\mathbf{E}e^{\delta \eta_t} = \mathbf{E}e^{\delta(\kappa t + \sigma W_t)} = e^{\delta t(\beta + \delta)\sigma^2/2} < 1,$$

and, therefore,

$$\mathbf{E}M_1^\delta = \int_0^\infty \mathbf{E}e^{\delta \eta_t} \alpha e^{-\alpha t} dt < 1. \quad (12)$$

According to ([8], p. 250) if a random variable ν is independent of W and has the exponential distribution with parameter α , then

$$\mathbf{E}e^{q \max_{v \leq \nu} (\mu v + W_v)} = \frac{\sqrt{2\alpha + \mu^2} - \mu}{\sqrt{2\alpha + \mu^2} - \mu - q}. \quad (13)$$

provided that

$$\sqrt{2\alpha + \mu^2} - \mu - q > 0.$$

Changing the variable and estimating the integrand by its maximal value we get that

$$\begin{aligned} \mathbf{E} \left(\int_0^{T_1} e^{\eta_{T_1} - \eta_v} dv \right)^\delta &= \mathbf{E} \left(\int_0^{T_1} e^{\kappa v + \sigma W_v} dv \right)^\delta \leq \mathbf{E} T_1^\delta e^{\delta \max_{v \leq T_1} (\kappa v + \sigma W_v)} \\ &= \int_0^\infty t^\delta \mathbf{E} e^{\delta \max_{v \leq t} (\kappa v + \sigma W_v)} \alpha e^{-\alpha t} dt \\ &\leq 2 \sup_{t \geq 0} \{t^\delta e^{-\alpha t/2}\} \mathbf{E} e^{\delta \max_{v \leq \nu'} (\kappa v + \sigma W_v)} \\ &= 2e^{\delta(\ln(2\delta/\alpha) - 1)} \mathbf{E} e^{\delta \max_{v \leq \nu'} (\kappa v + \sigma W_v)}, \end{aligned}$$

where ν' is an exponential random variable with parameter $\alpha/2$. In view of the equality (13)

$$\mathbf{E} e^{\delta \max_{v \leq \nu'} (\kappa v + \sigma W_v)} = \frac{\sqrt{\sigma^2 \alpha + \kappa^2} - \kappa}{\sqrt{\sigma^2 \alpha + \kappa^2} - \kappa - \delta \sigma^2},$$

provided that

$$\delta < \frac{\sqrt{\sigma^2 \alpha + \kappa^2} - \kappa}{\sigma^2},$$

i.e. for such $\delta > 0$ we have the bound

$$\mathbf{E} \left(\int_0^{T_1} e^{\eta_{T_1} - \eta_v} dv \right)^\delta \leq 2e^{\delta(\ln(2\delta/\alpha) - 1)} \frac{\sqrt{\sigma^2 \alpha + \kappa^2} - \kappa}{\sqrt{\sigma^2 \alpha + \kappa^2} - \kappa - \delta \sigma^2}. \quad (14)$$

Using these estimates and the assumption of the proposition we conclude that $\mathbf{E}|Q_1|^\delta < \infty$ for sufficiently small $\delta > 0$. Thus, the hypothesis of the ergodic theorem for autoregression with random coefficients is fulfilled (see Proposition 8). The latter claims that for any bounded uniformly continuous function f

$$\mathbf{P}\text{-}\lim_N \frac{1}{N} \sum_{k=1}^N f(Y_k) = \mathbf{E} f(\zeta), \quad (15)$$

where

$$\zeta = Q_1 + \sum_{k=2}^{\infty} Q_k \prod_{j=1}^{k-1} M_j.$$

Let us represent ζ in the form

$$\zeta = \xi_1 - \int_0^{T_1} e^{\eta_{T_1} - \eta_v} dv + e^{\eta_{T_1}} \zeta_1, \quad \zeta_1 := \sum_{k=2}^{\infty} Q_k \prod_{j=2}^{k-1} M_j.$$

Clearly, the random variables ξ , ζ_1 and $(\eta_{T_1}, \int_0^{T_1} e^{\eta_{T_1} - \eta_v} dv)$ are independent. Moreover, Lemma 1 given after the proof implies that the support of conditional distribution of the integral $\int_0^t e^{\eta_t - \eta_v} dv$ given $\eta_t = x$ is unbounded from above. From this we easily infer that the support of distribution of ζ is unbounded from below. Thus, for

$$f(x) = \mathbf{1}_{\{x \leq -1\}} + |x| \mathbf{1}_{\{-1 < x < 0\}}$$

the right-hand side of (15) is strictly positive and, therefore, $\mathbf{P}(\inf_{k \geq 1} Y_k < 0) = 1$. $\square$

Lemma 1 *Let $\sigma > 0$. Then the support of conditional distribution of the random variable*

$$I = \int_0^1 e^{\sigma W_v} dv$$

given $W_1 = y$ is unbounded from above.

Proof. It is well-known (see, e.g., [29]) that the conditional distribution of the Wiener process $(W_v)_{v \leq 1}$ given $W_1 = y$ coincides with the distribution of the Brownian bridge B^y with $B_t^y = W_t + t(y - W_1)$. Thus, the conditional distribution of I is the same as the unconditional distribution of

$$\tilde{I} := \int_0^1 e^{\sigma(W_v + v(y - W_1))} dv.$$

Since the Wiener measure has the full support in the space $C_0[0, 1]$ of continuous functions on $[0, 1]$ with zero initial value, the support of distribution of $\tilde{I}$ is unbounded from above. $\square$

Proposition 4 *Assume that $\beta = 0$ and $\mathbf{E} \xi_1^\delta < \infty$ for some $\delta > 0$. Then $\Psi(u) = 1$ for all $u > 0$.*

Proof. In the considered case the imbedded discrete-time process is defined by (9) with

$$M_k = e^{\sigma \Delta V_k} \quad \text{and} \quad Q_k = \xi_k - c \int_{T_{k-1}}^{T_k} e^{\sigma(W_{T_k} - W_v)} dv,$$

where $V_k = W_{T_k}$ and $\Delta V_k := V_k - V_{k-1}$. To study the asymptotic properties of the equation (9) we use the approach proposed in [27] for the non-life insurance models. To this end, define recursively the sequence of random variables putting $\theta_0 := 0$ and

$$\theta_n := \inf\{k > \theta_{n-1} : V_k - V_{\theta_{n-1}} < 0\}, \quad n \geq 1. \quad (16)$$

Note that $\theta_n = \sum_{j=1}^n \Delta \theta_j$ where $(\Delta \theta_j)_{j \geq 1}$ is a sequence of i.i.d. random variables distributed as θ_1 . It is known (see, e.g., XII.7, Theorem 1a in [11]) that

$$C := \sup_{n \geq 1} n^{1/2} \mathbf{P}(\theta_1 > n) < \infty. \quad (17)$$

Putting $y_k = Y_{\theta_k}$, we obtain from (9) that

$$y_k = a_k y_{k-1} + b_k, \quad y_0 = u, \quad (18)$$

where

$$a_k = \prod_{j=1}^{\Delta\theta_k} M_{\theta_{k-1}+j} = e^{\sigma(V_{\theta_k} - V_{\theta_{k-1}})}$$

and

$$b_k = \sum_{l=1}^{\Delta\theta_k} \left(\prod_{j=l+1}^{\Delta\theta_k} M_{\theta_{k-1}+j} \right) Q_{\theta_{k-1}+l}.$$

It is clear that $a_k < 1$ a.s. Moreover, the first condition in Theorem 2 and the inequality (14) with $\kappa = 0$ implies that $\mathbf{E}|Q_1|^\delta < \infty$ for any sufficiently small δ . Now, taking into account that

$$|b_1| \leq \sum_{l=1}^{\Delta\theta_1} \left(\prod_{j=l+1}^{\Delta\theta_1} M_j \right) |Q_l| = \sum_{l=1}^{\Delta\theta_1} \frac{a_1}{\prod_{j=1}^l M_j} |Q_l| \leq \sum_{l=1}^{\Delta\theta_1} |Q_l|,$$

we can get, for $r \in]0, 1[$ and an increasing sequence of integers l_n , that

$$\begin{aligned} \mathbf{E}|b_1|^r &\leq 1 + r \sum_{n \geq 1} \frac{1}{n^{1-r}} \mathbf{P}(|b_1| > n) \\ &\leq 1 + r \sum_{n \geq 1} \frac{1}{n^{1-r}} \mathbf{P}\left(\sum_{j=1}^{l_n} |Q_j| > n\right) + r \sum_{n \geq 1} \frac{1}{n^{1-r}} \mathbf{P}(|t_1| > l_n) \\ &\leq 1 + r \mathbf{E}|Q_1|^\delta \sum_{n \geq 1} \frac{l_n}{n^{1-r+\delta}} + r C \sum_{n \geq 1} \frac{1}{n^{1-r} l_n^{1/2}}. \end{aligned}$$

Putting here $l_n = \lfloor n^{4r} \rfloor$ we obtain that $\mathbf{E}|b_1|^r < \infty$ for any $r \in]0, \delta/5[$. Therefore, due to Proposition 8, we obtain that for any bounded uniformly continuous function f

$$\mathbf{P}\text{-}\lim_N \frac{1}{N} \sum_{k=1}^N f(y_k) = \mathbf{E}f(\zeta),$$

where

$$\zeta = b_1 + \sum_{k=2}^{\infty} b_k \prod_{j=1}^{k-1} a_j. \quad (19)$$

Now we show that

$$\mathbf{P}(\zeta < -x) > 0 \quad \text{for any } x > 0.$$

Indeed, the random variable (19) can be represented as

$$\zeta = b_1 + a_1 \zeta_1, \quad \zeta_1 = \sum_{k=2}^{\infty} b_k \prod_{j=2}^{k-1} a_j.$$

It is clear that the ζ_1 is independent on b_1 and a_1 . Note that on the set $\{\Delta\theta_1 = 1\}$ we have $a_1 = M_1$ and $b_1 = Q_1$. Therefore, for any $x > 0$

$$\mathbf{P}(\zeta < -x) \geq \mathbf{P}(b_1 + a_1 \zeta_1 < -x, \Delta\theta_1 = 1) = \mathbf{P}(Q_1 + M_1 \zeta < -x, w_{T_1} < 0)$$

and we conclude as in the previous proposition. $\square$

5 Regularity of the ruin probability

5.1 Integral representations

The proof of smoothness of a function H admitting an integral representation is based on a simple idea which merits to be explained.

First, we recall the classical result on differentiability of the integral $H(u) = \int f(u, z)dz$ where $f(u, \cdot) \in L^1$ for each u from open subset $U \in \mathbb{R}$. If $f(\cdot, z)$ is differentiable on an open interval $]u_0 - \varepsilon, u_0 + \varepsilon[$ for almost all z and, on this interval, $|\partial f(\cdot, z)/\partial u| \leq g(z)$ (a.e.) where $g \in L^1$. Then H is differentiable at u_0 and $H'(u_0) = \int \partial f(u_0, z)/\partial u dz$.

Suppose that we are given a bounded measurable function $h(z)$ and a Gaussian random variable $\zeta \sim N(0, 1)$. Let $H(u) = \mathbf{E}h(u + \zeta) = \int h(u + x)\varphi_{0,1}(x)dx$. Then H is differentiable and even of the class C^∞ . Of course, the above result cannot be applied directly. But using the change of variable we get the representation

$$H(u) = \int h(u + x)\varphi_{0,1}(x)dx = \int h(z)\varphi_{0,1}(z - u)dz.$$

Now the parameter u appears only in the function $\varphi_{0,1}$, the integrand is differentiable in u and we can apply the classical sufficient condition.

The issues here are: an integral representation, the smoothness of the density, and the integrability of its derivatives. In the case of survival probability Φ the integral representation is obtained from the strong Markov property. Unfortunately, the structure of the representation is rather complicated, the random variable standing for ζ is not a Gaussian one, and its density is not given by a simple formula. Nevertheless, the idea of using a change of variable to move the parameter from the unknown function on which we have only a limited information (essentially, the boundedness and measurability) to the density still does work. The main difficulty is to check the smoothness of the density and find appropriate bounds for its derivatives.

Theorem 3 *Assume that the distribution function of ξ_1 has a density f differentiable on $\mathbb{R}_+$ and such that $f' \in L^1(\mathbb{R}_+)$. Then $\Phi(u)$ is two times continuously differentiable on $]0, \infty[$.*

Proof. We again consider the process

$$Y_t^u = e^{\eta_t} (u - R_t), \quad (20)$$

where R_t is defined in (7). Put

$$\theta^u := \inf\{t \geq 0: Y_t^u \leq 0\}. \quad (21)$$

By virtue of the strong Markov property of X^u

$$\Phi(u) = \mathbf{E}\Phi(X_{\theta^u \wedge T_1}^u). \quad (22)$$

Note that the process Y^u is strictly positive before the time θ^u , zero at θ^u , and strictly negative afterwards. Due to the independence of the Wiener process and the instants of jumps, $\theta^u \neq T_1$ a.s. Thus, $\{Y_{T_1}^u > 0\} = \{\theta^u > T_1\}$ a.s. Taking into account that $\Phi(0) = 0$, we get that

$$\Phi(u) = \mathbf{E}I_{\{Y_{T_1}^u > 0\}}\Phi(X_{T_1}^u) = \mathbf{E}I_{\{Y_{T_1}^u > 0\}}\Phi(Y_{T_1}^u + \xi_1) = \Phi_1(u) + \Phi_2(u),$$

where

$$\Phi_1(u) := \alpha \int_0^2 \mathbf{E}G(Y_t^u) e^{-\alpha t} dt, \quad \Phi_2(u) := \alpha \int_2^\infty \mathbf{E}G(Y_t^u) e^{-\alpha t} dt$$

with

$$G(y) := \mathbf{1}_{\{y>0\}} \mathbf{E} \Phi(y + \xi_1) = \mathbf{1}_{\{y>0\}} \int_0^\infty \Phi(y+x) dF(x).$$

We analyze separately the smoothness of Φ_1 and Φ_2 using for this function appropriate integral representations.

5.2 Smoothness of Φ_2

We start with a simpler case of Φ_2 and show that this function is infinitely differentiable without any assumptions on the distributions of ξ_1 .

From the representation

$$Y_t^u = e^{\eta_t - \eta_1} Y_1^u - c \int_1^t e^{\eta_t - \eta_s} ds, \quad t \geq 1,$$

we obtain, using the independence of Y_1^u and the process $(\eta_s - \eta_1)_{s \geq 1}$, that

$$\mathbf{E}(G(Y_t^u) | Y_1^u) = G(t, Y_1^u),$$

where

$$G(t, y) := \mathbf{E}G\left(e^{\eta_t - \eta_1} y - c \int_1^t e^{\eta_t - \eta_s} ds\right).$$

Substituting the expression for Y_1^u given by (22) we have:

$$\Phi_2(u) = \mathbf{E} \int_2^\infty \mathbf{E}(G(Y_t^u) | Y_1^u) \alpha e^{-\alpha t} dt = \mathbf{E}H\left(e^{\kappa + \sigma W_1}(u - R_1)\right),$$

where H is a function taking values in $[0, 1]$ and given by the formula

$$H(y) := \alpha \int_2^\infty G(t, y) e^{-\alpha t} dt.$$

Taking into account that the conditional distribution of the process $(W_s)_{s \leq 1}$ given $W_1 = x$ is the same as of the Brownian bridge $B^x = (B_s^x)_{s \leq 1}$ with $B_s^x = W_s + s(x - W_1)$ we obtain the representation

$$\Phi_2(u) = \int \mathbf{E}H(e^{\kappa + \sigma x}(u - \zeta^x)) \varphi_{0,1}(x) dx \quad (23)$$

where

$$\zeta^x := c \int_0^1 e^{-(\kappa s + sx + \sigma(W_s - sW_1))} ds. \quad (24)$$

Lemma 2 below asserts that for every x the random variable ζ^x has a density $\rho(x, \cdot)$ on $]0, \infty[$ and we easily obtain from (23) by changing variable that

$$\Phi_2(u) = \int_0^u \int H(e^{\kappa + \sigma x} z) \rho(x, u - z) \varphi_{0,1}(x) dx dz. \quad (25)$$

Lemma 2 *The random variable ζ^x has a density $\rho(x, \cdot) \in C^\infty$ such that for any $n \geq 1$*

$$\sup_{y \geq 0} \left| \frac{\partial^n}{\partial y^n} \rho(x, y) \right| \leq C_n e^{C_n |x|} \quad (26)$$

with some constant C_n and $(\partial^n / \partial y^n) \rho(x, 0) = 0$.

Proof. We obtain the result using again the integral representation. Let us introduce the random process

$$\begin{aligned} D_s &:= ((W_s - 2sW_{1/2}) + s(W_{1/2} - W_1))I_{\{s \leq 1/2\}} \\ &\quad + ((1-s)(W_s - W_{1/2})) - s(W_1 - W_s))I_{\{s > 1/2\}}, \end{aligned}$$

and the piecewise linear function

$$\gamma_s := sI_{\{s \leq 1/2\}} + (1-s)I_{\{s > 1/2\}}, \quad s \in [0, 1].$$

The following identity is obvious:

$$W_s - sW_1 = D_s + \gamma_s W_{1/2}.$$

Since D_s and $W_{1/2}$ are independent random variables, for any bounded Borel function g we have:

$$\mathbf{E} g(\zeta^x) = \mathbf{E} \int g\left(c \int_0^1 e^{-(\kappa s + sx + \sigma D_s + \sigma \gamma_s v)} ds\right) \varphi_{0,1/2}(v) dv$$

Let $v(x, \cdot)$ be the inverse of the continuous strictly decreasing function

$$y \mapsto c \int_0^1 e^{-(\kappa s + sx + \sigma D_s + \sigma \gamma_s y)} ds$$

depending on the parameter x (and also on ω omitted as usual). Note that $v(x, 0+) = \infty$ and $v(x, \infty) = 0$. After the change of variable, we obtain, using the notation

$$K(x, z) := c\sigma \int_0^1 \gamma_s e^{-(\kappa s + sx + \sigma D_s + \sigma \gamma_s z)} ds,$$

that

$$\mathbf{E} g(\zeta^x) = \int_0^\infty g(y) \rho(x, y) dy,$$

where

$$\rho(x, \cdot) := \mathbf{E} \frac{\varphi_{0,1/2}(v(x, \cdot))}{K(x, v(x, \cdot))}. \quad (27)$$

Thus, $\rho(x, \cdot)$ is the density of distribution of the random variable ζ^x . It remains to check that it is infinitely differentiable and find appropriate bounds for its derivatives.

Put

$$Q^{(0)}(x, z) := \frac{\varphi_{0,1/2}(z)}{K(x, z)}, \quad Q^{(n)}(x, z) := -\frac{Q_z^{(n-1)}(x, z)}{K(x, z)}, \quad n \geq 1.$$

Then

$$\frac{\partial}{\partial y} Q^{(0)}(x, v(x, y)) = Q_z^{(0)}(x, v(x, y)) v_y(x, y) = -\frac{Q_z^{(0)}(x, v(x, y))}{K(x, v(x, y))} = Q^{(1)}(x, v(x, y))$$

and, similarly,

$$\frac{\partial^n}{\partial y^n} Q^{(0)}(x, v(x, y)) = -\frac{Q_z^{(n-1)}(x, v(x, y))}{K(x, v(x, y))} = Q^{(n)}(x, v(x, y)).$$

It is easily seen that

$$Q^{(n)}(x, z) = \frac{\varphi_{0,1/2}(z)}{K^{n+1}(x, z)} \sum_{k=0}^n P_k(z) R_{n-k}(x, z), \quad (28)$$

where $P_k(z)$ is a polynomial of order k and $R_{n-k}(x, z)$ is a linear combination of products of derivatives of $K(x, z)$ in variable z . Note that for any $x, y \in [0, 1]$ we have the bounds

$$-|x| - \sigma|z| - 3\sigma W_1^* \leq \kappa s + sx + \sigma D_s + \sigma \gamma_s z \leq \kappa + |x| + \sigma|z| + 3\sigma W_1^*$$

where $W_1^* = \sup_{s \leq 1} |W_s|$. It follows that there exists a constant $C_n > 0$ such that

$$|Q^{(n)}(x, z)| \leq C_n e^{C_n|x|} (1+z^n) e^{-z^2} e^{C_n W_1^*}. \quad (29)$$

Since $\mathbf{E} e^{C_n W_1^*} < \infty$, for each x, y and n the derivative $(\partial^n / \partial y^n) Q^{(0)}(x, v(x, y))$ admits a $\mathbf{P}$ -integrable bound. Thus, we can differentiate under the sign of expectation and obtain that

$$\frac{\partial^n}{\partial y^n} \rho(x, y) = \mathbf{E} \frac{\partial^n}{\partial y^n} \frac{\varphi_{0,1/2}(v(x, y))}{K(x, v(x, y))} = \mathbf{E} \frac{\partial^n}{\partial y^n} Q^{(0)}(x, v(x, y)) = \mathbf{E} Q^{(n)}(x, v(x, y)).$$

Moreover, the bound (29) ensures that, for some constant $\tilde{C}_n$,

$$\sup_{y \geq 0} \mathbf{E} \left| \frac{\partial^n}{\partial y^n} Q^{(0)}(x, v(x, y)) \right| \leq \mathbf{E} \sup_{z \in \mathbb{R}} |Q^{(n)}(x, z)| \leq \tilde{C}_n e^{C_n|x|}$$

and the bound (26) holds.

Since $v(x, 0+) = \infty$, the bound (29) implies that $(\partial^n / \partial y^n) \rho(x, 0) = 0$. $\square$

Remark 1 It is worth to trace in these arguments the dependence of the constant $\tilde{C}_n$ on the parameters c and σ when they are approaching zero. From the formula (28) it is clear that $\tilde{C}_n$ should be proportional to $(c\sigma)^{-n}$.

Proposition 5 *The function $\Phi_2(u)$ belongs to $C^\infty([0, \infty[)$.*

Proof. Putting

$$\tilde{H}(u, z) := \int H(e^{\kappa+\sigma x} z) \rho(x, u-z) \varphi_{0,1}(x) dx,$$

we rewrite the formula (25) as

$$\Phi_2(u) = \int_0^u \tilde{H}(u, z) dz. \quad (30)$$

Clearly, the function $\tilde{H}$ is continuous on $]0, \infty[\times]0, \infty[$. Using Lemma 2 we obtain that

$$\frac{\partial}{\partial u} \tilde{H}(u, z) = \int H(e^{\kappa+\sigma x} z) \frac{\partial}{\partial u} \rho(x, u-z) \varphi_{0,1}(x) dx$$

and

$$\sup_{u,z} \left| \frac{\partial}{\partial u} \tilde{H}(u, z) \right| < \infty.$$

By induction, for every $n \geq 1$,

$$\frac{\partial^n}{\partial u^n} \tilde{H}(u, z) = \int H(e^{\kappa+\sigma x} z) \frac{\partial^n}{\partial u^n} \rho(x, u-z) \varphi_{0,1}(x) dx$$

and

$$\sup_{u,z} \left| \frac{\partial^n}{\partial u^n} \tilde{H}(u, z) \right| < \infty.$$

By virtue of Lemma 2 $\rho(x, 0) = 0$, i.e. $\tilde{H}(u, u) = 0$. So,

$$\frac{d}{du} \Phi_2(u) = \tilde{H}(u, u) + \int_0^u \frac{\partial}{\partial u} \tilde{H}(u, z) dz = \int_0^u \frac{\partial}{\partial u} \tilde{H}(u, z) dz.$$

In the same way we check that

$$\frac{d^n}{du^n} \Phi_2(u) = \int_0^u \frac{\partial^n}{\partial u^n} \tilde{H}(u, z) dz.$$

for any $n \geq 1$. $\square$

5.3 Smoothness of Φ_1

Arguing in the same spirit as in the previous subsection but taking this time the conditional expectation with respect to W_t we obtain that

$$\mathbf{E} \mathbf{1}_{\{R_t < u\}} h(e^{\kappa t + \sigma W_t} (u - R_t)) = \frac{1}{\sqrt{t}} \mathbf{E} \int \mathbf{1}_{\{\zeta^{t,x} < u\}} h(u, t, x) \varphi_{0,1} \left(\frac{x}{\sqrt{t}} \right) dx$$

where we use the abbreviations

$$h(u, t, x) := h(e^{\kappa t + \sigma x} (u - \zeta^{t,x})), \quad h(y) = \mathbf{E} \Phi(y + \xi_1)$$

and

$$\zeta^{t,x} := c \int_0^t e^{-(sx/t + \kappa s + \sigma(W_s - (s/t)W_t))} ds.$$

It is easily seen that the random variable $\zeta^{t,x}$ has infinitely differentiable density (the same as of ζ^x defined in (24) but with the parameters ct , κt , and $\sigma t^{1/2}$). Unfortunately, derivatives of this density have non-integrable singularities as t tends to zero (see Remark at the end of previous subsection). By this reason we cannot use the strategy of proof used for Φ_2 . Nevertheless, the hypothesis on the distribution of ξ_1 allows us to establish the claimed result.

Note that the function $x \rightarrow \zeta^{t,x}$ is strictly decreasing and maps $\mathbb{R}$ onto $\mathbb{R}_+$. Let denote $z(t, \cdot)$ its inverse. The derivative of the latter is given by the formula

$$z_x(t, x) = -\frac{t}{L(t, z(t, x))},$$

where

$$L(t, z) = c \int_0^t s e^{-(sz/t + \kappa s + \sigma(W_s - (s/t)W_t))} ds. \quad (31)$$

Changing the variable we obtain that

$$\int \mathbf{1}_{\{\zeta^t, z < u\}} h(u, t, z) \varphi_{0,1} \left(\frac{z}{\sqrt{t}} \right) dz = t \int_0^u h(u, t, z(t, x)) D(t, z(t, x)) dx,$$

where

$$D(t, z) = \frac{\varphi_{0,1}(z/\sqrt{t})}{L(t, z)}.$$

Summarizing, we get that

$$\Phi_1(u) = \alpha \mathbf{E} \int_0^2 \sqrt{t} H(t, u) e^{-\alpha t} dt,$$

where

$$H(t, u) := \int_0^u h(u, t, z(t, x)) D(t, z(t, x)) dx. \quad (32)$$

Proposition 6 *Under the conditions of Theorem 3 the function $H(t, u)$ defined in (32) for any fixed $u_0 > 0$ satisfies the following inequality*

$$\sup_{t \in]0, 2]} \mathbf{E} \sup_{u \geq u_0} (|H_u(t, u)| + |H_{uu}(t, u)|) < \infty. \quad (33)$$

Proof. In virtue of the hypothesis the function $h(y) = \mathbf{E}\Phi(y + \xi_1)$ is differentiable. Differentiating (32) we get that

$$H_u(t, u) = h(0) D(t, z(t, u)) + \int_0^u h_u(u, z(t, x)) D(t, z(t, x)) dx, \quad (34)$$

where $h(0) = \mathbf{E}\Phi(\xi_1)$ and

$$h_u(u, t, z(t, x)) = h'(e^{\kappa t + \sigma z(t, x)}(u - x)) e^{\kappa t + \sigma z(t, x)}.$$

Note that

$$\frac{\partial}{\partial x} h(u, t, z(t, x)) = h_u(u, t, z(t, x)) [\sigma z_x(t, x)(u - x) - 1].$$

Therefore,

$$\int_0^u h_u(u, z(t, x)) D(t, z(t, x)) dx = - \int_0^u \frac{\varphi_{0,1}(z(t, x)/\sqrt{t})}{\sigma t(u - x) + L(t, z(t, x))} dx h(u, t, z(t, x)).$$

Integrating by parts and taking into account that $z(t, 0+) = \infty$ we get that

$$H_u(t, u) = \int_0^u h(u, t, z(t, x)) \Theta(u, t, z(t, x)) \varphi_{0,1}(z(t, x)/\sqrt{t}) dx, \quad (35)$$

where

$$\Theta(u, t, z) = \frac{z}{L(t, z)(\sigma t(u - \zeta^{t,z}) + L(t, z))} - \frac{t\sigma L(t, z) + tL_z(t, z)}{L(t, z)(\sigma t(u - \zeta^{t,z}) + L(t, z))^2}.$$

Inspecting the formula (31) defining $L(t, z)$ we conclude that there exist positive constants C_0 ("small") and C_1 ("large") such that

$$\max_{t \in]0, 2]} (L(t, z) + t|Lz(t, z)|) \leq C_1 e^{C_1(|z| + W_t^*)}$$

and for any $t \in]0, 2]$

$$|L(t, z)| \geq C_0 t^2 e^{-C_1(|z| + W_t^*)},$$

where $W_t^* := \max_{v \leq t} |W_v|$. Taking this into account we obtain that for some $C > 0$

$$|\Theta(u, t, z)| \leq C t^{-6} e^{C(|z| + W_t^*)}. \quad (36)$$

Using a generic notation C for a constant (which may vary even within a single formula) we obtain, for any $t \in]0, 2]$

$$\begin{aligned} |H_u(t, u)| &\leq C t^{-7} e^{C W_t^*} \int_{z(t, u)}^{\infty} e^{C|z| - \frac{z^2}{2t}} L(t, z) dz \\ &\leq C t^{-7} e^{C W_2^*} e^{-\frac{z^2(t, u)}{4t}} \int_0^{\infty} e^{C|z| - \frac{z^2}{4}} dz. \end{aligned}$$

So, we have the bound

$$|H_u(t, u)| \leq C t^{-7} e^{C W_2^*} e^{-\frac{z^2(t, u)}{4t}}.$$

For any $u \geq u_0 > 0$ and $t \in]0, 2]$

$$u_0 \leq u = \zeta^{t, z(t, u)} \leq c t e^{2|\kappa| + \sigma|z(t, u)| + \sigma W_t^*},$$

i.e.

$$|z(t, u)| \geq \frac{1}{\sigma} \ln \frac{e^{-2|\kappa|} u_0}{t c} - W_t^*.$$

Put

$$t_0 := \min \left(\frac{e^{-2|\kappa| - 3\sigma} u_0}{c}, 2 \right), \quad \Gamma := \{W_t^* \leq t^{1/4}\}.$$

Thus, for $t \in]0, t_0]$ on the set Γ we have the inequality

$$|z(t, u)| \geq 1.$$

Taking into account that $\mathbf{E} e^{a W_t^*} < \infty$ for any a and $t > 0$, we obtain that

$$\mathbf{E} \max_{t \in [t_0, 2]} \sup_{u \geq 0} |H_u(t, u)| \leq C t_0^{-7} \mathbf{E} e^{C W_2^*} < \infty.$$

For $t \in]0, t_0]$ we have

$$\begin{aligned} \mathbf{E} \max_{u \geq u_0} |H_u(t, u)| &\leq C t^{-7} \left(e^{-\frac{1}{4t}} + \mathbf{E} e^{C W_t^*} \mathbf{1}_{\Gamma^c} \right) \\ &\leq C t^{-7} \left(e^{-\frac{1}{4t}} + \sqrt{\mathbf{E} e^{C W_2^*}} \sqrt{\mathbf{P}(W_t^* \geq t^{1/4})} \right). \end{aligned}$$

By the Chebyshev inequality we have:

$$\mathbf{P}\left(W_t^* \geq t^{1/4}\right) \leq e^{-t^{-1/4}} \mathbf{E} e^{\frac{W_t^*}{\sqrt{t}}} = e^{-t^{-1/4}} \mathbf{E} e^{W_1^*},$$

that is

$$\sup_{t \geq 0} e^{t^{-1/4}} \mathbf{P}\left(W_t^* \geq t^{1/4}\right) < \infty.$$

This implies that

$$\max_{t \in]0, t_0]} \mathbf{E} \max_{u \geq u_0} |H_u(t, u)| < \infty.$$

Therefore,

$$\max_{t \in]0, 2]} \mathbf{E} \max_{u \geq u_0} |H_u(t, u)| < \infty.$$

Differentiating (35) we find that

$$H_{uu}(t, u) = h(0) \Theta(u, t, z(t, u)) \varphi\left(\frac{z(t, u)}{\sqrt{t}}\right) + \int_0^u \Upsilon(u, t, z(t, x)) \varphi\left(\frac{z(t, x)}{\sqrt{t}}\right) dx,$$

where

$$\Upsilon(u, t, z) = h_u(u, t, z) \Theta(u, t, z) + h(u, t, z) \Theta_u(u, t, z).$$

By assumption, the distribution function F has the density f whose derivative f' is a continuous function on $\mathbb{R}_+$ integrable with respect to the Lebesgue measure. Changing variable we get that

$$\begin{aligned} h'(y) &= \frac{d}{dy} \int_0^\infty \Phi(y+x) f(x) dx = \frac{d}{dy} \int_y^\infty \Phi(x) f(x-y) dx \\ &= -\Phi(y) f(0) - \int_y^\infty \Phi(z) f'(z-y) dz, \end{aligned}$$

i.e.

$$\sup_{x \geq 0} |h'(x)| < \infty.$$

Similarly to (36) we obtain that

$$|\Upsilon_t(u, z)| \leq C t^{-8} e^{C(|z|+W_t^*)}.$$

Therefore,

$$|H_{uu}(t, u)| \leq C t^{-9} e^{C W_t^*} e^{-\frac{z^2(t, u)}{4t}},$$

implying that

$$\max_{t \in]0, 2]} \mathbf{E} \max_{u \geq u_0} |H_{uu}(t, u)| < \infty.$$

Proposition 6 is proven. $\square$

5.4 Integro-differential equation for the survival probability

Proposition 7 *Suppose that $\Phi \in C^2$. Then Φ satisfies the equation (5).*

Proof. For $h > 0$ and $\epsilon > 0$ assumed to be small enough to ensure that $u \in]\epsilon, \epsilon^{-1}[$ we put

$$\tau_h^\epsilon := \inf \{t \geq 0: Y_t^u \notin [\epsilon, \epsilon^{-1}]\} \wedge h.$$

Using the Ito formula and taking into account that on the interval on $[0, T_1[$ the process X^u coincides with Y^u we obtain the representation

$$\begin{aligned} \Phi(X_{\tau_h^\epsilon \wedge T_1}^u) &= \Phi(u) + \sigma \int_0^{\tau_h^\epsilon \wedge T_1} \Phi'(Y_s^u) dW_s \\ &\quad + \int_0^{\tau_h^\epsilon \wedge T_1} \left(\frac{1}{2} \sigma^2 (Y_s^u)^2 \Phi''(Y_s^u) + (aY_s^u - c) \Phi'(Y_s^u) \right) ds \\ &\quad + (\Phi(Y_{T_1}^u + \xi_1) - \Phi(Y_{T_1}^u)) I_{\{T_1 \leq \tau_h^\epsilon\}}. \end{aligned}$$

Due to the strong Markov property $\Phi(u) = \mathbf{E} \Phi(X_{\tau_h^\epsilon \wedge T_1}^u)$. For every $\epsilon > 0$ the integrands above are bounded by constants and, hence, the expectation of the stochastic integral is zero. Moreover, $\tau_h^\epsilon \wedge T_1 = h$ when h is sufficiently small (the threshold below which we have this equality, of course, depends on ω).

It follows that, independently of ϵ ,

$$\frac{1}{h} \mathbf{E} \int_0^{\tau_h^\epsilon \wedge T_1} \left(\frac{1}{2} \sigma^2 (Y_s^u)^2 \Phi''(Y_s^u) + aY_s^u \Phi'(Y_s^u) - c \right) ds \rightarrow \frac{1}{2} \sigma^2 u^2 \Phi''(u) + (au - c) \Phi'(u).$$

Finally,

$$\frac{1}{h} \mathbf{E} (\Phi(Y_{T_1}^u + \xi_1) - \Phi(Y_{T_1}^u)) I_{\{T_1 \leq \tau_h^\epsilon\}} = \alpha \mathbf{E} \frac{1}{h} \int_0^{\tau_h^\epsilon} (\Phi(Y_t^u + \xi_1) - \Phi(Y_t^u)) e^{-\alpha t} dt.$$

The right-hand converges to $\alpha(\mathbf{E} \Phi(u + \xi_1) - \Phi(u))$ as $h \rightarrow 0$ in virtue of the Lebesgue theorem on dominated convergence. It follows that Φ satisfies the equation (5). $\square$

6 Proof of Theorem 1

Assume that the claims are exponentially distributed, i.e. $F(x) = 1 - e^{-x/\mu}$. Similarly to the classical case, this assumption allows us to obtain for the ruin probability an ordinary differential equation (but of a higher order). Indeed, now the integro-differential equation (5) has the form

$$\frac{1}{2} \sigma^2 u^2 \Phi''(u) + (au - c) \Phi'(u) - \alpha \Phi(u) + \frac{\alpha}{\mu} \int_0^\infty \Phi(u + y) e^{-y/\mu} dy = 0. \quad (37)$$

Notice that

$$\frac{d}{du} \int_0^\infty \Phi(u + y) e^{-y/\mu} dy = -\Phi(u) + \frac{1}{\mu} \int_0^\infty \Phi(u + y) e^{-y/\mu} dy.$$

Differentiating (37) and adding to it the obtained identity multiplied by μ we exclude the integral term and arrive at a third order ordinary differential equation. It does not contain

the function itself and, therefore, is reduced to a second order differential equation for the function $G = \Phi'$ which can be easily transformed to the form

$$G'' - p(u)G' + p_0(u)G = 0, \quad (38)$$

where

$$p(u) := \frac{1}{\mu} - 2 \left(1 + \frac{a}{\sigma^2} \right) \frac{1}{u} + \frac{2c}{\sigma^2} \frac{1}{u^2},$$

$$p_0(u) := -\frac{2a}{\mu\sigma^2} \frac{1}{u} + (a - \alpha + c/\mu) \frac{2}{\sigma^2} \frac{1}{u^2}.$$

The substitution $G(u) = R(u)Z(u)$ with

$$R(u) := \exp \left\{ \frac{1}{2} \int_1^u p(s) ds \right\}$$

eliminates the first derivative and leads to the equation

$$Z'' - q(u)Z = 0, \quad (39)$$

where

$$q(u) := \frac{1}{4\mu^2} + \left(\frac{a}{\sigma^2} - 1 \right) \frac{1}{u} + \sum_{i=2}^4 A_i \frac{1}{u^i}$$

with certain constants A_i which are of no importance in our asymptotic analysis. It is easy to check that

$$\int_{x_0}^{\infty} \left(\frac{|q''(x)|}{q^{3/2}(x)} + \frac{|q'(x)|^2}{q^{5/2}(x)} \right) dx < \infty,$$

where $x_0 = \sup\{x \geq 1: q(x) \leq 0\} + 1$.

According to [10], pp. 54-55, the equation (39) has two fundamental solutions Z_+ and Z_-

$$Z_{\pm}(x) = \sqrt{2\mu} \exp \left\{ \pm \int_{x_0}^x \sqrt{q(z)} dz \right\} (1 + o(1)), \quad x \rightarrow \infty,$$

i.e.

$$Z_{\pm} \sim \exp \left\{ \pm \left(\frac{x}{2\mu} + \frac{a - \sigma^2}{\sigma^2} \ln x \right) \right\}.$$

Since

$$R(x) \sim \exp \left\{ \frac{x}{2\mu} - \frac{a + \sigma^2}{\sigma^2} \ln x \right\},$$

we obtain that (38) admits, as solutions, functions with the following asymptotics:

$$G_+(x) \sim x^{-2} e^{\frac{1}{\mu}x}, \quad G_-(x) \sim x^{-2a/\sigma^2}.$$

The differential equation of the third order for Φ has three solutions: $\Phi_0(u) = 1$,

$$\Phi_+(u) = \int_{x_0}^u G_+(x) dx \quad \Phi_-(u) = - \int_u^{+\infty} G_-(x) dx.$$

The ruin probability $\Psi := 1 - \Phi$ is the linear combination of these functions, i.e.

$$\Psi(u) = C_0 + C_1 \Phi_+(u) + C_2 \Phi_-(u).$$

Since $\Phi_+(u) \rightarrow \infty$ as $u \rightarrow \infty$, we obtain immediately that $C_1 = 0$. For the case $\beta > 0$ we know from Proposition 1 that $\Psi(\infty) = 0$. Thus, $C_0 = 0$ and

$$\Psi(u) = C_2 \int_u^\infty x^{-2a/\sigma^2} (1 + \delta(x)) dx,$$

where $\delta(x) \rightarrow 0$ as $x \rightarrow \infty$ and $C_2 > 0$. The integral decreases at infinity as the power function $u^{-\beta}/\beta$ and we obtain Theorem 1. $\square$

7 Appendix: ergodic theorem for an autoregression with random coefficients

Let $(a_n, b_n)_{n \geq 1}$ be an i.i.d. sequence of random variable in $\mathbb{R}^2$ and let x_0 be an arbitrary constant. Define the sequence (x_n) recursively by the formula

$$x_n = a_n x_{n-1} + b_n, \quad n \geq 1.$$

Proposition 8 Assume that there exists $\delta \in]0, 1]$ such that

$$\mathbf{E} |a_1|^\delta < 1, \quad \mathbf{E} |b_1|^\delta < \infty.$$

Then for any bounded uniformly continuous function f

$$\mathbf{P}\text{-}\lim_N \frac{1}{N} \sum_{n=1}^N f(x_n) = \mathbf{E} f(\zeta),$$

where

$$\zeta = \sum_{k=1}^{\infty} b_k \prod_{j=1}^{k-1} a_j \quad \text{with} \quad \prod_{j=1}^0 a_j = 1.$$

The proof is given in [27].

Acknowledgements The research is funded by the grant of the Government of Russian Federation $n^\circ 14.A.12.31.0007$. The second author is partially supported by the department of Mathematics and Mechanics of National Research Tomsk State University.

References

1. Albrecher H., Badescu A., Landriault D. On the dual risk model with taxation, *Insurance: Mathematics and Economics*, **42** (2008), 1086–1094.
2. Asmussen S., Albrecher H. *Ruin Probabilities*. World Scientific, Singapore, 2010.
3. Avanzi B., Gerber H.U., Shiu E.S.W. Optimal dividends in the dual model. *Insurance: Mathematics and Economics*, **41** (2007), 111–123.
4. Bayraktar E., Egami M. Optimizing venture capital investments in a jump diffusion model. *Mathematical Methods of Operations Research*, **67** (2008), 1, 21–42.
5. Belkina T. Risky investments for insurers and sufficiency theorems for the survival probability. *Markov Processes Relat. Fields*, **20** (2014), 505–525.
6. Belkina T.A., Konyukhova N.B., Kurochkin S.V. Singular boundary value problem for the integrodifferential equation in an insurance model with stochastic premiums: Analysis and numerical solution. *Comp. Math. and Math. Physics*. **52** (2012), 10, 1384–1416.
7. Belkina T., Konyukhova N., Kurochkin S. Singular problems for integro-differential equations in dynamic insurance models. *Springer Proceedings in Mathematics. Proc. Intern. Conf. on Differential and Difference Equations and Applications (in honour of Prof. Ravi P. Agarwal)*, **47** (2013), 27–44.

8. Borodin A.N., Salminen P. *Handbook of Brownian Motion - Facts and Formulae*. Birkhäuser, 2002.
9. Dufresne, D. The distribution of a perpetuity, with application to risk theory and pension funding. *Scand. Actuarial J.*, (1990), 39–79.
10. Fedoryuk M.V. *Asymptotic Analysis: Linear Ordinary Differential Equations*. Springer, Berlin, 1993.
11. Feller, W. *An Introduction to Probability Theory and Its Applications*. Vol. 2, Wiley, New York, 1996.
12. Frolova A.G. Some mathematical models of risk theory. All-Russian School-Colloquium on Stochastic Methods in Geometry and Analysis. Abstracts, 1994, 117–118.
13. Frolova A., Kabanov Yu., Pergamenschikov S. In the insurance business risky investments are dangerous. *Finance and Stochastics*, **6** (2002), 227–235.
14. Gaier J., Granditz P. Ruin probabilities and investment under interest force in the presence of regularly varying tails. *Scand. Actuarial J.* (2004), 4, 256–278.
15. Goldie C.M. Implicit renewal theory and tails of solutions of random equation. *Ann. Appl. Probab.*, **1** (1991), 1, 126–166.
16. Grandell I. *Aspects of Risk Theory*. Springer, Berlin, 1990.
17. Granditz P. A Karamata-type theorem and ruin probabilities for an insurer investing proportionally in the stock market. *Insurance: Mathematics and Economics*, **34** (2004), 297–305.
18. Hult H., Lindskog F. Ruin probabilities under general investments and heavy-tailed claims. *Finance and Stochastics*, **15** (2011), 243–265.
19. Kalashnikov V. Ruin probability under random interest rate. Manuscript, 1999.
20. Kalashnikov V., Norberg R. Power tailed ruin probabilities in the presence of risky investments. *Stoch. Proc. Appl.*, **98** (2002), 211–228.
21. Matsumoto H., Yor M. Exponential functionals of Brownian motion, I: Probability laws at fixed time. *Probability Surveys*, **2** (2005), 312–347.
22. Nyrhinen H. On the ruin probabilities in a general economic environment. *Stoch. Proc. Appl.*, **83** (1999), 319–330.
23. Nyrhinen H. Finite and infinite time ruin probabilities in a stochastic economic environment. *Stoch. Proc. Appl.*, **92** (2001), 265–285.
24. Paulsen J. *Stochastic Calculus with Applications to Risk Theory*. Lecture Notes, Univ. of Bergen and Univ. of Copenhagen, 1996.
25. Paulsen J. Sharp conditions for certain ruin in a risk process with stochastic return on investments. *Stoch. Proc. Appl.*, **75** (1998), 135–148.
26. Paulsen J., Gjessing H.K. Ruin theory with stochastic return on investments. *Adv. Appl. Probab.*, **29** (1997), 4, 965–985.
27. Pergamenschikov S., Zeitouni O. Ruin probability in the presence of risky investments. *Stoch. Process. Appl.*, **116** (2006), 267–278.
28. Pergamenschikov S. Erratum to: "Ruin probability in the presence of risky investments" [*Stoch. Proc. Appl.*, **116** (2006), 267–278]. *Stoch. Proc. Appl.*, **119** (2009), 1, 305–306.
29. Revuz D., Yor M. *Continuous Martingales and Brownian Motion*. Springer, Berlin, 1999.
30. Saxén T. On the probability of ruin in the collective risk theory for insurance enterprises with only negative risk sums. *Scand. Actuarial J.* (1948), 1–2, 199–228.
31. Wang G., Wu R. Distributions for the risk process with a stochastic return on investments. *Stoch. Process. Appl.*, **95** (2001), 329–341.
32. Yor M. On some exponential functionals of Brownian motion. *Adv. Appl. Probab.*, **24** (1992), 3, 509–531.