The Generalized Franchetta Conjecture for Some Hyperkähler Fourfolds

Robert Laterveer

► To cite this version:

Robert Laterveer. The Generalized Franchetta Conjecture for Some Hyperkähler Fourfolds. Publications of the Research Institute for Mathematical Sciences, 2019, 55 (4), pp.859-893. 10.4171/PRIMS/55-4-8 . hal-02336792

HAL Id: hal-02336792

<https://hal.science/hal-02336792>

Submitted on 29 Oct 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THE GENERALIZED FRANCHETTA CONJECTURE FOR SOME HYPERKÄHLER FOURFOLDS

ROBERT LATERVEER

ABSTRACT. We obtain a “generalized Franchetta conjecture” type of statement for the Hilbert squares of low genus $K3$ surfaces, and for the Fano varieties of lines on certain cubic fourfolds.

1. INTRODUCTION

For a smooth projective variety X over $\mathbb{C}$, let $A^i(X) := CH^i(X)_{\mathbb{Q}}$ denote the Chow groups (i.e. the groups of codimension i algebraic cycles on X with $\mathbb{Q}$ -coefficients, modulo rational equivalence).

The world of Chow groups is still shrouded in mystery, its map containing vast underdeveloped regions only partially sketched in by conjectures [7], [21], [22], [23], [30], [45], [31]. One region on this map that holds particular interest is that of hyperkähler varieties (i.e. projective irreducible holomorphic symplectic manifolds [2], [1]). Here, recent years have seen significant progress in the understanding of Chow groups. Much of this progress centers around the following conjecture:

Conjecture 1.1 (Beauville, Voisin [5], [42]). *Let X be a hyperkähler variety. Let $D^*(X) \subset A^*(X)$ denote the $\mathbb{Q}$ -subalgebra generated by divisors and Chern classes of X . Then the cycle class maps induce injections*

$$D^i(X) \hookrightarrow H^{2i}(X, \mathbb{Q}) \quad \forall i.$$

(cf. [5], [42], [4], [8], [35], [10] for cases where conjecture 1.1 is satisfied.)

The “motivation” underlying conjecture 1.1 is that for a hyperkähler variety X , the Chow ring $A^*(X)$ is expected to have a bigrading $A^*_{[*]}(X)$, where the piece $A^i_{[j]}(X)$ corresponds to the graded $\mathrm{Gr}^j_F A^i(X)$ for the conjectural Bloch–Beilinson filtration. In particular, it is expected that the subring $A^*_{[0]}(X)$ injects into cohomology, and that $D^*(X) \subset A^*_{[0]}(X)$.

In addition to divisors and Chern classes, what other cycles should be in the subring $A^*_{[0]}(X)$ (assuming this subring exists) ? The following conjecture of Voisin provides more candidate members:

2010 *Mathematics Subject Classification*. Primary 14C15, 14C25, 14C30.

Key words and phrases. Algebraic cycles, Chow groups, motives, Bloch–Beilinson filtration, hyperkähler variety, Lagrangian subvariety, Fano variety of lines on cubic fourfold, $K3$ surface, Beauville’s splitting property, multiplicative Chow–Künneth decomposition, spread of algebraic cycles.

Author address. Institut de Recherche Mathématique Avancée, CNRS – Université de Strasbourg, 7 Rue René Descartes, 67084 Strasbourg CEDEX, FRANCE. robert.laterveer@math.unistra.fr.

Communicated by S. Mukai. Received February 15, 2017. Revised September 17, 2018.

Conjecture 1.2 (Voisin [47]). *Let X be a hyperkähler variety of dimension $n = 2m$. Let $Z \subset X$ be a codimension i subvariety swept out by i -dimensional constant cycle subvarieties. There exists a subring $A_{[0]}^*(X) \subset A^*(X)$ injecting into cohomology, and*

$$Z \in A_{[0]}^i(X) .$$

A *constant cycle subvariety* is by definition a closed subvariety $T \subset X$ such that the image of the natural map $A_0(T) \rightarrow A^n(X)$ has dimension 1. In particular, conjecture 1.2 stipulates that Lagrangian constant cycle subvarieties (i.e., constant cycle subvarieties of dimension m) should lie in $A_{[0]}^m(X)$.

Another conjecture concerns more generally m -dimensional subvarieties $Z \subset X$ that are *Lagrangian*, i.e. such that

$$\cup Z : H^{2,0}(X) \rightarrow H^{m+2,m}(X)$$

is the zero map. Since $H^{*,0}(X)$ is generated by $H^{2,0}(X)$, we have that

$$\cup Z : H^{j,0}(X) \rightarrow H^{m+j,m}(X)$$

is the zero map for all $j > 0$. Since conjecturally, the piece $A_{[j]}^j(X)$ is determined by $H^{j,0}(X)$, and the piece $A_{[j]}^{2m}(X)$ is determined by $H^{2m,2m-j}(X)$, we arrive at the following conjecture:

Conjecture 1.3. *Let X be a hyperkähler variety of dimension $2m$. Let $Z \subset X$ be a codimension m subvariety that is Lagrangian. Then the map*

$$A_{[j]}^j(X) \xrightarrow{\cdot Z} A^{m+j}(X) \xrightarrow{b} A^{2m}(X) \rightarrow A_{[j]}^{2m}(X)$$

is zero for all $j > 0$ and all $b \in A^{m-j}(X)$. (Here, the last arrow is projection to the piece $A_{[j]}^{2m}(X)$.)

The goal of this note is to provide some examples where conjecture 1.2 and 1.3 are satisfied, by looking at Hilbert squares of $K3$ surfaces. Here, thanks to the *Fourier decomposition* of Shen–Vial [38], the Chow groups of X split in a finite number of pieces $A_{(*)}^*(X)$.¹

The main result of this note is a “generalized Franchetta conjecture” type of statement for certain Hilbert squares X . The statement is that codimension 2 cycles that can be defined relatively live in the subring $A_{(0)}^*(X)$:

Theorem (=theorem 3.1). *Let $\mathcal{S} \rightarrow B$ denote the universal family of $K3$ surfaces of genus $g \in \{2, 3, 4, 5, 6, 8\}$. Let $\mathcal{X} = \mathcal{S}^{[2]} \rightarrow B$ denote the universal family of Hilbert squares of genus g $K3$ surfaces (so a fibre X_b is the Hilbert square of a genus g $K3$ surface). Let $\Gamma \in A^2(\mathcal{X})$ be a relative cycle. Then*

$$\Gamma|_{X_b} \in A_{(0)}^2(X_b) \quad \forall b \in B .$$

(Here $A_{(0)}^2()$ refers to the Fourier decomposition of Shen–Vial [38].)

¹NB: we will use the notation $A_{(*)}^*(\cdot)$ for the bigrading that is constructed unconditionally in [38] for certain hyperkähler fourfolds. The notation $A_{[*]}^*(\cdot)$, that occurs only in this introduction, refers to a *conjectural* bigrading with the property that $A_{[j]}^i(X)$ is related to the graded $\mathrm{Gr}_F^j A^i(X)$ for the conjectural Bloch–Beilinson filtration. Hence, the (unconditionally existing) bigrading $A_{(*)}^*(\cdot)$ is a candidate for the (only ideally existing) bigrading $A_{[*]}^*(\cdot)$.

A second result addresses conjecture 1.3:

Theorem (=theorem 3.2). *Let $\mathcal{X} \rightarrow B$ be as in theorem 3.1. Let $\Gamma \subset \mathcal{X}$ be a codimension 2 subvariety. Assume the restriction*

$$\Gamma_b := \Gamma|_{X_b} \in A^2(X_b)$$

is Lagrangian (i.e., $\cup \Gamma_b: H^{2,0}(X_b) \rightarrow H^{4,2}(X_b)$ is the zero map), for very general $b \in B$. Then

$$\cdot \Gamma_b: A_{hom}^2(X_b) \rightarrow A^4(X_b)$$

is the zero map, for all $b \in B$.

Section 4 presents some concrete applications of these results. Here is one of them:

Corollary (=corollary 4.1). *Let S be a general $K3$ surface of genus 5, and let $X = S^{[2]}$ be the Hilbert square. Let $A \subset X$ be a general fibre of the Lagrangian fibration $\phi: X \rightarrow \mathbb{P}^2$ of proposition 2.25. Let $b \in A^4(X)$ be a 0-cycle of the form*

$$b = A \cdot a + p \in A^4(X),$$

where $a \in A^2(X)$ and p is a sum of intersections of divisors and Chern classes of X . Then b is rationally trivial if and only if b is of degree 0.

The proof of theorems 3.1 and 3.2 is based on Voisin’s method of “spread” of algebraic cycles [43], [44], [45], [46], combined with results of Pavic–Shen–Yin on the generalized Franchetta conjecture for $K3$ surfaces [33].

It would be interesting to extend theorem 3.1 to Hilbert squares of $K3$ surfaces of higher genus, and also to other hyperkähler varieties.² For starters, we prove this for Fano varieties of cubic fourfolds containing two planes (theorem 5.1).

Conventions. *In this article, the word variety will refer to a reduced irreducible scheme of finite type over $\mathbb{C}$. A subvariety is a (possibly reducible) reduced subscheme which is equidimensional.*

All Chow groups will be with rational coefficients: *we will denote by $A_j(X)$ the Chow group of j -dimensional cycles on X with $\mathbb{Q}$ -coefficients; for X smooth of dimension n the notations $A_j(X)$ and $A^{n-j}(X)$ are used interchangeably.*

The notations $A_{hom}^j(X)$, $A_{AJ}^j(X)$ will be used to indicate the subgroups of homologically trivial, resp. Abel–Jacobi trivial cycles. For a morphism $f: X \rightarrow Y$, we will write $\Gamma_f \in A_(X \times Y)$ for the graph of f . The contravariant category of Chow motives (i.e., pure motives with respect to rational equivalence as in [37], [31]) will be denoted $\mathcal{M}_{rat}$.*

We will use $H^j(X)$ to indicate singular cohomology $H^j(X, \mathbb{Q})$.

2. PRELIMINARIES

2.1. The Fourier decomposition.

²Results of this kind can be found in [12], which was written after the present paper.

Theorem 2.1 (Shen–Vial [38]). *Let S be a K3 surface, and let $X = S^{[2]}$ be the Hilbert scheme of length 2 subschemes of S . There is a decomposition*

$$A^i(X) = \bigoplus_{\substack{0 \leq j \leq i \\ j \text{ even}}} A_{(j)}^i(X) ,$$

with the following properties:

- (i) $A_{(*)}^*(X)$ is a bigraded ring;
- (ii) $A_{(j)}^i(X) \subset A_{hom}^i(X)$ for $j > 0$.

Proof. This is essentially [38, Theorem 2], combined with the fact that there is a class $L \in A^2(X \times X)$ lifting the Beauville–Bogomolov class and satisfying certain equalities, which is [38, Part 2]. $\square$

Theorem 2.2 (Shen–Vial [38]). *Let $Y \subset \mathbb{P}^5(\mathbb{C})$ be a smooth cubic fourfold, and let $X = F(Y)$ be the Fano variety of lines in Y . There is a decomposition*

$$A^i(X) = \bigoplus_{\substack{0 \leq j \leq i \\ j \text{ even}}} A_{(j)}^i(X) ,$$

with the following properties:

- (i) $A_{(j)}^i(X) \subset A_{hom}^i(X)$ for $j > 0$;
- (ii) if Y is very general, $A_{(*)}^*(X)$ is a bigraded ring.

Proof. This follows again from [38, Theorem 2]. Point (ii) is [38, Theorem 3]. $\square$

2.2. MCK decomposition.

Definition 2.3 (Murre [30]). *Let X be a smooth projective variety of dimension n . We say that X has a CK decomposition if there exists a decomposition of the diagonal*

$$\Delta_X = \pi_0 + \pi_1 + \cdots + \pi_{2n} \quad \text{in } A^n(X \times X) ,$$

such that the π_i are mutually orthogonal idempotents in $A^n(X \times X)$ and $(\pi_i)_ H^*(X) = H^i(X)$.*

(NB: “CK decomposition” is shorthand for “Chow–Künneth decomposition”.)

Remark 2.4. *The existence of a CK decomposition for any smooth projective variety is part of Murre’s conjectures [30], [21], [22].*

Definition 2.5 (Shen–Vial [38]). *Let X be a smooth projective variety of dimension n . Let $\Delta_X^{sm} \in A^{2n}(X \times X \times X)$ be the class of the small diagonal*

$$\Delta_X^{sm} := \{(x, x, x) \mid x \in X\} \subset X \times X \times X .$$

An MCK decomposition is a CK decomposition $\{\pi_i^X\}$ of X that is multiplicative, i.e. it satisfies

$$\pi_k^X \circ \Delta_X^{sm} \circ (\pi_i^X \times \pi_j^X) = 0 \quad \text{in } A^{2n}(X \times X \times X) \quad \text{for all } i + j \neq k .$$

(NB: “MCK decomposition” is shorthand for “multiplicative Chow–Künneth decomposition”.)

A weak MCK decomposition is a CK decomposition $\{\pi_i^X\}$ of X that satisfies

$$\left(\pi_k^X \circ \Delta_X^{sm} \circ (\pi_i^X \times \pi_j^X) \right)_* (a \times b) = 0 \quad \text{for all } a, b \in A^*(X) \text{ and for all } i + j \neq k .$$

Remark 2.6. *The small diagonal (seen as a correspondence from $X \times X$ to X) induces the multiplication morphism*

$$\Delta_X^{sm}: h(X) \otimes h(X) \rightarrow h(X) \text{ in } \mathcal{M}_{\text{rat}}.$$

Suppose X has a CK decomposition

$$h(X) = \bigoplus_{i=0}^{2n} h^i(X) \text{ in } \mathcal{M}_{\text{rat}}.$$

By definition, this decomposition is multiplicative if for any i, j the composition

$$h^i(X) \otimes h^j(X) \rightarrow h(X) \otimes h(X) \xrightarrow{\Delta_X^{sm}} h(X) \text{ in } \mathcal{M}_{\text{rat}}$$

factors through $h^{i+j}(X)$.

If X has a weak MCK decomposition, then setting

$$A_{(j)}^i(X) := (\pi_{2i-j}^X)_* A^i(X),$$

one obtains a bigraded ring structure on the Chow ring: that is, the intersection product sends $A_{(j)}^i(X) \otimes A_{(j')}^{i'}(X)$ to $A_{(j+j')}^{i+i'}(X)$.

It is expected (but not proven !) that for any X with a weak MCK decomposition, one has

$$A_{(j)}^i(X) \stackrel{??}{=} 0 \text{ for } j < 0, \quad A_{(0)}^i(X) \cap A_{\text{hom}}^i(X) \stackrel{??}{=} 0;$$

this is related to Murre's conjectures B and D, that have been formulated for any CK decomposition [30].

The property of having an MCK decomposition is severely restrictive, and is closely related to Beauville's "(weak) splitting property" [5]. For more ample discussion, and examples of varieties with an MCK decomposition, we refer to [38, Section 8], as well as [41], [39], [13].

Lemma 2.7. *Let X, X' be birational hyperkähler varieties. Then X has an MCK decomposition if and only if X' has one.*

Proof. This is noted in [41, Introduction]; the idea is that Rieß's result [34] implies that X and X' have isomorphic Chow motives and the isomorphism is compatible with the multiplicative structure. $\square$

2.3. Relative MCK for $S^{[m]}$ and for S^m .

Theorem 2.8 (Vial [41]). *Let S be a projective K3 surface, and let $X = S^{[m]}$ be the Hilbert scheme of length m subschemes of S . Then X has a self-dual MCK decomposition $\{\Pi_i^X\}$. In particular, $A^*(X) = A_{(*)}^*(X)$ is a bigraded ring, where*

$$A^i(X) = \bigoplus_{j=2i-2n}^i A_{(j)}^i(X),$$

and $A_{(j)}^i(X) = 0$ for j odd. In case $m = 2$, the bigrading $A_{()}^*(X)$ coincides with the one given by the Fourier decomposition of theorem 2.1.*

Proof. This is [41, Theorems 1 and 2]. The last statement is [38, Theorem 15.8], plus the fact that for $m = 2$ the MCK decomposition of [41] coincides with the one of [38]. $\square$

Notation 2.9. Let $\mathcal{S} \rightarrow B$ be a family (i.e., a smooth projective morphism). For $r \in \mathbb{N}$, we write $\mathcal{S}^{r/B}$ for the relative r -fold fibre product

$$\mathcal{S}^{r/B} := \mathcal{S} \times_B \mathcal{S} \times_B \cdots \times_B \mathcal{S}$$

(r copies of $\mathcal{S}$).

Proposition 2.10. Let $\mathcal{S} \rightarrow B$ be a family of K3 surfaces. There exist relative correspondences

$$\Pi_j^{\mathcal{S}^{m/B}} \in A^{2m}(\mathcal{S}^{m/B} \times \mathcal{S}^{m/B}) \quad (j = 0, 2, 4, \dots, 4m),$$

such that for each $b \in B$, the restriction

$$\Pi_j^{(S_b)^m} := \Pi_j^{\mathcal{S}^{m/B}}|_{(S_b)^{2m}} \in A^4((S_b)^m \times (S_b)^m)$$

defines a self-dual MCK decomposition for $(S_b)^m$.

Proof. On any K3 surface S_b , there is the distinguished 0-cycle $\mathfrak{o}_{S_b}$ such that $c_2(S_b) = 24\mathfrak{o}_{S_b}$ [4]. Let $p_i: \mathcal{S}^{m/B} \rightarrow \mathcal{S}$, $i = 1, \dots, m$, denote the projections to the two factors. Let $T_{\mathcal{S}/B}$ denote the relative tangent bundle. The assignment

$$\begin{aligned} \Pi_0^{\mathcal{S}} &:= (p_1)^* \left(\frac{1}{24} c_2(T_{\mathcal{S}/B}) \right) \in A^2(\mathcal{S} \times_B \mathcal{S}), \\ \Pi_4^{\mathcal{S}} &:= (p_2)^* \left(\frac{1}{24} c_2(T_{\mathcal{S}/B}) \right) \in A^2(\mathcal{S} \times_B \mathcal{S}), \\ \Pi_2^{\mathcal{S}} &:= \Delta_{\mathcal{S}} - \Pi_0^{\mathcal{S}} - \Pi_4^{\mathcal{S}} \end{aligned}$$

defines (by restriction) an MCK decomposition for each fibre, i.e.

$$\Pi_j^{S_b} := \Pi_j^{\mathcal{S}}|_{S_b \times S_b} \in A^2(S_b \times S_b) \quad (j = 0, 2, 4)$$

is an MCK decomposition for any $b \in B$ [38, Example 8.17].

Next, we consider the m -fold relative fibre product $\mathcal{S}^{m/B}$. Let

$$p_{i,j}: \mathcal{S}^{2m/B} \rightarrow \mathcal{S}^{2/B} \quad (1 \leq i < j \leq 2m)$$

denote projection to the i -th and j -th factor. We define

$$\begin{aligned} \Pi_j^{\mathcal{S}^{m/B}} &:= \sum_{k_1+k_2+\dots+k_m=j} (p_{1,m+1})^*(\Pi_{k_1}^{\mathcal{S}}) \cdot (p_{2,m+2})^*(\Pi_{k_2}^{\mathcal{S}}) \cdot \dots \cdot (p_{m,2m})^*(\Pi_{k_m}^{\mathcal{S}}) \\ &\in A^{2m}(\mathcal{S}^{2m/B}), \quad (j = 0, 2, 4, \dots, 4m). \end{aligned}$$

By construction, the restriction to each fibre induces an MCK decomposition (the “product MCK decomposition”)

$$\begin{aligned} \Pi_j^{(S_b)^m} &:= \Pi_j^{\mathcal{S}^{m/B}}|_{(S_b)^{2m}} = \sum_{k_1+k_2+\dots+k_m=j} \Pi_{k_1}^{S_b} \times \Pi_{k_2}^{S_b} \times \dots \times \Pi_{k_m}^{S_b} \in A^{2m}((S_b)^{2m}), \\ &\quad (j = 0, 2, 4, \dots, 4m). \end{aligned}$$

$\square$

Remark 2.11. Let $\mathcal{S} \rightarrow B$ be a family of K3 surfaces. Let $\mathcal{X} \rightarrow B$ denote the relative Hilbert scheme (i.e., the blow-up of $\mathcal{S} \times_B \mathcal{S}$ along the relative diagonal, quotiented by the natural action of the symmetric group on 2 elements $\mathfrak{S}_2$). Since the above construction of the $\Pi_j^{\mathcal{S}^{2/B}}$ is $\mathfrak{S}_2$ -invariant, it induces relative correspondences

$$\Pi_j^{\mathcal{X}} \in A^4(\mathcal{X} \times_B \mathcal{X}) \quad (j = 0, 2, 4, 6, 8),$$

with the property that the restrictions

$$\Pi_j^{X_b} := \Pi_j^{\mathcal{X}}|_{X_b \times X_b} \in A^4(X_b \times X_b)$$

form an MCK decomposition for X_b , for all $b \in B$ (cf. [25, Proposition 2.16] for a detailed proof).

Proposition 2.12. Let $\mathcal{S} \rightarrow B$ be a family of K3 surfaces. There exist relative correspondences

$$\Theta'_1, \dots, \Theta'_m \in A^{2m}(\mathcal{S}^{m/B} \times_B \mathcal{S}), \quad \Xi'_1, \dots, \Xi'_m \in A^2(\mathcal{S} \times_B \mathcal{S}^{m/B})$$

such that for each $b \in B$, the composition

$$\begin{aligned} A_{(2)}^{2m}((S_b)^m) &\xrightarrow{((\Theta'_1|_{(S_b)^{m+1}})^*, \dots, (\Theta'_m|_{(S_b)^{m+1}})^*)} A^2(S_b) \oplus \dots \oplus A^2(S_b) \\ &\xrightarrow{((\Xi'_1 + \dots + \Xi'_m)|_{(S_b)^{m+1}})^*} A^{2m}((S_b)^m) \end{aligned}$$

is the identity.

Proof. As before, let

$$p_{i,j}: \mathcal{S}^{2m/B} \rightarrow \mathcal{S}^{2/B} \quad (1 \leq i < j \leq 2m)$$

denote projection to the i -th and j -th factor, and let

$$p_i: \mathcal{S}^{m/B} \rightarrow \mathcal{S} \quad (1 \leq i \leq m)$$

denote projection to the i -th factor.

We now claim that for each $b \in B$, there is equality

(1)

$$\begin{aligned} (\Pi_{4m-2}^{\mathcal{S}^{m/B}})|_{(S_b)^{2m}} &= \frac{1}{2^{4m-2}} \left({}^t\Gamma_{p_1} \circ \Pi_2^{\mathcal{S}} \circ \Gamma_{p_1} \circ ((p_{1,m+1})^*(\Delta_{\mathcal{S}}) \cdot \prod_{\substack{2 \leq j \leq 2m \\ j \neq m+1}} (p_j)^* c_2(T_{\mathcal{S}/B})) \right. \\ &\quad \left. + \dots + {}^t\Gamma_{p_m} \circ \Pi_2^{\mathcal{S}} \circ \Gamma_{p_m} \circ ((p_{m,2m})^*(\Delta_{\mathcal{S}}) \cdot \prod_{\substack{1 \leq j \leq 2m-1 \\ j \neq m}} (p_j)^* c_2(T_{\mathcal{S}/B})) \right) |_{(S_b)^{2m}} \\ &\quad \text{in } A^{2m}((S_b)^m \times (S_b)^m). \end{aligned}$$

Indeed, using Lieberman's lemma [14, 16.1.1], we find that

$$\begin{aligned} ({}^t\Gamma_{p_1} \circ \Pi_2^{\mathcal{S}} \circ \Gamma_{p_1})|_{(S_b)^{2m}} &= (({}^t\Gamma_{p_{1,m+1}})_*(\Pi_2^{\mathcal{S}}))|_{(S_b)^{2m}} = ((p_{1,m+1})^*(\Pi_2^{\mathcal{S}}))|_{(S_b)^{2m}}, \\ &\vdots \\ ({}^t\Gamma_{p_m} \circ \Pi_2^{\mathcal{S}} \circ \Gamma_{p_m})|_{(S_b)^{2m}} &= (({}^t\Gamma_{p_{m,2m}})_*(\Pi_2^{\mathcal{S}}))|_{(S_b)^{2m}} = ((p_{m,2m})^*(\Pi_2^{\mathcal{S}}))|_{(S_b)^{2m}}. \end{aligned}$$

Let us now (by way of example) consider the first summand of the right-hand side of (1). For brevity, let

$$P: (S_b)^{3m} \rightarrow (S_b)^{2m}$$

denote the projection on the first m and last m factors. Writing out the definition of composition of correspondences, we find that

$$\begin{aligned} & \frac{1}{24^{2m-2}} \left({}^t\Gamma_{p_1} \circ \Pi_2^S \circ \Gamma_{p_1} \circ ((p_{1,m+1})^*(\Delta_S) \cdot \prod_{\substack{2 \leq j \leq 2m \\ j \neq m+1}} (p_j)^* c_2(T_{S/B})) \right) |_{(S_b)^{2m}} = \\ & \frac{1}{24^{2m-2}} \left(((p_{1,m+1})^*(\Pi_2^{S_b})) \circ ((p_{1,m+1})^*(\Delta_{S_b}) \cdot \prod_{\substack{2 \leq j \leq 2m \\ j \neq m+1}} (p_j)^* c_2(T_{S_b})) \right) = \\ & P_* \left(((\Delta_{S_b})_{(1,m+1)} \times \mathfrak{o}_{S_b} \times \cdots \times \mathfrak{o}_{S_b} \times S_b \times \cdots \times S_b) \cdot \right. \\ & \quad \left. (S_b \times \cdots \times S_b \times (\Pi_2^{S_b})_{(m+1,2m+1)} \times S_b \times \cdots \times S_b) \right) = \\ & P_* \left(((\Delta_{S_b} \times S_b) \cdot (S_b \times \Pi_2^{S_b}))_{(1,m+1,2m+1)} \times \mathfrak{o}_{S_b} \times \cdots \times \mathfrak{o}_{S_b} \times S_b \times \cdots \times S_b \right) = \\ & \Pi_2^{S_b} \times \Pi_4^{S_b} \times \cdots \times \Pi_4^{S_b} \quad \text{in } A^{2m}((S_b)^m \times (S_b)^m). \end{aligned}$$

(Here, we use the notation $(C)_{(i,j)}$ to indicate that the cycle C lies in the i th and j th factor, and likewise for $(D)_{(i,j,k)}$.)

Doing the same for the other summands in (1), one convinces oneself that both sides of (1) are equal to the fibrewise product Chow–Künneth component

$$\Pi_{4m-2}^{(S_b)^m} = \Pi_2^{S_b} \times \Pi_4^{S_b} \times \cdots \times \Pi_4^{S_b} + \cdots + \Pi_4^{S_b} \times \cdots \times \Pi_4^{S_b} \times \Pi_2^{S_b} \in A^{2m}((S_b)^m \times (S_b)^m),$$

thus proving the claim.

Let us now define

$$\Theta'_i := \frac{1}{24^{2m-2}} \Gamma_{p_i} \circ ((p_{i,m+i})^*(\Delta_S) \cdot \prod_{\substack{j \in [1,2m] \\ j \notin \{i,m+i\}}} (p_j)^* c_2(T_{S/B})) \in A^{2m}((\mathcal{S}^{m/B}) \times_B \mathcal{S}),$$

$$\Xi'_i := {}^t\Gamma_{p_i} \circ \Pi_2^S \in A^2(\mathcal{S} \times_B (\mathcal{S}^{m/B})),$$

where $1 \leq i \leq m$. It follows from equation (1) that there is equality

$$(2) \quad \left((\Xi'_1 \circ \Theta'_1 + \cdots + \Xi'_m \circ \Theta'_m) |_{(S_b)^{2m}} \right)_* = (\Pi_{4m-2}^{(S_b)^m})_* : A_{(j)}^i((S_b)^m) \rightarrow A_{(j)}^i((S_b)^m) \quad \forall b \in B \quad \forall (i,j).$$

Taking $(i,j) = (2m, 2)$, this proves the proposition. $\square$

The following is a version of proposition 2.12 for the group $A_{(2)}^2((S_b)^m)$:

Proposition 2.13. *Let $\mathcal{S} \rightarrow B$ be a family of K3 surfaces. There exist relative correspondences*

$$\Theta_1, \dots, \Theta_m \in A^{2m}(\mathcal{S} \times_B (\mathcal{S}^{m/B})), \quad \Xi_1, \dots, \Xi_m \in A^2((\mathcal{S}^{m/B}) \times_B \mathcal{S})$$

such that for each $b \in B$, the composition

$$A_{(2)}^2((S_b)^m) \xrightarrow{((\Xi_1|_{(S_b)^{m+1}})_*, \dots, (\Xi_m|_{(S_b)^{m+1}})_*)} A^2(S_b) \oplus \dots \oplus A^2(S_b) \\ \xrightarrow{((\Theta_1 + \dots + \Theta_m)|_{(S_b)^{m+1}})_*} A^2((S_b)^m)$$

is the identity.

Proof. One may take

$$\Theta_i := {}^t\Theta'_i \in A^{2m}(\mathcal{S} \times_B (\mathcal{S}^{m/B})) , \\ \Xi_i := {}^t\Xi'_i \in A^2((\mathcal{S}^{m/B}) \times_B \mathcal{S}) \quad (i = 1, \dots, m) .$$

By construction, the product MCK decomposition $\{\Pi_i^{(S_b)^m}\}$ satisfies

$$\Pi_2^{(S_b)^m} = {}^t(\Pi_{4m-2}^{(S_b)^m}) \text{ in } A^{2m}((S_b)^m \times (S_b)^m) .$$

Hence, the transpose of equation (2) gives the equality

$$(\Pi_2^{(S_b)^m})_* = ({}^t(\Pi_{4m-2}^{(S_b)^m}))_* = ({}^t\Theta'_1 \circ {}^t\Xi'_1 + \dots + {}^t\Theta'_m \circ {}^t\Xi'_m)_* : \\ A_{(j)}^i((S_b)^m) \rightarrow A_{(j)}^i((S_b)^m) \quad \forall b \in B \quad \forall (i, j) .$$

Taking $(i, j) = (2, 2)$, this proves the proposition. $\square$

2.4. Relative CK for Fano varieties.

Theorem 2.14 (Shen–Vial [38]). *Let $Y \subset \mathbb{P}^5(\mathbb{C})$ be a smooth cubic, and let $X = F(Y)$ be the Fano variety of lines in X . The Fourier decomposition is induced by a CK decomposition $\{\Pi_i^X\}$, i.e. there is equality*

$$A_{(j)}^i(X) = (\Pi_{2i-j}^X)_* A^i(X) \quad \forall i, j .$$

In particular, for very general Y , the $\{\Pi_i^X\}$ are a weak MCK decomposition.

Proof. This is [38, Theorem 3.3]. The last statement follows from theorem 2.2(ii). $\square$

Notation 2.15. Let

$$\mathcal{Y} \rightarrow B$$

denote the universal family of smooth cubic fourfolds Y_b . Here B is a Zariski open in the parameter space $\mathbb{P}H^0(\mathbb{P}^5, \mathcal{O}_{\mathbb{P}^5}(3))$. Let

$$\mathcal{X} := \{(\ell, b) \mid \ell \subset Z_b\} \subset G(1, 5) \times B$$

be the corresponding family of Fano varieties of lines. (Here $G(1, 5)$ denotes the Grassmannian of lines in $\mathbb{P}^5$.) A fibre X_b of $\mathcal{X} \rightarrow B$ is a Fano variety of lines on a cubic Y_b .

Proposition 2.16. *Let $\mathcal{X} \rightarrow B$ be as in notation 2.15. There exist relative correspondences*

$$\Pi_i^{\mathcal{X}} \in A^4(\mathcal{X} \times_B \mathcal{X}) \quad (i = 2, 6) ,$$

with the property that for each $b \in B$, one has

$$((\Pi_2^{\mathcal{X}})|_{X_b \times X_b})_* = (\Pi_2^{X_b})_* : A^2(X_b) \rightarrow A^2(X_b) , \\ ((\Pi_6^{\mathcal{X}})|_{X_b \times X_b})_* = (\Pi_6^{X_b})_* : A^4(X_b) \rightarrow A^4(X_b) .$$

Here $\Pi_i^{X_b}$ is the Chow–Künneth decomposition of theorem 2.2(i).

Proof. The main point is that the Shen–Vial cycle $L \in A^2(X_b \times X_b)$ furnishing the Fourier decomposition [38] exists relatively: this is because by definition

$$L := \frac{1}{3}(g_1^2 + \frac{3}{2}g_1g_2 + g_2^2 - c_1 - c_2) - I \in A^2(X_b \times X_b)$$

[38, Equation (107)]. Here, $g := -c_1(\mathcal{E}_2) \in A^1(X_b)$ and $c := c_2(\mathcal{E}_2) \in A^2(X_b)$ (and $\mathcal{E}_2$ is the restriction of the rank 2 tautological bundle on the Grassmannian), and $g_i := (p_i)^*(g)$, $c_i := (p_i)^*(c)$ (where $p_i: X_b \times X_b \rightarrow X_b$ is projection on the i th factor), and $I \subset X_b \times X_b$ is the incidence correspondence. Since g_i, c_i and I obviously exist relatively, the same goes for L , i.e. there exists a relative correspondence

$$\mathcal{L} \in A^2(\mathcal{X} \times_B \mathcal{X})$$

with the property that for any $b \in B$ the restriction

$$\mathcal{L}|_{X_b \times X_b} \in A^2(X_b \times X_b)$$

is the Shen–Vial class L of [38]. This implies that the class $\ell \in A^2(X_b)$ of [38] (mentioned in theorem 2.17(ii) below) also exists relatively: it is defined as

$$\ell := (i_\Delta)^*(\mathcal{L}) \in A^2(\mathcal{X}),$$

where $i_\Delta: \mathcal{X} \rightarrow \mathcal{X} \times_B \mathcal{X}$ denotes the embedding along the relative diagonal. (NB: this makes sense because i_Δ is a regular embedding.) Next, the classes $\ell_i := (p_i)^*(\ell) \in A^2(X_b \times X_b)$ of [38] also exist relatively.

Armed with these facts, let us inspect the construction of the $\{\Pi_i^{X_b}\}$ in [38, Theorem 3.3].

As a first approach towards the construction of $\Pi_2^{X_b}$ and $\Pi_6^{X_b}$, Shen–Vial define

$$p_b := \frac{1}{25}L \cdot \ell_2 \in A^4(X_b \times X_b).$$

Again, by the above remarks the cycle p_b exists relatively (i.e. there is $\mathbf{p} \in A^4(\mathcal{X} \times_B \mathcal{X})$ which restricts to p_b on each fibre). We define $\Pi_6^{\mathcal{X}} := \mathbf{p} \in A^4(\mathcal{X} \times_B \mathcal{X})$. This does the job, for it is shown in [38, Proof of Theorem 3.3] that $(p_b)_*$ acts as the identity on $A_{(2)}^4(X_b)$ and acts as 0 on $A_{(0)}^4(X_b) \oplus A_{(4)}^4(X_b)$.

We define $\Pi_2^{\mathcal{X}}$ as the transpose $\Pi_2^{\mathcal{X}} := {}^t\Pi_6^{\mathcal{X}}$. This does the job, for it is shown in loc. cit. that $(p_b)^*$ acts as the identity on $A_{(2)}^2(X_b)$ and acts as 0 on $A_{(0)}^2(X_b)$. $\square$

2.5. A result in the Chow ring of $S^{[2]}$.

Theorem 2.17 (Shen–Vial [38]). *Let $X = S^{[2]}$ where S is a K3 surface. There exists a class $\ell \in A_{(0)}^2(X)$ inducing an isomorphism*

$$\cdot\ell: A_{(2)}^2(X) \xrightarrow{\cong} A_{(2)}^4(X).$$

The inverse isomorphism is given by

$$\frac{1}{25}L_*: A_{(2)}^4(X) \xrightarrow{\cong} A_{(2)}^2(X),$$

where $L \in A^2(X \times X)$ is the Shen–Vial class lifting the Beauville–Bogomolov form.

Proof. This is [38, Theorems 2.2 and 2.4] (combined with the fact that X satisfies the hypotheses of loc. cit., which is [38, Part 2]). $\square$

For later use, we remark that the Shen–Vial class L exists relatively:

Lemma 2.18. *Let $\mathcal{S} \rightarrow B$ be a family of K3 surfaces, and let $\mathcal{X} \rightarrow B$ be the family of Hilbert squares. There exists a class $\mathcal{L} \in A^2(\mathcal{X} \times_B \mathcal{X})$ such that for any $b \in B$, the restriction*

$$L_b := \mathcal{L}|_{X_b \times X_b} \in A^2(X_b \times X_b)$$

is the Shen–Vial class of theorem 2.17.

Proof. For a given K3 surface $S = S_b$, let

$$\begin{array}{ccc} Z & \xrightarrow{q} & S \\ \downarrow p & & \\ X & & \end{array}$$

denote the universal family of length 2 subschemes of S . Let

$$I := {}^t Z \circ Z \in A^2(X \times X)$$

denote the incidence correspondence. Let $\delta \in A^1(X)$ be the class supported on the exceptional divisor as in [38, Section 10], and let $\delta_i := (p_i)^*(\delta) \in A^1(X \times X)$, where $p_i: X \times X \rightarrow X$ is projection on the i th factor. Let $\mathfrak{o}_S \in A^2(S)$ denote the distinguished point [4], and let $S_{\mathfrak{o}} := p_* q^*(\mathfrak{o}_S) \in A^2(X)$.

By definition, the Shen–Vial class L is

$$L = I - 2(p_1)^*(S_{\mathfrak{o}}) - 2(p_2)^*(S_{\mathfrak{o}}) - \frac{1}{2}\delta_1\delta_2 \in A^2(X \times X)$$

[38, Equation (92)].

The distinguished point $\mathfrak{o}_S$ equals $\frac{1}{24}c_2(T_S)$ [4], hence it can be defined relatively. The universal family Z can also be defined relatively, hence the same holds for the incidence correspondence I . This implies that the class L can be defined relatively, i.e. there exists a class $\mathcal{L}$ as indicated. $\square$

2.6. A result in the Chow ring of the Fano variety.

Theorem 2.19 (Shen–Vial [38]). *Let $Z \subset \mathbb{P}^5(\mathbb{C})$ be a smooth cubic fourfold, and let X be the Fano variety of lines in Z . There is a distinguished class $l \in A_{(0)}^2(X)$ such that intersection induces an isomorphism*

$$\cdot l: A_{(2)}^2(X) \xrightarrow{\cong} A_{(2)}^4(X).$$

The inverse isomorphism is given by

$$\frac{1}{25}L_*: A_{(2)}^4(X) \xrightarrow{\cong} A_{(2)}^2(X),$$

where $L \in A^2(X \times X)$ is the Shen–Vial class lifting the Beauville–Bogomolov form.

Proof. This follows from [38, Theorems 2.2 and 2.4] (combined with the fact that X satisfies the hypotheses of loc. cit., which is [38, Part 3]. $\square$

The following is a reformulation of theorem 2.19:

Proposition 2.20. *Let $Y \subset \mathbb{P}^5(\mathbb{C})$ be a smooth cubic fourfold, and let X be the Fano variety of lines in Y . Let $I \in A^2(X \times X)$ be the incidence correspondence, and let $g = -c_1(\mathcal{E}_2) \in A^1(X)$ be the Plücker polarization. Then*

$$\cdot g^2: A_{(2)}^2(X) \rightarrow A_{(2)}^4(X)$$

is an isomorphism. The inverse isomorphism is given by

$$-\frac{1}{6}I_*: A_{(2)}^4(X) \rightarrow A_{(2)}^2(X).$$

Proof. This is implicit in the arguments of [38]. For any $a \in A_{hom}^4(X)$, there is equality

$$\ell \cdot L_*(a) = -\frac{25}{6}g^2 \cdot I_*(a) \quad \text{in } A^4(X).$$

(This follows from [38, Equations (107) and (108)], cf. the proof of [38, Proposition 19.4].) But for $a \in A_{(2)}^4(X)$, we know (theorem 2.17(ii)) that $\ell \cdot L_*(a) = 25a$, and so for any $a \in A_{(2)}^4(X)$ we get an equality

$$a = -\frac{1}{6}g^2 \cdot I_*(a) \quad \text{in } A^4(X).$$

Applying I_* to this equality, we obtain an equality (for any $a \in A_{(2)}^4(X)$)

$$(3) \quad I_*(a) = -\frac{1}{6}I_*(g^2 \cdot I_*(a)) \quad \text{in } A^2(X).$$

But we know that

$$A_{(2)}^2(X) = I_*A_{hom}^4(X) = I_*A_{(2)}^4(X).$$

(Here, the first equality is [38, Proof of Proposition 21.10], and the second equality follows from the fact that $I_*A_{(4)}^4(X) = 0$ [38, Theorem 20.5].) Equation (3) thus becomes the statement that for any $b \in A_{(2)}^2(X)$ there is equality

$$b = -\frac{1}{6}I_*(g^2 \cdot b) \quad \text{in } A^2(X).$$

This proves the proposition. $\square$

Corollary 2.21. *Let $Y \subset \mathbb{P}^5(\mathbb{C})$ be a smooth cubic fourfold, and let X be the Fano variety of lines in Y . There exist correspondences $P \in A^3(X \times Y)$, $Q \in A^5(Y \times X)$ such that*

$$\begin{aligned} ({}^tP \circ {}^tQ)_* &= \text{id}: A_{(2)}^2(X) \rightarrow A_{(2)}^2(X), \\ (Q \circ P)_* &= \text{id}: A_{(2)}^4(X) \rightarrow A_{(2)}^4(X). \end{aligned}$$

Moreover, let $\mathcal{Y} \rightarrow B$ and $\mathcal{X} \rightarrow B$ denote the universal families as in notation 2.29. Then there exist relative correspondences $\mathcal{P} \in A^3(\mathcal{X} \times_B \mathcal{Y})$ and $\mathcal{Q} \in A^5(\mathcal{Y} \times_B \mathcal{X})$ with the above property on each fibre: i.e., for each $b \in B$ we have

$$\begin{aligned} (({}^t\mathcal{P} \circ {}^t\mathcal{Q})|_{X_b \times X_b})_* &= (\Pi_2^{X_b})_*: A^2(X_b) \rightarrow A^2(X_b), \\ ((\mathcal{Q} \circ \mathcal{P})|_{X_b \times X_b})_* &= (\Pi_6^{X_b})_*: A^4(X_b) \rightarrow A^4(X_b). \end{aligned}$$

Proof. The correspondence $P \in A^3(X \times Y)$ is defined as the universal family of lines on Y . Letting $I \subset X \times X$ denote the incidence correspondence, we have

$$I = {}^tP \circ P \text{ in } A^2(X \times X)$$

[38, Lemma 17.2].

Proposition 2.20 states that the composition

$$A_{(2)}^2(X) \xrightarrow{\cdot g^2} A_{(2)}^4(X) \xrightarrow{-\frac{1}{6}({}^tP \circ P)_*} A_{(2)}^2(X)$$

is the identity, in other words

$$\left(-\frac{1}{6}{}^tP \circ P \circ \Gamma_{g^2}\right)_* = \text{id}: A_{(2)}^2(X) \rightarrow A_{(2)}^2(X).$$

(Here $\Gamma_{g^2} \in A^6(X \times X)$ can be defined as $\frac{1}{d} {}^t\Gamma_\tau \circ \Gamma_\tau$, where $\tau: R \rightarrow X$ denotes the inclusion of a smooth complete intersection of class dg^2). Defining

$${}^tQ := -\frac{1}{6}P \circ \Gamma_{g^2} \in A^5(X \times Y),$$

we obtain the first equality of corollary 2.21.

For the second equality, it suffices to take the transpose of the first equality, since we know that the CK decomposition of theorem 2.14 is self-dual.

As to the “moreover” part of the corollary: obviously both P and Γ_{g^2} exist relatively, and so the same goes for Q . $\square$

2.7. Spread.

Proposition 2.22 (Voisin [43]). *Let M be a smooth projective variety of dimension $r + 2$, and assume M has trivial Chow groups (i.e. $A_{hom}^*(M) = 0$). Let $L_1, \dots, L_r$ be very ample line bundles on M , and let*

$$\mathcal{Y} \rightarrow B$$

be the universal family of smooth complete intersections

$$Y_b = M \cap D_1 \cap \dots \cap D_r, \quad D_j \in |L_j|.$$

Let $R \in A^2(\mathcal{Y} \times_B \mathcal{Y})$ be a relative correspondence such that

$$R|_{Y_b \times Y_b} = 0 \in H^4(Y_b \times Y_b) \text{ for very general } b \in B.$$

Then there exists $\delta \in A^n(M \times M)$ such that

$$R|_{Y_b \times Y_b} = \delta|_{Y_b \times Y_b} \in A^2(Y_b \times Y_b) \quad \forall b \in B.$$

Proof. This follows from [43] (NB: as noted in loc. cit., the “Voisin standard conjecture” [43, Conjecture 1.6] is satisfied for $n = 2$, so is not needed as extra assumption).

(Alternatively, one could give a quick proof of proposition 2.22 along the lines of [44, Proposition 1.6], at least under the extra assumption that the surfaces Y_b have non-zero primitive cohomology, which is OK in all cases where we apply proposition 2.22 since we only consider $K3$ surfaces.) $\square$

2.8. Families of $K3$ surfaces.

Notation 2.23. Let $g \in \{2, 3, 4, 5, 6, 8\}$. Let

$$\mathbb{P}_g := \begin{cases} \mathbb{P}(1^3, 3) & \text{if } g = 2, \\ \mathbb{P}^g(\mathbb{C}) & \text{if } g = 3, 4, 5, \\ G(1, 4) & \text{if } g = 6, \\ G(1, 5) & \text{if } g = 8 \end{cases}$$

(here $\mathbb{P}(1^3, 3)$ denotes a weighted projective space, and $G(1, m)$ is the Grassmannian of lines in $\mathbb{P}^m(\mathbb{C})$). Consider the vector bundle U_g on $\mathbb{P}_g$ defined as

$$U_g := \begin{cases} \mathcal{O}(6) & \text{if } g = 2, \\ \mathcal{O}(4) & \text{if } g = 3, \\ \mathcal{O}(3) \oplus \mathcal{O}(2) & \text{if } g = 4, \\ \mathcal{O}(2)^{\oplus 3} & \text{if } g = 5, \\ \mathcal{O}(2) \oplus \mathcal{O}(1)^{\oplus 3} & \text{if } g = 6, \\ \mathcal{O}(1)^{\oplus 6} & \text{if } g = 8 \end{cases}$$

(here $\mathcal{O}(i)$ on a Grassmannian refers to the Plücker embedding).

Let $B_g \subset \mathbb{P}H^0(\mathbb{P}_g, U_g)$ denote the Zariski open parametrizing smooth sections, and let

$$\mathcal{S}_g := \{(x, s) \mid s(x) = 0\} \subset \mathbb{P}_g \times B_g$$

denote the universal family.

As shown by Mukai [29], a general $K3$ surface of genus $g \in \{2, 3, 4, 5, 6, 8\}$ is isomorphic to a fibre S_b of the family $\mathcal{S}_g \rightarrow B_g$.

Notation 2.24. Let $\mathcal{S}_g \rightarrow B_g$ be as in notation 2.23. The family

$$\mathcal{S}_g^{[2]} \rightarrow B_g$$

is defined as follows: take $\mathcal{S} \times_B \mathcal{S}$ and blow-up the relative diagonal, then take the quotient for the action of the symmetric group exchanging the two factors. The fibre of $\mathcal{S}_g^{[2]} \rightarrow B_g$ is the Hilbert square $(S_b)^{[2]}$ of the $K3$ surface S_b .

2.9. A Lagrangian fibration.

Proposition 2.25 (Mukai [28]). *Let S be a general K3 surface of genus 5, and let $X = S^{[2]}$ be the Hilbert scheme. There exists a Lagrangian fibration*

$$\phi: X \rightarrow \mathbb{P}^2.$$

Proof. The surface S can be defined as the intersection of three quadrics Q_1, Q_2, Q_3 in $\mathbb{P}^5(\mathbb{C})$. Let $N \cong (\mathbb{P}^2)^\vee$ be the net of quadrics spanned by Q_1, Q_2, Q_3 . Any length 2 subscheme ξ in S determines a line ℓ_ξ in $\mathbb{P}^5$. Quadrics in N containing the line ℓ_ξ form a pencil $P_\xi \cong \mathbb{P}^1$ inside N . Dually, this determines a point in $\mathbb{P}^2$, and so we obtain a morphism

$$\begin{aligned} \phi: X &\rightarrow \mathbb{P}^2, \\ \xi &\mapsto (P_\xi)^\vee. \end{aligned}$$

(This fibration ϕ is also described in [36, Section 2.1] and [9].) □

Remark 2.26. *Hassett–Tschinkel prove more generally that $X = S^{[2]}$ admits a Lagrangian fibration, for S a generic K3 surface of degree $2m^2$ with $m \geq 2$ [18, Proposition 7.1]. Generalizations to certain higher-dimensional Hilbert schemes $S^{[r]}$, $r > 2$ are given in [36] and [20].*

2.10. A relation of motives. Galkin–Shinder [15] have constructed a relation between a cubic and its Fano variety of lines in the Grothendieck ring of varieties. The following is a version of this relation on the level of Chow motives:

Theorem 2.27 ([24]). *Let $Y \subset \mathbb{P}^{n+1}(\mathbb{C})$ be a smooth cubic, and let $X = F(Y)$ be the Fano variety of lines in Y . There is an isomorphism of motives*

$$\Gamma: h(X)(2) \oplus \bigoplus_{i=0}^n h(Y)(i) \cong h(Y^{[2]}) \quad \text{in } \mathcal{M}_{\text{rat}}.$$

Proof. This is [24, Theorem 5]. □

In particular, this implies the following:

Corollary 2.28. *Let $Y \subset \mathbb{P}^5(\mathbb{C})$ be a smooth cubic fourfold, and let $X = F(Y)$ be the Fano variety of lines in Y . There exists a correspondence inducing an injection*

$$A_{\text{hom}}^i(X) \hookrightarrow A_{\text{hom}}^{i+2}(Y \times Y) \oplus \bigoplus A_{\text{hom}}^3(Y).$$

Moreover, this can be done relatively: let $\mathcal{Y} \rightarrow B$ be a family of smooth cubic fourfolds, and let $\mathcal{X} \rightarrow B$ be the family of associated Fano varieties. Then there exists a relative correspondence $\Psi_1 \in A^6(\mathcal{X} \times_B (\mathcal{Y} \times_B \mathcal{Y})) \oplus \bigoplus A^(\mathcal{X} \times_B \mathcal{Y})$, inducing an injection*

$$(\Psi_1|_b)_*: A_{\text{hom}}^i(X_b) \hookrightarrow A_{\text{hom}}^{i+2}(Y_b \times Y_b) \oplus \bigoplus A_{\text{hom}}^3(Y_b) \quad \forall b \in B.$$

(Here $\Psi_1|_b$ is shorthand for the restriction of Ψ_1 to the fibre $X_b \times (Y_b)^2 \cup \bigcup (X_b \times Y_b)$.)

Proof. The first statement follows immediately from theorem 2.27, for it is readily seen that there is a correspondence-induced injection

$$(4) \quad A^{i+2}(Y^{[2]}) \hookrightarrow A^{i+2}(Y \times Y) \oplus \bigoplus A^*(Y) .$$

Restricting to $A_{hom}^*(\cdot)$ (and remembering that $A_{hom}^i(Y) = 0$ for $i \neq 3$) gives the first statement.

To prove the “moreover” part, we need to understand how the correspondence Γ of theorem 2.27 is constructed. The construction is based on the existence of a birational map (found by Galkin–Shinder [15, Proof of Theorem 5.1])

$$\phi: Y^{[2]} \dashrightarrow W ,$$

where $W \rightarrow Y$ denotes the $\mathbb{P}^4$ -bundle whose fibre over $y \in Y$ is the set of lines in $\mathbb{P}^5$ passing through y . The indeterminacy locus $\tau: Z \hookrightarrow Y^{[2]}$ of ϕ is a $\mathbb{P}^2$ -bundle $p: Z \rightarrow X$, and it turns out [48, Proposition 2.9] that the indeterminacy of ϕ is resolved by the blow-up with center Z . The map from $h(X)(2)$ to $h(Y^{[2]})$ in theorem 3.1 is defined simply as

$$\Gamma_\tau \circ {}^t\Gamma_p: h(X)(2) \rightarrow h(Y^{[2]}) \text{ in } \mathcal{M}_{\text{rat}} .$$

This construction naturally extends to the family: inside $\mathcal{Y}^{[2]}$ there is a codimension 2 locus $\tau: \mathcal{Z} \hookrightarrow \mathcal{Y}^{[2]}$ (defined as those points $x \in (Y_b)^{[2]}$ for which the line L_x determined by x is contained in Y_b), and $\mathcal{Z}$ has the structure of a $\mathbb{P}^2$ -bundle $p: \mathcal{Z} \rightarrow \mathcal{X}$. One can thus define

$$\Psi := \Gamma_\tau \circ {}^t\Gamma_p \in A^6(\mathcal{X} \times_B \mathcal{Y}^{[2]}) .$$

The desired relative correspondence Ψ_1 is defined as the composition $\Psi_1 = \Psi_0 \circ \Psi$, where the correspondence

$$\Psi_0 \in A^4(\mathcal{Y}^{[2]} \times_B (\mathcal{Y} \times_B \mathcal{Y})) \oplus \bigoplus A^*(\mathcal{Y}^{[2]} \times_B \mathcal{Y})$$

induces the fibrewise injection (4); this Ψ_0 obviously exists relatively. $\square$

2.11. Cubics containing two planes.

Notation 2.29. Let $P_1, P_2 \subset \mathbb{P}^5(\mathbb{C})$ be two fixed disjoint planes, and consider smooth cubic fourfolds Y_b containing these two planes. Such cubics form a family

$$\mathcal{Y} \rightarrow B ,$$

where $B \subset \bar{B} \subset \mathbb{P}H^0(\mathbb{P}^5, \mathcal{O}_{\mathbb{P}^5}(3))$. Here B is Zariski open in $\bar{B}$, and $\bar{B}$ is the closed subset parametrizing equations of the form

$$f_1(u, v, w; x, y, z) + f_2(u, v, w; x, y, z) = 0 ,$$

where f_1, f_2 are bihomogeneous of bidegree $(1, 2)$ resp. $(2, 1)$ in the variables $[u, v, w]$ and $[x, y, z]$.

Let

$$\mathcal{X} \rightarrow B$$

be the universal family of Fano varieties of lines in cubic fourfolds containing two planes, i.e.

$$\mathcal{X} := \{(\ell, b) \mid \ell \subset Y_b\} \subset \text{Gr}(1, 5) \times B$$

(here $Gr(1, 5)$ is the Grassmannian of lines in $\mathbb{P}^5$). The fibre X_b (of $\mathcal{X} \rightarrow B$ over $b \in B$) is the Fano variety of lines in the cubic Y_b .

Let

$$\mathcal{S} \rightarrow B$$

be the universal family of K3 surfaces in $\mathbb{P}^2 \times \mathbb{P}^2$ defined by the bihomogeneous equations

$$f_1(u, v, w; x, y, z) = f_2(u, v, w; x, y, z) = 0.$$

For later use, we record two lemmas that express a relation between the cubic Y_b and the K3 surface S_b :

Lemma 2.30. *Notation as in notation 2.29. There exist relative correspondences*

$$\Psi_2 \in A^3(\mathcal{Y} \times_B \mathcal{S}), \quad \Xi_2 \in A^3(\mathcal{S} \times_B \mathcal{Y}),$$

with the property that

$$((\Xi_2 \circ \Psi_2)|_{Y_b \times Y_b})_* = \text{id}: A_{\text{hom}}^3(Y_b) \rightarrow A_{\text{hom}}^3(Y_b) \quad \forall b \in B.$$

Proof. The proof of the rationality of the cubics Y_b (cf. for instance [16, Section 5], [17, Section 1.2]) can be done by showing that for any $b \in B$ there exists a birational map

$$\rho_b: \mathbb{P}^2 \times \mathbb{P}^2 \dashrightarrow Y_b.$$

Taking coordinates $[u, v, w], [x, y, z]$ for $\mathbb{P}^2 \times \mathbb{P}^2$, the indeterminacy locus of ρ_b is exactly the K3 surface S_b , and the blow-up Z_b of $\mathbb{P}^2 \times \mathbb{P}^2$ along S_b admits a morphism $\tilde{\rho}_b: Z_b \rightarrow Y_b$ resolving the indeterminacy of ρ_b . Let $i_b: E_b \hookrightarrow Z_b$ denote the exceptional divisor of the blow-up, and let $p_b: E_b \rightarrow S_b$ denote the induced morphism. The usual blow-up exact sequence implies that the composition

$$A_{\text{hom}}^3(Y_b) \xrightarrow{(\tilde{\rho}_b)^*} A_{\text{hom}}^3(Z_b) \xrightarrow{(i_b)^*} A_{\text{hom}}^3(E_b) \xrightarrow{(p_b)_*} A_{\text{hom}}^2(S_b)$$

is injective, and a left-inverse is given by

$$A_{\text{hom}}^2(S_b) \xrightarrow{(p_b)^*} A_{\text{hom}}^2(E_b) \xrightarrow{(i_b)_*} A_{\text{hom}}^3(Z_b) \xrightarrow{(\tilde{\rho}_b)_*} A_{\text{hom}}^3(Y_b).$$

This defines fibrewise correspondences between Y_b and S_b . By looking at the blow-up of $\mathbb{P}^2 \times \mathbb{P}^2 \times B$ along $\mathcal{S}$, one sees that these correspondences exist relatively, as desired. $\square$

Lemma 2.31. *Notation as in notation 2.29. There exists a relative correspondence*

$$\Psi_3 \in A^6((\mathcal{Y} \times_B \mathcal{Y}) \times_B (\mathcal{S} \times_B \mathcal{S})) \oplus A^*(\mathcal{Y} \times_B \mathcal{Y}) \times_B \mathcal{S}^{\oplus 2},$$

with the property that

$$(\Psi_3|_b)_*: A_{\text{hom}}^4(Y_b \times Y_b) \rightarrow A_{\text{hom}}^2(S_b \times S_b) \oplus A_{\text{hom}}^2(S_b)^{\oplus 2}$$

is injective for very general $b \in B$.

Proof. Lemma 2.30 implies that the map of Chow motives

$$(\Psi_2, \Psi'): h(Y_b) \rightarrow h(S_b)(1) \oplus \bigoplus_{i=0}^4 \mathbb{L}(i) \quad \text{in } \mathcal{M}_{\text{rat}}$$

admits a left-inverse (i.e. the left-hand side is a direct summand of the right-hand side). This implies that there is an induced map

$$h(Y_b \times Y_b) \rightarrow h(S_b \times S_b)(2) \oplus \bigoplus_j h(S_b)(j) \oplus \bigoplus_k \mathbb{L}(k) \quad \text{in } \mathcal{M}_{\text{rat}}$$

admitting a left-inverse. In particular, taking $A_{\text{hom}}^4()$ on both sides, we find there is an injection

$$A_{\text{hom}}^4(Y_b \times Y_b) \hookrightarrow A_{\text{hom}}^2(S_b \times S_b) \oplus \bigoplus A_{\text{hom}}^2(S_b)$$

which is induced by various combinations of Ψ_2 and Ψ' . We have seen (lemma 2.30) that Ψ_2 exists relatively. As for Ψ' , this is obtained by considering the “trivial” cycles $h^i \in A^i(Y_b)$ coming from a hyperplane section $h \in A^1(\mathbb{P}^5)$, plus choosing a basis for $N^2 H^4(Y_b)$. The cycles h^i obviously exist relatively. Let $B_0 \subset B$ be the locus where $\dim N^2 H^4(Y_b)$ equals 3; this B_0 is the intersection of a countable number of Zariski opens in B . The correspondence Ψ' exists as a relative correspondence over B_0 . It follows there exists a relative correspondence Ψ_3 as indicated. $\square$

Remark 2.32. Let Y_b be a smooth cubic fourfold containing two planes. We will say that S_b is the related K3 surface. (We want to avoid the terminology “associated K3 surface”, which already has a different meaning [16], [17].)

2.12. The generalized Franchetta conjecture for certain K3 surfaces.

Proposition 2.33. Let $\mathcal{S} \rightarrow B$ be the family of K3 surfaces of notation 2.29. Let $\Gamma \in A^2(\mathcal{S})$ be a relative cycle such that

$$\Gamma|_{S_b} = 0 \quad \text{in } H^4(S_b) \quad \forall b \in B.$$

Then also

$$\Gamma|_{S_b} = 0 \quad \text{in } A^2(S_b) \quad \forall b \in B.$$

Proof. Let $\mathbb{P} := \mathbb{P}^2 \times \mathbb{P}^2$, and let $U \rightarrow \mathbb{P}$ be the rank 2 vector bundle

$$U := \mathcal{O}_{\mathbb{P}}(1, 2) \oplus \mathcal{O}_{\mathbb{P}}(2, 1).$$

Let

$$\bar{B} := \mathbb{P}H^0(\mathbb{P}, U), \quad \bar{\mathcal{S}} := \{(x, f) \mid f(x) = 0\} \subset \mathbb{P} \times \bar{B}.$$

Then B and $\mathcal{S}$ are Zariski opens in $\bar{B}$ resp. $\bar{\mathcal{S}}$. There is a diagram

$$\begin{array}{ccc} \bar{\mathcal{S}} & \xrightarrow{\nu} & \mathbb{P} \\ \downarrow \pi & & \\ \bar{B} & & \end{array}$$

Let $\phi_b: S_b \hookrightarrow \mathcal{S} \hookrightarrow \bar{\mathcal{S}}$ and $i_b: S_b \hookrightarrow \mathbb{P}$ denote the inclusion morphisms. We claim that there is equality

$$(5) \quad \text{Im}(A^2(\bar{\mathcal{S}}) \xrightarrow{(\phi_b)^*} A^2(S_b)) = \text{Im}(A^2(\mathbb{P}) \xrightarrow{(i_b)^*} A^2(S_b)).$$

This claim implies the proposition: indeed,

$$A^2(\mathbb{P}) = A^2(\mathbb{P}^2 \times \mathbb{P}^2) = \bigoplus_{k+\ell=2} A^k(\mathbb{P}^2) \otimes A^\ell(\mathbb{P}^2)$$

is a 3-dimensional $\mathbb{Q}$ -vector space with generators $h^2 \times \mathbb{P}^2$, $h \times h$, $\mathbb{P}^2 \times h^2$ (where $h \in A^1(\mathbb{P}^2)$ is the class of a hyperplane section). Thus every element in $A^2(\mathbb{P})$ is a sum of intersections of divisors. Since intersections of two divisors are a multiple of the distinguished class $\mathfrak{o}_{S_b}$ in $A^2(S_b)$ [4], the claim (applied to any lifting $\bar{\Gamma}$ of Γ to $A^2(\bar{\mathcal{S}})$) implies that

$$\Gamma|_{S_b} = (\phi_b)^*(\bar{\Gamma}) = m_b \mathfrak{o}_{S_b} \text{ in } A^2(S_b)$$

for some $m_b \in \mathbb{Q}$, and the proposition is proven.

To prove the claim, we apply the argument of [33, Lemma 2.1] (where (5) is proven in a very similar set-up). First, one observes that $U \rightarrow \mathbb{P}$ is globally generated and so $\nu: \bar{\mathcal{S}} \rightarrow \mathbb{P}$ is a projective bundle; the fibre over a point $x \in \mathbb{P}$ is

$$\mathbb{P}H^0(\mathbb{P}, U \otimes I_x)$$

(where I_x denotes the ideal sheaf of x). The projective bundle formula gives an equality

$$(6) \quad A^r(\bar{\mathcal{S}}) = \xi^r \cdot \nu^* A^0(\mathbb{P}) + \xi^{r-1} \cdot \nu^* A^1(\mathbb{P}) + \cdots + \nu^* A^r(\mathbb{P}),$$

where $\xi \in A^1(\bar{\mathcal{S}})$ denotes the relative hyperplane class. In particular, if $H \in A^1(\bar{B})$ denotes the hyperplane class, we can write

$$\pi^*(H) = a \cdot \xi + \nu^*(b) \text{ in } A^1(\bar{\mathcal{S}})$$

for some $a \in \mathbb{Q}$ and $b \in A^1(\mathbb{P})$. The constant a must be non-zero, for otherwise

$$\pi^*(H^{\dim \bar{B}}) = \nu^*(b^{\dim \bar{B}}) \text{ in } A^{\dim \bar{B}}(\bar{\mathcal{S}}),$$

which is absurd (the right-hand side is 0 since $\dim \bar{B} > 4 = \dim \mathbb{P}$, but the left-hand side is the pullback of an effective 0-cycle and is non-zero).

It follows that one may write

$$\xi = \frac{1}{a}(\pi^*(H) - \nu^*(b)) \text{ in } A^1(\bar{\mathcal{S}}).$$

Plugging this in equality (6) and taking $r = 2$, we find that

$$A^2(\bar{\mathcal{S}}) = (\pi^*(H) - \nu^*(b))^2 \cdot \nu^* A^0(\mathbb{P}) \oplus (\pi^*(H) - \nu^*(b)) \cdot \nu^* A^1(\mathbb{P}) \oplus \nu^* A^2(\mathbb{P}).$$

The class $\pi^*(H)$ vanishes when restricting to S_b , i.e. $(\phi_b)^* \pi^*(H) = 0$ in $A^2(S_b)$ for any $b \in B$, and so

$$\begin{aligned} \text{Im}(A^2(\bar{\mathcal{S}}) \xrightarrow{(\phi_b)^*} A^2(S_b)) &= \text{Im}\left((\nu^*(b^2) \cdot \nu^* A^0(\mathbb{P}) \oplus \nu^*(b) \cdot \nu^* A^1(\mathbb{P}) \oplus \nu^* A^2(\mathbb{P})) \xrightarrow{(\phi_b)^*} A^2(S_b)\right) \\ &= \text{Im}(\nu^* A^2(\mathbb{P}) \xrightarrow{(\phi_b)^*} A^2(S_b)) \\ &= \text{Im}(A^2(\mathbb{P}) \xrightarrow{(i_b)^*} A^2(S_b)), \end{aligned}$$

proving the claim. □

Remark 2.34. For any $b \in B$, let $\mathfrak{o}_{S_b} \in A^2(S_b)$ denote the distinguished 0-cycle of [4]. An equivalent way of stating proposition 2.33 is as follows: given any relative cycle $\Gamma \in A^2(\mathcal{S})$, the restriction

$$\Gamma|_{S_b} \in A^2(S_b)$$

is a multiple of $\mathfrak{o}_{S_b}$ in $A^2(S_b)$.

This statement is very similar to the “generalized Franchetta conjecture” for K3 surfaces, which was formulated by O’Grady [32, Section 5] and proven in certain cases by Pavic–Shen–Yin [33].

3. MAIN RESULTS

This section proves the two main results of this note, theorems 3.1 and 3.2. The proofs are based on the method of “spread” of cycles in nice families, as developed by Voisin [43], [44], [45], [46].

The first main result is a “generalized Franchetta conjecture” type of statement for certain Hilbert squares:

Theorem 3.1. *Let $g \in \{2, 3, 4, 5, 6, 8\}$. Let $\mathcal{X} \rightarrow B$ denote the universal family of Hilbert squares of genus g K3 surfaces (notation 2.24). Let $\Gamma \subset \mathcal{X}$ be a codimension 2 subvariety. Then the restriction $\Gamma|_{X_b} \in A^2(X_b)$ is in $A_{(0)}^2(X_b)$ for all $b \in B$.*

Proof. Let us fix an integer $g \in \{2, 3, 4, 5, 6, 8\}$, and write

$$\mathcal{S} \rightarrow B, \quad \mathcal{X} := \mathcal{S}^{[2]} \rightarrow B$$

for the universal family of genus g K3 surfaces (i.e., the family $\mathcal{S}_g \rightarrow B_g$ of subsection 2.8), resp. the universal Hilbert square of a genus g K3 surface (as in notation 2.24).

One knows (remark 2.11) that the MCK decomposition $\{\Pi_j^{X_b}\}$ for the X_b of [38] exists relatively, in particular there exists a relative correspondence $\Pi_2^{\mathcal{X}} \in A^4(\mathcal{X} \times_B \mathcal{X})$ such that

$$\Pi_2^{\mathcal{X}}|_{X_b \times X_b} = \Pi_2^{X_b} \quad \text{in } A^4(X_b \times X_b) \quad \forall b \in B.$$

Let us now consider the relative cycle

$$\Gamma_0 := (\Pi_2^{\mathcal{X}})_*(\Gamma) \in A^2(\mathcal{X}).$$

We know this is fibrewise homologically trivial:

$$(7) \quad \Gamma_0|_{X_b} = 0 \quad \text{in } H^4(X_b) \quad \forall b \in B.$$

(This is simply because $\Pi_2^{X_b} \circ \Pi_4^{X_b} = 0$ in $H^8(X_b \times X_b)$.)

What we desire to prove (in order to establish the theorem) is that there is a fibrewise rational equivalence

$$(8) \quad \Gamma_0|_{X_b} \stackrel{??}{=} 0 \quad \text{in } A^2(X_b) \quad \forall b \in B.$$

It is easier to move things to the family $\mathcal{S} \times_B \mathcal{S}$. To this end, let

$$\Psi \in A^4(\mathcal{X} \times_B (\mathcal{S} \times_B \mathcal{S}))$$

be the relative correspondence defined by the standard diagram

$$\begin{array}{ccc} \mathcal{X} := \mathcal{S}^{[2]} & \xleftarrow{\quad} & \widetilde{\mathcal{S} \times_B \mathcal{S}} \\ \downarrow & & \downarrow \\ \mathcal{S}^{(2)} := (\mathcal{S} \times_B \mathcal{S})/\mathfrak{S}_2 & \xleftarrow{\quad} & \mathcal{S} \times_B \mathcal{S} \end{array}$$

Let us consider the relative correspondence

$$\Gamma_1 := (\Psi)_*(\Gamma_0) \in A^2(\mathcal{S} \times_B \mathcal{S}).$$

We recall from equality (7) that $\Gamma_0|_{X_b}$ is in $A_{hom}^2(X_b) = A_{AJ}^2(X_b)$ for all $b \in B$. The restriction $\Psi_b := \Psi|_{X_b \times (S_b)^2}$ acts as

$$(\Psi_b)_*: A_{AJ}^2(X_b) \rightarrow A_{AJ}^2(\widetilde{S_b \times S_b}) \rightarrow A_{AJ}^2(S_b \times S_b),$$

where the first arrow is an injection and the second arrow is an isomorphism. Hence, in order to prove (8) it will suffice to prove

$$(9) \quad \Gamma_1|_{X_b} \stackrel{??}{=} 0 \quad \text{in } A^2(S_b \times S_b) \quad \forall b \in B.$$

But the family $\mathcal{S} \rightarrow B$ enters into the set-up of proposition 2.22 (with $M = \mathbb{P}_g$ and L_j as given in subsection 2.8), and (7) implies that the cycle $\Gamma_1 \in A^2(\mathcal{S} \times_B \mathcal{S})$ is fibrewise homologically trivial:

$$\Gamma_1|_{S_b \times S_b} = 0 \quad \text{in } H^4(S_b \times S_b) \quad \forall b \in B.$$

Proposition 2.22 allows us to conclude that there exists $\delta \in A^2(\mathbb{P}_g \times \mathbb{P}_g)$ such that there is fibrewise rational equivalence

$$(10) \quad \Gamma_1|_{S_b \times S_b} = \delta|_{S_b \times S_b} \quad \text{in } A^2(S_b \times S_b) \quad \forall b \in B.$$

Let us decompose

$$\delta = \delta_0 + \delta_1 + \delta_2 \in A^2(\mathbb{P}_g) \otimes A^0(\mathbb{P}_g) \oplus A^1(\mathbb{P}_g) \otimes A^1(\mathbb{P}_g) \oplus A^0(\mathbb{P}_g) \otimes A^2(\mathbb{P}_g),$$

where it is understood that δ_0 is in the first summand, and δ_1, δ_2 are in the second resp. last summand. (This decomposition is possible because $\mathbb{P}_g$ has trivial Chow groups, and so

$$A^*(\mathbb{P}_g \times \mathbb{P}_g) = A^*(\mathbb{P}_g) \otimes A^*(\mathbb{P}_g).)$$

As the restriction $\delta|_{S_b \times S_b}$ is homologically trivial (for each $b \in B$), and the $\delta_j|_{S_b \times S_b}$ land in different summands of the Künneth decomposition of cohomology, we have

$$(11) \quad \delta_j|_{S_b \times S_b} \in A_{hom}^2(S_b \times S_b) \quad \forall b \in B, \quad j = 0, 1, 2.$$

From this, we may conclude that

$$\delta_1|_{S_b \times S_b} = 0 \quad \text{in } A^2(S_b \times S_b) \quad \forall b \in B,$$

thanks to Voisin's proof of the Beauville–Voisin conjecture for $(S_b)^m$ with m small [42, Proposition 2.2].

As for the component δ_0 , let us take a relatively ample divisor $h \in A^1(\mathcal{S})$ (e.g., induced by an ample divisor on $\mathbb{P}_g$), and consider the relative cycle

$$\delta_{00} := (p_1)_*((\delta_0 \times B)|_{\mathcal{S} \times_B \mathcal{S}}) \cdot (p_2)^*(h^2) \in A^2(\mathcal{S}),$$

where $p_i: \mathcal{S} \times_B \mathcal{S} \rightarrow \mathcal{S}$ for $i = 1, 2$ denotes projection on the first resp. second factor. We have that

$$\delta_0|_{S_b \times S_b} = d \cdot (p_1)^*(\delta_{00}|_{S_b}) \quad \text{in } A^2(S_b \times S_b) \quad \forall b \in B,$$

for some non-zero integer d . We observe that (11) implies that the restriction $\delta_{00}|_{S_b}$ is of degree 0 for all $b \in B$. But then, reasoning exactly as in [33], we must have that $\delta_{00}|_{S_b}$ is rationally trivial. (More in detail: by construction, one has

$$\delta_{00}|_{S_b} \in \text{Im}(A^2(\mathbb{P}_g) \rightarrow A^2(S_b)) .$$

However, it is known that

$$\text{Im}(A^2(\mathbb{P}_g) \rightarrow A^2(S_b)) = \mathbb{Q}[\mathfrak{o}_{S_b}]$$

(where $\mathfrak{o}_{S_b}$ is the distinguished point of [4]); Pavic–Shen–Yin supply not one but two proofs for this [33, Proposition 3.1].)

It follows that also

$$\delta_0|_{S_b \times S_b} = 0 \quad \text{in } A^2(S_b \times S_b) \quad \forall b \in B .$$

The proof for δ_2 is the same as for δ_0 . We conclude that

$$\delta|_{S_b \times S_b} = 0 \quad \text{in } A^2(S_b \times S_b) \quad \forall b \in B ,$$

and so (10) gives the desired equality (9). $\square$

We now come to the second main result of this note:

Theorem 3.2. *Let $g \in \{2, 3, 4, 5, 6, 8\}$. Let $\mathcal{X} \rightarrow B$ denote the universal family of Hilbert squares of genus g K3 surfaces (notation 2.24). Let $\Gamma \subset \mathcal{X}$ be a codimension 2 subvariety. Assume the restriction*

$$\Gamma_b := \Gamma|_{X_b} \in A^2(X_b)$$

is Lagrangian (i.e., $\cup \Gamma_b: H^{2,0}(X_b) \rightarrow H^{4,2}(X_b)$ is the zero map), for very general $b \in B$. Then

$$\cdot \Gamma_b: A_{hom}^2(X_b) \rightarrow A^4(X_b)$$

is the zero map, for all $b \in B$.

Proof. Thanks to theorem 2.1, there is a splitting

$$A_{hom}^2(X_b) = A_{(2)}^2(X_b) \oplus (A_{(0)}^2(X_b) \cap A_{hom}^2(X_b)) ,$$

where the second summand is conjecturally zero. It is known that the intersection product map

$$A_{(0)}^2(X_b) \otimes (A_{(0)}^2(X_b) \cap A_{hom}^2(X_b)) \rightarrow A^4(X_b)$$

is the zero map [38, Theorem 3]. Since $\Gamma_b \in A_{(0)}^2(X_b)$ by theorem 3.1, we are reduced to proving that

$$\cdot \Gamma_b: A_{(2)}^2(X_b) \rightarrow A^4(X_b)$$

is the zero map.

Let us consider a resolution of singularities $f: \tilde{\Gamma} \rightarrow \Gamma$, and let $\tau: \tilde{\Gamma} \hookrightarrow \mathcal{X}$ denote the composition of f with the inclusion morphism $\Gamma \hookrightarrow \mathcal{X}$. Let us consider the relative correspondence

$$\Gamma_0 := \mathcal{L} \circ \Gamma_\tau \circ {}^t\Gamma_\tau \circ \Pi_2^\mathcal{X} \in A^4(\mathcal{X} \times_B \mathcal{X}) ,$$

where $\Pi_2^\mathcal{X}$ and $\mathcal{L}$ are as in remark 2.11 resp. lemma 2.18.

By construction, for any $b \in B$, the restriction $\Gamma_0|_{X_b \times X_b}$ acts on Chow groups as

$$A^i(X_b) \rightarrow (\Pi_2^{X_b})_* A^i(X_b) \xrightarrow{\cdot \Gamma_b} A^{i+2}(X_b) \xrightarrow{(L_b)_*} A^i(X_b) .$$

In particular, taking $i = 2$ and applying the fact that $\Gamma_b \in A_{(0)}^2(X_b)$ (theorem 3.1), we get an action

$$(\Gamma_0|_{X_b \times X_b})_*: A_{(2)}^2(X_b) \xrightarrow{\text{id}} A_{(2)}^2(X_b) \xrightarrow{\Gamma_b} A_{(2)}^4(X_b) \xrightarrow{(L_b)_*} A_{(2)}^2(X_b).$$

We know that the last arrow is an isomorphism (theorem 2.17). Also, $\Pi_2^{X_b}$ acts as the identity on $A_{(2)}^2(X_b)$. Thus, to prove theorem 3.2 we are reduced to proving that

$$(12) \quad ((\Pi_2^{\mathcal{X}} \circ \Gamma_0)|_{X_b \times X_b})_* \stackrel{??}{=} 0: A_{(2)}^2(X_b) \rightarrow A_{(2)}^2(X_b) \quad \forall b \in B.$$

The input we have at our disposition is that we know (from the coisotropic assumption) that

$$(13) \quad (\Gamma_0|_{X_b \times X_b})_* = 0: H^{2,0}(X_b) \rightarrow H^{2,0}(X_b) \quad \text{for very general } b \in B.$$

We observe that (13), combined with the Lefschetz (1,1) theorem, implies the following: for very general $b \in B$, there exist a curve $C_b \subset X_b$, a divisor D_b and a cycle γ_b supported on $C_b \times D_b \subset X_b \times X_b$, such that

$$\Gamma_0|_{X_b \times X_b} - \gamma_b = 0 \quad \text{in } H^8(X_b \times X_b).$$

Thanks to Voisin's key result [43, Proposition 3.7] (cf. also [44, Proposition 4.25]), it is possible to spread out these data. That is, there exist subvarieties $\mathcal{C} \subset \mathcal{X}$, $\mathcal{D} \subset \mathcal{X}$ of codimension 3 resp. 1, and a cycle $\gamma \in A^4(\mathcal{X} \times_B \mathcal{X})$ supported on $\mathcal{C} \times_B \mathcal{D}$ that does the job of the various γ_b , i.e. such that

$$(\Gamma_0 - \gamma)|_{X_b \times X_b} = 0 \quad \text{in } H^8(X_b \times X_b) \quad \text{for very general } b \in B.$$

In other words, the relative correspondence defined as

$$\Gamma_1 := \Gamma_0 - \gamma \in A^4(\mathcal{X} \times_B \mathcal{X})$$

has the property that

$$(14) \quad \Gamma_1|_{X_b \times X_b} = 0 \quad \text{in } H^8(X_b \times X_b) \quad \text{for very general } b \in B.$$

It is more convenient to move to correspondences living in $A^2(\mathcal{S} \times_B \mathcal{S})$, where $\mathcal{S} \rightarrow B$ is (as before) the family of K3 surfaces of genus g . That is, first we consider the relative correspondence

$$\Gamma_2 := \Psi \circ \Gamma_1 \circ {}^t\Psi \in A^4(\mathcal{S}^{2/B} \times_B \mathcal{S}^{2/B}),$$

where $\Psi \in A^4(\mathcal{X} \times_B \mathcal{S}^{2/B})$ is as in the proof of theorem 3.1. Next, we define relative correspondences

$$\Gamma_3^{i,j} := \Xi_i \circ \Gamma_2 \circ \Theta_j \in A^2(\mathcal{S} \times_B \mathcal{S}) \quad (1 \leq i, j \leq 2),$$

where Ξ_i, Θ_j are as in proposition 2.13. The relative correspondence Γ_1 being fibrewise homologically trivial (equation (14)), the same holds for Γ_2 and $\Gamma_3^{i,j}$:

$$\Gamma_3^{i,j}|_{S_b \times S_b} = 0 \quad \text{in } H^4(S_b \times S_b) \quad \text{for very general } b \in B \quad (1 \leq i, j \leq 2).$$

We can now apply proposition 2.22 to the $\Gamma_3^{i,j}$ (with $M = \mathbb{P}_g$ and L_r as given in subsection 2.8). The conclusion is that there is a fibrewise rational equivalence

$$\Gamma_3^{i,j}|_{S_b \times S_b} = 0 \quad \text{in } A^2(S_b \times S_b) \quad \forall b \in B \quad (1 \leq i, j \leq 2).$$

In particular, this implies that

$$\Pi_2^{\mathcal{S}^{2/B}} \circ \Gamma_2 \circ \Pi_2^{\mathcal{S}^{2/B}} = \sum_{i,j \in \{1,2\}} \Theta_i \circ \Gamma_3^{i,j} \circ \Xi_j \in A^4(\mathcal{S}^{2/B} \times_B \mathcal{S}^{2/B})$$

is fibrewise rationally trivial (the equality follows from proposition 2.13). Composing some more, we find that also

$${}^t\Psi \circ \Pi_2^{\mathcal{S}^{2/B}} \circ \Gamma_2 \circ \Pi_2^{\mathcal{S}^{2/B}} \circ \Psi \in A^4(\mathcal{X} \times_B \mathcal{X})$$

is fibrewise rationally trivial. In view of the definition of Γ_2 (plus the fact that $(\Psi_b)^*(\Psi_b)_*$ is a multiple of the identity on $A_{hom}^2(X_b)$), this implies that

$$(\Pi_2^{\mathcal{X}} \circ \Gamma_1 \circ \Pi_2^{\mathcal{X}})|_{X_b \times X_b} = 0: A_{hom}^2(X_b) \rightarrow A^2(X_b) \quad \forall b \in B.$$

As $A_{(2)}^2(X_b) \subset A_{hom}^2(X_b)$, this implies in particular that

$$A_{(2)}^2(X_b) \xrightarrow{\text{id}} A_{(2)}^2(X_b) \xrightarrow{(\Gamma_1|_{X_b \times X_b})_*} A^2(X_b) \xrightarrow{(\Pi_2^{X_b})_*} A_{(2)}^2(X_b)$$

is the zero map, for all $b \in B$.

For general $b \in B$, the restriction of the cycle δ to the fibre $X_b \times X_b$ will be supported on (curve) $\times$ (divisor), and so will act trivially on $A^2(X_b)$ for dimension reasons. That is, for general $b \in B$ we have equality

$$(\Gamma_1|_{X_b \times X_b})_* = (\Gamma_0|_{X_b \times X_b})_*: A^2(X_b) \rightarrow A^2(X_b).$$

The above thus implies that

$$A_{(2)}^2(X_b) \xrightarrow{(\Gamma_0|_{X_b \times X_b})_*} A^2(X_b) \xrightarrow{(\Pi_2^{X_b})_*} A_{(2)}^2(X_b)$$

is the zero map, for general $b \in B$. That is, we have proven the desired statement (12) for general $b \in B$.

To extend the statement to *all* $b \in B$, one notes that the construction of [43, Proposition 3.7] (which was used above to globalize the various γ_b) can be done locally around a given $b_0 \in B$. $\square$

Remark 3.3. Huybrechts ventures the following guess: “Any distinguished curve, i.e. a curve that is naturally defined in all generic K3 surfaces of fixed degree, should be a constant cycle curve” [19, Section 8.3 Footnote 5].

Similarly, in the set-up of theorem 3.1, could it be the case that for any codimension 2 subvariety $\Gamma \subset \mathcal{X}$ flat over B , the restriction $\Gamma|_{X_b}$ is a constant cycle surface provided it is irreducible? This would somehow “explain” that the restriction $\Gamma|_{X_b}$ is in $A_{(0)}^2(X_b)$, in view of conjecture 1.2.

Remark 3.4. Here is a slightly different way of proving theorem 3.2: Let X_b be the Hilbert square of a very general K3 surface S_b of genus $g \in \{2, 3, 4, 5, 6, 8\}$. Let $\Gamma_b \subset X_b$ be a subvariety of codimension 2 that is Lagrangian. It follows from a result of Voisin [47, Proposition 4.2] that there is a homological equivalence

$$(15) \quad \Gamma_b = \sum_i \lambda_i C_b^i \text{ in } H^4(X_b),$$

where $\lambda_i \in \mathbb{Q}$ and $C_b^i \subset X_b$ is a constant cycle surface.

Suppose we know in addition that $\Gamma_b \in A_{(0)}^2(X_b)$ is the restriction of a relative cycle $\Gamma \in A^2(\mathcal{X})$ as in theorem 3.2). Then, one may “spread out” the fibrewise surfaces C_b^i and use the argument of theorem 3.1 (or alternatively, the more recent [12, Theorems 1.4 and 1.5]) to conclude that there is a rational equivalence

$$(16) \quad \Gamma_b = \sum_i \lambda_i C_i \quad \text{in } A^2(X_b) .$$

This clearly implies theorem 3.2.

4. SOME COROLLARIES

Corollary 4.1. *Let S be a general K3 surface of genus 5, and let $X = S^{[2]}$ be the Hilbert square. Let $A \subset X$ be a general fibre of the Lagrangian fibration $\phi: X \rightarrow \mathbb{P}^2$ of proposition 2.25. Let $b \in A^4(X)$ be a 0-cycle of the form*

$$b = A \cdot a + p \quad \in A^4(X) ,$$

where $a \in A^2(X)$ and p is a sum of intersections of divisors and Chern classes of X . Then b is rationally trivial if and only if b is of degree 0.

Proof. The first thing to remark is that the Lagrangian surface A is in $A_{(0)}^2(X)$. This follows from theorem 3.1, but it is far easier to note that a point $p \in \mathbb{P}^2$ is an intersection of two divisors, and so $A = \phi^{-1}(p)$ is an intersection of two divisors.

Next, one observes that the rational equivalence class of the surface $A \subset X$ exists relatively (this is because the Lagrangian fibration of proposition 2.25 exists relatively). Thus, one can apply theorem 3.2 to conclude that

$$\cdot A: A_{hom}^2(X) \rightarrow A^4(X)$$

is the zero map. This implies that $A \cdot a$ is in $A_{(0)}^4(X)$, and so also b is in $A_{(0)}^4(X)$. Since $A_{(0)}^4(X) \cong \mathbb{Q}$, this proves the corollary. $\square$

Corollary 4.2. *Let $g \in \{2, 3, 4, 5, 6, 8\}$, and let $\mathcal{S} \rightarrow B$ be the universal family of genus g K3 surfaces. Let $\mathcal{C} \subset \mathcal{S}$ be a “relative curve” (i.e., a divisor flat over B), and let $C_b \subset S_b$ denote the restriction $\mathcal{C}|_{S_b}$. Let X be the Hilbert scheme $X = (S_b)^{[2]}$ for any $b \in B$, and let $Y \subset X$ be the Lagrangian surface $(C_b)^{(2)}$. Then*

$$\begin{aligned} Y &\in A_{(0)}^2(X) , \\ \cdot Y = 0: A_{hom}^2(X) &\rightarrow A^4(X) . \end{aligned}$$

Proof. This is immediate from theorem 3.1. $\square$

Remark 4.3. *In the set-up of corollary 4.2, it may be expected that $C_b \subset S_b$ is a constant cycle curve (this is Huybrechts’ guess, cf. remark 3.3). This would imply that $Y \subset X$ is a constant cycle surface, which would nicely explain the behaviour of Y exhibited in corollary 4.2.*

Corollary 4.4. *Let $S \subset \mathbb{P}^3(\mathbb{C})$ be a smooth quartic, and let $X = S^{[2]}$ be the Hilbert scheme. Let $T \subset X$ be the surface of bitangents. Then*

$$\begin{aligned} T &\in A_{(0)}^2(X), \\ \cdot T = 0: A_{hom}^2(X) &\rightarrow A^4(X). \end{aligned}$$

Proof. The surface $T \subset X$ exists relatively, and so this follows from theorem 3.1. $\square$

Remark 4.5. *When $S \subset \mathbb{P}^3(\mathbb{C})$ is a generic quartic, there is Beauville’s anti-symplectic involution $\iota: X \rightarrow X$ [1]. The fixed locus of ι is the surface of bitangents T [6, Example 3.5]. The first part of corollary 4.4 thus follows from the more general statement that*

$$A^2(X)^\iota \subset A_{(0)}^2(X),$$

which is proven in [25] (cf. also [12, Corollary 1.8]).

As for the second statement of corollary 4.4, it is perhaps possible to prove the stronger statement that $T \subset X$ is a constant cycle surface. (Since ι acts as minus the identity on $A_{(2)}^4(X)$ [25], we know that

$$\text{Im}(A_0(T) \rightarrow A^4(X)) \cap A_{(2)}^4(X) = 0.$$

To prove T is a constant cycle surface, it only remains to prove that

$$\text{Im}(A_0(T) \rightarrow A^4(X)) \cap A_{(4)}^4(X) \stackrel{??}{=} 0.$$

I don’t know how to do this.)

5. FANO VARIETIES OF CUBIC FOURFOLDS

In this final section, we prove a “generalized Franchetta conjecture” type of statement for the Fano variety of lines on a cubic fourfold containing two planes. The argument is similar to that of theorem 3.1.

Theorem 5.1. *Let $\mathcal{Y} \rightarrow B$ be the universal family of smooth cubic fourfolds containing two disjoint planes, and let $\mathcal{X} \rightarrow B$ be the family of associated Fano varieties of lines (cf. notation 2.29). Let $\Gamma \subset \mathcal{X}$ be a codimension 2 subvariety. The restriction $\Gamma|_{X_b} \in A^2(X_b)$ is in $A_{(0)}^2(X_b)$ for all $b \in B$.*

Proof. Let us consider the relative correspondence

$$\Gamma_0 := (\Pi_2^\mathcal{X})_*(\Gamma) \in A^2(\mathcal{X}),$$

where $\Pi_2^\mathcal{X}$ is the relative CK projector as in proposition 2.16. We know that there is fibrewise homological vanishing

$$(17) \quad \Gamma_0|_{X_b} = 0 \text{ in } H^4(X_b) \quad \forall b \in B.$$

What we need to prove (in order to have $\Gamma \in A_{(0)}^2(X_b)$) is that there is a fibrewise rational equivalence

$$(18) \quad \Gamma_0|_{X_b} \stackrel{??}{=} 0 \text{ in } A^2(X_b) \quad \forall b \in B.$$

To obtain this, we move things to the universal family of (squares of) related $K3$ surfaces. That is, we first consider the relative correspondence

$$\Gamma_1 := (\Psi_1)_*(\Gamma_0) \in A^4(\mathcal{Y} \times_B \mathcal{Y}) \oplus \bigoplus A^3(\mathcal{Y}),$$

where $\Psi_1 \in A^6(\mathcal{X} \times_B (\mathcal{Y} \times_B \mathcal{Y})) \oplus \bigoplus A^*(\mathcal{X} \times_B \mathcal{Y})$ is as in corollary 2.28. Let us write

$$\Gamma_1 = \Gamma_1^0 + \Gamma_1^1 + \cdots + \Gamma_1^m,$$

where Γ_1^0 is in $A^4(\mathcal{Y} \times_B \mathcal{Y})$ and Γ_1^i is in $A^3(\mathcal{Y})$ for $i = 1, \dots, m$.

Next, we consider the relative correspondence

$$\Gamma_2 := (\Psi_3)_*(\Gamma_1^0) + \sum_{i=1}^m (\Psi_2)_*(\Gamma_1^i) \in A^2(\mathcal{S} \times_B \mathcal{S}) \oplus \bigoplus_{j=1}^{m+2} A^2(\mathcal{S}),$$

where Ψ_2 and Ψ_3 are as in lemma 2.30 resp. lemma 2.31. Let us split this correspondence into two parts, writing

$$\Gamma_2 = \Gamma'_2 + \Gamma''_2$$

with

$$\Gamma'_2 \in A^2(\mathcal{S} \times_B \mathcal{S}), \quad \Gamma''_2 \in \bigoplus A^2(\mathcal{S}).$$

Property (17) implies there is a fibrewise homological vanishing

$$(19) \quad (\Gamma'_2 + \Gamma''_2)|_{S_b \times S_b} = 0 \quad \text{in } H^4(S_b \times S_b) \oplus \bigoplus H^4(S_b) \quad \forall b \in B.$$

The family $\mathcal{S} \rightarrow B$ fulfills the conditions of proposition 2.22 (with $M = \mathbb{P}^2 \times \mathbb{P}^2$ and $L_1 = \mathcal{O}_M(1, 2)$ and $L_2 = \mathcal{O}_M(2, 1)$). Applying proposition 2.22 to Γ'_2 , we find that there exists $\delta \in A^2(M \times M)$ such that

$$\Gamma'_2|_{S_b \times S_b} = \delta|_{S_b \times S_b} \quad \text{in } A^2(S_b \times S_b) \quad \forall b \in B.$$

Combined with (19), this implies in particular that

$$\delta|_{S_b \times S_b} \in A_{hom}^2(S_b \times S_b) \quad \forall b \in B.$$

But $A^*(M \times M) = A^*(M) \otimes A^*(M)$ is generated by divisors, and so the restriction $\delta|_{S_b \times S_b}$ is in the subring generated by divisors. It is known this subring injects into cohomology [42], and so

$$\delta|_{S_b \times S_b} = 0 \quad \text{in } A^2(S_b \times S_b) \quad \forall b \in B.$$

It follows that also

$$(20) \quad \Gamma'_2|_{S_b \times S_b} = 0 \quad \text{in } A^2(S_b \times S_b) \quad \forall b \in B.$$

Next, let us consider the part Γ''_2 . We have seen (equality (19)) that this cycle is fibrewise homologically trivial:

$$\Gamma''_2|_{(S_b)^{\oplus m+2}} = 0 \quad \text{in } \bigoplus_{j=1}^{m+2} H^4(\mathcal{S}) \quad \forall b \in B.$$

It follows from proposition 2.33 that there is a rational equivalence

$$(21) \quad \Gamma_2''|_{(S_b)^{\oplus m+2}} = 0 \quad \text{in} \quad \bigoplus_{j=1}^{m+2} A^2(\mathcal{S}) \quad \forall b \in B.$$

Combining (20) and (21), we find that

$$(22) \quad \Gamma_2|_{(S_b \times S_b) \cup (S_b)^{\oplus m+2}} = 0 \quad \text{in} \quad A^2(S_b \times S_b) \oplus \bigoplus_{j=1}^{m+2} A^2(S_b).$$

Let us now get back to the Fano varieties X_b . We know that

$$(\Psi_1): A_{hom}^2(X_b) \rightarrow A_{hom}^4(Y_b \times Y_b) \oplus \bigoplus_{i=1}^m A_{hom}^3(Y_b)$$

is injective (corollary 2.28). Likewise, we know that

$$(\Psi_2)_*: A_{hom}^3(Y_b) \rightarrow A^2(S_b),$$

$$(\Psi_3)_*: A_{hom}^4(Y_b \times Y_b) \rightarrow A^2(S_b \times S_b) \oplus A^2(S_b)^{\oplus 2}$$

are injective, at least for general $b \in B$ (lemmas 2.30 and 2.31).

Looking at the definition of Γ_2 , the vanishing (22) thus implies that

$$\Gamma_0|_{X_b} = 0 \quad \text{in} \quad A^2(X_b) \quad \text{for very general } b \in B.$$

In view of [45, Lemma 3.2], this implies that actually

$$\Gamma_0|_{X_b} = 0 \quad \text{in} \quad A^2(X_b) \quad \forall b \in B$$

as desired, proving equality (18) and hence theorem 5.1. $\square$

Remark 5.2. *One can also prove the analogue of theorem 3.2 for Fano varieties X_b as in theorem 5.1. The details will appear elsewhere.*

Acknowledgements. *Thanks to the referee for pertinent and helpful comments. Thanks to all the members of Het Lieve Team, in particular Kai-kun, Len-kun & Yoyo-chan.*

REFERENCES

- [1] A. Beauville, Some remarks on Kähler manifolds with $c_1 = 0$, in: Classification of algebraic and analytic manifolds (Katata, 1982), Birkhäuser Boston, Boston 1983,
- [2] A. Beauville, Variétés Kähleriennes dont la première classe de Chern est nulle, J. Differential Geom. 18 no. 4 (1983), 755—782,
- [3] A. Beauville and R. Donagi, La variété des droites d’une hypersurface cubique de dimension 4, C. R. Acad. Sci. Paris Sér. I Math. 301 no. 14 (1985), 703—706,
- [4] A. Beauville and C. Voisin, On the Chow ring of a $K3$ surface, J. Alg. Geom. 13 (2004), 417—426,
- [5] A. Beauville, On the splitting of the Bloch–Beilinson filtration, in: Algebraic cycles and motives (J. Nagel et al., eds.), London Math. Soc. Lecture Notes 344, Cambridge University Press 2007,

- [6] A. Beauville, Antisymplectic involutions of holomorphic symplectic manifolds, *Journal of Topology* 4 no. 2 (2011), 300—304,
- [7] S. Bloch, *Lectures on algebraic cycles*, Duke Univ. Press Durham 1980,
- [8] A. Ferretti, The Chow ring of double EPW sextics, *Algebra and Number Theory* 6(3) (2009),
- [9] B. Fu, Abelian fibrations on $S^{[n]}$, *C. R. Math. Acad. Sci. Paris* 337 no. 9 (2003), 593—596,
- [10] L. Fu, Beauville–Voisin conjecture for generalized Kummer varieties, *Int. Math. Res. Notices* 12 (2015), 3878—3898, 2014,
- [11] L. Fu, On the action of symplectic automorphisms on the CH_0 -groups of some hyper-Kähler fourfolds, *Math. Z.* 280 (2015), 307—334,
- [12] L. Fu, R. Laterveer and Ch. Vial, The generalized Franchetta conjecture for some hyper-Kähler varieties (with an appendix joint with M. Shen), arXiv:1708.02919v2, to appear in *Journal de Math. Pures et Appliquées*,
- [13] L. Fu, Z. Tian and C. Vial, Motivic hyperkähler resolution conjecture: I. Generalized Kummer varieties, to appear in *Geom. Topol.*,
- [14] W. Fulton, *Intersection theory*, Springer–Verlag Ergebnisse der Mathematik, Berlin Heidelberg New York Tokyo 1984,
- [15] S. Galkin and E. Shinder, The Fano variety of lines and rationality problem for a cubic hypersurface, arXiv:1405.5154,
- [16] B. Hassett, Some rational cubic fourfolds, *J. Alg. Geom.* 8(1) (1999), 103—114,
- [17] B. Hassett, Cubic fourfolds, $K3$ surfaces and rationality questions, in: *Rationality problems in algebraic geometry* (R. Pardini et al., eds.), Springer Lecture Notes in Mathematics, Springer 2016,
- [18] B. Hassett and Y. Tschinkel, Abelian fibrations and rational points on symmetric products, *International Journal of Mathematics* 11(9) (2000), 1163—1176,
- [19] D. Huybrechts, Curves and cycles on $K3$ surfaces, *Algebraic Geometry* 1 (2014), 69—106,
- [20] A. Iliev and K. Ranestad, The abelian fibration on the Hilbert cube of a $K3$ surface of genus 9, *International Journal of Mathematics* 18(1) (2007),
- [21] U. Jannsen, Motivic sheaves and filtrations on Chow groups, in: *Motives* (U. Jannsen et al., eds.), *Proceedings of Symposia in Pure Mathematics* Vol. 55 (1994), Part 1,
- [22] U. Jannsen, On finite-dimensional motives and Murre’s conjecture, in: *Algebraic cycles and motives* (J. Nagel et al., eds.), Cambridge University Press, Cambridge 2007,
- [23] S. Kimura, Chow groups are finite dimensional, in some sense, *Math. Ann.* 331 (2005), 173—201,
- [24] R. Laterveer, A remark on the motive of the Fano variety of lines of a cubic, *Ann. Math. Québec* 41 no. 1 (2017), 141—154,
- [25] R. Laterveer, Bloch’s conjecture for some hyperkähler fourfolds, to appear in *Pure and Applied Math. Quarterly*,
- [26] H. Lin, On the Chow groups of zero-cycles of a generalized Kummer variety, *Advances in Mathematics* 298 (2016), 448—472,
- [27] H. Lin, Lagrangian constant cycle subvarieties in Lagrangian fibrations, to appear in *Int. Math. Res. Notices*,
- [28] S. Mukai, Symplectic structure of the moduli space of sheaves on an abelian or $K3$ surface., *Invent. Math.* 77 no. 1 (1984), 101—116,
- [29] S. Mukai, Curves, $K3$ surfaces and Fano 3-folds of genus ≤ 10 , in: *Algebraic geometry and commutative algebra* (H. Hijikata et al., eds.), Kinokuniya Tokyo 1988,
- [30] J. Murre, On a conjectural filtration on the Chow groups of an algebraic variety, parts I and II, *Indag. Math.* 4 (1993), 177—201,
- [31] J. Murre, J. Nagel and C. Peters, *Lectures on the theory of pure motives*, Amer. Math. Soc. University Lecture Series 61, Providence 2013,
- [32] K. O’Grady, Moduli of sheaves and the Chow group of $K3$ surfaces, *Journal de Math. Pures et Appliquées* 100 no. 5 (2013), 701—718,

- [33] N. Pavic, J. Shen and Q. Yin, On O’Grady’s generalized Franchetta conjecture, *Int. Math. Res. Notices* 2016 (2016), 1—13,
- [34] U. Rieß, On the Chow ring of birational irreducible symplectic varieties, *Manuscripta Math.* 145 (2014), 473—501,
- [35] U. Rieß, On Beauville’s conjectural weak splitting property, *Int. Math. Res. Notices* 2016 no. 20 (2016), 6133—6150,
- [36] J. Sawon, Lagrangian fibrations on Hilbert schemes of points on $K3$ surfaces, *Journal of Alg. Geom.* 16(3) (2007), 477—497,
- [37] T. Scholl, Classical motives, in: *Motives* (U. Jannsen et alii, eds.), *Proceedings of Symposia in Pure Mathematics* Vol. 55 (1994), Part 1,
- [38] M. Shen and C. Vial, The Fourier transform for certain hyperKähler fourfolds, *Memoirs of the AMS* 240 (2016), no.1139,
- [39] M. Shen and C. Vial, The motive of the Hilbert cube $X^{[3]}$, *Forum Math. Sigma* 4 (2016),
- [40] C. Vial, Remarks on motives of abelian type, *Tohoku Math. J.* 69 (2017), 195—220,
- [41] C. Vial, On the motive of some hyperkähler varieties, *J. für Reine u. Angew. Math.* 725 (2017), 235—247,
- [42] C. Voisin, On the Chow ring of certain algebraic hyperkähler manifolds, *Pure Appl. Math. Q.* 4 no. 3 part 2 (2008), 613—649,
- [43] C. Voisin, The generalized Hodge and Bloch conjectures are equivalent for general complete intersections, *Ann. Sci. Ecole Norm. Sup.* 46, fascicule 3 (2013), 449—475,
- [44] C. Voisin, The generalized Hodge and Bloch conjectures are equivalent for general complete intersections, II, *J. Math. Sci. Univ. Tokyo* 22 (2015), 491—517,
- [45] C. Voisin, *Chow Rings, Decomposition of the Diagonal, and the Topology of Families*, Princeton University Press, Princeton and Oxford, 2014,
- [46] C. Voisin, Hodge structures, Coniveau and Algebraic Cycles, in: “The Legacy of Bernhard Riemann After One Hundred and Fifty Years”, *ALM35*, Higher Education Press and International Press, Beijing Boston 2016,
- [47] C. Voisin, Remarks and questions on coisotropic subvarieties and 0–cycles of hyper–Kähler varieties, in: *K3 Surfaces and Their Moduli*, *Proceedings of the Schiermonnikoog conference 2014* (C. Faber et al., eds.), *Progress in Maths* 315, Birkhäuser 2016,
- [48] C. Voisin, On the universal CH_0 group of cubic hypersurfaces, *Journal Eur. Math. Soc.* 19 no. 6 (2017), 1619—1653.

INSTITUT DE RECHERCHE MATHÉMATIQUE AVANCÉE, CNRS – UNIVERSITÉ DE STRASBOURG, 7 RUE RENÉ DESCARTES, 67084 STRASBOURG CEDEX, FRANCE.

E-mail address: robert.laterveer@math.unistra.fr